1 **Title**
2
3 # Impact of different mutations on Kelch13 protein levels, ART
4 # resistance and fitness cost in *Plasmodium falciparum* parasites
5

6 **Authors**
7 Hannah M. Behrens[1#], Sabine Schmidt[1#], Domitille Peigney[1], Ricarda Sabitzki[1], Isabelle Henshall[1],
8 Jürgen May[2,3], Oumou Maïga-Ascofaré[2,3], Tobias Spielmann[1]*
9
10
11 **Affiliations**
12 [1]Malaria Cell Biology, Molecular Biology and Immunology, Bernhard Nocht Institute for Tropical
13 Medicine, Bernhard-Nocht-Str. 74, 20359 Hamburg, Germany.
14
15 [2]Infectious Disease Epidemiology Department, Epidemiology and Diagnostics, Bernhard Nocht
16 Institute for Tropical Medicine, Bernhard-Nocht-Str. 74, 20359 Hamburg, Germany.
17
18 [3]German Centre for Infection Research (DZIF), Partner Site Hamburg-Luebeck-Borstel-Riems,
19 Hamburg, Germany.
20
21 [#]equal contribution
22
23 *Tobias Spielmann
24 **Email:** spielmann@bnitm.de

25
26

Abstract
Reduced susceptibility to ART, the first-line treatment against malaria, is common in South East Asia
(SEA). It is caused by point mutations, mostly in *kelch13* (*k13*) but also in other genes, like *upb1*. K13
and its compartment neighbors (KICs), including UBP1, are involved in endocytosis of host cell cytosol.
We tested 135 mutations in KICs but none conferred ART resistance. Double mutations of *k13*C580Y
with *k13*R539T or *k13*C580Y with *ubp1*R3138H, did also not increase resistance. In contrast, *k13*C580Y
parasites subjected to consecutive RSAs did, but the *k13* sequence was not altered. Using isogenic
parasites with different *k13* mutations, we found correlations between K13 protein amount,
resistance and fitness cost. Titration of K13 and KIC7 indicated that the cellular levels of these proteins
determined resistance through the rate of endocytosis. While fitness cost of *k13* mutations correlated
with ART resistance, *ubp1*R3138H caused a disproportionately higher fitness cost.

38
Significance
ART resistance is only a partial resistance with a proportion of ring stages surviving physiological
ART levels. The correlation of resistance with fitness cost in isogenic lines indicates that fitness cost
likely prevents resistance levels permitting survival of all ring stages under physiological ART
concentrations. We also found no indication that double mutations in *k13,* including the two most
common resistance mutations in SEA, or with non-*k13* genes, are a threat to lead to higher
resistance. However, repeated ART exposure increased resistance by mechanisms not altering *k13*
gene sequence, indicating changes in the background of these parasites. The disproportionally high

fitness cost of *ubp1*R3138H may explain why *kic* mutations affecting resistance are rare and
highlights the unique property of K13 to influence endocytosis only in ring stages.
49
50
51   **Introduction**
52
Malaria kills approximately half a million people per year, most of them children in sub-Saharan Africa
(*1*, *2*). Treatment heavily relies on artemisinin and its derivatives (ARTs) which are typically used in
artemisinin-based combination therapies (ACTs) with a partner drug (*3*). More than 10 years ago
reduced susceptibility to artemisinin was observed in low malaria incidence settings such as Asia,
Oceania and South America. This manifests as delayed parasite clearance after treatment (*3*, *4*),
caused by some ring stage parasites surviving the ART-pulse, which is brief (a few hours) because of
the low half-life of ART drugs (*5*) and can lead to treatment failure (*6*). *In vitro* this reduced
susceptibility is measured by the ring stage survival assay (RSA) where survival of >1% of parasites
indicates what we call *in vitro* ART resistance (*7*).
62
The main cause of reduced ART susceptibility are mutations in the *Pfkelch13* gene (*k13*) encoding the
K13 protein (*8*, *9*), which was recently found to be involved in endocytosis of host cell cytosol in the
ring stage of the parasite (*10*). Ten mutations affecting the C-terminal propeller domain of K13 are
confirmed to cause reduced susceptibility as evident from delayed clearance in patients (F446I,
N458Y, M476I, Y493H, R539T, I543T, P553L, R561H, P574L, C580Y) and others have been associated
with reduced susceptibility (*3*). Several other proteins, mostly from the K13-compartment, confer *in*
vitro ART resistance in RSA either when downregulated or when mutated (*10–14*). One of them is
ubp1, for which the R3138H was shown to result in moderate resistance in RSA (*10*) and which was
associated with reduced susceptibility in patient samples from Asia, although it occurred only rarely
(*15*). The contribution of mutations affecting K13 compartment proteins (other than K13) to reduced
ART susceptibility in endemic countries is largely unknown.
74
75   Resistant clinical isolates and laboratory parasites carrying the resistance-associated mutations *k13*
76   R539T and *k13* C580Y contain reduced amounts of K13 (*10*, *16–18*). Artificially modulating K13 levels
77   indicated that the reduced K13 levels lead to resistance (*10*, *16*, *19*). It is currently unclear whether
78   other *k13* mutations, like R561H, confer resistance through the same mechanism or affect the
79   functionality of K13.
80
While ART-resistance mutations are frequent in South East Asia, up to 100% in some locations (*20–*
22), they are rare in Africa (*21*, *23*, *24*). Six validated resistance-conferring *k13* mutations (M476I,
P553L, R561H, P574L, C580Y and A675V) have occasionally been reported in African countries at
moderate to low frequency of 4.1% or less (*25–27*). Two exceptions are the R561H mutation in
Rwanda, which was reported more frequently since 2019, most recently up to 16% (*28–30*) and the
11% of A675V mutation in Uganda in 2019 (*27*). It is not clear whether further resistance mutations
exist in Africa. Most studies focused on mutations occurring in Asia and there is no systematic *k13*
surveillance in Africa. In agreement with the rare occurrence of resistance mutations, delayed
parasite clearance occurs at less than 2% in Africa (*31*), with the exception of the Masaka region in
Rwanda at 16%, which is where *k13* R561H occurs (*30*). It was postulated that resistance would
emerge later in Africa than in Asia because of the higher transmission rate and higher level of
immunity (*32*, *33*).
93
The reduced endocytosis in ART resistant parasites deprives the parasite of amino acids (*34*) and
there is clear evidence for a fitness cost of ART resistance (*9*, *35*). Usually considerably less than 50%
of parasites from resistant patient isolates survive in the RSA (*8*) and the delayed clearance rather
than full loss of susceptibility in patients could indicate that it is possible for parasites with higher
levels of ART resistance to arise. However, it is at present unclear if and to what extent the fitness

cost impedes higher levels of resistance. It is also unclear if combinations of mutations could lead to
hyper-resistant parasites.

Here we test 135 mutations in proteins from the K13-compartment including 4 mutations in K13 itself
and find that none of them confers significant levels of resistance over GFP-tagged wild type (WT)
K13. However, some of the *k13* mutations resulted in a significantly lower susceptibility to ART than
the 3D7 background. Together with known high resistance mutations this provided us with 3D7-based
laboratory lines with different resistance levels. A comparison of parasites with different resistance
levels showed a correlation between resistance, fitness cost and K13 protein abundance of a given
variant, suggesting a constraint of resistance levels by the fitness cost. This was confirmed using
parasites artificially selected for higher resistance. We also assess the threat of hyper-resistance, the
impact on fitness cost on this and the potential impact of *k13* double mutations and combinations
with resistance mutations outside *k13*.

Results

Tested mutations in K13-compartment proteins do not cause *in vitro* ART resistance
We aimed to assess whether further mutations in addition to the already identified resistance
mutations could cause reduced ART susceptibility. As not only *k13*, but also genes of K13
compartment proteins such as *ubp1* can cause reduced susceptibility, we analyzed non-synonymous
mutations in *k13* and genes of other K13 compartment proteins from African patient samples
available from WWARN (*36*), the Ghanaian Fever Without Source study (*37*) and MalariaGen (*38*).
These included mutations in the genes encoding K13, KIC1, KIC2, KIC4, KIC5, KIC7, KIC9, UBP1,
EPS15and MyoF (previously annotated as MyoC) (list of mutations in Table S1). In addition we
selected one untested mutation in AP2α, which was reported to be present in *in vitro* ART-selected
parasites that displayed a K13-independent reduced ART-susceptibility (*39*). Due to the large number
of mutations in the K13 compartment proteins (a total of 131 mutations in ten genes) we created
parasites carrying multiple mutations (multi-mutants) by introducing a recodonized sequence of the
target gene that encodes several mutations in the same gene into 3D7 parasites using selection-linked
integration (SLI, (*40*)) (Table S1). In total 125 mutations were included in 8 multi-mutants and 6
mutations were tested individually (Figure 1A,B). RSAs (using dihydroartemisinin (DHA)) with multi-
mutants of compartment members of K13 showed no change in susceptibility to ART, suggesting that
none of the mutations in the genes encoding KIC1, KIC2, KIC4, KIC5, KIC7, KIC9, UBP1, EPS15 and MyoF
resulted in ART resistance (Figure 1A,B). One exception was AP2α H817P which significantly increased
RSA survival, although this was not above 1% which is considered the cut-off for clinically relevant
resistance (*7*). Yet this increase might explain the occurrence of this mutation in parasites that had
been selected for ART-resistance through ART exposure *in vitro* but that lacked K13 mutations (*39*).

For *k13* itself we investigated the mutations V520A, V589I and E612K that have been found with low
to moderate prevalence (0.8%-5.0%) in different malaria endemic regions in Africa and were not
previously tested for ART resistance *in vitro* (Figure 1C)*.* All mutations originated from studies where
the samples had been taken after ACT started to be used in the respective country, with exception of
one study in Kenya in 2002 (Dataset S1). We noticed that *k13* V520A and V589I occurred in regions
with above average and average malaria incidence, respectively, while *k13* E612K, C580Y, R539T and
k13 R561H, which was recently detected in Rwanda, occurred in regions with below average malaria
incidences (Figure S1, Dataset S2). E252Q occurred in Thailand and Myanmar between 2005 and 2013
where it was previously associated with slow parasite clearance (*36*) but not tested *in vitro.* Although
only detected in Asia, we included this mutation because it might have been missed in Africa as
typically only the propeller domain (amino acid residues 444–721) is sequenced when screening for
k13 mutations. The respective mutations were introduced into the *k13* locus of *P. falciparum* 3D7
parasites together with fluorescent protein GFP to result in a GFP-K13 fusion as done previously (*10*).
The susceptibility of these parasites to ART was tested by RSA. Introduction of the GFP to the *k13*

locus already led to small but significant reduction in ART-susceptibility (0.2±0.2% vs 0.6% ±0.3%)
(K13-WT, Figure 1D), suggesting that addition of the GFP to the protein's N-terminus resulted in a
very mild reduction in K13 activity. All three mutations caused a small further decrease in ART-
susceptibility although this was not significantly different over *k13* WT but for *k13* V520A and E252Q
was significant over the 3D7 parent (Figure 1D). We conclude that these mutations have likely no
relevance for ART-resistance in endemic settings but provide us with parasites with a mildly reduced
ART-susceptibility compared with 3D7.
158
159 For comparison, we also tested *k13* C580Y, R561H and R539T, which were previously shown to be
160 resistance mutations. Parasites with C580Y (16.3% ±6.1%), R561H (11.1% ±8.0%) and R539T (38.4%
161 ±13.2%) *k13* showed higher RSA survival (Figure 1C), similar to what was previously observed for
162 C580Y and R539T in 3D7 (*8*).
163
164 **Double mutations in *k13* do not significantly increase ART resistance**
165
To anticipate how *k13* resistance phenotypes could further develop in endemic areas, we investigated
whether combination of the *k13* C580Y mutation (which is highly prevalent in SE Asia) with the V520A
mutation might potentiate resistance e.g. through a synergetic effect. Further we investigated
whether the two known resistance mutations *k13* C580Y and R539T could combine to result in hyper-
resistance. To obtain the corresponding cell lines, we again modified the endogenous *kelch13* locus
of 3D7 parasites using SLI so that the double-mutated K13 was fused to GFP. The K13$^{V520A+C580Y}$
parasites did not display an increased resistance level in RSAs (11.5% survival) compared with
parasites with only the C580Y mutation (16.3% survival) (Figure 2A), indicating that there is no
synergistic effect. Similarly, K13$^{R539T+C580Y}$ parasites showed no significant change in RSA survival
(33.4% survival) compared with parasites harboring R539T alone (38.4% survival) but significantly
more survival than the C580Y alone, suggesting no additive or synergistic effect. In conclusion, —at
least with the tested mutations — there is no indication for additive or synergistic effects.
178
179 **Selection by consecutive RSAs significantly increases ART resistance in *k13* C580Y parasites**
180
In a further attempt to anticipate increased resistance, we tested whether parasites with a common
resistance mutation can be selected for increased resistance by performing consecutive RSAs. For
this, an RSA was carried out with K13^{C580Y} parasites and surviving parasites were subjected to a
consecutive RSA as soon as they reached sufficient parasitemia following the previous RSA (Figure
2B). After 29 iterations of RSAs, we obtained the parasite line K13^{C580Y}-29th. These parasites displayed
significantly increased survival in RSA (39.2%) compared with the original K13^{C580Y} parasites at the
start of this experiment (12.4%) (Figure 2C), showing that repeated ART-treatment of *k13* C580Y-
harbouring parasites can increase ART resistance. Sequencing of *k13* in these parasites showed that
this was not due to further mutations in *k13* in addition to the C580Y already present at the beginning
of the experiment (Figure S2).
191
192
193 **Cellular levels of K13 and KIC7 proteins determine ART-resistance**
194
Previously, we and others showed that the level of K13 in the K13^{C580Y} and K13^{R539T} parasites is lower
than in K13^{wt} parasites and that reducing K13 abundance resulted in resistance (*10*, *16*, *18*). We
assessed the cellular K13 levels in the parasites for all *k13* mutations in this study, to find whether a
change in K13 level generally accompanies resistance mutations. We previously used either Western
blots or direct microscopy-based measurement of GFP fluorescence in the cells to determine K13
levels (*10*). Here we used direct microscopy-based measurement of K13 levels because in comparison
to Western blots, which are only semi-quantitative, this procedure is not susceptible to influence of
cell lysis and the multiple steps required for blotting and detection. These experiments showed no

significant reduction in the amount of K13 with the V520A (94% of WT) and the V589I (91% of WT) mutations (Figure 3A). The E252Q mutation showed 105% of the amount (non-significant) of WT (Figure 3A). The resistance mutations *k13* C580Y, *k13* R561H and *k13* R539T had 52%, 51% and 51% of the K13 amount of WT, respectively (Figure 3A). As expected from their RSA resistance behavior, parasites with the *k13* V520A C580Y double mutation harbored similar levels of K13 (48% of WT) as parasites with the *k13* C580Y alone, and parasites with *k13* R539T and C580Y (57% of WT) had similar levels as both *k13* C580Y and *k13* R539T individually. The K13 levels in K13[C580Y]-29th parasites showed a further reduction (32% of WT) when compared with K13[C580Y] parasites, even though this reduction was not significant when comparing means (Figure 3A), only when comparing all data points (Figure S3). We plotted the cellular K13 levels against RSA survival (*in vitro* resistance). Cellular K13 levels correlated with *in vitro* resistance (Pearson's r=-0.81, p=0.005) (Figure 3B). We note that this was largely because the mutations with low K13 levels clustered into a group conferring significant *in vitro* resistance over GFP-K13-WT and the other mutations into a group with high K13 levels (Figure 3B).

To test whether RSA-resistance not just correlates with but is determined by the amount of K13 in its cellular location, we used K13 WT 1xNLS parasites (*10*). In this line K13 protein is removed into the nucleus by a nuclear localization signal (NLS) upon addition of rapalog (termed knock sideways), resulting in conditional reduction of K13 at its cellular location. Previously, 250 nM rapalog was used to achieve a knock sideway of K13 (*40*). Here, we used different concentrations of rapalog to result in different levels of K13 protein remaining at its site of action. ART-resistance, measured by RSA, decreased as the rapalog concentration and hence K13 knock sideways decreased, confirming that RSA-survival is a function of K13 levels in the parasites (Figure 3C). These results confirm observations made on parasites in which the K13 abundance was titrated on mRNA level through the glmS system (*19*). To confirm that this was due to a reduction in endocytosis, we performed a second titration using KIC7, a K13-compartment protein, which was also knocked aside using the same decreased rapalog concentrations as for K13 (Figure 3D). Again, RSA survival was dependent on the level of knock sideways of KIC7 (i.e. amount of KIC7 at the K13 compartment), indicating that the activity of the endocytic process determines resistance and that this occurs through the available amount of some of the proteins involved in this process.

Fitness cost correlates with ART-resistance

Several studies modelling fitness and within-host competition have highlighted the importance of parasite fitness and propose that resistant parasites with reduced fitness are less likely to establish themselves in high transmission areas like most of Africa (*33*, *41*, *42*). Some K13 resistance mutations, including C580Y and R561H, have been shown to have a fitness costs (*22*, *43*). To measure the impact of *k13* mutations on fitness of our parasites with isogenic backgrounds in this work, we first monitored *in vitro* growth over 96 h. This confirmed the fitness cost of *k13* C580Y and suggested that generally, fitness cost might correlate with resistance but the variation between experiments was too high for reliable conclusions (Figure 4A,B).

For better fitness cost measurements, parasite proportions were determined in a mixed culture of a mutant and the 3D7 parent parasite line, as previously described (*22*, *44*). Parasites were mixed 1:1 with the 3D7 parasites and the proportion of mutated (fluorescent) parasites was tracked until it made up less than 5% of the parasite population (Figure 4C ,S3). The parasites with K13 mutations (V520A, V589I, E252Q, C580Y and C580Y-29th) declined in proportion compared to 3D7 in all 6 experiments whereas GFP-K13 WT did not or only to a small extent (Figure S3). Similar to what was observed when parasites growth was tracked in individual cultures, there was a trend that the RSA-survival caused by a mutation correlated with the fitness cost inflicted by the mutation (Figure 4D). However, this trend was not significant (Pearson's r=0.69, p=0.13) because *k13* V520A behaved as an outlier in half of the experimental repeats, in which its growth slowed severely after the start of the experiment (Figure S4), while in the other half it grew in line with the trend of the other mutations

255　(Pearson's r=0.90, p=0.04 when V520A was entirely excluded) (Figure 4C,D). Together with the
256　findings on K13 abundance (Figure 3), it is therefore likely that both fitness cost and ART-resistance
257　are caused by the reduced amount of K13 in the cell as a result of the destabilization of the K13
258　protein through the given mutation.

259
260
261　***ubp1* R3138H has a disproportional fitness cost that is further aggravated in K13^{C580Y} parasites**
262
After assessing the effect of the *k13* double mutations V520A+C580Y and R539T+C580Y, we
wondered whether there are combinations of mutations or gene disruptions beyond *k13* that harbour
the risk of very high resistance levels. We tested this by combining *k13* C580Y with *ubp1* R3138H and
the resulting K13^{C580Y}+UBP1^{R3138H} parasites did not display a significantly increased resistance level
(17.3% mean survival) compared with parasites with the C580Y mutation alone (Figure 5A). We also
attempted to disrupt both *kic4* and *kic5*, two genes we were previously able to disrupt individually
(*10*), or combine their disruption with UBP1^{R3138H} but were unsuccessful despite six attempts per
combination, which may indicate that the fitness cost of combining these prevented us from
obtaining these parasite lines.

272
We further characterized the K13^{C580Y}+UBP1^{R3138H} parasites and found that they displayed a similar
K13 abundance to resistance level ratio (Figure 5B,C) as the *k13* mutation (Figure 3B). The growth of
K13^{C580Y}+UBP1^{R3138H} parasites was slower than that of K13^{C580Y} parasites and UBP1^{R3138H} parasites,
which each confer a fitness cost, both when measured in individual cultures and direct competition
assays (Figure 5D-G). The ratio of fitness cost to resistance level was higher than for *k13* mutations
alone (Figure 5E,G). To determine whether just of the combination of the two mutations led to such
a reduced growth rate or whether it was due to *ubp1* R3138H by itself, we determined the growth
rate of UBP1^{R3138H} parasites. Interestingly, already the *ubp1* R3138H mutation alone caused a
disproportionally low growth compared with the level of ART resistance it affords when compared to
all tested *k13* mutations (Figure 5E). This is congruent with our previous finding that in contrast to
K13 which is only required in rings, UBP1 is also important for hemoglobin endocytosis in trophozoites
(*10*) and could explain why we were not able to combine this mutation with *kic4* and *kic5* TGDs and
why this mutation has rarely been observed in patient samples. Overall, due to its disproportionally
high fitness cost and lacking synergism with *k13* mutations, *ubp1* R3138H and possible mutations in
other K13 compartment proteins important for endocytosis in trophozoites, are therefore unlikely to
form combinations with *k13* mutations in endemic countries.

289
290　**Discussion**
291
In this study we tested 131 mutations in ten genes and found that none of the tested mutations in
the genes encoding KIC1, KIC2, KIC4, KIC5, KIC7, KIC9, UBP1, EPS15 and MyoF resulted in significantly
decreased susceptibility to ART. We also tested K13 mutations. We found that N-terminally fusing a
tag to K13, even without any mutations, very mildly reduced the susceptibility to ART. This effect is
negligible compared with the larger effects of common resistance mutations such as C580Y, R561H
and R539T and likely does not generally impact experiments with these parasites but needs to
considered when assessing mutations with small effect sizes. Thus, we found that the four previously
unstudied *k13* mutations did not cause significant resistance in the RSA compared with K13-WT. The
small non-significant decrease in susceptibility that was observed for *k13* V520A, V589I and E252Q
likely has no clinical impact but we can't exclude that these variants have some small advantage in
the population that is difficult to assess. However, all of these mutations that were tested in
competition assays were consistently outcompeted by 3D7 whereas GFP-K13 WT was not, indicating
that they have a small disadvantage over WT. Despite likely not being clinically relevant, parasites
with a GFP-fused K13 with or without the V520A, E252Q and E612K were significantly less susceptible

to ART in RSA compared to 3D7 and together with the high resistance parasites offered us a tool to
explore the properties of parasites with different levels of *in vitro* ART resistance.

We also found that the combination of *k13* C580Y with *k13* V520A, or of two resistance mutations
with each other, like *k13* C580Y and *k13* R539T, and *k13* C580Y with *ubp1* R3138H did not increase
ART resistance. These are the first combinations of C580Y with other naturally occuring mutations
generated in isogenic backgrounds. A similar non-additive result was obtained in a recent study for
coronin R100K and E107V, two *in vitro* selected mutations, combined with *k13* C580Y (*14*).

Previous work has shown that the genetic background can influence the resistance level of parasites
with identical *k13* resistance alleles (*9*, *22*), indicating that mutations outside the K13 compartment
exist that modulate the ART susceptibility of parasites with *k13* resistance mutations and remain to
be determined. Repeated ART-treatment in consecutive RSAs of *k13* C580Y-harbouring parasites
increased ART resistance, but this was not due to additional *k13* mutations apart from C580Y.
Parasites from such selections might thus be a promising tool to study factors outside of *k13* that
influence resistance levels in parasites with a resistance-conferring allel of *k13*.

We found that the amount of K13 and KIC7 at the K13 compartment in the cell determine the level of
in vitro resistance, confirming that the endocytic process is important for ART resistance and
suggesting that the amount of protein directly influences the rate of endocytosis and level of ART
susceptibility. The same effect on resistance was observed in a *k13* mRNA titration recently (*19*) and
a role of K13 protein abundance rather than a specific functional change in resistance is also
supported by the finding that increasing levels of K13 C580Y reverts resistant parasite back to
sensitive (*10*). The finding that KIC7 titration had a similar effect strengthens this conclusion and
further supports that resistance is due to altered endocytosis levels. We found that all tested
resistance mutations had low K13 levels and all other tested mutations had high K13 levels. The
protein levels in the K13 double mutants were comparable to that of the single resistance mutation,
indicating that at least in these combinations, multiple mutations do not further destabilize K13. In
conclusion, for all tested *k13* resistance mutations, the amount of K13 in the parasite, likely through
destabilization of K13, was a factor contributing to or causing the resulting resistance.

We further showed that fitness cost of *k13* mutations generally correlated with resistance. This agrees
with a recent study looking at *k13* M579I, C580Y and R561H in several African strains including 3D7
although no such correlation was observed in the Asian strain Dd2 (*22*). It is unclear what caused
these differences between African strains and Dd2. Nonetheless, it showed that the genetic
background in which the mutations occur can influence the fitness cost and the resistance level (*22*,
35, *45*). Many differences have been observed in ART resistant parasites compared with sensitive
parasites (*17*, *20*, *46–51*) and some of these might correspond to such changes in the background that
alter the effect of K13 mutations, although some of these changes likely also are downstream effects
of the resistance mechanism (*34*). The data presented here originates from a set of isogenic 3D7-
derived parasites which differ from each other only in the studied mutations. While this limits
extrapolation of the conclusions to other parasite strains, it presents the advantage of separating
effects caused by the studied mutations from effects of the genetic background. Due to its long
presence in culture, 3D7 also avoids variations that may arise in more recently culture adapted lines
that often show reduced growth levels and may still acquire further changes during continued culture
to adapt for better growth in culture.

K13 E252Q was previously observed to inflict a fitness cost that is smaller than that of *k13* C580Y,
however, these experiments were not performed using parasites with isogenic backgrounds (*44*). Our
results confirm this finding in an isogenic background. The E252Q mutation caused low but significant
in vitro resistance over 3D7 but not over WT in our study. It was previously associated with delayed
parasite clearance in patients, suggesting that either E252Q can only confer resistance in a specific

parasite background or that it was not the cause of the delayed clearance. Regarding K13^{C580Y}-29th, it
should also be noted that the long duration of the competition experiments may permit the K13^{C580Y}-
29th to revert back to a state similar to the K13^{C580Y} parasites, as the continuous RSA selection pressure
had to be lifted in the competition assay.
362
We observed a disproportionally high fitness cost in parasites with the *ubp1* mutation conferring ART
resistance. The high fitness cost is likely due to its importance for hemoglobin endocytosis in
trophozoites. This is in agreement with the finding that lowering endocytosis by conditional
inactivation or disruption of K13 compartment proteins impairs parasite growth (*10*, *52*) and that
parasites with a *k13* ART resistance mutation are hypersensitive to low amino acids (*34*). The small
fitness cost of *k13* mutations compared to other K13-compartment proteins is likely due to its role in
endocytosis exclusively in rings (*10*). Our findings therefore highlight the unique property of K13
which stands in contrast to resistance-conferring changes in KICs that also affect endocytosis in later
stage parasites, thus incurring a higher fitness cost. This is a likely reason why *k13* is the predominant
gene mutated to cause delayed clearance in patients. Due to their high fitness cost, changes outside
k13 are less likely to arise and the resistance level, as observed with the few found in the field (*15*,
53, *54*), is low. The high fitness cost of mutations of proteins involved in endocytosis outside *k13* also
indicates that combinations of them with *k13* resistance mutations are unlikely to lead to hyper-
resistant parasites in the field. We also did not detect any indication for additive or synergistic effects
on *in vitro* ART resistance when the *ubp1* resistance mutation and *k13* C580Y were both present in
the parasite or if two *k13* mutations were combined.
379
Our findings also indicate that there seems to be no danger of hyper resistant parasites due to a
combination of the two most common resistance mutations *k13* C580Y and *k13* R539T, although we
cannot exclude that other combinations of mutations lead to increased resistance. Based on the one
tested combination it would also indicate that mutations with minimal increase in ART resistance such
as V520A are not a stepping stone for potentiated resistance when combined with current resistance
mutations. The only means by which we were able to increase *in vitro* ART resistance was a constant
RSA selection regime. As in our isogenic parasites, higher resistance was accompanied by higher
fitness costs, it is possible that fitness cost limits the level of ART resistance parasites can reach.
388
389
390   **Materials and Methods**
391
392   **Plasmid construction**
393
All primers used in this study are listed in Dataset S3. The mutations tested in *k13* and *k13*
compartment proteins were chosen to include only mutations that were present in a maximum of 5%
of samples taken in the sampling study. For the multi-mutants the pSLI-TGD plasmid (*40*) was used.
The homology region was directly linked to a synthesized functional codon-changed version of the
corresponding candidate containing all selected mutations (Genscript). The fragments of the
homology region and the mutated recodonized sequence of all candidates were cloned by Gibson
assembly into the pSLI-TGD vector via NotI/MluI, resulting in the vectors pSLI-KIC1mutpool, pSLI-
KIC2mutpool, pSLI-KIC4mutpool, pSLI-KIC5mutpool, pSLI-KIC7mutpool, pSLI-KIC9mutpool, pSLI-
UBP1mutpool and pSLI-MyosinFmutpool. Sequencing was performed to confirm absence of
undesired mutations.
404
405   The N-terminal SLI plasmid of K13, pSLI-N-GFP-2xFKBP-K13-loxP (*40*) was modified for the Kelch13
406   mutant parasites as follows: the mutation was obtained by amplifying the codon-changed synthesized
407   version of *k13* using primers encoding the mutation. The resulting two fragments were cloned into
408   the pSLI-N-GFP-2xFKBP-K13-loxP via AvrII/StuI.
409

To generate K13[V520A+C580Y] and K13[R539T+C580Y], pSLI-N-GFP-2xFKBP-K13_C580Y-loxP (*40*) was modified by amplifying *k13* from this plasmid using primers to introduce the mutation and ligating the resulting fragments into the same vector at the AvrII/XhoI sites.

The *k13* C580Y mutation was introduced into UBP1[R3138H] parasites using a SLI2a plasmid based on the pSLI-N plasmid (*40*) but containing the BSD resistance gene instead of the hDHFR gene (*55*). The hDHFR was excised using BamHI and HindIII. KIC4-TGD and KIC5-TGD plasmids were created by inserting the respective homology regions (*10*) into SLI2a plasmids based on the pSLI-TGD plasmid (*55*).

Parasite culturing and transfection

P. falciparum 3D7 parasites (*56*) were cultivated at 37°C in 0+ erythrocytes in RPMI complete medium with 0.5% Albumax (Life Technologies) with 5% hematocrit and transfected as previously described (*57*, *58*). All parasite lines used in this study are listed (Table S2). Transgenic parasites were selected using 4 nM WR99210 (Jacobus Pharmaceuticals) or 2.5 µg-ml BSD (Invitrogen). For selection-linked integration, parasites were selected as previously described using 0.9 µM DSM1 (BEI resources) or 400 µg/ml G418 (Merck) (*40*). Correct integration was confirmed by PCR as described (*40*) (Figure S5). Parasite line-specific integration check primers are listed in Dataset S3.

***In-vitro* ring-stage survival assay[0-3h] (RSA) and consecutive RSAs**

All RSAs were performed according to the standard procedure described previously (*59*). 0-3 h old rings were treated with 700 nM DHA for 6 h and cultivated for another 66 h at 37°C. Giemsa smears were taken and parasite survival rate determined by comparing the parasitemia of viable parasites after DHA against the parasitemia of the untreated control. Parasites were defined as resistant when mean survival rate exceeded the cut-off value of 1% (*59*).

For the consecutive RSA, K13[C580Y] parasites (*10*) were used. 66 h after the DHA pulse, Giemsa smears were taken and the surviving parasites of the DHA treated sample were re-cultivated in a new Petri dish. After sufficient parasitemia was reached, the parasites originating from the RSA survivors were subjected to a new RSA. This procedure was continuously repeated for 30 rounds. The *k13* gene of the resulting parasites K13[C580Y]-29[th] was sequenced and showed no changed compared to the starting cell line which harbored a recodonized *k13* with the C580Y mutation (*10*).

Fluorescence microscopy

Microscopy was performed as described earlier (*60*). A Zeiss Axio Imager M1 or M2 provided with a Hamamatsu Orca C4742-95 camera was used for imaging. Zeiss Plan-apochromat 63x or 100x oil immersion objectives with 1.4 numerical aperture were used. Images were edited using Corel Photo Paint X8 and brightness and intensity were adjusted. Images that were used for quantification were not adjusted for brightness and intensity.

Measurement of protein amount by fluorescence intensity

GFP-K13 parasites were synchronized two times using 5% sorbitol at intervals of two days. After the second sorbitol synchronization, the cell lines were cultivated at 37°C for 2 more hours and then GFP signal of the ring-stage parasites was detected by fluorescence microscopy using the 63x oil immersion objective. GFP-K13 WT parasites were always imaged alongside parasites carrying mutations and were used to normalize the signal of mutation-harboring parasites. Parasites were selected based on DIC and then exposed for 200 ms to image green fluorescence. Total intensity of the GFP signal in foci was measured and background signal subtracted using ImageJ (ImageJ2 2018, (*61*)).

Growth assessment

The parasitemia of a mixed-stage parasite culture was measured by flow cytometry (*40*) and based on this the parasitemia was adjusted to 0.05 to 0.1% parasitemia in 2.5% hematocrit. The parasitemia was then measured again by flow cytometry to determine the start parasitemia. These parasites were cultivated for 96 h and the medium was changed every 24 h. After 96 h parasitemia was measured again by flow cytometry and divided by the starting parasitemia to obtain the fold change in parasitemia.

Competition assay

Schizonts were isolated using 60% percoll purification, washed once with medium and cultured for 6 hours to allow invasion of merozoites into new red blood cells. Remaining schizonts were removed by a 10-min incubation in 5% sorbitol solution. The resulting 0-6 h old parasites were cultured at 37°C for 20-24 h after which the parasitemia was measured by flow cytometry (*40*). Based on the determined parasitemia, the mutant K13 cell lines were co-cultivated in a 1:1 ratio with 3D7 control in a 5 mL Petri dish. The proportion of GFP-positive parasites was assessed by fluorescence microscopy until one parasite line reached 95% of parasite proportion.

Statistical analysis and malaria incidence data

Unpaired or paired two-tailed t-tests were performed as indicated and Pearson's r calculated using GraphPad Prism 9.0.2. Linear regressions were fit to log transformed data (GraphPad Prism). All error bars shown are standard deviations. Malaria incidence data was downloaded from https://ghdx.healthdata.org/gbd-results-tool.

References

1. Institute for Health Metrics and Evaluation (IHME), Global Burden of Disease Collaborative Network. Global Burden of Disease Study 2019 (GBD 2019) Results. (2021).
2. World Health Organization, *World malaria report 2021* (2021; https://www.who.int/teams/global-malaria-programme/reports/world-malaria-report-2021).
3. World Health Organization, *Report on antimalarial drug efficacy, resistance and response: 10 years of surveillance (2010-2019)* (2020; https://www.who.int/publications/i/item/9789240012813).
4. H. Noedl, Y. Se, K. Schaecher, B. L. Smith, D. Socheat, M. M. Fukuda, Evidence of artemisinin-resistant malaria in Western Cambodia. *N. Engl. J. Med.* **359**, 2619–2620 (2008).
5. P. J. de Vries, T. K. Dien, Clinical Pharmacology and Therapeutic Potential of Artemisinin and its Derivatives in the Treatment of Malaria. *Drugs*. **52**, 818–836 (1996).
6. D. Menard, A. Dondorp, Antimalarial Drug Resistance: A Threat to Malaria Elimination. *Cold Spring Harb. Perspect. Med.* **7**, a025619 (2017).
7. B. Witkowski, C. Amaratunga, N. Khim, S. Sreng, P. Chim, S. Kim, P. Lim, S. Mao, C. Sopha, B. Sam, J. M. Anderson, D. Socheat, C. M. Chuor, W. R. J. Taylor, S. Suon, O. Mercereau-Puijalon, R. M. Fairhurst, D. Menard, Novel phenotypic assays for the detection of artemisinin-resistant Plasmodium falciparum malaria in Cambodia: in-vitro and ex-vivo drug-response studies. *Lancet Infect Dis.* **13**, 1043–1049 (2013).
8. F. Ariey, B. Witkowski, C. Amaratunga, J. Beghain, A.-C. Langlois, N. Khim, S. Kim, V. Duru, C. Bouchier, L. Ma, P. Lim, R. Leang, S. Duong, S. Sreng, S. Suon, C. M. Chuor, D. M. Bout, S. Ménard, W. O. Rogers, B. Genton, T. Fandeur, O. Miotto, P. Ringwald, J. Le Bras, A. Berry, J.-C. Barale, R. M. Fairhurst, F. Benoit-Vical, O. Mercereau-Puijalon, D. Ménard, A molecular

marker of artemisinin-resistant Plasmodium falciparum malaria. *Nature*. **505**, 50–55 (2014).

9.   J. Straimer, N. F. Gnädig, B. Witkowski, C. Amaratunga, V. Duru, A. P. Ramadani, M. Dacheux,
N. Khim, L. Zhang, R. M. Fairhurst, D. Ménard, D. A. Fidock, K13-propeller mutations confer
artemisinin resistance in Plasmodium falciparum clinical isolates. *Science.* **347**, 428–431
(2015).

10.   J. Birnbaum, S. Scharf, S. Schmidt, E. Jonscher, W. A. M. Hoeijmakers, S. Flemming, C. G.
Toenhake, M. Schmitt, R. Sabitzki, B. Bergmann, U. Fröhlke, P. Mesén-Ramírez, A. B. Soares,
H. Herrmann, R. Bártfai, T. Spielmann, A Kelch13-defined endocytosis pathway mediates
artemisinin resistance in malaria parasites. *Science.* **367**, 51–59 (2020).

11.   N. V. Simwela, K. R. Hughes, A. B. Roberts, M. T. Rennie, M. P. Barrett, A. P. Waters,
Experimentally Engineered Mutations in a Ubiquitin Hydrolase, UBP-1, Modulate In Vivo
Susceptibility to Artemisinin and Chloroquine in Plasmodium berghei. *Antimicrob. Agents*
Chemother. **64**, e02484-19 (2020).

12.   R. C. Henrici, D. A. van Schalkwyk, C. J. Sutherland, Modification of pfap2μ and pfubp1
Markedly Reduces Ring- Stage Susceptibility of Plasmodium falciparum to Artemisinin In
Vitro. *Antimicrob. Agents Chemother.* **64**, e01542-19 (2019).

13.   A. R. Demas, A. I. Sharma, W. Wong, A. M. Early, S. Redmond, S. Bopp, D. E. Neafsey, S. K.
Volkman, D. L. Hartl, D. F. Wirth, Mutations in plasmodium falciparum actin-binding protein
coronin confer reduced artemisinin susceptibility. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 12799–
12804 (2018).

14.   A. I. Sharma, S. H. Shin, S. Bopp, S. K. Volkman, D. L. Hartl, D. F. Wirth, Genetic background
and PfKelch13 affect artemisinin susceptibility of PfCoronin mutants in Plasmodium
falciparum. *PLOS Genet.* **16**, e1009266 (2020).

15.   G. C. Cerqueira, I. H. Cheeseman, S. F. Schaffner, S. Nair, M. McDew-White, A. P. Phyo, E. A.
Ashley, A. Melnikov, P. Rogov, B. W. Birren, F. Nosten, T. J. C. Anderson, D. E. Neafsey,
Longitudinal genomic surveillance of Plasmodium falciparum malaria parasites reveals
complex genomic architecture of emerging artemisinin resistance. *Genome Biol.* **18**, 78
(2017).

16.   T. Yang, L. M. Yeoh, M. V. Tutor, M. W. Dixon, P. J. McMillan, S. C. Xie, J. L. Bridgford, D. L.
Gillett, M. F. Duffy, S. A. Ralph, M. J. McConville, L. Tilley, S. A. Cobbold, Decreased K13
Abundance Reduces Hemoglobin Catabolism and Proteotoxic Stress, Underpinning
Artemisinin Resistance. *Cell Rep.* **29**, 2917-2928.e5 (2019).

17.   N. F. Gnädig, B. H. Stokes, R. L. Edwards, G. F. Kalantarov, K. C. Heimsch, M. Kuderjavy, A.
Crane, M. C. S. Lee, J. Straimer, K. Becker, I. N. Trakht, A. R. Odom John, S. Mok, D. A. Fidock,
Insights into the intracellular localization, protein associations and artemisinin resistance
properties of Plasmodium falciparum K13. *PLOS Pathog.* **16**, e1008482 (2020).

18.   G. Siddiqui, A. Srivastava, A. S. Russell, D. J. Creek, Multi-omics based identification of specific
biochemical changes associated with PfKelch13-mutant artemisinin-resistant plasmodium
falciparum. *J. Infect. Dis.* **215**, 1435–1444 (2017).

19.   R. Schumann, E. Bischoff, S. Klaus, S. Möhring, J. Flock, S. Keller, K. Remans, M. Ganter, M.
Deponte, Protein abundance and folding rather than the redox state of Kelch13 determine
the artemisinin susceptibility of Plasmodium falciparum. *Redox Biol.* **48**, 102177 (2021).

20.   A. Xiong, P. Prakash, X. Gao, M. Chew, I. J. J. Tay, C. J. Woodrow, B. P. Engelward, J. Han, P. R.
Preiser, K13-Mediated Reduced Susceptibility to Artemisinin in Plasmodium falciparum Is
Overlaid on a Trait of Enhanced DNA Damage Repair. *Cell Rep.* **32**, 107996 (2020).

21.   D. Ménard, N. Khim, J. Beghain, A. A. Adegnika, M. Shafiul-Alam, O. Amodu, G. Rahim-Awab,
C. Barnadas, A. Berry, Y. Boum, M. D. Bustos, J. Cao, J. H. Chen, L. Collet, L. Cui, G. Das
Thakur, A. Dieye, D. Djallé, M. A. Dorkenoo, C. E. Eboumbou-Moukoko, F. E. C. J. Espino, T.
Fandeur, M. F. Ferreira-Da-Cruz, A. A. Fola, H. P. Fuehrer, A. M. Hassan, S. Herrera, B.
Hongvanthong, S. Houzé, M. L. Ibrahim, M. Jahirul-Karim, L. Jiang, S. Kano, W. Ali-Khan, M.
Khanthavong, P. G. Kremsner, M. Lacerda, R. Leang, M. Leelawong, M. Li, K. Lin, J. B.
Mazarati, S. Ménard, I. Morlais, H. Muhindo-Mavoko, L. Musset, K. Na-Bangchang, M.

Nambozi, K. Niaré, H. Noedl, J. B. Ouédraogo, D. R. Pillai, B. Pradines, B. Quang-Phuc, M.
Ramharter, M. Randrianarivelojosia, J. Sattabongkot, A. Sheikh-Omar, K. D. Silué, S. B. Sirima,
C. Sutherland, D. Syafruddin, R. Tahar, L. H. Tang, O. A. Touré, P. Tshibangu-Wa-Tshibangu, I.
Vigan-Womas, M. Warsame, L. Wini, S. Zakeri, S. Kim, R. Eam, L. Berne, C. Khean, S. Chy, M.
Ken, K. Loch, L. Canier, V. Duru, E. Legrand, J. C. Barale, B. Stokes, J. Straimer, B. Witkowski,
D. A. Fidock, C. Rogier, P. Ringwald, F. Ariey, O. Mercereau-Puijalon, A worldwide map of
Plasmodium falciparum K13-propeller polymorphisms. *N. Engl. J. Med.* **374**, 2453–2464
(2016).
22.   B. H. Stokes, S. K. Dhingra, K. Rubiano, S. Mok, J. Straimer, N. F. Gnädig, I. Deni, K. A.
Schindler, J. R. Bath, K. E. Ward, J. Striepen, T. Yeo, L. S. Ross, E. Legrand, F. Ariey, C. H.
Cunningham, I. M. Souleymane, A. Gansané, R. Nzoumbou-Boko, C. Ndayikunda, A. M.
Kabanywanyi, A. Uwimana, S. J. Smith, O. Kolley, M. Ndounga, M. Warsame, R. Leang, F.
Nosten, T. J. Anderson, P. J. Rosenthal, D. Ménard, D. A. Fidock, Plasmodium falciparum K13
mutations in Africa and Asia impact artemisinin resistance and parasite fitness. *Elife*. **10**,
e66277 (2021).
23.   R. Amato, O. Miotto, C. J. Woodrow, J. Almagro-Garcia, I. Sinha, S. Campino, D. Mead, E.
Drury, M. Kekre, M. Sanders, A. Amambua-Ngwa, C. Amaratunga, L. Amenga-Etego, V.
Andrianaranjaka, T. Apinjoh, E. Ashley, S. Auburn, G. A. Awandare, V. Baraka, A. Barry, M. F.
Boni, S. Borrmann, T. Bousema, O. Branch, P. C. Bull, K. Chotivanich, D. J. Conway, A. Craig, N.
P. Day, A. Djimdé, C. Dolecek, A. M. Dondorp, C. Drakeley, P. Duffy, D. F. Echeverry, T. G.
Egwang, R. M. Fairhurst, A. Faiz, C. I. Fanello, T. T. Hien, A. Hodgson, M. Imwong, D.
Ishengoma, P. Lim, C. Lon, J. Marfurt, K. Marsh, M. Mayxay, P. Michon, V. Mobegi, O. A.
Mokuolu, J. Montgomery, I. Mueller, M. P. Kyaw, P. N. Newton, F. Nosten, R. Noviyanti, A.
Nzila, H. Ocholla, A. Oduro, M. Onyamboko, J. B. Ouedraogo, A. P. P. Phyo, C. Plowe, R. N.
Price, S. Pukrittayakamee, M. Randrianarivelojosia, P. Ringwald, L. Ruiz, D. Saunders, A.
Shayo, P. Siba, S. Takala-Harrison, T. N. N. Thanh, V. Thathy, F. Verra, J. Wendler, N. J. White,
H. Ye, V. J. Cornelius, R. Giacomantonio, D. Muddyman, C. Henrichs, C. Malangone, D. Jyothi,
R. D. Pearson, J. C. Rayner, G. McVean, K. A. Rockett, A. Miles, P. Vauterin, B. Jeffery, M.
Manske, J. Stalker, B. Macinnis, D. P. Kwiatkowski, Genomic epidemiology of artemisinin
resistant malaria. *Elife*. **5**, e08714 (2016).
24.   O. Miotto, R. Amato, E. A. Ashley, B. MacInnis, J. Almagro-Garcia, C. Amaratunga, P. Lim, D.
Mead, S. O. Oyola, M. Dhorda, M. Imwong, C. Woodrow, M. Manske, J. Stalker, E. Drury, S.
Campino, L. Amenga-Etego, T.-N. N. Thanh, H. T. Tran, P. Ringwald, D. Bethell, F. Nosten, A.
P. Phyo, S. Pukrittayakamee, K. Chotivanich, C. M. Chuor, C. Nguon, S. Suon, S. Sreng, P. N.
Newton, M. Mayxay, M. Khanthavong, B. Hongvanthong, Y. Htut, K. T. Han, M. P. Kyaw, M. A.
Faiz, C. I. Fanello, M. Onyamboko, O. A. Mokuolu, C. G. Jacob, S. Takala-Harrison, C. V Plowe,
N. P. Day, A. M. Dondorp, C. C. A. Spencer, G. McVean, R. M. Fairhurst, N. J. White, D. P.
Kwiatkowski, Genetic architecture of artemisinin-resistant Plasmodium falciparum. *Nat.*
Genet. **47**, 226–234 (2015).
25.   N. K. Kayiba, D. M. Yobi, E. Tshibangu-Kabamba, V. P. Tuan, Y. Yamaoka, B. Devleesschauwer,
D. M. Mvumbi, E. Okitolonda Wemakoy, P. De Mol, G. L. Mvumbi, M. Hayette, A. Rosas-
Aguirre, N. Speybroeck, Spatial and molecular mapping of Pfkelch13 gene polymorphism in
Africa in the era of emerging Plasmodium falciparum resistance to artemisinin: a systematic
review. *Lancet Infect. Dis.* **21**, e82–e92 (2021).
26.   L. Ndwiga, K. M. Kimenyi, K. Wamae, V. Osoti, M. Akinyi, I. Omedo, D. S. Ishengoma, N. Duah-
Quashie, B. Andagalu, A. Ghansah, A. Amambua-Ngwa, S. Tukwasibwe, S. K. Tessema, C.
Karema, A. A. Djimde, A. M. Dondorp, J. Raman, R. W. Snow, P. Bejon, L. I. Ochola-Oyier, A
review of the frequencies of Plasmodium falciparum Kelch 13 artemisinin resistance
mutations in Africa. *Int. J. Parasitol. Drugs Drug Resist.* **16**, 155–161 (2021).
27.   B. Balikagala, N. Fukuda, M. Ikeda, O. T. Katuro, S.-I. Tachibana, M. Yamauchi, W. Opio, S.
Emoto, D. A. Anywar, E. Kimura, N. M. Q. Palacpac, E. I. Odongo-Aginya, M. Ogwang, T. Horii,
T. Mita, Evidence of Artemisinin-Resistant Malaria in Africa. *N. Engl. J. Med.* **385**, 1163–1171

(2021).

28.   World Health Organization, "Artemisinin resistance and artemisinin-based combination
620   therapy efficacy" (2019), (available at
621   https://www.who.int/malaria/publications/atoz/artemisinin-resistance-august2018/en/).

622   29.   A. Uwimana, E. Legrand, B. H. Stokes, J. L. M. Ndikumana, M. Warsame, N. Umulisa, D.
623   Ngamije, T. Munyaneza, J. B. Mazarati, K. Munguti, P. Campagne, A. Criscuolo, F. Ariey, M.
624   Murindahabi, P. Ringwald, D. A. Fidock, A. Mbituyumuremyi, D. Menard, Emergence and
625   clonal expansion of in vitro artemisinin-resistant Plasmodium falciparum kelch13 R561H
626   mutant parasites in Rwanda. *Nat. Med.* **26**, 1602–1608 (2020).

627   30.   A. Uwimana, N. Umulisa, M. Venkatesan, S. S. Svigel, Z. Zhou, T. Munyaneza, R. M.
628   Habimana, A. Rucogoza, L. F. Moriarty, R. Sandford, E. Piercefield, I. Goldman, B. Ezema, E.
629   Talundzic, M. A. Pacheco, A. A. Escalante, D. Ngamije, J.-L. N. Mangala, M. Kabera, K.
630   Munguti, M. Murindahabi, W. Brieger, C. Musanabaganwa, L. Mutesa, V. Udhayakumar, A.
631   Mbituyumuremyi, E. S. Halsey, N. W. Lucchi, Association of Plasmodium falciparum kelch13
632   R561H genotypes with delayed parasite clearance in Rwanda: an open-label, single-arm,
633   multicentre, therapeutic efficacy study. *Lancet Infect. Dis.* **21**, 1120–1128 (2021).

634   31.   K. G.-P. S. WWARN, P. Dahal, U. D'Alessandro, G. Dorsey, P. J. Guerin, C. Nsanzabana, R. N.
635   Price, C. H. Sibley, K. Stepniewska, A. O. Talisuna, Clinical determinants of early
636   parasitological response to ACTs in African patients with uncomplicated falciparum malaria: a
637   literature review and meta-analysis of individual patient data. *BMC Med.* **13**, 212 (2015).

638   32.   N. Scott, R. Ataide, D. P. Wilson, M. Hellard, R. N. Price, J. A. Simpson, F. J. I. Fowkes,
639   Implications of population-level immunity for the emergence of artemisinin-resistant
640   malaria: a mathematical model. *Malar. J.* **17**, 279 (2018).

641   33.   E. Y. Klein, D. L. Smith, R. Laxminarayan, S. Levin, Superinfection and the evolution of
642   resistance to antimalarial drugs. *Proc. R. Soc. B Biol. Sci.* **279**, 3834–3842 (2012).

643   34.   P. Mesén-Ramírez, B. Bergmann, M. Elhabiri, L. Zhu, H. von Thien, C. Castro-Peña, T.-W.
644   Gilberger, E. Davioud-Charvet, Z. Bozdech, A. Bachmann, T. Spielmann, The parasitophorous
645   vacuole nutrient channel is critical for drug access in malaria parasites and modulates the
646   artemisinin resistance fitness cost. *Cell Host Microbe*. **29**, 1774-1787.e9 (2021).

647   35.   S. Nair, X. Li, G. A. Arya, M. McDew-White, M. Ferrari, T. Anderson, Nutrient Limitation
648   Magnifies Fitness Costs of Antimalarial Drug Resistance Mutations. *Antimicrob. Agents*
649   *Chemother.* **66**, e00605-18 (2022).

650   36.   WWARN K13 Genotype-Phenotype Study Group, Association of mutations in the
651   Plasmodium falciparum Kelch13 gene (Pf3D7_1343700) with parasite clearance rates after
652   artemisinin-based treatments—a WWARN individual patient data meta-analysis. *BMC Med.*
653   **17**, 1 (2019).

654   37.   B. Hogan, D. Eibach, R. Krumkamp, N. Sarpong, D. Dekker, B. Kreuels, O. Maiga-Ascofaré, K.
655   G. Boahen, C. W. Akenten, Y. Adu-Sarkodie, E. Owusu-Dabo, J. May, Malaria Coinfections in
656   Febrile Pediatric Inpatients: A Hospital-Based Study from Ghana. *Clin. Infect. Dis.* **66**, 1838–
657   1845 (2018).

658   38.   MalariaGEN, MalariaGen (2021), (available at https://www.malariagen.net/).

659   39.   F. Rocamora, L. Zhu, K. Y. Liong, A. Dondorp, O. Miotto, S. Mok, Z. Bozdech, Oxidative stress
660   and protein damage responses mediate artemisinin resistance in malaria parasites. *PLoS*
661   *Pathog.* **14** (2018), doi:10.1371/journal.ppat.1006930.

662   40.   J. Birnbaum, S. Flemming, N. Reichard, A. B. Soares, P. Mesén-ramírez, E. Jonscher, B.
663   Bergmann, T. Spielmann, A genetic system to study Plasmodium falciparum protein function.
664   *Nat. Methods*. **14**, 450–456 (2017).

665   41.   M. Bushman, R. Antia, V. Udhayakumar, J. C. de Roode, Within-host competition can delay
666   evolution of drug resistance in malaria. *PLOS Biol.* **16**, e2005712 (2018).

667   42.   M. Legros, S. Bonhoeffer, A combined within-host and between-hosts modelling framework
668   for the evolution of resistance to antimalarial drugs. *J. R. Soc. Interface*. **13**, 20160148 (2016).

669   43.   S. Nair, X. Li, G. A. Arya, M. McDew-White, M. Ferrari, T. Anderson, Nutrient Limitation

670		Magnifies Fitness Costs of Antimalarial Drug Resistance Mutations. *Antimicrob. Agents*
671		*Chemother.* (2022), doi:10.1128/aac.01529-21.
672	44.	A. R. Tirrell, K. M. Vendrely, L. A. Checkley, S. Z. Davis, M. McDew-White, I. H. Cheeseman, A.
673		M. Vaughan, F. H. Nosten, T. J. C. Anderson, M. T. Ferdig, Pairwise growth competitions
674		identify relative fitness relationships among artemisinin resistant Plasmodium falciparum
675		field isolates. *Malar. J.* **18**, 295 (2019).
676	45.	J. Straimer, N. F. Gnädig, B. H. Stokes, M. Ehrenberger, A. A. Crane, D. A. Fidock, Plasmodium
677		falciparum K13 Mutations Differentially Impact Ozonide Susceptibility and Parasite Fitness In
678		Vitro. *MBio.* **8**, e00172-17 (2017).
679	46.	S. Mok, B. H. Stokes, N. F. Gnädig, L. S. Ross, T. Yeo, C. Amaratunga, E. Allman, L. Solyakov, A.
680		R. Bottrill, J. Tripathi, R. M. Fairhurst, M. Llinás, Z. Bozdech, A. B. Tobin, D. A. Fidock,
681		Artemisinin-resistant K13 mutations rewire Plasmodium falciparum's intra-erythrocytic
682		metabolic program to enhance survival. *Nat. Commun.* **12**, 530 (2021).
683	47.	S. Mok, E. A. Ashley, P. E. Ferreira, L. Zhu, Z. Lin, T. Yeo, K. Chotivanich, M. Imwong, S.
684		Pukrittayakamee, M. Dhorda, C. Nguon, P. Lim, C. Amaratunga, S. Suon, T. T. Hien, Y. Htut,
685		M. A. Faiz, M. A. Onyamboko, M. Mayxay, P. N. Newton, R. Tripura, C. J. Woodrow, O.
686		Miotto, D. P. Kwiatkowski, F. Nosten, N. P. J. Day, P. R. Preiser, N. J. White, A. M. Dondorp, R.
687		M. Fairhurst, Z. Bozdech, Population transcriptomics of human malaria parasites reveals the
688		mechanism of artemisinin resistance. *Science.* **347**, 431–435 (2015).
689	48.	C. Dogovski, S. C. Xie, G. Burgio, J. Bridgford, S. Mok, J. M. McCaw, K. Chotivanich, S. Kenny,
690		N. Gnädig, J. Straimer, Z. Bozdech, D. A. Fidock, J. A. Simpson, A. M. Dondorp, S. Foote, N.
691		Klonis, L. Tilley, Targeting the Cell Stress Response of Plasmodium falciparum to Overcome
692		Artemisinin Resistance. *PLOS Biol.* **13**, e1002132 (2015).
693	49.	S. Bhattacharjee, I. Coppens, A. Mbengue, N. Suresh, M. Ghorbal, Z. Slouka, I. Safeukui, H. Y.
694		Tang, D. W. Speicher, R. V. Stahelin, N. Mohandas, K. Haldar, Remodeling of the malaria
695		parasite and host human red cell by vesicle amplification that induces artemisinin resistance.
696		*Blood.* **131**, 1234–1247 (2018).
697	50.	M. Zhang, J. Gallego-Delgado, C. Fernandez-Arias, N. C. Waters, A. Rodriguez, M. Tsuji, R. C.
698		Wek, V. Nussenzweig, W. J. Sullivan, Inhibiting the Plasmodium eIF2α Kinase PK4 Prevents
699		Artemisinin-Induced Latency. *Cell Host Microbe.* **22**, 766–776 (2017).
700	51.	J. Gibbons, K. A. Button-Simons, S. R. Adapa, S. Li, M. Pietsch, M. Zhang, X. Liao, J. H. Adams,
701		M. T. Ferdig, R. H. Y. Jiang, Altered expression of K13 disrupts DNA replication and repair in
702		Plasmodium falciparum. *BMC Genomics.* **19**, 849 (2018).
703	52.	S. Schmidt, J. S. Wichers-misterek, H. M. Behrens, J. Birnbaum, I. Henshall, E. Jonscher, S.
704		Flemming, T. Spielmann, The Kelch13 compartment is a hub of highly divergent vesicle
705		trafficking proteins in malaria parasites. *bioRxiv [Preprint]* (2022),
706		doi:https://doi.org/10.1101/2022.12.15.520209.
707	53.	S. Borrmann, J. Straimer, L. Mwai, A. Abdi, A. Rippert, J. Okombo, S. Muriithi, P. Sasi, M. M.
708		Kortok, B. Lowe, S. Campino, S. Assefa, S. Auburn, M. Manske, G. Maslen, N. Peshu, D. P.
709		Kwiatkowski, K. Marsh, A. Nzila, T. G. Clark, Genome-wide screen identifies new candidate
710		genes associated with artemisinin susceptibility in Plasmodium falciparum in Kenya. *Sci. Rep.*
711		**3**, 3318 (2013).
712	54.	G. Henriques, R. L. Hallett, K. B. Beshir, N. B. Gadalla, R. E. Johnson, R. Burrow, D. A. Van
713		Schalkwyk, P. Sawa, S. A. Omar, T. G. Clark, T. Bousema, C. J. Sutherland, Directional selection
714		at the pfmdr1, pfcrt, pfubp1, and pfap2mu loci of Plasmodium falciparum in Kenyan children
715		treated with ACT. *J. Infect. Dis.* **210**, 2001–2008 (2014).
716	55.	A. B. Soares, J. Stäcker, S. Schwald, W. Hoijmakers, N. G. Metwally, H. Schoeler, S. Flemming,
717		K. Höhn, U. Fröhlke, P. Mesén-Ramírez, B. Bergmann, M. Khosh-Naucke, I. Bruchhaus, R.
718		Bártfai, T. Spielmann, *bioRxiv [Preprint]*, in press (available at
719		https://www.biorxiv.org/content/10.1101/2021.12.03.471078v1?ct=).
720	56.	D. Walliker, I. Quakyi, T. Wellems, T. McCutchan, A. Szarfman, W. London, L. Corcoran, T.
721		Burkot, R. Carter, Genetic analysis of the human malaria parasite Plasmodium falciparum.

Science. **236**, 1661–1666 (1987).
57.   W. Trager, J. Jensen, Human malaria parasites in continuous culture. *Science.* **193**, 673–675
(1976).
58.   R. W. Moon, J. Hall, F. Rangkuti, Y. S. Ho, N. Almond, G. H. Mitchell, A. Pain, A. A. Holder, M.
J. Blackman, Adaptation of the genetically tractable malaria pathogen Plasmodium knowlesi
to continuous culture in human erythrocytes. *Proc. Natl. Acad. Sci.* **110**, 531–536 (2013).
59.   B. Witkowski, C. Amaratunga, N. Khim, S. Sreng, P. Chim, S. Kim, P. Lim, S. Mao, C. Sopha, B.
Sam, J. M. Anderson, D. Socheat, C. M. Chuor, W. R. J. Taylor, S. Suon, O. Mercereau-
Puijalon, R. M. Fairhurst, D. Menard, Novel phenotypic assays for the detection of
artemisinin- resistant Plasmodium falciparum malaria in Cambodia: in-vitro and ex-vivo drug-
response studies. *Lancet Infect Dis.* **13**, 1043–1049 (2013).
60.   C. Grüring, A. Heiber, F. Kruse, J. Ungefehr, T.-W. Gilberger, T. Spielmann, Development and
host cell modifications of Plasmodium falciparum blood stages in four dimensions. *Nat.*
Commun. **2**, 165 (2011).
61.   C. A. Schneider, W. S. Rasband, K. W. Eliceiri, NIH Image to ImageJ: 25 years of image
analysis. *Nat. Methods*. **9**, 671–675 (2012).

738
739

Acknowledgments

We thank Jacobus Pharmaceuticals for supplying WR99210. DSM1 (MRA-1161) was received from
MR4/BEI Resources, NIAID, NIH. We thank Birgit Förster for sequencing and Ralf Krumpkamp for
helpful discussions. This research made use of PlasmoDB.org. HMB acknowledges funding by the
Ortrud Mührer Fellowship of the Vereinigung der Freunde des Tropeninstituts Hamburg e.V.. This
study was in part funded by the German Center for Infection Research (TTU03.806). OMA was
supported by the DELGEME fellowship funded by DELTAS-Africa 107740/Z/15/Z from Wellcome
Trust. TS, IH and RS acknowledge funding by the European Research Council (ERC, grant 101021493).

Author contributions: TS, OMA and JM conceived the project. TS supervised research. SSchm, HMB,
RS, IH and DP performed experiments. SSchm and HMB analyzed data and prepared figures. HMB and
TS wrote the manuscript draft. All authors approved the manuscript.

Competing interests: The authors declare no competing interests.

Data and materials availability: All data are available in the main text or the supplementary materials.

Figures and Tables

761
Figure 1. 135 tested mutations in K13-compartment proteins do not cause ART resistance. (A)
Mutations introduced in the multi-mutant of KIC1 are exemplarily shown in a schematic and
fluorescence images of the resulting parasites harboring multi-mutant-KIC1 are shown.
Scale bar 5 μm. RSA survival for all multi-mutants is displayed (% survival compared to
control without DHA 66 h after 6 h DHA treatment in RSA). MyoF multi-mutant was
compared with 3D7 as matching WT was not available. **(B)** RSA survival for single mutations
in K13 compartment proteins. Eps15 K447R was compared with 3D7 as matching WT was
not available. **(C)** Position of *k13* mutations as well as BTB domain and propeller domain are
shown on K13. **(D)** RSA of different *k13* mutant cell lines. Error bars show standard
deviations. WT, wild type; red bars indicate the mean parasite survival rate of the respective
cell line, each dot represents an independent experiment, green dashed line represents the
1% cut-off value defining clinically-relevant ART resistance. Position of *k13* mutations is
shown on K13. Error bars show standard deviations. P values derive from two-tailed
unpaired t-tests.

778
Figure 2. DHA-selection but not combination of two mutations increases resistance. (A) RSA
survival of the different *k13* mutant cell lines shown. WT, wild type; each point represents
an independent RSA; red bars indicate the mean parasite survival rate of the respective cell
line. Position of *k13* mutations are shown on K13. Domains are colored as in Figure 1. Red
bars show mean; error bars show standard deviations. P values derive from two-tailed
unpaired t-tests. **(B)** Scheme of experimental procedure of consecutive RSA cycles
performed with DHA survivors of the respective prior cycle. **(C)** Parasite survival rate of
K13^{C580Y} (% survival compared to control without DHA) 66 h after 6 h DHA treatment in RSA.
Six experiments per cycle were performed, consisting of six replicates (see color code).
Replicates 1, 2, 3 (orange) and 4, 5, 6 (green) were each started on the same days. Black
bars show mean. P value is indicated, two-tailed paired t-test.
790

791
792

793
Figure 3. Resistance is inversely correlated with K13- and KIC7-abundance. (A) K13-abundance
measured by the GFP fluorescence intensity of the single focus observed in ring stage
parasites with the indicated K13 expressed from the endogenous locus and fused to GFP,
normalized to the fluorescence in parasites with the identically modified endogenous locus
but with a WT GFP-K13. Each small dot represents the measured value from one focus in
one parasite. Large dots of the same color as small dots represent mean of the respective
measured values and each large dot represents one biological replicate. Each biological
replicate (large dot) consists of 20 individual measurements (small dots). P-values derive
from comparing means (large dots) by one-way ANOVA. Example fluorescence microscopy
images are shown. Scale bar 5 μm. **(B)** Mean of parasite survival in RSA plotted against
mean of K13-abundance. Error bars show standard deviations. **(C+D)** RSA survival of (C) K13
1xNLS parasites or (D) KIC7 1xNLS parasites grown in the presence of 12.5 nM, 25 nM, 125
nM or 250 nM rapalog for 3 h before and 6 h during the DHA exposure of the RSA. Red bar,
mean; standard deviation indicated; each dot derives from an independent experiment.
808
809
810
811

Figure 4. Fitness cost correlates with resistance: (A) Parasite growth, calculated as fold-change of parasitemia after 96 h compared to parasitemia at 0 h, normalized to WT growth for different cell lines with indicated k13 mutations. Error bars show standard deviation. P values derive from two-tailed unpaired t-tests. **(B)** Mean of growth of strains harboring different *k13* mutations plotted against mean of parasite survival in RSA. Error bars show standard deviation. **(C)** Fitness cost of parasites harboring different *k13* mutations grown in competition with 3D7 parasites given as loss of parasite proportion/generation. Error bars show standard deviation. P values derive from two-tailed unpaired t-tests. **(D)** Mean of fitness cost plotted against mean of parasite survival in RSA. All error bars show standard deviations.

824
Figure 5. Double mutation of *k13* C580Y and *ubp1* R3138H result in a high fitness cost. (A) RSA of
K13^{C580Y} and UBP1^{R3138H} single and double mutant cell lines. Red bars indicate the mean
parasite survival rate of the respective cell line. Data for UBP1^{R3138H} was previously
published in (*10*). Position of *k13* and *ubp1* mutation is depicted in K13 and UBP1. K13
domains are colored as in Figure 1. Error bars show standard deviation. P values derive from
two-tailed unpaired t-tests. **(B)** K13-abundance measured GFP fluorescence intensity of the
single focus observed in ring stage parasites of the indicated K13 expressed from the

endogenous locus and fused to GFP, normalized to the fluorescence in parasites with the
identically modified endogenous locus but with a WT GFP-K13. Small dots represent
measured values for individual parasites. Large dots of the same color as small dots
represent mean of the respective measured values and each large dot represents one
biological replicate. Dots of the same color were obtained as part of the same experimental
repeat. P values derive from two-tailed unpaired t-tests. **(C)** K13 abundance plotted against
parasite survival in RSA for K13C580Y+UBP1R3138H parasites (yellow). Grey dots show data
of k13 mutations from Figure 3B for comparison. Error bars show standard deviation. **(D)**
Growth, calculated as fold-change of parasitemia after 96 h normalized compared to
parasitemia at 0 h for different cell lines with indicated *k13* or *ubp1* mutations. Error bars
show standard deviation. P values derive from two-tailed unpaired t-tests. **(E)** Growth of
strains harboring *ubp1* R3138H with (yellow) and without (orange) *k13* C580Y plotted
against parasite survival in RSA. Grey dots show data of *k13* mutations from Figure 4B for
comparison. Error bars show standard deviation. **(F)** Fitness cost for different cell lines with
indicated *k13* or *ubp1* mutations grown in mixed cultures with 3D7. Error bars show
standard deviation. P values derive from two-tailed unpaired t-tests. **(G)** Fitness cost
K13^{C580Y}+UBP1^{R3138H} parasites (yellow) plotted against parasite survival in RSA. Grey dots
show data of *k13* mutations from Figure 4D for comparison. All error bars show standard
deviations.