Comparative Transcriptome Analysis of Hypocotyls During the Developmental Transition of C₃ cotyledons to C₄ Leaves in *Halimocnemis mollissima* Bunge

Mahdis Zolfaghar[1], Twan Rutten[2], Mohammad Reza Ghaffari[*3], Ali Mohammad Banaei-Moghaddam[*1]

[1] Laboratory of Genomics and Epigenomics (LGE), Institute of Biochemistry and Biophysics, University of Tehran, Tehran, Iran

[2] Department of Physiology and Cell Biology, Leibniz Institute of Plant Genetics and Crop Plant Research (IPK), Stadt Seeland/OT Gatersleben, Germany

[3] Systems Biology Department, Agricultural Biotechnology Research Institute of Iran (ABRII), Agricultural Research, Education, and Extension Organization (AREEO), Karaj, Iran

[*] **Correspondence:**

Mohammad Reza Ghaffari

ghaffari@abrii.ac.ir (M.R.G)

Ali Mohammad Banaei-Moghaddam

 am_banaei@ut.ac.ir (A.M.B-M)

ORCID ID:

Mahdis Zolfaghar 0000-0001-8762-3603

Twan Rutten 0000-0001-5891-6503

Mohammad Reza Ghaffari 0000-0002-7139-8613

Ali Mohammad Banaei-Moghaddam 0000-0003-1298-3748

Abstract

Identification of signaling pathways that control C₄ photosynthesis development is essential for introducing the C₄ pathway into C₃ crops. Species with dual photosynthesis in their life cycle are interesting models to study such regulatory mechanisms. The species used here *Halimocnemis mollissima* Bunge, belonging to the Caroxyleae tribe, displays C₃ photosynthesis in its cotyledons and a NAD-ME subtype of C₄ photosynthesis in the First leaves

(FLs) onwards. We explored the long-distance signaling pathways that are probably implicated in the shoot-root coordination associated with the manifestation of the C_4 traits, including efficient resource usage by comparing the mRNA content of hypocotyls before and after the C_4 first leave's formation. Histological examination showed the presence of C_3 anatomy in cotyledons and C_4 anatomy in the FLs. Our transcriptome analyses verified the performance of the NAD-ME subtype of C_4 in FLs and revealed differential transcript abundance of several potential mobile regulators and their associated receptors or transporters in two developmentally different hypocotyls of *H. mollissima* Bunge. These differentially expressed genes (DEGs) belong to diverse functional groups, including various transcription factor (TF) families, phytohormones metabolism, and signaling peptides, part of which could be related to hypocotyl development. Our findings support the higher nitrogen and water use efficiency associated with C_4 photosynthetic and provide insights into the coordinated above- and under-ground tissue communication during the developmental transition of C_3 to C_4 photosynthesis in this species.

Keywords: C_4 photosynthesis, *Halimocnemis mollissima* Bunge, long-distance signaling, NAD-ME subtype, RNA sequencing, Shoot-root coordination

Abbreviations

TF	Transcription factor
DEG	Differentially expressed gene
FLs	First leaves
PEPC	Phosphoenolpyruvate carboxylase
BCA	Beta-carbonic anhydrase
PPdK	pyruvate orthophosphate dikinase
Asp-AT	Aspartate aminotransferas
Ala-AT	Alanine aminotransferase
NAD-ME	NAD-dependent malic enzyme
NADP-ME	NADP-dependent malic enzyme
TPT	Triosephosphate translocator
PPase6	Pyrophosphatase
PPdK-RP1	Pyruvate orthophosphate dikinase related protein1
PPT2	PEP/phosphate translocator
BASS4	Bile acid:sodium symporter family protein
BASS2	Bile acid:sodium symporter family protein
AMK2	Adenosine monophosphate kinase
HY5	Elongated hypocotyl 5

57   SHR   Short-Root

Introduction

As an adaptation to the hot and dry environment, C_4 photosynthesis is a carbon concentration mechanism that

reduces the counteracting photorespiration by suppressing the oxygenase activity of the carboxylating enzyme

RuBisCO (Ribulose-1,5-bisphosphate carboxylase/oxygenase). The leaves of C_4 plants have a unique structure

called Kranz anatomy, enabling them to spatially separate two phases of photosynthesis into the mesophyll and

bundle sheath cells. The Kranz anatomy precedes evolving C_4 biochemistry (McKown and Dengler 2007). A

universal feature of C_4 plants is their higher usage efficiency of resources such as nitrogen, water, and radiation

(Vogan and Sage 2011; Fatima *et al.* 2018). Nitrogen and water are uptaken by roots and are largely controlled

by the demands in the shoots. The indispensable shoot-root communication is coordinated by long-distance

signaling molecules such as phytohormones, signaling peptides, and TFs (Ko and Helariutta 2017).

69   Generally, in the C_4 pathway, the carbon fixation begins in the mesophyll, where carbon dioxide is

converted into bicarbonate and then to four-carbon (C_4) organic acids. The C_4 compounds are transported to the

bundle sheath and decarboxylated to release the CO_2 at the site of RuBisCO for assimilation (Sage 2004; Gowik

and Westhoff 2011). Based on the primary decarboxylating enzymes, NADP-ME and NAD-ME are two main

biochemical subtypes of the C_4 pathway (Rao and Dixon 2016). As a striking example of convergent evolution,

the conversion of C_3 into C_4 photosynthesis has evolved independently in more than 60 lineages of angiosperms

(Slewinski 2013). The polyphyletic origin of C_4 syndrome implies that the transition of C_3 into C_4 is not as

genetically complicated as inferred from the associated anatomical and biochemical features. It is suggested that

the pre-existing C_3 gene regulatory networks are recruited in the C_4 pathway with relatively few modifications

(Hibberd and Covshoff 2010; Reyna-Llorens and Hibberd 2017).

79   In addition to efforts distinguishing genes encoding C_4 enzymes, other studies have tried to identify the

genetic architecture and regulation of initiating the C_4 metabolism (Gowik *et al.* 2011; Wang, Vlad and Langdale

2016; Cui, 2021). For this transcriptome profiles were compared of C_3 and C_4 tissues of close relative (C_3 and C_4

Flaveria) (Gowik *et al.* 2011) or distantly related (C_3 rice and C_4 maize) (Wang *et al.* 2014) species. Most of the

omic analyses on C_4 photosynthesis have been performed in C_4 plants belonging to the NADP-ME subtype (Gowik

et al. 2011; Lauterbach *et al.* 2017). However, to comprehend the nature and commonality of regulatory features

of C_4 syndrome it is essential to investigate both biochemical subtypes. The nature and mechanism of distant

signaling molecules act in C_4 plants to coordinate the above- and under-ground tissues to maintain their

homeostasis and establish C_4 features, including Kranz anatomy, nitrogen, and water use efficiency, remains

largely obscure and has hampered the long-standing goal of engineering C_4 metabolism into the C_3 crops (Gerlich

et al. 2018).

Recently, species such as *Haloxylon ammodendron* (Y. Li *et al.* 2015) and *Salsola soda* (Lauterbach,

Billakurthi et al. 2017) have been found to display two photosynthetic mechanisms in their life cycle. As such

they provide an excellent model for studying the molecular regulatory mechanisms underlying the coordination

of roots and modified shoots during the C_3 to C_4 conversion while eliminating phylogenetic noises that hindered

earlier models. Unlike the C_3 pathway in cotyledons, which coincides with moderate temperature, the C_4 pathway

in these plants, which belongs to the NADP-ME subtype of C_4 photosynthesis, outperforms at higher light and

temperature conditions concurrent with the formation of the first leaf stage onwards (Lauterbach, Billakurthi et

al. 2017).

H. mollissima Bunge used in the present study belongs to the Caroxyleae tribe and performs both C_3 (in

cotyledons) and C_4 photosynthesis (NAD-ME subtype, in FLs onwards) (Akhani *et al.* 2009). To unravel the

signaling pathways associated with the C_3 to C_4 transition in *H. mollissima* Bunge, we used RNA sequencing to

generate transcriptome profiles of hypocotyls before and after forming the C_4 FLs alongside the FLs. Genes

associated with several phytohormone metabolisms, TFs, and signal peptides were differentially expressed in

hypocotyls before and after forming C_4 FLs. Our results indicated that, besides those associated with hypocotyl

development, some of these genes show homology to known shoot-to-root and vice-versa signals and their

receptors or transporters that are possibly implicated in establishing unique C_4 traits, especially in the efficient

consumption of resources such as nitrogen and water.

Material and Methods

Plant Cultivation and Sampling

Seeds of *H. mollissima* Bunge were collected at Mahdasht, Karaj, Iran ($35^o43^{'}35.4^{"}$N; $50^o44^{'}04.5^{"}$E) in November

2019 and stored at 4^{oC}. After removal of the hard seed coats, seeds were disinfected with ethanol 75%, washed

with water and germinated on the wet, sterilized filter paper in Petri dishes. After root formation, at 3 days after

incubation (DAI), seedlings were transferred into the plastic pots (5 cm diameter, 7 cm height) filled with sand,

perlite, and compost. *H. mollissima* Bunge Plants were grown in a greenhouse at the Institute of Biochemistry and

Biophysics, University of Tehran, at 25-30°C, ~200 µmol m^{-2} s^{-1} illumination and 16/8-hours light/dark cycle.

For sampling, two series of *H. mollissima* Bunge were cultivated. In the first series, expanded cotyledons,

hypocotyls, and roots (Series A) (Fig. 1(a)) were harvested before FLs were formed. In the second series, FLs,

cotyledons, hypocotyls, and roots (Series B) (Fig. 1(b)) were harvested after formation of FLs. Tissues of interest

were sampled in January and February 2020 between 11:00 and 13:00, shock-frozen in liquid nitrogen and stored

at -80°C. Two biological replicates of hypocotyls and FLs and three of all sampled tissues were used for RNA

sequencing and RT-qPCR, respectively, each containing pooled tissue from 5 independent plants.

Light Microscopy

For histological studies, C_3 cotyledon and C_4 first leaf were fixated for 16h in 50 mM Phosphate Puffer, pH 7.0,

containing 1 % (v/v) glutaraldehyde and 4 % (v/v) formaldehyde at 8°C. After 2 x 10 min washing with distilled

water and subsequent dehydration in an ascending ethanol series (30, 40, 50, 60, 75, 90 % and 2 × 100 % ethanol,

10 min each), probes were infiltrated with Spurr resin and polymerized in an oven at 70°C for 24 h. Semithin

sections of the middle regions of fully expanded cotyledons and FLs were made on a Reichert Ultracut S (Leica

Microsystems, Wetzlar, Germany) and stained with 1 % (w/v) methylene blue and 1 % (w/v) Azur II. Images

were taken on a Zeiss Axiovert 135 microscope (Carl Zeiss, Jena, Germany) and stored as TIFF files.

RNA Extraction and Sequencing

The guanidine/ phenol-based method (Chomczynski and Sacchi 1987) was used to isolate total RNA from 100-

150 mg of the homogenized samples. Quality, quantity, and integrity of isolated RNA were checked by Agarose

gel electrophoresis, Nanodrop (Thermo Fisher Scientific), and Agilent Bioanalyzer 2100 system (Agilent

Technologies Co. Ltd., Beijing, China), respectively. High-quality RNA (RIN value > 5.2) was used for cDNA

preparation and sequencing by Illumina Hiseq 2500 platform at the Novogene Bioinformatics Institute (Beijing,

China). One hundred fifty bp paired-end reads were obtained, and their adapter-contained and low-quality

(Qscore< = 5) base-contained were removed.

De novo Assembly and Functional Annotation

Raw   reads   were   evaluated   for   quality   by   FastQC   tool

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Due to removing adapters and low-quality reads by

the Novogene Bioinformatics Institute, there was no need for trimming. High-quality reads were assembled by

Trinity v2.4.0 (Grabherr *et al.* 2011) using default settings. First, probable open reading frames (ORF) were

predicted using the Transdecoder tool (Haas and Papanicolaou, 2016) with a minimum length of 100 amino acids.

To obtain Gene Ontology (GO) terms of the transcripts, Blast2GO software was used with $1e^3$ as an e-value cut-

off to perform BlastX against a non-redundant protein database. The online KEGG Automatic Annotation Server

(KAAS) (http://www.genome.jp/kegg/kaas) and Mapman V4.0 (https://www.plabipd.de/portal/mercator4)

(Schwacke *et al.* 2019) were used for pathway analysis (p-value ≤ 0.05). Also, the Arabidopsis TAIR locus IDs

were assigned to each transcript during Mapman analysis (https://www.plabipd.de/portal/web/guest/mercator-

sequence-annotation) (Lohse *et al.* 2014).

Determination of Genes with Different Expression

RSEM tool (http://deweylab.biostat.wisc.edu/RSEM) (Li and Dewey 2011) was used to estimate the abundance

of the transcripts for each sample. The high-quality raw reads were separately mapped back onto the assembled

transcripts using Bowtie 2.0. Differential expression analysis was performed by edgeR (Robinson, McCarthy and

Smyth 2010) in R after normalization using the trimmed mean of M-values (TMM) method. Statistical tests for

the pairwise comparisons were Benjamini-Hochberg correction FDR ≤ 0.05, $\log_2 FC \geq 1.5$ or ≤ -1.5 to set a

significant threshold for determining DEGs. Principal component analysis (PCA) with $\log_2$ transformed read count

(Transcript Per Million TPM) was performed in RStudio to visualize the expression pattern of the samples.

Furthermore, hierarchical clustering of the transcript abundance of selected known C_4 proteins using Pearson

correlation, average linkage method was performed (http://www.heatmapper.ca/expression/).

The DEGs most likely associated with the hypocotyl development of *H. mollissima* Bunge were excluded

from further analysis by comparing them with DEGs involved in the hypocotyl development in other plants of

flax   and   hemp   (Roach   and   Deyholos   2008;   Behr   *et   al.*   2018)

(https://bioinformatics.psb.ugent.be/webtools/Venn/).

Determination of Active Biological Functions

Pathways in each tissue of interest were explored using the percentage of total transcripts (normalized to

Transcript Per Million (TPM)) of each particular bincode in Mapman V4.0. According to the further emphasis of

this study on the survey of probable long-distance signaling, gene enrichment analysis was conducted by PageMan

software (Usadel *et al.* 2006) through Fisher's exact test followed by the Benjamini-Hochberg correction to

identify significant differences of Mapman bins among the two developmentally different hypocotyls. Moreover,

Mapman V3.6.0RC1 was used to visualize selected DEGs (through the generation of a custom MapMan pathway

image) encoding phytohormone synthesizing and degrading enzymes, transporter ("Phytohormone action bin"

and "Solute transporter"), and also genes related to different signal transduction pathways ("Multi-process

regulation. Calcium-dependent signaling", "Protein modification. Protein kinase" and "Redox homeostasis" bins).

Identification of TFs and Prediction of Cis-acting elements in the Promoter of Predicted Mobile

TFs

To predict differentially expressed TFs between hypocotyl before and after the formation of the FLs, the homology

of related amino acid sequences against the plant TFs database (PlantTFDB V5.0, http://planttfdb.gao-

lab.org/prediction.php) was implemented. Afterward, TFs with mobile mRNA, based on the information about

the genes with "cell-to-cell mobile mRNA" in The Arabidopsis Information Resource (TAIR) database, were

predicted.

Since the *H. mollissima* Bunge genome is not yet available, the promoter region of the orthologs of

predicted mobile TFs in Arabidopsis was obtained from the Arabidopsis Gene Regulatory Information Server

(AGRIS) database (https://agris-knowledgebase.org/AtcisDB/), and their cis-acting were predicted using

PlantCARE (http://bioinformatics.psb.ugent.be/webtools/plantcare/html/).

Real-time quantitative PCR (RT-qPCR)

To confirm RNA seq data, RT-qPCR was performed for eight genes. Total RNA was extracted by the above-

mentioned method (Chomczynski and Sacchi, 1987) and treated with DNase I (Thermo Fisher). RNA was reverse

transcribed with the AddScript cDNA Synthesis Kit (add bio), according to manufacturer's instructions, using 1

μg of RNA. RT-qPCR was performed with a 6plex real-time PCR system (Qiagen, Rotor Gen Q, Germany) using

the RealQ Plus 2x Master Mix Green kit, without ROX (AMPLIQON) using 1 μL of 1:40-fold diluted template

cDNAs. Run conditions were: 10 min of initial denaturation at 95°C, 95°C for 30s, 60°C for 30s, and 72°C for

30s (40 cycles). Melting curve analysis and agarose gel electrophoresis were done to verify the primer specificity.

Primer efficiency was estimated for each primer pair by four consecutive 10-fold dilutions of the cDNAs as a

template. The relative expression of genes was calculated using the $2^{-\Delta\Delta Ct}$ method (Livak and Schmittgen 2001)

using the Polyubiquitin ortholog from *H. mollissima* Bunge as an endogenous control. Three biological replicates

were quantified for each sample.

Results

Cotyledons and FLs of *H. mollissima* Bunge have Different Anatomical Features in Accordance with Their Photosynthesis Strategy

C_3 cotyledons of *H. mollissima* Bunge are flattened, bifacial, and their cross-section (Fig. 2 (a)) doesn't show any Kranz anatomy. They have several layers of chlorenchyma cells around the vascular bundles (VBs) and intercellular airspaces across the blades (Fig. 2 (a)). While C_4 first leaf is terete, linear, and covered by hairs, and its anatomy is salsoloid. The leaf cross section (Fig. 2 (b) and (c)) shows a large central water storage tissue which is respectively surrounded by the Kranz layer, palisade mesophyll cells, and a single-layered epidermis, while there is no space to be seen in the microscopic leaf structure.

Processing of Raw Reads and Statistics of De novo Assembly

The mRNA of three different tissues, including hypocotyls before (hypocotyl A) and after (hypocotyl B) the formation of FLs, and the FLs themselves (FL), were sequenced. The quality of reads was checked, and based on FastQC results (Phred score= 36) all reads had high quality (Additional file 1(A)) circumventing the need for further trimming. Raw data is available in the ? database under accession number ? (Additional file 2). To evaluate the Trinity results, N50 and total assembly length were checked by Trinitystat.pl (Additional file 1 (B)). PCA analysis (Fig. 3), where the first principal component contained 69.57% and the second principal component had 21.52% of the variance, demonstrated that the analyzed tissues have a distinct expression patterns. Nevertheless, the replicates' expression data were similar and grouped. As expected, the same tissues (hypocotyl A and hypocotyl B) had more similar expression patterns than the FLs.

Determination of Active Functional Pathways in analyzed tissues

To evaluate how the transcriptome as whole changes during the transition of photosynthesis strategy, the functional classes of each tissue transcriptome were determined as the percentage of all transcripts (normalized to Transcript Per Million (TPM)) assigned to each Mapman bincodes (Fig. 4(a), Additional file 3). MapMan analysis showed "Not assigned, annotated" and "Not assigned, not annotated" with 19.41 and 45.86% of all transcripts being the most significant functional gene categories in each sample. While with 15.44% the "Photosynthesis" bin was the third-highest abundant functional class in the FLs, in the hypocotyl A and B it accounted for only 3.25% and 2.47%, respectively. Out of 16 transcripts associated with known C_4 proteins, 14 transcripts, including Beta-carbonic anhydrase (BCA3), PEP carboxylase1 (PEPC1), pyruvate orthophosphate dikinase (PPdK),

aspartate aminotransferase (Asp-AT), alanine aminotransferase (Ala-AT), NAD-dependent malic enzyme (NAD-

ME), NADP-dependent malic enzyme (NADP-ME), Triosephosphate translocator (TPT), Pyrophosphatase

(PPase6), pyruvate orthophosphate dikinase related protein1 (PPdK-RP1), PEP/phosphate translocator (PPT2),

Bile acid:sodium symporter family protein (BASS4), Bile acid:sodium symporter family protein (BASS2),

Adenosine monophosphate kinase (AMK2) were more abundant in FLs than hypocotyls (Fig. 4(b), Additional

file 4). As expected, NAD-ME is more abundant than NADP-ME in FLs.

"Protein biosynthesis" and "Protein homeostasis" were the third- and fourth-highest abundant functional

categories in hypocotyl A (4.68 and 3.96%, respectively) and B (4.67 and 4%, respectively) (Fig. 4(a)). "RNA

biosynthesis" (1.92-2.34%) and "solute transport" (1.87-2.09%) were also abundant in all three samples

(Additional file 3). Although the abundance of most functional categories is similar in hypocotyl A and B, the

"RNA biosynthesis" (2.04 and 2.34 %, respectively) and "Phytohormone action" (1.2 and 1.31%, respectively)

classes are somewhat more abundant in the hypocotyl B.

The Pairwise comparison between the samples revealed several DEGs (FDR$\leq$ 0.05) among three

different developmental tissues (Additional file 5). In order to identify possible signaling pathways involved in

the C_3 to C_4 transition, pathways enrichment analysis was conducted for 8424 DEGs in the hypocotyl B vs.

hypocotyl A comparison. Among the pathways that according to the overrepresentation analysis appeared to be

up-regulatedly enriched in hypocotyl B were those involved in "Protein biosynthesis", "Protein homeostasis and

the ubiquitin-proteasome system" (Fig. 5).

Next, to explore possible long-distance regulators, we focus on the various regulatory mechanisms by

further analyzing DEGs in the two developmentally different hypocotyls. To differentiate between genes

controlling the transition of photosynthesis strategy and those involved in hypocotyl development, we compared

them (based on the assigned Arabidopsis IDs) with previously identified genes associated with hypocotyls

development (Roach and Deyholos, 2008; Behr *et al.* 2018). This way 190 genes were excluded (Additional file

6, Additional file 7). We then focused on the unique genes which do not overlap with related developmental genes.

DEGs Associated with Classical Phytohormones Metabolism

Mapping of unique DEGs of hypocotyl B vs. hypocotyl A in Mapman V3.6.0RC1 (Fig. 6, Additional file 8)

indicated several genes involved in the biosynthesis, degradation, transport, and signal transduction of plant

hormones (Phytohormone bin in Mapman) that are up- or down-regulated after the establishment of C_4

photosynthesis in the FLs.

Several transcripts encoding membrane bound auxin transporters, such as PIN-FORMED (PIN) (PIN2; At5g57090), and membrane-bound ATP-binding cassette-B (ABCB) auxin transporter (ABCB19; At3g28860), which are involved in long-distance auxin signaling proved to be less abundant in hypocotyl B compared to hypocotyl A. Other auxin-responsive genes like NITRATE TRANSPORTER 1 (NRT1; AT1G12110), an auxin-responsive transporter, were down-regulated in hypocotyl B after C_4 establishment, as was the flavin-dependent monooxygenase (YUCCA) involved in auxin biosynthesis. In contrast, transcripts of several PIN-Like transporters (PILS), which are distributed in the intracellular compartments and mediate intracellular auxin homeostasis, were more abundant in hypocotyl B. Also transcripts assigned to AUX/ IAA (AT5G43700, AT3G04730, and AT4G29080) and ARF7/ARF19 (AT1G19220) auxin signal transduction genes were up-regulated in hypocotyl B. Apparently, the biosynthesis and long-distance transport of auxin are reduced in hypocotyls after the formation of the FLs. In contrast to this, almost all transcripts associated with different cytokinin pathways bin (biosynthesis, signaling and transport) were up-regulated in hypocotyl B (Fig. 6). For example, type-B ARABIDOPSIS RESPONSE REGULATOR (ARR) and type-A ARR, involved in the cytokinin signaling, were up- and down-regulated in the hypocotyl B, respectively. Also, transcripts of ENT (EQUILIBRATIVE NUCLEOTIDE TRANSPORTER 1) (at4g05120, at1g70330), importers to take up apoplastic nucleosides of cytokinins, were more expressed in the hypocotyl B. DEGs associated with brassinosteroids were down-regulated.

Several long-range hormone transporters were more expressed in hypocotyl B (Fig. 6, Additional file 8), such as ABCG30 (AT4G15230), ABCG40 (AT1G15520), LHT1 (LYSINE-HISTIDINE TRANSPORTER) (at5g40780). NPF3.1 (Nitrate transport1/Peptide transporter family) (at1g68570), which is a multihormone transporter involved in the GA/ABA antagonism (Hirner *et al.* 2006), was down-regulated in hypocotyl B.

Differentially Expressed Signaling Peptides

Overrepresentation analysis of DEGs in hypocotyls before and after the formation of the FLs revealed that in contrast to the CRP classes of plant peptides (showing an overrepresentation of down-regulated genes), transcripts related to NCRP did not show enrichment in hypocotyls after the formation of the FLs (Fig. 5). Transcript abundance of different groups of these classes, based on Mapman bins, was then compared between analyzed tissues which showed a complex pattern of up and down-regulation (Fig. 7, Additional file 4). In general, transcripts belonging to different groups of the CRP categories were less abundant in hypocotyls after the establishment of C_4 photosynthesis in the FLs (Fig. 7 (a)). In this regard, transcripts assigned to several GASA

group members, including GASA1 (at1g74670), GASA8 (AT2G39540), GASA13 (AT3G10185), GASA14 (AT5G14920), and GASA 10 (AT5G59845) as well as RALF signaling peptides like RALF33 (AT4G15800) and its receptor CrRLK1L were down-regulated in hypocotyl after C_4 manifestation. Transcripts encoding peptides belonging to NCRP classes, including plant natriuretic peptide (PNP), CLAVATA3-EMBRYO-SURROUNDING REGION (CLE), Casparian strip integrity factor (CIF), PAMP-INDUCED SECRETED PEPTIDE (PIP) were more abundant in the hypocotyl after the formation of FLs (Fig. 6, Fig. 7 (b)). However, transcripts that are assigned to the receptor of C-terminally Encoded Peptides (CEP) such as CEPR2 (AT1G72180), acting in the nitrogen uptake signaling and Root meristem growth factor, were down-regulated in the hypocotyl after FLs formation.

DEGs Taken Part in Various Signal Transduction Pathways

The binding of the ligands to their receptors at the cell surface triggers intracellular reaction cascades to amplify the input signals, eventually altering gene expression in various pathways. We found that transcripts of genes involved in the transduction pathways were differentially regulated in the hypocotyls after forming the FLs (Fig. 6). Transcripts of several protein kinase genes belonging to the casein kinase (CK) and sterility (STE) protein kinase superfamilies, which are implicated in responses to many signals, were more abundant in hypocotyl after forming the FLs. Most genes belonging to the AGC (PKA-PKG-PKC), and CMGC (Cyclin-dependent kinases, Mitogen-activated protein kinases (MAPK), Glycogen synthase kinases, and Cyclin-dependent like kinases) superfamilies were more expressed in hypocotyl B, as were transcripts associated with receptor-like cytoplasmic protein kinases (RLCKs).

Also several genes involved in redox signaling and also those encoding for calmodulin-like proteins (CMLs) involved in plants Ca2+ signaling processes were up-regulated in the hypocotyl after the formation of the FLs (Fig. 6).

Differentially Expressed TFs

According to PlantTFDB V5.0, 290 out of 9640 ORFs from differentially expressed transcripts between hypocotyl before and after the formation of the FLs were annotated as different TF families with the best fit in *A. thaliana* locus IDs (Additional file 9). The mRNA levels of 71 and 219 of these TFs were down- and up-regulated, respectively, in hypocotyl after the formation of the FLs. In this comparison, C2H2, bHLH, and NAC families had a larger number of differentially regulated transcripts. Almost all transcripts belonging to C2H2, C3H, GRAS,

MYB, TCP, Trihelix, NF-Y, and WRKY TFs had more expression in hypocotyl after the formation of the FLs.

SHORT ROOT (SHR) (AT4G37650), a member of the GRAS family, and ELONGATED HYPOCOTYL 5

(HY5) (AT5G11260, AT3G17609), a member of the basic leucine zipper (bZIP) family, were up-regulated in

hypocotyl after the formation of the FLs. Notably, both SHR and HY5 are shown to have mobility and could

affect the expression of other genes (Dolan 2001; Chen *et al.* 2016).

Among 290 differentially expressed transcripts that were annotated as TFs based on PlantTFDB, 12 of

them had orthologous with cell-to-cell mobile mRNA, according to the TAIR database (Table 1). Eight of these

TFs were more abundant in hypocotyl after the formation of the FLs, including SCL1 (GRAS family,

AT1G21450), HAT22 (HD-ZIP family, AT4G37790), HSFA1E (HSF family, AT3G02990), CDC5

(MYB_related, AT1G09770), MYBH (MYB_related family, AT5G47390), NAC053 (NAC family,

AT3G10500), NAP (NAC family, AT1G69490) and ATWRKY40 (WRKY family, AT1G808$_4$0).

To predict if specific cis-regulatory elements are overrepresented in the regulatory region of candidate

mobile TFs with differentially expression levels in hypocotyls, PlantCARE (Lescot *et al.* 2002) was used to

analyze the upstream promoter elements of the Arabidopsis orthologs of these transcripts. The results showed that

Arabidopsis orthologs of differentially regulated mobile TFs carry several different cis-elements responsive to

phytohormones and environmental stimuli such as light and temperature. Among these, Box 4 (part of a conserved

DNA module involved in light responsiveness), AE-box (part of a module for light response), G-Box (cis-acting

regulatory element involved in light responsiveness), and GT1-motif (light-responsive element) were more

common light-responsive elements in the upstream of the mobile TFs (Additional file 10). Besides, the ABA-

responsive element (ABRE) was predicted in the promoter of BOA, SCL1, HAT22, MYBH, NAC053, ATAF1,

NAP, and WRKY40 TFs. Other motifs related to stress responses, including the MYB binding site involved in

drought-inducibility (MBS), a cis-acting element involved in defense and stress responsiveness (TC-rich repeat),

and a cis-acting regulatory element involved in the MeJA-responsiveness (CGTCA-motif and TGACG-motif),

were predicted to be present in the upstream region of the orthologs of differentially regulated mobile TFs genes

after the formation of the FLs.

To confirm RNA sequencing results, RT-qPCR was performed for eight target genes. The results were

correlated with the RNA seq analysis (R^2=0.81; Additional file 11). The RT-qPCR data analysis confirmed a

higher abundance of the C$_4$ enzymes, PEPC, BCA, and NAD-ME encoding genes in the FLs compared to the

cotyledons (Fig. 8(a)). The expression pattern of the analyzed genes somewhat varied between developmentally

different cotyledons and hypocotyls (Fig. 8(b), 8(d)). PEPC, BCA, NAD-ME, CEPR2, AUX/ IAA, ABCG40,

WRKY40, and NRT1.1 were less expressed in the roots after the formation of the FLs (Fig. 8(c)).

Discussion

Species with Switching Mechanisms are Valuable Models for studying C$_4$ Photosynthesis

We investigated the transcriptome profile of *H. mollissima* Bunge hypocotyls before and after first leaf formation

to identify possible components of long-distance signaling pathways involved in the unique C$_4$ traits through the

connection between the above- and under-ground tissues. *H. mollissima* Bunge belongs to the Caroxyleae tribe

and has two photosynthetic mechanisms in its annual life cycle; C$_3$ in cotyledons and NAD-ME subtype of C$_4$ in

the FLs onwards. Anatomical studies of cotyledons and FLs (Fig. 2) reveal features related to C$_3$ and C$_4$

(Salsoloid) (Akhani *et al.* 2009; Freitag and Kadereit 2014), respectively. In leaves the proportion of water storage

tissue was larger compared to the cotyledons while air space volume was less. This correlates with the

development of functional and efficient C$_4$ photosynthesis. Akhani et al. (Akhani *et al.* 2009) who determined the

carbon isotope composition of cotyledons and leaves of *H. mollissima* Bunge, found that cotyledons and leaves

have C$_3$ and C$_4$ type isotope values, respectively. This fits with our observations of typical C$_4$ genes (PEPC, BCA,

and NAD-ME) being higher expressed in the FLs (Fig. 8(A)).

A C$_3$ to C$_4$ switching mechanism has been reported for several species in the Salsoloideae (Caroxyleae

and Salsoleae tribes) and Suaedoideae subfamilies (Rudov *et al.* 2020). Thus, their germination usually coincides

with low-temperature conditions that possibly favor C$_3$ photosynthesis in the cotyledons. As these species grow

and the temperature rises, C$_4$ photosynthesis is encouraged in the leaves (Rudov *et al.* 2020). Plants expressing

such C$_3$ to C$_4$ switching may be appropriate models to explore the C$_4$ pathway and its regulatory mechanisms due

their lack of phylogenetic noises and interspecies differences. Transcriptome analysis of *Haloxylon ammodendron*

(Y. Li *et al.* 2015) and *Salsola soda* (Lauterbach, Billakurthi et al. 2017) both belonging to the *Salsoleae* tribe and

performing a NADP-ME subtype of C$_4$ photosynthesis in true leaves, revealed differentially expressed C$_4$ pathway

genes, such as PEPC between C$_3$ cotyledons and C$_4$ true leaves, but paid less attention to possible differences in

regulatory pathways. In the present study we investigated potential shoot-root communications during the C$_3$ to

NAD-ME subtype of C$_4$ transition by focusing on the DEGs in hypocotyls before and after the C$_4$ first leave's

formation.

Novel Consistent Data Provided by *De novo* Assembly of *H. mollissima* Bunge Transcriptome

Most transcriptome data analyses in the field of C$_4$ photosynthesis, with a few exceptions, have used a reference-

based approach to annotate their data using the Arabidopsis data (Gowik and Westhoff 2011; Lauterbach,

Billakurthi *et al.* 2017; Lauterbach, Schmidt *et al.* 2017; Siadjeu, Lauterbach and Kadereit 2021). As a non-model

species from the Caroxyleae tribe whose genome is not sequenced, we used RNA-seq to de novo assemble and

characterize the *H. mollissima* Bunge hypocotyls and FLs transcriptome.

PCA (Fig. 3) and analysis of functional classes of annotated genes (Fig. 4(a)) confirm the sequencing

data's reasonable and comparable relationship. Based on functional annotation, FLs expressed the NAD-ME

subtype of the C$_4$ photosynthesis pathway. The expression of photosynthesis-related genes was reduced in

hypocotyls after establishing the photosynthetic leaves. The formation of FLs also coincides with increased

translation processes in the hypocotyls an example of this being the upregulation of the ubiquitin-proteasome

system, a critical regulator of many plants signaling pathways in respond to internal and external cues

(Sadanandom *et al.* 2012).

Possible Regulatory Mechanisms Implicated in the Unique C$_4$ Features through Adjustment of

Shoot and Root Communication

Several regulatory systems with overlapping functions control the developmental processes in plants. These

include among others phytohormones, signaling peptides, and complex network of TFs. Long-range regulators

are an essential part of these systems. C$_4$ photosynthesis which is seen as a response to a reduced atmospheric

CO$_2$, requires a coordination between the root and the modified shoot. Similarly, the onset of the C$_4$ pathway in

leaves, followed by increased photosynthesis rate and more efficient use of resources, requires coordination

between roots and the modified shoots via signaling molecules. Interspecies grafting experiments between C$_3$

Flaveria robusta and C$_4$ *Flaveria bidentis* indicated that the root preliminary controls sulfur allocation between

roots and shoots and sulfur homeostasis in C$_4$ *Flaveria* (Gerlich *et al.* 2018). The communication ways between

above- and below-ground plant parts are very diverse. They can be physical (e.g. propagating Ca^{2+} waves),

chemical (e.g. signaling peptides and hormones), or molecular (e.g. RNA and proteins) (Shabala *et al.* 2015).

These signals can move through vasculature systems or cell-by-cell via stems or stem-like tissue (Bartusch and

Melnyk 2020). Also, the phloem comprises several mobile developmental signals (mRNAs, proteins, peptides,

e.g.), which can interplay in the different stages of development and growth (Koenig and Hoffmann-Benning

2020). Long-range movement of regulatory molecules has already been demonstrated, for auxin (Bellstaedt *et al.*

2019), cytokinin, abscisic acid (Kiba *et al.* 2011), various signaling peptides such as CEP, specific transporters or

receptors and some TFs (Shabala *et al.* 2015). By comparing transcriptome profiles of hypocotyls during the C_3

to C_4 transition, we revealed that transcripts of genes encoding components of different phytohormone

metabolisms, signaling peptides, and TFs (Fig. 6, Additional file 8) were differentially expressed in hypocotyls

after establishing the C_4 leaves. In addition, the expression levels of several transporters, receptors, and responsive

genes of root- or leaf-derived signals were also affected, including those of PIN1, CEPR, and ARF7/ARF19 (Fig.

6, Additional file 8). Some of these regulators are most likely implicated in the normal development of hypocotyls.

Therefore, to distinguish true C_4-related genes, we used a literature survey to exclude those genes involved in the

hypocotyl development (Additional file 6, Additional file 7). The mobility of mRNA as long-distance signal is

supposed to be a complex process while elucidating its physiological roles remains challenging (Xia *et al.* 2018).

Nitrogen Use Efficiency: The Reduction of Shoot-to-Root Nitrogen Status Signals after the

Formation of the C_4 FLs

Nitrogen assimilation by roots can be controlled by a number of signals representing the nitrogen status in the

shoots (Ruffel *et al.* 2008; Forde 2002). These signals include phytohormones such as auxin and cytokinin (Kiba

et al. 2011). Our results showed that several genes involved in the biosynthesis, transport, and biodegradation of

phytohormones were differentially expressed in the hypocotyls after formation of the FLs. Cytokinin signaling

increased after the establishment of C_4 photosynthesis. Shoot-derived cytokinin is a signal of nitrogen satiety in

well-supplied nitrogen conditions, suppressing the nitrogen uptake by roots (Kiba *et al.* 2011). Our findings

suggest that long-range cytokinin signaling can be a candidate pathway for controlling nitrogen uptake in C_4

plants. No particular cytokinin transporters for this shoot-to-root movement through the phloem has been

identified yet, but ENTs are likely involved (Liu, Zhao and Zhang 2019). Type-B ARR is the positive regulator

of cytokinin signaling (Argyros *et al.* 2008) and type-A ARR is the negative regulator of cytokinin signaling (To

et al. 2004) which have their corresponding transcripts (AT1G67710 and AT3G57040) up- and down-regulated,

respectively, in the hypocotyl B.

Due to its role in lateral root initiation, auxin has long been known as a shoot-to-root nitrogen signal (Forde 2002;

Fukaki and Tasaka 2009). In the C_4 crop maize, auxin concentration in phloem exudates decreases under high

nitrate concentrations (Tian *et al.* 2008). In the same line, our data showed that cell-to-cell auxin transporters,

responsible for auxin's long-distance movement and an enzyme catalyzing the rate-limiting step in auxin

biosynthesis, were down-regulated in hypocotyls B. In contrast, the intracellular auxin transporters that modulate

the nuclear abundance of auxin by transporting auxin into the ER lumen (Sauer and Kleine-Vehn 2019) were up-

regulated. These findings imply that long-distance movement of auxin is reduced in hypocotyls after C_4 leaf

formation. At the same time, the NRT1.1 transporter becomes down-regulated in hypocotyls B. As a dual-affinity

transporter, NRT1.1 acts as a nitrate sensor that regulates the expression level of genes involved in nitrate transport

(Sun and Zheng 2015). It is located in roots, vascular cells, and shoots and displays an auxin-responsive activity.

The reduced expression in hypocotyl B is consistent with other findings in this study, implying a general reduction

of auxin content, either derived from leaves or the hypocotyl itself, in the hypocotyl B. Furthermore, NRT1.1 has

a significant role in the nitrogen uptake in roots affected by shoot-derived signals. Thus, its lower expression in

roots after formation of the first leaves (Fig. 8(c)) may be a response to a decrease in the nitrogen uptake by roots

after C_4 establishment. Other reports claim, however, that local increases of auxin biosynthesis and transport in

developing leaves is necessary for manifestation of C_4 key property of higher vein density (Huang *et al.* 2017).

A further mobile regulator mediating nitrogen assimilation in roots under nitrogen deficiency condition,

is CEP which has a shoot orientation from the roots. CEP receptor, like CEPR2, was down-regulated in hypocotyl

B (Fig. 8(d)), most likely due to the efficient use of nitrogen by C_4 plants. This likely reflects the reduced effect

of root-derived CEP signal after C_4 leave's formation. Indeed, this receptor implicates nitrogen uptake signaling

by enhancing the expression of NRT1(Tabata *et al.* 2014; Ohkubo *et al.* 2017). The observed lower NRT1

expression in roots after C_4 first leave's formation (Fig. 8(c)) is thus consistent with possible reduced root-derived

CEP. Notably, CEPR2 expression was higher in the cotyledon B (C_3) compared to the FLs (C_4) (Fig. 8(a)),

indicating a higher level of nitrogen consumption by C_3 cotyledons than by C_4 leaves.

Taken together, gene expression changes after emerging C_4 photosynthesis in the FLs indicate modulated

auxin and cytokinin metabolism and signaling alongside a reduced CEP signaling in hypocotyl B and FLs. This

supports the notion of an enhanced nitrogen use efficiency and reduction in total nitrogen requirement in C_4 plants

(Jobe *et al.* 2020).

Drought Resistance and Water Use Efficiency: Several Pathways involved in the High-Temperature Resistance after the Formation of the C_4 FLs

Through different mechanisms, C_4 plants produce more efficiently than C_3 under higher temperatures. One

mechanism is preventing dehydration through stomatal closure. Stomatal closure under drought stress is regulated

by root-derived abscisic acid (Kuromori, Seo and Shinozaki 2018). Some studies claim that stomata of C_4 plants

are more sensitive to intercellular CO_2 concentration (Ci) compared to C_3 (Huxman and Monson 2003), and that

abscisic acid can increase this sensitivity (Dubbe, Farquhar and Raschke 1978). We observed that the expression

level of several abscisic acid influx genes (ABCG30, ABCG40) (Zhang *et al.* 2014; Kang *et al.* 2015) was up-

regulated in hypocotyls B compared to hypocotyls A (Fig. 6, Fig. 8(d)). Our finding for up-regulation of ABCG40

in C_4 leaves compared to C_3 cotyledons (Fig. 8(a)) is consistent with its functions in shoots as a root-derived-

abscisic acid importer and its essential role in stomatal closure (Kang *et al.* 2010). The differential regulation of

ABCG40 in roots (Fig. 8(d)) likely is due the interaction between abscisic acid and nitrate uptake (Harris and

Ondzighi-Assoume 2017). Moreover, studies on the amphibious *Eleocharis vivipara* have revealed the significant

role of abscisic acid and hormonal control in the manifestation of Kranz anatomy (Ueno 1998).

Stress- and Light-Responsive Differentially expressed Mobile TFs in Hypocotyls are Possibly

Involved in the Formation of C_4 Unique Features

Many of the TFs that are up-regulated in the hypocotyls after the formation of the FLs are implicated in the

regulatory networks controlling responses to the different plant development and growth conditions as well as

abiotic stresses. The appearance of C_4 leaves in *H. mollissima* Bunge, and other halophytes or xerohalophytes

switching chenopods is coincides with an increase in environmental temperature (Rudov *et al.* 2020).

Transcriptome analysis of C_3 cotyledons and C_4 leaves in *Salsola soda* demonstrated that 55 and 32 TFs were

increased and decreased in C_4 leaves compared to C_3 cotyledons, respectively (Lauterbach, Billakurthi et al. 2017).

Not surprisingly, a greater number of the up-regulated TFs in our study were stress-responsive, including NAC

(Shao, Wang and Tang, 2015), WRKY (Wang *et al.* 2018), and NF-Y (Zhao *et al.* 2017).

Among the differentially expressed TFs in hypocotyls before and after the formation of the C_4 FLs, SHR

and HY5 TFs are known to be mobile although at protein level (Dolan 2001; Chen *et al.* 2016). SHR is implicated

in modulating endodermis (known as bundle-sheath) differentiation in leaves and apparently plays a role in the

Kranz anatomy formation in maize (Slewinski *et al.* 2012). Moreover, Siadjeu et al. (Siadjeu, Lauterbach and

Kadereit 2021) who compared transcriptome data of C_3, C_2, and C_4 species from *Camphorosmeae*, found that,

BBX15, SHR, SCZ, and LBD41 are more expressed and co-regulated in the C_4 than C_3 species. Interestingly, out

of 38 differentially expressed Arabidopsis orthologous TFs in the comparative transcriptomic analysis between

Kranz (foliar leaf blade) and non-Kranz (husk leaf sheath) maize leaves, we identified four TFs that express more

in hypocotyl after C_4 leave's formation; including SHR (AT4G37650), BHLH96 (AT1G72210), GATA7

(AT4G36240) and WRKY12 (AT2G44745).

Most of the up-regulated TFs after C_4 manifestation belonged to C2H2 and bZIP families, including

HY5. HY5 is a master regulator of transcription (Gangappa and Botto 2016), a phloem-mediated shoot-to-root

signal for lateral root formation, and thus the carbon-nitrogen balance adjustment in Arabidopsis (Burko *et al.*

2020). HY5 mediates photomorphogenesis so that it is located downstream of phytochrome B, the sensor of light

and temperature (Legris *et al.* 2016). Thus, in C_4 plants where shoots are exposed to high light intensity and

temperature, and since the function of HY5 is conserved in many plants (Yamawaki *et al.* 2011; Huai, Jing and

Lin 2020), HY5 can be an appropriate candidate for synchronizing shoots and roots. Moreover, HY5 is one of the

light-responsive TFs essential in inhibiting hypocotyl elongation during seedling (Oyama, Shimura and Okada

1997). Therefore, HY5 is a shared regulator between several various pathways.

The mRNA of 12 differentially expressed TFs in the *H. mollissima* Bunge hypocotyls have cell-to-cell

mobility, according to the TAIR database (Table 1). We also predicted the probable cis-elements present in the

promoter region of their orthologous in *A. thaliana*. Several of the predicted cis-elements in the mobile TFs were

light- and stress-responsive (Additional file 10). This indicates that expression and transport of these TFs is

influenced by abiotic stresses. One of these TFs is WRKY40, an orthologue of which, ZmWRKY40, is a drought-

responsive gene in *Zea mays* (Wang *et al.* 2018). HSFA1e was more expressed in hypocotyl B, which is implicated

in the tolerance to heat shock stress via transcriptional regulation of HsfA2 function (Nishizawa-Yokoi *et al.*

2011). Overexpression of maize ZmHsf06 in Arabidopsis, improves drought-stress tolerance (H. Li *et al.* 2015).

The findings of this study pave the way for identifying possible regulators underlying the coordination of

developmental changes associated with functional C_4 photosynthesis. Also, by investigating a plant with a NAD-

ME subtype of C_4 photosynthesis our data fills the gap caused by previous studies solely concentrating on the

NADP-ME subtype of photosynthesis. Our data provides a valuable source for further functional genomics and

genetic studies to identify master regulators of C_4-related traits and engineering C_4 features into C_3 crops.

Funding

A.M.B-M gratefully acknowledges the financial support from the Iran National Science Foundation (INSF)

[funding reference number 95838484].

Author Contribution

MZ performed the experiments, analyzed data, and wrote the original draft. TR performed the microscopic

analysis and critically revised the manuscript. MRG analyzed data. AMB-M conceived and designed the research,

supervised the experiment, and critically revised the manuscript. All authors contributed to the article and

approved the submitted version.

Acknowledgment

We thank Stanislav Kopriva (University of Cologne) for the critical reading of the manuscript. We also would like to thank Alexander Rudov and Hossein Akhani (University of Tehran) for their support during seed collection.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Additional information

Supplementary Information

Supplementary file 1 Summary of A) sequencing information B) De novo transcriptome assembly results by Trinity

Supplementary file 2. The accession number of uploaded raw sequences to ??

Supplementary file 3. Active functional classes in the First leaves, hypocotyl A and hypocotyl B

Supplementary file 4. Abundance (total mean TPM) of C_4-related proteins and genes categorized in the CRP and NCRP signaling peptide groups

Supplementary file 5. Transcript abundance, statistics of differentially expressed genes, and their annotations.

Additional file 6. Venn diagram analysis of DEGs of two developmental *H.mollissima*'s hypocotyls and identified genes related to the hypocotyl developments in flax (Roach and Deyholos, 2008) and hemp (Behr *et al.* 2018) based on their assigned Arabidopsis ID

Supplementary file 7. List of the unique and shared genes compared to DEGs of two developmental *H.mollissima*'s hypocotyls and identified genes related to the hypocotyl developments in flax (Roach and Deyholos, 2008) and hemp (Behr *et al.* 2018) based on their assigned Arabidopsis ID.

Supplementary file 8. Functional analysis of the Cell signaling genes using Mapman

Supplementary file 9. Differentially transcription factor genes between hypocotyl A and hypocotyl B using PlantTFDB v5.0

535   **Supplementary file 10.** Cis-elements located upstream of orthologous transcription factor genes with mobile

536   mRNA in Arabidopsis

537   **Additional file 11** Validation of RNA-Seq results using quantitative RT-PCR (qRT-PCR)

References:

Akhani, H. *et al.* (2009) 'Does Bienertia cycloptera with the single-cell system of C4 photosynthesis exhibit a
seasonal pattern of δ13C values in nature similar to co-existing C4 Chenopodiaceae having the dual-cell (Kranz)
system?', *Photosynthesis Research*, 99(1), pp. 23–36.

Argyros, R.D. *et al.* (2008) 'Type B response regulators of Arabidopsis play key roles in cytokinin signaling and
plant development', *The Plant Cell*, 20(8), pp. 2102–2116.

Bartusch, K. and Melnyk, C.W. (2020) 'Insights into plant surgery: an overview of the multiple grafting
techniques for Arabidopsis Thaliana', *Frontiers in Plant Science*, p. 2000.

Behr, M. *et al.* (2018) 'Insights into the molecular regulation of monolignol-derived product biosynthesis in the
growing hemp hypocotyl', *BMC plant biology*, 18(1), pp. 1–18.

Bellstaedt, J. *et al.* (2019) 'A mobile auxin signal connects temperature sensing in cotyledons with growth
responses in hypocotyls', *Plant Physiology*, 180(2), pp. 757–766.

Burko, Y. *et al.* (2020) 'Local HY5 activity mediates hypocotyl growth and shoot-to-root communication', *Plant
communications*, 1(5), p. 100078.

Chen, X. *et al.* (2016) 'Shoot-to-root mobile transcription factor HY5 coordinates plant carbon and nitrogen
acquisition', *Current Biology*, 26(5), pp. 640–646.

Chomczynski, P. and Sacchi, N. (1987) 'Single-step method of RNA isolation by acid guanidinium thiocyanate-
phenol-chloroform extraction', *Analytical biochemistry*, 162(1), pp. 156–159.

Cui, H. (2021) 'Challenges and Approaches to Crop Improvement Through C3-to-C4 Engineering', *Frontiers in
Plant Science*, p. 1851.

Dolan, L. (2001) 'Root patterning: SHORT ROOT on the move', *Current biology*, 11(23), pp. R983–R985.

Dubbe, D.R., Farquhar, G.D. and Raschke, K. (1978) 'Effect of abscisic acid on the gain of the feedback loop
involving carbon dioxide and stomata', *Plant Physiology*, 62(3), pp. 413–417.

Fatima, Z. *et al.* (2018) 'Resource use efficiencies of C3 and C4 cereals under split nitrogen regimes', *Agronomy*,
8(5), p. 69.

Forde, B.G. (2002) 'Local and long-range signaling pathways regulating plant responses to nitrate', *Annual
Review of Plant Biology*, 53(1), pp. 203–224.

Freitag, H. and Kadereit, G. (2014) 'C3 and C4 leaf anatomy types in Camphorosmeae (Camphorosmoideae,
Chenopodiaceae)', *Plant Systematics and Evolution*, 300(4), pp. 665–687.

Fukaki, H. and Tasaka, M. (2009) 'Hormone interactions during lateral root formation', *Plant molecular biology*,
69(4), pp. 437–449.

Gangappa, S.N. and Botto, J.F. (2016) 'The multifaceted roles of HY5 in plant growth and development',
Molecular plant, 9(10), pp. 1353–1365.

Gerlich, S.C. *et al.* (2018) 'Sulfate metabolism in C4 Flaveria species is controlled by the root and connected to
serine biosynthesis', *Plant physiology*, 178(2), pp. 565–582.

Gowik, U. *et al.* (2011) 'Evolution of C4 photosynthesis in the genus Flaveria: how many and which genes does
it take to make C4?', *The Plant Cell*, 23(6), pp. 2087–2105.

Gowik, U. and Westhoff, P. (2011) 'The path from C3 to C4 photosynthesis', *Plant Physiology*, 155(1), pp. 56–
63.

Grabherr, M.G. *et al.* (2011) 'Full-length transcriptome assembly from RNA-Seq data without a reference
genome', *Nature biotechnology*, 29(7), pp. 644–652.

Haas, B. and Papanicolaou, A. (2016) 'TransDecoder (find coding regions within transcripts)', *Google Scholar*
[Preprint].

Harris, J.M. and Ondzighi-Assoume, C.A. (2017) 'Environmental nitrate signals through abscisic acid in the root
tip', *Plant signaling & behavior*, 12(1), p. e1273303.

Hibberd, J.M. and Covshoff, S. (2010) 'The regulation of gene expression required for C4 photosynthesis', *Annual*
review of plant biology, 61, pp. 181–207.

Hirner, A. *et al.* (2006) 'Arabidopsis LHT1 is a high-affinity transporter for cellular amino acid uptake in both
root epidermis and leaf mesophyll', *The Plant Cell*, 18(8), pp. 1931–1946.

Huai, J., Jing, Y. and Lin, R. (2020) 'Functional analysis of ZmCOP1 and ZmHY5 reveals conserved light
signaling mechanism in maize and Arabidopsis', *Physiologia Plantarum*, 169(3), pp. 369–379.

Huang, C.-F. *et al.* (2017) 'Elevated auxin biosynthesis and transport underlie high vein density in C4 leaves',
Proceedings of the National Academy of Sciences, 114(33), pp. E6884–E6891.

Huxman, T.E. and Monson, R.K. (2003) 'Stomatal responses of C3, C3-C4 and C4Flaveria species to light and
intercellular CO2 concentration: implications for the evolution of stomatal behaviour', *Plant, Cell & Environment*,
26(2), pp. 313–322.

Jobe, T.O. *et al.* (2020) 'Ensuring nutritious food under elevated CO2 conditions: A case for improved C4 crops',
Frontiers in Plant Science, 11, p. 1267.

Kang, J. *et al.* (2010) 'PDR-type ABC transporter mediates cellular uptake of the phytohormone abscisic acid',
Proceedings of the National Academy of sciences, 107(5), pp. 2355–2360.

Kang, J. *et al.* (2015) 'Abscisic acid transporters cooperate to control seed germination', *Nature communications*,
6(1), pp. 1–10.

Kiba, T. *et al.* (2011) 'Hormonal control of nitrogen acquisition: roles of auxin, abscisic acid, and cytokinin',
Journal of experimental botany, 62(4), pp. 1399–1409.

Ko, D. and Helariutta, Y. (2017) 'Shoot–root communication in flowering plants', *Current Biology*, 27(17), pp.
R973–R978.

Koenig, A.M. and Hoffmann-Benning, S. (2020) 'The interplay of phloem-mobile signals in plant development
and stress response', *Bioscience Reports*, 40(10).

Kuromori, T., Seo, M. and Shinozaki, K. (2018) 'ABA transport and plant water stress responses', *Trends in plant*
science, 23(6), pp. 513–522.

Lauterbach, M., Billakurthi, K., *et al.* (2017) 'C3 cotyledons are followed by C4 leaves: intra-individual
transcriptome analysis of Salsola soda (Chenopodiaceae)', *Journal of experimental botany*, 68(2), pp. 161–176.

Lauterbach, M., Schmidt, H., *et al.* (2017) 'De novo transcriptome assembly and comparison of C3, C3-C4, and
C4 species of tribe Salsoleae (Chenopodiaceae)', *Frontiers in plant science*, 8, p. 1939.

Legris, M. *et al.* (2016) 'Phytochrome B integrates light and temperature signals in Arabidopsis', *Science*,
354(6314), pp. 897–900.

Lescot, M. *et al.* (2002) 'PlantCARE, a database of plant cis-acting regulatory elements and a portal to tools for
in silico analysis of promoter sequences', *Nucleic acids research*, 30(1), pp. 325–327.

Li, B. and Dewey, C.N. (2011) 'RSEM: accurate transcript quantification from RNA-Seq data with or without a
reference genome', *BMC bioinformatics*, 12(1), pp. 1–16.

Li, H. *et al.* (2015) 'Expression of maize heat shock transcription factor gene ZmHsf06 enhances the
thermotolerance and drought-stress tolerance of transgenic Arabidopsis', *Functional Plant Biology*, 42(11), pp.
1080–1091.

Li, Y. *et al.* (2015) 'Developmental genetic mechanisms of C4 syndrome based on transcriptome analysis of C3
cotyledons and C4 assimilating shoots in Haloxylon ammodendron', *PLoS One*, 10(2), p. e0117175.

Liu, C.-J., Zhao, Y. and Zhang, K. (2019) 'Cytokinin transporters: multisite players in cytokinin homeostasis and
signal distribution', *Frontiers in Plant Science*, p. 693.

Livak, K.J. and Schmittgen, T.D. (2001) 'Analysis of relative gene expression data using real-time quantitative
PCR and the 2− ∆∆CT method', *methods*, 25(4), pp. 402–408.

Lohse, M. *et al.* (2014) *M ercator: a fast and simple web server for genome scale functional annotation of plant*
sequence data. Wiley Online Library.

McKown, A.D. and Dengler, N.G. (2007) 'Key innovations in the evolution of Kranz anatomy and C4 vein pattern
in Flaveria (Asteraceae)', *American journal of botany*, 94(3), pp. 382–399.

Nishizawa-Yokoi, A. *et al.* (2011) 'HsfA1d and HsfA1e involved in the transcriptional regulation of HsfA2
function as key regulators for the Hsf signaling network in response to environmental stress', *Plant and Cell*
Physiology, 52(5), pp. 933–945.

Ohkubo, Y. *et al.* (2017) 'Shoot-to-root mobile polypeptides involved in systemic regulation of nitrogen
acquisition', *Nature plants*, 3(4), pp. 1–6.

Oyama, T., Shimura, Y. and Okada, K. (1997) 'The Arabidopsis HY5 gene encodes a bZIP protein that regulates
stimulus-induced development of root and hypocotyl', *Genes & development*, 11(22), pp. 2983–2995.

Rao, X. and Dixon, R.A. (2016) 'The differences between NAD-ME and NADP-ME subtypes of C4
photosynthesis: more than decarboxylating enzymes', *Frontiers in plant science*, 7, p. 1525.

Reyna-Llorens, I. and Hibberd, J.M. (2017) 'Recruitment of pre-existing networks during the evolution of C4
photosynthesis', *Philosophical Transactions of the Royal Society B: Biological Sciences*, 372(1730), p. 20160386.

Roach, M.J. and Deyholos, M.K. (2008) 'Microarray analysis of developing flax hypocotyls identifies novel
transcripts correlated with specific stages of phloem fibre differentiation', *Annals of botany*, 102(3), pp. 317–330.

Robinson, M.D., McCarthy, D.J. and Smyth, G.K. (2010) 'edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data', *Bioinformatics*, 26(1), pp. 139–140.

Rudov, A. *et al.* (2020) 'A review of C4 plants in southwest Asia: An ecological, geographical and taxonomical
analysis of a region with high diversity of C4 eudicots', *Frontiers in plant science*, 11, p. 1374.

Ruffel, S. *et al.* (2008) 'Systemic signaling of the plant nitrogen status triggers specific transcriptome responses
depending on the nitrogen source in Medicago truncatula', *Plant physiology*, 146(4), pp. 2020–2035.

Sadanandom, A. *et al.* (2012) 'The ubiquitin–proteasome system: central modifier of plant signalling', *New*
Phytologist, 196(1), pp. 13–28.

Sage, R.F. (2004) 'The evolution of C4 photosynthesis', *New phytologist*, 161(2), pp. 341–370.

Sauer, M. and Kleine-Vehn, J. (2019) 'PIN-FORMED and PIN-LIKES auxin transport facilitators', *Development*,
146(15), p. dev168088.

Schwacke, R. *et al.* (2019) 'MapMan4: a refined protein classification and annotation framework applicable to
multi-omics data analysis', *Molecular plant*, 12(6), pp. 879–892.

Shabala, S. *et al.* (2015) 'Root-to-shoot signalling: integration of diverse molecules, pathways and functions',
Functional Plant Biology, 43(2), pp. 87–104.

Shao, H., Wang, H. and Tang, X. (2015) 'NAC transcription factors in plant multiple abiotic stress responses:
progress and prospects', *Frontiers in plant science*, 6, p. 902.

Siadjeu, C., Lauterbach, M. and Kadereit, G. (2021) 'Insights into regulation of C2 and C4 photosynthesis in
Amaranthaceae/Chenopodiaceae using RNA-Seq', *International journal of molecular sciences*, 22(22), p. 12120.

Slewinski, T.L. *et al.* (2012) 'Scarecrow plays a role in establishing Kranz anatomy in maize leaves', *Plant and
Cell Physiology*, 53(12), pp. 2030–2037.

Slewinski, T.L. (2013) 'Using evolution as a guide to engineer Kranz-type C4 photosynthesis', *Frontiers in plant
science*, 4, p. 212.

Sun, J. and Zheng, N. (2015) 'Molecular mechanism underlying the plant NRT1. 1 dual-affinity nitrate
transporter', *Frontiers in Physiology*, 6, p. 386.

Tabata, R. *et al.* (2014) 'Perception of root-derived peptides by shoot LRR-RKs mediates systemic N-demand
signaling', *Science*, 346(6207), pp. 343–346.

Tian, Q. *et al.* (2008) 'Inhibition of maize root growth by high nitrate supply is correlated with reduced IAA levels
in roots', *Journal of plant physiology*, 165(9), pp. 942–951.

To, J.P.C. *et al.* (2004) 'Type-A Arabidopsis response regulators are partially redundant negative regulators of
cytokinin signaling', *The Plant Cell*, 16(3), pp. 658–671.

Ueno, O. (1998) 'Induction of Kranz anatomy and C4-like biochemical characteristics in a submerged amphibious
plant by abscisic acid', *The Plant Cell*, 10(4), pp. 571–583.

Usadel, B. *et al.* (2006) 'PageMan: an interactive ontology tool to generate, display, and annotate overview graphs
for profiling experiments', *BMC bioinformatics*, 7(1), pp. 1–8.

Vogan, P.J. and Sage, R.F. (2011) 'Water-use efficiency and nitrogen-use efficiency of C3-C4 intermediate
species of Flaveria Juss.(Asteraceae)', *Plant, cell & environment*, 34(9), pp. 1415–1430.

Wang, C.-T. *et al.* (2018) 'The maize WRKY transcription factor ZmWRKY40 confers drought resistance in
transgenic Arabidopsis', *International journal of molecular sciences*, 19(9), p. 2580.

Wang, L. *et al.* (2014) 'Comparative analyses of C4 and C3 photosynthesis in developing leaves of maize and
rice', *Nature biotechnology*, 32(11), pp. 1158–1165.

Wang, P., Vlad, D. and Langdale, J.A. (2016) 'Finding the genes to build C4 rice', *Current Opinion in Plant
Biology*, 31, pp. 44–50.

Xia, C. *et al.* (2018) 'Elucidation of the mechanisms of long-distance mRNA movement in a Nicotiana
benthamiana/tomato heterograft system', *Plant physiology*, 177(2), pp. 745–758.

Yamawaki, S. *et al.* (2011) 'Functional characterization of HY5 homolog genes involved in early light-signaling
in Physcomitrella patens', *Bioscience, biotechnology, and biochemistry*, 75(8), pp. 1533–1539.

Zhang, H. *et al.* (2014) 'A DTX/MATE-type transporter facilitates abscisic acid efflux and modulates ABA
sensitivity and drought tolerance in Arabidopsis', *Molecular plant*, 7(10), pp. 1522–1532.

Zhao, H. *et al.* (2017) 'The Arabidopsis thaliana nuclear factor Y transcription factors', *Frontiers in plant science*,
7, p. 2045.

CAPTION of Table:

Table 1. List of differentially expressed transcription factors between hypocotyl A and hypocotyl B with mobile

mRNAs based on TAIR database (https://www.arabidopsis.org/)

Arabidopsis ID	Transcription factor name	Transcription factor family	Log$_2$FC	FDR
AT3G10500	NAC053	NAC	10/19	7/19E-05
AT1G09770	CDC5	MYB_related	9/78	0/001280562
AT3G02990	HSFA1E	HSF	3/16	2/80E-05
AT1G69490	NAP	NAC	2/43	6/08E-06
AT1G21450	SCL1	GRAS	2/03	7/52E-10
AT1G80840	WRKY40	WRKY	1/32	2/32E-12
AT4G37790	HAT22	HD-ZIP	1/55	8/04E-06
AT5G47390	MYBH	MYB_related	1/52	1/21E-09
AT5G62020	HSFB2A	HSF	-1/84	2/75E-06
AT5G60690	REV	HD-ZIP	-3/12	0/030388143
AT5G59570	BOA	G2-like	-4/08	0/013424766
AT1G01720	ATAF1	NAC	-11/88	2/42E-21

CAPTION of Figures:

Fig. 1 Sampled tissues from *H. mollissima* Bunge in the current study a) Series A b) Series B

Fig. 2 Leaf and cotyledon cross-sections of *H. mollissima* Bunge a) Cross-section of the cotyledon (C$_3$), b) Cross-

section of the entire leaf (C$_4$, Salsoloid type Kranz anatomy), and c) close-up of the leaf Kranz anatomy

Abbreviations: E, Epidermis; PM, palisade mesophyll; KC, Kranz cell; PVB, peripheral vascular bundle, MV,

mid-vein, WS, water-storage tissue; AS, air space; CH, chlorenchyma. Bars = 100 μm (a, b), 25 μm (c)

Fig. 3 PCA analysis which explains 91.09% of the total variation. Principal component analysis of the read counts

separated the samples based on the tissue type and indicated the consistency and validation of our normalized data

Fig. 4 a) Active functional classes of each sample according to all percentages of all transcripts belonged to Mapman categories **b)** Relative transcript abundance of C_4 cycle proteins: most C_4 proteins are more expressed in leaves than in hypocotyls. Beta-carbonic anhydrase (BCA3), PEP carboxylase1 (PEPC1), pyruvate orthophosphate dikinase (PPdK), aspartate aminotransferase (Asp-AT), alanine aminotransferase (Ala-AT), NAD-dependent malic enzyme (NAD-ME), NADP-dependent malic enzyme (NADP-ME), Triosephosphate translocator (TPT), Pyrophosphatase (PPase6), pyruvate orthophosphate dikinase related protein1 (PPdK-RP1), PEP/phosphate translocator (PPT2), Bile acid:sodium symporter family protein (BASS4), Bile acid:sodium symporter family protein (BASS2), Adenosine monophosphate kinase (AMK2), Dicarboxylate translocator1 (Dit1), phosphoenolpyruvate carboxykinase (PEPC-K)

Fig. 5 Overrepresentation analysis of up- and down-regulated genes in pairwise comparison of hypocotyl B vs. hypocotyl A within functional gene classes defined by MapMan bins (MapMan, v3.6, available online at https://www.plabipd.de/portal/mercator4) in Pageman software. Fisher's exact test, followed by the Benjamini Hochberg correction and cut-off value one, was used for functional enrichment of DEGs between three binary comparisons

Blue, up- or downregulated genes are significantly overrepresented; red, up- or downregulated genes are significantly underrepresented

Fig. 6 "Cell signaling" using MapMan. Visualization of transcriptional changes in genes involved in the various signaling pathways (FDR<0.05, log2FC>1.5 <-1.5) at hypocotyl B Vs. hypocotyl A (DEGs encoding phytohormone synthesizing and degrading enzymes, transporter ("Phytohormone action bin" and "Solute transporter"), and different signal transduction pathways ("Multi-process regulation. Calcium-dependent signalling", "Protein modification. Protein kinase" and "Redox hemoestasis" bins))**;** Red, down-regulated gene, and green, up-regulated genes

Fig. 7 Relative transcript abundance of transcripts associated with a) CRP peptide activities b) NCRP peptide activities

Fig. 8 RT-qPCR analysis of selected genes in the tissues of interest a) (CotA, CotB Vs, FL) b) CotB Vs. CotA c) RootB Vs. RootA d) Hypocotyl BVs. Hypocotyl A

Fig. 1

Fig. 2

Fig. 3

a)   **b)**

Fig. 4

758   **Fig. 5**

760　

761　**Fig. 6**

764　**Fig. 7**

Fig. 8