Dendrite branching receptor HPO-30 uses two novel mechanisms to regulate actin cytoskeletal remodeling

Daniel A. Kramer[1], Heidy Y. Narvaez-Ortiz[2], Rebecca Shi[3,4], Kang Shen[3,5], Julien Roche[1], Brad J. Nolen[2], and Baoyu Chen[1#]

Affiliations:

[1]Roy J Carver Department of Biochemistry, Biophysics, and Molecular Biology, Iowa State University, Ames, IA 50014, USA.

[2]Department of Chemistry and Biochemistry, Institute of Molecular Biology, University of Oregon, Eugene, OR 97403, USA.

[3]Department of Biology, Stanford University, Stanford, CA, 94305, USA

[4]Neurosciences IDP, Stanford University, Stanford, CA, 94305, USA

[5]Howard Hughes Medical Institute, Stanford University, Stanford, CA, 94305, USA

[#]Correspondence to: stone@iastate.edu

Abstract

Dendrite morphogenesis is essential to neural circuit formation, but the molecular

mechanisms controlling the growth of complicated dendrite branches are not well understood.

Prior studies using the highly branched *C. elegans* PVD sensory neuron identified a membrane

co-receptor complex that transmits extracellular cues to intracellular actin remodeling machinery

to promote high-order dendrite branching. In this complex, the transmembrane protein HPO-30

recruits the WAVE regulatory complex (WRC) from the cytosol to dendrite branching sites,

where WRC stimulates the Arp2/3 complex to polymerize actin. Here we report biochemical and

structural characterization of this interaction, revealing that the intracellular domain (ICD) of

HPO-30 uses two novel mechanisms to regulate the actin cytoskeleton. First, the unstructured

HPO-30 ICD likely undergoes dimerization and induced folding to bind the WRC, with the

binding simultaneously promoting WRC activation by the GTPase Rac1. Second, the dimerized

HPO-30 ICD directly binds to both the sides and barbed end of actin filaments. The barbed end

binding activity resembles that of the actin capping protein CapZ and prevents both actin

polymerization and depolymerization. The novel dual functions of this dendrite receptor provide

an intriguing model of how membrane proteins can use distinct mechanisms to fine-tune local

actin dynamics.

Introduction

While neurons have vastly different shapes, they all share a similar architecture—one

cell body, a long primary axon, and many branching protuberances called dendrites, along which

are numerous micron-sized projections called spines (Burianek and Soderling, 2013; Jan and Jan,

2003; Lefebvre et al., 2015). The formation of dendrites and spines is critical for the

development of the nervous system, as they form the majority of post-synaptic connections in

animals and directly determine the complex wiring of neural circuits (Hotulainen and

Hoogenraad, 2010; Scott and Luo, 2001; Tavosanis, 2012). Dendrite development requires actin

and microtubule filaments to drive the initiation and outgrowth of newly formed neurites (Cheng

and Poo, 2012; Zhao et al., 2017). Defects in actin cytoskeletal regulation alter dendrite

morphology and neural connections and contribute to many neurodevelopmental disorders, such

as autism, mental retardation, and schizophrenia (Yan et al., 2016).

A central mechanism by which neurons control actin polymerization is through the

Arp2/3 complex, a seven-protein complex that binds to the sides of existing actin filaments and

initiates the formation of branched actin filament networks (Goley and Welch, 2006; Machesky

et al., 1994; Pollard, 2007). Intrinsically inactive, the Arp2/3 complex requires activation by a

group of proteins called nucleation promoting factors (NPFs). The Wiskott-Aldrich Syndrome

Protein (WASP) family proteins are a major group of NPFs (Alekhina et al., 2017; Kramer et al.,

2022; Pollitt and Insall, 2009; Takenawa and Suetsugu, 2007). They all contain a conserved

WCA (WH2-central-acidic) sequence at their C terminus, which can directly bind to and activate

Arp2/3, while their N-terminal sequences vary greatly and define their regulatory mechanism

and cellular function (Alekhina et al., 2017; Kramer et al., 2022; Machesky and Insall, 1998).

Among the WASP-family proteins, WAVE (Wiskott-Aldrich Verprolin homology) exists

in a multi-protein complex named the WAVE regulatory complex (WRC), which contains five

subunits: Sra1, Nap1, Abi2, HSPC300, and WAVE (or their corresponding orthologs in

vertebrates) (Chen et al., 2010; Eden et al., 2002; Gautreau et al., 2004; Polesskaya et al., 2021;

Rottner et al., 2021). Enriched in neurons, the WRC and Arp2/3 complex promote actin

polymerization downstream of membrane signaling to drive various neuronal activities,

including growth cone formation, axon branching, dendrite branching, synapse formation, and

axon guidance and projection (Chia et al., 2014; Chou and Wang, 2016; Pilpel and Segal, 2005;

Racz and Weinberg, 2008; Soderling et al., 2007; Stephan et al., 2011; Tahirovic et al., 2010;

Yamazaki et al., 2005; Yokoyama et al., 2011). Disrupting the WRC in animals profoundly

impacts the nervous system, leading to altered spine morphology and density, intellectual

disability, as well as embryonic death (Dahl et al., 2003; Soderling et al., 2003). Mutations in

WAVE and other subunits of the WRC lead to a variety of neurodevelopmental disorders in

human patients, including neurodevelopmental disorder with absent language and variable

seizures (NEDALVS), developmental and epileptic encephalopathy-65 (DEE-65), and

Alzheimer's disease (Begemann et al., 2021; Conway et al., 2018; Ito et al., 2018; Kirkpatrick et

al., 2017; Kramer et al., 2022; Kumar et al., 2013; Olive et al., 2020; Rottner et al., 2021;

Shimojima Yamamoto et al., 2021; Sims et al., 2017; Srivastava et al., 2021; Zhao et al., 2021;

Zweier et al., 2019).

The WRC keeps WAVE basally inhibited by sequestering its C-terminal WCA to a

conserved surface formed by both the Sra1 and WAVE subunits (Chen et al., 2017, 2010a; Eden

et al., 2002; Kramer et al., 2022) . Through direct interactions, various molecules can recruit the

WRC to the membrane and/or simultaneously activate it, which releases the WCA to promote

Arp2/3-mediated actin polymerization. These ligands include small GTPases (e.g., Rac1 and

Arf), acidic phospholipids (e.g., PIP_3), various adaptor proteins, and over 100 different

membrane proteins that contain a 6 amino acid peptide motif named the WIRS motif (WRC

interacting receptor sequence, defined as Φ-x-T/S-F-x-x, where Φ is a bulky hydrophobic

residue and x is any residues) (Chen et al., 2017, 2014a, 2010a; Eden et al., 2002; Kobayashi et

al., 1998; Koronakis et al., 2011; Lebensohn and Kirschner, 2009; Rottner et al., 2021). Many

WIRS-containing membrane proteins, such as SYG-1, Robo, Neogenin, TMEM132, neuroligins,

and various protocadherins, are important neuronal receptors and have been shown to rely on the

WIRS-WRC interaction to regulate various processes in neural development (Chaudhari et al.,

2021; Chia et al., 2014; Fan et al., 2018; Lee et al., 2016; Wang et al., 2021; Xing et al., 2018).

Previous studies identified the claudin-like dendrite receptor HPO-30 as a novel WRC

binding protein, and this interaction was essential to higher-order dendrite branching. HPO-30

acts as a co-receptor of the cell adhesion molecule DMA-1 in the PVD sensory neuron in *C.

elegans*. The extracellular domain of DMA-1 forms a multi-ligand complex with the secreted

protein LECT-2 and the extracellular domains of the epidermis cell receptors SAX-7 and MNR-1

(Zou et al., 2016). These interactions provide the spatial cue for initiating dendrite branching

(Dong et al., 2013; Zou et al., 2016, 2018). The intracellular domain (ICD) of HPO-30 directly

binds the WRC, while the ICD of DMA-1 binds the Rac guanine nucleotide exchange factor

(GEF) TIAM-1, which can increase the local concentration of activated Rac1—the canonical

activator of WRC (Tang et al., 2019). Together, HPO-30 and DMA-1 organize a multi-protein

complex to spatiotemporally coordinate extracellular cues with the intracellular actin remodeling

essential to high-order dendrite branching (Zou et al., 2018).

It is unclear how the HPO-30 ICD interacts with the WRC. It is also unknown if, in

addition to binding to the WRC, the HPO-30 ICD has other functions. The HPO-30 ICD does not

contain a WIRS motif and must use a non-WIRS mechanism to interact with the WRC. Previous

studies showed that deleting the C-terminal part of the HPO-30 ICD strongly affected WRC

binding *in vitro* and high-order dendrite branching *in vivo*, but it was unclear whether the HPO-

30 ICD used a linear peptide motif analogous to WIRS to bind the WRC (Zou et al., 2018).

Furthermore, although HPO-30 is homologous to the tight junction claudin proteins in mammals,

its ICD sequence is only conserved in nematode worms. Nevertheless, this ICD binds to both *C.*

elegans and human WRC, suggesting that the interaction mechanism is conserved in other

animals (Zou et al., 2018). It is likely that humans have an unidentified membrane protein that

uses the same interaction surface to regulate the WRC. Thus, it is important to understand the

mechanism underlying the interaction between HPO-30 and WRC.

Here we report biochemical and structural analysis of the HPO-30-WRC interaction. We

find that, unlike other WRC-interacting receptors known to date, the HPO-30 ICD requires

dimerization and folding into a three-dimensional structure to bind the WRC. Furthermore, to

our surprise, we find the HPO-30 ICD directly interacts with actin filaments, also in a

dimerization-dependent manner. The dimeric form binds to both the side and barbed end of actin

filaments and inhibits both actin polymerization and depolymerization, resembling the activity of

the actin capping protein CapZ. The dual activities of HPO-30 ICD provide an intriguing

example of how a membrane receptor can regulate actin dynamics by simultaneously controlling

the localization of a central actin nucleation factor and modulating local actin networks to

promote an important biological process.

Results

HPO-30 ICD may use a folded structure, instead of a short peptide motif, to bind the WRC

The HPO-30 ICD (or ICD for short hereafter) does not contain a WIRS motif. Therefore,

we first hypothesized it might use a distinct linear motif to bind the WRC. To identify this

sequence motif, we mutated every five residues to alanine throughout the ICD and used GST

pull-down assays to evaluate how they disrupted WRC binding (**Figure 1A**). Interestingly, we

found that mutating nearly any five amino acids (a.a.), except for the fourth and tenth region,

significantly reduced the binding to WRC, with the middle region of the ICD showing the

strongest effect (Δ5, 6, 7 in **Figure 1A,B**, lanes 7-9). This is consistent with the previous

qualitative data obtained in a less ideal pull-down condition (Zou et al., 2018). It was intriguing

that none of these 5-a.a. mutations completely abolished the binding to WRC, unlike WIRS-

mediated binding, where mutating the WIRS motif readily diminished the interaction (Chen et

al., 2014a). This is consistent with our quantitative measurement of the binding affinity using

equilibrium pull-down (EPD) assays, in which the wild type (WT) ICD had a dissociation

constant (K_D) of ~1.69 μM (**Figure 1D,** black). Alanine mutant Δ5, which had the strongest

effect in the non-equilibrium pull-down assays (**Figure 1A,B,** lane 7), increased the K_D only

mildly to ~5.06 μM, still maintaining significant binding (**Figure 1D**, purple) (Chen et al., 2017;

Kuzmic, 1996; Pollard, 2010). By contrast, replacing either the entire N-terminal or C-terminal

sequences, or two consecutive 5-a.a. mutations with a (GGS) sequence almost completely

abolished the binding (**Figure 1A,C** Δ11-14). Note that while most mutants contain a GST-tag at

the N-terminus, the 5-a.a. mutations located at the C-terminus of the ICD contain a GST-tag at

the C-terminus (Δ8-10 in **Figure 1**, lanes 10-13 in **Figure 1B**), which we found was necessary to

protect the mutant ICDs from degradation (data not shown).

Nearly all of the 5-a.a. alanine scan mutants affected WRC binding. This is distinct from

WIRS-containing proteins, where only mutations in the WIRS motif disrupted binding (Chen et

al., 2014a). We hypothesized that, instead of using a short linear peptide motif like WIRS, either

the HPO-30 ICD folds into a three-dimensional structure and uses the folded structure to bind the

WRC, or the entire sequence binds as a long linear peptide that interacts extensively with the

WRC. We favor the first hypothesis because it is supported by both secondary structure analysis

from various programs, including JPred, PredictProtein, PSIPred, and SABLE, and *ab initio*

tertiary structure prediction by many different algorithms, including LOMETS, QUACK,

RaptorX, Rosetta, AlphaFold 2.0, and RoseTTAFold, (**Figure 1E; Figure 1—figure**

supplement 1) (Adamczak et al., 2004; Bernhofer et al., 2021; Bonneau et al., 2001;

Drozdetskiy et al., 2015; Jumper et al., 2021; Källberg et al., 2012; McGuffin et al., 2000;

Minkyung et al., 2021; Wu and Zhang, 2007). Nearly all predictions suggest the ICD contains a

short alpha helix at the N-terminus followed by two beta strands, which together could adopt

various three-dimensional structures. The predicted structures vary depending on the program

used and the context of ICD sequence (isolated vs. within the full-length HPO-30), suggesting

the ICD structure is likely unstable and can adopt diverse conformations (**Figure 1E; Figure 1—**

figure supplement 1). Together, the above mutagenesis and structural analysis suggest that

HPO-30 ICD likely uses a folded tertiary structure, instead of a linear peptide motif, to bind to

the WRC.

Figure 1. HPO-30 binds to WRC likely using a folded domain, rather than a linear peptide motif. (A)
Left: cartoon representation of HPO-30 as a four-pass transmembrane protein, with the ICD binding to the
WRC. Right: annotation of HPO-30 ICD sequence and mutants used in this study. Each mutant replaces the
corresponding amino acids to alanines (for Δ1-10) or (GGS)n (for Δ11-14). **(B)** Representative Coomassie
blue-stained SDS PAGE gel (left) and quantification (right) of three independent experiments showing GST-
ICD (lane 2-9, 200 pmol) and ICD-GST (lane 10-13, 200 pmol) pulling down WRC$^{230\Delta WCA}$ (150 pmol).
Sra1/Nap1 band intensity was used to quantify the pull-down signals of WRC. Signals from GST-ICD or ICD-
GST pull-downs were normalized to corresponding ICD WT (lane 2 and 13, respectively). Error bars represent
standard error, * $p < 0.05$, ** $p < 0.005$ from Student's paired t-test. **(C)** Coomassie blue-stained SDS PAGE
gel showing GST-ICD Δ11-14 mutants (200 pmol) pulling down WRC$^{217\Delta WCA}$ (300 pmol). **(D)** EPD assay to
measure the dissociation constant (K_D) of the interaction between ICD-GST and WT. Left: data pooled from
three independent repeats were fitted to a one-site binding model using DynaFit. Right: representative
Coomassie blue-stained SDS PAGE gels used for quantification. **(E)** Consensus secondary structure (top) and
tertiary structures of HPO-30 ICD predicted by indicated programs (bottom, same color scheme as the
secondary structure annotation). "FL" indicates structural predictions of ICD in the context of full length (FL)
HPO-30.

Figure 1—figure supplement 1. Top five predicted structures of HPO-30 ICD by the indicated programs.

HPO-30 ICD is disordered in solution

We tried several approaches to determine the structure of the HPO-30 ICD in isolation. The ICD did not grow crystals either by itself or when attached to solubility tags, or as a co-crystal with the WRC. However, due to its small size and our improved protocol to purify the isotope-labeled ICD in high concentration, we were able to use solution nuclear magnetic resonance (NMR) spectroscopy to obtain well-resolved ^{1}H-^{15}N HSQC spectra of untagged ^{15}N-labeled ICD (**Figure 2A**). From the spectra of the untagged ICD, we could readily identify 50 distinct amide crosspeaks out of the 51 we predicted to observe. The narrow ^{1}H chemical shift dispersion suggests that the HPO-30 ICD is predominantly disordered in solution, with little indications of secondary structure formation. To further increase the stability and yield of untagged ICD, we switched to using GB1-ICD (GB1 is a monomeric, small soluble tag commonly used in NMR) (Zhou and Wagner, 2010). Comparison of the 2D ^{1}H-^{15}N TROSY-HSQC spectra collected for GB1-ICD with the isolated ICD (**Figure 2B**, blue**)** and with the isolated GB1 (**Figure 2C**, red**)** showed no significant chemical shift changes. This suggests that HPO-30 ICD remains largely disordered even when it is linked to a well-folded protein like GB1. Furthermore, the NMR spectra were nearly identical at different protein concentrations (50 – 700 µM) and different temperatures (283 K – 298 K), suggesting the ICD alone does not undergo structural changes in a concentration-dependent manner (data not shown).

This result was consistent with the circular dichroism (CD) spectrum of the ICD, in which no major peaks at either the beta-sheet wavelength (positive at 195 nm and negative at 217 nm) or the alpha-helix wavelengths (positive at 193 nm and negative at 218 and 222 nm) were observed, in contrast to the spectrum of bovine serum albumin (BSA) obtained in the same conditions, which shows clear peaks associated with both structural elements (**Figure 2E**). Addition of an osmolyte, trimethylamine N-oxide (TMAO), which is commonly used to promote protein folding, did not cause a significant change to the CD spectrum or enhance the ICD-WRC interaction (**Figure 2—figure supplement 1**) (Baskakov et al., 1999). Together, these data suggest the purified ICD is unstructured in solution.

Figure 2. HPO-30 ICD alone is unstructured. (A-D) 2D TROSY spectra of untagged HPO-30 ICD (A), GB1-ICD (B), GB1 (C) and GST-ICD (D) in the same buffer condition (100 mM NaCl, 10 mM HEPES pH 7.0, and 5% glycerol). (E) CD spectrum of 2 mg/mL untagged HPO-30 ICD or BSA in 50KMEH5Gd (see Methods and Materials).

Figure 2—figure supplement 1. (A) CD spectrum of 2 mg/mL HPO-30 ICD in the absence or presence of 0.1 M TMAO. (B-C) Coomassie blue-stained SDS PAGE gels showing GST-ICD (200 pmol) pulling down WRC$^{230\Delta WCA}$ (150 pmol) (B) or WRC-(MBP)$_2$ (60 pmol) pulling down GST-ICD (600 pmol) (C) with and without TMAO.

Dimerization is required for HPO-30 ICD to bind WRC

How can the unstructured HPO-30 ICD use a folded structure to bind the WRC? While attempting to solve this conundrum, we realized that the GST-tagged constructs used in our pull-

down assays would dimerize the ICD, as GST is a constitutive dimer (Askelöf et al., 1975). We

then hypothesized that the HPO-30 ICD may need to be dimerized for efficient binding to the

WRC. To test this hypothesis, we repeated the pull-down assay after switching the GST tag to

the monomeric tag MBP (maltose binding protein) (**Figure 3A**). Consistent with our hypothesis,

the MBP-tagged ICD indeed showed much weaker binding to the WRC (**Figure 3B**, right).

Further supporting the observation that monomeric ICD does not bind to the WRC effectively,

GST-ICD binding could not be competed off by a chemically synthesized peptide covering the

whole HPO-30 ICD, even when the peptide was added in 5000-fold excess (**Figure 3—figure**

supplement 1A).

We next tested several different strategies to further validate if dimerization of the ICD is

required for WRC binding. First, we immobilized the WRC to the affinity beads by tethering a

dual MBP tag, $(MBP)_2$, to the WRC to pull down different monomeric vs. dimeric forms of

HPO-30 ICD. To our surprise, the WRC containing an $(MBP)_2$ tag at the N-terminus of

HSPC300, which was previously used to immobilize the WRC and pull down WIRS-containing

proteins (Chen et al., 2014a), substantially reduced HPO-30 ICD binding (**Figure 3—figure**

supplement 1C, lane 6 vs. 8). Similarly, an $(MBP)_2$ tag at the N-terminus of Abi2 also prevented

HPO-30 ICD binding (**Figure 3—figure supplement 1B**). These data suggest tagging $(MBP)_2$ at

the N-terminus of HSPC300 or Abi2 may interfere with the HPO-30 ICD binding surface (but

not the WIRS binding surface). We eventually used sortase-mediated protein ligation (Chen et

al., 2011) to tether an $(MBP)_2$ tag to the C-terminus of Abi2, which produced a WRC that

efficiently bound to amylose beads and effectively retained HPO-30 ICD binding **(Figure 3—**

figure supplement 1C). Using this construct, we observed robust binding of all dimeric

constructs we tested, including GST-ICD, ICD-GST, and DLC8-ICD (DLC8 is a constitutive

dimer from the dynein light chain) (Wang et al., 2003). By contrast, all monomeric constructs,

including GB1-ICD, sumo-ICD, and mEGFP-ICD showed very weak, if any, binding (**Figure**

3A,C).

Figure 3. HPO-30 ICD binding to the WRC is enhanced by dimerization. (A) Schematic of HPO-30 ICD

constructs used in this study. **(B)** Coomassie blue-stained SDS PAGE gels showing GST-ICD or MBP-ICD (200

pmol) pulling down WRC$^{230\Delta WCA}$ (200 pmol). **(C)** Coomassie blue-stained SDS PAGE gels showing WRC-(MBP)$_2$

or (MBP)$_2$-WRC (60 pmol) pulling down various monomeric and dimeric HPO-30 ICD (600 pmol) indicated in (A).

Where it is noted, 5 µM rapamycin (vs. equal volume of DMSO as a negative control) was used to induced

dimerization. White arrowheads indicate pull-down signals. For clarity, GB1-FKBP-ICD and GB1-FRB-ICD were

referred to as FKBP-ICD and FRB-ICD, respectively. **(D)** Coomassie blue-stained SDS PAGE gels showing WRC-

(MBP)$_2$ (60 pmol) pulling down different dimeric HPO-30 ICD (600 pmol) in the presence of GST-ICD as a

competitor (600 to 6000 pmol). **(E)** Schematic of the assembly of homo- and heterodimers of HPO-30 ICD (left) and

Coomassie blue-stained SDS PAGE gels (right) showing WRC-(MBP)$_2$ (60 pmol) pulling down indicated homo-

and heterodimers. In homo- and heterodimers, 1200 pmol of ICD monomeric unit was included to ensure the total

amount of ICD remained the same. **(F)** Schematic of full-length HPO-30 constructs co-expressed in *Drosophila* S2

cells (left) and Western blot images (right) showing co-immunoprecipitation of 3xHA-tagged HPO-30 (pink

arrowhead) with 6xMyc-tagged HPO-30 (orange arrowhead). Asterisks indicate light and heavy chain of antibody.

Figure 3—figure supplement 1. Additional data supporting HPO-30 ICD binding to WRC is enhanced by

dimerization. (A) Coomassie blue-stained SDS PAGE gel showing GST-ICD (200 pmol) pulling down

WRC$^{230\Delta WCA}$ (150 pmol) in the presence of a chemically synthesized HPO-30 ICD peptide as a competitor (18-100

nmol). **(B)** Schematic of removing (MBP)$_2$ tag from WRC$^{(MBP)2-Abi2}$ by thrombin cleavage (left) and Coomassie blue-

stained SDS PAGE gels (right) comparing GST-ICD (200 pmol) pulling down WRC$^{(MBP)2-Abi2}$ (150 pmol) before or

after removal of the (MBP)$_2$ tag. **(C)** Schematic of (MBP)$_2$-tagged WRC constructs (left) and Coomassie blue-

stained SDS PAGE gels showing indicated (MBP)$_2$-tagged WRC (60 pmol) pulling down GST-ICD (600 pmol).

White arrowheads indicated pull-down signals. **(D)** Top: gel filtration chromatograms of equimolar GB1-FKBP-ICD

and GB1-FRB-ICD in the presence or absence of 5 µM rapamycin (or equal volume of DMSO). Bottom: Coomassie

blue-stained SDS-PAGE gels of the indicated gel filtration fractions. **(E)** Coomassie blue-stained SDS PAGE gel

showing WRC-(MBP)₂ (60 pmol) pulling down ICD-GST (600 pmol) in the presence of FKBP/FRB-ICD as a

competitor (600 – 6000 pmol). **(F)** Coomassie blue-stained SDS PAGE gels showing WRC-(MBP)₂ (60 pmol)

pulling down different combinations of homo- and heterodimers of WT and Δ5 HPO-30 ICD (in which WT ICD

monomeric unit was kept at 1200 pmol to ensure the same amount of the WT ICD was included in the reactions).

Figure 3—figure supplement 2. HPO-30 can potentially form a dimer. (A) AlphaFold Multimer predictions of

the isolated HPO-30 ICD dimer. Promoter A is colored using the same scheme shown on the right (and in Figure

1E), while protomer B is shown in lighter colors. **(B)** Two distinct dimer conformations of the full-length HPO-30

predicted by AlphaFold Multimer. **(C)** Coomassie blue-stained SDS PAGE gels showing GST-ICD (200 pmol)

pulling down MBP-ICD (6000 pmol) in indicated buffer conditions.

To further exclude the possibility that the affinity tags nonspecifically affected protein

interaction, we introduced the chemically inducible dimerization (CID) tags FKBP and FRB to

the ICD, with the GB1 tag at the N-terminus to improve protein expression and solubility

(Banaszynski et al., 2005) (**Figure 3A**). In the presence of the dimerizing agent rapamycin, GB1-

FKBP-ICD and GB1-FRB-ICD can readily form a tight heterodimer (**Figure 3A**), as confirmed

by size exclusion chromatography in the presence and absence of rapamycin (**Figure 3—figure**

supplement 1D). Consistent with results from the above constitutive dimeric tags, the addition

of rapamycin to dimerize GB1-FKBP/FRB-ICD (F/F-ICD for short hereafter) clearly promoted

HPO-30 ICD binding (**Figure 3C**, lanes 8 vs. 9).

To ensure that the ICDs that contain different dimerization tags bind to the WRC using

the same mechanism, we used competition pull-down assays to compare different dimeric

constructs. We found GST-ICD effectively competed off both F/F-ICD and DLC8-ICD binding

to the WRC in a dose-dependent manner (**Figure 3D**, lanes 4-6 and 8-10). Complementarily,

F/F-ICD effectively competed off GST-ICD binding to WRC only in the presence of rapamycin,

while in the absence of rapamycin the competition efficiency was significantly reduced (**Figure**

3—figure supplement 1E, lanes 4 and 5). These data further corroborate the notion that

dimerized ICD promotes WRC binding, and that all dimeric constructs tested so far bind WRC

using the same mechanism.

As an alternative way to validate that dimerization of the HPO-30 ICD is required for

WRC binding, we mixed different GB1-FKBP/FRB constructs to produce various heterodimers

of the ICD. We found that a heterodimer containing either only one ICD (by mixing with empty

GB1-FKBP or FRB tag) or having one WT ICD and an alanine Δ5 mutant ICD failed to support

binding to the WRC to the same level as a homodimer containing two WT ICDs (**Figure 3E**;

Figure 3—figure supplement 1F). Note that in this experiment we kept the total concentration

of the WT ICD monomeric unit the same between heterodimers and homodimer. Therefore, the

lack of binding from heterodimers was due to lack of the WT ICD dimer, but not reduced WT

ICD concentration.

If the ICD needs to be dimerized in order to bind the WRC, we speculated HPO-30

should be able to form dimers in the cell. To test this hypothesis, we expressed 3xHA-tagged and

6xMyc-tagged full-length HPO-30 in *Drosophila* S2 cells and used co-immunoprecipitation to

test if HPO-30 interacted with itself (**Figure 3F)**. We found 3xHA-tagged HPO-30 robustly

retained 6xMyc-tagged HPO-30, suggesting that full-length HPO-30 is capable of dimerization

(or oligomerization) in cells, which could be the functional form of HPO-30. With recent

advances in protein structural prediction by AlphaFold, we used AlphaFold Multimer (AFM) to

predict whether and how the isolated ICD or the full-length HPO-30 can form a dimer (**Figure**

3—figure supplement 2A,B) (Evans et al., 2022; Jumper et al., 2021). We found the ICD alone

was consistently predicted to form a homodimer. Out of 25 prediction solutions, 20 solutions

show the same conformation in which dimerization is mediated by a beta sheet formed by the

two beta strands from each ICD (**Figure 3—figure supplement 2A**). Consistent with this

prediction, purified GST-ICD was able to pull down MBP-tagged ICD in a pH and salt

dependent manner, which suggests the dimerization of ICD is mediated through polar

interactions (**Figure 3—figure supplement 2C**). The full-length HPO-30 was also predicted to

form dimers, but with two distinct conformations. In the first conformation, dimerization was

mediated by the interactions between the transmembrane helix 2 (H2) and the beta strand 4 (β4)

in the extracellular domain, while in the second conformation, dimerization was mediated by

transmembrane helix 3 (H3) and the first helix of the ICD (**Figure 3—figure supplement 2B**).

In both conformations, the ICD was predicted to form a long helix, without two beta strands.

This contradicts the secondary structural predictions and tertiary structural predictions of the

isolated ICD by all other methods, likely because the global prediction of the full-length HPO-30

somehow influenced the prediction of the ICD due to the dominant structures of the N-terminal

regions (**Figure 1E**; **Figure 1—figure supplement 1**; **Figure 3—figure supplement 2A**).

Taken together, our data suggest that HPO-30 ICD has the potential to dimerize both *in*

vitro and in cells, and the dimerization is required for efficient binding to the WRC.

Dimerization is not sufficient to induce HPO-30 ICD folding

As the dimeric ICD binds to the WRC more strongly than the monomeric form, we used

NMR to determine if dimerization could induce structure changes in the HPO-30 ICD. Among

various dimerized ICD constructs, only GST-ICD could produce protein at a high enough

concentration for NMR measurement. With this material, we managed to obtain well-resolved

^{1}H-^{15}N TROSY-HSQC spectra, which showed ^{1}H chemical shift dispersion nearly identical to

the untagged ICD, indicating that dimerization alone was not enough to induce significant ICD

folding (**Figure 2D**). Note that we could not see any spectral peaks for the GST tag in this

condition due to its large size (~54 kDa as a constitutive dimer) and consequent slow tumbling.

Collecting the above evidence, we posit that the HPO-30 ICD needs to undergo both

dimerization and induced folding in order to bind to the WRC. This induced folding has been

proposed for many DNA-binding proteins and cell-signaling molecules, such as the interaction

between E-cadherin and ß-catenin (Dyson and Wright, 2002; Huber et al., 2001; Huber and

Weis, 2001; Turjanski et al., 2008). High-resolution structures of the ICD bound to WRC will be

necessary to confirm this hypothesis.

HPO-30 ICD inhibits actin polymerization, but promotes Rac1-WRC activation

367   Previous studies have shown that Rac1 is the canonical activator of the WRC, and that

various WIRS-containing receptors could further fine tune WRC activity, likely through

secondary, weak interactions between the WRC and the receptor sequences flanking the WIRS

motif (Chen et al., 2014a). After establishing that dimeric ICD robustly binds the WRC, we

tested if the interaction could similarly influence WRC activity in promoting Arp2/3-mediated

actin polymerization (Cooper et al., 1983; Doolittle et al., 2013a; Kouyama and Mihashi, 1981).

While conducting the pyrene-actin polymerization assay, we noticed that buffer conditions had a

significant impact on the ICD-WRC interaction. The commonly used pyrene-actin assay buffer

for reactions involving WRC, 50KMEI20Gd (50 mM KCl, 2 mM $MgCl_2$, 1 mM EGTA pH 8.0,

10 mM Imidazole pH 7.0, 20% [w/v] glycerol, 1 mM DTT), reduced HPO-30 ICD binding to the

WRC. Lowering the concentration of glycerol to 5% and replacing imidazole with HEPES at the

same pH rescued binding (**Figure 4—figure supplement 1A**). Using this optimized buffer

(50KMEH5Gd, containing 50 mM KCl, 2 mM $MgCl_2$, 1 mM EGTA pH 8.0, 10 mM HEPES pH

7.0, 5% [w/v] glycerol, 1 mM DTT), we tested the effect of monomeric and dimeric HPO-30

ICD on actin polymerization using our chemically inducible dimerization (CID) constructs

(shown in **Figure 3A)**, which allowed us to compare the monomeric vs. dimeric ICD directly by

switching between DMSO and rapamycin instead of using different solubility tags (**Figure 3—**

supplement figure 1D). To our surprise, we observed a strong, dose-dependent inhibition of

Rac1-WRC-mediated actin polymerization by HPO-30 ICD (**Figure 4A**, blue vs. orange curves),

with the dimerized ICD showing much stronger inhibition (**Figure 4A**, solid vs. dashed curves).

By contrast, the dimerized FKBP/FRB tag at the highest concentration only had a mild effect,

indicating the inhibition was due to the HPO-30 ICD itself (**Figure 4A**, grey curve). Rapamycin

alone had no effect on actin polymerization (**Figure 4—figure supplement 1B**).

390   Interestingly, we noticed the ICD reduced actin polymerization to a level lower than actin

alone. This suggests the inhibitory effect was likely not because of the ICD binding to the WRC

and thereby inhibiting WRC activity. Indeed, we found HPO-30 ICD similarly inhibited actin

polymerization induced by the isolated, constitutively active WCA peptide, with the dimerized

ICD again showing more potent inhibition (**Figure 4B**). This suggests the inhibitory effect of

ICD was not related to WRC, but likely directly related to actin (see below).

Figure 4. HPO-30 ICD inhibits actin polymerization but promotes Rac1-WRC activation. (A-E) Pyrene-actin

polymerization assays of indicated conditions. Reactions contain 2 μM actin (10% pyrene-labeled), 10 nM Arp2/3

complex, 100 nM WRC^{230WCA} or isolated WCA, Rac1^{QP}, indicated concentrations of equimolar GB1-FKBP-ICD

and GB1-FRB-ICD, and 5 μM rapamycin or equal volume of DMSO. **(F)** Coomassie blue-stained SDS PAGE gels

(left) and quantification from three independent repeats (right) showing equilibrium pull-down of WRC^{230WCA} by

two concentrations of ICD-GST in the presence or absence of 40 μM Rac1^{QP}. Error bars represent standard error, *

$p < 0.05$, from Student's paired t-test.

Figure 4—figure supplement 1. Buffer optimization of HPO-30 ICD pyrene-actin polymerization assay. (A)
Coomassie blue-stained SDS PAGE gels showing GST-ICD (200 pmol) pulling down WRC$^{230\Delta WCA}$ (150 pmol) in
indicated buffer conditions, with 50PDB being the buffer used in the majority of pull-down reactions herein and
50KMEI20G commonly used in pyrene-actin assembly assays in previous studies of the WRC. All buffers contained
1 mM DTT and 0.05% Triton X-100. (B) Pyrene-actin polymerization assays showing rapamycin and DMSO did
not affect WRC activity. Reactions contain 4 µM actin (5% pyrene-labeled), 10 nM Arp2/3 complex, 100 nM
WRC^{230WCA}, Rac1^{QP}, and 5 µM rapamycin or equal volume of DMSO.

       The strong, WRC-independent inhibitory effect of the ICD on actin polymerization
prevented us from directly examining how ICD binding influenced WRC activity. To overcome
this challenge, we reduced the ICD concentration to 1 µM in the actin assays to minimize its
inhibitory effect. Interestingly, we found that 1 µM monomeric ICD slightly, but robustly,
increased the WRC-mediated polymerization when WRC was activated to a low level by an
intermediate concentration of Rac1 (**Figure 4C**, dashed blue vs. solid pink curves). At the same
concentration, the dimerized ICD still showed a dominant inhibitory effect (**Figure 4C**, solid
blue curve). We hypothesized that in the absence of rapamycin, a small amount of HPO-30 ICD
existed in equilibrium as a dimer, which could bind the WRC to promote activation. This is
supported by both structural predictions and *in vitro* pulldown results (**Figure 3—figure
supplement 2A,C**). To test this hypothesis, we further reduced HPO-30 ICD to a much lower

concentration (0.1 µM) and included rapamycin to ensure dimer formation. At this low

concentration, the dimerized ICD similarly promoted Rac1-WRC-medicated actin

polymerization (**Figure 4C**, solid green curve). Although we were limited by the strong

inhibitory activity of the ICD on actin polymerization and therefore could only measure a mild

activating effect by using low ICD concentrations, this effect was specific to Rac1-activated

WRC, as neither the WRC in the absence of Rac1 nor the isolated WCA could be further

activated by ICD (**Figure 4 D,E**). This effect is similar to the WIRS-containing ICD from the

cell adhesion membrane protein protocadherin 10 (PCDH10), in which PCDH10 ICD by itself

had no effect on WRC activity, but cooperatively increased WRC activity when WRC was

slightly activated by intermediate concentrations of Rac1 (Chen et al., 2014a). This cooperativity

effect suggests the HPO-30 ICD should prefer to bind the WRC activated by Rac1. Consistent

with this hypothesis, our equilibrium pull-down (EPD) assay showed GST-ICD could indeed

pull down more WRC in the presence of saturating concentrations of Rac1 (**Figure 4F**, lane 2 vs.

4—note that in EPD assays, the supernatant of the pull-down reactions was used to quantify the

WRC that was not retained by immobilized GST-bait). Interestingly, within the HPO-30-DMA-1

co-receptor complex, DMA-1 directly binds to TIAM-1, a Rac GEF, which could act

synergistically with the direct effect of HPO-30 to promote WRC activation by increasing Rac1

activity (Zou et al., 2018).

HPO-30 ICD binds to F-actin and inhibits actin depolymerization, similar to CapZ

We first wondered if the WRC-independent inhibitory effect of HPO-30 ICD was due to

the ICD binding to and sequestering free G-actin. However, we observed no detectable binding

between GST-ICD or dimerized F/F-ICD and G-actin (**Figure 5—figure supplement 1**). We

next wondered if the ICD inhibited actin polymerization by binding to F-actin. In our F-actin co-

pelleting assay (Heier et al., 2017), we found that significantly more F/F-ICD indeed bound to F-

actin, compared to BSA, F/F tag, and the capping protein CapZ (which binds to the barbed ends

of F-actin) (**Figure 5A,B**). Interestingly, both the monomeric and dimeric F/F-ICD showed

significant and similar binding to F-actin (**Figure 5A,B**, +DMSO vs. +Rap), although the

dimeric ICD showed slightly stronger binding. This suggests that the monomeric ICD, while not

as effective in inhibiting actin polymerization as the dimeric form, is also capable of binding to

actin filaments. Together, these data led us to a hypothesis that HPO-30 ICD may bind to actin

filaments to prevent actin polymerization.

Figure 5. HPO-30 ICD binds to F-actin and inhibits actin depolymerization. (A) Representative Coomassie

blue-stained SDS PAGE gels showing F-actin co-pelleting assays of indicated F/F-ICD and CapZ in the presence or

absence of 5 µM rapamycin (or equal volume of DMSO). S: supernatant, P: pellet. **(B)** Quantification of (A) from

three independent repeats, showing percentage of proteins in the pellet. Bars represent standard error, *** $p < 0.001$,

ANOVA with Tukey test. **(C)** Representative F-actin depolymerization assay fluorescence curves of indicated

conditions. Each reaction contained 5 µM pre-polymerized actin (70% pyrene labeled), diluted 20-fold into

depolymerization buffer containing indicated proteins in the presence or absence of 5 µM rapamycin (or equal

volume of DMSO). **(D)** Quantification of the initial velocity of fluorescence curves shown in (C). Initial velocity

was normalized to F-actin alone in the presence of DMSO. Error bars represent standard error, n = 3 independent

repeats, *** $p < 0.001$, ANOVA with Tukey test.

Figure 5—figure supplement 1. HPO-30 ICD does not bind to G-actin. (A-B) Coomassie blue-stained SDS

PAGE gels showing GST-ICD (200 pmol) (A) or His6-F/F-ICD (B) (300 pmol) pulling down actin or pyrene

labeled actin (500 pmol) in G buffer (2 mM Tris-HCl pH 8, 0.5 mM DTT, 0.2 mM ATP, 0.1 mM $CaCl_2$, 1 mM

NaN_3), in the presence of 5 μM rapamycin (or equal volume of DMSO).

One group of proteins well-known to inhibit actin polymerization are called capping

proteins (e.g., CapZ), which bind to the barbed ends of actin filaments to inhibit both actin

polymerization and depolymerization from the barbed end (Caldwell et al., 1989; Weeds and

Maciver, 1993). To test if the HPO-30 ICD may act as a capping protein, we used a

depolymerization assay that has been commonly used to measure CapZ activity (Caldwell et al.,

1989; Cooper and Pollard, 1985). We found that, similar to CapZ, HPO-30 ICD inhibited actin

depolymerization (**Figure 5C,D**). Note that this activity was both concentration- and

dimerization-dependent, similar to the ICD binding to WRC, suggesting dimerization is also

involved in this capping-like activity. The ability to cap filaments also explains why dimeric

HPO-30 ICD is capable of inhibiting actin polymerization more strongly than monomeric ICD.

HPO-30 ICD binds both the side and barbed end of actin filaments

In the above bulk solution assays, the dimerization-independent binding to F-actin

(**Figure 5A,B**) versus the dimerization-dependent inhibition of actin polymerization and

depolymerization (**Figure 4A,B, 5C,D**) suggest HPO-30 ICD may have two distinct activities on

actin filaments, with only the dimerization-dependent activity resembling that of CapZ. This is

also supported by the observation that F-actin pelleted a more significant amount of HPO-30

ICD than CapZ, as CapZ only binds to the barbed end of F-actin (albeit with high affinity)

(**Figure 5A,B**) (Caldwell et al., 1989). Therefore, the ICD may bind to not only the barbed end

(in a dimerization-dependent manner), but also other locations of the actin filament (in a

dimerization-independent manner).

To directly validate these F-actin binding activities, we used single-molecule total

internal reflection fluorescence (smTIRF) microscopy to visualize how HPO-30 ICD binds to

actin filaments. For this assay, we prepared fluorophore-labeled proteins by fusing a SNAP tag to

the N-terminus of the CapZ β-subunit, FKBP-ICD, FRB-ICD, FKBP tag, and FRB tag, which

allowed us to label proteins using different SNAP-Surface® AlexaFluor® dyes (hereafter

referred to as AF followed by the corresponding excitation wavelength) through the SNAP tag

and minimize the effect of fluorophore-labeling on protein activity. Actin was labeled with

AF647 using previously established methods (Hansen et al., 2013). The SNAP-tagged F/F-ICDs

were capable of binding the WRC in a rapamycin-dependent manner and to a level similar to the

GB1-tagged F/F-ICD, suggesting that the SNAP-tag did not affect ICD binding to WRC (**Figure**

6—figure supplement 1A, lane 1-4). The labeled CapZ and F/F-ICD proteins exhibited capping

activity similar to their counterparts used in bulk solution assays (except AF647-labeled F/F-

ICD, which we decided to exclude from smTIRF assays) (**Figure 6—figure supplement 1B,C**).

The smTIRF experiments clearly showed SNAP^{AF488}-CapZ (at 5 nM) bound to the

barbed end of filaments and stopped their growth (**Figure 6A** top; **6C** green; **Video 1,2**),

consistent with the high affinity and very slow off-rate of CapZ at the barbed end shown in

previous studies (Caldwell et al., 1989). Distinct from CapZ, SNAP^{AF546}-F/F-ICD (15 nM, with

8-fold molar excess of rapamycin) clearly revealed HPO-30 ICD mainly bound to the side of

actin filaments, instead of the barbed end (**Figure 6A** bottom; **6B**; **Video 3-4**) (see below).

Figure 6A bottom shows an example of binding of HPO-30 ICD near the barbed end of the

filament (time 36.5 s), after which the filament continued to grow while the ICD molecule stayed

bound to the same position. Note that this HPO-30 ICD molecule was bound to the filament, not

the PEG-coated surface, because it moved together with the filament (Video 3,4). We applied

stringent criteria to prevent misidentification of nonspecific background signals as binding

events (see Materials and Methods) and found that 33% of filaments had at least one binding

event over the course of the 15-minute duration of the experiment, which was significantly

higher than the number of events for the F/F tag in identical conditions (**Figure 6B**).

Figure 6. HPO-30 ICD binds both the side and barbed end of actin filaments. **(A)** Examples of time lapse images from smTIRF experiments. Top: a capping event of SNAP[AF488]-CapZ (green, 5 nM) and actin[AF647] (red). Bottom: a side-binding event of SNAP[AF546]-F/F-ICD (cyan, 15 nM) and actin[AF647] (red). Note that in smTIRF time-lapses, sometimes a lag was observed between signals at the two channels. For example, in the top row, the filament and SNAP[AF488]-CapZ puncta were displaced at 69.5 s, but aligned again at 70.5 s. The lagging was due to image acquisition conditions, where, to prevent photobleaching of AF647, images in the 640 nm-channel for actin[AF647] were taken every 20 acquisitions of the 488 nm-channel for SNAP[AF488]-CapZ (see Materials and Methods) **(B)** Quantification of the frequency of side binding events from smTIRF imaging. Left: comparison of overall side binding events. Right: comparison of the percentage of filaments with 0, 1, and 2 events. Data were from three independent repeats, at least 15 filaments randomly selected per video. Bars represent standard error. * $p < 0.05$, ** $p < 0.01$, Student's paired t-test. **(C)** Violin plot showing quantification of elongation rate of actin filaments measured by smTIRF imaging. Total number of filaments pooled from the number of independent repeats for each sample are indicated in the paratheses on top of each column, respectively. *** $p < 0.001$, ANOVA with Tukey test. **(D)** Representative Coomassie blue-stained SDS PAGE gel (left) and quantification of pelleted CapZ from three independent repeats (right) showing F-actin (2 μM) co-pelleting assay of CapZ (0.6 μM) with the indicated F/F-ICD or F/F tag as a competitor (5 μM) in the presence or absence of 5 μM rapamycin (or equivalent volume DMSO). S: supernatant, P: pellet. Bars represent standard error, ** $p < 0.01$, *** $p < 0.001$, ANOVA with Tukey test.

Figure 6—figure supplement 1. SNAP tag and fluorophore-labeling did not affect HPO-30 ICD function.
(A) Coomassie blue-stained SDS PAGE gel showing WRC-(MBP)$_2$ (60 pmol) pulling down GB1-tagged vs.
SNAP-tagged HPO-30 ICD (600 pmol) in the presence or absence of 5 µM rapamycin. **(B)** SDS PAGE gels of
SNAP-tag labeling of CapZ, FKBP-ICD, and FKBP tag with indicated SNAP-Surface© AlexaFluor© dyes.
Left: UV light with Alexa-488 filter. Right: Coomassie blue stain. **(C)** Actin depolymerization assays of
SNAP-Alexa labeled proteins in conditions identical to Figure 5C,D.

Since we could not further increase the concentration of fluorophore-labeled ICD in
smTIRF experiments to promote the chance of observing barbed end binding events without
causing high background, we switched to using unlabeled ICD at a higher concentration to
examine if the ICD inhibited actin filament growth as we observed in bulk solution assays in
Figure 5C,D (Figure 6C; **Video 5)**. We found that at 1 µM concentration, dimerized F/F-ICD
significantly reduced actin elongation rate from 2.3 ± 0.5 subunits/sec for actin alone to 1.5 ± 0.3
subunits/sec (Kuhn and Pollard, 2005) (**Figure 6C**, black vs. solid blue). The same concentration
of monomeric F/F-ICD also significantly slowed down actin elongation, but to a lesser extent
$(2.0 \pm 0.4$ subunits/sec) (**Figure 6C**, open blue). This effect may result from the side-binding

activity of ICD slowing down actin polymerization or, more likely, a small population of dimeric

ICD in equilibrium with the monomeric ICD, which could bind to the barbed end (see below).

Importantly, dimerized F/F tag barely affected actin polymerization (2.1 ± 0.4 subunits/sec)

(**Figure 6C**, grey). We were limited by sample addition to the microfluid chamber from high-

concentration protein stocks and, as such, we could not further increase the unlabeled ICD

concentration to higher than 1 µM in order to observe an even stronger inhibitory effect. By

comparison, 5 nM CapZ strongly inhibited actin polymerization (0.3 ± 0.1 subunits/sec).

Together, these data suggest that HPO-30 ICD and CapZ similarly inhibit actin filament

elongation, although the activity of HPO-30 ICD is less potent.

In the above smTIRF experiments, we were unable to capture stable binding of the ICD

to the barbed end as we were for CapZ, likely because of multiple technical challenges: 1) the

low concentration of ICD that we had to use in single molecule assays (nM vs. µM that we used

in bulk solution assays), 2) possibly low affinity and fast off-rate of the ICD at the barbed end

(reflected by the higher concentration than CapZ that was required for inhibiting actin

depolymerization in bulk solution assays), 3) the complications of the side-binding activity

(which, due to light diffraction limit, made it difficult to distinguish real barbed end binding

events from side binding near the barbed end), 4) the amount of dimeric HPO-30 is likely to be

very low, as the affinity of the FKBP-rapamycin complex for FRB (which is responsible for

dimerization of HPO-30 in this system) is ~12 nM (Banaszynski et al., 2005), and 5) the speed

limitation of data acquisition (50 or 100 ms exposure time, which could miss fast

binding/dissociation events). In order to determine if the ICD can indeed bind to the barbed end,

we used a competition co-pelleting assay to examine if ICD binding can compete off CapZ

binding to the barbed end of F-actin (**Figure 6D**). In this assay, due to the high affinity of CapZ

to the barbed end, we used 10 times the concentration of F/F-ICD to compete against CapZ. We

found that the dimerized F/F-ICD effectively reduced CapZ binding by over 60%, whereas the

monomeric F/F-ICD or the dimerized F/F tag had no effect (**Figure 6D**). This result suggests

dimeric, but not monomeric ICD binds to the barbed end, which can block CapZ binding,

consistent with the result that dimeric ICD is more effective in inhibiting actin polymerization

(**Figure 4A,B**) and depolymerization (**Figure 5C,D**). Alternatively, dimeric HPO-30 ICD could

bind near, instead of directly to, the barbed end, which could allosterically destabilize CapZ

binding by altering the conformation of the actin filament. High-resolution structural information

will be necessary to distinguish between these two scenarios.

593   Combining the above results, we conclude that HPO-30 ICD monomers preferentially

594 bind to the side of actin filaments, while the dimers can bind to both the sides and the barbed end

595 (or near the barbed end) of filaments. Only the (near-) barbed end binding by dimers can inhibit

596 actin filament growth in a manner similar to the capping protein CapZ.

598 **Discussion**

599   HPO-30 is a claudin-like membrane protein required for higher order dendrite branching

in *C. elegans* PVD neurons. Previous studies showed this function required the HPO-30 ICD to

bind to the WRC, which in turn could activate Arp2/3 to generate branched actin networks to

deform the membrane and promote new dendrite formation (Zou et al., 2018). Here, our

biochemical, biophysical, structural, and single molecule analysis uncovers two distinct, novel

mechanisms used by the HPO-30 ICD to regulate local actin dynamics. In the first mechanism,

the unstructured ICD undergoes dimerization and induced folding to bind the WRC, which, in

addition to recruiting the WRC to local dendrite branching sites, can also cooperate with Rac1 to

promote WRC activation (**Figure 7**, step 3). In parallel, the co-receptor of HPO-30, DMA-1, was

shown to bind the Rac-GEF, TIAM1, which would provide an additional layer of control to

promote Rac1 activation and drive WRC-Arp2/3-mediated actin polymerization (Zou et al.,

2018). In the second mechanism, the ICD can directly bind to actin filaments to modulate actin

dynamics. Both monomeric and dimeric forms of the ICD can bind to the side of actin filaments,

but only the dimeric ICD can bind to the barbed end to inhibit both actin polymerization and

depolymerization, similar to the activity of the capping protein CapZ (**Figure 7**, step 1). These

two seemingly contradictory actin regulatory mechanisms offered by the same membrane

receptor could provide exquisite tuning of local actin dynamics to prepare for dendrite branching

(**Figure 7**, and see discussion below).

Figure 7. Model showing HPO-30 ICD integrates two distinct mechanisms to regulate actin dynamics and promote dendrite branching. In (1), HPO-30 on the dendrite membrane undergoes dimerization (or oligomerization). Both monomeric and dimeric forms of HPO-30 ICD can bind to the side of actin filaments, but only the dimeric ICD can bind to the fast-growing barbed end. In (2), dimeric ICD binding to the barbed end acts as a capping protein, which stops long filament growth and helps reserve G-actin for the Arp2/3 complex to nucleate actin for producing more short, branched actin filaments. In (3), the dimeric ICD undergoes induced folding to bind the WRC. This interaction recruits the WRC to the membrane and simultaneously promotes WRC activation by Rac1, which in turn stimulates Arp2/3 to produce branched actin filaments. In (4), the dual actions of HPO-30 ICD cooperatively promote the formation of branched actin networks, which can cause the "swelling" of dendrite observed in previous studies (Shi et al., 2021), an important prerequisite for the outgrowth of a new dendrite branch.

Many neuronal receptors, including SYG-1, Robo, Neogenin, TMEM132, neuroligins, and various protocadherins, contain a short WIRS peptide motif in their ICD, which allows them to recruit the WRC to their sites of action at membranes to regulate local actin polymerization in diverse cellular processes (Chaudhari et al., 2021; Chia et al., 2014; Fan et al., 2018; Lee et al., 2016; Wang et al., 2021; Xing et al., 2018). The HPO-30 ICD does not contain a WIRS motif. Instead, our structural-function analysis, structural predictions, and biophysical measurements reveal that HPO-30 ICD is unstructured, requires dimerization, and likely undergoes induced folding in order to bind to the WRC. This binding mechanism should be conserved throughout animals, as the HPO-30 ICD is conserved only in nematodes, but can bind to both *C. elegans* and human WRCs (Zou et al., 2018). Therefore, HPO-30 represents a new class of transmembrane proteins, which interact with the WRC using a noncanonical mechanism distinct from the previously identified WIRS peptide-containing receptors (Chen et al., 2014a). Exactly how HPO-30 ICD dimerizes and folds into a structure to bind the WRC is currently unknown, but the knowledge from our study has laid the ground for high-resolution structural determination of the HPO-30 ICD bound to the WRC, which will not only reveal the interaction mechanism and

identify the interaction surface on the WRC, but also facilitate the identification of human

homologs of HPO-30 and revelation of new biological functions of this interaction in neural

development (and/or other processes).

Our study suggests that HPO-30 binding to the WRC not only provides a membrane

localization mechanism to recruit the WRC to the dendrite branching site, but also can

simultaneously promote WRC activation by Rac1. The dual, synergistic actions of both binding

and activating the WRC have been observed in several WIRS-containing receptors, such as

PCDH10, in which WIRS peptide binding is responsible for membrane recruitment, while weak,

secondary interactions from the flanking sequences in receptor ICD promotes WRC activation

(Chen et al., 2014a). The HPO-30 ICD, however, is distinct from WIRS-containing receptors in

that the ICD binding can directly promote WRC activation. Due to the strong inhibitory effect of

HPO-30 ICD in pyrene-actin polymerization assays, it will be important to develop new assays

to directly measure WRC activation (e.g. by tracking WCA release) to understand how HPO-30

ICD binding enhances WRC activation by Rac1. Rac1 activates the WRC by binding to two

distinct sites located on the Sra1 subunit (Chen et al., 2017; Ding et al., 2022). It is possible

HPO-30 ICD promotes WRC activation by Rac1 by directly stabilizing Rac1 binding to either

site or by destabilizing the sequestration of WCA.

HPO-30 to our knowledge represents the first transmembrane protein that possesses actin

capping, or capping-like, activity. The abilities of HPO-30 to promote WRC membrane

recruitment and activation (which stimulates Arp2/3-mediated actin polymerization) and to

simultaneously cap actin filaments and inhibit actin polymerization seem to be conflicting

functions. However, such capping activity, which is usually provided by cytosolic capping

proteins like CapZ, are known to be essential for Arp2/3-based actin dynamics in both bead

motility assays *in vitro* and many processes *in vivo* (Akin and Mullins, 2008; Miyoshi et al.,

2006). It is believed that capping proteins can reserve G-actin by blocking it from accessing

barbed ends and simultaneously prevent the barbed end from sequestering the WH2 region of

WCA, which collectively stimulates the formation of new branches through Arp2/3. Thus,

capping proteins are critical for dynamically modulating the actin architecture to generate force

against the surface where actin is localized (Akin and Mullins, 2008; Funk et al., 2021) (**Figure**

7, step 2). The capping activity of HPO-30 ICD in principle should similarly stimulate the

formation of a highly branched actin network. Although HPO-30 ICD apparently has lower

capping activity compared to the canonical capping protein CapZ, it has the advantage of

concentrating the capping activity next to membranes to provide a more focused control of actin

dynamics.

The requirement of dimerization of HPO-30 ICD for both binding the WRC and

providing the capping activity suggests an intriguing regulatory mechanism of membrane

proteins. Clustering of membrane receptors is a common mechanism to increase local

concentration of membrane signaling (Johannes et al., 2018). Enrichment of HPO-30 at

developing PVD neuron dendrites is correlated with the high level of F-actin in these branches

(Zou et al., 2018). In addition to increasing HPO-30 density, clustering of HPO-30 would

facilitate dimerization of its ICD, which can act as a functional switch to allow the ICD to adopt

two functions: binding to the WRC and capping actin filaments. This dimerization-mediated

functional switch can help the cell to distinguish signal from noise and achieve switch-like

spatiotemporal control of actin polymerization in response to upstream stimuli. Given the distinct

surfaces of the actin barbed end and the WRC, it is possible the dimerized HPO-30 ICD uses

distinct structural mechanisms to bind the two targets. Such structural plasticity has been

observed for signaling molecules with versatile functions (Bürgi et al., 2016; Dishman and

Volkman, 2018). Resolving the high-resolution structures and identifying the key residues

responsible for each binding event would allow us to differentiate the two functions of HPO-30,

both in *in vitro* and *in vivo*.

Together, our data establish HPO-30 as the first membrane receptor that, upon

dimerization, can directly integrate the activities of Arp2/3-mediated actin polymerization

(resembling WIRS-containing receptors, albeit through a distinct mechanism) and actin filament

capping (resembling canonical barbed end cappers like CapZ, although the exact mechanism

could be different). These two distinct functions by the same membrane receptor provide an

exciting model to explain how HPO-30 regulates local actin dynamics to facilitate dendrite

branching (**Figure 7**). Synergistic action of both functions of HPO-30 would promote the

formation of highly branched actin networks, giving rise to dendritic "swellings" observed in

PVD neurons prior to dendrite branch outgrowth (**Figure 7**, step 4) (Shi et al., 2021).

Materials and Methods

Protein expression and purification

HPO-30 ICD proteins (and their associated alanine-scan mutants), including GST-ICD,

ICD-GST, DLC8-ICD-His6, GB1-ICD-His6, His9-sumo-ICD, MBP-ICD, GST-ICD-mEGFP,

GB1-FKBP-ICD-His6, GB1-FRB-ICD-His6, GB1-FKBP-His6, GB1-FRB-His6, SNAP-FKBP-

ICD-His6, SNAP-FRB-ICD-His6, SNAP-FKBP-His6, and SNAP-FRB-His6, as well as other

related proteins, including MBP-WAVE1 (1-230), MBP-WAVE1 (1-230)-WCA, MBP-WAVE1

(1-178), MBP-Abi2 (1-158), MBP-HSPC300, $(MBP)_2$-Abi2 (1-158), $(MBP)_2$-HSPC300, and

$Rac1^{Q61L/P29S}$ $\Delta4$ (herein referred to as $Rac1^{QP}$), were individually expressed in Arctic Express

(DE3) RIL (Agilent) or BL21 (DE3) $T1^R$ (Sigma) cells after induction with 0.75 mM IPTG at 10

°C or 18 °C for 16 hours. His10-SNAP-CapZ β and CapZ α were expressed together from the

pCDF Duet vector in Arctic Express (DE3) RIL (Agilent) cells after induction with 0.75 mM

IPTG at 10 °C for 16 hours. GST-ICD or ICD-GST (and alanine-scan mutants) and GST-

mEGFP were purified through Glutathione Sepharose beads (GE Healthcare), followed by cation

exchange chromatography using a Source 15S column (GE Healthcare) at pH 7.0. MBP-ICD

was purified through amylose resin (New England Biolabs), followed by cation exchange

chromatography using a Source 15S column (GE Healthcare) at pH 7.0. The GST tag from GST-

ICD-mEGFP and the MBP tag from MBP-ICD were removed using TEV cleavage at 4 °C

overnight, followed by cation exchange chromatography using a Source 15S column (GE

Healthcare) at pH 7.0. DLC8-ICD-His6, GB1-ICD-His6, His9-sumo-ICD, GB1-FKBP-ICD-His6

(and associated alanine mutant), GB1-FRB-ICD-His6 (and associated alanine mutants), GB1-

FKBP-His6, GB1-FRB-His6, SNAP-FKBP-ICD-His6, SNAP-FRB-ICD-His6, SNAP-FKBP-

His6, and SNAP-FRB-His6 were purified through Ni-NTA Agarose resin (Qiagen), followed by

cation exchange chromatography using a Source 15S column (GE Healthcare) at pH 7.0. GST-

ICD, ICD-GST, DLC8-ICD-His6, GB1-ICD-His6, His6-sumo-ICD, ICD-mEGFP, untagged

ICD, GB1-FKBP-ICD-His6 (and associated alanine mutants), GB1-FRB-ICD-His6 (and

associated alanine mutants), GB1-FKBP-His6, GB1-FRB-His6, SNAP-FKBP-ICD-His6, SNAP-

FRB-ICD-His6, SNAP-FKBP-His6, SNAP-FRB-His6 were further purified through a Superdex

75 column (GE Healthcare). His10-SNAP-CapZ β and CapZ α were purified through Ni-NTA

Agarose resin (Qiagen), followed by anion exchange chromatography using a Source 15Q

column (GE Healthcare) at pH 8.0 and size exclusion chromatography on a Superdex 200

column (GE Healthcare). Rac1^{QP}Δ4 was purified by an SP-Sepharose Fast Flow column (GE

Healthcare) at pH 7.0 followed by size exclusion chromatography through a Superdex 75

column.

MBP-WAVE1, MBP-Abi2, MBP-HSPC300, (MBP)$_2$-Abi2, and (MBP)$_2$-HSPC300

proteins were purified through amylose beads (New England Biolabs). His6-Sra1 and Nap1

dimer were co-expressed in Tni insect cells (Expression systems), and the dimer was purified

through Ni-NTA agarose resin (Qiagen) followed by anion exchange chromatography using a

Source 15Q column at pH 8.0. Pentameric WRC was assembled and purified following

previously described protocols (Chen et al., 2014b, 2010). Briefly, individually purified

WAVE1, Abi2, and HSPC300 subunits were mixed at equimolar ratio in the presence of 1%

(w/v) NP40 and incubated on ice for 48 hours. The assembled trimer was then purified by anion

exchange chromatography through a Source 15Q column at pH 8.0 and cation exchange

chromatography by a Source 15S column at pH 6.0. Dimer and trimer were mixer at equimolar

ratio and incubated on ice for 30 min. The assembled pentamer was purified on amylose beads

(NEB), after which the MBP and His6 tags were cleaved using TEV protease incubation

overnight. The pentamer was further purified using anion exchange chromatography through a

Source 15Q column at pH 8.0 and size exclusion chromatography using a Superdex 200 column.

Actin was purified as previously described from rabbit muscle acetone powder from Pel-

Freeze (Spudich and Watt, 1971). Actin was labeled by pyrene or Alexa Fluor® 647 after

polymerization at 4 °C, using a 10-fold or 2-fold excess dye, respectively. Actin and pyrene actin

for actin polymerization and depolymerization assays were kept in continuous dialysis at 4 °C,

with biweekly buffer changes. Actin and Alexa Fluor® 647-labeled actin for smTIRF

experiments were kept in closed tubes for two weeks.

Generation of WRC-(MBP)$_2$

To create WRC-(MBP)$_2$, MBP-Abi2 (1-158) had the sortase ligation sequence, LPGTG,

genetically fused to the C-terminus. Meanwhile, a TEV site was added to the N-terminus of an

(MBP)$_2$ tag, which exposes a Gly after Tev cleavage. MBP-Abi2 (1-158)-LPGTG was

expressed, purified, and incorporated into the WRC as described above to create WRC-LPGTG.

GG-2MBP was expressed in Arctic Express (DE3) RIL (Agilent) cells after induction with 0.75

mM IPTG at 10 °C for 16 hours, purified on amylose resin, and subjected to TEV cleavage

overnight, followed by anion exchange chromatography using a Source 15Q column (GE

Healthcare). Sortase5M (sortase A pentamutant) was a gift from David Liu (Addgene plasmid #

75144), expressed in Arctic Express (DE3) RIL (Agilent) cells, purified over Ni-NTA agarose

resin, followed by cation exchange using a Source 15S column (GE Healthcare) and size

exclusion chromatography using a Superdex 75 column (GE Healthcare) (Chen et al., 2011). 1

µM WRC-LPGTG was mixed with 25 µM GG-MBP and 10 µM sortase in 50 mM Tris pH 7.5,

150 mM NaCl, and 10 mM $CaCl_2$ and left at room temperature for two hours. The reaction was

quenched by adding 25 mM EGTA, and the WRC-$(MBP)_2$ was purified over a Superdex 200

column to separate the WRC-$(MBP)_2$ from unligated products.

Regular pull-down assay

GST pull-down assays were performed as previously described (Shi et al., 2021). Briefly,

20 µL of GSH-Sepharose beads were mixed with bait protein and prey protein in 1 mL of pull-

down buffer (50 mM NaCl, 10 mM HEPES pH 7.0, 5% (w/v) glycerol, 5 mM 2-

mercaptoethanol, and 0.05% Triton X-100). The samples were mixed at 4 °C for 30 minutes,

washed three times with 1 mL of pull-down buffer, and eluted with 40 µL of elution buffer

containing 30 mM reduced glutathione and 100 mM Tris pH 8.5. MBP pull-down assays were

performed like GST pull-down assays, but used 20 µL of amylose resin and elution buffer that

was pull-down buffer supplemented with 2% [w/v] maltose. His-tagged pull-down assays were

performed as above, used 20 µL of Ni-NTA agarose resin, G-Buffer (2 mM Tris HCl pH 8.0,

200 µM ATP, 0.5 mM DTT, 0.1 mM $CalCl_2$, and 1mM NaN_3) as the wash buffer, and elution

buffer containing 500 mM imidazole pH 7.0. In all pull-down assays, the eluant was examined

by SDS-PAGE and Coomassie blue staining. In all pull-down assays using FKBP and FRB,

rapamycin was added to 5 µM final concentration. As controls, the same volume of DMSO was

added in place of rapamycin. For the alanine scan pull-down quantification, the intensity of the

Sra1 and Nap1 bands were quantified using ImageJ. The intensity from the GST control lane was

subtracted from the alanine protein lane, and the corrected intensity was divided by the intensity

of the wild type HPO-30 lane. A Student's paired t-test was used to compare the wild type and

each mutant separately.

Equilibrium pull-down (EPD) assay

Equilibrium pull-down assays were performed as previously described (Chen et al.,

2017). Briefly, 60 µL of GSH-Sepharose beads (50% slurry equilibrated in pull-down buffer)

were mixed with 0.1 µM WRC and various amounts of GST-tagged protein (from 0.01 µM to 30

µM) and brought to 100 µL with pull-down buffer (composition the same as in GST pull-down

assays, above). The reactions were allowed to mix for 30 min at 4 °C, and four reactions at a

time were spun at 15 krpm for 15 seconds. The supernatant was removed and examined by SDS-

PAGE and Coomassie blue staining. Each assay was repeated 3 times. The Sra1/Nap1 intensity

was quantified using ImageJ to calculate the fractional occupancy. The data was fitted in

DynaFit using a single binding site model (Kuzmic, 1996).

Size exclusion chromatography analysis

GB1-FKBP-ICD and GB1-FRB-ICD were mixed at equimolar ratio and loaded onto a

24-mL Superdex 200 column (GE healthcare) equilibrated in 100 mM NaCl, 10 mM HEPES pH

7.0, 5% (w/v) glycerol, and 1 mM DTT, with or without 5 µM rapamycin.

Pyrene-actin polymerization assay

Pyrene-actin polymerization assays were performed as previously described (Doolittle et

al., 2013a). Actin was purified and pyrene-labeled as described above and kept in continuous

dialysis in G-Buffer (2 mM Tris-HCl pH 8, 0.5 mM DTT, 0.2 mM ATP, 0.1 mM $CaCl_2$, 1 mM

NaN_3) that is changed twice a week. Arp2/3 was purified following existing protocols and kept

aliquoted at -80°C (Doolittle et al., 2013b). All proteins except for the WRC and Arp2/3 were

purified into 50KMEH5Gd (50 mM KCl, 1 mM $MgCl_2$, 1 mM EGTA, 10 mM HEPES pH 7.0,

5% [w/v] glycerol, 1 mM DTT) and stored at -80°C. WRC230VCA was purified into

100KMEI20Gd (100 mM KCl, 10 mM Imidazole pH 7.0, 20% [w/v] glycerol) and kept at -80°C.

Unless otherwise noted, a typical reaction contained 2 µM actin with 10% pyrene labeled, 10 nM

Arp2/3, 100 nM WRC[230WCA] or free WCA, and /or 0.4 µM Rac1[QP]Δ4, and/or additional ICD

ligands to be analyzed, with or without 5 µM rapamycin or an equivalent volume of DMSO. The

excitation and emission wavelengths were set to 365 nm and 407 nm, respectively. Data were

collected on a TECAN SPARK plate reader.

Actin depolymerization assay

Actin depolymerization assays were performed as previously described, with some

modifications (Heiss and Cooper, 1991). Actin at ~20 µM and 70% pyrene-labeling was pre-

polymerized at room temperature overnight by addition of 1 mM $MgCl_2$ and 50 mM KCl. For

each depolymerization reaction, actin was diluted to 5 µM in one pooled mixture, either alone or

with proteins to be tested, and left for 3 minutes at room temperature to allow for protein

binding. Protein at the same concentration was prepared in a second, separate mixture. After 3

minutes, the actin was further diluted 20-fold by the addition of the second mixture. All proteins

were diluted into depolymerization buffer, which is 3 parts G-Buffer and 1 part 50KMEH5Gd

(see above for buffer composition). The excitation and emission wavelengths were set to 365 nm

and 407 nm, respectively. Data were collected on a TECAN SPARK plate reader. To calculate

the relative initial velocity, the slope for the first 30 seconds of the reaction was calculated and

divided by the slope of the actin control. ANOVA on Ranks and Dunn-Tukey tests were

performed to determine significance.

Circular dichroism (CD) measurement

Untagged HPO-30 ICD was purified into 100 mM NaCl, 10 mM HEPES pH 7.0, 5%

(w/v) glycerol, and 1 mM DTT. The same buffer was used to dissolve BSA powder (Fisher Cat #

BP1600-100) and as a blank for the CD spectrometer. The blank buffer was degassed by

sonication for 3 minutes at 30% power. Data were collected on a MOS-500 fluorimeter using an

ALX250 lamp. Data were collected for wavelengths between 190 nm and 260 nm, with a 1 nm

step, 0.5 s acquisition period, and averaged over three repeats. Mean residue ellipticity was

calculated as described previously (Greenfield, 2006). Data for BSA and HPO-30 were collected

at the same concentration of protein. TMAO was dissolved to 3 M in the same buffer as HPO-30

purification and was added to the appropriate concentration before measurement. Separate buffer

blanks containing the same concentration of TMAO were measured to ensure TMAO alone did

not contritube to the spectrum.

Nuclear magnetic resonance (NMR) spectroscopy

Isotopically labeled proteins were expressed and purified as described for non-labeled

proteins, using minimal media containing N^{15} NH_4Cl instead of traditional media. Proteins were

purified into 100 mM NaCl, 10 mM HEPES pH 7.0, 5% (w/v) glycerol, 1 mM DTT and were

supplemented with 10% D_2O. Protein concentration ranged from 70 μM (GB1 tag) to 850 μM

(GB1-ICD). GB1 tag was produced by thrombin cleavage of GB1-ICD and removal of ICD

using nickel-NTA resin (Qiagen). NMR spectra were collected on a Bruker 700 MHz

spectrometer at Iowa State University equipped with z-shielded gradient triple resonance 5 mm

TCI cryoprobe. 2D ^1H-^{15}N TROSY-HSQC and ^1H-^{15}N HSQC experiments were recorded with a

time domain matrix consisting of 100* (t_1, ^{15}N) × 1024* (t_2, ^1H) complex points with acquisition

time of 50 ms (t_1) and 91.8 ms (t_2) using 16 scans per FID and 1.5 s interscan delay. Spectral

widths for ^1H and ^{15}N dimensions were set to 15.9 and 28.2 ppm, respectively, with carriers set

at 4.821 ppm (^1H) and 119.138 ppm (^{15}N).

Fluorophore labeling of proteins

SNAP-tagged proteins were labeled with SNAP-Surface® Alexa Fluor® 488, SNAP-

Surface® Alexa Fluor® 546, and SNAP-Surface® Alexa Fluor® 647 (New England Biolabs). 5

μM protein and 10 μM dye were mixed and allowed to react in 50KMEH5Gd at room

temperature for two hours, followed by desalting into 50KMEH5Gd buffer and concentration.

Dye extinction coefficients were calculated from a standard curve and are as follows: Alexa 488

at 495 nm, 95000 M^{-1}*cm^{-1}; Alexa 546 at 556 nm, 120000 M^{-1}*cm^{-1}; Alexa 647 at 650 nm,

255000 M^{-1}*cm^{-1}. Protein labeling efficiency was calculated by dividing protein concentration

by dye concentration—for Alexa Fluor® 488 the labeling efficiency was estimated at ~100%, for

Alexa Fluor® 546 the labeling efficiency was estimated at ~60%.

smTIRF data collection

All time lapses were collected on a Nikon TE2000-E inverted microscope equipped with

a 100x 1.49 NA TIRF objective and a TIRF Quad filter cube (Chroma C141789), using an Andor

iXon3 EM-CCD (DU-897-CS0) camera, with a GATACA iLas system to prevent uneven

illumination. Coverslips were prepared as described previously with slight modifications

(Narvaez-Ortiz and Nolen, 2022). Briefly, glass coverslips (VWR Cat # 48393-241) were

cleaned with 2% (w/v) Hellmanex, acetone, and 1 M KOH solutions with sonication, and rinsed

extensively with DI water before each step and after the KOH treatment. Coverslips were rinsed

with methanol and dried using a N_2 gas stream. GOPTES (Fisher Cat # G0210100G) was added

to the coverslips, which were then baked at 75°C for 30 minutes. Coverslips were rinsed with

microscope grade acetone and dried with N_2 gas stream. A 2% (w/w) Biotin-PEG3000/NH2-

PEG3000 mixture of powder (Rapp Polymere GmbH) was prepared, placed on top of the

coverslips, and the coverslips baked overnight at 75°C. After overnight baking, the coverslips

were washed with water and dried with N_2 gas. Individual wells were made by placing

functionalized coverslips on Ibidi® sticky-Slide IV 0.4 slides (Ibidi Cat # 80608). Individual

lanes were prepared for microscopy by incubating with 5% (w/v) Pluronic F-127 for 10 min at

RT, followed by either 100 nM (for elongation rate analysis) or 35 nM (for side binding analysis)

neutravidin incubation (in 50 mM Tris pH 7.5, 200 mM NaCl, referred to as LS TBS for short)

for 10 min at RT, and either 10 nM (for elongation rate analysis) or 3.5 nM (for side binding

analysis) biotinylated inactivated heavy meromyosin (in LS TBS) incubation for 10 min at RT.

The chambers were washed with 20 mg/mL BSA in 50 mM Tris pH 7.5, 600 mM NaCl, and

incubated with 20 mg/mL BSA in LS TBS for 10 min at RT. Actin was diluted to a final

concentration of 2.1 μM with 20% Alexa-647 labeling in 1X TIRF buffer (final concentrations:

50 mM KCl, 2 mM $MgCl_2$, 2 mM EGTA pH 8.0, 10 mM Imidazole pH 7.0, 25 mM glucose, 1

mM Trolox, 0.5% methylcellulose (400 cP), 20 mM 2-mercaptoethanol, 0.4 mM ATP, 20

mg/mL BSA, 0.02 mg/mL catalase, 0.1 mg/mL glucose oxidase, 1 mM 4-nitrobenzyl alcohol,

and 0.5 mM propyl gallate) and allowed to polymerize on the slide for 5 minutes at room

temperature. Excess actin was removed by two washes, each time using 40 μL of 1X TIRF

buffer. A separate mixture of 1 μM actin with 20% Alexa-647 labeling containing desired

proteins in 1X TIRF buffer was then added to the wells to start data acquisition. Time lapse

images were acquired using the following setups. Experiments involving unlabeled HPO-30 ICD

and CapZ for elongation rate analysis of actin[AF647]: 640-nm laser, 5% power, 50 ms exposure

time, and a 5-s interval between exposures; experiments involving CapZ[AF488]: 488-nm laser

(15% power, 50 ms exposure time) and 640-nm laser (5% power, 50 ms exposure time),

alternating between 20 consecutive exposures in the 488-nm channel for CapZ[AF488] and one

exposure in the 640-nm channel for actin[AF647], with a 500-ms interval between exposures;

experiments involving HPO-30 ICD[AF546]: 561-nm laser (10% power, 50 or 100 ms exposure

time) and 640-nm laser (5% power, 50 ms exposure time), alternating between 20 consecutive

exposures in the 561-nm channel for ICD[AF546] and one exposure in the 640-nm channel for

actin[AF647], with a 500-ms interval between exposures.

smTIRF data processing - actin elongation rate measurement

Time lapses were opened in ImageJ and the background removed using a rolling ball radius of 10

pixels. The length of actin filaments was calculated using the Filament Length ImageJ plugin,

kindly provided by Jeff Kuhn, using a Gaussian-Radius of 1.5-2, determined for each individual

movie (Kuhn and Pollard, 2005). The length (in microns) was converted to subunits using the

established value of 370 actin subunits / µm (Huxley and Brown, 1967). Time points were taken

from NIS Elements software. Filaments were only selected if they were present within the first

10 frames of the movie and did not leave the frame during the course of the video. 10 time points

were selected and the length at each point was calculated, and the average slope was used for the

actin elongation rate. ANOVA with Dunn-Tukey tests were used to determine significance.

smTIRF data processing - side-binding/capping analysis

Time lapses were opened in ImageJ and the background removed using a rolling ball radius

of 10 pixels. Only filaments present at the beginning of the videos and those that did not leave

the frame during the duration of the video were selected. Analysis was performed in a single-

blinded manner. Side binding events were determined if they met the following criteria: 1) the

ICD/empty tag puncta must be present for more than one frame; 2) the ICD/empty tag puncta

must move with the filament at least once; 3) the filament must not move away from the

ICD/empty tag puncta; 4) the HPO-30/vector puncta must be smaller than a circle with a radius

of 4 pixels. Capping events were confirmed by the absence of growth with puncta present and, if

available, growth from the end capped after the puncta leaves the filament. A Student's t-test was

used to determine significance between ICD and tag reactions.

Actin pelleting assay

Actin pelleting assays were performed based on (Heier et al., 2017) with modifications.

Actin was pre-polymerized at room temperature overnight by addition of 1X 50KMEH5Gd.

Reactions (60 µL) were assembled by mixing 2 µM actin and 5 µM protein (in the same

50KMEH5Gd buffer), which were then allowed to bind at room temperature for 30 minutes.

Reactions were centrifuged at 100,000 g at 4 °C for 30 min in a Type 42.2 Ti rotor in a Beckman

ultracentrifuge. 40 µL of the supernatant was removed and mixed with SDS, and the remaining

~15 µL was removed and discarded. The pellet was dissolved by the addition of 40 µL of G-

buffer, followed by brief pipetting and vortexing, and allowed to sit at room temperature for 5

min before the liquid was removed and mixed with SDS PAGE loading buffer. The intensity of

the supernatant and pellet bands on SDS PAGE gels were measured using ImageJ. The total

intensity of the supernatant and pellet bands, and the percentage of intensity from the pellet and

the supernatant were calculated. The percentage pelleted protein was calculated by subtracting

the percentage of intensity from the pelleted protein in the absence of actin from the percentage

of intensity from the pelleted protein in the presence of actin. For example, in Figure 5A, the

intensity of BSA from lanes 1 and 2 was summed and the percentage of intensity from 1 and

from 2 were calculated. This was repeated for lanes 3 and 4, then the percentage intensity of lane

2 was subtracted from lane 4. ANOVA with Dunn-Tukey tests were used to determine

significance.

Actin pelleting competition assays

Actin pelleting competition assays were performed nearly identically to the actin

pelleting assay described above, except that after 30 minutes of incubation of 5 µM HPO-30 ICD

proteins with 2 µM actin, 600 nM CapZ was added and incubated for 5 minutes before

ultracentrifugation. The intensity of the top bands for CapZ in the pellet were calculated for all

reactions and corrected for the intensity of CapZ pelleted without actin. The relative intensity

was calculated by dividing the intensity of the lane by CapZ with actin alone. For example, in

Figure 6D, the intensity of lane 2 was subtracted from lanes 4, 6, 8, and 10. Then, the corrected

intensity of lane 4 was divided by that of lane 2. ANOVA with Dunn-Tukey tests were used to

determine significance.

S2 cell culture and transfection

Drosophila S2 cells were grown in Schneider's media (Thermo Fisher) supplemented with 10%

heat-inactivated FBS (Life Technologies) and 50 U/mL penicillin-streptomycin (Thermo Fisher).

Cells were transfected with Effectene (Qiagen) and 1ug of total plasmid (either Pactin>HPO-

30:6xMyc (pXD384) or both Pactin>HPO-30:6xMyc (pXD384) and Pactin>HPO-30:HA

(pXD226)).

Co-immunoprecipitation

S2 cells were harvested 72 hours after transfection. Cells were lysed in RIPA buffer (Thermo

Fisher) with 1x Halt Protease Inhibitor Cocktail (Thermofisher) for 30 minutes on ice. Cell

lysates were incubated with anti-HA affinity gel beads (Sigma E6779-1ML) for 1 hour at 4℃

with rotation. Proteins were eluted at 80℃ in NuPAGE LDS Sample Buffer (Life Technologies)

supplemented with DTT (GoldBio) and detected using Western blot with mouse antibody to HA

(1:1000, Sigma H3663), rabbit antibody to Myc (1:1000, Santa Cruz Biotechnology sc-789), and

HRP-conjugated goat antibodies to mouse (1:20,000, Jackson Immuno Research).

Acknowledgements

We thank ResearchIT at Iowa State University for installing AlphaFold 2 and AlphaFold

Multimer; Daniel Boesch for help with submitting AlphaFold prediction tasks; Scott Nelson at

Iowa State for guidance with CD spectroscopy; Xuefeng Wang at Iowa State for the initial setup

of smTIRF experiments; Aubrey Sijo-Gonzales, Finlan Rhodes, Leyuan Loh, Ganesh Prasad,

Simanta Mitra, and the ResearchIT at Iowa State University for establishing a web-based

application (*https://biochempy.bb.iastate.edu*) for data visualization; and Michael Rosen at UT

Southwestern for sharing FKBP/FRB constructs. This work was supported by funding from the

National Institutes of Health (R35-GM128786) and start-up funds from the Iowa State University

and the Roy J. Carver Charitable Trust to B.C., National Institutes of Health (R35-GM136319) to

B.J.N., National Institutes of Health (R01-GM132561) to J.R., Howard Hughes Medical Institute

and the National Institute of Neurological Disorders and Stroke (1R01NS082208) to K.S., and the

National Science Foundation Graduate Research Fellowship and a training grant from the National

Institutes of Health to R.S.

1008

Author contributions

B.C. conceived and oversaw the project. B.N. oversaw single molecule experiments. J.R. oversaw

NMR experiments. D.A.K. purified proteins and performed biochemical experiments. D.A.K.

performed AlphaFold predictions. D.A.K. and J.R. performed NMR measurements. D.A.K. and

H.Y.N-O. performed single molecule TIRF experiments. R.S. performed co-immunoprecipitation

experiments under the supervision of K.S. D.A.K. and B.C. drafted the manuscript and prepared

the figures with assistance from all the authors.

1016

1017 **Competing interests**

1018 The authors declare no competing interests.

1019

1020 **Additional Information**

1021 Correspondence and request for materials should be addressed to stone@iastate.edu

1022

1023 **Supplemental materials**

1024 **Video 1. CapZ capping event example 1.** Related to Figure 6A, top. Time lapse of SNAP[AF488]-

1025 CapZ (green) bound to the end of actin filaments (red) obtained by smTIRF microscopy. Video

1026 length of 1200 frames at 120 frames per second. The filament in the top left corner is the one

1027 showcased in Figure 6A, top.

1028

1029 **Video 2. CapZ capping event example 2.** Related to Figure 6A, top. Time lapse of SNAP[AF488]-

1030 CapZ (green) bound to the end of an actin filament (red) obtained by smTIRF microscopy. Video

1031 length of 1800 frames at 120 frames per second.

1032

1033 **Video 3. HPO-30 ICD side binding event example 1.** Related to Figure 6A, bottom.

1034 SNAP[AF546]-F/F-ICD (cyan) bound to the side of an actin filament (red) obtained by smTIRF

1035 microscopy. Video length of 1000 frames at 100 frames per second.

1036

1037 **Video 4. HPO-30 ICD side binding event example 2.** Related to Figure 6A, bottom.

1038 SNAP[AF546]-F/F-ICD (cyan) bound to the side of an actin filament (red) obtained by smTIRF

1039 microscopy. Video length of 350 frames at 35 frames per second.

1040

1041 **Video 5. Actin elongation rate comparison.** Related to Figure 6C. Representative smTIRF time

1042 lapses used to quantify the elongation rate of actin filaments. Video shows a random selection of

1043 actin filaments from each condition tested. Video length of 15 minutes and 181 frames was

1044 compressed to 9 seconds.

1045

1046

1047

1048 **Supplemental Table 1: Sequences of recombinant proteins used in this study.**

Construct name	Description	Plasmid Identity	Sequence	Source or reference
GST-ICD	GST-thrombin-TEV-ceHPO-30 ICD (a.a. 229-279)	pDK079	MSPILGYWKIKGLVQPTRLLLEYLEEKYEEHLYERDEGDKWRNKKFELGLEFPNLPY YIDGDVKLTQSMAIIRYIADKHNMLGGCPKERAEISMLEGAVLDIRYGVSRIAYSKD FETLKVDFLSKLPEMLKMFEDRLCHKTYLNGDHVTHPDFMLYDALDVVLYMDPMCLD AFPKLVCFKKRIEAIPQIDKYLKSSKYIAWPLQGWQATFGGGDHPPKSDLVPRGSEN LYFQGHM<u>T</u>SKHAHDVCCTSRKEYREQTKWKNNGLILKTGRVNHQSHRPFVVIDDDSS M	Zou, 2018
GST-ICD Δ1	GST-thrombin-TEV-ceHPO-30 ICD Ala 229-233	pDK325	MSPILGYWKIKGLVQPTRLLLEYLEEKYEEHLYERDEGDKWRNKKFELGLEFPNLPY YIDGDVKLTQSMAIIRYIADKHNMLGGCPKERAEISMLEGAVLDIRYGVSRIAYSKD FETLKVDFLSKLPEMLKMFEDRLCHKTYLNGDHVTHPDFMLYDALDVVLYMDPMCLD AFPKLVCFKKRIEAIPQIDKYLKSSKYIAWPLQGWQATFGGGDHPPKSDLVPRGSEN LYFQGHM<u>A</u>AAAAHDVCCTSRKEYREQTKWKNNGLILKTGRVNHQSHRPFVVIDDDSS M	Zou, 2018
GST-ICD Δ2	GST-thrombin-TEV-ceHPO-30 ICD Ala 234-238	pDK326	MSPILGYWKIKGLVQPTRLLLEYLEEKYEEHLYERDEGDKWRNKKFELGLEFPNLPY YIDGDVKLTQSMAIIRYIADKHNMLGGCPKERAEISMLEGAVLDIRYGVSRIAYSKD FETLKVDFLSKLPEMLKMFEDRLCHKTYLNGDHVTHPDFMLYDALDVVLYMDPMCLD AFPKLVCFKKRIEAIPQIDKYLKSSKYIAWPLQGWQATFGGGDHPPKSDLVPRGSEN LYFQGHM<u>T</u>SKHAAAAAATSRKEYREQTKWKNNGLILKTGRVNHQSHRPFVVIDDDSS M	Zou, 2018
GST-ICD Δ3	GST-thrombin-TEV-ceHPO-30 ICD Ala 239-243	pDK327	MSPILGYWKIKGLVQPTRLLLEYLEEKYEEHLYERDEGDKWRNKKFELGLEFPNLPY YIDGDVKLTQSMAIIRYIADKHNMLGGCPKERAEISMLEGAVLDIRYGVSRIAYSKD FETLKVDFLSKLPEMLKMFEDRLCHKTYLNGDHVTHPDFMLYDALDVVLYMDPMCLD AFPKLVCFKKRIEAIPQIDKYLKSSKYIAWPLQGWQATFGGGDHPPKSDLVPRGSEN LYFQGHM<u>T</u>SKHADVCCAAAAAYREQTKWKNNGLILKTGRVNHQSHRPFVVIDDDSS M	Zou, 2018
GST-ICD Δ4	GST-thrombin-TEV-ceHPO-30 ICD Ala 244-248	pDK328	MSPILGYWKIKGLVQPTRLLLEYLEEKYEEHLYERDEGDKWRNKKFELGLEFPNLPY YIDGDVKLTQSMAIIRYIADKHNMLGGCPKERAEISMLEGAVLDIRYGVSRIAYSKD FETLKVDFLSKLPEMLKMFEDRLCHKTYLNGDHVTHPDFMLYDALDVVLYMDPMCLD AFPKLVCFKKRIEAIPQIDKYLKSSKYIAWPLQGWQATFGGGDHPPKSDLVPRGSEN LYFQGHM<u>T</u>SKHAHDVCCTSRKEAAAAAKWKNNGLILKTGRVNHQSHRPFVVIDDDSS M	Zou, 2018
GST-ICD Δ5	GST-thrombin-TEV-ceHPO-30 ICD Ala 249-253	pDK329	MSPILGYWKIKGLVQPTRLLLEYLEEKYEEHLYERDEGDKWRNKKFELGLEFPNLPY YIDGDVKLTQSMAIIRYIADKHNMLGGCPKERAEISMLEGAVLDIRYGVSRIAYSKD FETLKVDFLSKLPEMLKMFEDRLCHKTYLNGDHVTHPDFMLYDALDVVLYMDPMCLD AFPKLVCFKKRIEAIPQIDKYLKSSKYIAWPLQGWQATFGGGDHPPKSDLVPRGSEN LYFQGHM<u>T</u>SKHAHDVCCTSRKEYREQTAAAAAGLILKTGRVNHQSHRPFVVIDDDSS M	Zou, 2018
GST-ICD Δ6	GST-thrombin-TEV-ceHPO-30 ICD Ala 254-258	pDK330	MSPILGYWKIKGLVQPTRLLLEYLEEKYEEHLYERDEGDKWRNKKFELGLEFPNLPY YIDGDVKLTQSMAIIRYIADKHNMLGGCPKERAEISMLEGAVLDIRYGVSRIAYSKD FETLKVDFLSKLPEMLKMFEDRLCHKTYLNGDHVTHPDFMLYDALDVVLYMDPMCLD AFPKLVCFKKRIEAIPQIDKYLKSSKYIAWPLQGWQATFGGGDHPPKSDLVPRGSEN LYFQGHM<u>T</u>SKHAHDVCCTSRKEYREQTKWKNNAAAAATGRVNHQSHRPFVVIDDDSS M	Zou, 2018
GST-ICD Δ7	GST-thrombin-TEV-ceHPO-30 ICD Ala 259-263	pDK331	MSPILGYWKIKGLVQPTRLLLEYLEEKYEEHLYERDEGDKWRNKKFELGLEFPNLPY YIDGDVKLTQSMAIIRYIADKHNMLGGCPKERAEISMLEGAVLDIRYGVSRIAYSKD FETLKVDFLSKLPEMLKMFEDRLCHKTYLNGDHVTHPDFMLYDALDVVLYMDPMCLD AFPKLVCFKKRIEAIPQIDKYLKSSKYIAWPLQGWQATFGGGDHPPKSDLVPRGSEN LYFQGHM<u>T</u>SKHAHDVCCTSRKEYREQTKWKNNGLILKAAAAAHQSHRPFVVIDDDSS M	Zou, 2018
ICD-GST	ceHPO-30 ICD (a.a. 229-279)-GG-3C-GST	pDK238	M<u>T</u>SKHAHDVCCTSRKEYREQTKWKNNGLILKTGRVNHQSHRPFVVIDDDSS<u>M</u>GGLEV LFQGPMSPILGYWKIKGLVQPTRLLLEYLEEKYEEHLYERDEGDKWRNKKFELGLEF PNLPYYIDGDVKLTQSMAIIRYIADKHNMLGGCPKERAEISMLEGAVLDIRYGVSRI AYSKDFETLKVDFLSKLPEMLKMFEDRLCHKTYLNGDHVTHPDFMLYDALDVVLYMD PMCLDAFPKLVCFKKRIEAIPQIDKYLKSSKYIAWPLQGWQATFGGGDHPPK	This paper
ICD Δ8 - GST	ceHPO-30 ICD Ala 264-268	pDK257	M<u>T</u>SKHAHDVCCTSRKEYREQTKWKNNGLILKTGRVNAAAAAPFVVIDDDSS<u>M</u>GGLEV LFQGPMSPILGYWKIKGLVQPTRLLLEYLEEKYEEHLYERDEGDKWRNKKFELGLEF PNLPYYIDGDVKLTQSMAIIRYIADKHNMLGGCPKERAEISMLEGAVLDIRYGVSRI AYSKDFETLKVDFLSKLPEMLKMFEDRLCHKTYLNGDHVTHPDFMLYDALDVVLYMD PMCLDAFPKLVCFKKRIEAIPQIDKYLKSSKYIAWPLQGWQATFGGGDHPPK	This paper
ICD Δ9 - GST	ceHPO-30 ICD Ala 269-273	pDK258	M<u>T</u>SKHAHDVCCTSRKEYREQTKWKNNGLILKTGRVNHQSHRAAAAADDDSS<u>M</u>GGLEV LFQGPMSPILGYWKIKGLVQPTRLLLEYLEEKYEEHLYERDEGDKWRNKKFELGLEF PNLPYYIDGDVKLTQSMAIIRYIADKHNMLGGCPKERAEISMLEGAVLDIRYGVSRI AYSKDFETLKVDFLSKLPEMLKMFEDRLCHKTYLNGDHVTHPDFMLYDALDVVLYMD PMCLDAFPKLVCFKKRIEAIPQIDKYLKSSKYIAWPLQGWQATFGGGDHPPK	This paper

ICD Δ10 - GST	ceHPO-30 ICD Ala 274-279	pDK259	MTSKHAHDVCCTSRKEYREQTKWKNNGLILKTGRVNHQSHRPFVVIAAAAAAGGGLEV LFQGPMSPILGYWKIKGLVQPTRLLLEYLEEKYEEHLYERDEGDKWRNKKFELGLEF PNLPYYIDGDVKLTQSMAIIRYIADKHNMLGGCPKERAEISMLEGAVLDIRYGVSRI AYSKDFETLKVDFLSKLPEMLKMFEDRLCHKTYLNGDHVTHPDFMLYDALDVVLYMD PMCLDAFPKLVCFKKRIEAIPQIDKYLKSSKYIAWPLQGWQATFGGGDHPPK	This paper
GST-ICD Δ11	GST-thrombin-TEV-ceHPO-30 ICD GGS 249-258	pDK332	MSPILGYWKIKGLVQPTRLLLEYLEEKYEEHLYERDEGDKWRNKKFELGLEFPNLPY YIDGDVKLTQSMAIIRYIADKHNMLGGCPKERAEISMLEGAVLDIRYGVSRIAYSKD FETLKVDFLSKLPEMLKMFEDRLCHKTYLNGDHVTHPDFMLYDALDVVLYMDPMCLD AFPKLVCFKKRIEAIPQIDKYLKSSKYIAWPLQGWQATFGGGDHPPKSDLVPRGSEN LYFQGHMTSKHAHDVCCTSRKEYREQTGGSGGSGGSGSGTGRVNHQSHRPFVVIDDDSSM	This paper
GST-ICD Δ12	GST-thrombin-TEV-ceHPO-30 ICD GGS 264-273	pDK333	MSPILGYWKIKGLVQPTRLLLEYLEEKYEEHLYERDEGDKWRNKKFELGLEFPNLPY YIDGDVKLTQSMAIIRYIADKHNMLGGCPKERAEISMLEGAVLDIRYGVSRIAYSKD FETLKVDFLSKLPEMLKMFEDRLCHKTYLNGDHVTHPDFMLYDALDVVLYMDPMCLD AFPKLVCFKKRIEAIPQIDKYLKSSKYIAWPLQGWQATFGGGDHPPKSDLVPRGSEN LYFQGHMTSKHAHDVCCTSRKEYREQTKWKNNAAAAAATGRVNGGSGGSGGSGDDDSS M	This paper
GST-ICD Δ13	GST-thrombin-TEV-ceHPO-30 ICD GGS 264-275	pDK334	MSPILGYWKIKGLVQPTRLLLEYLEEKYEEHLYERDEGDKWRNKKFELGLEFPNLPY YIDGDVKLTQSMAIIRYIADKHNMLGGCPKERAEISMLEGAVLDIRYGVSRIAYSKD FETLKVDFLSKLPEMLKMFEDRLCHKTYLNGDHVTHPDFMLYDALDVVLYMDPMCLD AFPKLVCFKKRIEAIPQIDKYLKSSKYIAWPLQGWQATFGGGDHPPKSDLVPRGSEN LYFQGHMTSKHAHDVCCTSRKEYREQTGGSGGSGGSGSGSGGSGGSGSGGSGGSGGSDSS M	This paper
GST-ICD Δ14	GST-thrombin-TEV-ceHPO-30 ICD GGS 229-248	pDK335	MSPILGYWKIKGLVQPTRLLLEYLEEKYEEHLYERDEGDKWRNKKFELGLEFPNLPY YIDGDVKLTQSMAIIRYIADKHNMLGGCPKERAEISMLEGAVLDIRYGVSRIAYSKD FETLKVDFLSKLPEMLKMFEDRLCHKTYLNGDHVTHPDFMLYDALDVVLYMDPMCLD AFPKLVCFKKRIEAIPQIDKYLKSSKYIAWPLQGWQATFGGGDHPPKSDLVPRGSEN LYFQGHMGGSGGSGGSGSGGSGGSGGSGGKWKNNGLILKTGRVNHQSHRPFVVIDDDSS M	This paper
ICD-mEGFP	GST-TEV-ceHPO-30 ICD 229-279 – GGS2-mEGFP, GST tag removed by TEV cleavage	pDK184	MSPILGYWKIKGLVQPTRLLLEYLEEKYEEHLYERDEGDKWRNKKFELGLEFPNLPY YIDGDVKLTQSMAIIRYIADKHNMLGGCPKERAEISMLEGAVLDIRYGVSRIAYSKD FETLKVDFLSKLPEMLKMFEDRLCHKTYLNGDHVTHPDFMLYDALDVVLYMDPMCLD AFPKLVCFKKRIEAIPQIDKYLKSSKYIAWPLQGWQATFGGGDHPPKSDLVPRGSEN LYFQGHMTSKHAHDVCCTSRKEYREQTKWKNNGLILKTGRVNHQSHRPFVVIDDDSS MGGSGGSMVSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTLKFICT TGKLPVPWPTLVTTLTYGVQCFSRYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNY KTRAEVKFEGDTLVNRIELKGIDFKEDGNILGHKLEYNYNSHNVYIMADKQKNGIKV NFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPDNHYLSTQSKLSKDPNEKRDHMVL KEKVTAAGITLGMDELYK	This paper
ICD	MBP-TEV-ceHPO-30 ICD 229-279, MBP tag removed by TEV cleavage	pDK092	MKIEEGKLVIWINGDKGYNGLAEVGKKFEKDTGIKVTVEHPDKLEEKFPQVAATGDG PDIIFWAHDRFGGYAQSGLLAEITPDKAFQDKLYPFTWDAVRYNGKLIAYPIAVEAL SLIYNKDLLPNPPKTWEEIPALDKELKAKGKSALMFNLQEPYFTWPLIAADGGYAFK YENGKYDIKDVGVDNAGAKAGLTFLVDLIKNKHMNADTDYSIAEAAFNKGETAMTIN GPWAWSNIDTSKVNYGVTVLPTFKGQPSKPFVGVLSAGINAASPNKELAKEFLENYL LTDEGLEAVNKDKPLGAVALKSYEEELAKDPRIAATMENAQKGEIMPNIPQMSAFWY AVRTAVINAASGRQTVDEALKDAQTNSSSNNNNNNNNNNLGIEGRISEFENLYFQGH TSKHAHDVCCTSRKEYREQTKWKNNGLILKTGRVNHQSHRPFVVIDDDSSM	This paper
DLC8-ICD	DLC8-TEV-ceHPO-30 ICD 229-279-His6	pDK268	MSDRKAVIKNADMSEEMQQDAVDCATQALEKYNIEKDIAAYIKKEFDKKYNPTWHCI VGRNFGSYVTHETRHFIYFYLGQVAILLLFKSGGSENLYFQGHMTSKHAHDVCCTSRK EYREQTKWKNNGLILKTGRVNHQSHRPFVVIDDDSSMHHHHHH	This paper
GB1-ICD	GB1-thrombin-ceHPO-30 ICD 229-279-His6	pDK239	MQYKLALNGKTLKGETTTEAVDAATAEKVFKQYANDNGVDGEWTYDDATKTFTVTEL VPRGSTSKHAHDVCCTSRKEYREQTKWKNNGLILKTGRVNHQSHRPFVVIDDDSSMH HHHHH	This paper
sumo-ICD	His10-Sumo-GGS-ceHPO-30 ICD 229-279	pDK219	MGHHHHHHHHHHSSGHIEGRHMASMSDSEVNQEAKPEVKPETHINLKVSDGSS EIFFKIKKTTPLRRLMEAFAKRQGKEMDSLRFLYDGIRIQADQTPEDLDMEDNDIIE AHREQIGGSTSKHAHDVCCTSRKEYREQTKWKNNGLILKTGRVNHQSHRPFVVIDDD SSM	This paper
GB1-FKBP-ICD	GB1-thrombin-FKBP-GGS3-ceHPO-30 ICD 229-279 – His6	pDK264	MQYKLALNGKTLKGETTTEAVDAATAEKVFKQYANDNGVDGEWTYDDATKTFTVTEL VPRGSHMVQVETISPGDGRTFPKRGQTCVVHYTGMLEDGKKFDSSRDRNKPFKFMLG KQEVIRGWEEGVAQMSVGQRAKLTISPDYAYGATGHPGIIPPHATLVFDVELLKLEG GSGGSGGSMTSKHAHDVCCTSRKEYREQTKWKNNGLILKTGRVNHQSHRPFVVIDDD SSMHHHHHH	This paper
GB1-FRB-ICD	GB1-thrombin-FRB-GGS3-ceHPO-30	pDK265	MQYKLALNGKTLKGETTTEAVDAATAEKVFKQYANDNGVDGEWTYDDATKTFTVTEL VPRGSHMELIRVAILWHEMWHEGLEEASRLYFGERNVKGMFEVLEPLHAMMERGPQT LKETSFNQAYGRDLMEAQEWCRKYMKSGNVKDLTQAWDLYYHVFRRISKQGGSGGSG	This paper

	ICD 229-279 – His6		GSMTSKHAHDVCCTSRKEYREQTKWKNNGLILKTGRVNHQSHRPFVVIDDDSSMHHH HHH	
GB1-FKBP	GB1-thrombin-FKBP-GGS3-His6	pDK270	MQYKLALNGKTLKGETTTEAVDAATAEKVFKQYANDNGVDGEWTYDDATKTFTVTEL VPRGSHMVQVETISPGDGRTFPKRGQTCVVHYTGMLEDGKKFDSSRDRNKPFKFMLG KQEVIRGWEEGVAQMSVGQRAKLTISPDYAYGATGHPGIIPPHATLVFDVELLKLEG GSGGSGGSHHHHHH	This paper
GB1-FRB	GB1-thrombin-FRB-GGS3-His6	pDK271	MQYKLALNGKTLKGETTTEAVDAATAEKVFKQYANDNGVDGEWTYDDATKTFTVTEL VPRGSHMELIRVAILWHEMWHEGLEEASRLYFGERNVKGMFEVLEPLHAMMERGPQT LKETSFNQAYGRDLMEAQEWCRKYMKSGNVKDLTQAWDLYYHVFRRISKQGGSGGSG GSHHHHHH	This paper
GB1-FKBP-ICD Δ5	GB1-thrombin-FKBP-GGS3-ceHPO-30 ICD Ala 249-253– His6	pDK274	MQYKLALNGKTLKGETTTEAVDAATAEKVFKQYANDNGVDGEWTYDDATKTFTVTEL VPRGSHMVQVETISPGDGRTFPKRGQTCVVHYTGMLEDGKKFDSSRDRNKPFKFMLG KQEVIRGWEEGVAQMSVGQRAKLTISPDYAYGATGHPGIIPPHATLVFDVELLKLEG GSGGSGGSMTSKHAHDVCCTSRKEYREQTAAAAAGLILKTGRVNHQSHRPFVVIDDD SSMHHHHHH	This paper
GB1-FRB-ICD Δ5	GB1-FRB-thrombin-GGS3-ceHPO-30 ICD Ala 249-253– His6	pDK275	QYKLALNGKTLKGETTTEAVDAATAEKVFKQYANDNGVDGEWTYDDATKTFTVTELV PRGSHMELIRVAILWHEMWHEGLEEASRLYFGERNVKGMFEVLEPLHAMMERGPQTL KETSFNQAYGRDLMEAQEWCRKYMKSGNVKDLTQAWDLYYHVFRRISKQGGSGGSGG SMTSKHAHDVCCTSRKEYREQTAAAAAGLILKTGRVNHQSHRPFVVIDDDSSMHHHH HH	This paper
SNAP-FKBP-ICD	SNAP-FKBP-GGS3-ceHPO-30 ICD 229-279-His6	pDK283	MDKDCEMKRTTLDSPLGKLELSGCEQGLHEIKLLGKGTSAADAVEVPAPAAVLGGPE PLMQATAWLNAYFHQPEAIEEFPVPALHHPVFQQESFTRQVLWKLLKVVKFGEVISY QQLAALAGNPAATAAVKTALSGNPVPILIPCHRVVSSSGAAGGYEGGLAVKEWLLAH EGHRLGKPGLGPAGIGAPGSMVQVETISPGDGRTFPKRGQTCVVHYTGMLEDGKKFD SSRDRNKPFKFMLGKQEVIRGWEEGVAQMSVGQRAKLTISPDYAYGATGHPGIIPPH ATLVFDVELLKLEGGSGGSGGSMTSKHAHDVCCTSRKEYREQTKWKNNGLILKTGRV NHQSHRPFVVIDDDSSMHHHHHH	https://ww w.addgene. org/101137 / and this paper
SNAP-FRB-ICD	SNAP-FRB-GGS3-ceHPO-30 ICD 229-279-His6	pDK281	MDKDCEMKRTTLDSPLGKLELSGCEQGLHEIKLLGKGTSAADAVEVPAPAAVLGGPE PLMQATAWLNAYFHQPEAIEEFPVPALHHPVFQQESFTRQVLWKLLKVVKFGEVISY QQLAALAGNPAATAAVKTALSGNPVPILIPCHRVVSSSGAVGGYEGGLAVKEWLLAH EGHRLGKPGLGPAGIGAPGSMELIRVAILWHEMWHEGLEEASRLYFGERNVKGMFEV LEPLHAMMERGPQTLKETSFNQAYGRDLMEAQEWCRKYMKSGNVKDLTQAWDLYYHV FRRISKQGGSGGSGGSMTSKHAHDVCCTSRKEYREQTKWKNNGLILKTGRVNHQSHR PFVVIDDDSSMHHHHHH	https://ww w.addgene. org/101137 / and this paper
SNAP-FKBP	SNAP-FKBP-GGS3-His6	pDK282	MDKDCEMKRTTLDSPLGKLELSGCEQGLHEIKLLGKGTSAADAVEVPAPAAVLGGPE PLMQATAWLNAYFHQPEAIEEFPVPALHHPVFQQESFTRQVLWKLLKVVKFGEVISY QQLAALAGNPAATAAVKTALSGNPVPILIPCHRVVSSSGAVGGYEGGLAVKEWLLAH EGHRLGKPGLGPAGIGAPGSMVQVETISPGDGRTFPKRGQTCVVHYTGMLEDGKKFD SSRDRNKPFKFMLGKQEVIRGWEEGVAQMSVGQRAKLTISPDYAYGATGHPGIIPPH ATLVFDVELLKLEGGSGGSGGSHHHHHH	https://ww w.addgene. org/101137 / and this paper
SNAP-FRB	SNAP-FRB-GGS3-His6	pDK280	MDKDCEMKRTTLDSPLGKLELSGCEQGLHEIKLLGKGTSAADAVEVPAPAAVLGGPE PLMQATAWLNAYFHQPEAIEEFPVPALHHPVFQQESFTRQVLWKLLKVVKFGEVISY QQLAALAGNPAATAAVKTALSGNPVPILIPCHRVVSSSGAVGGYEGGLAVKEWLLAH EGHRLGKPGLGPAGIGAPGSMELIRVAILWHEMWHEGLEEASRLYFGERNVKGMFEV LEPLHAMMERGPQTLKETSFNQAYGRDLMEAQEWCRKYMKSGNVKDLTQAWDLYYHV FRRISKQGGSGGSGGSHHHHHH	https://ww w.addgene. org/101137 / and this paper
SNAP-CapZ	His9-SNAP-CapZ β / CapZ α, from *Gallus gallus*	pDK288	>His9-SNAP-CapZ β MGHHHHHHHHHENLYFQGSEFMDKDCEMKRTTLDSPLGKLELSGCEQGLHEIKLLGK GTSAADAVEVPAPAAVLGGPEPLMQATAWLNAYFHQPEAIEEFPVPALHHPVFQQES FTRQVLWKLLKVVKFGEVISYQQLAALAGNPAATAAVKTALSGNPVPILIPCHRVVS SSGAVGGYEGGLAVKEWLLAHEGHRLGKPGLGPAGIGAPGSMSDQQLDCALDLMRRL PPQQIEKNLSDLIDLVPSLCEDLLSSVDQPLKIARDKVVGKDYLLCDYNRDGDSYRS PWSNKYDPPLEDGAMPSARLRKLEVEANNAFDQYRDLYFEGGVSSVYLWDLDHGFAG VILIKKAGDGSKKIKGCWDSIHVVEVQEKSSGRTAHYKLTSTVMLWLQTNKTGSGTM NLGGSLTRQMEKDETVSDSSPHIANIGRLVEDMENKIRSTLNEIYFGKTKDIVNGLR SIDAIPDNQKYKQLQRELSQVLTQRQIYIQPDN

>CapZ α MADFEDRVSDEEKVRIAAKFITHAPPGEFNEVFNDVRLLLNNDNLLREGAAHAFAQY NMDQFTPVKIEGYDDQVLITEHGDLGNGRFLDPRNKISFKFDHLRKEASDPQPEDTE SALKQWRDACDSALRAYVKDHYPNGFCTVYGKSIDGQQTIIACIESHQFQPKNFWNG RWRSEWKFTITPPTAQVAAVLKIQVHYYEDGNVQLVSHKDIQDSVQVSSDVQTAKEF IKIIENAENEYQTAISENYQTMSDTTFKALRRQLPVTRTKIDWNKILSYKIGKEMQN A | https://ww w.addgene. org/101137 / and this paper |
| Sra1 | His6-TEV-hSra1 (1-1253, FL). | pDK116 | MSYYHHHHHHDYDIPTTENLYFQGAMAAQVTLEDALSNVDLLEELPLPDQQPCIEPP PSSLLYQPNFNTNFEDRNAFVTGIARYIEQATVHSSMNEMLEEGQEYAVMLYTWRSC SRAIPQVKCNEQPNRVEIYEKTVEVLEPEVTKLMNFMYFQRNAIERFCGEVRRLCHA | (Ismail et al., 2009) |

	His6 tag removed by TEV cleavage		ERRKDFVSEAYLITLGKFINMFAVLDELKNMKCSVKNDHSAYKRAAQFLRKMADPQS IQESQNLSMFLANHNKITQSLQQQLEVISGYEELLADIVNLCVDYYENRMYLTPSEK HMLLKVMGFGLYLMDGSVSNIYKLDAKKRINLSKIDKYFKQLQVVPLFGDMQIELAR YIKTSAHYEENKSRWTCTSSGSSPQYNICEQMIQIREDHMRFISELARYSNSEVVTG SGRQEAQKTDAEYRKLFDLALQGLQLLSQWSAHVMEVYSWKLVHPTDKYSNKDCPDS AEEYERATRYNYTSEEKFALVEVIAMIKGLQVLMGRMESVFNHAIRHTVYAALQDFS QVTLREPLRQAIKKKKNVIQSVLQAIRKTVCDWETGHEPFNDPALRGEKDPKSGFDI KVPRRAVGPSSTQLYMVRTMLESLIADKSGSKKTLRSSLEGPTILDIEKFHRESFFY THLINFSETLQQCCDLSQLWFREFFLELTMGRRIQFPIEMSMPWILTDHILETKEAS MMEYVLYSLDLYNDSAHYALTRFNKQFLYDEIEAEVNLCFDQFVYKLADQIFAYYKV MAGSLLLDKRLRSECKNQGATIHLPPSNRYETLLKQRHVQLLGRSIDLNRLITQRVS AAMYKSLELAIGRFESEDLTSIVELDGLLEINRMTHKLLSRYLTLDGFDAMFREANH NVSAPYGRITLHVFWELNYDFLPNYCYNGSTNRFVRTVLPFSQEFQRDKQPNAQPQY LHGSKALNLAYSSIYGSYRNFVGPPHFQVICRLLGYQGIAVVMEELLKVVKSLLQGT ILQYVKTLMEVMPKICRLPRHEYGSPGILEFFHHQLKDIVEYAELKTVCFQNLREVG NAILFCLLIEQSLSLEEVCDLLHAAPFQNILPRVHVKEGERLDAKMKRLESKYAPLH LVPLIERLGTPQQIAIAREGDLLTKERLCCGLSMFEVILTRIRSFLDDPIWRGPLPS NGVMHVDECVEFHRLWSAMQFVYCIPVGTHEFTVEQCFGDGLHWAGCMIIVLLGQQR RFAVLDFCYHLLKVQKHDGKDEIIKNVPLKKMVERIRKFQILNDEIITILDKYLKSG DGEGTPVEHVRCFQPPIHQSLASS	
Nap1	hNap1 (1-1128, FL)	pDK149	MSRSVLQPSQQKLAEKLTILNDRGVGMLTRLYNIKKACGDPKAKPSYLIDKNLESAV KFIVRKFPAVETRNNNQQLAQLQKEKSEILKNLALYYFTFVDVMEFKDHVCELLNTI DVCQVFFDITVNFDLTKNYLDLIITYTTLMILLSRIEERKAIIGLYNYAHEMTHGAS DREYPRLGQMIVDYENPLKKMMEEFVPHSKSLSDALISLQMVYPRRNLSADQWRNAQ LLSLISAPSTMLNPAQSDTMPCEYLSLDAMEKWIIFGFILCHGILNTDATALNLWKL ALQSSSCLSLFRDEVFHIHKAAEDLFVNIRGYNKRINDIRECKEAAVSHAGSMHRER RKFLRSALKELATVLSDQPGLLGPKALFVFMALSFARDEIIWLLRHADNMPKKSADD FIDKHIAELIFYMEELRAHVRKYGPVMQRYYVQYLSGFDAVVLNELVQNLSVCPEDE SIIMSSFVNTMTSLSVKQVEDGEVFDFRGMRLDWFRLQAYTSVSKASLGLADHRELG KMMNTIIFHTKMVDSLVEMLVETSDLSIFCFYSRAFEKMFQQCLELPSQSRYSIAFP LLCTHFMSCTHELCPEERRHHIGDRSLSLCNMFLDEMAKQARNLITDICTEQCTLSDQ LLPKHCAKTISQAVNKKSKKQTGKKGEPEREKPGVESMRKNRLVVTNLDKLHTALSE LCFSINYVPNMVVWEHTFTPREYLTSHLEIRFTKSIVGMTMYNQATQEIAKPSELLT SVRAYMTVLQSIENYVQIDITRVFNNVLLQQTQHLDSHGEPTITSLYTNWYLETLLR QVSNGHIAYFPAMKAFVNLPTENELTFNAEEYSDISEMRSLSELLGPYGMKFLSESL MWHISSQVAELKKLVVENVDVLTQMRTSFDKPDQMAALFKRLSSVDSVLKRMTIIGV ILSFRSLAQEALRDVLSYHIPFLVSSIEDFKDHIPRETDMKVAMNVYELSSAAGLPC EIDPALVVALSSQKSENISPEEEYKIACLLMVFVAVSLPTLASNVMSQYSPAIEGHC NNIHCLAKAINQIAAALFTIHKGSIEDRLKEFLALASSSLLKIGQETDKTTTRNRES VYLLLDMIVQESPFLTMDLLESCFPYVLLRNAYHAVYKQSVTSSA	(Ismail et al., 2009)
WAVE1²³⁰Δ WCA	MBP-TEV-hWAVE(1) 1-230, MBP tag removed by TEV cleavage	pDK071	MKIEEGKLVIWINGDKGYNGLAEVGKKFEKDTGIKVTVEHPDKLEEKFPQVAATGDG PDIIFWAHDRFGGYAQSGLLAEITPDKAFQDKLYPFTWDAVRYNGKLIAYPIAVEAL SLIYNKDLLPNPPKTWEEIPALDKELKAKGKSALMFNLQEPYFTWPLIAADGGYAFK YENGKYDIKDVGVDNAGAKAGLTFLVDLIKNKHMNADTDYSIAEAAFNKGETAMTIN GPWAWSNIDTSKVNYGVTVLPTFKGQPSKPFVGVLSAGINAASPNKELAKEFLENYL LTDEGLEAVNKDKPLGAVALKSYEEELAKDPRIAATMENAQKGEIMPNIPQMSAFWY AVRTAVINAASGRQTVDEALKDAQTNSSSNNNNNNNNNLGIEGRISEFENLYFQGH MPLVKRNIDPRHLCHTALPRGIKNELECVTNISLANIIRQLSSLSKYAEDIFGELFN EAHSFSFRVNSLQERVDRLSVSVTQLDPKEEELSLQDITMRKAFRSSTIQDQQLFDR KTLPIPLQETYDVCEQPPPLNILTPYRDDGKEGLKFYTNPSYFFDLWKEKMLQDTED KRKEKRKQKQKNLDRPHEPEKVPRAPHDRRREWQKLAQGPELAEDDANLLHKHIEVA NG	(Chen et al., 2017)
WAVE1²³⁰W CA	MBP-TEV-hWAVE(1) 1-230 – GGS6 – VCA, MBP tag removed by TEV cleavage	pDK081	MKIEEGKLVIWINGDKGYNGLAEVGKKFEKDTGIKVTVEHPDKLEEKFPQVAATGDG PDIIFWAHDRFGGYAQSGLLAEITPDKAFQDKLYPFTWDAVRYNGKLIAYPIAVEAL SLIYNKDLLPNPPKTWEEIPALDKELKAKGKSALMFNLQEPYFTWPLIAADGGYAFK YENGKYDIKDVGVDNAGAKAGLTFLVDLIKNKHMNADTDYSIAEAAFNKGETAMTIN GPWAWSNIDTSKVNYGVTVLPTFKGQPSKPFVGVLSAGINAASPNKELAKEFLENYL LTDEGLEAVNKDKPLGAVALKSYEEELAKDPRIAATMENAQKGEIMPNIPQMSAFWY AVRTAVINAASGRQTVDEALKDAQTNSSSNNNNNNNNNLGIEGRISEFENLYFQGH MPLVKRNIDPRHLCHTALPRGIKNELECVTNISLANIIRQLSSLSKYAEDIFGELFN EAHSFSFRVNSLQERVDRLSVSVTQLDPKEEELSLQDITMRKAFRSSTIQDQQLFDR KTLPIPLQETYDVCEQPPPLNILTPYRDDGKEGLKFYTNPSYFFDLWKEKMLQDTED KRKEKRKQKQKNLDRPHEPEKVPRAPHDRRREWQKLAQGPELAEDDANLLHKHIEVA NGGGSGGSGGSGGSGGSGGSKRHPSTLPVISDARSVLLEAIRKGIQLRKVEEQREQE AKHERIENDVATILSRRIAVEYSDSEDDSEFDEVDWLE	(Chen et al., 2017)
Abi2 (1-158)	MBP-TEV-hAbi(2) 1-158, MBP tag removed by TEV cleavage	pDK075	MKIEEGKLVIWINGDKGYNGLAEVGKKFEKDTGIKVTVEHPDKLEEKFPQVAATGDG PDIIFWAHDRFGGYAQSGLLAEITPDKAFQDKLYPFTWDAVRYNGKLIAYPIAVEAL SLIYNKDLLPNPPKTWEEIPALDKELKAKGKSALMFNLQEPYFTWPLIAADGGYAFK YENGKYDIKDVGVDNAGAKAGLTFLVDLIKNKHMNADTDYSIAEAAFNKGETAMTIN GPWAWSNIDTSKVNYGVTVLPTFKGQPSKPFVGVLSAGINAASPNKELAKEFLENYL LTDEGLEAVNKDKPLGAVALKSYEEELAKDPRIAATMENAQKGEIMPNIPQMSAFWY	(Ismail et al., 2009)

			AVRTAVINAASGRQTVDEALKDAQTNSSSNNNNNNNNNNLGIEGRISEFENLYFQGH<u>MAELQMLLEEEIPGGRRALFDSYTNLERVADYCENNYIQSADKQRALEETKAYTTQS</u>LASVAYLINTLANNVLQMLDIQASQLRRMESSINHISQTVDIHKEKVARREIGILTTNKNTSRTHKIIAPANLERPVRYIRKPIDYTILDDIGHGVKVSTQ	
Abi2 (1-158)-sortase	MBP-TEV-hAbi(2) 1-158-LPGTG, MBP tag removed by TEV cleavage	pDK255	MKIEEGKLVIWINGDKGYNGLAEVGKKFEKDTGIKVTVEHPDKLEEKFPQVAATGDGPDIIFWAHDRFGGYAQSGLLAEITPDKAFQDKLYPFTWDAVRYNGKLIAYPIAVEALSLIYNKDLLPNPPKTWEEIPALDKELKAKGKSALMFNLQEPYFTWPLIAADGGYAFKYENGKYDIKDVGVDNAGAKAGLTFLVDLIKNKHMNADTDYSIAEAAFNKGETAMTINGPWAWSNIDTSKVNYGVTVLPTFKGQPSKPFVGVLSAGINAASPNKELAKEFLENYLLTDEGLEAVNKDKPLGAVALKSYEEELAKDPRIAATMENAQKGEIMPNIPQMSAFWYAVRTAVINAASGRQTVDEALKDAQTNSSSNNNNNNNNNNLGIEGRISEFENLYFQGH<u>MAELQMLLEEEIPGGRRALFDSYTNLERVADYCENNYIQSADKQRALEETKAYTTQS</u>LASVAYLINTLANNVLQMLDIQASQLRRMESSINHISQTVDIHKEKVARREIGILTTNKNTSRTHKIIAPANLERPVRYIRKPIDYTILDDIGHGVKVSTQGGLPGTGG	This paper
(MBP)₂-Abi2	2MBP-Thrombin-StrepII-hAbi(2) 1-158	pDK119	MKIEEGKLVIWINGDKGYNGLAEVGKKFEKDTGIKVTVEHPDKLEEKFPQVAATGDGPDIIFWAHDRFGGYAQSGLLAEITPDKAFQDKLYPFTWDAVRYNGKLIAYPIAVEALSLIYNKDLLPNPPKTWEEIPALDKELKAKGKSALMFNLQEPYFTWPLIAADGGYAFKYENGKYDIKDVGVDNAGAKAGLTFLVDLIKNKHMNADTDYSIAEAAFNKGETAMTINGPWAWSNIDTSKVNYGVTVLPTFKGQPSKPFVGVLSAGINAASPNKELAKEFLENYLLTDEGLEAVNKDKPLGAVALKSYEEELAKDPRIAATMENAQKGEIMPNIPQMSAFWYAVRTAVINAASGRQTVDEALKDAQTNSSSLEWSHPQFEKAGGMKIEEGKLVIWINGDKGYNGLAEVGKKFEKDTGIKVTVEHPDKLEEKFPQVAATGDGPDIIFWAHDRFGGYAQSGLLAEITPDKAFQDKLYPFTWDAVRYNGKLIAYPIAVEALSLIYNKDLLPNPPKTWEEIPALDKELKAKGKSALMFNLQEPYFTWPLIAADGGYAFKYENGKYDIKDVGVDNAGAKAGLTFLVDLIKNKHMNADTDYSIAEAAFNKGETAMTINGPWAWSNIDTSKVNYGVTVLPTFKGQPSKPFVGVLSAGINAASPNKELAKEFLENYLLTDEGLEAVNKDKPLGAVALKSYEEELAKDPRIAATMENAQKGEIMPNIPQMSAFWYAVRTAVINAASGRQTVDEALKDAQTNSSSNNNNNNNNNNLGEFLVPRGSWSHPQFEKAGGH<u>MAELQMLLEEEIPGGRRALFDSYTNLERVADYCENNYIQSADKQRALEETKAYTTQS</u>LASVAYLINTLANNVLQMLDIQASQLRRMESSINHISQTVDIHKEKVARREIGILTTNKNTSRTHKIIAPANLERPVRYIRKPIDYTILDDIGHGVKVSTQ	This paper
HSPC300	MBP-TEV-hHSPC300, MBP tag removed by TEV cleavage	pDK069	MKIEEGKLVIWINGDKGYNGLAEVGKKFEKDTGIKVTVEHPDKLEEKFPQVAATGDGPDIIFWAHDRFGGYAQSGLLAEITPDKAFQDKLYPFTWDAVRYNGKLIAYPIAVEALSLIYNKDLLPNPPKTWEEIPALDKELKAKGKSALMFNLQEPYFTWPLIAADGGYAFKYENGKYDIKDVGVDNAGAKAGLTFLVDLIKNKHMNADTDYSIAEAAFNKGETAMTINGPWAWSNIDTSKVNYGVTVLPTFKGQPSKPFVGVLSAGINAASPNKELAKEFLENYLLTDEGLEAVNKDKPLGAVALKSYEEELAKDPRIAATMENAQKGEIMPNIPQMSAFWYAVRTAVINAASGRQTVDEALKDAQTNSSSNNNNNNNNNNLGIEGRISEFENLYFQGH<u>MGAAMAGQEDPVQREIHQDWANREYIEIITSSIKKIADFLNSFDMSCRSRLATLNEK</u>LTALERRIEYIEARVTKGETLT	(Ismail et al., 2009)
(MBP)₂-HSPC300	2MBP-Thrombin-StrepII-hHSPC300	pDK118	MKIEEGKLVIWINGDKGYNGLAEVGKKFEKDTGIKVTVEHPDKLEEKFPQVAATGDGPDIIFWAHDRFGGYAQSGLLAEITPDKAFQDKLYPFTWDAVRYNGKLIAYPIAVEALSLIYNKDLLPNPPKTWEEIPALDKELKAKGKSALMFNLQEPYFTWPLIAADGGYAFKYENGKYDIKDVGVDNAGAKAGLTFLVDLIKNKHMNADTDYSIAEAAFNKGETAMTINGPWAWSNIDTSKVNYGVTVLPTFKGQPSKPFVGVLSAGINAASPNKELAKEFLENYLLTDEGLEAVNKDKPLGAVALKSYEEELAKDPRIAATMENAQKGEIMPNIPQMSAFWYAVRTAVINAASGRQTVDEALKDAQTNSSSLEWSHPQFEKAGGMKIEEGKLVIWINGDKGYNGLAEVGKKFEKDTGIKVTVEHPDKLEEKFPQVAATGDGPDIIFWAHDRFGGYAQSGLLAEITPDKAFQDKLYPFTWDAVRYNGKLIAYPIAVEALSLIYNKDLLPNPPKTWEEIPALDKELKAKGKSALMFNLQEPYFTWPLIAADGGYAFKYENGKYDIKDVGVDNAGAKAGLTFLVDLIKNKHMNADTDYSIAEAAFNKGETAMTINGPWAWSNIDTSKVNYGVTVLPTFKGQPSKPFVGVLSAGINAASPNKELAKEFLENYLLTDEGLEAVNKDKPLGAVALKSYEEELAKDPRIAATMENAQKGEIMPNIPQMSAFWYAVRTAVINAASGRQTVDEALKDAQTNSSSNNNNNNNNNNLGEFLVPRGSWSHPQFEKAGGH<u>MGAAMAGQEDPVQREIHQDWANREYIEIITSSIKKIADFLNSFDMSCRSRLATLNEK</u>LTALERRIEYIEARVTKGETLT	(Chen et al., 2014)
GG-(MBP)₂	MKI-GGS-TEV-GG-2MBP-TEV, N-terminal and C-terminal regions removed after TEV cleavage	pDK256	MKIGGSENLYFQG<u>G</u>GKTEEGKLVIWINGDKGYNGLAEVGKKFEKDTGIKVTVEHPDKLEEKFPQVAATGDG<u>P</u>DIIFWAHDRFGGYAQSGLLAEITPDKAFQDKLYPFTWDAVRYNGKLIAYPIAVEALSLIYNKDLLPNPPKTWEEIPALDKELKAKGKSALMFNLQEPYFTWPLIAADGGYAFKYENGKYDIKDVGVDNAGAKAGLTFLVDLIKNKHMNADTDYSIAEAAFNKGETAMTINGPWAWSNIDTSKVNYGVTVLPTFKGQPSKPFVGVLSAGINAASPNKELAKEFLENYLLTDEGLEAVNKDKPLGAVALKSYEEELAKDPRIAATMENAQKGEIMPNIPQMSAFWYAVRTAVINAASGRQTVDEALKDAQTNSSSNNGGSGGSGGSKTEEGKLVIWINGDKGYNGLAEVGKKFEKDTGIKVTVEHPDKLEEKFPQVAATGDGPDIIFWAHDRFGGYAQSGLLAEITPDKAFQDKLYPFTWDAVRYNGKLIAYPIAVEALSLIYNKDLLPNPPKTWEEIPALDKELKAKGKSALMFNLQEPYFTWPLIAADGGYAFKYENGKYDIKDVGVDNAGAKAGLTFLVDLIKNKHMNADTDYSIAEAAFNKGETAMTINGPWAWSNIDTSKVNYGVTVLPTFKGQPSKPFVGVLSAGINAASPNKELAKEFLENYLLTDEGLEAVNKDKPLGAVALKSYEEELAKDPRIAATMENAQKGEIMPNIPQMSAFWYAVRTAVINAASGRQTVDEALKDAQTNSSSNNNNNNNNNNLGIEGRISEFENLYF<u>Q</u>GHMLEE	This paper

			FGSSRVDLQASLALAVVLQRRDWENPGVTQLNRLAAHPPFASWRNSEEARTDRPSQQ LRSLNGEWQLGCFGG	
Sortase 5M	Sortase 5M-His6	pDK085	MQAKPQIPKDKSKVAGYIEIPDADIKEPVYPGPATREQLNRGVSFAEENESLDDQNI SIAGHTFIDRPNYQFTNLKAAKKGSMVYFKVGNETRKYKMTSIRNVKPTAVEVLDEQ KGKDKQLTLITCDDYNEETGVWETRKIFVATEVKLEHHHHHH	https://ww w.addgene. org/75144/
Rac1^{QP}	hRac1$^{Q61L/P29S}$ Δ4	pDK077	MQAIKCVVVGDGAVGKTCLLISYTTNAFSGEYIPTVFDNYSANVMVDGKPVNLGLWD TAGLEDYDRLRPLSYPQTDVFLICFSLVSPASFENVRAKWYPEVRHHCPNTPIILVG TKLDLRDDKDTIEKLKEKKLTPITYPQGLAMAKEIGAVKYLECSALTQRGLKTVFDE AIRAVLCPPPVKKRKRK	This paper

1049

References

Adamczak, R., Porollo, A., Meller, J., 2004. Accurate prediction of solvent accessibility using neural networks-based regression. Proteins Struct. Funct. Genet. 56, 753–767. https://doi.org/10.1002/prot.20176

Akin, O., Mullins, R.D., 2008. Capping Protein Increases the Rate of Actin-Based Motility by Promoting Filament Nucleation by the Arp2/3 Complex. Cell 133, 841–851. https://doi.org/10.1016/j.cell.2008.04.011

Alekhina, O., Burstein, E., Billadeau, D.D., 2017. Cellular functions of WASP family proteins at a glance. J. Cell Sci. 130, 2235–2241. https://doi.org/10.1242/jcs.199570

Askelöf, P., Guthenberg, C., Jakobson, I., Mannervik, B., 1975. Purification and characterization of two glutathione S-aryltransferase activities from rat liver. Biochem. J. 147, 513–522. https://doi.org/10.1042/bj1470513

Banaszynski, L.A., Liu, C.W., Wandless, T.J., 2005. Characterization of the FKBP·Rapamycin·FRB Ternary Complex. J. Am. Chem. Soc. 127, 4715–4721. https://doi.org/10.1021/ja043277y

Baskakov, I. V, Kumar, R., Srinivasan, G., Ji, Y., Bolen, D.W., Thompson, E.B., 1999. Trimethylamine N-Oxide-induced Cooperative Folding of an Intrinsically Unfolded Transcription-activating Fragment of Human Glucocorticoid Receptor*. J. Biol. Chem. 274, 10693–10696. https://doi.org/https://doi.org/10.1074/jbc.274.16.10693

Begemann, A., Sticht, H., Begtrup, A., Vitobello, A., Faivre, L., Banka, S., Alhaddad, B., Asadollahi, R., Becker, J., Bierhals, T., Brown, K.E., Bruel, A.L., Brunet, T., Carneiro, M., Cremer, K., Day, R., Denommé-Pichon, A.S., Dyment, D.A., Engels, H., Fisher, R., Goh, E.S., Hajianpour, M.J., Haertel, L.R.M., Hauer, N., Hempel, M., Herget, T., Johannsen, J., Kraus, C., Le Guyader, G., Lesca, G., Mau-Them, F.T., McDermott, J.H., McWalter, K., Meyer, P., Õunap, K., Popp, B., Reimand, T., Riedhammer, K.M., Russo, M., Sadleir, L.G., Saenz, M., Schiff, M., Schuler, E., Syrbe, S., Van der Ven, A.T., Verloes, A., Willems, M., Zweier, C., Steindl, K., Zweier, M., Rauch, A., 2021. New insights into the clinical and molecular spectrum of the novel CYFIP2-related neurodevelopmental disorder and impairment of the WRC-mediated actin dynamics. Genet. Med. 23, 543–554. https://doi.org/10.1038/s41436-020-01011-x

Bernhofer, M., Dallago, C., Karl, T., Satagopam, V., Heinzinger, M., Littmann, M., Olenyi, T.,

Qiu, J., Schütze, K., Yachdav, G., Ashkenazy, H., Ben-Tal, N., Bromberg, Y., Goldberg, T., Kajan, L., O'Donoghue, S., Sander, C., Schafferhans, A., Schlessinger, A., Vriend, G., Mirdita, M., Gawron, P., Gu, W., Jarosz, Y., Trefois, C., Steinegger, M., Schneider, R., Rost, B., 2021. PredictProtein - Predicting protein structure and function for 29 years. Nucleic Acids Res. 49, W535–W540. https://doi.org/10.1093/nar/gkab354

Bonneau, R., Tsai, J., Ruczinski, I., Chivian, D., Rohl, C., Strauss, C.E., Baker, D., 2001. Rosetta in CASP4: progress in ab initio protein structure prediction. Proteins Suppl 5, 119–126. https://doi.org/10.1002/prot.1170

Bürgi, J., Xue, B., Uversky, V.N., van der Goot, F.G., 2016. Intrinsic Disorder in Transmembrane Proteins: Roles in Signaling and Topology Prediction. PLoS One 11, e0158594–e0158594. https://doi.org/10.1371/journal.pone.0158594

Burianek, L.E., Soderling, S.H., 2013. Under lock and key: Spatiotemporal regulation of WASP family proteins coordinates separate dynamic cellular processes. Semin. Cell Dev. Biol. 24, 258–266. https://doi.org/10.1016/j.semcdb.2012.12.005

Caldwell, J.E., Heiss, S.G., Mermall, V., Cooper, J.A., 1989. Effects of CapZ, an Actin Capping Protein of Muscle, on the Polymerization of Actin. Biochemistry 28, 8506–8514. https://doi.org/10.1021/bi00447a036

Chaudhari, K., Gorla, M., Chang, C., Kania, A., Bashaw, G.J., 2021. Robo recruitment of the wave regulatory complex plays an essential and conserved role in midline repulsion. Elife 10, 1–35. https://doi.org/10.7554/ELIFE.64474

Chen, B., Brinkmann, K., Chen, Z., Pak, C.W., Liao, Y., Shi, S., Henry, L., Grishin, N. V., Bogdan, S., Rosen, M.K., 2014a. The WAVE regulatory complex links diverse receptors to the actin cytoskeleton. Cell 156, 195–207. https://doi.org/10.1016/j.cell.2013.11.048

Chen, B., Chou, H.-T., Brautigam, C.A., Xing, W., Yang, S., Henry, L., Doolittle, L.K., Walz, T., Rosen, M.K., 2017. Rac1 GTPase activates the WAVE regulatory complex through two distinct binding sites. Elife 6, 1–22. https://doi.org/10.7554/elife.29795

Chen, B., Padrick, S.B., Henry, L., Rosen, M.K., 2014b. Biochemical reconstitution of the WAVE regulatory complex. Methods Enzymol. 540, 55–72. https://doi.org/10.1016/B978-0-12-397924-7.00004-2

Chen, I., Dorr, B.M., Liu, D.R., 2011. A general strategy for the evolution of bond-forming enzymes using yeast display. Proc. Natl. Acad. Sci. U. S. A. 108, 11399–11404.

https://doi.org/10.1073/pnas.1101046108

Chen, Z., Borek, D., Padrick, S.B., Gomez, T.S., Metlagel, Z., Ismail, A.M., Umetani, J.,

Billadeau, D.D., Otwinowski, Z., Rosen, M.K., 2010. Structure and control of the actin

regulatory WAVE complex. Nature 468, 533–538. https://doi.org/10.1038/nature09623

Cheng, P., Poo, M., 2012. Early events in axon/dendrite polarization. Annu. Rev. Neurosci. 35,

181–201. https://doi.org/10.1146/annurev-neuro-061010-113618

Chia, P.H., Chen, B., Li, P., Rosen, M.K., Shen, K., 2014. Local F-actin network links synapse

formation and axon branching. Cell 156, 208–220.

https://doi.org/10.1016/j.cell.2013.12.009

Chou, F.-S., Wang, P.-S., 2016. The Arp2/3 complex is essential at multiple stages of neural

development. Neurogenesis 3, e1261653. https://doi.org/10.1080/23262133.2016.1261653

Conway, O.J., Carrasquillo, M.M., Wang, X., Bredenberg, J.M., Reddy, J.S., Strickland, S.L.,

Younkin, C.S., Burgess, J.D., Allen, M., Lincoln, S.J., Nguyen, T., Malphrus, K.G., Soto,

A.I., Walton, R.L., Boeve, B.F., Petersen, R.C., Lucas, J.A., Ferman, T.J., Cheshire, W.P.,

van Gerpen, J.A., Uitti, R.J., Wszolek, Z.K., Ross, O.A., Dickson, D.W., Graff-Radford,

N.R., Ertekin-Taner, N., 2018. ABI3 and PLCG2 missense variants as risk factors for

neurodegenerative diseases in Caucasians and African Americans. Mol. Neurodegener. 13,

1–12. https://doi.org/10.1186/s13024-018-0289-x

Cooper, J.A., Pollard, T.D., 1985. Effects of Capping Protein on the Kinetics of Actin

Polymerization. Biochemistry 793–799.

Cooper, J.A., Walker, S.B., Pollard, T.D., 1983. Pyrene actin: documentation of the validity of a

sensitive assay for actin polymerization. J. Muscle Res. Cell Motil. 4, 253–262.

https://doi.org/10.1007/BF00712034

Dahl, J.P., Wang-Dunlop, J., Gonzales, C., Goad, M.E.P., Mark, R.J., Kwak, S.P., 2003.

Characterization of the WAVE1 knock-out mouse: Implications for CNS development. J.

Neurosci. 23, 3343–3352. https://doi.org/10.1523/jneurosci.23-08-03343.2003

Ding, B., Yang, S., Schaks, M., Liu, Y., Brown, A., Rottner, K., Chowdhury, S., Chen, B., 2022.

Structures reveal a key mechanism of WAVE Regulatory Complex activation by Rac1

GTPase. BioRxiv [Preprint], May 10. doi: 10.1101/2022.05.10.491380.

Dishman, A.F., Volkman, B.F., 2018. Unfolding the Mysteries of Protein Metamorphosis. ACS

Chem. Biol. 13, 1438–1446. https://doi.org/10.1021/acschembio.8b00276

Dong, X., Liu, O.W., Howell, A.S., Shen, K., 2013. An extracellular adhesion molecule complex patterns dendritic branching and morphogenesis. Cell 155, 296. https://doi.org/10.1016/j.cell.2013.08.059

Doolittle, L.K., Rosen, M.K., Padrick, S.B., 2013a. Measurement and analysis of in vitro actin polymerization. Methods Mol. Biol. 1046, 273–293. https://doi.org/10.1007/978-1-62703-538-5_16

Doolittle, L.K., Rosen, M.K., Padrick, S.B., 2013b. Purification of native Arp2/3 complex from bovine thymus. Methods Mol. Biol. 1046, 231–250. https://doi.org/10.1007/978-1-62703-538-5_14

Drozdetskiy, A., Cole, C., Procter, J., Barton, G.J., 2015. JPred4: a protein secondary structure prediction server. Nucleic Acids Res. 43, W389–W394. https://doi.org/10.1093/nar/gkv332

Dyson, H.J., Wright, P.E., 2002. Coupling of folding and binding for unstructured proteins. Curr. Opin. Struct. Biol. 12, 54–60. https://doi.org/10.1016/S0959-440X(02)00289-0

Eden, S., Rohatgi, R., Podtelejnikov, A. V., Mann, M., Kirschner, M.W., 2002. Mechanism of regulation of WAVE1-induced actin nucleation by Rac1 and Nck. Nature 418, 790–793. https://doi.org/10.1038/nature00859

Evans, R., O'Neill, M., Pritzel, A., Antropova, N., Senior, A., Green, T., Žídek, A., Bates, R., Blackwell, S., Yim, J., Ronneberger, O., Bodenstein, S., Zielinski, M., Bridgland, A., Potapenko, A., Cowie, A., Tunyasuvunakool, K., Jain, R., Clancy, E., Kohli, P., Jumper, J., Hassabis, D., 2022. Protein complex prediction with AlphaFold-Multimer. bioRxiv 2021.10.04.463034. https://doi.org/10.1101/2021.10.04.463034

Fan, L., Lu, Y., Shen, X., Shao, H., Suo, L., Wu, Q., 2018. Alpha protocadherins and Pyk2 kinase regulate cortical neuron migration and cytoskeletal dynamics via rac1 GTPase and WAVE complex in mice. Elife 7, 1–26. https://doi.org/10.7554/eLife.35242

Funk, J., Merino, F., Schaks, M., Rottner, K., Raunser, S., Bieling, P., 2021. A barbed end interference mechanism reveals how capping protein promotes nucleation in branched actin networks. Nat. Commun. 12. https://doi.org/10.1038/s41467-021-25682-5

Gautreau, A., Ho, H. -y. H., Li, J., Steen, H., Gygi, S.P., Kirschner, M.W., 2004. Purification and architecture of the ubiquitous Wave complex. Proc. Natl. Acad. Sci. 101, 4379–4383. https://doi.org/10.1073/pnas.0400628101

Goley, E.D., Welch, M.D., 2006. The ARP2/3 complex: An actin nucleator comes of age. Nat.

Rev. Mol. Cell Biol. 7, 713–726. https://doi.org/10.1038/nrm2026

Greenfield, N.J., 2006. Using circular dichroism spectra to estimate protein secondary structure.

Nat. Protoc. 1, 2876–2890. https://doi.org/10.1038/nprot.2006.202

Hansen, S.D., Zuchero, J.B., Mullins, R.D., 2013. Cytoplasmic Actin: Purification and Single

Molecule Assembly Assays BT - Adhesion Protein Protocols, in: Coutts, A.S. (Ed.), .

Humana Press, Totowa, NJ, pp. 145–170. https://doi.org/10.1007/978-1-62703-538-5_9

Heier, J.A., Dickinson, D.J., Kwiatkowski, A. V, 2017. Measuring Protein Binding to F-actin by

Co-sedimentation. J. Vis. Exp. 55613. https://doi.org/10.3791/55613

Heiss, S.G., Cooper, J.A., 1991. Regulation of CapZ, an Actin Capping Protein of Chicken

Muscle, by Anionic Phospholipids. Biochemistry 30, 8753–8758.

https://doi.org/10.1021/bi00100a006

Hotulainen, P., Hoogenraad, C.C., 2010. Actin in dendritic spines: Connecting dynamics to

function. J. Cell Biol. 189, 619–629. https://doi.org/10.1083/jcb.201003008

Huber, A.H., Stewart, D.B., Laurents, D. V, Nelson, W.J., Weis, W.I., 2001. The cadherin

cytoplasmic domain is unstructured in the absence of beta-catenin. A possible mechanism

for regulating cadherin turnover. J. Biol. Chem. 276, 12301–12309.

https://doi.org/10.1074/jbc.M010377200

Huber, A.H., Weis, W.I., 2001. The structure of the beta-catenin/E-cadherin complex and the

molecular basis of diverse ligand recognition by beta-catenin. Cell 105, 391–402.

https://doi.org/10.1016/s0092-8674(01)00330-0

Huxley, H.E., Brown, W., 1967. The low-angle X-ray diagram of vertebrate striated muscle and

its behaviour during contraction and rigor. J. Mol. Biol. 30, 383–434.

https://doi.org/10.1016/S0022-2836(67)80046-9

Ito, Y., Carss, K.J., Duarte, S.T., Hartley, T., Keren, B., Kurian, M.A., Marey, I., Charles, P.,

Mendonça, C., Nava, C., Pfundt, R., Sanchis-Juan, A., van Bokhoven, H., van Essen, A.,

van Ravenswaaij-Arts, C., Aitman, T., Bennett, D., Caulfield, M., Chinnery, P., Gale, D.,

Koziell, A., Kuijpers, T.W., Laffan, M.A., Maher, E., Markus, H.S., Morrell, N.W.,

Ouwehand, W.H., Perry, D.J., Raymond, F.L., Roberts, I., Smith, K.G.C., Thrasher, A.,

Watkins, H., Williamson, C., Woods, G., Ashford, S., Bradley, J.R., Fletcher, D.,

Hammerton, T., James, R., Kingston, N., Penkett, C.J., Stirrups, K., Veltman, M., Young,

T., Brown, M., Clements-Brod, N., Davis, J., Dewhurst, E., Dolling, H., Erwood, M., Frary,

A., Linger, R., Martin, J.M., Papadia, S., Rehnstrom, K., Stark, H., Allsup, D., Austin, S.,

Bakchoul, T., Bariana, T.K., Bolton-Maggs, P., Chalmers, E., Collins, J., Collins, P., Erber,

W.N., Everington, T., Favier, R., Freson, K., Furie, B., Gattens, M., Gebhart, J., Gomez, K.,

Greene, D., Greinacher, A., Gresele, P., Hart, D., Heemskerk, J.W.M., Henskens, Y.,

Kazmi, R., Keeling, D., Kelly, A.M., Lambert, M.P., Lentaigne, C., Liesner, R., Makris, M.,

Mangles, S., Mathias, M., Millar, C.M., Mumford, A., Nurden, P., Payne, J., Pasi, J.,

Peerlinck, K., Revel-Vilk, S., Richards, M., Rondina, M., Roughley, C., Schulman, S.,

Schulze, H., Scully, M., Sivapalaratnam, S., Stubbs, M., Tait, R.C., Talks, K., Thachil, J.,

Toh, C.H., Turro, E., Van Geet, C., De Vries, M., Warner, T.Q., Watson, H., Westbury, S.,

Furnell, A., Mapeta, R., Rayner-Matthews, P., Simeoni, I., Staines, S., Stephens, J., Watt,

C., Whitehorn, D., Attwood, A., Daugherty, L., Deevi, S.V.V., Halmagyi, C., Hu, F.,

Matser, V., Meacham, S., Megy, K., Shamardina, O., Titterton, C., Tuna, S., Yu, P., von

Ziegenweldt, J., Astle, W., Bleda, M., Carss, K.J., Gräf, S., Haimel, M., Lango-Allen, H.,

Richardson, S., Calleja, P., Rankin, S., Turek, W., Anderson, J., Bryson, C., Carmichael, J.,

McJannet, C., Stock, S., Allen, L., Ambegaonkar, G., Armstrong, R., Arno, G., Bitner-

Glindzicz, M., Brady, A., Canham, N., Chitre, M., Clement, E., Clowes, V., Deegan, P.,

Deshpande, C., Doffinger, R., Firth, H., Flinter, F., French, C., Gardham, A., Ghali, N.,

Gissen, P., Grozeva, D., Henderson, R., Hensiek, A., Holden, S., Holder, M., Holder, S.,

Hurst, J., Josifova, D., Krishnakumar, D., Kurian, M.A., Lees, M., MacLaren, R., Maw, A.,

Mehta, S., Michaelides, M., Moore, A., Murphy, E., Park, S.M., Parker, A., Patch, C.,

Paterson, J., Rankin, J., Reid, E., Rosser, E., Sanchis-Juan, A., Sandford, R., Santra, S.,

Scott, R., Sohal, A., Stein, P., Thomas, E., Thompson, D., Tischkowitz, M., Vogt, J.,

Wakeling, E., Wassmer, E., Webster, A., Ali, Sonia, Ali, Souad, Boggard, H.J., Church, C.,

Coghlan, G., Cookson, V., Corris, P.A., Creaser-Myers, A., DaCosta, R., Dormand, N.,

Eyries, M., Gall, H., Ghataorhe, P.K., Ghio, S., Ghofrani, A., Gibbs, J.S.R., Girerd, B.,

Greenhalgh, A., Hadinnapola, C., Houweling, A.C., Humbert, M., in't Veld, A.H.,

Kennedy, F., Kiely, D.G., Kovacs, G., Lawrie, A., Ross, R.V.M., Machado, R., Masati, L.,

Meehan, S., Moledina, S., Montani, D., Othman, S., Peacock, A.J., Pepke-Zaba, J., Pollock,

V., Polwarth, G., Ranganathan, L., Rhodes, C.J., Rue-Albrecht, K., Schotte, G., Shipley, D.,

Soubrier, F., Southgate, L., Scelsi, L., Suntharalingam, J., Tan, Y., Toshner, M., Treacy,

C.M., Trembath, R., Noordegraaf, A.V., Walker, Sara, Wanjiku, I., Wharton, J., Wilkins,

M., Wort, S.J., Yates, K., Alachkar, H., Antrobus, R., Arumugakani, G., Bacchelli, C.,

Baxendale, H., Bethune, C., Bibi, S., Booth, C., Browning, M., Burns, S., Chandra, A.,

Cooper, N., Davies, S., Devlin, L., Drewe, E., Edgar, D., Egner, W., Ghurye, R., Gilmour,

K., Goddard, S., Gordins, P., Grigoriadou, S., Hackett, S., Hague, R., Harper, L., Hayman,

G., Herwadkar, A., Huissoon, A., Jolles, S., Kelleher, P., Kumararatne, D., Lear, S.,

Longhurst, H., Lorenzo, L., Maimaris, J., Manson, A., McDermott, E., Murng, S.,

Nejentsev, S., Noorani, S., Oksenhendler, E., Ponsford, M., Qasim, W., Quinti, I., Richter,

A., Samarghitean, C., Sargur, R., Savic, S., Seneviratne, S., Sewell, C., Staples, E., Stauss,

H., Thaventhiran, J., Thomas, M., Welch, S., Willcocks, L., Yeatman, N., Yong, P., Ancliff,

P., Babbs, C., Layton, M., Louka, E., McGowan, S., Mead, A., Roy, N., Chambers, J.,

Dixon, P., Estiu, C., Hague, B., Marschall, H.U., Simpson, M., Chong, S., Emmerson, I.,

Ginsberg, L., Gosal, D., Hadden, R., Horvath, R., Mahdi-Rogers, M., Manzur, A., Marshall,

A., Matthews, E., McCarthy, M., Reilly, M., Renton, T., Rice, A., Themistocleous, A.,

Vale, T., Van Zuydam, N., Walker, Suellen, Ormondroyd, L., Hudson, G., Wei, W., Yu

Wai Man, P., Whitworth, J., Afzal, M., Colby, E., Saleem, M., Alavijeh, O.S., Cook, H.T.,

Johnson, S., Levine, A.P., Wong, E.K.S., Tan, R., Boycott, K.M., MacKenzie, A.,

Majewski, J., Brudno, M., Bulman, D., Dyment, D., Boycott, K.M., Kernohan, K.D.,

Dyack, S., Raymond, F.L., 2018. De Novo Truncating Mutations in WASF1 Cause

Intellectual Disability with Seizures. Am. J. Hum. Genet. 103, 144–153.

https://doi.org/10.1016/j.ajhg.2018.06.001

Jan, Y.-N., Jan, L.Y., 2003. The Control of Dendrite Development. Neuron 40, 229–242.

https://doi.org/https://doi.org/10.1016/S0896-6273(03)00631-7

Johannes, L., Pezeshkian, W., Ipsen, J.H., Shillcock, J.C., 2018. Clustering on Membranes:

Fluctuations and More. Trends Cell Biol. 28, 405–415.

https://doi.org/10.1016/j.tcb.2018.01.009

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool,

K., Bates, R., Žídek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S.A.A., Ballard,

A.J., Cowie, A., Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J., Back, T., Petersen, S.,

Reiman, D., Clancy, E., Zielinski, M., Steinegger, M., Pacholska, M., Berghammer, T.,

Bodenstein, S., Silver, D., Vinyals, O., Senior, A.W., Kavukcuoglu, K., Kohli, P., Hassabis,

D., 2021. Highly accurate protein structure prediction with AlphaFold. Nature 596, 583–

589. https://doi.org/10.1038/s41586-021-03819-2

Källberg, M., Wang, H., Wang, S., Peng, J., Wang, Z., Lu, H., Xu, J., 2012. Template-based

protein structure modeling using the RaptorX web server. Nat. Protoc. 7, 1511–1522.

https://doi.org/10.1038/nprot.2012.085

Kirkpatrick, S.L., Goldberg, L.R., Yazdani, N., Babbs, R.K., Wu, J., Reed, E.R., Jenkins, D.F.,

Bolgioni, A.F., Landaverde, K.I., Luttik, K.P., Mitchell, K.S., Kumar, V., Johnson, W.E.,

Mulligan, M.K., Cottone, P., Bryant, C.D., 2017. Cytoplasmic FMR1-Interacting Protein 2

Is a Major Genetic Factor Underlying Binge Eating. Biol. Psychiatry 81, 757–769.

https://doi.org/10.1016/j.biopsych.2016.10.021

Kobayashi, K., Kuroda, S., Fukata, M., Nakamura, T., Nagase, T., Nomura, N., Matsuura, Y.,

Yoshida-Kubomura, N., Iwamatsu, A., Kaibuchi, K., 1998. p140Sra-1 (specifically Rac1-

associated protein) is a novel specific target for Rac1 small GTPase. J. Biol. Chem. 273,

291–295. https://doi.org/10.1074/jbc.273.1.291

Koronakis, V., Hume, P.J., Humphreys, D., Liu, T., Horning, O., Jensen, O.N., McGhie, E.J.,

2011. WAVE regulatory complex activation by cooperating GTPases Arf and Rac1. Proc.

Natl. Acad. Sci. 108, 14449–14454. https://doi.org/10.1073/pnas.1107666108

Kouyama, T., Mihashi, K., 1981. Fluorimetry study of N-(1-pyrenyl)iodoacetamide-labelled F-

actin. Local structural change of actin protomer both on polymerization and on binding of

heavy meromyosin. Eur. J. Biochem. 114, 33–38.

Kramer, D.A., Piper, H.K., Chen, B., 2022. WASP family proteins : Molecular mechanisms and

implications in human disease. Eur. J. Cell Biol. 101.

https://doi.org/10.1016/j.ejcb.2022.151244

Kuhn, J.R., Pollard, T.D., 2005. Real-time measurements of actin filament polymerization by

total internal reflection fluorescence microscopy. Biophys. J. 88, 1387–1402.

https://doi.org/10.1529/biophysj.104.047399

Kumar, V., Kim, K., Joseph, C., Kourrich, S., Yoo, S.-H., Huang, H.C., Vitaterna, M.H., de

Villena, F.P.-M., Churchill, G., Bonci, A., Takahashi, J.S., 2013. C57BL/6N Mutation in

Cytoplasmic FMRP interacting protein 2 Regulates Cocaine Response. Science (80-.). 342,

1508–1512.

Kuzmic, P., 1996. Program DYNAFIT for the analysis of enzyme kinetic data: application to

HIV proteinase. Anal. Biochem. 237, 260–273. https://doi.org/10.1006/abio.1996.0238

Lebensohn, A.M., Kirschner, M.W., 2009. Activation of the WAVE Complex by Coincident

Signals Controls Actin Assembly. Mol. Cell 36, 512–524.

https://doi.org/10.1016/j.molcel.2009.10.024

Lee, N.K., Fok, K.W., White, A., Wilson, N.H., O'Leary, C.J., Cox, H.L., Michael, M., Yap,

A.S., Cooper, H.M., 2016. Neogenin recruitment of the WAVE regulatory complex

maintains adherens junction stability and tension. Nat. Commun. 7.

https://doi.org/10.1038/ncomms11082

Lefebvre, J.L., Sanes, J.R., Kay, J.N., 2015. Development of Dendritic Form and Function.

Annu. Rev. Cell Dev. Biol. 31, 741–777. https://doi.org/10.1146/annurev-cellbio-100913-

013020

Machesky, L.M., Atkinson, S.J., Ampe, C., Vandekerckhove, J., Pollard, T.D., Vandekercldaove,

J., 1994. Purification of a cortical complex containing two unconventional actins from

Acanthamoeba by affinity chromatography on profilin-agarose. J. Cell Biol. 127, 107–115.

https://doi.org/10.1083/jcb.127.1.107

Machesky, L.M., Insall, R.H., 1998. Scar1 and the related Wiskott-Aldrich syndrome protein,

WASP, regulate the actin cytoskeleton through the Arp2/3 complex. Curr. Biol. 8, 1347–

1356. https://doi.org/10.1016/S0960-9822(98)00015-3

McGuffin, L.J., Bryson, K., Jones, D.T., 2000. The PSIPRED protein structure prediction server

. Bioinformatics 16, 404–405. https://doi.org/10.1093/bioinformatics/16.4.404

Minkyung, B., Frank, D., Ivan, A., Justas, D., Sergey, O., Rie, L.G., Jue, W., Qian, C., N., K.L.,

Dustin, S.R., Claudia, M., Hahnbeom, P., Carson, A., R., G.C., Andy, D., H., P.J., V., R.A.,

A., van D.A., C., E.A., J., O.D., Theo, S., Christoph, B., Tea, P.-K., K., R.M., Udit, D., K.,

Y.C., E., B.J., Christopher, G.K., V., G.N., D., A.P., J., R.R., David, B., 2021. Accurate

prediction of protein structures and interactions using a three-track neural network. Science

(80-.). 373, 871–876. https://doi.org/10.1126/science.abj8754

Miyoshi, T., Tsuji, T., Higashida, C., Hertzog, M., Fujita, A., Narumiya, S., Scita, G., Watanabe,

N., 2006. Actin turnover-dependent fast dissociation of capping protein in the dendritic

nucleation actin network: Evidence of frequent filament severing. J. Cell Biol. 175, 947–

955. https://doi.org/10.1083/jcb.200604176

Narvaez-Ortiz, H.Y., Nolen, B.J., 2022. Unconcerted conformational changes in Arp2/3 complex

integrate multiple activating signals to assemble functional actin networks. Curr. Biol. 32,

975-987.e6. https://doi.org/10.1016/j.cub.2022.01.004

Olive, C., Ibanez, L., Farias, F.H.G., Wang, F., John, P., Norton, J.B., Gentsch, J., Morris, J.C.,

Li, Z., Del-aguila, J., Bergmann, K., Bradley, J., Benitez, B.A., Harari, O., Fagan, A.,

Ances, B., Cruchaga, C., Victoria, M., 2020. Examination of the Effect of Rare Variants in

TREM2, ABI3, and PLCG2 in LOAD Through Multiple Phenotypes. J. Alzheimers Dis. 77,

1469–1482. https://doi.org/10.3233/JAD-200019.Examination

Pilpel, Y., Segal, M., 2005. Rapid WAVE dynamics in dendritic spines of cultured hippocampal

neurons is mediated by actin polymerization. J. Neurochem. 95, 1401–1410.

https://doi.org/10.1111/j.1471-4159.2005.03467.x

Polesskaya, A., Boutillon, A., Yang, S., Wang, Y., Romero, S., Liu, Y., Lavielle, M., Vacher, S.,

Schnitzler, A., Molinie, N., Rocques, N., Fokin, A., Guérois, R., Bièche, I., Chen, B.,

David, N.B., Gautreau, A.M., 2021. CYFIP2-containing WAVE complexes inhibit cell

migration by a competition mechanism. bioRxiv 2020.07.02.184655.

Pollard, T.D., 2010. A Guide to Simple and Informative Binding Assays. Mol. Biol. Cell 21,

4061–4067. https://doi.org/10.1091/mbc.e10-08-0683

Pollard, T.D., 2007. Regulation of Actin Filament Assembly by Arp2/3 Complex and Formins.

Annu. Rev. Biophys. Biomol. Struct. 36, 451–477.

https://doi.org/10.1146/annurev.biophys.35.040405.101936

Pollitt, A.Y., Insall, R.H., 2009. WASP and SCAR/WAVE proteins: the drivers of actin

assembly. J. Cell Sci. 122, 2575–2578. https://doi.org/10.1242/jcs.023879

Racz, B., Weinberg, R.J., 2008. Organization of the Arp2/3 Complex in Hippocampal Spines. J.

Neurosci. 28, 5654–5659. https://doi.org/10.1523/jneurosci.0756-08.2008

Rottner, K., Stradal, T.E.B., Chen, B., 2021. WAVE regulatory complex. Curr. Biol. 31, R512–

R517. https://doi.org/10.1016/j.cub.2021.01.086

Scott, E.K., Luo, L., 2001. How do dendrites take their shape? Nat. Neurosci. 4, 359–365.

https://doi.org/10.1038/86006

Shi, R., Kramer, D.A., Chen, B., Shen, K., 2021. A two-step actin polymerization mechanism

drives dendrite branching. Neural Dev. 16, 1–16. https://doi.org/10.1186/s13064-021-

00154-0

Shimojima Yamamoto, K., Yanagishita, T., Yamamoto, H., Miyamoto, Y., Nagata, M., Ishihara,

Y., Miyashita, Y., Asano, Y., Sakata, Y., Yamamoto, T., 2021. Recurrent de novo

pathogenic variant of WASF1 in a Japanese patient with neurodevelopmental disorder with

absent language and variable seizures. Hum. Genome Var. 8, 7–9.

https://doi.org/10.1038/s41439-021-00176-4

Sims, R., Van Der Lee, S.J., Naj, A.C., Bellenguez, C., Badarinarayan, N., Jakobsdottir, J.,

Kunkle, B.W., Boland, A., Raybould, R., Bis, J.C., Martin, E.R., Grenier-Boley, B.,

Heilmann-Heimbach, S., Chouraki, V., Kuzma, A.B., Sleegers, K., Vronskaya, M., Ruiz,

A., Graham, R.R., Olaso, R., Hoffmann, P., Grove, M.L., Vardarajan, B.N., Hiltunen, M.,

Nöthen, M.M., White, C.C., Hamilton-Nelson, K.L., Epelbaum, J., Maier, W., Choi, S.H.,

Beecham, G.W., Dulary, C., Herms, S., Smith, A.G.A. V., Funk, C.C., Derbois, C.,

Forstner, A.J., Ahmad, S., Li, H., Bacq, D., Harold, D., Satizabal, C.L., Valladares, O.,

Squassina, A., Thomas, R., Brody, J.A., Qu, L., Sánchez-Juan, P., Morgan, T., Wolters, F.J.,

Zhao, Y., Garcia, F.S., Denning, N., Fornage, M., Malamon, J., Naranjo, M.C.D., Majounie,

E., Mosley, T.H., Dombroski, B., Wallon, D., Lupton, M.K., Dupuis, J., Whitehead, P.,

Fratiglioni, L., Medway, C., Jian, X., Mukherjee, S., Keller, L., Brown, K., Lin, H.,

Cantwell, L.B., Panza, F., McGuinness, B., Moreno-Grau, S., Burgess, J.D., Solfrizzi, V.,

Proitsi, P., Adams, H.H., Allen, M., Seripa, D., Pastor, P., Cupples, L.A., Price, N.D.,

Hannequin, D., Frank-García, A., Levy, D., Chakrabarty, P., Caffarra, P., Giegling, I.,

Beiser, A.S., Giedraitis, V., Hampel, H., Garcia, M.E., Wang, X., Lannfelt, L., Mecocci, P.,

Eiriksdottir, G., Crane, P.K., Pasquier, F., Boccardi, V., Henández, I., Barber, R.C., Scherer,

M., Tarraga, L., Adams, P.M., Leber, M., Chen, Y., Albert, M.S., Riedel-Heller, S.,

Emilsson, V., Beekly, D., Braae, A., Schmidt, R., Blacker, D., Masullo, C., Schmidt, H.,

Doody, R.S., Spalletta, G., Jr, W.T.L., Fairchild, T.J., Bossù, P., Lopez, O.L., Frosch, M.P.,

Sacchinelli, E., Ghetti, B., Yang, Q., Huebinger, R.M., Jessen, F., Li, S., Kamboh, M.I.,

Morris, J.C., Sotolongo-Grau, O., Katz, M.J., Corcoran, C., Dunstan, M., Braddel, A.,

Thomas, C., Meggy, A., Marshall, R., Gerrish, A., Chapman, J., Aguilar, M., Taylor, S.,

Hill, M., Fairén, M.D., Hodges, A., Vellas, B., Soininen, H., Kloszewska, I., Daniilidou, M.,

Uphill, J., Patel, Y., Hughes, J.T., Lord, J., Turton, J., Hartmann, A.M., Cecchetti, R.,

Fenoglio, C., Serpente, M., Arcaro, M., Caltagirone, C., Orfei, M.D., Ciaramella, A.,

Pichler, S., Mayhaus, M., Gu, W., Lleó, A., Fortea, J., Blesa, R., Barber, I.S., Brookes, K.,

Cupidi, C., Maletta, R.G., Carrell, D., Sorbi, S., Moebus, S., Urbano, M., Pilotto, A.,

Kornhuber, J., Bosco, P., Todd, S., Craig, D., Johnston, J., Gill, M., Lawlor, B., Lynch, A.,

Fox, N.C., Hardy, J., Albin, R.L., Apostolova, L.G., Arnold, S.E., Asthana, S., Atwood,

C.S., Baldwin, C.T., Barnes, L.L., Barral, S., Beach, T.G., Becker, J.T., Bigio, E.H., Bird,

T.D., Boeve, B.F., Bowen, J.D., Boxer, A., Burke, J.R., Burns, J.M., Buxbaum, J.D., Cairns,

N.J., Cao, C., Carlson, C.S., Carlsson, C.M., Carney, R.M., Carrasquillo, M.M., Carroll,

S.L., Diaz, C.C., Chui, H.C., Clark, D.G., Cribbs, D.H., Crocco, E.A., Decarli, C., Dick, M.,

Duara, R., Evans, D.A., Faber, K.M., Fallon, K.B., Fardo, D.W., Farlow, M.R., Ferris, S.,

Foroud, T.M., Galasko, D.R., Gearing, M., Geschwind, D.H., Gilbert, J.R., Graff-Radford,

N.R., Green, R.C., Growdon, J.H., Hamilton, R.L., Harrell, L.E., Honig, L.S., Huentelman,

M.J., Hulette, C.M., Hyman, B.T., Jarvik, G.P., Abner, E., Jin, L.W., Jun, G., Karydas, A.,

Kaye, J.A., Kim, R., Kowall, N.W., Kramer, J.H., Laferla, F.M., Lah, J.J., Leverenz, J.B.,

Levey, A.I., Li, G., Lieberman, A.P., Lunetta, K.L., Lyketsos, C.G., Marson, D.C.,

Martiniuk, F., Mash, D.C., Masliah, E., McCormick, W.C., McCurry, S.M., McDavid,

A.N., McKee, A.C., Mesulam, M., Miller, B.L., Miller, C.A., Miller, J.W., Murrell, J.R.,

Myers, A.J., O'Bryant, S., Olichney, J.M., Pankratz, V.S., Parisi, J.E., Paulson, H.L., Perry,

W., Peskind, E., Pierce, A., Poon, W.W., Potter, H., Quinn, J.F., Raj, A., Raskind, M.,

Reisberg, B., Reitz, C., Ringman, J.M., Roberson, E.D., Rogaeva, E., Rosen, H.J.,

Rosenberg, R.N., Sager, M.A., Saykin, A.J., Schneider, J.A., Schneider, L.S., Seeley,

W.W., Smith, A.G.A. V., Sonnen, J.A., Spina, S., Stern, R.A., Swerdlow, R.H., Tanzi, R.E.,

Thornton-Wells, T.A., Trojanowski, J.Q., Troncoso, J.C., Van Deerlin, V.M., Van Eldik,

L.J., Vinters, H. V., Vonsattel, J.P., Weintraub, S., Welsh-Bohmer, K.A., Wilhelmsen, K.C.,

Williamson, J., Wingo, T.S., Woltjer, R.L., Wright, C.B., Yu, C.E., Yu, L., Garzia, F.,

Golamaully, F., Septier, G., Engelborghs, S., Vandenberghe, R., De Deyn, P.P., Fernadez,

C.M., Benito, Y.A., Thonberg, H., Forsell, C., Lilius, L., Kinhult-Stählbom, A., Kilander,

L., Brundin, R., Concari, L., Helisalmi, S., Koivisto, A.M., Haapasalo, A., Dermecourt, V.,

Fievet, N., Hanon, O., Dufouil, C., Brice, A., Ritchie, K., Dubois, B., Himali, J.J., Keene,

C.D., Tschanz, J., Fitzpatrick, A.L., Kukull, W.A., Norton, M., Aspelund, T., Larson, E.B.,

Munger, R., Rotter, J.I., Lipton, R.B., Bullido, M.J., Hofman, A., Montine, T.J., Coto, E.,

Boerwinkle, E., Petersen, R.C., Alvarez, V., Rivadeneira, F., Reiman, E.M., Gallo, M.,

O'Donnell, C.J., Reisch, J.S., Bruni, A.C., Royall, D.R., Dichgans, M., Sano, M.,

Galimberti, D., St George-Hyslop, P., Scarpini, E., Tsuang, D.W., Mancuso, M.,

Bonuccelli, U., Winslow, A.R., Daniele, A., Wu, C.K., Peters, O., Nacmias, B.,

Riemenschneider, M., Heun, R., Brayne, C., Rubinsztein, D.C., Bras, J., Guerreiro, R., Al-Chalabi, A., Shaw, C.E., Collinge, J., Tsolaki, M., Clarimón, J., Sussams, R., Lovestone, S., O'Donovan, M.C., Owen, M.J., Behrens, T.W., Mead, S., Uitterlinden, A.G., Holmes, C., Cruchaga, C., Ingelsson, M., Bennett, D.A., Powell, J., Golde, T.E., Graff, C., De Jager, P.L., Morgan, K., Ertekin-Taner, N., Combarros, O., Psaty, B.M., Passmore, P., Younkin, S.G., Berr, C., Gudnason, V., Rujescu, D., Dickson, D.W., Dartigues, J.F., Destefano, A.L., Ortega-Cubero, S., Hakonarson, H., Campion, D., Boada, M., Kauwe, J.K., Farrer, L.A., Van Broeckhoven, C., Ikram, M.A., Jones, L., Haines, J.L., Tzourio, C., Tzourio, C., Escott-Price, V., Mayeux, R., Deleuze, J.F., Amin, N., Goate, A.M., Pericak-Vance, M.A., Amouyel, P., Van Duijn, C.M., Ramirez, A., Wang, L.S., Lambert, J.C., Seshadri, S., Williams, J., Schellenberg, G.D., 2017. Rare coding variants in PLCG2, ABI3, and TREM2 implicate microglial-mediated innate immunity in Alzheimer's disease. Nat. Genet. 49, 1373–1384. https://doi.org/10.1038/ng.3916

Soderling, S.H., Guire, E.S., Kaech, S., White, J., Zhang, F., Schutz, K., Langeberg, L.K., Banker, G., Raber, J., Scott, J.D., 2007. A WAVE-1 and WRP Signaling Complex Regulates Spine Density, Synaptic Plasticity, and Memory. J. Neurosci. 27, 355–365. https://doi.org/10.1523/jneurosci.3209-06.2006

Soderling, S.H., Langeberg, L.K., Soderling, J.A., Davee, S.M., Simerly, R., Raber, J., Scott, J.D., H., S.S., K., L.L., A., S.J., M., D.S., Richard, S., Jacob, R., D., S.J., 2003. Loss of WAVE-1 causes sensorimotor retardation and reduced learning and memory in mice. Proc. Natl. Acad. Sci. 100, 1723–1728. https://doi.org/10.1073/pnas.0438033100

Spudich, J.A., Watt, S., 1971. The Regulation of Rabbit Skeletal Muscle Contraction. J. Biol. Chem. 246, 4866–4871. https://doi.org/10.1016/s0021-9258(18)62016-2

Srivastava, S., Macke, E.L., Swanson, L.C., Coulter, D., Klee, E.W., Mullegama, S. V., Xie, Y., Lanpher, B.C., Bedoukian, E.C., Skraban, C.M., Villard, L., Milh, M., Leppert, M.L.O., Cohen, J.S., 2021. Expansion of the genotypic and phenotypic spectrum of wasf1-related neurodevelopmental disorder. Brain Sci. 11, 1–10. https://doi.org/10.3390/brainsci11070931

Stephan, R., Gohl, C., Fleige, A., Klämbt, C., Bogdan, S., 2011. Membrane-targeted WAVE mediates photoreceptor axon targeting in the absence of the WAVE complex in Drosophila. Mol. Biol. Cell 22, 4079–4092. https://doi.org/10.1091/mbc.E11-02-0121

Tahirovic, S., Hellal, F., Neukirchen, D., Hindges, R., Garvalov, B.K., Flynn, K.C., Stradal, T.E., Chrostek-Grashoff, A., Brakebusch, C., Bradke, F., 2010. Rac1 regulates neuronal polarization through the WAVE complex. J. Neurosci. 30, 6930–6943. https://doi.org/10.1523/JNEUROSCI.5395-09.2010

Takenawa, T., Suetsugu, S., 2007. The WASP-WAVE protein network: Connecting the membrane to the cytoskeleton. Nat. Rev. Mol. Cell Biol. 8, 37–48. https://doi.org/10.1038/nrm2069

Tang, L.T., Diaz-Balzac, C.A., Rahman, M., Ramirez-Suarez, N.J., Salzberg, Y., Lázaro-Peña, M.I., Bülow, H.E., 2019. TIAM-1/GEF can shape somatosensory dendrites independently of its GEF activity by regulating F-actin localization. Elife 8, 1–27. https://doi.org/10.7554/elife.38949

Tavosanis, G., 2012. Dendritic structural plasticity. Dev. Neurobiol. 72, 73–86. https://doi.org/10.1002/dneu.20951

Turjanski, A.G., Gutkind, J.S., Best, R.B., Hummer, G., 2008. Binding-Induced Folding of a Natively Unstructured Transcription Factor. PLOS Comput. Biol. 4, e1000060.

Wang, W., Lo, K.W.-H., Kan, H.-M., Fan, J.-S., Zhang, M., 2003. Structure of the monomeric 8-kDa dynein light chain and mechanism of the domain-swapped dimer assembly. J. Biol. Chem. 278, 41491–41499. https://doi.org/10.1074/jbc.M307118200

Wang, X., Jiang, W., Luo, S., Yang, X., Wang, C., Wang, B., Dang, Y., Shen, Y., Ma, D.K., 2021. The C. elegans homolog of human panic-disorder risk gene TMEM132D orchestrates neuronal morphogenesis through the WAVE-regulatory complex. Mol. Brain 14, 54. https://doi.org/10.1186/s13041-021-00767-w

Weeds, A., Maciver, S., 1993. F-actin capping proteins. Curr. Opin. Cell Biol. 5, 63–69. https://doi.org/10.1016/S0955-0674(05)80009-2

Wu, S., Zhang, Y., 2007. LOMETS: a local meta-threading-server for protein structure prediction. Nucleic Acids Res. 35, 3375–3382. https://doi.org/10.1093/nar/gkm251

Xing, G., Li, M., Sun, Y., Rui, M., Zhuang, Y., Lv, H., Han, J., Jia, Z., Xie, W., 2018. Neurexin–neuroligin 1 regulates synaptic morphology and functions via the WAVE regulatory complex in Drosophila neuromuscular junction. Elife 7, 1–23. https://doi.org/10.7554/eLife.30457

Yamazaki, D., Fujiwara, T., Suetsugu, S., Takenawa, T., 2005. A novel function of WAVE in

lamellipodia: WAVE1 is required for stabilization of lamellipodial protrusions during cell

spreading. Genes Cells 10, 381–392. https://doi.org/10.1111/j.1365-2443.2005.00845.x

Yan, Z., Kim, E., Datta, D., Lewis, D.A., Soderling, S.H., 2016. Synaptic Actin Dysregulation, a

Convergent Mechanism of Mental Disorders? J. Neurosci. 36, 11411–11417.

https://doi.org/10.1523/jneurosci.2360-16.2016

Yokoyama, K., Tezuka, T., Kotani, M., Nakazawa, T., Hoshina, N., Shimoda, Y., Kakuta, S.,

Sudo, K., Watanabe, K., Iwakura, Y., Yamamoto, T., 2011. NYAP: A phosphoprotein

family that links PI3K to WAVE1 signalling in neurons. EMBO J. 30, 4739–4754.

https://doi.org/10.1038/emboj.2011.348

Zhao, A., Zhou, R., Gu, Q., Liu, M., Zhang, B., Huang, J., Yang, B., Yao, R., Wang, J., Lv, H.,

Wang, J., Shen, Y., Wang, H., Chen, X., 2021. Trio exome sequencing identified a novel de

novo WASF1 missense variant leading to recurrent site substitution in a Chinese patient

with developmental delay, microcephaly, and early-onset seizures: a mutational hotspot

p.Trp161 and literature review. Clin. Chim. Acta. https://doi.org/10.1016/j.cca.2021.08.030

Zhao, B., Meka, D.P., Scharrenberg, R., König, T., Schwanke, B., Kobler, O., Windhorst, S.,

Kreutz, M.R., Mikhaylova, M., Calderon de Anda, F., 2017. Microtubules Modulate F-actin

Dynamics during Neuronal Polarization. Sci. Rep. 7, 9583. https://doi.org/10.1038/s41598-

017-09832-8

Zhou, P., Wagner, G., 2010. Overcoming the solubility limit with solubility-enhancement tags:

successful applications in biomolecular NMR studies. J. Biomol. NMR 46, 23–31.

https://doi.org/10.1007/s10858-009-9371-6

Zou, W., Dong, X., Broederdorf, T.R., Shen, A., Kramer, D.A., Shi, R., Liang, X., Miller, D.M.,

Xiang, Y.K., Yasuda, R., Chen, B., Shen, K., 2018. A Dendritic Guidance Receptor

Complex Brings Together Distinct Actin Regulators to Drive Efficient F-Actin Assembly

and Branching. Dev. Cell 45, 362-375.e3. https://doi.org/10.1016/j.devcel.2018.04.008

Zou, W., Shen, A., Dong, X., Tugizova, M., Xiang, Y.K., Shen, K., 2016. A multi-protein

receptor-ligand complex underlies combinatorial dendrite guidance choices in C. elegans.

Elife 5, 1–25. https://doi.org/10.7554/elife.18345

Zweier, M., Begemann, A., McWalter, K., Cho, M.T., Abela, L., Banka, S., Behring, B., Berger,

A., Brown, C.W., Carneiro, M., Chen, J., Cooper, G.M., Finnila, C.R., Guillen Sacoto, M.J.,

Henderson, A., Hüffmeier, U., Joset, P., Kerr, B., Lesca, G., Leszinski, G.S., McDermott,

J.H., Meltzer, M.R., Monaghan, K.G., Mostafavi, R., Õunap, K., Plecko, B., Powis, Z.,

Purcarin, G., Reimand, T., Riedhammer, K.M., Schreiber, J.M., Sirsi, D., Wierenga, K.J.,

Wojcik, M.H., Papuc, S.M., Steindl, K., Sticht, H., Rauch, A., 2019. Spatially clustering de

novo variants in CYFIP2, encoding the cytoplasmic FMRP interacting protein 2, cause

intellectual disability and seizures. Eur. J. Hum. Genet. 27, 747–759.

https://doi.org/10.1038/s41431-018-0331-z

1521

1522

	Overlay	1	2	3	4	5
LOMETS						
QUARK						
RaptorX						
Rosetta						
AlphaFold 2						
RoseTTAFold						
AlphaFold 2 (HPO-30 FL)						
RoseTTAFold (HPO-30 FL)						

A

5 representative models   top ranked model

protomer A

protomer B

TSKHAHDVCCTSRKEYREQTKWKNNGLILKTGRVNHQSHRPFVVIDDDSSM

ICD:

Δ1 Δ2 Δ3 Δ4 Δ5 Δ6 Δ7 Δ8 Δ9 Δ10

B

Conformation #1   Conformation #2

5 representative models   top ranked model   5 representative models   top ranked model

β4

protomer A protomer B   protomer A protomer B

membrane

H2   H3

cytosol

C

pH 7.0,   pH 8.0,
50 mM NaCl   100 mM NaCl

MBP-ICD: + +   + +

kDa: 150
100
75

50 ◄ MBP-ICD

◄ GST-ICD

25 ◄ GST

1 2 3 4   5 6 7 8   9

GST Pulldown   Loading Control

A

Buffer: (all pH 7.0 and containing 1 mM MgCl₂, 1 mM EGTA, 1 mM DTT)

	50PDB	50PDB20G	50PDBI	50PDBI20G	**50KMEI20G**	50KMEI5G	50KMEH20G	50KMEH5G
	10 mM HEPES	10 mM HEPES	10 mM Imidazole	10 mM Imidazole	10 mM Imidazole	10 mM Imidazole	10 mM HEPES	10 mM HEPES
	50 mM NaCl	50 mM NaCl	50 mM NaCl	50 mM NaCl	50 mM KCl	50 mM KCl	50 mM KCl	50 mM KCl
	5% glycerol	20% glycerol	5% glycerol	20% glycerol	20% glycerol	5% glycerol	20% glycerol	5% glycerol

Gel lanes labeled 1–32.

Markers (kDa): 150, 100, 75, 50, 37, 25

Protein bands: Sra1, Nap1, GST-ICD, GST

B

Fluorescence assay plot:
- Y-axis: Fluoresence (A.U.), 0 to 100
- X-axis: Time (s), 0 to 1800
- ‑‑‑‑ + DMSO
- —— + Rapamycin
- + WRC + 0.5 μM Rac1 (orange)
- + WRC + 0.4 μM Rac1 (magenta)
- + WRC (purple)
- Actin + Arp2/3 (black)