Novel Mechanism for Tubular Injury in Nephropathic Cystinosis

Swastika Sur[1], Maggie Kerwin[1], Silvia Pineda[1,2], Poonam Sansanwal[1], Tara K. Sigdel[1], Marina Sirota[2] and Minnie M. Sarwal[1]

[1]Division of Multi-Organ Transplantation, Department of Surgery, University of California, San Francisco

[2]Bakar Computational Health Sciences Institute, University of California San Francisco, San Francisco, California, USA.

Corresponding Author: Minnie M. Sarwal, MD, PhD, Medical Director, Kidney and Pancreas Transplant Program, Division of Multi-Organ Transplantation, Department of Surgery, University of California, San Francisco, 513 Parnassus Ave, Med Sciences Bldg, Room S1268, San Francisco, CA 94143 (minnie.sarwal@ucsf.edu)

Running Title: ATP6V0A1 in Nephropathic Cystinosis

Key words: Cystinosis, kidney disease, genetic renal, ESRD, autophagy, CKD

Abstract

Understanding the unique susceptibility of the human kidney to pH dysfunction and injury in cystinosis is paramount to developing new therapies to preserve renal function. Renal proximal tubular epithelial cells (RPTECs) and fibroblasts isolated from patients with cystinosis were transcriptionally profiled. Lysosomal fractionation, immunoblotting, confocal microscopy, intracellular pH, TEM, mitochondrial stress test, and membrane integrity assays were performed for validation. CRISPR, $CTNS^{-/-}$ RPTECs were generated. Alterations in cell stress, pH, autophagic turnover, and mitochondrial energetics highlighted key changes in the vacuolar (V)-ATPases in patient-derived and $CTNS^{-/-}$ RPTECs. ATP6V0A1 was significantly downregulated in cystinosis and highly co-regulated with loss of $CTNS$. Correction of ATP6V0A1 rescued cell stress and mitochondrial function. Treatment of $CTNS^{-/-}$ RPTECs with antioxidants astaxanthin (ATX) induced ATP6V0A1 expression and improved autophagosome turnover and mitochondrial integrity.

In conclusion, our exploratory transcriptional and in vitro cellular and functional studies confirm that loss of cystinosin in RPTECs, results in a reduction in ATP6V0A1 expression, with changes in intracellular pH, mitochondrial integrity, mitochondrial function, and autophagosome-lysosome clearance. The novel findings are ATP6V0A1's role in cystinosis-associated renal pathology and among other antioxidants, ATX specifically upregulated ATP6V0A1, improved autophagosome turnover or reduced autophagy and mitochondrial integrity. This is a pilot study highlighting a novel mechanism of tubular injury in cystinosis and requires further study in animal models to clarify its utility in clinical settings.

INTRODUCTION

Lysosomal storage diseases (LSD) form a significant subgroup of inherited metabolic disorders(1-5), with an incidence of more than 1:5,000 live births (6). Cystinosis is a rare autosomal recessive LSD, caused by mutations in the *CTNS* gene, encoding lysosomal membrane transporter Cystinosin (7). Deletion of 57-kb of the *CTNS* gene is the most common mutation that accounts for approximately 75% of the affected alleles in northern Europe (8, 9). A deficiency of Cystinosin results in lysosomal cystine accumulation and cystine crystal formation in virtually all tissues and organs (7). The kidney is the first organ affected functionally despite lysosomal cystine loading in multiple tissues and organ systems. Based on severity and age at onset of renal injury, cystinosis is phenotypically classified into nephropathic/infantile (OMIM219800) (8), juvenile (OMIM 219900) (9) and ocular (OMIM 219750) (10). The most severe form is nephropathic/infantile cystinosis, characterized by the development of renal Fanconi syndrome and glomerular dysfunction, resulting in end-stage renal disease (ESRD) by 10 years of age, extendable to the second decade of life with cystine depletion therapy (11, 12). Nevertheless, persistent Fanconi and progressive renal failure remains a reality for these patients despite robust adherence to cystine depletion therapies (13). These observations suggest that Cystinosin or other key molecular perturbations impact renal tubular integrity (14-17) and function besides cystine transport (18).

Though associative functional roles for cystinosin have been identified, including inactive mammalian target of rapamycin (mTOR) signaling (19-22), defective autophagy, flawed clearance of damaged mitochondria by disrupted mitophagy (23-25), lysosomal biogenesis (22), abnormal tight junction-associated signaling (26), vesicular transport defects, and increased endoplasmic reticulum (ER) stress (24), the underlying drivers for these injuries have not been well identified, resulting in a paucity of new therapies to improve outcomes in nephropathic cystinosis. Furthermore, we and others have shown that renal biopsies from patients with nephropathic cystinosis reveal morphologically abnormal mitochondria and mitophagy with autophagic vacuoles, reduced mitochondrial numbers (26, 27), increased ROS production (28) and reduced lysosomal enzyme activity (26).

We hypothesized that defective lysosomal clearance may be a pivotal cause of the altered function of renal proximal tubular epithelial cells (RPTECs). Given lysosomes ubiquitous presence in all tissues and organ-systems, we hypothesized that a comparison of transcriptional profiles of renal and extra-renal tissues from patients with nephropathic cystinosis would shed light on the unique lysosomal changes in the human kidney responsible for both the renal tubular injury in Fanconi and the progressive renal tubular injury resulting in ESRD. With this in mind, we investigated whole-genome expression profiles in paired RPTECs (29, 30) (isolated from patient urine samples) and fibroblasts (31) (isolated from the same patient skin biopsies) from patients with cystinosis and healthy age- and gender-matched controls. We identified dysregulation of the vacuolar (V)-ATPase multigene family, specifically ATP6V0A1, only in cystinotic RPTECs and not in normal RPTECs and fibroblasts or cystinotic fibroblasts. To further study the structural and functional consequences of ATP6V0A1 deficiency, a CRISPR/Cas9-mediated *CTNS*-knock-out (*CTNS*$^{-/-}$) immortalized RPTEC line was generated as an *in-vitro* model that mimicked structural and functional changes seen in patient-derived RPTECs. The *CTNS*$^{-/-}$ RPTECs also demonstrated ATP6V0A1 downregulation, resulting in significant functional (loss of acidic lysosomal pH, decreased autophagic flux, compromised mitochondrial ATP-production) and structural consequences (increased vacuolization, lack of mitochondrial cristae, reduced number of mitochondria, ER-stress, and intracellular lipid droplet (LD) formation). Further, correction of ATP6V0A1 in *CTNS*$^{-/-}$ RPTECs and treatment with antioxidants specifically, astaxanthin (ATX) increased the production of cellular ATP6V0A1, identified from a custom FDA-drug database generated by our group, partially rescued the nephropathic RPTEC phenotype (32). ATX is a xanthophyll carotenoid occurring in a wide variety of organisms. ATX is reported to have the highest known antioxidant activity (33) and has proven to have various anti-inflammatory, anti-tumoral, immunomodulatory, anti-cancer, and cytoprotective activities both *in vivo* and *in vitro* (18, 31-37). We are the first to show that ATX can induce the expression of ATP6V0A1 and has a protective effect against cystinosis-induced autophagosome formation and mitochondrial dysfunction. We have used other antioxidants, such as, Cysteamine and Vitamin E, but both had no effect on ATP6V0A1 protein expression. Our findings lead us to believe that ATP6V0A1 is not only a potential target for therapy

in cystinosis, but that ATX has the potential to be repurposed as a ATP6V0A1-inducer and has potential to be used as a combination treatment with cysteamine for the renal pathology in nephropathic cystinosis that persists despite current cystine depletion therapies (14-16).

RESULTS

Figure 1A summarizes the study design. Briefly, the study is divided into three steps. First, transcriptional study with genome microarray in RPTE and fibroblast cells. Second, protein study and functional assays in primary and lab generated RPTECs. Third, correcting the error by using plasmid or inducers to express ATP6V0A1 and checking its effect on the downstream markers. **Figure 1B** represents the schematic overview of the established (black) and hypothesized (blue) mechanisms in normal and cystinotic RPTECs. In summary, we show that compared to healthy, cystinotic RPTECs have dysfunctional *CTNS* that affects the expression of v-ATPases and SLCs, which reduces mTORC1 activity, increases autophagy, compromises mitochondrial function, and causes defective autophago-lysosomal clearance.

Global Transcriptional Changes in Cystinotic RPTEC Affect Lysosomal and Mitochondrial Pathways. Performed genome wide transcriptional profiling of normal and cystinotic RPTECs and skin fibroblasts with and without cystine dimethylester (CDME), used to load lysosomes with cystine to mimic the basic defect in cystinosis (38, 39) (**Figure 2**). The genetic anomaly in cystinosis (mutation of the cystine: proton transporter) is quite heterogeneous with more than 140 identified mutations (18). This heterogeneity affects the phenotype of the disease quite significantly. With this in mind all 8 patients from whom the cells were obtained had the same 57kb mutation (40). Normal CDME loaded RPTECs, and fibroblasts did not transcriptionally mimic the cystinotic phenotype suggesting that CDME loading is a poor surrogate (41) *in vitro* model to study cystinosis tissue injury despite much of earlier research in cystinosis using CDME loading **(Figure 2).** Specific transcriptional signatures are observed in cystinotic skin-fibroblasts and RPTECs obtained from the same individual with cystinosis versus their healthy counterparts (**Figure 2B and C**). These differences between cell types, at the transcriptional level highlight tissue-specific changes

in cystinosis. Nevertheless, some overlapping genes are significantly dysregulated in both cystinotic RPTECs (n=1,926; FDR<0.05) and fibroblasts (n=745; FDR<0.05), with 219 overlapping genes associated with DNA integrity loss and damage. Further analysis identified that certain molecular pathways were highly enriched (42) only in the kidney, and 11 significant pathways were found to be unique to cystinotic RPTECs alone (**Table 1, Supplemental Table S1**). Metabolic pathways, oxidative phosphorylation and acid secretion pathways are some of the significantly affected pathways in cystinosis-RPTECs. Using K-mean clustering on the genes in these significantly enriched pathways, we identified 2 distinct clusters (data not shown). One, is enriched in nucleus-encoded mitochondrial genes crucial for energy production, and the other is enriched in v-ATPases family, which are crucial for lysosomes and kidney tubular acid secretion. Ten lysosomal v-ATPases (**Table 2**) were downregulated in cystinotic RPTECs, five of which are significantly downregulated and some of which play important roles in proximal tubule (PT) H^+ secretion to support reabsorption of luminal HCO_3^-, apical endocytosis (43, 44) and may cause PT acidosis (45). The most significantly perturbed member of the V-ATPase gene family that was found to be downregulated in cystinosis RPTECs is ATP6V0A1 (**Table 2**), hence further attention was focused on characterization of the role of this particular gene in a human in vitro model of cystinosis.

CRISPR-Cas9 Mediated Immortalized *CTNS$^{-/-}$* RPTECs

We and others have found that CDME loading is a poor surrogate *in vitro* model; hence there is a need to develop a more robust *in vitro* model to study the renal tubular injury in cystinosis. Over time, it is likely that the primary cell lines generated from cystinotic patient urine samples may have undergone transformation and the development of a stable, robust in vitro model of renal proximal tubular epithelial cell (RPTEC) injury was recognized as an unmet need in the study of nephropathic cystinosis. Hence, we generated and validated a CRISPR/cas9 based *CTNS* gene knockout model of RPTEC injury. Ribonucleoprotein (RNP)-mediated CRISPR genome editing (**Figure 3A**) was adapted to design a guide RNA that binds to exon 3 of the *CTNS* gene in human immortalized (HuIm) RPTECs and successfully knockout the gene (*CTNS$^{-/-}$*) (**Figure 3B**) with 94% efficiency. We isolated the genomic DNA, PCR

amplified the *CTNS*-region with suitable primers, ran the amplified PCR product in a 2% gel to confirm the primer set that worked best, and finally submitted the amplified DNA with primers to QuintaraBio (SF, CA) for Sanger DNA sequencing. The chromatogram obtained after Sanger-seq was analyzed with TIDE webtool online (**Figure 3A**). As shown in **figure 3B**, TIDE analysis identified an estimated percentage of insertions or deletions (indels) in the *CTNS* gene and showed that the efficiency of the *CTNS* knockout is 94% in the knockout RPTECs compared to the control RPTECs. To confirm the *CTNS* knockout at the mRNA level, we isolated total RNA, prepared cDNA, and performed qPCR (**Figure 3C**). We designed two primers targeting different *CTNS* exons – 2-3 (CTNS#1) and 9-10 (CTNS#2). Since our CRSPR-guide was targeted on exon 3 of the *CTNS* gene, we still could observe some *CTNS* mRNA expression with primer #1; however, *CTNS* expression was undetectable with primer #2. Further, at the functional level, we performed HPLC-MS/MS and showed a significant increase in cystine accumulation in *CTNS*[-/-] RPTECs (**Figure 3D**), similar to intracellular cystine accumulation found in cystinosis patients (5-6 nmol/mg protein).

Knock out of Cystinosin in RPTECs Downregulated ATP6V0A1 Expression

Compared to controls, ATP6V0A1 protein expression was significantly decreased in both primary cystinotic and *CTNS*[-/-] RPTECs with an eight-fold reduction in isolated lysosomal fractions (**Figure 4A-C**). Lysosomes were isolated from control and *CTNS*[-/-] RPTECs, and the purity of the isolation is shown by the presence of lysosomal marker LAMP2 and absence of GAPDH, expressed in cytoplasm (**Figure 4C**). Confocal imaging showed reduced immunopositivity for both LAMP2 (lysosomal marker) and ATP6V0A1 in cystinotic RPTECs (**Figure 4D, Supplementary Figure S2**), with a clear expression of both in normal lysosomes. As shown by other studies (11, 46), reduced immunopositivity for LAMP2

due to its impaired localization caused by defective *CTNS* expression in cystinotic RPTECs. V-ATPases are highly conserved, ubiquitously expressed, and essential for organelle acidification and ATP-driven proton transport. **Figure 4E-F** shows a significant reduction in intracellular acidic pH in cystinotic (pH=5 vs normal RPTEC pH of 4.2) and *CTNS*$^{-/-}$ RPTECs (pH=6.6 vs control human immortalized RPTECs pH of 5.8). An increase of ~0.8 pH units in cystinotic RPTEC translates into a 6.3-fold difference in hydrogen ions, large enough to disturb the cellular function of normal RPTECs.

Reduced Autophagosome Turnover and mTORC1 Activity in Cystinotic RPTECs

By confocal microscopy, we showed increased immunopositivity to LC3B puncta and active or phosphorylated p70S6K in cystinotic RPTECs compared to the control (**Figure 5A and B**). We performed immunoblotting to quantitatively show that both primary and CTNS$^{-/-}$ RPTECs, have reduced phosphorylated and total p70S6 kinase protein expression in the cystinotic RPTECs than its control (**Figure 5C and D**).

Compromised Mitochondrial Function in Cystinotic RPTECs

The cell mitochondria-stress test with Agilent Seahorse XFe instrument assessed mitochondrial (mt) function in *CTNS*$^{-/-}$ and cystinotic RPTECs. Both cystinotic and *CTNS*$^{-/-}$ RPTECs exhibited similar patterns of compromised mitochondrial function (**Figure 6**), with significantly decreased basal, maximal and ATP-linked respiration compared to their respective controls. The oxygen consumption rate (OCR) linked to proton leak, cells spare respiratory capacity and non-mitochondrial oxygen consumption was significantly lost in cystinotic cells. There was a substantial increase in OCR linked to proton leak in normal cells, indicating that normal cells, when injected with mt-electron chain inhibitors, have higher proton leak than the diseased cells with inhibitors. This observation indicates damaged mitochondrial inner membrane in cystinotic cells to begin with, as injecting inhibitors did not make much difference to already compromised mitochondria. The findings suggest that decline in total ATP-linked respiration can be largely attributed to decreased mitochondrial ATP production.

Correction of ATP6V0A1 Expression Improved Autophagosome Turnover and Mitochondrial Function

Since most genes crucial for mitochondrial ATP-generation were downregulated in cystinotic RPTEC, we hypothesized that even without functional cystinosin, correcting ATP6V0A1 expression would positively affect mitochondrial function and autophagosome turnover. We expressed ATP6V0A1via a plasmid in *CTNS*[-/-] RPTECs and observed reduced LC3-II accumulation (**Figure 7A**) indicating either reduced autophagy or increased autophagosome turnover. In addition, study of mitochondrial function (seahorse) revealed increased mitochondrial basal, maximal and ATP-linked respiration - indicating an increased energy demand. ATP6V0A1 also improved non-mitochondrial oxygen consumption. However, the correction had no effect on the proton leak and significantly reduced the spare respiratory capacity (**Figure 7B**). Additionally, we showed that correcting ATP6V0A1 expression with the plasmid expressing the gene in *CTNS*[-/-] cell lines did not have any effect on intracellular cystine level (**Figure 7C**). Overall, ATP6V0A1 correction in *CTNS*[-/-] cells partially improved mitochondrial function compared to the *CTNS*[-/-] RPTECs transfected with control plasmid. These results suggest that ATP6V0A1 plays a critical role in autophagosome turnover and mitochondrial function in *CTNS*[-/-] RPTECs, however, more work is needed to fully understand how ATP6V0A1 regulates these mechanisms in *CTNS*[-/-] RPTEC.

Correction of ATP6V0A1 Rescued Morphologic Renal Tubular Alterations

We performed TEM to characterize cellular morphologic aberrations in *CTNS*[-/-] compared to normal RPTECs and if correcting the ATP6V0A1 expression in *CTNS*[-/-] rescued the diseased RPTEC phenotype (**Figure 7D).** We observed a significant increase in autophagic vacuoles (AV), decrease in mitochondrial number with few or no cristae, swollen ER and the presence of LD within *CTNS*[-/-] RPTECs. Similar to control RPTECs, ATP6V0A1 correction significantly reduced AV (**Figure 7E-G**) and increased mitochondria number with well-preserved cristae. Nevertheless, correction did not affect intracellular LD accumulation.

ATX but Not Other Antioxidants Increases ATP6V0A1 Expression in *CTNS*[-/-] RPTECs

Since there is an increased production of reactive oxygen species in cystinotic renal epithelial cells (22), we evaluated the effect of various antioxidants – Cysteamine, Vitamin E, and ATX. Both RPTECs with and without the *CTNS* were treated with 100 uM of Cysteamine for 24 hours and three different concentrations (20, 50, 100 uM) of vitamin E for 48 hours, and 20 uM of ATX for 48 hours (**Supp.Fig.3**). Both Cysteamine and Vitamin E had no significant effect on the ATP6V0A1 protein expression in *CTNS*[-/-] RPTECs (**Figure 8A, B**). Interestingly, only the ATX pretreatment increased ATP6V0A1 protein expression in *CTNS*[-/-] RPTECs (**Figure 8C**). To evaluate whether ATX could improve autophagic clearance via upregulating ATP6V0A1, LC3-II levels were measured and showed significant reduction of LC3-II with ATX (**Figure 8C**). Similar findings between ATP6V0A1 correction and ATX treatment suggest ATX may ameliorate the impaired autophagy phenotype occurring in cystinotic RPTEC via regulating ATP6V0A1 expression. **Figure 8D** shows significantly increased ROS production in *CTNS*[-/-] RPTEC (p<0.0001) which was rescued with ATX treatment, suggesting ATX's potent antioxidant capacity and ability to improve mitochondrial function. Using the JC-1 probe, we measured mt-membrane potential ($\Delta\Psi$m) since its decrease is a quintessential event in the early stages of apoptosis. Compared to control, *CTNS*[-/-] RPTEC had a significant decrease (p<0.04) in the JC-1 ratio of aggregates (590nm) to monomers (530nm) indicating a significant loss in $\Delta\Psi$m, whereas ATX treatment rescued $\Delta\Psi$m (**Figure 8E**), suggesting ATX's protective role in maintaining mitochondrial integrity in *CTNS*[-/-] RPTEC. MitoSox, a mitochondrial selective ROS detector, was used to measure mitochondrial-mediated ROS production and WST-1 was used to normalize cell viability.

DISCUSSION

Despite whole-body cystine loading, cystinosis predominantly affects the human kidney, causing renal Fanconi syndrome soon after birth and progressive renal damage. These processes are slowed but not evaded by cystine-depleting therapies (11, 26). Over the years, multiple studies have shown some potential

for cysteamine-combined treatments, both in *in-vivo* and *in-vitro* models. But none are yet to be approved by the FDA (14-17). In our study, transcriptional microarray analysis identified that the majority of the genes affected in cystinosis belong to one large cluster, which is crucial for normal lysosomal and mitochondrial function (**Table 1**). We identified a list of vacuolar (v)-ATPases and ATP6V0A1 (**Table 2**) as the most downregulated genes in cystinotic RPTECs, which were also found to have a role in multiple cystinosis-related dysregulated pathways in our dataset (**Table 1 and 2**). Thus, we identified a unique disruption in lysosomal pathways of RPTECs harvested from patients with nephropathic cystinosis. This was not observed in normal RPTECs, nor in paired skin fibroblasts from the same cystinosis patients.

To assess and cross-validate the structural and functional impact of these lysosomal changes in human cystinotic RPTECs we generated a *CTNS⁻/⁻ in vitro,* HuIm RPTEC cystinosis model (16, 47). We found an association between the knocking out of the *CTNS* and the downregulation of v-ATPases. We selected *ATP6V0A1,* the most downregulated v-ATPase, crucial for lysosomal acidification. and investigated its association with the cystinosis phenotype. We aimed to find out if this relationship could explain the early occurrence of lysosomal acidification defect in renal Fanconi syndrome and progressive structural and functional loss of renal reserve. Cystinotic cells are known to have an increased autophagy or reduced autophagosome turnover (19, 20), hence it is essential to determine autophagic flux in any study of cellular cystinosis, which is determined by inhibiting lysosomal degradation with a v-ATPase inhibitor, bafilomycin. However, we found that, even in the absence of bafilomycin, cystinotic RPTECs had LC3B-II accumulation (**Figure 5A**), an autophagosomal marker. Thus, we inferred that lysosomal degradation is innately inhibited in cystinotic RPTECs, and the increment in LC3B-II puncta formation in cystinotic RPTECs could imply an innate compromise of v-ATPase function in these cells. Similar to the study of Andrzejewska et al. (21), we have also observed reduced immunopositivity and expression of phosphorylated-p70S6 kinase protein, a direct downstream substrate of mTORC1 (**Figure 5B-D**) and a common marker for mTORC activity. Our study aligns with others (19-21, 48) who have demonstrated that Cystinosin deficiency is associated with perturbed mTORC1 signaling, which is further reduced with

starvation. In cystinosis, reduction in total p70S6k expression is associated with impaired endocytosis (29, 49), whereas reduction in its phosphorylated form is directly linked to decreased mTORC activity (50). Further, we identified drugs with strong antioxidant activity by using a collated FDA-drug functional database (51): Cysteamine, vitamin E, and ATX were selected from the literature based on their known antioxidant properties. However, ATX was the only agent that upregulated ATP6V0A1 and provided recovery of the specific RPTEC injury phenotype in cystinosis.

Our data indicates that the absence of functional Cystinosin downregulates ATP6V0A1 expression (**Figure 4**), resulting in loss of intracellular acidic pH, decreased autophagy flux, increased autophagosome accumulation (**Figure 7**), loss of mTORC1 activity (**Figure 5**), and a compromised mitochondrial structure and function (**Figure 6 and 7**); moreover, collectively the results suggest a possible mechanistic explanation for the PT dysfunction in cystinosis (**Figure 1B**). Interestingly, all these pathophysiologies known to be compromised in cystinotic RPTECs, and are not recovered by Cysteamine (52, 53).

V-ATPases play a critical role in the acidification of organelles within the endocytic, lysosomal, and secretory pathways and are required for ATP-driven proton transport across membranes (54). Hence, the measurement of lysosomal pH was important. However, none of the available methods to measure lysosomal pH (55-57) worked for cystinotic RPTECs as these cells are too sensitive; therefore, we used BCECF-AM dye, which is a cell-membrane permeable dye that can enter the cell and its organelles. Intracellular pH measured using BCECF-AM represents the overall pH of the cytoplasm and its organelles. We showed that the absence of functional Cystinosin results in less acidic or more basic intracellular-organelle pH in cystinotic RPTECs (**Figure 4E and F**). From published literature, we gathered that an elevation by 0.2-0.3 pH units in lysosomes was linked to a reduction in the pH-dependent cleavage and chronic changes in autophagy and degradation. Re-acidification of the lysosomes with cAMP reversed these above-mentioned changes (51). In yeast and humans, lack of an acidic pH within lysosomes impairs their function of degradation, resulting in autophagosome-filled lysosomes (58, 59) and also reduces pH-dependent amino acid storage in the vacuolar lumen, causing mitochondrial dysfunction (60). Our work

shall be the first to demonstrate, in cystinotic RPTECs, that the absence of Cystinosin affects v-ATPase expression, imbalances acidic pH in the endolysosomal system (**Figure 4**), and disrupts mitochondrial function (**Figure 6**). Upon successful transfection of a plasmid carrying *ATP6V0A1* gene in *CTNS*[-/-] RPTECs the v-ATPase expression levels were corrected (**Figure 7A**). At the functional level, this correction of ATP6V0A1 expression in *CTNS*[-/-] RPTECs reduced LC3-II protein expression, indicating decreased autophagy or improved autophagosome turnover (**Figure 7A**), and partially improved mitochondrial function (**Figure 7B**). But it had no effect on the cellular cystine levels (**Figure 7C**). At the structural level, this correction of ATP6V0A1 expression in *CTNS*[-/-] RPTECs increased the number of mitochondria and improved its structure (**Figure 7E-F**), decreased the number of autophagosome or vacuoles by either improving the autophago-lysosomal clearance or decreasing autophagy (**Figure 7D**).

Alteration in mitochondrial functions (**Figure 6**) likely also results in ER stress or vice versa; ER stress was previously demonstrated in nephropathic cystinosis. Our study revealed increased ER wrapping of mitochondria in *CTNS*[-/-] RPTECs (**Figure 7F**), a phenomenon referred to as mitochondria-ER-associated membranes (MAMs). This observation is known to occur in ER that is stressed (61). MAMs mediate apoptosis, which is known to be higher in nephropathic cystinosis. However, curiously, ATP6V0A1 correction did not rescue ER-wrapping of mitochondria or intracellular LDs (**Figure 7G**) in *CTNS*[-/-] RPTECs. LDs are known to be induced by inflammation and ROS and have been linked with neurodegenerative disorders (52). Previously, LDs have been shown in muscle biopsies from patients with cystinosis (62, 63); however, further studies are needed to understand LD's role in renal pathology in cystinosis. V-ATPases are needed for ATP-driven proton transport across membranes (64), which explains the significant downregulation of several amino acid and metabolite transporters, including SLC17A1, SLC17A3, SLC17A5, SLC3A1 and SLC7A7 (**Supplemental Figure 1**) in cystinotic RPTECs. These findings were a little surprising as there is no clear explanation as to why sialic acid transporters would be affected. Moreover, to gauge the clinical impact of this perturbation we would require an exclusive study.

Another notable finding of our study is the specific effect of antioxidant ATX on $CTNS^{-/-}$ RPTEC, which corrects the ATP6V0A1 levels, enhances autophagosomal turnover, improves cystinosis-induced mitochondrial dysfunction, and rescues mitochondrial membrane potential. The rescue pattern of ATX in $CTNS^{-/-}$ RPTECs is in many ways similar to the metabolic correction of $CTNS$ knock out cells by plasmids bearing ATP6V0A1. This suggests the unique ability of antioxidant ATX to specifically increase ATP6V0A1 expression and, in turn, improve autophagy turnover and mitochondrial function; some of the processes compromised in $CTNS^{-/-}$ RPTECs. However, further studies are warranted to understand how ATX regulates ATP6V0A1 expression and its other protective mechanism in $CTNS^{-/-}$ RPTECs. In various disease models, ATX is shown to regulate mitogen-activated protein kinase (MAPK) by inhibiting JNK1/2 activation (65) (66), affect AMP-activated protein kinase (AMPK) by inhibiting mTOR pathway (67), affect Cerulein mediated increase in LC3 expression—causing reduced apoptosis, autophagy and improved cell potency (68-72)— and impedes inflammation by inhibiting the JAK/STAT3 pathway (73). ATX is also known to ameliorate oxidative stress, ER stress, and mitochondrial dysfunction (74-79). Cumulatively, all these studies support ATX as a promising therapeutic agent for the treatment of a wide variety of diseases. We are currently planning additional *in vivo* experiments to study ATX's effects on lysosomal degradation.

There are a few limitations to our study, which are inherent to any life science studies involving humans, wherein experimental controls cannot be tightly imposed. Firstly, since cysteamine treatment does not reverse Fanconi syndrome or inhibit ESRD, we hypothesized the presence of other confounding genes that could be affected in cystinosis other than $CTNS$. These would be those genes, whose expression or lack of it, cannot be corrected by cystine-depleting therapy. However, there are a few reports of preservation of some renal tubular function when cysteamine was initiated early after birth, but even these patients succumb to ESRD in their second decade (26, 28). Typically, Cysteamine treatment is initiated at one year of age, and maybe, even at this young age, lysosomal cystine-accumulation has started, and at the cellular level irreversible tubular cell damage is progressing to renal Fanconi or ESRD. Second, we also acknowledge that our RPTECs were exposed to CDME for only 30 minutes, after which they began to lose cystine from

their lysosomes because of their normal contingent of Cystinosin (79). Regardless of how long the cells were harvested after loading, the exposure of the cells to increased lysosomal cystine cannot be compared to longstanding exposure of the knock-out cells to lysosomal cystine. This also explains the similarity in transcriptional profiles of CDME-loaded cells to normal and cystinotic cells. Third, we also noted the loss of some epithelial markers in the patient derived RPTECs, which may have occurred with time but is less relevant to this paper as we are looking at a focused pathway in nephropathic cystinosis. We recognized the need for other cell lines that closely mimic the cystinosis-mediated renal pathology; hence, we created the CRISPR-mediated *CTNS*[-/-] RPTECs. Nevertheless, we believe that urine RPTECs harvested from Cystinosis patients and the *CTNS*[-/-] model system are closer mimics of the human disease phenotype and, hence, suit our pursuit—the understanding of molecular injury pathways in cystinotic RPTECs. Fourth, BCECF-AM measures whole cell rather than lysosomal pH, but due to the fragility of the cystinotic RPTEC, direct lysosomal pH measurement was not possible. We acknowledge that additional studies needed to understand the interplay between transcriptional regulation of ATP6V0A1 in cystinosis and if *CTNS*[-/-] cells, chronically treated with cysteamine, acquire any further changes to ATP6V0A1 expression. An important observation in our paper is that the abnormalities present in RPTECs are absent in fibroblasts; we believe that some of these differences are due to the unique changes in v-ATPases, many of which are tissue-specific in expression. A thorough study of other tissues that are also affected by cystinosis, such as the retina, esophagus, skeletal muscle, CNS, and endocrine tissues may provide a more valuable information. It is possible that minor and slower onset of functional perturbations in extra-renal organs is largely due to the intra-cellular cystine accumulation in lysosomes and not due to V-ATPase changes, as seen in the kidney. More specialized and targeted studies are warranted to study V-ATPase as a confounding gene in cystinosis and to confirm it as the pathophysiological variable behind the slow evolution of renal Fanconi and the progression of renal tubular and renal cortical damage, resulting in renal failure by the second decade of life. Understanding this mechanism of injury will be critical also for early detection of kidney damage, as renal injury detection at a functional level is delayed as compensatory mechanisms in

the kidney result in a delayed rise in the serum creatinine, used to calculate the renal reserve by the eGFR formulae (80).

In summary, the novel findings of this study are ATP6V0A1's role in cystinosis-associated renal pathology and, among other antioxidants, ATX specifically upregulated ATP6V0A1, improved autophagosome turnover, reduced autophagy, and secured mitochondrial integrity. Although this is only a pilot study, our finding that ATX, *in vitro,* can ameliorate cystinosis-associated dysfunctional pathways is of paramount importance and requires further study in cystinotic animal models, to clarify its utility in clinical settings.

METHODS

Study design and Samples:

Human RPTECs and fibroblasts were isolated from 8 unique individuals with a biochemically, clinically and genetically confirmed diagnosis of nephropathic cystinosis (provided by Dr. Gahl) (81, 82); in addition, similar cell types, RPTEC (Cambrex Biosciences, East Rutherford, NJ) and fibroblast (Coriell Cell Repositories, Camden, NJ) cell lines were commercially obtained as normal control cells. The patient derived RPTECs has its limitations, which are highlighted in the discussion section. Both normal RPTECs and fibroblasts were treated with cysteine dimethyl ester (CDME) to artificially load lysosomes with cystine, as this method has been used previously as a disease model for cystine RPTEC loading, though it is unclear how accurate this model is for functional analysis of molecular changes in cystinosis patient derived RPTECs (12, 83). In addition, we utilized a human immortalized (HuIm) RPTEC line (Clone TH1, passage 8, Cat No. ECH001, Kerafast, Boston, MA), as control, and used CRISPR-Cas9 to generate a *CTNS*-/- knock out HuIm RPTECs that was then structurally and functionally characterized to evaluate it as an in vitro model system to study human cystinosis RPTEC injury. This purchased HuIm RPTECs are derived from primary human renal proximal tubule epithelial cells (RPTECs) immortalized by two lentiviral vectors carrying the human telomerase and the SV40 T antigen. Details of the study design and its findings are shown in **Figure 1**.

All RPTECs were cultured in renal epithelial growth medium (REGM) (Lonza, Bend, OR); fibroblast cells were cultured in Minimum Essential Media (MEM) with Earl's salts, supplemented with 15% FBS, 2mM L- glutamine, 2X conc. of non-essential AA, 100 µg/ml Penicillin, 100 U/ml Streptomycin and 0.5 µg/ml Fungizone (Invitrogen Corporation, Carlsbad, CA) at 37°C in a 5% CO2 atmosphere. The medium was changed every alternate day, and cultured cells were harvested with 0.05% Trypsin/EDTA (Lonza, Bend, OR) and passaged. All cells were cultured in a 95% air/5% CO2 Thermo Forma incubator (Thermo Scientific, Waltham, MA) at 37°C. All the experiments with cystinotic fibroblasts and RPTECs were performed between passage numbers 2-7, and normal immortalized RPTECs purchased from the company were used even at later passages as long as the cells looked healthy under microscope. For CDME loading, cells were treated with 1mM CDME (Sigma-Aldrich, St. Louis, MO) for 30 min. Cells were pretreated with 20 µM ATX, purchased from Millipore Sigma, Burlington, MA (Cat. No SML0982) for 48 hr. Cysteamine was purchased from Millipore Sigma, Burlington, MA (Cat. No M9768) and cells were pretreated with Cysteamine for 24 hrs. Vitamin E was purchased from Selleckchem, Houston TX (Cat. No S4686) and cells were treated with it for 48 hrs. For each of the treatments, the control well of cells was treated with the same solvent where the compound was dissolved in.

Adaptation of CRISPR-Cas9 Method to Generate *CTNS*^{-/-} RPTECs

Since variations in genetic background and characteristics of the patient and normal RPTECs can result in experimental variations, independent of the *CTNS* mutation, immortalized healthy RPTEC cell lines (Kerafast, Boston, MA) were used to generate isogenic *CTNS*^{-/-} cell lines. CRISPR-Cas9 ribonucleoproteins (crRNPs) were synthesized in vitro by the incubation of *CTNS*-specific guide RNA, trans-activating crRNA (tracrRNA), and Cas9 protein (Dharmacon, Lafayette, CO). These preformed complexes were then delivered to immortalized RPTECs by nucleofection for editing. We used benchling to design the guide RNA to specifically cut at exon 3 (84). Though this is not a known cystinosis-causing mutation but such a knockout causes complete inactivation of cystinosin, which is the common output of all the known mutations associated with nephropathic cystinosis. Since efficiency of gene-knockout varies from one cell

to other, we sorted single cells by FACS and created a pure culture from a single cell. , amplified over the cut-site using touch-down PCR amplification, and then submitted the PCR products for Sanger sequencing using both the forward and reverse TIDE oligos originally used for amplification. Once the chromatograms were returned, we chose to populate and use cells with at least 95% allelic editing, which provided a rough estimate of knock-out percentages (**Figure 3**). Briefly, we cultured these single cells and isolated the DNA from both, the control RPTECs and *CTNS*$^{-/-}$ RPTECs, and then performed PCR amplification over the cut site from their genomic DNA. We designed single-stranded DNA oligos to serve as PCR primers for *CTNS* over the targeted cut site. We then amplified over the cut site using a touch-down PCR amplification strategy with appropriate annealing temperatures for the *CTNS*-specific primers. After this step we submitted the amplified DNA along with both the forward and reverse oligos originally used for amplification the *CTNS* gene for Sanger sequencing. If the nucleofection was successful, then it will be evident in the chromatogram obtained after Sanger. An estimated percentage of Indels was generated by uploading the experimental and control chromatograms to the TIDE webtool online.

Validation of the Generated *CTNS*$^{-/-}$ RPTECs at the Functional level by HPLC-MS/MS

To detect the phenotype of these newly developed *CTNS*$^{-/-}$ cell lines, we performed a functional assay measuring intracellular cystine levels using an HPLC-MS/MS method (UCSD Biochemical Genetics, San Diego). Briefly, we prepared the sample by trypsinizing the adherent cells, washed the cell pellet with ice cold 1mL distilled PBS, centrifuged at 500 g for 5min, resuspended the cell pellet in 150 µl ice cold 650ug/mL N-Ethylmaleimide (NEM; Sigma-Aldrich, St. Louis, MO) in PBS solution, then performed cell-dissociation, followed by adding 50uL of 15% Sulfosalicylic Acid (SSA; Sigma-Aldrich, St. Louis, MO), centrifuged, saved the cell pellet for protein estimation and collect the supernatant separately, bring the volume up to 0.5 ml. The samples were stored at -80°C until transfer (on dry ice) to the UCSD Biochemical Genetics lab. The pellet was resuspended in 0.5 mL 0.1N NaOH to the cell protein pellets to solubilize, pulse vortexed, placed the tube on a rocker with gentle agitation overnight, then protein concentration was calculated by using a standard Pierce BCA Protein Assay Kit (Fisher Scientific, Hampton, NH). The

standards for the BCA assay were resuspended and diluted in 0.1N NaOH instead of water. High levels of intracellular cystine in the generated *CTNS*[-/-] cell line compared to control cells confirmed the successful knockout of the *CTNS* gene. The level of intracellular cystine accumulation in *CTNS*[-/-] cell lines were comparable to cystine levels in cystinosis patients (12, 83) (**Figure 3C**).

Validation of the Generated *CTNS*[-/-] RPTECs at the Transcript level by qPCR

We isolated total RNA from control and *CTNS*[-/-] RPTECs by using RNeasy Mini kit (Cat# 74104) (Qiagen, Hilden, Germany). We followed the standardized protocol provided by the company. The RNA was stored in -80° C for long-term storage. We used 200 μg of the RNA for the complementary-DNA (cDNA) preparation or reverse transcription. Briefly, we added VILO master mix (Thermos Fisher, Waltham, MA) that contains all the reaction components in a pre-mixed formulation and nuclease-free water to the RNA for cDNA synthesis in a thermal cycler (Eppendorf) using lab standardized cDNA synthesis method. The cDNA was stored at 4° C to be used next day for quantitative Polymerase Chain Reaction (qPCR). For qPCR, we designed two primers targeting two specific exons on *CTNS* gene – primer#1 targets between exon 2-3 and primer#2 targets between exon 9-10. These primers were connected to Taqman MGB probe (Thermos Fisher, Waltham, MA). We diluted the cDNA and used 1.25 ng for the qPCR reaction. Briefly, we added master mix (Applied Bioscience, Waltham, MA) that contains all the reaction components, nuclease-free water, and primers to the cDNA and loaded the 384 PCR plate to the thermal cycle (Applied Bioscience, Waltham, MA) using lab standardized qPCR template. Raw Ct data normalized using the delta delta Ct method against 18S and a human universal reference RNA was uploaded into Partek Genomics Suite v.6.6 (Partek Inc., St. Louis, MO, USA). Data were analyzed with Student's t-test to determine any statistically significant differences between groups. All data are presented as mean ± SD. All statistical analyses were performed in Partek Genomics Suite v.6.6., GraphPadPrim v.8. (GraphPad Software Inc.) and in Microsoft Excel (Microsoft, USA).

RNA Isolation for Microarray

Cells were grown until 70-80% confluence and processed for total RNA extraction using RNeasy Midi Kit® (Qiagen Inc., Germantown, MD). Total RNA concentration was measured by NanoDrop® ND-1000 (NanoDrop Technologies, Wilmington, DE) and the integrity of the extracted total RNA was assessed with the Agilent 2100 Bioanalyzer using RNA Nano Chips (Agilent Technologies, Santa Clara, CA). Total RNA was stored at -80°C until preparation for the microarray experiments.

Microarray Experiments to Characterize RPTEC and Fibroblast Transcriptional Profiles in Nephropathic Cystinosis

Hybridization of samples was conducted on Agilent Whole Human Genome 4×44-k 60-mer oligonucleotide arrays (G4112F, Agilent Technologies, Santa Clara, CA), using 150ng of total RNA as template/sample. The arrays were scanned on an Agilent scanner and further processed using Agilent Feature Extraction Software (Agilent Technologies, Santa Clara, CA).

Lysosomal Fractionation

Lysosomal fractions were isolated from cultured cells by density gradient separation using the lysosome enrichment kit (Pierce Biotechnology, Waltham, MA) for both tissue and cultured cells, following the protocol provided by Pierce. Fraction purity was assessed by western blot using lysosome-specific antibody against LAMP2.

Western Blot

Whole cells and lysosomal extracts were prepared, and an equal amount of protein (15ug) was subjected to SDS-PAGE. All primary antibody incubations were done in PBS supplemented with 0.1% Tween-20 (vol/vol) and 5% milk (wt/vol) overnight followed by washing with PBS-Tween (PBS supplemented with 0.1% Tween). The primary antibodies used were: ATP6V0A1 (Cat. No. 109002, Synaptic Systems, Goettingen, Germany), LAMP2 (Cat. No. sc18822, Santa Cruz Biotechnology, Santa Cruz, CA), LC3B (Cat. No. 3868, Cell Signaling, Danvers, MA), phospho-p70 S6 Kinase (Thr389) (Cat. No. MAB S82,

Millipore Sigma, Burlington, MA) GAPDH (Cat. No. 97166S, Cell Signaling, Danvers, MA), beta-tubulin (Cat. No. 2128S, Cell Signaling, Danvers, MA), and p53 (Cat. No. A5761, Abclonal, Woburn, MA). The Peroxidase-conjugated secondary antibodies were diluted 1:2000 in PBS-Tween, incubated with the blot for a minimum of 1 hour at room temperature, and then washed with PBS-Tween and developed using Amersham ECL Plus Detection Reagent (RPN2124, Millipore Sigma, Burlington, MA). Loading levels were normalized using 1:2000 anti-GAPDH or beta-tubulin and anti-LAMP2 Abs. Band quantification was performed using the ImageLab software (National Institutes of Health).

Measurement of pH

To measure lysosomal pH, we tried two methods – 1) pHLARE [**PMID: 33237838**] and 2) LysoSensor [**PMID:**]. But due the fragility of the cystinotic cells, optimum fluorescence level of the biosensor, pHLARE could not be reached. Again, lysoSensor treatment to measure the pH killed the cystinotic RPTECs rapidly even at a very low concentrations. Finally, the conversion of non-fluorescent 2',7'-bis-(2-carboxyethyl)-5-(and-6)-carboxyfluorescein acetoxymethyl ester (BCECF AM) (Invitrogen, Waltham, MA) into a pH sensitive fluorescent indicator by the intracellular esterase was used to measure the pH, which represents an overall cytoplasmic and organelle pH and is not specific to the lysosomes. Briefly, we seeded a fixed number of cells in a clear-bottom black 96-well plate and incubated overnight in a CO_2 incubator for the cells to attach. Next day, we incubated the cells with 2uM of BCECF for 30 min, washed the plate with Hank's Balanced Salt Solution (HBSS) then BCECF fluorescence was measured by using fluorescence microplate reader and pH was calculated. The fluorescence ratio was acquired using the SpectraMax iD3 plate reader (Molecular Devices, San Jose, CA) (excitation = 490, 440 nm; emission = 535 nm). The ratio of BCECF fluorescence at 490/440 nm is a function of pH.

Measurement of mitochondrial oxygen consumption rate

The Agilent Seahorse XFe analyzer allows for real-time measurements of cellular metabolic function in cultured cells. The oxygen consumption rate (OCR) was measured by the extracellular flux analyzer XF24

(Seahorse Bioscience, Santa Clara, CA) following optimization of cell number per well. RPTECs were plated at 4 x 10^5 cells/well in a Seahorse 24-well V7 microplate (Seahorse Bioscience, Santa Clara, CA) and cultured in complete renal epithelial growth medium for 16-18 h in a 5% CO_2 incubator at 37°C. Cells were counted carefully and an equivalent optimum cell density (4x 105 cells/ well) was used to always seed the same number of cells on the same plate. Additionally, background correction wells (ie, wells that have not been seeded with cells) were included in the assay to normalize the data to background plate noise. Prior to the assay, the cells were washed and incubated with assay media (Agilent Technologies, Santa Clara, CA) supplemented with 1 mM glucose (Agilent Technologies, Santa Clara, CA), 1 mM pyruvate (Agilent Technologies), and 2 mM glutamine (Agilent Technologies, Santa Clara, CA) at 37°C without CO_2 for 45 min. Mitochondrial function was measured using Seahorse XF Cell Mito Stress test (Agilent Technologies, Santa Clara, CA). Mitochondrial complex inhibitors (1.5 μM of oligomycin, 0.5 μM of FCCP, 0.5 μM of rotenone and antimycin A) were freshly prepared in XF assay media prior to each experiment and were distributed in ports surrounding the sensor which were sequentially injected to each well.

OCR following serial injection of various probes was used as an indicator of mitochondrial function. Oligomycin, an ATP synthase inhibitor, was utilized as a probe for ATP-linked oxygen consumption; carbonyl cyanide-4-(trifluoromethoxy)phenylhydrazone (FCCP), an oxidative phosphorylation uncoupling agent, was used to induce maximum oxygen consumption and the resultant OCR was used to calculate spare respiratory capacity. A mixture of rotenone and antimycin-A inhibited complex I and complex III was used to result in complete inhibition of mitochondrial respiration and determination of non-mitochondrial oxygen consumption. We compared the pattern observed after injection of each inhibitor in cystinotic RPTECs.

Plasmid-Mediated ATP6V0A1 expression in *CTNS* $^{-/-}$ RPTECs

Myc-DDK-tagged ATP6V0A1 expression plasmid (RC226206, Origene, Rockville, MD) for ATP6V0A1 induction and pCMV6-Entry, mammalian vector with C-terminal Myc-DDK Tag (PS100001, Origene, Rockville, MD) as control were used. Briefly, 3×10^5 *CTNS*^{-/-} and control RPTECs were seeded in six-well plates and incubated for 24 hours prior to transfection. TurboFectin 8.0 Transfection Reagent (F81001, Origene, Rockville, MD) was used at a final concentration of 4 µg/ml for transduction. The correction of ATP6V0A1 expression was verified with western blotting analysis.

Confocal Microscopy

For immunofluorescence, RPTECs were plated in 4-well Chamber Slide with removable wells (ThermoFisher, Waltham, MA), fixed in 100% chilled methanol (5 minutes), permeabilized with PBS containing 0.25% Triton X-100 (10 minutes), and washed three times in PBS. Cells were incubated in 10% normal goat serum blocking solution (Invitrogen, Waltham, MA) for 1 hour followed by overnight primary antibody incubation in a 4°C humidified chamber. For co-immunostaining, we added both primary antibodies raised in different host species at the same time. The next day, slides were washed three times in PBS and followed by secondary antibody incubation in 1% BSA for 1 h at room temperature in the dark. After washing with PBS, the cells were counterstained with DAPI for 5 min and then washed. The 1.5mm thick coverslip was then mounted with a drop of ProLong Glass Antifade Mountant (ThermoFisher, Waltham, MA). Primary antibodies used were: LAMP2 (Santa Cruz Biotechnology, Santa Cruz, CA), ATP6V1B2 (Abcam, Cambridge, United Kingdom), ATP6V0A1 (Synaptic Systems, Goettingen, Germany), LC3B (Cell Signaling Technology, Danvers, MA), phosphor-p70S6 Kinase (Thr398) (Millipore Sigma-Aldrich, St. Louis, MO). Secondary antibodies, donkey anti-Rabbit IgG Alexa Fluor 488 and goat anti-Mouse IgG Alexa Fluor 555 (ThermoFisher, Waltham, MA) were used to detect bound primary antibody. Slides were viewed using a Leica SP5 Confocal Laser Scanning Microscope, and the images were analyzed by Leica Confocal software.

Transmission Electron Microscopy (TEM)

3×10^5 RPTECs were seeded in each well of a six well plate and maintained for 24 h. On the day of the experiment, fresh 2% glutaraldehyde was generated from an 8% stock glutaraldehyde in complete REGM culture media and was added to each well so that cells were fully covered with fixative. After 15 min, the fixed cells were scrapped gently and transferred to a microcentrifuge tube. The samples were centrifuged, and the supernatant was discarded followed by the quick addition of 1ml fresh fixative (2% glutaraldehyde in 0.1M Cacodylate buffer pH 7.2). At this stage, the cells were stored at 4°C and were then handed over to the Electron Microscope Laboratory (EML) imaging core at the University of California, Berkeley for sectioning and imaging. Sections were cut at 80 nm, stained with lead citrate and uranyl acetate, and examined under an FEI Tecnai12 electron microscope (FEI). The electron micrographs obtained from multiple distinct low-powered fields were used to count the number of mitochondria and autophagic vacuoles per cell in at least eight different view fields for each cell culture sample, and the average number of mitochondria or autophagic vacuole per cell culture was calculated.

Mitochondrial Membrane Potential Assay

The mitochondrial membrane potential ($\Delta\Psi$m) was measured with the JC-1 mitochondrial membrane potential assay kit (ab113580, Abcam, Cambridge, United Kingdom) according to manufacturer's instructions. Briefly, RPTEC were seeded at 12,000 cells/well and allowed to adhere overnight in a black clear-bottom 96 well plate. Cells were treated with or without 10 or 20uM ATX for 48hr and then washed once with 1X dilution buffer prior to incubation with 20uM JC-1 dye for 10 min at 37°C. Following incubation, cells were washed twice with 1X dilution buffer and fluorescence intensity was determined for red aggregates (excitation = 535 nm)/emission = 590 nm) and green monomers (excitation = 475 nm/emission = 530 nm) with the SpectraMax iD3 plate reader (Molecular Devices, San Jose, CA). The ratio of JC-1 aggregates (590nm) to JC-1 monomers (530 nm) was calculated. A decrease in aggregate fluorescent count is indicative of depolarization whereas an increase is indicative of hyperpolarization.

Mitochondrial ROS Production and Cell Viability

Mitochondrial ROS production was assessed using MitoSox Red superoxide indicator (ThermoFisher, Waltham, MA). Briefly, cells plated at 10,000 cells/well in 96-well plates were washed with Hanks balanced salt solution (HBSS) (ThermoFisher, Waltham, MA) and treated with 5uM MitoSOX for 15 min at 37C and 5% CO2, protected from light. After staining, cells were washed twice with HBSS to remove background fluorescence. Fluorescence was read (excitation= 510nm, emission= 580nm) with the SpectraMax iD3 plate reader (Molecular Devices, San Jose, CA). Following MitoSox assay, cells were washed with HBSS and WST-1 (Abcam, Cambridge, United Kingdom) was added to the plate and incubated for 30min at 37°C and 5% CO_2 to measure the cell viability. Absorbance was read at 440nm to assess cell viability. Mitochondrial ROS production was normalized to cell viability.

Statistics

Agilent array data were processed and normalized using LOWESS in Gene Spring GX7.3 (Agilent Technologies). The LOWESS normalized data were further analyzed using significance analysis of microarrays (SAM) for two-class unpaired data to detect expression differences based on q-values (<5%)(84). The input for SAM was gene expression measurements from a set of microarray experiments, as well as a response variable from each experiment. SAM used simple median centering of the arrays is an unbiased statistical technique for finding significant genes in a set of microarray experiments. SAM uses repeated permutations of the data to determine whether the expression of each gene is significantly related to the response. Significance levels were set at a q-value of 5%. We used a cutoff of the absolute value of log_2 red channel/green channel >0.5. Data were analyzed using GraphPad Prism software. P-values were calculated using Student's t-test or One-way ANOVA and Tukey' test. Results were expressed as mean ± SD (number of experiments) and considered to be statistically significant when P < 0.05.

MATERIALS AVAILABILITY

We did not use human or animal models, but we have used human cells for this study. Cystinosis RPTE and fibroblast cells are gifts from Dr. Gahl and Dr. Racusen. In addition, the cell lines are commercially

purchased. CRISPR-edited renal cell line is available in Sarwal Lab. Please email: Minnie.sarwal@ucsf.edu or Swastika.sur@ucsf.edu. The study was controlled by institutional review board approvals from the National Institute of Health, Stanford University and the Regents, University of California.

DATA AVAILABILITY

The transcriptomic data was submitted to GEO. The GEO Accession number is GSE190500.

AUTHOR CONTRIBUTIONS

S.S. and M.S. designed the study; S.S., M.K. and P.S. carried out the experiments; S.S., M.K., S.P., M.S. analyzed the data; S.S. and M.K. made the figures; S.S., M.K., T.K.S., M.S. drafted and revised the paper; all authors approved the final version of the manuscript.

ACKNOWLEDGMENTS

We thank Dr. William Gahl and Dr. Lorraine Racusen for the generous gift of cystinosis RPTE and fibroblast cells and Reena Zalpuri at the University of California Berkeley Electron Microscope Laboratory for advice and assistance in electron microscopy sample preparation and data collection. In addition, we would also like to thank Jon Gangoiti at the UCSD Biochemical Genetics Laboratory for advice on sample preparations and performing HPLC-MS/MS method to measure intracellular cystine levels. This work was supported by grants from the Health Research Board, Ireland and The Cystinosis Foundation, Ireland.

TABLES

Table 1: Pathway analysis was performed with all the significantly regulated genes, and 11 pathways were found to be significantly affected in cystinotic RPTECs.

Significantly Enriched Pathways	Adjusted P-value
DNA replication	0.0000
Metabolic pathways	0.0000
Oxidative phosphorylation	0.0014
Cell cycle	0.0014
Fanconi anemia pathway	0.0014
Valine, leucine and isoleucine degradation	0.0049
Collecting duct acid secretion	0.0049
Aminoacyl-tRNA biosynthesis	0.0337
Pyrimidine metabolism	0.0330
Nucleotide excision repair	0.0337
Homologous recombination	0.0337

Table 2: List of all the v-ATPases that are significantly downregulated in cystinosis RPTECs compared to normal.

Gene Name	Fold Change	q-value
ATP6V0A1	0.5867	0.0407
ATP6V1C1	0.5782	0.0000
ATP6V1B2	0.5736	0.0000
ATP6VIE1	0.5489	0.4074
ATP6V1H	0.4616	0.0000
ATP6V0E1	0.6500	0.7087
ATP6V1E2	0.6757	0.2956
ATP6AP1	0.7300	0.4474
ATP6V1A	0.7401	1.4208
ATP6V0D1	0.7987	1.4208

FIGURES

Fig. 1: Overall study design, workflow, and schematic overview. (**A**) Transcriptome profiling with gene microarray performed on cystinotic, normal, and CDME-treated RPTECs and fibroblasts (**Step 1**). Bioinformatic analysis allowed the identification of genes that are differentially regulated in cystinosis versus normal in both the cell types. Pathway analysis was performed with all the significantly regulated genes, and 11 pathways were found to be significantly affected in cystinotic RPTECs. Most of these pathways and the genes in these pathways are crucial for lysosomal acidic pH and mitochondrial ATP production. Dysfunctional mitochondria and compromised lysosomal pH were validated in **step 2**. Our CRISPR-mediated $CTNS^{-/-}$ immortalized renal cell line mimicked cystinosis patient primary cells isolated from their urine. Further, we rescued (**Step 3**) the disease phenotype of the cell by over-expressing ATP6V0A1, the most significantly downregulated gene among other V-ATPases. RNP, Ribonucleoprotein; V-ATPase, vacuolar ATPase; RPTEC, renal proximal tubular epithelial cells. (**B**) A schematic overview of normal and cystinotic RPTECs. We previously showed that loss of function of cystinosin in cystinotic cells inactivates the mTORC pathway, and induces autophagy, mitophagy, and clusterin protein expression. In this study, we show that downregulation of *CTNS* gene also downregulates V-ATPase expression resulting in the loss of lysosomal acidity (pH). The basic lysosomal pH thus blocks its clearance after autophagosome-lysosome fusion. Inhibited autophagy flux due to lysosomal acidity loss may explain why cystinotic cells have increased LC3B-II expression. Loss of acidic lysosomal pH also impairs proton-dependent import of amino acids and other metabolites (noted in our microarray data) into the lysosome lumen, and inhibits conversion of these imported large heterodimeric amino acids into usable form. This increases the presence of metabolites in the cytoplasm that may compromise mitochondrial function and increase ROS production through an unspecified mechanism.

Fig. 2: Microarray study design and transcriptional changes for cystinotic, normal, and CDME-treated RPTECs or fibroblasts. (**A**) Study design: Two types of cells were used- RPTECs and fibroblasts. For each cell type there were three groups – normal that serves as control - commercially obtained, cystinotic – obtained from individuals with cystinosis, and CDME-treated normal cells. (**B**) PCA plot further highlights the unique injury mediated gene profile in cystinotic RPTEC versus normal. Gene expression patterns of cystinotic RPTECs and fibroblasts were found to have no commonality, with cystinostic fibroblasts similar to normal. Gene expression patterns in cystinotic RPTECs are distinct and negatively correlated with normal RPTECs. Whereas gene expression in normal RPTECs laden with CDME are similar to the normal rather than disease phenotype, and hence have a positive correlation with each other. (**C**) Microarray data from both cell types are represented as a correlation heatmap. Cystinotic RPTECs showed markedly different expression profiles from both the normal and CDME loaded normal RPTEC, which was more similar to normal than diseased. Gene expression patterns in normal, cystinotic, and CDME-treated fibroblasts were found to be similar to each other depicting that nephropathic cystinosis-

related   changes   are   highly   specific   to   renal   cells.

Fig. 3: **Experimental overview for the development of immortalized CTNS[-/-] RPTECs using CRISPR-Cas9 RNPs.** (**A**) Outline of RNP CRISPR editing. The *CTNS* gene is located on chromosome 17p13.3 and consists of 12 exons, of which the first 2 are non-coding. Therefore, the guide RNA was targeted towards exon 3 to completely knockout the functional *CTNS* gene. CRISPR-Cas9 ribonucleoproteins (crRNPs) were synthesized in vitro to knockout *CTNS* gene and delivered to immortalize RPTECs by nucleofection. These cells were expanded for molecular validation of gene editing and downstream functional assays. (B)

The guide RNA target sequence and associated PAM are highlighted on the right. We performed Sanger sequencing and the TIDE output calculating percent indels from chromatograms is depicted in the bar-graph. (C) Validation of *CTNS* knockout (-/-) in immortalized RPTEC by using two primers targeting different exons – *CTNS*#1 targets between exon 2-3 and *CTNS*#2 targets between exon 9-10. We have shown that in the CRISPR/Cas9 *CTNS*-/- RPTEC there is a significant reduced cystinosin RNA levels with both the primers. (D) Validation of *CTNS* $^{-/-}$ in immortalized RPTEC. Increased intracellular accumulation of cystine is shown by HPLC-MS/MS method in control and *CTNS*$^{-/-}$ RPTECs.

Fig. 4: **Vacuolar-ATPase, ATP6V0A1, is downregulated in both, cystinotic and *CTNS* -/- RPTECs, further confirming that our in vitro cellular model behaves like primary cystinotic renal cells. This highlights that in addition to the loss of lysosomal cystinosin there are also other key biological changes in the lysosomes in nephropathic cystinosis**. (A-B) Immunoblot analysis of the expression of ATP6V0A1 in cystinotic and *CTNS* $^{-/-}$ RPTECs with their respective controls. (4C) Immunoblot analysis of the expression of ATP6V0A1 in lysosomal fraction isolated from control and *CTNS* $^{-/-}$ RPTECs. Absence

of GAPDH expression depicts successful isolation of pure lysosomes from RPTECs. Results are represented as mean ± SD. The statistically significant differences between the two groups are indicated in the figure. Student's t-test (* *P*≤0.05; ***P*≤0.01; ****P*≤0.001). There was no difference between LAMP2 expression in control and cystinotic RPTECs, making this marker a good control for the experiment. (D) Immunofluorescence and confocal microscopy showing endogenous LAMP2 and ATP6V0A1 distribution in normal and cystinotic RPTECs. Colocalization of ATP6V0A1 and LAMP2 is shown in the right panel. Results are representative of at least 3 separate experiments. Cystinotic and *CTNS* [-/-] RPTECs have an intracellular basic pH compared to their respective controls. Cells were stained with BCECF-AM and measured with a microplate fluorimeter in triplicate wells. The pH-dependent spectral shifts exhibited by BCECF allow calibration of the pH response in terms of the ratio of fluorescence intensities measured at two different excitation wavelengths (490, 440 nm). (E) RPTECs isolated from cystinosis patient urine had a more basic pH than normal individuals (**P≤0.01). (F) Similarly, to the primary cells, our in vitro cellular model, *CTNS*-KO RPTEC, also had a more basic pH than control. The unpaired t-test indicated significant differences (***P≤0.001) between control and diseased cells. Data are representative of three biological replicates. Values represent mean ±SD for three independent experiments.

Fig. 5: Increased immunopositivity to LC3B-II, or autophagy, and decreased immunopositivity to phospho-p70S6 kinase, or mTORC1 activity. (**A**) Immunofluorescence and confocal microscopy showing endogenous LC3B and LAMP2 distribution in normal and cystinotic RPTECs. Colocalization of LC3B and LAMP2 is shown in the right panel. (**B**) Immunofluorescence and confocal microscopy showing endogenous LC3B and phospho-p70S6 kinase (marker of mTOR activity) distribution in normal and cystinotic RPTECs. Colocalization of LC3B and phosphor-p70S6 kinase is shown in the right panel. (**C-D**) Immunoblot analysis of the expression of phosphorylated-p70S6 kinase and total p70S6 kinase in cystinotic and *CTNS* $^{-/-}$ RPTECs with their respective controls. Since there is a change in total p70S6K expression in with and without the *CTNS*, we normalized both, the phosphorylated and total protein, to GAPDH. mTORC1 plays a central role in cell growth, proliferation, survival, and autophagy inhibition via AMPK, therefore, the presence of lower activity of mTORC1 in cystinosis supports the observations of increased cell death and autophagy as shown by increased LC3B puncta. Results are representative of at least 3 separate experiments (biological replicates). Results are represented as mean ± SD. The statistically significant differences between the two groups are indicated in the figure. Student's t-test (* *P*≤0.05; ***P*≤0.01; ****P*≤0.001).

Fig. 6: The Agilent Seahorse Mitochondria Stress Test detected mitochondrial defects in primary cystinotic and *CTNS^{-/-}* RPTECs compared to their respective controls. Top Center) XF Cell Mito Stress Test assay design and standard output parameters. (**A**) XF Cell Mito Stress Test shows that primary cystinotic RPTECs have diminished basal and maximal respiration, ATP-linked respiration, spare respiratory capacity, non-mitochondrial oxygen consumption, and proton leak compared with its control (normal RPTEC) when both were treated with specific electron transport chain (ETC) inhibitors. (**B**) Similar to the primary diseased cells, the *CTNS^{-/-}* also demonstrated diminished mitochondrial activity. Both cystinotic and *CTNS^{-/-}* RPTECs exhibited similar patterns of compromised mitochondrial function, with significantly (Normal vs. Cystinosis: p<0.025, p<0.0019, p<0.023; human immortalized RPTEC (HuIm) vs. *CTNS^{-/-}*: p=0.006, p=0.001, p=0.0002, respectively) decreased basal and maximal respiration and ATP-linked respiration compared to controls. The OCR linked to proton leak and cells spare respiratory capacity was significantly (Normal vs. Cystinosis: p<0.007, p<0.0007; HuIm vs. *CTNS^{-/-}*:

p<0.0001, p=0.002 respectively) lost in cystinotic cells. Non-mitochondrial oxygen consumption did not significantly change in cystinotic or *CTNS*[-/-] RPTECs. There was a substantial (Normal vs. Cystinosis: p<0.007; HuIm vs. *CTNS*[-/-]: p<0.0001) increase in OCR linked to proton leak. Values represent mean ± SD for three independent experiments (* *P*≤0.05; ***P*≤0.01; ****P*≤0.001). OCR, oxygen consumption rate.

Fig. 7: Correction of ATP6V0A1, by transfecting the *CTNS*^{-/-} RPTECs with a Myc-DDK-tagged human ATP6V0A1-plasmid, partially rescued the disease cell phenotype. (**A**) Immunoblot showing successful correction of ATP6V0A1 protein in the *CTNS*^{-/-} cells. Correction of ATP6V0A1 significantly reduced the LC3-II protein expression. (**B**) The XF Cell Mito Stress Test shows that correction of ATP6V0A1 in *CTNS*^{-/-} cells significantly improved basal respiration, and increased maximal respiration and ATP-linked respiration. Overall, ATP6V0A1 correction in *CTNS*^{-/-} cells most closely resemble the normal human immortalized (HuIm) RPTECs compared to *CTNS*^{-/-} or *CTNS*^{-/-} with control plasmid. Electron microscopic evaluation of *CTNS*^{-/-} RPTECs, and its partial recovery by overexpressing ATP6V0A1. (**C**) Correction of ATP6V0A1 in *CTNS*^{-/-} cells had no effect on its cystine level. (**D**) TEM of control, *CTNS*^{-/-}, and ATP6V0A1 overexpressed *CTNS*^{-/-} (*CTNS*^{-/-} +V0A1) RPTECs at low magnification. Asterisks indicate autophagic vacuoles (AV). Number of AV per cell from 8 different sections of each sample was counted and depicted as a bar graph. Significant increase in AV (p=0.006) is observed in *CTNS*^{-/-} compared to that of RPTECs and *CTNS*^{-/-} +V0A1 groups. (**E**) TEM of control, *CTNS*^{-/-}, and *CTNS*^{-/-} +V0A1 RPTECs at low magnification. Asterisks indicate mitochondria. Number of mitochondria per cell from 8 different sections of each sample was counted and depicted as a bar graph. Significant decrease (p=0.0005)

in mitochondria is observed in *CTNS*[-/-] compared to that of RPTECs and *CTNS*[-/-] +V0A1 groups. Both figures A and B indicates rescue of the diseased phenotype by the correction of ATP6V0A1 (p=0.019; p=0.02, respectively). **(F)** Structurally abnormal ER-wrapped mitochondria with lesser cristae was observed in *CTNS*[-/-] compared to that of RPTECs. Though, structurally normal mitochondria with cristae was observed in *CTNS*[-/-] +V0A1 group, but majority of the mitochondria are wrapped by ER. Scale = 0.5um. **(G)** Intracellular LD was observed in more number of cells in *CTNS*[-/-] and *CTNS*[-/-] +V0A1 samples than control RPTECs. LDs have a unique structure, which is delimited by a monolayer of phospholipids differing from the classical bilayer structural organization. Arrows indicate endoplasmic reticulum (ER). Scale = 0.2um. One way ANOVA or t-test as suitable. Values represent mean ± SD for at least three independent experiments (* *P*≤0.05; ***P*≤0.01; ****P*≤0.001).

Fig. 8: Astaxanthin (ATX) but not Cysteamine or Vitamin E upregulates ATP6V0A1 and rescued the cystinosis RPTEC phenotype. (**A**) Immunoblot showing that 100uM of cysteamine treatment for 24 hours did not increase the expression of ATP6V0A1 in *CTNS*[-/-] RPTECs. (**B**) Immunoblots showing VitaminE has no effect on ATP6V0A1 expression in *CTNS*[-/-] RPTECs. (**C**) Immunoblots showing ATX upregulates

ATP6V0A1, reduced the LC3-II (p=0.04) protein expression. Values represent mean ± SD for three independent experiments (* *P*≤0.05; ***P*≤0.01; ****P*≤0.001). (**D**) ATX reduced (p=0.0002) mitochondrial ROS normalized to cell viability measured by WST-1. (**E**) ATX improved mitochondrial membrane potential (p=0.002, p<0.0001 at both 10uM and 20uM, respectively) in *CTNS*[-/-] as shown by the JC1 ratio. Values represent mean ± SD for atleast three independent experiments (* *P*≤0.05; ***P*≤0.01; ****P*≤0.001).

Supplementary Materials:

Supplemental Table S1: List of the 10 v-ATPases showing no significant changes in its expression in cystinosis fibroblasts, CDME-treated fibroblasts and CDME-treated RPTECs compared to their respective controls.

Gene Name	Fold Change (Fibroblasts)	q-value (Fibroblasts)	Fold Change (CDME treated Fibroblast)	q-value (CDME treated Fibroblast)	Fold Change (CDME treated RPTECs)	q-value (CDME treated RPTECs)
ATP6V0A1			-	-	-	-
ATP6V1C1			-	-	-	-
ATP6V1B2	0.39	0.00	-	-	-	-
ATP6VIE1			-	-	-	-
ATP6V1H			-	-	-	-
ATP6V0E1	1.66	0.52	-	-	1.58	0.45
ATP6V1E2			-	-	1.25	2.79
ATP6AP1	0.72	4.50	-	-	-	-
ATP6V1A			-	-	-	-
ATP6V0D1			-	-	-	-

Supplementary Figures:

Supplementary Figure S1: Amino Acids & Sialic Acid Transporters. Microarray identified additional transcriptional changes in amino acid and sialic acid transporters in cystinotic RPTEC. Significant downregulation of SLC17A1, SLC17A3, SLC17A5, SLC3A1, and SLC7A7. (* $P \leq 0.05$; **$P \leq 0.01$; ***$P \leq 0.001$).

Supplementary Figure S2: Immunofluorescence and confocal microscopy showing endogenous LAMP2 and ATP6V0A1 distribution in human immortalized RPTEC and CTNS$^{-/-}$ RPTECs. Colocalization of ATP6V0A1 and LAMP2 is shown in the right panel. Decreased immunopositivity to ATP6V0A1 is noted

in CTNS-/- RPTECs compared to its control. The pattern is similar to primary RPTECs isolated from individuals with cystinosis.

Supplementary Figure S3: ATX Dose Response Curve. We used three concentrations of ATX 10, 20, 50 uM out of which 50 uM was toxic to RPTECs as it killed the cells. However, ATX improved mitochondrial membrane potential (p=0.002, p<0.0001 at both 10uM and 20uM, respectively) in CTNS$^{-/-}$ as shown by the JC1 ratio. Values represent mean ± SD for at least three independent experiments (* *P*≤0.05; ***P*≤0.01; ****P*≤0.001).

References

1.   Anikster Y, Shotelersuk V, Gahl WA. CTNS mutations in patients with cystinosis. Human mutation. 1999;14(6):454-8. PubMed PMID: 10571941.

2.   Fuller M, Meikle PJ, Hopwood JJ. Epidemiology of lysosomal storage diseases: an overview. Fabry disease: perspectives from 5 years of FOS: Oxford PharmaGenesis; 2006.

3.   Touchman JW, Anikster Y, Dietrich NL, Maduro VV, McDowell G, Shotelersuk V, et al. The genomic region encompassing the nephropathic cystinosis gene (CTNS): complete sequencing of a 200-kb

segment and discovery of a novel gene within the common cystinosis-causing deletion. Genome Res. 2000 Feb;10(2):165-73. PubMed PMID: 10673275. Pubmed Central PMCID: PMC310836. Epub 2000/02/15.

4.   Burton BK. Inborn errors of metabolism in infancy: a guide to diagnosis. Pediatrics. 1998 Dec;102(6):E69. PubMed PMID: 9832597.

5.   Ballabio A, Gieselmann V. Lysosomal disorders: from storage to cellular damage. Biochim Biophys Acta. 2009 Apr;1793(4):684-96. PubMed PMID: 19111581.

6.   Gholami Yarahmadi S, Sarlaki F, Morovvati S. Cystinosis and two rare mutations in CTNS gene: two case reports. J Med Case Rep. 2022 May 6;16(1):181. PubMed PMID: 35513889. Pubmed Central PMCID: PMC9074344. Epub 2022/05/06.

7.   Kalatzis V, Cherqui S, Antignac C, Gasnier B. Cystinosin, the protein defective in cystinosis, is a H(+)-driven lysosomal cystine transporter. The EMBO journal. 2001 Nov 1;20(21):5940-9. PubMed PMID: 11689434. Pubmed Central PMCID: 125690.

8.   Town M, Jean G, Cherqui S, Attard M, Forestier L, Whitmore SA, et al. A novel gene encoding an integral membrane protein is mutated in nephropathic cystinosis. Nature genetics. 1998 Apr;18(4):319-24. PubMed PMID: 9537412.

9.   Attard M, Jean G, Forestier L, Cherqui S, van't Hoff W, Broyer M, et al. Severity of phenotype in cystinosis varies with mutations in the CTNS gene: predicted effect on the model of cystinosin. Human molecular genetics. 1999 Dec;8(13):2507-14. PubMed PMID: 10556299.

10.   Anikster Y, Lucero C, Guo J, Huizing M, Shotelersuk V, Bernardini I, et al. Ocular nonnephropathic cystinosis: clinical, biochemical, and molecular correlations. Pediatric research. 2000 Jan;47(1):17-23. PubMed PMID: 10625078.

11.   Cherqui S, Courtoy PJ. The renal Fanconi syndrome in cystinosis: pathogenic insights and therapeutic perspectives. Nature reviews Nephrology. 2017 Feb;13(2):115-31. PubMed PMID: 27990015. Pubmed Central PMCID: 5657490.

12.	Elmonem MA, Veys KR, Soliman NA, van Dyck M, van den Heuvel LP, Levtchenko E. Cystinosis: a review. Orphanet journal of rare diseases. 2016 Apr 22;11:47. PubMed PMID: 27102039. Pubmed Central PMCID: 4841061.

13.	Wilmer MJ, Schoeber JP, van den Heuvel LP, Levtchenko EN. Cystinosis: practical tools for diagnosis and treatment. Pediatric nephrology. 2011 Feb;26(2):205-15. PubMed PMID: 20734088. Pubmed Central PMCID: 3016220.

14.	Bellomo F, De Leo E, Taranta A, Giaquinto L, Di Giovamberardino G, Montefusco S, et al. Drug Repurposing in Rare Diseases: An Integrative Study of Drug Screening and Transcriptomic Analysis in Nephropathic Cystinosis. International journal of molecular sciences. 2021 Nov 27;22(23). PubMed PMID: 34884638. Pubmed Central PMCID: 8657658.

15.	Hollywood JA, Przepiorski A, D'Souza RF, Sreebhavan S, Wolvetang EJ, Harrison PT, et al. Use of Human Induced Pluripotent Stem Cells and Kidney Organoids To Develop a Cysteamine/mTOR Inhibition Combination Therapy for Cystinosis. Journal of the American Society of Nephrology : JASN. 2020 May;31(5):962-82. PubMed PMID: 32198276. Pubmed Central PMCID: 7217405.

16.	Jamalpoor A, van Gelder CA, Yousef Yengej FA, Zaal EA, Berlingerio SP, Veys KR, et al. Cysteamine-bicalutamide combination therapy corrects proximal tubule phenotype in cystinosis. EMBO molecular medicine. 2021 Jul 7;13(7):e13067. PubMed PMID: 34165243. Pubmed Central PMCID: 8261496.

17.	De Leo E, Elmonem MA, Berlingerio SP, Berquez M, Festa BP, Raso R, et al. Cell-Based Phenotypic Drug Screening Identifies Luteolin as Candidate Therapeutic for Nephropathic Cystinosis. Journal of the American Society of Nephrology : JASN. 2020 Jul;31(7):1522-37. PubMed PMID: 32503896. Pubmed Central PMCID: 7351012.

18.	David D, Princiero Berlingerio S, Elmonem MA, Oliveira Arcolino F, Soliman N, van den Heuvel B, et al. Molecular Basis of Cystinosis: Geographic Distribution, Functional Consequences of Mutations in the CTNS Gene, and Potential for Repair. Nephron. 2019;141(2):133-46. PubMed PMID: 30554218.

19.   Ivanova EA, van den Heuvel LP, Elmonem MA, De Smedt H, Missiaen L, Pastore A, et al. Altered mTOR signalling in nephropathic cystinosis. Journal of inherited metabolic disease. 2016 May;39(3):457-64. PubMed PMID: 26909499.

20.   Nevo N, Thomas L, Chhuon C, Andrzejewska Z, Lipecka J, Guillonneau F, et al. Impact of Cystinosin Glycosylation on Protein Stability by Differential Dynamic Stable Isotope Labeling by Amino Acids in Cell Culture (SILAC). Molecular & cellular proteomics : MCP. 2017 Mar;16(3):457-68. PubMed PMID: 28082515. Pubmed Central PMCID: 5341006.

21.   Sansanwal P, Sarwal MM. p62/SQSTM1 prominently accumulates in renal proximal tubules in nephropathic cystinosis. Pediatric nephrology. 2012 Nov;27(11):2137-44. PubMed PMID: 22714671.

22.   Sansanwal P, Sarwal MM. Abnormal mitochondrial autophagy in nephropathic cystinosis. Autophagy. 2010 Oct;6(7):971-3. PubMed PMID: 20729635. Pubmed Central PMCID: 3039743.

23.   Zhang W, Li X, Wang S, Chen Y, Liu H. Regulation of TFEB activity and its potential as a therapeutic target against kidney diseases. Cell death discovery. 2020;6:32. PubMed PMID: 32377395. Pubmed Central PMCID: 7195473.

24.   Luciani A, Festa BP, Chen Z, Devuyst O. Defective autophagy degradation and abnormal tight junction-associated signaling drive epithelial dysfunction in cystinosis. Autophagy. 2018;14(7):1157-9. PubMed PMID: 29806776. Pubmed Central PMCID: 6103718.

25.   Johnson JL, Napolitano G, Monfregola J, Rocca CJ, Cherqui S, Catz SD. Upregulation of the Rab27a-dependent trafficking and secretory mechanisms improves lysosomal transport, alleviates endoplasmic reticulum stress, and reduces lysosome overload in cystinosis. Molecular and cellular biology. 2013 Aug;33(15):2950-62. PubMed PMID: 23716592. Pubmed Central PMCID: 3719680.

26.   Sansanwal P, Yen B, Gahl WA, Ma Y, Ying L, Wong LJ, et al. Mitochondrial autophagy promotes cellular injury in nephropathic cystinosis. Journal of the American Society of Nephrology : JASN. 2010 Feb;21(2):272-83. PubMed PMID: 19959713. Pubmed Central PMCID: 2834547.

27.   Raggi C, Luciani A, Nevo N, Antignac C, Terryn S, Devuyst O. Dedifferentiation and aberrations of the endolysosomal compartment characterize the early stage of nephropathic cystinosis. Human molecular genetics. 2014 May 1;23(9):2266-78. PubMed PMID: 24319100.

28.   Sansanwal P, Li L, Sarwal MM. Inhibition of intracellular clusterin attenuates cell death in nephropathic cystinosis. Journal of the American Society of Nephrology : JASN. 2015 Mar;26(3):612-25. PubMed PMID: 25071085. Pubmed Central PMCID: 4341467.

29.   Le BL, Iwatani S, Wong RJ, Stevenson DK, Sirota M. Computational discovery of therapeutic candidates for preventing preterm birth. JCI insight. 2020 Feb 13;5(3). PubMed PMID: 32051340. Pubmed Central PMCID: 7098786.

30.   Naguib YM. Antioxidant activities of astaxanthin and related carotenoids. J Agric Food Chem. 2000 Apr;48(4):1150-4. PubMed PMID: 10775364. Epub 2000/04/25.

31.   Ambati RR, Phang SM, Ravi S, Aswathanarayana RG. Astaxanthin: sources, extraction, stability, biological activities and its commercial applications--a review. Mar Drugs. 2014 Jan 7;12(1):128-52. PubMed PMID: 24402174. Pubmed Central PMCID: PMC3917265. Epub 2014/01/10.

32.   Petri D, Lundebye AK. Tissue distribution of astaxanthin in rats following exposure to graded levels in the feed. Comp Biochem Physiol C Toxicol Pharmacol. 2007 Mar;145(2):202-9. PubMed PMID: 17257901. Epub 2007/01/30.

33.   Guo SX, Zhou HL, Huang CL, You CG, Fang Q, Wu P, et al. Astaxanthin attenuates early acute kidney injury following severe burns in rats by ameliorating oxidative stress and mitochondrial-related apoptosis. Mar Drugs. 2015 Apr 13;13(4):2105-23. PubMed PMID: 25871290. Pubmed Central PMCID: PMC4413202. Epub 2015/04/15.

34.   Guo S, Guo L, Fang Q, Yu M, Zhang L, You C, et al. Astaxanthin protects against early acute kidney injury in severely burned rats by inactivating the TLR4/MyD88/NF-kappaB axis and upregulating heme oxygenase-1. Sci Rep. 2021 Mar 23;11(1):6679. PubMed PMID: 33758309. Pubmed Central PMCID: PMC7988001. Epub 2021/03/25.

35.   Qiu X, Fu K, Zhao X, Zhang Y, Yuan Y, Zhang S, et al. Protective effects of astaxanthin against ischemia/reperfusion induced renal injury in mice. J Transl Med. 2015 Jan 27;13:28. PubMed PMID: 25623758. Pubmed Central PMCID: PMC4323259. Epub 2015/01/28.

36.   Ben-Nun A, Bashan N, Potashnik R, Cohen-Luria R, Moran A. Cystine loading induces Fanconi's syndrome in rats: in vivo and vesicle studies. Am J Physiol. 1993 Dec;265(6 Pt 2):F839-44. PubMed PMID: 8285217. Epub 1993/12/01.

37.   Foreman JW, Benson L. Effect of cystine loading and cystine dimethylester on renal brushborder membrane transport. Biosci Rep. 1990 Oct;10(5):455-9. PubMed PMID: 2282372. Epub 1990/10/01.

38.   Gahl WA, Balog JZ, Kleta R. Nephropathic cystinosis in adults: natural history and effects of oral cysteamine therapy. Ann Intern Med. 2007 Aug 21;147(4):242-50. PubMed PMID: 17709758. Epub 2007/08/22.

39.   Reimand J, Isserlin R, Voisin V, Kucera M, Tannus-Lopes C, Rostamianfar A, et al. Pathway enrichment analysis and visualization of omics data using g:Profiler, GSEA, Cytoscape and EnrichmentMap. Nature protocols. 2019 Feb;14(2):482-517. PubMed PMID: 30664679. Pubmed Central PMCID: 6607905.

40.   Wagner CA, Finberg KE, Breton S, Marshansky V, Brown D, Geibel JP. Renal vacuolar H+-ATPase. Physiological reviews. 2004 Oct;84(4):1263-314. PubMed PMID: 15383652.

41.   Sumayao R, McEvoy B, Martin-Martin N, McMorrow T, Newsholme P. Cystine dimethylester loading promotes oxidative stress and a reduction in ATP independent of lysosomal cystine accumulation in a human proximal tubular epithelial cell line. Experimental physiology. 2013 Oct;98(10):1505-17. PubMed PMID: 23813804.

42.   Hennings JC, Picard N, Huebner AK, Stauber T, Maier H, Brown D, et al. A mouse model for distal renal tubular acidosis reveals a previously unrecognized role of the V-ATPase a4 subunit in the proximal tubule. EMBO molecular medicine. 2012 Oct;4(10):1057-71. PubMed PMID: 22933323. Pubmed Central PMCID: 3491836.

43.   Zhang J, Johnson JL, He J, Napolitano G, Ramadass M, Rocca C, et al. Cystinosin, the small GTPase Rab11, and the Rab7 effector RILP regulate intracellular trafficking of the chaperone-mediated autophagy receptor LAMP2A. J Biol Chem. 2017 Jun 23;292(25):10328-46. PubMed PMID: 28465352. Pubmed Central PMCID: PMC5481548. Epub 2017/05/04.

44.   Zhang J, He J, Johnson JL, Rahman F, Gavathiotis E, Cuervo AM, et al. Chaperone-Mediated Autophagy Upregulation Rescues Megalin Expression and Localization in Cystinotic Proximal Tubule Cells. Front Endocrinol (Lausanne). 2019;10:21. PubMed PMID: 30774622. Pubmed Central PMCID: PMC6367655. Epub 2019/02/19.

45.   Vaisbich MH, Pache de Faria Guimaraes L, Shimizu MH, Seguro AC. Oxidative stress in cystinosis patients. Nephron Extra. 2011 Jan;1(1):73-7. PubMed PMID: 22470381. Pubmed Central PMCID: PMC3290844. Epub 2011/01/01.

46.   Forestier L, Jean G, Attard M, Cherqui S, Lewis C, van't Hoff W, et al. Molecular characterization of CTNS deletions in nephropathic cystinosis: development of a PCR-based detection assay. American journal of human genetics. 1999 Aug;65(2):353-9. PubMed PMID: 10417278. Pubmed Central PMCID: 1377934.

47.   Hollywood JA, Kallingappa PK, Cheung PY, Martis RM, Sreebhavan S, Atiola RD, et al. Cystinosin-deficient rats recapitulate the phenotype of nephropathic cystinosis. American journal of physiology Renal physiology. 2022 Aug 1;323(2):F156-F70. PubMed PMID: 35695380.

48.   Andrzejewska Z, Nevo N, Thomas L, Chhuon C, Bailleux A, Chauvet V, et al. Cystinosin is a Component of the Vacuolar H+-ATPase-Ragulator-Rag Complex Controlling Mammalian Target of Rapamycin Complex 1 Signaling. Journal of the American Society of Nephrology : JASN. 2016 Jun;27(6):1678-88. PubMed PMID: 26449607. Pubmed Central PMCID: PMC4884097. Epub 2015/10/10.

49.   Gahl WA, Thoene JG, Schneider JA. Cystinosis. N Engl J Med. 2002 Jul 11;347(2):111-21. PubMed PMID: 12110740. Epub 2002/07/12.

50.   Brodin-Sartorius A, Tete MJ, Niaudet P, Antignac C, Guest G, Ottolenghi C, et al. Cysteamine therapy delays the progression of nephropathic cystinosis in late adolescents and adults. Kidney Int. 2012 Jan;81(2):179-89. PubMed PMID: 21900880. Epub 2011/09/09.

51.   Coffey EE, Beckel JM, Laties AM, Mitchell CH. Lysosomal alkalization and dysfunction in human fibroblasts with the Alzheimer's disease-linked presenilin 1 A246E mutation can be reversed with cAMP. Neuroscience. 2014 Mar 28;263:111-24. PubMed PMID: 24418614. Pubmed Central PMCID: PMC4028113. Epub 2014/01/15.

52.   Jefferies KC, Cipriano DJ, Forgac M. Function, structure and regulation of the vacuolar (H+)-ATPases. Arch Biochem Biophys. 2008 Aug 1;476(1):33-42. PubMed PMID: 18406336. Pubmed Central PMCID: PMC2543942. Epub 2008/04/15.

53.   Zoncu R, Bar-Peled L, Efeyan A, Wang S, Sancak Y, Sabatini DM. mTORC1 senses lysosomal amino acids through an inside-out mechanism that requires the vacuolar H(+)-ATPase. Science. 2011 Nov 4;334(6056):678-83. PubMed PMID: 22053050. Pubmed Central PMCID: PMC3211112. Epub 2011/11/05.

54.   Hughes AL, Gottschling DE. An early age increase in vacuolar pH limits mitochondrial function and lifespan in yeast. Nature. 2012 Dec 13;492(7428):261-5. PubMed PMID: 23172144. Pubmed Central PMCID: PMC3521838. Epub 2012/11/23.

55.   Webb BA, Aloisio FM, Charafeddine RA, Cook J, Wittmann T, Barber DL. pHLARE: a new biosensor reveals decreased lysosome pH in cancer cells. Molecular biology of the cell. 2021 Jan 15;32(2):131-42. PubMed PMID: 33237838. Pubmed Central PMCID: 8120692.

56.   DePedro HM, Urayama P. Using LysoSensor Yellow/Blue DND-160 to sense acidic pH under high hydrostatic pressures. Analytical biochemistry. 2009 Jan 15;384(2):359-61. PubMed PMID: 18976626.

57.   Ma L, Ouyang Q, Werthmann GC, Thompson HM, Morrow EM. Live-cell Microscopy and Fluorescence-based Measurement of Luminal pH in Intracellular Organelles. Frontiers in cell and developmental biology. 2017;5:71. PubMed PMID: 28871281. Pubmed Central PMCID: 5566985.

58.   Marchi S, Patergnani S, Pinton P. The endoplasmic reticulum-mitochondria connection: one touch, multiple functions. Biochim Biophys Acta. 2014 Apr;1837(4):461-9. PubMed PMID: 24211533. Epub 2013/11/12.

59.   Farmer BC, Walsh AE, Kluemper JC, Johnson LA. Lipid Droplets in Neurodegenerative Disorders. Front Neurosci. 2020;14:742. PubMed PMID: 32848541. Pubmed Central PMCID: PMC7403481. Epub 2020/08/28.

60.   Gahl WA, Bernardini I, Finkelstein JD, Tangerman A, Martin JJ, Blom HJ, et al. Transsulfuration in an adult with hepatic methionine adenosyltransferase deficiency. J Clin Invest. 1988 Feb;81(2):390-7. PubMed PMID: 3339126. Pubmed Central PMCID: PMC329581. Epub 1988/02/01.

61.   Bernardini I, Rizzo WB, Dalakas M, Bernar J, Gahl WA. Plasma and muscle free carnitine deficiency due to renal Fanconi syndrome. J Clin Invest. 1985 Apr;75(4):1124-30. PubMed PMID: 3988933. Pubmed Central PMCID: PMC425435. Epub 1985/04/01.

62.   Sakai S, Nishida A, Ohno M, Inatomi O, Bamba S, Sugimoto M, et al. Astaxanthin, a xanthophyll carotenoid, prevents development of dextran sulphate sodium-induced murine colitis. J Clin Biochem Nutr. 2019 Jan;64(1):66-72. PubMed PMID: 30705514. Pubmed Central PMCID: PMC6348411. Epub 2019/02/02.

63.   Chitchumroonchokchai C, Bomser JA, Glamm JE, Failla ML. Xanthophylls and alpha-tocopherol decrease UVB-induced lipid peroxidation and stress signaling in human lens epithelial cells. J Nutr. 2004 Dec;134(12):3225-32. PubMed PMID: 15570017. Epub 2004/12/01.

64.   Yang JP, Shin JH, Seo SH, Kim SG, Lee SH, Shin EH. Effects of Antioxidants in Reducing Accumulation of Fat in Hepatocyte. International journal of molecular sciences. 2018 Aug 29;19(9). PubMed PMID: 30158441. Pubmed Central PMCID: PMC6164327. Epub 2018/08/31.

65.   Wang CM, Cai XL, Wen QP. Astaxanthin reduces isoflurane-induced neuroapoptosis via the PI3K/Akt pathway. Mol Med Rep. 2016 May;13(5):4073-8. PubMed PMID: 27035665. Epub 2016/04/02.

66.   Qiao J, Rong L, Wang Z, Zhang M. Involvement of Akt/GSK3beta/CREB signaling pathway on chronic omethoate induced depressive-like behavior and improvement effects of combined lithium chloride

and astaxanthin treatment. Neurosci Lett. 2017 May 10;649:55-61. PubMed PMID: 28366776. Epub 2017/04/04.

67.   Deng X, Wang M, Hu S, Feng Y, Shao Y, Xie Y, et al. The Neuroprotective Effect of Astaxanthin on Pilocarpine-Induced Status Epilepticus in Rats. Front Cell Neurosci. 2019;13:123. PubMed PMID: 30983975. Pubmed Central PMCID: PMC6449650. Epub 2019/04/16.

68.   Kim JH, Nam SW, Kim BW, Choi W, Lee JH, Kim WJ, et al. Astaxanthin improves stem cell potency via an increase in the proliferation of neural progenitor cells. International journal of molecular sciences. 2010;11(12):5109-19. PubMed PMID: 21614195. Pubmed Central PMCID: PMC3100832. Epub 2010/01/01.

69.   Kim JH, Nam SW, Kim BW, Kim WJ, Choi YH. Astaxanthin improves the proliferative capacity as well as the osteogenic and adipogenic differentiation potential in neural stem cells. Food Chem Toxicol. 2010 Jun;48(6):1741-5. PubMed PMID: 20385192. Epub 2010/04/14.

70.   Zhang H, Yang W, Li Y, Hu L, Dai Y, Chen J, et al. Astaxanthin ameliorates cerulein-induced acute pancreatitis in mice. Int Immunopharmacol. 2018 Mar;56:18-28. PubMed PMID: 29328945. Epub 2018/01/13.

71.   Yan T, Zhao Y, Zhang X, Lin X. Astaxanthin Inhibits Acetaldehyde-Induced Cytotoxicity in SH-SY5Y Cells by Modulating Akt/CREB and p38MAPK/ERK Signaling Pathways. Mar Drugs. 2016 Mar 10;14(3). PubMed PMID: 26978376. Pubmed Central PMCID: PMC4820310. Epub 2016/03/16.

72.   Ikeda Y, Tsuji S, Satoh A, Ishikura M, Shirasawa T, Shimizu T. Protective effects of astaxanthin on 6-hydroxydopamine-induced apoptosis in human neuroblastoma SH-SY5Y cells. J Neurochem. 2008 Dec;107(6):1730-40. PubMed PMID: 19014378. Epub 2008/11/19.

73.   Wang HQ, Sun XB, Xu YX, Zhao H, Zhu QY, Zhu CQ. Astaxanthin upregulates heme oxygenase-1 expression through ERK1/2 pathway and its protective effect against beta-amyloid-induced cytotoxicity in SH-SY5Y cells. Brain Res. 2010 Nov 11;1360:159-67. PubMed PMID: 20828541. Epub 2010/09/11.

74.   Lin X, Zhao Y, Li S. Astaxanthin attenuates glutamate-induced apoptosis via inhibition of calcium influx and endoplasmic reticulum stress. Eur J Pharmacol. 2017 Jul 5;806:43-51. PubMed PMID: 28400209. Epub 2017/04/13.

75.   Bhuvaneswari S, Yogalakshmi B, Sreeja S, Anuradha CV. Astaxanthin reduces hepatic endoplasmic reticulum stress and nuclear factor-kappaB-mediated inflammation in high fructose and high fat diet-fed mice. Cell Stress Chaperones. 2014 Mar;19(2):183-91. PubMed PMID: 23852435. Pubmed Central PMCID: PMC3933623. Epub 2013/07/16.

76.   Kitahara A, Takahashi K, Morita N, Murashima T, Onuma H, Sumitani Y, et al. The Novel Mechanisms Concerning the Inhibitions of Palmitate-Induced Proinflammatory Factor Releases and Endogenous Cellular Stress with Astaxanthin on MIN6 beta-Cells. Mar Drugs. 2017 Jun 20;15(6). PubMed PMID: 28632169. Pubmed Central PMCID: PMC5484135. Epub 2017/06/21.

77.   Kleta R, Bernardini I, Ueda M, Varade WS, Phornphutkul C, Krasnewich D, et al. Long-term follow-up of well-treated nephropathic cystinosis patients. The Journal of pediatrics. 2004 Oct;145(4):555-60. PubMed PMID: 15480385.

78.   Nesterova G, Williams C, Bernardini I, Gahl WA. Cystinosis: renal glomerular and renal tubular function in relation to compliance with cystine-depleting therapy. Pediatric nephrology. 2015 Jun;30(6):945-51. PubMed PMID: 25526929.

79.   Wilmer MJ, Willems PH, Verkaart S, Visch HJ, de Graaf-Hess A, Blom HJ, et al. Cystine dimethylester model of cystinosis: still reliable? Pediatric research. 2007 Aug;62(2):151-5. PubMed PMID: 17597653.

80.   Bjornsson TD. Use of serum creatinine concentrations to determine renal function. Clinical pharmacokinetics. 1979 May-Jun;4(3):200-22. PubMed PMID: 383355.

81.   Moran A, Ben-Nun A, Potashnik R, Bashan N. Renal cells in culture as a model for cystinosis. Journal of basic and clinical physiology and pharmacology. 1990 Jan-Dec;1(1-4):357-72. PubMed PMID: 2085526.

82.	Uniyal AP, Mansotra K, Yadav SK, Kumar V. An overview of designing and selection of sgRNAs for precise genome editing by the CRISPR-Cas9 system in plants. 3 Biotech. 2019 Jun;9(6):223. PubMed PMID: 31139538. Pubmed Central PMCID: 6529479.

83.	Schneider JA, Bradley K, Seegmiller JE. Increased cystine in leukocytes from individuals homozygous and heterozygous for cystinosis. Science. 1967 Sep 15;157(3794):1321-2. PubMed PMID: 6038997. Epub 1967/09/15.

84.	Tusher VG, Tibshirani R, Chu G. Significance analysis of microarrays applied to the ionizing radiation response. Proceedings of the National Academy of Sciences of the United States of America. 2001 Apr 24;98(9):5116-21. PubMed PMID: 11309499. Pubmed Central PMCID: 33173.