1 **Near-chromosomal *de novo* assembly of Bengal tiger genome reveals genetic hallmarks**
2 **of apex-predation**
3

4 Harsh Shukla[1,*], Kushal Suryamohan[2,3,*], Anubhab Khan[1*], Krishna Mohan[4], Rajadurai C. Perumal[4],
5 Oommen K. Mathew[4], Ramesh Menon[5], Mandumpala Davis Dixon[4], Megha Muraleedharan[4], Boney
6 Kuriakose[4], Saju Michael[4], Sajesh P. Krishnankutty[4], Arun Zachariah[3,6], Somasekar Seshagiri[3,@],
7 and Uma Ramakrishnan[1,@]
8

[1]National Centre for Biological Sciences, Tata Institute of Fundamental Research, Bangalore
560065, India; [2]Research and Development Department, MedGenome Inc., Foster City, CA 94404,
USA; [3]SciGenom Research Foundation, 3rd Floor, Narayana Nethralaya Building, Narayana Health
City, #258/A, Bommasandra, Hosur Road, Bangalore, Karnataka, 560099, India; [4]AgriGenome Labs
Private Ltd, Kochi, Kerala 682030, India; [5]MedGenome Labs Ltd., 3rd Floor, Narayana Nethralaya
Building, Narayana Health City, #258/A, Bommasandra, Hosur Road, Bangalore, Karnataka 560099,
India; [6]Wayanad Wildlife Sanctuary, Sultan Bathery, Kerala 673592, India.
16

17 * equal contribution
18 [@] corresponding author (SS: sekar@sgrf.org; UR: uramakri@ncbs.res.in)
19

**Abstract**

The tiger, a poster child for conservation, remains an endangered apex predator. Continued

survival and recovery will require a comprehensive understanding of their genetic diversity and

the use of such information for population management. A high-quality tiger genome assembly

will be an important tool for conservation genetics, especially for the Indian tiger, the most

abundant subspecies in the wild. Here, we present high-quality near-chromosomal genome

assemblies of a female and a male wild Indian tiger (*Panthera tigris tigris*). Our assemblies had

a scaffold N50 of >140 Mb, with 19 scaffolds, corresponding to the 19 numbered chromosomes,

containing 95% of the genome. Our assemblies also enabled detection of longer stretches of

runs of homozygosity compared to previous genome assemblies which will improve estimates of

genomic inbreeding. Comprehensive genome annotation identified 26,068 protein-coding

genes, including several gene families involved in key morphological features such as the teeth,

claws, vision, olfaction, taste and body stripes. We also identified 301 microRNAs, 365 small

nucleolar RNAs, 632 tRNAs and other noncoding RNA elements, several of which are predicted

to regulate key biological pathways that likely contribute to tiger's apex predatory traits. We

identify signatures of positive selection in the tiger genome that are consistent with the *Panthera*

lineage. Our high-quality genome will enable use of non-invasive samples for comprehensive

assessment of genetic diversity, thus supporting effective conservation and management of wild

tiger populations.

**Introduction**

Tigers are among the most iconic and recognizable species in the world.  Despite being apex predators, they are among the most endangered animals with an estimated global population of 3,900 tigers remaining in the wild compared to over 100,000 at the turn of the 20th century (https://www.worldwildlife.org/species/tiger).  Historically, wild tigers roamed large swaths of the planet that included a range spanning present-day Armenia in Eastern Asia to Indonesia in Southeast Asia and from the Russian Far East to the Southern tip of India (Luo et al., 2019).  Anthropogenic activities such as hunting, urbanization, expansion of agriculture and deforestation, leading to both loss of habitat and prey have resulted in a >95% decline in the wild tiger populations (Ripple et al., 2014).

The Bengal tiger or the Indian tiger, *Panthera tigris tigris*, is a subspecies native to the Indian subcontinent and is highly endangered.  It is estimated that over 50,000 tigers inhabited the unbroken forests of India (Mondol et al., 2009).  Much like global tiger population decline, Bengal tiger population in India also began to decline more than a century ago, leaving fewer than 2,000 tigers in the wild by 1970.  In 1973, India declared the tiger its national animal and set up 'Project Tiger' to conserve this majestic animal (Task Force, 1972).  With over 50 tiger reserves, the project has given the tiger a chance for continued survival.  Today there are ~2,900 wild tigers in these reserves in India, accounting for ~60% of the global wild tiger population (Jhala et al., 2020).  The Bengal tiger population is estimated to be genetically the most diverse and hence represents the best gene pool reservoir important for conservation (Armstrong et al., 2021; Luo et al., 2019).  While the genetic diversity for Bengal tigers continues to increase, small and fragmented reserves averaging <1,500 square kilometers in India will likely lead to isolated population groups susceptible to inbreeding depression and genetic bottlenecks, limiting the success of conservation efforts (Armstrong et al., 2021; Khan et al., 2021; Luo et al., 2019).  In this context, continued monitoring of the genetic diversity of these populations will be important for conservation.  Obtaining high quality samples from the animals in the wild to conduct genetic studies is not practical.  A high-quality reference genome will allow the use of DNA from non-invasive samples scat samples to study, monitor and manage wild populations (Khan et al., 2020).

The evolution of genomics technologies over the last decade (Giani et al., 2020; Phillippy, 2017) enable high-quality *de novo* genome assemblies of endangered and threatened species (Fan et al., 2019; Ge et al., 2019; Murchison et al., 2012; Suryamohan et al., 2020; Sutton et al., 2018). These include short- and long-read sequencing, optical mapping and

chromosome conformation capture sequencing technologies that together allow for the

generation of near-chromosomal genome assembly.

The Bengal tiger is a diploid organism with a chromosome number 2n = 38, which

includes 16 pairs of metacentric/submetacentric autosomes, two pairs of acrocentric

autosomes, a meta-centric X, and Y sex chromosome.  To date, none of the published tiger

genomes are near-chromosomal assemblies.  The Amur tiger (*Panthera tigris altaica*) genome

assembly, generated using short-read data was highly fragmented (Cho et al., 2013).  Recently,

a genome assembly for a captive-wild-caught Malayan tiger (*Panthera tigris jacksoni*) using 10x

linked-reads has been reported (Armstrong et al., 2021).  Additionally, although several tiger

resequencing studies have produced additional sequence data  (Khan et al., 2020; Liu et al.,

2018), none have resulted in a contiguous high-quality reference genome.

Here, we report near-chromosomal *de novo* reference genome assemblies of a male

and a female Bengal tiger from the wild. The female animal is a famous Bengal tigress named

Machali from the Ranthambore National Park, India, crowned the Queen Mother of Tigers and is

the most widely photographed tiger in the world (https://en.wikipedia.org/wiki/Machali_(tigress)).

Machali, a reproductively successful founder, gave birth to seven females and four males

between 1999 to 2006.

We have performed a comprehensive annotation of our high-quality genomes and

identified several genes that contribute to key traits that make the tiger an apex predator.  This

included genes involved in skin patterning, tooth development, vision, endurance and olfaction.

We also show that the contiguous genomes generated here enable identification of longer

stretches of runs of homozygosity (ROH) than previously possible with fragmented genomes.

Further, we used the reference quality gene annotation models to analyze signals of positive

selection and evolution of big cat-specific traits.

**Results**

**Genome sequencing and chromosome-level assembly**

We generated high-quality genome assembly using long-read and short-read

sequencing data, chromatin conformation (Hi-C) sequence and/or optical mapping (Bionano)

data (**Supplementary Table 1a-b**, materials and methods) (Dudchenko et al., 2017; Lieberman-

Aiden et al., 2009).  The Machali (MC) draft assembly generated using long-read sequence data

contained 2,766 contigs, spanning 2.4 Gb and had a contig N50 of 3.26 Mb (**Table 1a;** see

materials and methods).  Following error correction using PacBio and Illumina data, we obtained

the PanTigT.MC.v1 assembly that served as input for Hi-C data scaffolding (Putnam et al.,

2016).  This resulted in a near-chromosomal assembly, PanTigT.MC.v2 containing 1,054

scaffolds and a scaffold N50 of 145.22 Mb.  This represented a ~50x improvement in genome

contiguity (see materials and methods).  The BUSCO completeness score for this assembly for

detection of conserved mammalian single-copy orthologs (n= 4,104 genes; **Table 2**) was 95.4%

(Simao et al., 2015; Waterhouse et al., 2018).

Similarly, the initial draft assembly of the Southern Indian (SI) animal obtained using

long-read data consisted of 845 contigs with a contig N50 of 29 Mb (**Table 1b**, **Supplementary**

**Table 1a-b;** see materials and methods).  After polishing with Illumina short-read sequencing,

we used the resulting assembly, PanTigT.SI.v1, as input for scaffolding with Chicago chromatin

interaction mapping data and ordered and oriented the contigs, corrected mis-joined and

merged overlaps.  Integration with Chicago data led to a resolved assembly N50 of 2.3 Mb

(PanTigT.SI.v2; **Table 1b**). This is consistent with the finding that Chicago data captures short

range chromatin interactions and corrects for misjoins in assemblies obtained using long-read

sequencing data alone (Putnam et al., 2016).  Integration of this assembly with *de novo*

assembled Bionano optical map data (PanTigT.SI_BNG) resulted in an assembly,

PanTigT.SI.v3, containing 3,811 scaffolds and a scaffold N50 of 141.0 Mb.  The BUSCO

completeness score for this assembly was 92.6% (**Table 1b and 2**).  Interestingly, further

incorporation of Hi-C data did not result in an improvement of the SI genome contiguity (see

materials and methods).

To assess for any ordering/orientation errors in the PanTigT.MC.v2 assembly after Hi-C

scaffolding, we used the chromosome-level reference genome of domestic cat (*Felis catus*;

Genbank: GCA_013340865.1, *Felis catus 9.0*) (Buckley et al., 2020) as well as the

PanTigT.SI.v3 assembly to delineate the errors from actual structural changes (see materials

and methods).  Given the highly conserved karyotype among Felidae species (Wurster-Hill and

Gray, 1973), we used the domestic cat reference assembly to merge scaffolds into

chromosomes and assign chromosomes based on synteny giving rise to the final near-

chromosomal assemblies, PanTigT.MC.v3 and PanTigT.SI.v3 (**Figure 1a-b**).  The largest

scaffold from PanTigT.MC.v3 and PanTigT.SI.v3 spanned over 230 Mb.  Further, using the

male PanTigT.SI.v3 assembly, we were also able to identify 3 Y-chromosome-linked scaffolds

that spanned ~3 Mb (**Figure 1c**; see materials and methods).  Overall, both genome assemblies

were highly contiguous, with >93% of the genome contained in the 19 near-chromosome-level

scaffolds and were syntenic with each other (**Supplementary Figure 1**).  Also, consistent with

the evolutionary history, both genomes were highly collinear with the domestic cat reference

genome (**Figure 1d, Table 2**).

Comparison of the MC and SI genome assemblies to other published Felidae genomes

showed that the overall contiguity was better than the domestic cat reference genome.  For

comparison, the scaffold N50 of the Bengal tiger genomes were ~1.7× (83.8 versus 145-147

Mb) longer than that of the domestic cat genome (Felis_catus_9.0, Genbank:

GCA_013340865.1), a gold standard feline genome.  Unlike the tiger genome assembled *de*

*novo* in this study, the domestic cat genome was assembled using physical mapping and

sequencing.  Also, it was improved iteratively since its publication in 2007 (Pontius et al., 2007).

A comparison of our genome assemblies to that of the recently published Malayan tiger genome

(*Panthera tigris jacksoni*) showed that they were ~7x (21.2 versus 145-147 Mb) better as

assessed by scaffold N50 (Armstrong et al., 2021; Cho et al., 2013)(MalTig1.0, Genbank:

GCA_019348595.1) (**Table 3**). Also, a comparison of our genome assemblies to that of the

Amur tiger genome showed that they were ~16× more contiguous (8.8 versus 145-147 Mb

scaffold N50, respectively) (Armstrong et al., 2021; Cho et al., 2013) (PanTig1.0, Genbank:

GCA_000464555.1) (**Table 3**).  Although, genomes of other big cats such as jaguar and leopard

from the *Panthera* genus have been published (Figueiro et al., 2017; Kim et al., 2016), none are

near chromosomal reference assemblies (**Table 3**). Given our assembly included a male Bengal

tiger, we were able to identify Y chromosome-associated scaffolds that provides an important a

reference for Y-haplotyping of tiger populations.

**Runs of homozygosity analysis**

High-quality near-chromosomal genome assemblies can greatly help in several types of

population genetic analysis, especially of threatened and endangered species. Measuring

inbreeding due to shared ancestry of parental gametes is one of them. Typically, this is

measured by identifying runs of homozygosity (ROH) (Ceballos et al., 2018; Kardos et al., 2016;

Pemberton et al., 2012).  We aligned short-read Illumina data from four zoo-bred Individuals

with known pedigree inbreeding coefficients ranging from 0.21 to 0.28 (Sagar et al., 2021) to

MC, SI and the published Malayan tiger genomes (Armstrong et al. 2021) to assess the

effectiveness of these assemblies in detecting ROH (**Materials and methods**). We observed on

an average 2,895, 2,583 and 2,176 ROH regions that were >100 kb in length in the zoo-bred

individuals when using the MC, SI and Malayan tiger genome assemblies, respectively

(**Supplementary Tables 1c-e**).  Further, we estimated cumulative inbreeding in individuals due

to common ancestry dating back to 455, 45 and 4 generations ago (**Materials and methods**).

We also observed that all three genomes performed comparably when estimating shorter ROH

that represent older ancestry (Khan et al. 2021) (**Figure 1e**) while the MC and SI assemblies

performed better at detection of ROH longer than 1 Mb that represents recent inbreeding,

compared to the Malayan tiger genome (n= 177 for MC, n= 106 for SI vs n= 94 for Malayan tiger

on average) (**Figure 1f**). All estimates of recent inbreeding from ROH longer than 1Mb

correlated with the pedigree inbreeding coefficients of the individuals, thus demonstrating the

need for near-chromosomal genome assemblies in conservation genomics.

**Genome Annotation**

Our genome assemblies, PanTigT.MC.v3 and PanTigT.SI.v3, both contained 19

scaffolds >10 Mb corresponding to the 19 chromosomes (**Figure. 1a-b**). These scaffolds

accounted for ~93% of the genome.  The average DNA base (GC) content of the assemblies

was about 41%.  Analysis of the repeat content revealed that ~36% of the genome was

repetitive (~870 Mb; **Figure 1a-b** and **Supplementary Table 1f** and **Supplementary Figure 2**)

with long interspersed nuclear elements (LINEs) being the dominant family of repeats

(**Supplementary Table 1f**).

We also annotated the genomes for noncoding RNA elements and created a database

of micro RNAs (miRNAs; n=299 in MC and 301 in SI), small nucleolar RNAs (SnoRNA; n=365 in

MC and 371 in SI), tRNAs (n=548 and 632 in MC and SI) and other noncoding RNA elements

(**Supplementary Table 1g**).  We next searched for predicted miRNAs targets using miRanda

(Enright et al., 2003).  Several miRNAs were predicted to target genes involved in critical

biological processes including angiogenesis (n=132 target genes; GO:0001525 and

GO:0045765-0045766), brain (n=78; GO:0007420, 1990403), bone (n=73; GO:0060348,

0048539, 0030282, 0060349, 0030500) and eye (n=63; GO:0002088, 0043010, 0001654,

0048593) and heart development (n = 43; GO:0035904, 0060347, 0007507, 0007512, 0060914,

0060973) (**Supplementary Tables 1h-i**).  Functional validation studies involving the predicted

miRNAs (Ambros, 2004; Lee et al., 1993) shall lead to a better understanding of the post-

transcriptional gene expression regulation in tigers and big cats during pre- and post-

development.

Next, we used the MAKER pipeline (Campbell et al., 2014; Cantarel et al., 2008) to

annotate the genomes using protein homology and RNA-seq expression data.  We predicted

19,931 genes that correspond to 22,718 transcripts in the PanTigT.MC.v3 genome assembly.

In the male SI individual PanTigT.SI.v3, we detected 21,126 protein-coding genes that mapped

to 24,074 transcripts (**Figure 2a and Supplementary Table 1j**).  Overall, the annotation yielded

26,068 unique protein-coding between the MC and SI genomes.  About 95% of all annotated

genes were located on the 19 largest scaffolds corresponding to the numbered chromosomes in

the genomes. Using a previously developed annotation pipeline (Suryamohan et al., 2020), we

functionally classified the protein coding genes (**Supplementary Table 1j**). A total of 19,640

(86%) and 20,412 (85%) predicted proteins contained a canonical start and stop codon in the

PanTigT.MC.v3 and PanTigT.SI.v3 assemblies.  We found that ~99% of these genes had a

corresponding ortholog in either the Human Gene Nomenclature Committee database, NCBI's

non-redundant database or the TrEMBL (https://www.ebi.ac.uk/uniprot) database

(**Supplementary Tables 1k-n**).  High-level gene function classification of the identified proteins

revealed 179 solute carriers as the most abundant type of proteins followed by 78 zinc finger

proteins (**Supplementary Figure 3** and **Supplementary Tables 1l and 1n**).

A tiger typically consumes at least one deer-sized animal each week for survival.  Its

metabolism, sensory functions and other adaptations are key to it evolutionary success as an

apex predator.  Using our high-contiguity genomes, we annotated several full-length amino-acid

transport (n=7; GO: 0006865) and protein, cholesterol and fatty acid metabolism-related genes

(n=21; GO: 0008203, 0019538, 0006631) that likely are crucial for the successful adaptation to

its carnivorous diet (**Supplementary Tables 1kand 1m**) (Kim et al., 2016).  Further, we

identified multiple genes involved in G-protein coupled signaling (n=270) and olfactory receptor

activity (n=107) that likely have an important role in sensory functions such as smell, vision and

hearing, nervous system development, mate selection and hunting (Barnett et al., 2020).

The evolutionary versatility of teeth is an important factor contributing to the success of

carnivores.  Diet-dependent changes in complexity of dental patterns, tooth morphology,

number, function and diet are interlinked (Aristotle et al., 1937).  One unique morphological

feature that has evolved independently numerous times within the theriodont lineage that

includes mammalia, is the saber-tooth morphology of permanent upper canines (Simpson,

1941).  Tigers have the largest upper canines of all the big cats and is key for its apex predatory

status. We identified homologs of gene families known to be involved in tooth development

including the *Wnt* family members and homeobox genes (*Hox*) (Sharpe, 1995) *Msx-1*, *Msx-*

*2*,(Vastardis et al., 1996), *Dlx-1*, *Dlx-2* and *Barx-1.* We also identified *Pax9* known to be

important for molar development. Additionally, our analysis identified several *Fibroblast growth*

*factor* (*Fgf*)*,* and *LIM homeodomain transcription factor* (*Lhx*) family members (Zhang et al.,

2005) involved in regulating enamel and dentin levels (**Figure 2b**; **Supplementary Tables 1k**

**and 1m**). Mouse genetic analysis showed that the *fibroblast growth factor 10* (*Fgf10*) to be a

critical factor in determining the length of the teeth (Yokohama-Tamaki et al., 2006). In addition

to *Fgf genes*, we also identified *Matrix metalloproteinase-20* (*Mmp-20*) that has been shown to

be involved in tooth development (**Figure 2b**; **Supplementary Tables 1k and 1m**). (Thomas

and Sharpe, 1998; Thomas et al., 1997).

The retractable sharp tiger claws are an important tiger adaptation critical for predation.

Claws are a variation in the distal limb integumentary appendages of certain mammals,

including most carnivorous animals. Several epithelial-mesenchymal signaling molecules

involved in patterning ectodermal derivatives such as teeth, hair, and feathers are also involved

in patterning distal epidermal appendages such as claws and nails (Vickaryous and Sire, 2009).

Mutations in several *Hox family member* genes have previously been shown to affect claw/nail

and limb development in mice and lead to disorders such as brachydactyly in humans (Davis

and Capecchi, 1996; Hamrick, 2001; Johnson et al., 2003; Luo et al., 2019; Montague et al.,

2014). In the tiger genome besides the several homeobox genes including *Homeobox* family

genes (*Hox*), *Msx-1* and *Msx-2,* we identified bone morphogenic protein (*Bmp*) family members

known to be involved in early distal limb development (e.g., *Bmp4*, posterior *Hox* genes) and

induction of the claw growth, and/or proliferation (**Figure 2b**; **Supplementary Tables 1k and**

**1m**).

It is estimated that about two-thirds of all extant carnivorous mammals are mainly

nocturnal (Roots, 2006). Nocturnality is characterized by an expansion in the number of genes

responsible for dim/low-light vision. The retina is a light-sensitive layer of eye tissue and

consists of six types of neurons which include light-sensitive photoreceptor cells (rods and

cones). Rod cells are responsible for discerning shapes and are highly sensitive to low light

while cone cells confer bright light sensitivity as well as color vision. Tigers have more rods in

their eyes than cones (Ahnelt et al., 2006). We identified several genes critical for low-light

vision, including Neural retina leucine zipper (*Nrl*), S-antigen visual arrestin (*Sag*), G Protein

Subunit Alpha Transducin 1 (*Gnat1*), G Protein Subunit Beta 3 (*Gnb3*), Phosphodiesterase

genes *Pde6a, Pde6b and Pd36g,* and Cyclic Nucleotide Gated Channel Subunit Alpha 1

(*Cnga1*) and Beta 1 (*Cngb1*) (**Figure 2b**; **Supplementary Tables 1k and 1m**).  Knockout

mutation of *Nrl* leads to the loss of rod cells (Ohsawa and Kageyama, 2008) while mutations in

*Gnb3*, *Cnga1* and *Cngb1* have been shown to lead to night blindness in humans and retinal

degeneration in chickens (Bareil et al., 2001; Katagiri et al., 2014; Tummala et al., 2006; Vincent

et al., 2016).

273            While the tiger's sense of smell, though not critical for hunting, is used for

communication between animals particularly, for marking territory, courtship and reproduction.

The vomeronasal organ, also called the Jacobson's organ, is primarily responsible for

pheromone detection in tigers (Poddar-Sarkar and Brahmachary, 2014). We identified twenty

one vomeronasal type-1 receptor genes (V1R), of which twelve were incomplete with truncated

sequences, indicating potential pseudogenization, consistent with findings in other carnivorous

species (Young et al., 2010).  In contrast to the ~400 olfactory receptor genes found in humans

(Hasin-Brumshtein et al., 2009), we identified 69 olfactory receptor (OR) genes based on

homology to other OR genes (**Figure 2b**; **Supplementary Tables 1k and 1m**) (Montague et al.,

2014).

283            Mammalian taste is mainly mediated by receptor cells organized in taste buds on the

tongue (Bachmanov et al., 2014).  It has been established that the tiger can taste salt, bitter and

acidic flavors and to a lesser degree sweetness (Jiang et al., 2012; Li et al., 2005).  We

identified a fully intact copy of *Tas1r1* (type 1 taste receptor), responsible for sweet taste

perception, and two Tas2r (type 2 taste receptor) genes involved in bitter taste perception

(Chandrashekar et al., 2000) (**Figure 2b**; **Supplementary Tables 1k and 1m**).  Notably, we did

not identify a full-length homolog of *Tas1r2*, a sweet taste receptor gene, consistent with

observations in other carnivorous mammals (Jiang et al., 2012; Li et al., 2009b).

291            Tigers are the only striped cat of the genus *Panthera* and are most recognizable for their

pattern of black vertical stripes on reddish-orange fur.  Our high-quality assembly and

annotation identified several genes associated with morphometric variation in domestic cats

such as pigmentation, coat patterns and other phenotypes. Included among these were

homologs of the atrial natriuretic peptide-converting enzyme (*Corin*), membrane-associated

transporter protein (*Slc45a2)*, *Agouti-signaling protein* (*Asip*), *Tyrosinase-related protein 1*

(*Tyrp1*), *Tyrosinase-related protein 2* (*Tyrp2*), *Tyrosinase* (*Tyr*), *Melanophilin* (*Mlph*), *Tyrosine-*

*Protein Kinase Kit* (*Kit*), *Keratin 71* (*Krt71*) and *Aminopeptidase N* (*Laeverin*; *Anpep*).  Mutations

in these genes can lead to a variety of phenotypes including variation in coat and feet color,

stripe patterns and the color of stripes, tail shapes (**Figure 2c-d**) (Gandolfi et al., 2013; Ishida et

al., 2006; Kaelin et al., 2012; Luo et al., 2019; Lyons et al., 2005; Xu et al., 2017).  Of note, our

high-quality genome enabled accurate annotation of long genes with their complete intron-exon

structure including that of *Slc45a2,* a ~33kb gene consisting of 7 exons.  Using a combination of

transcriptome (RNA-seq) and genome data, we were also able to annotate a ~249 kb long full-

length copy of *Corin* made of 22 exons.  These fully annotated genes will serve as a high-quality

resource for understanding coat patterning genes in tigers and other animals.

**Phylogeny and positive selection analysis**

To compare the annotations of the genomes from this study to other big cat species

including the Amur tiger, we constructed a phylogeny using annotated protein sequences from

13 selected species, namely, dog, Amur tiger, Bengal tigers (this study), lion, leopard, jaguar,

clouded leopard, domestic cat, Canada lynx, cougar, horse, and rabbit.  The proteins from these

species clustered into 26,169 orthogroups (representing 97.2% of the input protein sequences)

of which 4,576 were single copy orthologs.  The remaining protein sequences did not cluster

(854 proteins (3.5%) in SI and 489 genes (2.2%) in MC, respectively).  Next, we used the 4,576

single-copy orthologs to test for signals of positive selection in big cats using site models in

PAML (Yang, 2007).  Using Likelihood Ratio Tests (LRT) for models M1-M2 and M7-M8, we

identified 1,484 single copy orthologs to be positively selected (p-value <0.05) (**Supplementary**

**Tables 1o-q**).  Included in this list of positively selected genes, were genes involved in muscle

development *Atp1b4*, angiogenesis (*Srpx2*), fatty acid and cholesterol metabolism (*Slc4a9,*

*Apod*), mitochondrial respiration (*Cq10a*) and heart development/endurance-related genes (e.g.,

*Nkx25*), indicating that these genes are evolving at a faster rate, consistent with the

physiological needs of big cats (**Figure 3b** and **Supplementary Tables 1o-q**) (Barnett et al.,

2020; Cho et al., 2013; Figueiro et al., 2017; Kim et al., 2016).  Further, pathway enrichment

analysis of the positively selected genes confirmed enrichment of several metabolism-related

pathways including cholesterol metabolism (GO:0008203) and fatty acid metabolism

(GO:0006631) that are critical obligate carnivores (**Figure 3c**).

**Discussion**

We have de novo assembled and annotated genomes of a female and a male Bengal

tiger.  The resulting assemblies were 17x more contiguous than the published Amur tiger

genome and 1.7x more contiguous than the domestic cat genome with a scaffold N50 of over

140 Mb, making it the most contiguous near-chromosomal wild felid genomes assembled to

date (**Table 3**).

We provide here a comprehensive annotation of 26,068 protein-coding genes from the

tiger genomes.  Additionally, we identified over 3,000 non-coding genes including for the first

time a genome-wide analysis of micro-RNAs and their putative target genes in a tiger genome.

Functional assignment identified genes and signaling pathways involved in endurance,

neurotransmission, teeth, vision, claw and skin patterning, all of which are important for the

tiger's adaption as an apex predator.  We have performed extensive annotation of genes

involved in variations in coat color, stripe patterns and other morphometric variations.

Our highly contiguous genome along with the annotations is an important resource for

studies on the genetics of Felidae and in general carnivoran development and evolution.  For

example, it is now feasible to establish homology between tooth components between different

big cat species by using the annotation generated in this study. Our annotated gene sets for the

Bengal tiger genomes provides a high-quality reference that can be used to support

resequencing efforts aimed at studying tiger populations.  The genome assemblies presented

here demonstrate their utility in population and conservation genetic studies of tigers. They

increase the accuracy of the estimates of important population genetic parameters like

inbreeding for assessing the threats to a population. Our reference genome along with

annotation-guided quantitative studies on wild and inbred, zoo-bred individuals will help in

understanding the possible phenotypic effects of inbreeding depression through assessment of

ROH.  The high-quality genomes present in this study will be important resources that will

enable analysis of sequencing data obtained from DNA present in non-invasive scat samples

obtained from the wild.  Such analysis will be important for understanding and monitoring tiger

population genetic diversity in the wild and support informed conservation efforts.

**356**    **Methods**

**357**    **Samples and nucleic acid preps**

**358**    Bengal tiger blood samples used in the study are reported in **Supplementary Table 1a**).

**359**    Genomic DNA from MC and SI was extracted from whole blood using the MagAttract HMW

**360**    DNA extraction kit (Qiagen). Sample quality was checked using Qubit 3.0, Nanodrop 8000,

**361**    Agilent TapeStation 4200 and pulse field gel electrophoresis (Sage Science).  Total RNA was

**362**    isolated from the blood using Trizol (ThermoFisher). A total of 500ng of RNA was used for

**363**    library preparation.

**364**

**365**    **Sequencing**

**366**    Sequencing libraries for PacBio, Oxford Nanopore (ONT) and Illumina sequencing were

**367**    constructed as per manufacturer's instructions.  Considering the genome size of Panthera Tigris

**368**    at ~2.4 Gb, we generated ~79.6 Gb of Pacbio long-read data (~33x; N50 = 10.3 Kb) and ~18.28

**369**    Gb of ONT long-read data (~7.5x, N50 = 8.14 kb) for the female tiger (MC). In addition, short-

**370**    read Illumina sequencing data 117 Gb, (50x; 250 bp library), 35 Gb (15x; 300 bp library 35 Gb

**371**    (15x; 500 bp library), and 35 Gb (15x; 800 bp library) was also generated.  For the male tiger

**372**    (SI), a total of 160.3 Gb (64x) ONT data, and ~60 Gb (25x) of Illumina short-read data was

**373**    generated (**Supplementary Table 1b**). For both animals, the long-read data was filtered using

**374**    BLAST against the 16s rRNA database (https://ncbiinsights.ncbi.nlm.nih.gov/2020/02/21/rrna-

**375**    databases/) to identify and remove any potential contaminant reads prior to genome assembly.

**376**    In addition, blood from an offspring of Machali, ST3 was also obtained.

**377**    RNA-seq short read (Illumina) data was obtained by sequencing PolyA RNA-sequencing

**378**    (RNA-seq) libraries prepared from ST3 blood using the Illumina TruSeq stranded messenger

**379**    RNA kit (**Supplementary Table 1b**).

**380**

**381**    **Chicago library preparation and sequencing**

**382**    Chicago library was prepared as described previously (Putnam et al., 2016). Briefly, for

**383**    each library, ~500ng of HMW gDNA (mean fragment length of 75 kb) was reconstituted into

**384**    chromatin *in vitro* and fixed with formaldehyde. Fixed chromatin was digested with DpnII, the 5'

**385**    overhangs filled in with biotinylated nucleotides, and then free blunt ends were ligated.  After

**386**    ligation, crosslinks were reversed and the DNA purified from protein.  Purified DNA was treated

**387**    to remove biotin that was not internal to ligated fragments.  The DNA was then sheared to ~350

**388**    bp mean fragment size and sequencing libraries were generated using NEBNext Ultra enzymes

**389**    and Illumina-compatible adapters.  Biotin-containing fragments were isolated using streptavidin

beads before PCR enrichment of each library.  The libraries were sequenced on an Illumina

platform.  A total of 155 million read pairs (2 x 150 bp) was generated for this library providing

~20x physical coverage of the genome (1-50 kb pairs) (**Supplementary Table 1b**).

**Bionano optical mapping data generation**

High-molecular weight genomic DNA extracted from whole blood from SI was embedded

in a thin agarose layer and was labeled and counterstained following the Direct Label and Stain

(DLS) Reagent Kit protocol (Bionano Genomics). The labeled sample was then loaded onto a

Saphyr chip and run on the Saphyr imaging instrument (BioNano Genomics).  A total of 473.7

Gb (~190x) optical map data was generated. *De novo* genome assembly using the Bionano

Access software for the DLS optical map data produced a 2.63 Gb assembly (PanTigT_BNG)

consisting of 182 scaffolds (N50 =145.93 Mb) (**Supplementary Table 1b**).

**Hi-C library preparation and sequencing**

Hi-C library (Dovetail Genomics), was prepared for the two samples (SI and ST3) as

described previously (Lieberman-Aiden et al., 2009).  Briefly, chromatin was fixed in place with

formaldehyde in the nucleus and then extracted.  Fixed chromatin was digested with DpnII, the

5' overhangs filled in with biotinylated nucleotides, and then free blunt ends were ligated.  After

ligation, crosslinks were reversed and the DNA purified from protein. Purified DNA was treated

to remove biotin that was not internal to ligated fragments. The DNA was then sheared to ~350

bp mean fragment size and sequencing libraries were generated using NEBNext Ultra enzymes

and Illumina-compatible adapters and sequenced (**Supplementary Table 1b**).

**Genome assembly**

We used Canu (v1.7.1; *corMinCoverage* = 0, *corMhapSensitivity* = *high and*

*correctedErrorRate*=0.12 (MC)/0.15 (SI)) to generate an initial draft assembly for both the study

animals.  The primary MC draft assembly (PanTigT.MC.v1) consisted of 2.4 Gb spanning 2,766

contigs (contig N50 = 3.26 Mb) (**Table 1**) and the SI draft assembly consisted of 845 contigs

totaling 2.38 Gb (contig N50 = 29.03 Mb). Next, both assemblies were polished to correct for

base call errors using raw PacBio data (MC only) and Illumina data (MC and SI).  Briefly,

PacBio reads were aligned to the PanTigT.MC.v1 using *pbalign* (pbalign version: 0.3.1)

resulting alignment bam file was used as input for polishing using *Arrow*

(https://github.com/pacificbiosciences/genomicconsensus/). A total of two rounds of polishing

were done using *Arrow.* Illumina short-read libraries generated for MC and SI (**Supplementary**

**Table 1b**) were used for error correction and conflict resolution using Pilon (v1.22) with the

following parameters (*--changes --diploid --fix all*) to produce polished assemblies

PanTig.T.MC.v1 and PanTigT.SI.v1 (**Table 1**).

Next, PanTigT.MC.v1 was used as the input for scaffolding with Hi-C sequence data

using HiRise resulting in an assembly PanTigT.MC.v2 containing 1,054 scaffolds (scaffold N50=

145.7 Mb). Similarly, PanTigT.SI.v1 was used as the input for scaffolding with Chicago

sequence data using HiRise.  This resulted in an assembly PanTigT.SI.v2 containing 6,889

scaffolds (scaffold N50 = 2.3 Mb).  PanTigT.SI.v2 was then scaffolded with Bionano optical map

data to yield PanTigT.SI.v3 assembly comprising of 3,886 scaffolds (scaffold N50 = 141.3 Mb).

Interestingly, incorporation of Hi-C data did not improve the contiguity of PanTigT.SI.v3 and did

not result in any breaks or joins to the assembly.  We also assessed if the order of integration of

the different sequencing data types would affect assembly quality. While scaffolding

PanTigT.SI.v1 with Hi-C data alone (no Bionano and Chicago) resulted in an assembly with

fewer scaffolds (n=692) but lower N50 (125.3 Mb), scaffolding of PanTigT.SI.v1 with only

Chicago and Hi-C data resulted in an assembly with 6,461 scaffolds and lower scaffold N50 of

48.2 Mb. Thus, we retained the Chicago-scaffolded assembly (PanTigT.SI.v2) for downstream

analysis including chromosome assignment.  Using the domestic cat reference assembly

(Felis_catus_9.0), we merged scaffolds into chromosomes, corrected orientation errors and

assigned chromosomes based on synteny using RaGOO (https://github.com/malonge/RaGOO)

resulting in the final SI assembly PanTigT.SI.v3 and PanTigT.MC.v3.

Next, error-corrected reads obtained from long-read data were used in conjunction with

RAILS (v1.4.1) (https://github.com/bcgsc/RAILS) to fill gaps in PanTigT.MC.v3 and

PanTigT.SI.v3 using Cobbler with the following parameters: -d 1000 -i 0.95. A total of 21 out of

1700 gaps (1.24%) were filled in PanTigT.MC.v3 with the largest gap closed being ~10,025 bp

while no significant improvement was observed for PanTigT.SI.v3. A final round of polishing with

Racon (v1.3.3) was done on PanTigT.MC.v3.

**Y-chromosome-linked scaffold identification**

To identify the male Y-linked scaffolds, we separately aligned Illumina reads obtained

from the female (MC) and the male animal (SI) to PanTigT.SI.v3 assembly using BWA (Li and

Durbin, 2009) allowing for two mismatches and one indel.  Scaffolds with less than 80%

alignment coverage were excluded from further analysis. Then, single-base depths were

calculated using SAMtools (Li et al., 2009a) following which coverage and mean depth for each

scaffold was calculated. Using the average coverage across a scaffold using either the male or

female reads, we identified 3 Y-linked scaffolds Sc4gRnz_35;HRSCAF=35_pilon,

Sc4gRnz_53;HRSCAF=53_pilon and Sc4gRnz_8;HRSCAF=8_pilon.

**Identification of mitochondrial contig**

In order to identify the mitochondrial contig, whole mitochondrial sequence

(NC_010642.1) for *Panthera tigris tigris* was downloaded from NCBI and all contigs from MC

and SI were mapped to it using minimap2. Two distinct mitochondrial contigs (MT) were

identified and retained in the primary assemblies.

**Chromosome painting**

Chromosome painting of SI or MC chromosomes with the domestic cat chromosomes

was performed using the SatsumaSynteny2 script with default parameters (Grabherr et al.,

2010).

**Runs of Homozygosity (ROH) analysis**

ROH were identified as described in Armstrong et al. (2021).  We aligned reads from short-

read sequencing (**Supplementary Table 1b**) from four zoo-bred tigers with known inbreeding

coefficients (Sagar et al., 2021) to MC, SI and Malayan tiger genome assembly (Armstrong et

al. 2021).  After sequence alignment, variant calling and filtering we used BCFtools/ROH at

default settings to determine the allele frequencies. We categorized ROH into three categories

of stretches longer than 100Kb, 1Mb and 10Mb.

**Repeat element identification**

We identified the repetitive elements in the genome by combining both homology-based

and de novo predictions. Briefly, we used RepeatModeler (v.1.0.11)

(http://www.repeatmasker.org/RepeatModeler.html) to construct the species-specific repeat

sequence libraries for both SI and MC genome assemblies, and then used these as a query to

identify repetitive elements using RepeatMasker.

**Noncoding RNA Annotation**

tRNA-encoding genes were predicted by tRNAscan-SE (version 1.3.1) (Lowe and Eddy,

1997) with default parameters. Ribosomal RNA (rRNA) fragments were predicted by aligning to

mammalian rRNA sequences by using BLASTN (version 2.2.26) with an E-value cutoff of 1e-10.

MicroRNA (miRNA) and small nuclear RNA (snRNA) genes were annotated using INFERNAL

(version 1.1.3) (Nawrocki and Eddy, 2013) by searching against the Rfam database (release

13.0) (Kalvari et al., 2018). Whole-genome target gene prediction for miRNAs was performed

using miRanda (version 3.3a) (Enright et al., 2003) using default parameters. Gene ontology

terms for the target genes were obtained from the functional annotation for the genomes.

**Genome Annotation**

Gene prediction was performed on the assemblies PanTigT.SI.v3 and PanTigT.MC.v3

using the genome annotation tool, MAKER2 (v2.31.10) (Cantarel et al., 2008) in an iterative

process.  First, *ab initio* gene prediction was performed by the programs SNAP (v2006- 07-28)

(Korf, 2004) and Augustus (v3.2.3) (Stanke and Morgenstern, 2005) using publicly available

transcriptome data from the Siberian tiger (SRA091968), a de novo assembled transcriptome

from ST3 blood RNA-seq data using Trinity (Grabherr et al., 2011) and a combined protein

database consisting of proteins from the UniProt/Swiss-Prot and NCBI non-redundant

databases of reviewed proteins. A total of three iterative runs of MAKER was used to refine the

gene models and produce the final gene set with an annotation edit distance (AED) cutoff of 0.6.

Genome annotation quality was assessed by BUSCO analysis using the conserved core set of

mammalian genes (79.3% and 76.9% BUSCO completeness score for MC and SI assemblies,

respectively).  Next, using the set of complete predicted protein sequences, we performed

functional annotation as described previously (Suryamohan et al., 2020) (**Supplementary Table**

**1g**).

To identify gene families specific to tooth development, claw, taste, neurotransmitter,

olfaction and coat color/pattern, we performed a literature-based search (Figueiro et al., 2017;

Hamrick, 2001; Luo et al., 2019; Montague et al., 2014) to identify genes potentially involved in

these developmental pathways. Completeness of gene models for each trait-specific gene

where present (predicted by MAKER) was confirmed by aligning each gene against its homolog

using NCBI BLAST.  Genes that were partial or incompletely annotated were manually curated.

Briefly, for each incomplete gene, the corresponding homolog obtained from UniProt was

aligned to the genome (SI/MC) using the 'est2genome' parameter in the Exonerate program

with default parameters.  The aligned genome coordinates of trait-specific genes were then

checked against the annotation file to validate the gene prediction.

**Phylogeny and positive selection analysis**

Proteomes from 12 selected species, namely, dog, Bengal tiger (this study), Amur tiger,

lion, leopard, jaguar, clouded leopard, domestic cat, Canada lynx, cougar, horse, and rabbit

were obtained from Ensembl Biomart (www.ensembl.org).  Next, Orthofinder (v2.5.2) was used

to cluster the proteins into orthogroups.  Of note, the snow leopard was not included in this

analysis due to limited annotation models and lack of a reference genome.  Next, single copy

orthologs were extracted for evolutionary divergence estimation and positive selection analyses.

The single copy orthologs were aligned with muscle (v3.8.31) and alignments trimmed and

converted to codon level alignments using trimal (v1.4rev15).  For each of these alignments

PAML/codeml (v.4.10) (Yang, 2007) was run pairwise to obtain four-fold degenerate bases

which are concatenated into a super-matrix based on gene partitions. The dN/dS ratios are also

obtained in this process.  The resulting supermatrix was used to generate a species tree using

iqtree2 (v2.0.3) with GTR+F+I+G4 models along with ultrabootstrap (1000 iterations).  Further,

PAML/mcmctree was used to estimate evolutionary divergence among the species considered

with 95% confidence intervals with a burn-in of 2,000 and sampling 20,000.  The evolutionary

divergence calibrations between the following species pairs were extracted from timetree.org:

*Felis catus* and *Lynx canadensis* (7.9 - 13.1 MYA), *Panthera concolor* and *Felis catus* (8.8 -

13.9 MYA), *Panthera leo* and *Panthera pardus* (2.6 - 4.74 MYA), *Neofelis nebulosa* and

*Panthera onca* (7.4 - 14 MYA).

**Site model tests**

The single copy orthologs were fit to the following codon substitution models (M0, M1, M2, M7,

M8) of PAML/codeml (Yang, 2007). Likelihood Ratio Test (LRT) between M1 vs M2 and M8 vs

M7 pairs were used to determine positive selection at Chi-square pvalue <= 0.05. Multiple

testing using Benjamini-Hodgberg (BH) method was then performed to determine FDR (False

Dicovery Rate) for the p-values obtained above in order to determine significantly positively

selected genes among the species. Bayesian Empirical Bayes (BEB) (Yang et al., 2005) results

were used to identify the sites of positive selection at >=95% and >=99% confidence intervals.

**Software URLs**

Canu - https://github.com/marbl/canu

RAILS - https://github.com/bcgsc/RAILS

RaGOO - https://github.com/malonge/RaGOO

PAML – http://abacus.gene.ucl.ac.uk/software/paml.html

Orthofinder - https://github.com/davidemms/OrthoFinder

Trimal – http://trimal.cgenomics.org/

Iqtree - http://www.iqtree.org/

Muscle - https://www.ebi.ac.uk/Tools/msa/muscle/

Vcftools - http://vcftools.sourceforge.net/

VEP – https://www.ensembl.org/vep

MAKER - http://www.yandell-lab.org/software/maker.html

BUSCO - https://busco.ezlab.org/

Exonerate - https://www.ebi.ac.uk/about/vertebrate-genomics/software/exonerate

Trinity - https://github.com/trinityrnaseq/trinityrnaseq

Augustus - http://bioinf.uni-greifswald.de/augustus/

SNAP - https://github.com/KorfLab/SNAP

Repeatmasker - https://www.repeatmasker.org/

RepeatModeler - https://www.repeatmasker.org/

D-Genies - http://dgenies.toulouse.inra.fr/

Satsuma - https://github.com/bioinfologics/satsuma2

Pilon - https://github.com/broadinstitute/pilon

miRanda - http://cbio.mskcc.org/miRNA2003/miranda.html

INFERNAL - http://eddylab.org/infernal

PLINK - http://zzz.bwh.harvard.edu/plink/

BWA - https://github.com/lh3/bwa

Samtools - http://www.htslib.org/

Sentieon - https://github.com/Sentieon

Racon - https://github.com/isovic/racon

**Acknowledgements**

This work was partially funded by Senior Fellowship DBT Wellcome Trust India Alliance grant to

UR (IA/S/16/2/502714); NCBS data cluster used is supported under project no. 12-R&D-TFR-

5.04-0900, Department of Atomic Energy, Government of India). All samples used here from

Rajasthan were collected for research presented in Khan et al., 2021, and permissions are

detailed there. We thank the Rajasthan Forest Department for their support.

**Data Accessibility**

Raw sequencing data (DNA and RNA) as well as both genome assemblies can be accessed at

NCBI under BioProject accession nos. PRJNA732096 and PRJNA796358.

**Table 1: Hierarchical genome assembly statistics for (a) MC and (b) SI genomes**

a)

| Assembly | Size (Gb) | # of contigs | # of scaffolds | Contig/scaffold N50 (Mb) | Longest scaffold (Mb) |
|---|---|---|---|---|---|
| **PanTigT.MC.v1 (LR + SR)** | 2.41 | 2,766 | - | 3.26 | 21.20 |
| **PanTigT.MC.v2 (v1 + Hi-C)** | 2.41 | - | 1,054 | 145.47 | 237.64 |

b)

| Assembly | Size (Gb) | # of contigs | # of scaffolds | Contig/scaffold N50 (Mb) | Longest scaffold (Mb) |
|---|---|---|---|---|---|
| **PanTigT.SI.v1 (LR + SR)** | 2.38 | 845 | - | 29 | 131.97 |
| **PanTigT.SI.v2 (v1 + Chicago)** | 2.53 | - | 6,889 | 2.10 | 29.57 |
| **PanTigT.SI_BNG (BNG)** | 2.63 | - | 182 | 145.93 | - |
| **PanTigT.SI.v3 (v2 + BNG)** | 2.53 | - | 3,811 | 141.30 | 238.76 |
| **ONT+Chicago+Bionano+Hi-C** | 2.53 | - | 3,811 | 141.30 | 238.76 |

**Table 2: Genome assembly statistics**

| Assembly | Machali | SI |
|---|---|---|
| Total sequence length (Gb) | 2.41 | 2.53 |
| Contig N50 (Mb) | 3.27 | 29.00 |
| Number of scaffolds | 1054 | 3,811 |
| Number of scaffolds > 10 Mb (% of assembly) | 19 (94.1) | 19 (93.3) |
| Largest scaffold (Mb) | 237.6 | 238.8 |
| Scaffold N50 (Mb) | 145.47 | 147.0 |
| Number of gaps | 1700 | 5708 |
| BUSCO (%) | 95.4 | 92.6 |

**Table 3: Comparison of MC and SI assemblies to other Felid genomes**

| Assembly | Domestic cat | Leopard | Jaguar | Lion | Amur Tiger | Malayan Tiger | Bengal Tiger ♀ (PanTigT.MC.v3) | Bengal Tiger ♂ (PanTigT.SI.v3) |
|---|---|---|---|---|---|---|---|---|
| Total sequence length (Gb) | 2.52 | 2.57 | 2.50 | 2.50 | 2.42 | 2.42 | 2.41 | 2.53 |
| Contig N50 (Mb) | 41.91 | 0.02 | 0.06 | 0.31 | 0.03 | 0.102 | 3.27 | 29.00 |
| #of scaffolds | 4,525 | 50,377 | 438,792 | 8,061 | 1,479 | 10,077 | 1054 | 3,811 |
| #of scaffolds > 10 Mb (% of assembly) | 19 (96.4) | 84 (75.8) | 0 | 18 (99.2) | 65 (40.9) | 77 (80.0) | 19 (94.1) | 19 (93.3) |
| Largest scaffold (Mb) | 242.1 | 84.1 | 1.4. | 238.9 | 41.6 | 79.7 | 237.6 | 238.8 |
| Scaffold N50 (Mb) | 83.8 | 21.7 | 116.0 | 136.0 | 8.8 | 21.3 | 145.5 | 147.0 |
| #of chromosomes (n) | 19 | 19 | 19 | 19 | 19 | 19 | 19 | 19 |
| Reference | (Buckley et al., 2020) | (Kim et al., 2016) | (Figueiro et al., 2017) | (Armstrong et al., 2020) | (Cho et al., 2013) | (Armstrong et al., 2021) | This study | This study |

**Figure legends**

**Figure 1: Bengal Tiger Genome assembly.** (**a-b**) Circos plot representing the near-
chromosome-level assembly of female (MC) **(a)** and male (SI) (**b**) Bengal tiger genomes.  The
two outermost tracks represent the chromosome length (Mb) and ID.     Repeat content per 100
kb window per chromosome (red line represents the mean genome-wide mean repeat content
~36 %) is shown in black.  Gene density (green bars indicate 1-5 genes per 100 kb window per
chromosome, orange 5-10, and red > 10 genes) is displayed next.  The innermost track
represents GC% per 100 kb window per chromosome (blue line represents mean GC% -
~41%). (**c**) Y-linked scaffolds from the male tiger genome assembly.  Circos plot track as in (a-
b). (**d**) Chromosome painting showing synteny between domestic cat genome (FelCat9.0;
Genbank: GCA_000181335.4) and the female Bengal tiger (MC) scaffolds. (**e**) Genomic
inbreeding coefficients ($F_{ROH}$) derived from runs of homozygosity (ROH) of different lengths
(>100 Kb, >1 Mb and >10 Mb) and, (**f**) box plot of binned distribution of ROH lengths in the four
zoo-bred tigers as derived when using the MC, SI and Malayan tiger genomes as the reference
genome.

**Figure 2: Genome Annotation and functional genes.** (**a**), Bar plot of the number of predicted
genes and corresponding transcripts observed in MC genome assembly and putative Y-linked
scaffolds in SI genome assembly. (**b**) Putative genes involved in various key biological
pathways important for apex predatory traints. (**c**) Complete *Slc45a* gene structure including the
predicted transmembrane (TM) domains.  Multiple sequence alignment of Scl45a c-terminal
region showing the A477>V mutation observed in white tiger (**d**) Schematic diagram of full-
length *Corin* gene.  Multi-species alignment of the Corin LDL receptor domain depicting the
known H>Y mutation associated with golden tabby pelage pattern (*Cell Res. 27(7):954–57*).

**Figure 3: Phylogeny and positive selection analysis.** (**a**) Phylogeny tree constructed using
single copy orthologs from selected Panthera and outgroup species. (**b**) Pathway analysis of
single copy orthologs within the Panthera lineage showing gene network interaction models. **(c)**
Scatter plot of evolutionary rates of single copy key pattern and pelage-related gene orthologs
identified in the Bengal tiger genome. cfa: *Canis familiaris* (dog)*,* pts: south Indian Bengal tiger
(male), ptm: Machali Bengal tigress (female), pta: *Panthera tigris altaica* (Amur tiger), ple:
*Panthera leo* (lion), ppa: *Panthera pardus* (leopard), pon: *Panthera onca* (jaguar), nne: *Neofelis*
*nebulosa* (clouded leopard), fca: *Felis catus* (domestic cat), lca: *Lynx canadensis* (Canada lynx),
pco: *Puma concolor* (cougar), eca: *Equus caballus* (domesticated horse), ocu: *Oryctolagus*
*cuniculus* (rabbit)

**References**

Ahnelt, P. K., Schubert, C., Kubber-Heiss, A., Schiviz, A., and Anger, E. (2006). Independent variation of retinal S and M cone photoreceptor topographies: A survey of four families of mammals. Vis Neurosci *23*, 429-435.

Ambros, V. (2004). The functions of animal microRNAs. Nature *431*, 350-355.

Aristotle, Forster, E. S., and Peck, A. L. (1937). Parts of animals Movement of animals ; Progression of animals / Aristotle ; with an English translation by A.L. Peck and E.S. Forster,  (Cambridge, MA: Harvard University Press).

Armstrong, E. E., Khan, A., Taylor, R. W., Gouy, A., Greenbaum, G., Thiery, A., Kang, J. T., Redondo, S. A., Prost, S., Barsh, G.*, et al.* (2021). Recent Evolutionary History of Tigers Highlights Contrasting Roles of Genetic Drift and Selection. Mol Biol Evol *38*, 2366-2379.

Bachmanov, A. A., Bosak, N. P., Lin, C., Matsumoto, I., Ohmoto, M., Reed, D. R., and Nelson, T. M. (2014). Genetics of taste receptors. Curr Pharm Des *20*, 2669-2683.

Bareil, C., Hamel, C. P., Delague, V., Arnaud, B., Demaille, J., and Claustres, M. (2001). Segregation of a mutation in CNGB1 encoding the beta-subunit of the rod cGMP-gated channel in a family with autosomal recessive retinitis pigmentosa. Hum Genet *108*, 328-334.

Barnett, R., Westbury, M. V., Sandoval-Velasco, M., Vieira, F. G., Jeon, S., Zazula, G., Martin, M. D., Ho, S. Y. W., Mather, N., Gopalakrishnan, S.*, et al.* (2020). Genomic Adaptations and Evolutionary History of the Extinct Scimitar-Toothed Cat, Homotherium latidens. Curr Biol *30*, 5018-5025 e5015.

Buckley, R. M., Davis, B. W., Brashear, W. A., Farias, F. H. G., Kuroki, K., Graves, T., Hillier, L. W., Kremitzki, M., Li, G., Middleton, R. P.*, et al.* (2020). A new domestic cat genome assembly based on long sequence reads empowers feline genomic medicine and identifies a novel gene for dwarfism. PLoS Genet *16*, e1008926.

Cabanettes, F., and Klopp, C. (2018). D-GENIES: dot plot large genomes in an interactive, efficient and simple way. PeerJ *6*, e4958.

Campbell, M. S., Holt, C., Moore, B., and Yandell, M. (2014). Genome Annotation and Curation Using MAKER and MAKER-P. Curr Protoc Bioinformatics *48*, 4 11 11-39.

Cantarel, B. L., Korf, I., Robb, S. M., Parra, G., Ross, E., Moore, B., Holt, C., Sanchez Alvarado, A., and Yandell, M. (2008). MAKER: an easy-to-use annotation pipeline designed for emerging model organism genomes. Genome Res *18*, 188-196.

Ceballos, F. C., Joshi, P. K., Clark, D. W., Ramsay, M., and Wilson, J. F. (2018). Runs of homozygosity: windows into population history and trait architecture. Nat Rev Genet *19*, 220-234.

Chandrashekar, J., Mueller, K. L., Hoon, M. A., Adler, E., Feng, L., Guo, W., Zuker, C. S., and Ryba, N. J. (2000). T2Rs function as bitter taste receptors. Cell *100*, 703-711.

Cho, Y. S., Hu, L., Hou, H., Lee, H., Xu, J., Kwon, S., Oh, S., Kim, H. M., Jho, S., Kim, S.*, et al.* (2013). The tiger genome and comparative analysis with lion and snow leopard genomes. Nat Commun *4*, 2433.

Davis, A. P., and Capecchi, M. R. (1996). A mutational analysis of the 5' HoxD genes: dissection of genetic interactions during limb development in the mouse. Development *122*, 1175-1185.

Dudchenko, O., Batra, S. S., Omer, A. D., Nyquist, S. K., Hoeger, M., Durand, N. C., Shamim, M. S., Machol, I., Lander, E. S., Aiden, A. P., and Aiden, E. L. (2017). De novo assembly of the Aedes aegypti genome using Hi-C yields chromosome-length scaffolds. Science *356*, 92-95.

Enright, A. J., John, B., Gaul, U., Tuschl, T., Sander, C., and Marks, D. S. (2003). MicroRNA targets in Drosophila. Genome Biol *5*, R1.

Fan, H., Wu, Q., Wei, F., Yang, F., Ng, B. L., and Hu, Y. (2019). Chromosome-level genome assembly for giant panda provides novel insights into Carnivora chromosome evolution. Genome Biol *20*, 267.

Figueiro, H. V., Li, G., Trindade, F. J., Assis, J., Pais, F., Fernandes, G., Santos, S. H. D., Hughes, G. M., Komissarov, A., Antunes, A.*, et al.* (2017). Genome-wide signatures of complex introgression and adaptive evolution in the big cats. Sci Adv *3*, e1700299.

Gandolfi, B., Alhaddad, H., Joslin, S. E., Khan, R., Filler, S., Brem, G., and Lyons, L. A. (2013). A splice variant in KRT71 is associated with curly coat phenotype of Selkirk Rex cats. Sci Rep *3*, 2000.

Ge, H., Lin, K., Shen, M., Wu, S., Wang, Y., Zhang, Z., Wang, Z., Zhang, Y., Huang, Z., Zhou, C., *et*
*al.* (2019). De novo assembly of a chromosome-level reference genome of red-spotted grouper
(Epinephelus akaara) using nanopore sequencing and Hi-C. Mol Ecol Resour *19*, 1461-1469.
Giani, A. M., Gallo, G. R., Gianfranceschi, L., and Formenti, G. (2020). Long walk to genomics:
History and current approaches to genome sequencing and assembly. Comput Struct Biotechnol J
*18*, 9-19.
Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., Adiconis, X., Fan,
L., Raychowdhury, R., Zeng, Q., *et al.* (2011). Full-length transcriptome assembly from RNA-Seq
data without a reference genome. Nat Biotechnol *29*, 644-652.
Grabherr, M. G., Russell, P., Meyer, M., Mauceli, E., Alfoldi, J., Di Palma, F., and Lindblad-Toh, K.
(2010). Genome-wide synteny through highly sensitive sequence alignment: Satsuma.
Bioinformatics *26*, 1145-1151.
Hamrick, M. W. (2001). Development and evolution of the mammalian limb: adaptive diversification
of nails, hooves, and claws. Evol Dev *3*, 355-363.
Hasin-Brumshtein, Y., Lancet, D., and Olender, T. (2009). Human olfaction: from genomic variation
to phenotypic diversity. Trends Genet *25*, 178-184.
Ishida, Y., David, V. A., Eizirik, E., Schaffer, A. A., Neelam, B. A., Roelke, M. E., Hannah, S. S.,
O'Brien, S. J., and Menotti-Raymond, M. (2006). A homozygous single-base deletion in MLPH
causes the dilute coat color phenotype in the domestic cat. Genomics *88*, 698-705.
Jhala, Y. V., Qureshi, Q., and Nayak, A. K. (2020). Status of tigers, copredators and prey in
India, 2018. . In,  (National Tiger Conservation Authority,
Government of India, New Delhi, and Wildlife Institute
of India, Dehradun.).
Jiang, P., Josue, J., Li, X., Glaser, D., Li, W., Brand, J. G., Margolskee, R. F., Reed, D. R., and
Beauchamp, G. K. (2012). Major taste loss in carnivorous mammals. Proc Natl Acad Sci U S A *109*,
4956-4961.
Johnson, D., Kan, S. H., Oldridge, M., Trembath, R. C., Roche, P., Esnouf, R. M., Giele, H., and
Wilkie, A. O. (2003). Missense mutations in the homeodomain of HOXD13 are associated with
brachydactyly types D and E. Am J Hum Genet *72*, 984-997.
Kaelin, C. B., Xu, X., Hong, L. Z., David, V. A., McGowan, K. A., Schmidt-Kuntzel, A., Roelke, M. E.,
Pino, J., Pontius, J., Cooper, G. M., *et al.* (2012). Specifying and sustaining pigmentation patterns in
domestic and wild cats. Science *337*, 1536-1541.
Kalvari, I., Argasinska, J., Quinones-Olvera, N., Nawrocki, E. P., Rivas, E., Eddy, S. R., Bateman, A.,
Finn, R. D., and Petrov, A. I. (2018). Rfam 13.0: shifting to a genome-centric resource for non-coding
RNA families. Nucleic Acids Res *46*, D335-D342.
Kardos, M., Taylor, H. R., Ellegren, H., Luikart, G., and Allendorf, F. W. (2016). Genomics advances
the study of inbreeding depression in the wild. Evol Appl *9*, 1205-1218.
Katagiri, S., Akahori, M., Sergeev, Y., Yoshitake, K., Ikeo, K., Furuno, M., Hayashi, T., Kondo, M.,
Ueno, S., Tsunoda, K., *et al.* (2014). Whole exome analysis identifies frequent CNGA1 mutations in
Japanese population with autosomal recessive retinitis pigmentosa. PLoS One *9*, e108721.
Khan, A., Patel, K., Bhattacharjee, S., Sharma, S., Chugani, A. N., Sivaraman, K., Hosawad, V.,
Sahu, Y. K., Reddy, G. V., and Ramakrishnan, U. (2020). Are shed hair genomes the most effective
noninvasive resource for estimating relationships in the wild? Ecol Evol *10*, 4583-4594.
Khan, A., Patel, K., Shukla, S., Viswanathan, A., van der Valk, T., Borthakur, U., Nigam, P.,
Zachariah, A., Jhala, Y., Kardos, M., and Ramakrishnan, U. (2021). Genomic evidence for
inbreeding depression and purging of deleterious genetic variation in Indian tigers. bioRxiv.
Kim, S., Cho, Y. S., Kim, H. M., Chung, O., Kim, H., Jho, S., Seomun, H., Kim, J., Bang, W. Y., Kim,
C., *et al.* (2016). Comparison of carnivore, omnivore, and herbivore mammalian genomes with a new
leopard assembly. Genome Biol *17*, 211.
Korf, I. (2004). Gene finding in novel genomes. BMC Bioinformatics *5*, 59.
Lee, R. C., Feinbaum, R. L., and Ambros, V. (1993). The C. elegans heterochronic gene lin-4
encodes small RNAs with antisense complementarity to lin-14. Cell *75*, 843-854.

Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics *25*, 1754-1760.
Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin,
R., and Genome Project Data Processing, S. (2009a). The Sequence Alignment/Map format and
SAMtools. Bioinformatics *25*, 2078-2079.
Li, X., Glaser, D., Li, W., Johnson, W. E., O'Brien, S. J., Beauchamp, G. K., and Brand, J. G.
(2009b). Analyses of sweet receptor gene (Tas1r2) and preference for sweet stimuli in species of
Carnivora. J Hered *100 Suppl 1*, S90-100.
Li, X., Li, W., Wang, H., Cao, J., Maehashi, K., Huang, L., Bachmanov, A. A., Reed, D. R., Legrand-
Defretin, V., Beauchamp, G. K., and Brand, J. G. (2005). Pseudogenization of a sweet-receptor
gene accounts for cats' indifference toward sugar. PLoS Genet *1*, 27-35.
Lieberman-Aiden, E., van Berkum, N. L., Williams, L., Imakaev, M., Ragoczy, T., Telling, A., Amit, I.,
Lajoie, B. R., Sabo, P. J., Dorschner, M. O*., et al.* (2009). Comprehensive mapping of long-range
interactions reveals folding principles of the human genome. Science *326*, 289-293.
Liu, Y. C., Sun, X., Driscoll, C., Miquelle, D. G., Xu, X., Martelli, P., Uphyrkina, O., Smith, J. L. D.,
O'Brien, S. J., and Luo, S. J. (2018). Genome-Wide Evolutionary Analysis of Natural History and
Adaptation in the World's Tigers. Curr Biol *28*, 3840-3849 e3846.
Lowe, T. M., and Eddy, S. R. (1997). tRNAscan-SE: a program for improved detection of transfer
RNA genes in genomic sequence. Nucleic Acids Res *25*, 955-964.
Luo, S. J., Liu, Y. C., and Xu, X. (2019). Tigers of the World: Genomics and Conservation. Annu Rev
Anim Biosci *7*, 521-548.
Lyons, L. A., Foe, I. T., Rah, H. C., and Grahn, R. A. (2005). Chocolate coated cats: TYRP1
mutations for brown color in domestic cats. Mamm Genome *16*, 356-366.
Mondol, S., Karanth, K. U., and Ramakrishnan, U. (2009). Why the Indian subcontinent holds the
key to global tiger recovery. PLoS Genet *5*, e1000585.
Montague, M. J., Li, G., Gandolfi, B., Khan, R., Aken, B. L., Searle, S. M., Minx, P., Hillier, L. W.,
Koboldt, D. C., Davis, B. W*., et al.* (2014). Comparative analysis of the domestic cat genome reveals
genetic signatures underlying feline biology and domestication. Proc Natl Acad Sci U S A *111*,
17230-17235.
Murchison, E. P., Schulz-Trieglaff, O. B., Ning, Z., Alexandrov, L. B., Bauer, M. J., Fu, B., Hims, M.,
Ding, Z., Ivakhno, S., Stewart, C*., et al.* (2012). Genome sequencing and analysis of the Tasmanian
devil and its transmissible cancer. Cell *148*, 780-791.
Nawrocki, E. P., and Eddy, S. R. (2013). Infernal 1.1: 100-fold faster RNA homology searches.
Bioinformatics *29*, 2933-2935.
Ohsawa, R., and Kageyama, R. (2008). Regulation of retinal cell fate specification by multiple
transcription factors. Brain Res *1192*, 90-98.
Pemberton, T. J., Absher, D., Feldman, M. W., Myers, R. M., Rosenberg, N. A., and Li, J. Z. (2012).
Genomic patterns of homozygosity in worldwide human populations. Am J Hum Genet *91*, 275-292.
Phillippy, A. M. (2017). New advances in sequence assembly. Genome Res *27*, xi-xiii.
Poddar-Sarkar, M., and Brahmachary, R. L. (2014). Pheromones of Tiger and Other Big Cats. In
Neurobiology of Chemical Communication, C. Mucignat-Caretta, ed. (Boca Raton (FL)).
Pontius, J. U., Mullikin, J. C., Smith, D. R., Agencourt Sequencing, T., Lindblad-Toh, K., Gnerre, S.,
Clamp, M., Chang, J., Stephens, R., Neelam, B*., et al.* (2007). Initial sequence and comparative
analysis of the cat genome. Genome Res *17*, 1675-1689.
Putnam, N. H., O'Connell, B. L., Stites, J. C., Rice, B. J., Blanchette, M., Calef, R., Troll, C. J.,
Fields, A., Hartley, P. D., Sugnet, C. W*., et al.* (2016). Chromosome-scale shotgun assembly using
an in vitro method for long-range linkage. Genome Res *26*, 342-350.
Ripple, W. J., Estes, J. A., Beschta, R. L., Wilmers, C. C., Ritchie, E. G., Hebblewhite, M., Berger, J.,
Elmhagen, B., Letnic, M., Nelson, M. P*., et al.* (2014). Status and ecological effects of the world's
largest carnivores. Science *343*, 1241484.
Roots, C. (2006). Nocturnal animals,  (Westport, Conn.: Greenwood Press).

Sagar, V., Kaelin, C. B., Natesh, M., Reddy, P. A., Mohapatra, R. K., Chhattani, H., Thatte, P.,
Vaidyanathan, S., Biswas, S., Bhatt, S., *et al.* (2021). High frequency of an otherwise rare phenotype
in a small and isolated tiger population. Proc Natl Acad Sci U S A *118*.
Sharpe, P. T. (1995). Homeobox genes and orofacial development. Connect Tissue Res *32*, 17-25.
Simao, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., and Zdobnov, E. M. (2015).
BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs.
Bioinformatics *31*, 3210-3212.
Simpson, G. G. (1941). The function of saber-like canines in carnivorous mammals, Vol 1130, (New
York City: The American Museum of Natural History).
Stanke, M., and Morgenstern, B. (2005). AUGUSTUS: a web server for gene prediction in
eukaryotes that allows user-defined constraints. Nucleic Acids Res *33*, W465-467.
Suryamohan, K., Krishnankutty, S. P., Guillory, J., Jevit, M., Schroder, M. S., Wu, M., Kuriakose, B.,
Mathew, O. K., Perumal, R. C., Koludarov, I., *et al.* (2020). The Indian cobra reference genome and
transcriptome enables comprehensive identification of venom toxins. Nat Genet *52*, 106-117.
Sutton, J. T., Helmkampf, M., Steiner, C. C., Bellinger, M. R., Korlach, J., Hall, R., Baybayan, P.,
Muehling, J., Gu, J., Kingan, S., *et al.* (2018). A High-Quality, Long-Read De Novo Genome
Assembly to Aid Conservation of Hawaii's Last Remaining Crow Species. Genes (Basel) *9*.
Task Force, I. B. f. W. L. (1972). Project Tiger: A Proposal for Preservation of the Tiger (Panthera
tigris tigris Linn.) in India. In, (New Delhi, India: Ministry of Agriculture, Government of India), p. p4.
Thomas, B. L., and Sharpe, P. T. (1998). Patterning of the murine dentition by homeobox genes. Eur
J Oral Sci *106 Suppl 1*, 48-54.
Thomas, B. L., Tucker, A. S., Qui, M., Ferguson, C. A., Hardcastle, Z., Rubenstein, J. L., and
Sharpe, P. T. (1997). Role of Dlx-1 and Dlx-2 genes in patterning of the murine dentition.
Development *124*, 4811-4818.
Tummala, H., Ali, M., Getty, P., Hocking, P. M., Burt, D. W., Inglehearn, C. F., and Lester, D. H.
(2006). Mutation in the guanine nucleotide-binding protein beta-3 causes retinal degeneration and
embryonic mortality in chickens. Invest Ophthalmol Vis Sci *47*, 4714-4718.
Vastardis, H., Karimbux, N., Guthua, S. W., Seidman, J. G., and Seidman, C. E. (1996). A human
MSX1 homeodomain missense mutation causes selective tooth agenesis. Nat Genet *13*, 417-421.
Vickaryous, M. K., and Sire, J. Y. (2009). The integumentary skeleton of tetrapods: origin, evolution,
and development. J Anat *214*, 441-464.
Vincent, A., Audo, I., Tavares, E., Maynes, J. T., Tumber, A., Wright, T., Li, S., Michiels, C.,
Consortium, G. N. B., Condroyer, C., *et al.* (2016). Biallelic Mutations in GNB3 Cause a Unique
Form of Autosomal-Recessive Congenital Stationary Night Blindness. Am J Hum Genet *98*, 1011-
1019.
Waterhouse, R. M., Seppey, M., Simao, F. A., Manni, M., Ioannidis, P., Klioutchnikov, G.,
Kriventseva, E. V., and Zdobnov, E. M. (2018). BUSCO Applications from Quality Assessments to
Gene Prediction and Phylogenomics. Mol Biol Evol *35*, 543-548.
Wurster-Hill, D. H., and Gray, C. W. (1973). Giemsa banding patterns in the chromosomes of twelve
species of cats (Felidae). Cytogenet Cell Genet *12*, 388-397.
Xu, X., Dong, G. X., Schmidt-Kuntzel, A., Zhang, X. L., Zhuang, Y., Fang, R., Sun, X., Hu, X. S.,
Zhang, T. Y., Yang, H. D., *et al.* (2017). The genetics of tiger pelage color variations. Cell Res *27*,
954-957.
Yang, Z. (2007). PAML 4: phylogenetic analysis by maximum likelihood. Mol Biol Evol *24*, 1586-
1591.
Yang, Z., Wong, W. S., and Nielsen, R. (2005). Bayes empirical bayes inference of amino acid sites
under positive selection. Mol Biol Evol *22*, 1107-1118.
Yokohama-Tamaki, T., Ohshima, H., Fujiwara, N., Takada, Y., Ichimori, Y., Wakisaka, S., Ohuchi,
H., and Harada, H. (2006). Cessation of Fgf10 signaling, resulting in a defective dental epithelial
stem cell compartment, leads to the transition from crown to root formation. Development *133*, 1359-
1366.
Young, J. M., Massa, H. F., Hsu, L., and Trask, B. J. (2010). Extreme variability among mammalian
V1R gene families. Genome Res *20*, 10-18.

Zhang, Y. D., Chen, Z., Song, Y. Q., Liu, C., and Chen, Y. P. (2005). Making a tooth: growth factors,
transcription factors, and stem cells. Cell Res *15*, 301-316.
854

**a**

**b**

① Endurance — ADCY5, MEF2C, PCK1, SLC5A7, IL15, PGC-1α, TFE3, TFEB, TRPV1

② Vision — NRL, SAG, GNAT1, GNB3, PDE6A, PDE6B, PDE6G, CNGA1, CNGB1

③ Olfaction — CNGA2, CNGA4, OR2B11, OR13H1, OR10K1, OR10V1, VN1R4

④ Neurotransmitter — NPFFR1, NPFFR2, PCDHB4, PCDHB6, PCDHB14, GRIN2C, GRIA, NPY, PY1R, NTF2, NTF4

⑥ Taste — TAS1R1, TAS2R1, TAS2R3, TAS2R5, TAS2R7, TAS2R38, TAS2R42, TAS2R346

⑧ Tooth — BMP2, BMP4, WNT1, WNT2, WNT3A, WNT4, WNT5B, WNT7A, WNT10A, WNT10B, WNT11, PIT1, PIT2, BARX1, BARX2, PAX1, PAX9, PAX7, MSX-1, MSX-2, FGF4, FGF6, FGF9, FGF10, FGF11, FGF13, FGF16, FGF18, FGF19, FGF20, FGF22, FGFBP1, LHX3, LHX4, LHX5, LHX6, LHX8, LHX9

⑤ Patterning — SLC45A2, MLPH, TYR2, ASIP, KIT, KRT71, TYR1, ANPEP, CORIN

⑦ Claw — HOXA2, HOXA4, HOXA9, HOXA10, HOXA11, HOXA13, MSX-1, MSX-2

**c**

transmembrane ——  inside ——  outside ——

**d**

**a**

**b**

**c**

Supplementary Figures

**Supplementary Figure 1:** Synteny between MC and SI tiger genome assemblies

**Supplementary Figure 2.** Repeat element distribution in MC (blue bars) and SI (orange bars) genomes. LINE: long interspersed nuclear element; LTR: long terminal repeat; SINE: short interspersed nuclear element, SSR: simple short repeat

**Supplementary Figure 3.** Treemap plot of gene families identified in MC genome