Chromatin-modifying enzymes as modulators of nuclear size during lineage differentiation

Lingjun Meng[1,*], Evgeny Zatulovskiy[2], Yohei Shibuya[1], Katie Marie Schaukowitch[1], Gernot

Neumayer[1], Jan M. Skotheim[2], Marius Wernig[1].

[1]Institute for Stem Cell Biology and Regenerative Medicine, Department of Pathology, Stanford

University, CA 94305, USA

[2]Department of Biology, Stanford University, Stanford, CA 94305, USA.

*Correspondence: lingjun@stanford.edu, menglingjundl@gmail.com

Key word: histone demethylases, nuclear size, differentiation, cancer.

Abstract

The mechanism of cell size determination and alteration during normal lineage development

and cancer pathologies which is not fully understood. As recently reported, chromatin modification

can change nuclear morphology. Therefore, we screened a range of pharmacological chemical

compounds that impact the activity of chromatin-modifying enzymes, in order to get a clue of the

specific types of chromatin-modifying enzymes that remarkably effect nuclear shape. We used

three assays to measure the nuclear size and cell size. The novel coulter counter was presented

here and validated with using flow cytometry and confocal fluorescence microscopy to measure

size. We found that interrupted activity of chromatin-modifying enzymes is associated with nuclear

shape abnormalities. Furthermore, the activity of chromatin-modifying enzymes perturbs cell fate

determination in cellular maintenance and lineage commitment. Our results present a new idea in

understanding the relation of chromatin-modifying enzymes and nuclear size during lineage

differentiation.

Introduction

As reported, nuclear volume is an important morphological parameter that can change during

cancer progression [1]. After confocal microscopy was discovered and broadly used to observe the

detailed structure of specific objects within the cell, the most fascinated cellular organelle - nucleus

was researched more carefully. Recent studies have also highlighted that nuclear shape can

influence fate decisions [2]. Embryonic stem cells (ESCs) differentiation is accompanied by an

increase in chromatin condensation [3] leading to an increase in nuclear stiffness [4] and a decrease in

nuclear size. As known, the nuclear size of stem cells is extremely bigger than terminal

differentiated cells. Cell differentiation is corresponding with chromatin marks reset, which raises

the question of whether chromatin's contribution to nuclear mechanics affects nuclear morphology

as well as cell fate. It is therefore critical to identify the contributions of various chromatin-

modifying enzymes that are relevant to cell fate decisions, and consequently display nuclear size

alteration.

While previous studies have focused on utilizing microscopy examination to achieve the

visualization of nucleus via its two-dimensional structure, here we employed two novel assays,

such as flow cytometry [5,6] and coulter counter [7] to measure the nuclear and cell size via three-

dimensional structure. We used flow cytometry to measure forward scatter (FSC) of individual

cells [8], cell volume was identified by forward scatter area (FSC-A), and nuclear volume was

identified by Hoechst-W (width) after staining with Hoechst. By using coulter counter, cells flow

through a measurement chamber where they displace an isotonic solution to cause a volume-

proportional change in electrical impedance [7]. Coulter counter measures cell number and size in

high throughput as well as flow cytometry. The high throughput quantitative analysis of 3D nuclear

architecture facilitates better investigation of a great variability of nuclear shapes, in order to fully

understand the nuclear contour related cell fate decisions also including cancer cell fate decisions.

Nuclear architecture is impacted via chromosome structure which is determined by histone

modifications. The cellular differentiation from stem cells to differentiated cells which is

corresponding with nuclear size reduction as well as chromosome condense. Histone demethylases

have also been reported to be critical enzymes that regulate the balance between self-renewal and

differentiation in hESCs (human embryonic stem cells) [9-12]. Accordingly, we broadly utilized

chromatin-modifying enzymes inhibitors to treat stem cells such as hESCs, NPCs (neural

progenitor cells), and later a cancer cell line such as leukemia cells. We demonstrated that the

irregularities of nuclear contour were highly related to pharmacological inhibition of chromatin-

modifying enzyme activity. As known, upregulation or downregulation of histone demethylases is

observed in various types of cancer and is associated with higher rates of malignancy and relapse

after treatment [13]. In particular, GASC1 (gene amplified in squamous cell carcinoma, also known

as KDM4C, a histone demethylase) promotes chromosomal instability and transcription initiation,

therefore, plays a causative role in cancer cells driven by its dysregulation [14]. GASC1 was found

aberrant expression in esophageal squamous carcinomas [15], medulloblastomas [16], glioma [17], breast

cancer [18], and prostate cancer [19]. In addition, GASC1 plays an essential role in the transition of

mouse ESCs (mESCs) to endothelial cells [20]. Consistently, we demonstrated that upregulation of

GASC1 impairs the differentiation of undifferentiated hESCs to terminal differentiated neural cells.

Moreover, upregulation of GASC1 impairs the proliferation of leukemia cells due to cell cycle

arrest in the G0/G1 phase and indicated misshapen nucleus. Taken together, epigenetic

perturbations are likely to be the cause for misregulation of nuclear size and of certain cancer cell

fates [21].

Result

The nuclear size is associated with cell fate decision and chromatin-modifying enzyme

activity

Size and shape are distinctive aspects of nuclear structure. Nuclear size impacts cell function,

and is altered according to the specific status of cell types during cell development [22]. It is unclear

which modulators effect the altered nuclear size during cell fate transformation. To investigate the

distinct nuclear size of various cell types, we employed highly efficient differentiation methods to

interrogate the size change of nucleus during lineage differentiation. To differentiate NPCs from

hESCs, we used two molecular compounds, SB431542 and LDN193189, to inhibit the TGF-β

(transforming growth factor beta) and BMP (bone morphogenetic protein) pathways, which are

also called Dual SMAD inhibitors [22,23]. In addition, as we previously used, overexpression of Ngn2

(neuronal gene neurogenin-2) could converts hESCs into iN cells (induced neuronal cells) with

nearly 100% yield in less than two weeks. The hESCs-derived iN cells grow long axon that conduct

electrical impulses away from the neuron's cell body [24]. We utilized 300 nM DAPI to stain hESCs,

hESC-derived NPCs, hESC-derived iN cells for 5 min, then examining the dynamic change of the

nuclear size via measuring the area of nucleus under a fluorescence microscope (Fig. 1a).

Fluorescence images of nucleus stained with DAPI were acquired through imaging at 40×

resolution. Notably, it was shown that there is a reduced size of approximately 40% from hESCs

to NPCs, and 24% decrease from NPCs to iN cells (Fig. 1b). Consistently as reported, the nuclear

size of hESCs is bigger than NPCs, and the nuclear size of NPCs is bigger than iN cells. In addition,

it was shown that the nuclear size of hESCs and NPCs are in a wider range than iN cells, which

indicates that there is a potential of stem cells in nuclear dynamics. The different sizes of nuclei

may correspond with hESCs, NPCs, and neural cells characterized with distinct unique chromatin

architecture. During the reprogramming of cell identity and cell fate determination,

heterochromatin plays a critical role. H3K9me3 (Tri-methylation of histone H3 lysine 9) is a

histone mark associated with heterochromatin, has emerged as a key player in repressing lineage-

inappropriate genes, and shielding them from activation by transcription factors. We investigated

the H3K9me3 expression in hESCs and hESC-derived NPCs by immunofluorescence staining. It

was shown that there are high levels of expression of H3K9me3 in the nucleus (Fig. 1c). To directly

address the role of chromatin in nuclear morphology, we employed a range of commercially

available chemical compound inhibitors of the enzymes that modulate histone modification state

to manipulate chromatin alterations to treat hESCs and hESC-derived NPCs (Fig. 1d, e). These

available inhibitors of chromatin-modifying enzymes that were represented the various functions

of chromatin regulation, which helped us to investigate broadly to chromatin-modifying enzymes

associated with nuclear alteration. We purchased these inhibitors from tocris as method shown. As

reported, GSK-J1 is a highly potent inhibitor of the H3K27 histone demethylases JMJD3 (KDM6B)

and UTX (KDM6A), likewise inhibits KDM5B, KDM5C, and KDM5A to increase H3K27me3.

Histone demethylases inhibitor increased heterochromatin levels and chromatin-based nuclear

stiffness. Heterochromatin is highly condensed DNA that leads to gene silencing [25]. R2-HG is

oncometabolite produced by mutant isocitrate dehydrogenase 1 (IDH1) and IDH2, inhibits

KDM4A in murine embryonic fibroblasts as well as increases H3K9me3, H3K36me3. 5-Aza is a

DNA methyltransferase inhibitor that induces demethylation. GSK-LSD1 is a potent and selective

lysine specific demethylase 1 (LSD1) inhibitor, increasing H3K4me3, H3K9me3, H3K27me3.

IOX1 is a histone demethylase JMJD inhibitor, increasing H3K36me3, H3K9me3, H3K27me3.

JIB04 is a pan Jumonji histone demethylase inhibitor that inhibits H3K4 demethylases, decreases

H3K36me3, H3K9me3. NSC636819 is a KDM4A/KDM4B inhibitor, increasing H3K9me3. PBIT

is a JARID1 (Jumonji AT-Rich Interactive Domain 1) inhibitor that increases H3K4me3. C646 is

a inhibit histone acetyltransferase, which corresponds predominantly to decompacted euchromatin,

allowing transcription factors to bind to regulatory sites on DNA, in order to cause transcriptional

activation [26]. Through the treatment with compounds function as chromatin-modifying enzyme

inhibitors, we found that 5-Aza treatment enlarges the nuclear size of both hESCs and hESC-

derived NPCs remarkably (Fig. 1d, e), also some other inhibitors. In contrast, 5-Aza treatment

slightly reduced the nuclear size of hESC-derived iN cells (Fig. 1f). The great increase in nuclear

size of hESCs from 5-Aza treatment that indicated that modulation of chromatin-modifying

enzymes is enabled to change nuclear size (Fig. 1g). It suggested that perturbation of chromatin-

modifying enzyme activity alters nuclear size in normal cells.

Inhibition of chromatin-modifying enzyme enlarges hESCs nuclear size

To investigate the nuclear size and cell size in various methods, we employed flow cytometry

and coulter counter to measure the nuclear size and cell size, then compared with confocal

fluorescence microscopy by DAPI staining used above. Flow cytometry optical scatter was used

to measure the size of live cells. This parameter is a measurement of the amount of the laser beam

that passes around the cell. This gives a relative size for the cell. Therefore, the forward scatter

area (FSC-A) parameter was used as a readout for cell size. Hoechst 33342 is a dye that binds to

AT-rich sequences of double-stranded DNA. So, we used Hoechst-W (width) indicates the nuclear

size (Fig. 2a, b, c). From the flow cytometric size measurements of hESCs after inhibitors

treatment for 7 days, which was shown that cell size distributions of hESCs were affected by these

chromatin-modifying enzymes inhibitors, especially in 5-Aza and IOX1 treatment compared to

DMSO control (Fig. 2d, e). The nuclear size indicated by Hoechst-W was increased 13% in 5-Aza

treatment, and the cell size indicated by FSC-A was increased 20%, which suggested the major

contributor to cell size change is the nucleus. As above shown, we also measured the nucleus size

of hESCs by DAPI staining after inhibitors treatment for 7 days (Fig. 1d). It was shown that 5-Aza

significantly enhances the nuclear size of hESCs (Fig. 1d). Treating with high concentrations of

IOX1 which revealed the cytotoxic effects on hESCs. Therefore, we tested a series of

concentrations of compounds for hESCs treatment. The high concentrations of inhibitors caused

cell death of hESCs, such as 100 mM GSK-J1, 100 mM IOX1, 1 mM JIB04, 10 mM PBIT, 25 mM

C646, especially, 10 μM, 1 mM, 10 mM 5-Aza induced numerous of hESCs death. The high

concentrations of inhibitors have cytotoxic effects on cells and impact cells alive. However, hESCs

grow well in 60 μM, 1 mM, 10 mM GSK-J1, 100 mM R2HG, 10 mM and 100 mM GSK-LSD1,

1 mM and 10 mM IOX1, 10 μM and 100 μM JIB04, 10 μM and 10 mM NSC636819, 100 μM and

1 mM PBIT, 250 μM and 2.5 mM C646 treatment. Furthermore, we performed coulter counter

analysis to measure the corresponding cell volume of hESCs in a dynamics pattern after inhibitors

treatment for 6 days. Coulter counter is an apparatus for counting and sizing particles suspended

in electrolytes, which provides a relative estimate of the regularity of each nucleus. After broadly

measuring the volume of each cell with treatment, it demonstrated that cellular volume of hESCs

measured as FL (Femtoliter) was increased in 5-Aza and IOX1 treatment compared with DMSO

control, which was consistent with flow cytometry measurement (Fig. 2f). The three different

assays were all indicated that compounds function as chromatin-modifying enzyme inhibitors

which effect nuclear size alteration, possibly through modulating histone dynamics. It had been

shown that utilizing histone methyltransferase inhibitors to decrease heterochromatin results in a

softer nucleus and nuclear blebbing, without perturbing lamins. Histone demethylase inhibitors

increase heterochromatin and chromatin nuclear rigidity [27]. Given that measurements in these

reports demonstrate that there is evidence that nuclear size abnormalities can be caused by

alterations to chromatin regulatory proteins.

Inhibition of chromatin-modifying enzyme effects nuclear size and cell identification of NPCs

To gain further insights into whether these inhibitors not only impact cell death also effect

additional features, such as cell fate decisions, we treated the hESC-derived NPCs with DMSO, 5-

Aza, IOX1, JIB04, and PBIT. The concentration we used that was tested and same concentration

was used as hESCs treatment. Then imaged hESC-derived NPCs after staining for Sox1 (green),

PI (red), and DAPI (blue) (Fig. 3a). As known, Sox1 is a neural stem cell marker. It was observed

that the expression of Sox1 was significantly reduced in a large number of NPCs in 5-Aza, JIB04,

and PBIT treatment. To eliminate the possibility of Sox1 expression decrease caused by cell death,

we performed propidium iodide (PI) staining to NPCs after inhibitors treatment. As known, PI is

not permanent to live cells but is used to detect dead cells in a population. It was shown that Sox1

silence was not caused by cell death due to PI staining being negative. Except only IOX1 increased

PI staining in NPCs which suggested that 10 mM IOX1 treatment induces cytotoxic responses of

NPCs occur as a consequence of cell death. It suggested that the compound inhibitors of chromatin-

modifying enzymes perturb cell identification. We also measured the cell size of hESC-derived

NPCs by the coulter counter after inhibitors treatment for 9 days (Fig. 3b). The treatment of 5-Aza,

GSK-LSD1, IOX1, and JIB04 enhanced the nuclear size of NPCs. Collectively, inhibition of

chromatin-modifying enzymes activity enabled to impact nuclear size in both hESCs and NPCs

which consequently leads cell fate to aberrant states.

Inhibition of chromatin-modifying enzyme effects leukemia cell nuclear architecture

As above discussed, chromatin-modifying enzyme activity is highly associated with nuclear

size alteration in normal stem cells during lineage differentiation. As reported, cancer cells are

known to develop larger nuclei as they become more malignant and increase the nuclear-to-

cytoplasmic ratio [28]. The mechanisms that underlie the increase in nuclear size in malignant cells

remain largely unknown. Abnormal nuclear morphology has been used for cancer diagnoses for

nearly a century in tests such as the Pap smear [29]. The novel assay for detecting nuclear shape

alteration of 3D structure is helpful to better understand. Therefore, we utilized flow cytometric

size measurements to measure the nuclear size of K562 leukemia cells, in order to investigate the

distinctive features of chromatin-modifying enzyme activity in cancer cells compared to normal

stem cells. We performed flow cytometry to measure the nuclear size and cell size of K562

leukemia cells after compound inhibitors treatment for 5 days (Fig. 4a, b, c). We tested a series of

concentrations of compound inhibitors for K562 cells treatment. Consistent with hESCs, high

concentrations of inhibitors caused cell death of K562 cells. Especially, high dose and long-time

of 5-Aza and PBIT treatment induce cell death. In addition, we photographed the DAPI stained

nucleus of K562 cells treated with DMSO as control and 5-Aza, IOX1, GSK-LSD1, and JIB04. It

was shown that K562 cells treated with inhibitors have a bigger size of nucleus and twisted shape

as clover, especially in IOX1, GSK-LSD1, and JIB04 treatment (Fig. 4d). It demonstrated that

there is the irregularity of nuclear contour in inhibitors treatment from the visible fluorescent

images. The most studied types of nuclear morphological abnormalities are protrusions from the

normally ellipsoidal nucleus, termed "blebs." K562 cells showed a "blebs" shape and seem prefer

to have a twisted nuclear shape but not die as quickly as hESCs and NPCs did. We measured the

nuclear size by DAPI staining after K562 cells were treated with inhibitors, it was shown that the

increased percentage of nuclear size of K562 cells is 5% ~ 44% in inhibitors treatment compared

to DMSO control, especially higher in 5-Aza and IOX1 treatment (Fig. 4e). It all suggested that

not only nuclear size change but also nucleus misshapen are indicated the pathological change of

K562 leukemia cells induced by inhibition of chromatin-modifying enzymes activity. Furthermore,

to measure nuclear size and cell size of cancer cells in 3D, we utilized flow cytometry, coulter

counter, and DAPI staining to measure the nuclear size and cell size of K562 leukemia cells. The

size of cancer cells was more similar to hESCs but bigger than NPCs (Fig. 4f). After inhibitors

treatment for 7 days (10 μM GSK-J1, 100 μM R2HG, 1 μM 5-Aza, 10 μM GSK-LSD1, 10 μM

IOX1, 100 μM JIB04, 10 μM NSC636819, 1 μM PBIT, 2.5 μM C646), compared to DMSO control,

treatment of 5-Aza and IOX1 markedly lead to the enlargement of the nuclear size and cell size of

K562 leukemia cells (Fig. 4f). Comparing these quantitative approaches, which we validated with

measurement by confocal microscopy, it reveals that all assays support chromatin-modifying

enzyme activity directly effects nuclear size. Except for lamina perturbation affecting nuclear

abnormalities [30], it also suggested that enlarged and twisted nuclear size effect the survival of K562

cells which is associated with chromatin-modifying enzyme activity.

Upregulated GASC1 impairs differentiation from hESCs to iN cells

As we previously discussed, the nuclear size dramatically reduced from hESCs to NPCs, and

to iN cells during lineage differentiation. Additionally, we want to validate the specificity of the

targeted chromatin-modifying enzyme instead of using chemical compounds. So, we selected three

represented histone enzymes such as GASC1, DNMT3A, and KDM4A, based on the screening of

inhibitors of chromatin-modifying enzyme activity, to investigate the impact of chromatin-

modifying enzymes in lineage differentiation. GASC1 (lysine-specific demethylase 4C, KDM4C)

is a histone demethylase as well as KDM4A (lysine-specific demethylase 4A), they both belong to

the families of lysine demethylases (KDM). DNMT3A (DNA methyltransferase 3A) catalyzes 5-

methylcytosine methylation. 5-Aza inhibits DNA methyltransferase (DNMT). We used the

doxycycline (dox)-inducible lentiviral vector system to transduce empty vector plasmid as control

and GASC1, DNMT3A, KDM4A into the hESCs H1 cell line, separately. As mentioned above,

we used the same approach to induce neuronal cells from differentiating hESCs by Ngn2

overexpression [24]. It was shown that there is a significantly high efficiency of wt hESCs

differentiating to iN cells as nearly 100%. The iN cells display a mature neural morphology with

axon extension and packaged smaller cell nucleus (Fig. 5a). We stained hESC-iN cells with a

neuronal marker Tuj1 (tubulin beta-III) and a pluripotent marker Sox2, it demonstrated that iN

cells differentiated from wt hESCs showed Tuj1 positive expression and long axon extension, no

Sox2 indicated hESCs left, which also occurred in KDM4A overexpressed hESC-iN cells. In

contrast, hESC-iN cells with induced overexpression of GASC1 and DNMT3A that grew few

extend axons. In addition, a large number of cells were Sox2 positive during differentiation (Fig.

5a). After measurement analysis of neurite length at 14 days of iN cell differentiation, we found

that hESC-iN cells with GASC1 and DNMT3A overexpression grew remarkably shorter neurite

compared to wt and KDM4A overexpressed iN cells (Fig. 5b). Perturbation of chromatin-

modifying enzymes activity alters the molecular and cellular identity of stem cells. Overall, these

data suggested that chromatin-modifying enzymes can drive lineage specification as well as assist

nuclear reconstruction, in order to facilitate a smoother transition from stem cells to differentiated

cells.

GASC1 promotes aberrant nuclear structure and cell cycle arrest of leukemia cells

As mentioned above, these chromatin-modifying enzymes effect lineage differentiation of

hESCs. These hESC-iN cells with GASC1 and DNMT3A aberrant expression retain their

proliferation capability, fail differentiation, and acquire resistance to cell death. As previously

reported, IDH1 mutated cells generate R(2)-2-hydroxyglutarate (2HG) [31], which causes GASC1

mediated hypermethylation [19,32]. IDH1 gene mutations occur frequently in AML (acute

lymphoblastic leukemia). Accordingly, we investigated the role of GASC1 in K562 leukemia cells

differentiation. Basically, healthy and as cultured K562 cells are spherical in shape without any

structural irregularities and distortions, and grow in suspension (Fig. 6a). After being differentiated

by Ngn2 overexpression for 14 days, it was shown that a portion of K562 cells grew spike, and

Tuj1 was not abundantly expressed (Fig. 6a). K562 cells are hard to differentiate. When adding

dox in cell culture for inducing GASC1 overexpression, the majority of K562 leukemia cells began

to change their morphology from smooth surfaces of spherical shape to twisted cellular shape

compared with no dox-induced K562 cells (Fig. 6b, c). The abnormal nuclear morphology and

protrusions termed "blebs" are both associated with chromatin alterations [27]. To validate the cell

shape change corresponding to nucleus change, we stain K562 cells with DAPI. Under microscope

examination of GASC1 overexpressed K562 cells with or without dox, we found that it is

consistent with cell shape change in dox-induced overexpressed K562 cells, that showed nucleus

enlarges and shapes as clover (Fig. 6d, e). To investigate the twisted shape consequently provides

benefits of growth of the K562 cells or not, we employed the Brdu cell proliferation assay to

interrogate the cell cycle of K562 cells growth. The dox-induced K562-GASC1 cells significantly

proliferate slower than the no dox-induced K562-GASC1 cells (Fig. 6f, g). Meanwhile, we count

the cell number of K562 cells with or without dox-induced GASC1 overexpression, it reveals that

there is a significant reduction in the growth of K562 cells with GASC1 overexpression (Fig. 6h).

To explore the mechanism of K562 cells with GASC1 expression was able to stop proliferation at

the genetic level, we selected a range of key regular genes function as cell cycle effectors, such as

Rbl2, PRDM1, p21, p53, E2F4, lin9, and CDK2 [33,34]. Rbl2 (RB Transcriptional Corepressor Like

2) is a key regulator of entry into cell division [35], directly involved in heterochromatin formation

by stabilizing histone methylation, thus maintaining overall chromatin structure. In particular, Rbl2

acts as a transcription repressor by recruiting chromatin-modifying enzymes to promoters [35].

PRDM1 regulates the expression of genes involved in cell cycle. p21 is a cyclin-dependent kinase

(CDK) inhibitor that promotes cell cycle arrest, also acts as a gene downstream of p53 with linking

DNA damage to cell cycle arrest [33]. E2F4 is a key regulator to achieve and maintain cellular

quiescence by repressing cell cycle-regulated genes and plays an important role in the suppression

of proliferation-associated genes. Depletion of Lin9 results in an accumulation of cells in G2 and

M phases. CDK2 (Cyclin-dependent kinase 2) is a key cell cycle regulator, inactivating

phosphorylation of the RB1 and controlling both G1/S and G2/M transitions [34]. Based on the qPCR

results, we found that there is a remarkably increased transcription level of Rbl2, PRDM1, p21,

E2F4, lin9, CDK2 in the dox-induced GASC1 expression K562 cells than no dox cells (Fig. 6i).

p53 mRNA level was reduced markedly in the dox-induced GASC1 expression K562 cells. As

known, depletion of p53 in cells causes G1 arrest and apoptosis. Taken together, modulation of

GASC1 expression in leukemia cells leads to cell cycle arrest in the G0/G1 phase and shows

misshapen nuclei. Through regulating the activity of chromatin-modifying enzymes, that lead to

nuclear rupture and blebs, we sought to address a novel strategy to regulate cancer cell growth.

Discussion

The objective of this study was to investigate the manner in which both cellular and nuclear

size change during differentiation and how such changes might relevant to chromatin-modifying

enzymes. However, to date, there have not been many investigations of chromatin contribution to

nuclear morphology, although altered histone modifications and chromatin decompaction are

consistently present in both nuclei with abnormal morphology and nuclear blebs.

Lamins are fibrous proteins as well as nuclear structural proteins. Mouse ES cells express high

levels of A-type lamin during cell differentiation [36,37]. Forced expression of the progerin mutant

isoform of lamin A/C (LMAC) increases osteogenic differentiation and reduces adipogenic

differentiation [37]. Additionally, deregulation of lamin expression can lead to various cancer and be

used as diagnostic biomarkers. Lamin is a genetic reason for nuclear misshapen, however

epigenetic mechanism of nuclear shape change in cell fate determination is still obscure. Here, we

aim to understand whether activity alterations of chromatin-modifying enzymes affect nuclear

contour, and what mechanisms contribute to cellular identification of cell fate. We employed

histone-modifying enzyme inhibitors to get some clues of whether certain types of enzymes or all

histone-modifying enzymes are able to effect nuclear size alteration. We used stem cells such as

hESCs and NPCs to investigate the impact of histone-modifying enzyme inhibitors at the stage of

pluripotency and multipotency maintenance, which may have some distinctions. Also, we

investigated the impact of histone-modifying enzyme inhibitors on cancer cells such as leukemia

cells. We observed that there is enlarged nuclear size and misshapen nucleus via chromatin-

modifying enzyme inhibition. As known, heterochromatin appears larger, fewer, and decondensed

in ES cells, numerous and condensed in ES cell-derived neuronal progenitor cells, and appear

fewer but more compact in terminally differentiated neurons [38,39]. Additionally, cancer cells have

global alterations in chromatin organization, mostly a reduction in heterochromatin [40]. Therefore,

it suggested that the original histone states identify the specific type of cell, and impact the outcome

of histone-modifying enzymes inhibitors treatment.

In addition, when treated with histone-modifying enzyme inhibitors, the nuclear size was able

to alter, and the nuclear shape becomes aberrant. It demonstrated that chromatin state is a major

factor determining the morphology of the nucleus. The increase of nuclear blebbing in nucleus

which indicated that a consequence of increasing euchromatin or decreasing heterochromatin, and

decreasing nuclear rigidity. In addition, during dysregulation of histone by treatment of histone-

modifying enzyme inhibitors, some cells were unable to endure the stimulation and die, but some

cells with a misshapen nucleus survived. Is that the reconstructive chromatin provides certain

benefits for cell survival during chromatin-modifying enzyme dysregulation? After treatment of

histone-modifying enzyme inhibitors in NPCs, it was shown that the decreased expression of genes

who play a role in stem cell identification. We also observed the nuclear size become smaller and

smaller when cells were differentiated from hESCs to NPCs, then to iN cells. Simultaneously,

chromatin appears decondensed in ES cells, then appears more compact in differentiated neurons.

Collectively, it implies that the major function of chromatin-modifying enzyme is to determine cell

fate, nuclear size alteration is a phenomenon accompanied by the occurrence of cell lineage

commitment in normal cells.

As known, histone modifications correlate with ESC differentiation. Hundreds of promoters

of pluripotent genes gain their epigenetic marks, whereas many others lose it, during the transition

from ESC to NPCs, and from NPCs to differentiated neurons [41,42]. Through modulating chromatin-

modifying enzymes that are able to control a huge number of epigenetic marks on the promoters

of stem cell genes. Therefore, we ask for is that possible to modulate chromatin-modifying

enzymes to alter cancer cell fate. We selected three representative chromatin-modifying enzymes.

Unlike KDM4A-B, KDM4C (GASC1) is associated with chromatin during mitosis, and triggers

the demethylation of H3K9me3 mark [14]. Ectopic expression of GASC1 or other JMJD2 members

markedly decreases H3K9me3/me2 levels, increases H3K9me1 levels, delocalizes HP1

(heterochromatin protein 1), and reduces heterochromatin in vivo [19]. GASC1 is an Oct4 target in

mouse ES cells, and acts as a positive regulator of Nanog, also is preferentially expressed in

undifferentiated ES cells [12,43]. Depletion of GASC1 in ES cells induced differentiation with a

global increase in H3K9me3, confirming that the histone demethylase activity of GASC1 is linked

to the maintenance of pluripotency. The expression levels of GASC1 were also significantly related

to the grade of basal-like breast cancers which is the most aggressive type [18]. It suggested that

histone-modifying enzymes impact the cell fate of normal cells as well as cancer cells. To

investigate the impact of GASC1 in cancer cells, we manipulated GASC1 expression in a

commonly used leukemia cell line K562. We found that overexpression GASC1 impairs the cell

cycle process, deprives K562 cells proliferation capacity, and shows nuclear abnormalities. It

indicated that chromatin-modifying enzymes are associated with nuclear size determination and

cell fate decision of normal cells as well as cancer cells. Through modulating chromatin-modifying

enzymes activity which enables to deprive the aggressive proliferation capacity of cancer cells,

arrest cells in the cell cycle to die, and as a consequence, show nucleus misshapen.

Methods

Cell culture and media.

All cell types were maintained at 37°C in a humidified 5% (vol/vol) CO2 environment during

culture. H1 ES cells were obtained from WiCell Research Resources (WiCell, WI). ES cells were

maintained as feeder-free cells in mTeSR™1 medium (Stem Cell Technologies). Cells were

allowed to spread overnight prior to transduction. We used a doxycycline (dox)-inducible lentiviral

vector system to transduce empty vector plasmid as a control and GASC1, DNMT3A, KDM4A

into the H1 ES cells, separately. Selection was carried out for 7 days in mTeSR™1 medium with

50 μg/ml hygromycin.

Generation of iN cells from human ES cells

H1 human ESCs were dissociated using accutase and plated as single cells in mTeSR™1 medium

supplemented with ROCK inhibitor Y-27632 (10 μM, StemCell Technologies, Inc.) on matrigel to

a final density of 20,000-30,000 cells per cm^2. On the second day, attached cells were infected with

lentivirus for Ngn2-puro and rtTA expression. After 24 hr (day 0), transgene expression was

induced by replacing the medium with N2/B27 medium (DMEM/F12, N2 supplement (Life

Technologies), B27 supplement (Life Technologies), 10 μg/ml insulin (Life Technologies)), and

2 μg/ml Doxycycline. Selection was carried out for 14 days (day 1 to day 14) in N2/B27 medium

with 2 μg/ml puromycin. From day 2, induced neuronal cells were maintained in N2/B27 medium

and changed medium every 2 days.

hESCs differentiate to neural progenitor cells

Growing hESCs in mTeSR media on matrigel coated dish until confluent, then enzymatically

passage with accutase, resuspend in Neurobasal-A medium, DMEM/F12, 1% N2, 1% B27 and

insulin media, with Y-27632, a specific inhibitor of Rho kinase (ROCK) activity, cultured in the

same size of the dish without matrigel coated. Overnight, colonies will form floating spherical

clusters termed "embryoid bodies" (EBs). Wait until the EB size grows to 200～400 μm, gently

move it to matrigel coated dish. The next day, it will attach to the bottom. Changing medium with

N2/B27 media supplemented with dual SMAD inhibitors (0.1 μM LDN193189 and 10 μM

SB431542). Feed EBs every second day with N2/B27 media supplemented with 0.1 μM

LDN193189 and 10 μM SB431542. After several days, the neural rosettes begin to appear,

characterized as round clusters of neuroepithelial cells with apicobasal polarity. Staining Sox1 to

indicate the neural progenitor cells.

Virus generation

Lentiviruses were produced as described [44] in HEK293T cells (ATCC, VA) by co-transfection with

three helper plasmids (pRSV-REV, pMDLg/pRRE and vesicular stomatitis virus G protein

expression vector) (12 μg of lentiviral vector DNA and 6 μg of each of the helper plasmid DNA

per 75 cm^2 culture area) using calcium phosphate [8]. Lentiviruses were harvested with the medium

24 and 46 hr after transfection. Viral supernatant was filtered through a 0.45 μm filter, pelleted by

centrifugation (21,000 × g for 2 hours), resuspended in DMEM, aliquoted, and snap-frozen in

liquid N_2. Only virus preparations with >90% infection efficiency as assessed by EGFP expression

or puromycin resistance were used for experiments.

Lentiviral infections

Complementary DNAs for candidate genes were cloned into doxycycline-inducible lentiviral

vectors, as described previously [45]. Lentiviral production and cell infection was performed as

described previously [46]. Cells were infected with concentrated lentivirus and treated with

doxycycline (2 mg/ul) 16-24 h later.

Immunofluorescence

Cultured iN cells were fixed in 4% paraformaldehyde in PBS for 15 min at room temperature,

washed three times with PBS, and incubated in 0.2% Triton X-100 in PBS for 10 min at room

temperature. Cells were blocked in PBS containing 10% BSA for 3 hours at 4°C. Primary

antibodies were applied overnight. Cells were washed in PBS three times. Secondary antibodies

were applied for 3 hours, washed. 300 nM DAPI staining for indication of nucleus. Mount.

Antibody

The following primary antibodies with indicated dilution in blocking buffer were used: Rabbit

anti-Tuj1 (Covance, BioLegend, MRB-435P, 1:1,000), Mouse anti-Tuj1 (Covance, BioLegend,

MMS-435P, 1:1,000), Rabbit anti-human Sox1 (Millipore, AB15766), Goat anti-Sox2 (Santa Cruz,

1:200). After staining with corresponding secondary antibodies in PBS plus 1% BSA, coverslips

were washed four times with PBS, each for 10 min, mounted with the mounting medium onto glass

slides, and examined under Olympus FluoView FV1000.

Inhibitors of chromatin-modifying enzymes

Purchase from tocris. GSK J1 (Cat. No. 4593), (R)-2-Hydroxyglutaric acid disodium salt (R2-HG,

Cat. No. 6124), GSK-LSD 1 dihydrochloride (Cat. No. 5361), IOX 1 (Cat. No. 4464/10), JIB04

(Cat. No. 4972/10), NSC636819 (Cat. No. 5287/10), PBIT (Cat. No. 5930), C646 (Cat. No.

4200/10), 5-azacytidine (5-Aza, Cat. No. 3842).

qRT-PCR

RNA was isolated through standard methods following manufacturer's instructions (QIAGEN),

and reverse transcribed with Superscript III (Life Technology). qRT-PCR was performed with Fast

Start universal SYBR green master mix (Roche) along with gene specific primers on a 7900HT

real-time PCR machine (Applied Biosystems). Statistical significance was determined by Students

t test based on triplicated experiments.

Flow cytometry

Flow cytometry was used to sort live cells by their size and cell cycle phase. The cells were

harvested by trypsinization, then stained with 20 μM Hoechst 33342 which is a DNA dye in PBS

for 15 minutes at 37°C and then sorted on a BD FACSAria flow cytometer. DNA content was used

to determine cell cycle phase, and the forward scatter area parameter was used as a readout for cell

size. Flow cytometry data was prepared for presentation using FlowJo2.

Coulter counter

Coulter counter is an apparatus for counting and sizing particles suspended in electrolytes. We use

the Z2 Coulter counter (Beckman). After collect the cells by trypsinization and spin down on

centrifuge, then resuspend in PBS, it is ready to run them through the counter (no staining needed).

Turn on the coulter counter and setup calibration as user's manual. Add your cell samples in

accuvette and gently mix then enter the accuvette on counting platform. Then, setting up the

aperture diameter, current Kd, and lines for selecting units and set the upper and lower size

expected particle sizes. Run the cell samples. Export the data for analysis later. The size

distribution of the sorted cells was determined using the Z2 Coulter Counter and Multisizer III

software (Beckman Coulter).

Statistical Tests

Microsoft Excel was used for data analysis. Experiments were performed in a blinded fashion; i.e.,

the experimenter was unaware of the genotype of the samples being studied. Cumulative plots

were generated from parameters collected from individual cells under similar experimental

conditions. Average data are presented as bar graphs indicating means ± SD. Statistical

comparisons between bar graphs were made using the unpaired, one-tail, Student t test (*$P < 0.05$,

**$P < 0.01$, and ***$P < 0.001$, all versus control).

Acknowledgements

We would like to thank all our laboratory members and collaborators for their comments and dedicated work that have greatly contributed to the ideas presented here. L.M. was supported by the California Institute of Regenerative Medicine (CIRM). This study was supported by the Howard Hughes Medical Institute Faculty Scholar Award (M.W.), and the NIH R01 grant.

Author Contributions

L.M., E.Z. and M.W. conceived and designed the study; L.M., E.Z., planned and performed the experiments; L.M., E.Z., performed FACS and Coulter counter; J.S. and M.W. provided critical advice on experimental designs and data interpretations; K.S. and G.N. provided experimental material; K.S. revised the manuscript and thank for helpful comments on the manuscript. L.M. wrote the manuscript.

Declaration of Interests

The authors declare no conflicts of interest in the presented study.

FIGURE LEGENDS

Fig. 1 Nuclear size is associated with cell fate decision and chromatin-modifying enzyme

activity.

a, Immunofluorescence staining cell nucleus of hESCs, hESC-derived NPCs, hESC-derived iN

cells by DAPI (blue).

b, Measurements of cell nucleus size of hESCs, hESC-derived NPCs, hESC-derived iN cells by

DAPI staining. ***$p < 0.001$. White scale bars= 50 μm.

c, Brightfield and fluorescent images of hESCs and hESC-derived NPCs immunofluorescence

staining for H3K9me3 (red) and DAPI (blue).

d, Measurements of nuclear size after hESCs treated with histone demethylase inhibitors (GSK J1,

(R)-2-Hydroxyglutaric acid disodium salt, GSK-LSD1, IOX1, JIB04, NSC636819, PBIT), DNA

methyltransferase inhibitor (5-Azacytidine), and histone acetyltransferase inhibitor (C646) by

DAPI staining. Statistically significant differences are marked with asterisks. *$P < 0.05$, **$p <$

0.01, ***$p < 0.001$.

e, Measurements of nuclear size by DAPI staining after hESC-derived NPCs treated with

chromatin-modifying enzymes inhibitors.

f, Measurements of nuclear size by DAPI staining after hESC-derived iN cells treated with DMSO

as control and 5-Aza.

g, Representative immunocytochemistry images of hESCs stained with DAPI. hESCs treated with

DMSO as control and 5-Aza. It was shown that the bigger nuclei occurred at 5-Aza treatment. The

images were taken at 40× on a confocal microscope. Red scale bars= 20 μm.

Fig. 2 Inhibition of chromatin-modifying enzyme enlarges hESCs nuclear size.

a, b, c, Using flow cytometry assay to measure cell size and nuclear size after chromatin-modifying

enzymes inhibitors treatment of hESCs for 7 days. FSC-A indicates cell size. Hoechst-W (width)

indicates nuclear size.

d, e, After inhibitors treatment, measuring cell size and nuclear size by flow cytometry. FSC-A

indicates cell size. Hoechst-W (width) indicates nuclear size. Each group normalizes to DMSO

control. *$P < 0.05$.

f, The cell volume of hESCs was measured with coulter counter after inhibitors treatment for 6

days. The curve was calculated from the coulter counter measurements of asynchronous size

distribution. FL (Femtoliter). *$P < 0.05$.

Fig. 3 Inhibition of chromatin-modifying enzyme effects nuclear size and cell identification

of NPCs.

a, Fluorescent images of hESC-derived NPCs immunofluorescence staining for Sox1 (green), PI

(red), and DAPI (blue) after treatment with inhibitors of chromatin-modifying enzymes. White

scale bars= 50 μm.

b, Histogram showing the cell size of hESC-derived NPCs measured by coulter counter after

inhibitors treatment for 9 days compared with DMSO control. Two-tailed t-test, Mean ± SD. *$P <$

0.05.

Fig. 4 Inhibition of chromatin-modifying enzyme effects leukemia cell nuclear architecture.

a, b, c, Using flow cytometry to measure cell size and nuclear size of K562 leukemia cells after

inhibitors treatment for 5 days. FSC-A indicates cell size. Hoechst-W (width) indicates nuclear

size.

d, Representative images of DAPI staining for K562 cells treated with DMSO as control and 5-

Aza, IOX1, GSK-LSD1, and JIB04. It was shown that the bigger size and twisted shape of nuclei

in inhibitors treatment. The images were taken at 40× on a confocal microscope. White scale bars=

25 μm.

e, Measurements of the nuclear size using DAPI staining and fluorescence microscopy after K562

cells treated with inhibitors of chromatin-modifying enzymes. Two-tailed t-test, Mean ± SD. **p

< 0.01, ***p < 0.001.

f, Measuring cell size and nuclear size of K562 cells through flow cytometry, DAPI staining, and

coulter counter after inhibitors treatment for 7 days. Using flow cytometry to measure cell size and

nuclear size. Measuring nuclear size after DAPI staining by confocal microscope. Coulter counter

measures cell size. The average values for each treatment were normalized to DMSO control. ***p

< 0.001.

Fig. 5 Upregulated GASC1 impairs differentiation from hESCs to iN cells.

a, Brightfield and fluorescent images of the hESC-derived iN cells with wt, dox-induced

overexpression of GASC1, DNMT3A, and KDM4A stained for the pluripotent marker Sox2

(green), the neural cells marker Tuj1 (red), and nucleus staining DAPI (blue) after 14 days of iN

cells differentiation.

b, Neurite length of the hESC-derived iN cells was measured at 14 days of iN cells differentiation.

p-values were calculated by unpaired t-test, error bars represent standard deviation, * $p < 0.05$, ***

$p < 0.001$.

Fig. 6 GASC1 promotes aberrant nuclear structure and cell cycle arrest of leukemia cells.

a, Brightfield and fluorescent images show that K562 leukemia cells differentiate to iN cells with

Ngn2 overexpression after 14 days differentiation and stain with the neuronal marker Tuj1 (red).

The yellow arrows indicate the cells of morphological changes.

b, c, Brightfield images of the K562 leukemia cells with induced expression of GASC1 with or

without doxycycline (dox). The no dox-induced GASC1 overexpression K562 cells show spherical

morphology, same as normal cells. The dox-induced GASC1 overexpression K562 cells show

morphology change from spherical to twisted shape and nuclear blebs. The red arrow indicates the

cells with twisted shapes. White scale bars= 25 μm.

d, e, Fluorescent images by DAPI staining demonstrate that the dox-induced GASC1

overexpression K562 cells show the twisted nuclear shape and bigger nuclei compared with the no

dox-induced GASC1 overexpression K562 cells. The red arrow indicates the enlarged and twisted

nucleus.

f, g, Flow cytometric analysis of K562 cells were double-stained with BrdU and PI. The dox-

induced GASC1 overexpression K562 cells proliferate significantly slower than the no dox-

induced GASC1 overexpression K562 cells by Brdu assay.

h, Cell number counting after induction by dox for 7 days. The cell number is decreased in dox-

induced GASC1 overexpression K562 cells than no dox-induced GASC1 overexpression K562

cells. ***$p < 0.001$. n=3.

i, The genes play a role in cell-cycle arrest detected by qPCR in the dox-induced GASC1

overexpression K562 cells and no dox-induced GASC1 overexpression K562 cells. Unpaired two-

tailed Student t-test. Data are mean ± SD. $p < 0.001$.

REFERENCES

Tamashunas, A. C. *et al.* High-throughput gene screen reveals modulators of nuclear shape. *Mol Biol Cell* **31**, 1392-1402 (2020).

McBeath, R., Pirone, D. M., Nelson, C. M., Bhadriraju, K. & Chen, C. S. Cell shape, cytoskeletal tension, and RhoA regulate stem cell lineage commitment. *Dev Cell* **6**, 483-495 (2004).

Bartova, E., Krejci, J., Harnicarova, A. & Kozubek, S. Differentiation of human embryonic stem cells induces condensation of chromosome territories and formation of heterochromatin protein 1 foci. *Differentiation* **76**, 24-32 (2008).

Pajerowski, J. D., Dahl, K. N., Zhong, F. L., Sammak, P. J. & Discher, D. E. Physical plasticity of the nucleus in stem cell differentiation. *Proc Natl Acad Sci U S A* **104**, 15619-15624 (2007).

Tzur, A., Kafri, R., LeBleu, V. S., Lahav, G. & Kirschner, M. W. Cell growth and size homeostasis in proliferating animal cells. *Science* **325**, 167-171 (2009).

Tzur, A., Moore, J. K., Jorgensen, P., Shapiro, H. M. & Kirschner, M. W. Optimizing optical flow cytometry for cell volume-based sorting and analysis. *PLoS One* **6**, e16053 (2011).

Gregg, E. C. & Steidley, K. D. Electrical counting and sizing of mammalian cells in suspension. *Biophys J* **5**, 393-405 (1965).

Pang, Z. P., Cao, P., Xu, W. & Sudhof, T. C. Calmodulin controls synaptic strength via presynaptic activation of calmodulin kinase II. *J Neurosci* **30**, 4132-4142 (2010).

9   Shen, H., Xu, W. & Lan, F. Histone lysine demethylases in mammalian embryonic

development. *Exp Mol Med* **49**, e325 (2017).

10   He, J. *et al.* Kdm2b maintains murine embryonic stem cell status by recruiting PRC1

complex to CpG islands of developmental genes. *Nat Cell Biol* **15**, 373-384 (2013).

11   Pereira, J. D. *et al.* Ezh2, the histone methyltransferase of PRC2, regulates the balance

between self-renewal and differentiation in the cerebral cortex. *Proc Natl Acad Sci U S A*

107, 15957-15962 (2010).

12   Loh, Y. H., Zhang, W., Chen, X., George, J. & Ng, H. H. Jmjd1a and Jmjd2c histone H3

Lys 9 demethylases regulate self-renewal in embryonic stem cells. *Genes Dev* **21**, 2545-

2557 (2007).

13   Kottakis, F. *et al.* NDY1/KDM2B functions as a master regulator of polycomb complexes

and controls self-renewal of breast cancer stem cells. *Cancer Res* **74**, 3935-3946 (2014).

14   Kupershmit, I., Khoury-Haddad, H., Awwad, S. W., Guttmann-Raviv, N. & Ayoub, N.

KDM4C (GASC1) lysine demethylase is associated with mitotic chromatin and regulates

chromosome segregation during mitosis. *Nucleic Acids Res* **42**, 6168-6182 (2014).

15   Yang, Z. Q. *et al.* Identification of a novel gene, GASC1, within an amplicon at 9p23-24

frequently detected in esophageal cancer cell lines. *Cancer Res* **60**, 4735-4739 (2000).

16   Northcott, P. A. *et al.* Multiple recurrent genetic events converge on control of histone

lysine methylation in medulloblastoma. *Nat Genet* **41**, 465-472 (2009).

17   Xiao, Z. *et al.* GASC1 promotes glioma progression by enhancing NOTCH1 signaling.

Mol Med Rep **23** (2021).

18   Liu, G. *et al.* Genomic amplification and oncogenic properties of the GASC1 histone

demethylase gene in breast cancer. *Oncogene* **28**, 4491-4500 (2009).

19   Cloos, P. A. *et al.* The putative oncogene GASC1 demethylates tri- and dimethylated lysine

9 on histone H3. *Nature* **442**, 307-311 (2006).

20   Wu, L. *et al.* Histone Demethylases KDM4A and KDM4C Regulate Differentiation of

Embryonic Stem Cells to Endothelial Cells. *Stem Cell Reports* **5**, 10-21 (2015).

21   Richly, H., Aloia, L. & Di Croce, L. Roles of the Polycomb group proteins in stem cells

and cancer. *Cell Death Dis* **2**, e204 (2011).

22   Heo, S. J. *et al.* Differentiation alters stem cell nuclear architecture, mechanics, and

mechano-sensitivity. *Elife* **5** (2016).

23   Chambers, S. M. *et al.* Highly efficient neural conversion of human ES and iPS cells by

dual inhibition of SMAD signaling. *Nat Biotechnol* **27**, 275-280 (2009).

24   Zhang, Y. *et al.* Rapid single-step induction of functional neurons from human pluripotent

stem cells. *Neuron* **78**, 785-798 (2013).

25   Volpe, T. A. *et al.* Regulation of heterochromatic silencing and histone H3 lysine-9

methylation by RNAi. *Science* **297**, 1833-1837 (2002).

26   Lee, K. K. & Workman, J. L. Histone acetyltransferase complexes: one size doesn't fit all.

Nat Rev Mol Cell Biol **8**, 284-295 (2007).

27   Stephens, A. D. *et al.* Chromatin histone modifications and rigidity affect nuclear

morphology independent of lamins. *Mol Biol Cell* **29**, 220-233 (2018).

Fischer, E. G. Nuclear Morphology and the Biology of Cancer Cells. *Acta Cytol* **64**, 511-519 (2020).

Papanicolaou, G. N. & Traut, H. F. The diagnostic value of vaginal smears in carcinoma of the uterus. 1941. *Arch Pathol Lab Med* **121**, 211-224 (1997).

Sullivan, T. *et al.* Loss of A-type lamin expression compromises nuclear envelope integrity leading to muscular dystrophy. *J Cell Biol* **147**, 913-920 (1999).

Dang, L. *et al.* Cancer-associated IDH1 mutations produce 2-hydroxyglutarate. *Nature* **462**, 739-744 (2009).

Lu, C. *et al.* IDH mutation impairs histone demethylation and results in a block to cell differentiation. *Nature* **483**, 474-478 (2012).

Koivomagi, M., Swaffer, M. P., Turner, J. J., Marinov, G. & Skotheim, J. M. G1 cyclin-Cdk promotes cell cycle entry through localized phosphorylation of RNA polymerase II. *Science* **374**, 347-351 (2021).

Coffman, J. A. Cell cycle development. *Dev Cell* **6**, 321-327 (2004).

Litovchick, L. *et al.* Evolutionarily conserved multisubunit RBL2/p130 and E2F4 protein complex represses human cell cycle-dependent genes in quiescence. *Mol Cell* **26**, 539-551 (2007).

Galiova, G., Bartova, E., Raska, I., Krejci, J. & Kozubek, S. Chromatin changes induced by lamin A/C deficiency and the histone deacetylase inhibitor trichostatin A. *Eur J Cell*

Biol **87**, 291-303 (2008).

37   Scaffidi, P. & Misteli, T. Lamin A-dependent misregulation of adult stem cells associated

with accelerated ageing. *Nat Cell Biol* **10**, 452-459 (2008).

38   Meshorer, E. *et al.* Hyperdynamic plasticity of chromatin proteins in pluripotent embryonic

stem cells. *Dev Cell* **10**, 105-116 (2006).

39   Meshorer, E. & Misteli, T. Chromatin in pluripotent embryonic stem cells and

differentiation. *Nat Rev Mol Cell Biol* **7**, 540-546 (2006).

40   Gurrion, C., Uriostegui, M. & Zurita, M. Heterochromatin Reduction Correlates with the

Increase of the KDM4B and KDM6A Demethylases and the Expression of Pericentromeric

DNA during the Acquisition of a Transformed Phenotype. *J Cancer* **8**, 2866-2875 (2017).

41   Bernstein, B. E. *et al.* A bivalent chromatin structure marks key developmental genes in

embryonic stem cells. *Cell* **125**, 315-326 (2006).

42   Mikkelsen, T. S. *et al.* Genome-wide maps of chromatin state in pluripotent and lineage-

committed cells. *Nature* **448**, 553-560 (2007).

43   Katoh, Y. & Katoh, M. Comparative integromics on JMJD2A, JMJD2B and JMJD2C:

preferential expression of JMJD2C in undifferentiated ES cells. *Int J Mol Med* **20**, 269-273

(2007).

44   Brambrink, T. *et al.* Sequential expression of pluripotency markers during direct

reprogramming of mouse somatic cells. *Cell Stem Cell* **2**, 151-159 (2008).

45   Stadtfeld, M., Maherali, N., Breault, D. T. & Hochedlinger, K. Defining molecular

cornerstones during fibroblast to iPS cell reprogramming in mouse. *Cell Stem Cell* **2**, 230-

240 (2008).

46   Chen, C. & Okayama, H. High-efficiency transformation of mammalian cells by plasmid

DNA. *Mol Cell Biol* **7**, 2745-2752 (1987).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

a

b

Neurite length

Figure 6

a Untreated   Ngn2 induction

25 um   50 um   25 um   Tuj1

K562-GASC1 cells

b no dox   **c** dox

25 um

d no dox DAPI   **e** dox DAPI

25 um

f no dox   **g** dox

BrdU 38.1   BrdU 4.54

h

i GASC1 causes cell-cycle arrest in the G0/G1 phase.

no dox   dox