Chromatin-modifying enzymes as modulators of nuclear size during lineage differentiation

Lingjun Meng[1, 2, *]

[1]Department of Pathology,

[2]Institute for Stem Cell Biology and Regenerative Medicine,

Stanford University School of Medicine, Stanford, CA 94305, USA

*Correspondence: lingjun@stanford.edu, menglingjundl@gmail.com.

Key word: histone demethylases, nuclear size, differentiation, cancer.

Abstract

The mechanism of nuclear size determination and alteration during normal lineage

development and cancer pathologies which is not fully understood. As recently reported, chromatin

modification can change nuclear morphology. Therefore, we screened a range of pharmacological

chemical compounds that impact the activity of chromatin-modifying enzymes, in order to get a

clue of the specific types of chromatin-modifying enzymes that remarkably effect nuclear size and

shape. We found that interrupted activity of chromatin-modifying enzymes is associated with

nuclear shape abnormalities. Furthermore, the activity of chromatin-modifying enzymes perturbs

cell fate determination in cellular maintenance and lineage commitment. Our results indicated that

chromatin-modifying enzyme regulates cell fate decision during lineage differentiation and is

associate with nuclear size alteration.

Significance

Here we described for the first time the modulation of chromatin-modifying enzymes

remarkably effects nuclear shape, and perturbs cell fate determination in cellular maintenance and

lineage fate commitment in normal stem cells and leukemia cancer cells. We found that the

irregularities of nuclear contour were highly related to pharmacological inhibition of chromatin-

modifying enzyme activity. After manipulating a histone demethylase named GASC1, we found

that upregulation of GASC1 impairs the differentiation of hESCs to terminally differentiated

neural cells. Moreover, upregulation of GASC1 impairs the proliferation of leukemia cells due to

cell cycle arrest in the G0/G1 phase and indicates misshapen nuclei. This study suggested that

chromatin-modifying enzyme regulates the nuclear contour related cell fate decision also including

cancer cell fate determination.

Introduction

As reported, nuclear volume is an important morphological parameter that can change during

cancer progression (1). After confocal microscopy was discovered and broadly used to observe the

detailed structure of specific objects within the cell, the most fascinated cellular organelle - nucleus

was researched more carefully. Recent studies have also highlighted that nuclear shape can

influence fate decisions (2). Embryonic stem cells (ESCs) differentiation is accompanied by an

increase in chromatin condensation (3) leading to an increase in nuclear stiffness (4) and a decrease

in nuclear size. As known, the nuclear size of stem cells is extremely bigger than terminal

differentiated cells. Cell differentiation is corresponding with chromatin marks reset, which raises

the question of whether chromatin's contribution to nuclear mechanics affects nuclear morphology

as well as cell fate. It is therefore critical to identify the contributions of various chromatin-

modifying enzymes that are relevant to cell fate decisions, and consequently display nuclear size

alteration.

Nuclear architecture is impacted via chromosome structure which is determined by histone

modifications. The cellular differentiation from stem cells to differentiated cells which is

corresponding with nuclear size reduction as well as chromosome condense. Histone demethylases

have also been reported to be critical enzymes that regulate the balance between self-renewal and

differentiation in hESCs (human embryonic stem cells) (5-8). Accordingly, we broadly utilized

chromatin-modifying enzymes inhibitors to treat stem cells such as hESCs, NPCs (neural

progenitor cells), and later cancer cells such as leukemia cells. We demonstrated that the

irregularities of nuclear contour were highly related to pharmacological inhibition of chromatin-

modifying enzyme activity. As known, upregulation or downregulation of histone demethylases is

observed in various types of cancer and is associated with higher rates of malignancy and relapse

after treatment (9). In particular, GASC1 (gene amplified in squamous cell carcinoma, also known

as KDM4C, a histone demethylase) promotes chromosomal instability and transcription initiation,

therefore, plays a causative role in cancer cells driven by its dysregulation (10). GASC1 was found

aberrant expression in esophageal squamous carcinomas (11), medulloblastomas (12), glioma (13),

breast cancer (14), and prostate cancer (15). In addition, GASC1 plays an essential role in the

transition of mouse ESCs (mESCs) to endothelial cells (16). Consistently, we demonstrated that

upregulation of GASC1 impairs the differentiation of undifferentiated hESCs to terminally

differentiated neural cells. Moreover, leukemia cells with upregulated GASC1 expression lost their

proliferation ability, show cell cycle arrest in the G0/G1 phase, and appear misshapen nucleus.

Taken together, epigenetic perturbations are likely to be the cause for misregulation of nuclear size

and of certain cancer cell fates (17).

Result

The nuclear size is associated with cell fate decision and chromatin-modifying enzyme

activity

Size and shape are distinctive aspects of nuclear structure. Nuclear size impacts cell function,

and is altered according to the specific status of cell types during cell development (18). It is

unclear which modulators effect the altered nuclear size during cell fate transformation. To

investigate the distinct nuclear size of various cell types, we employed highly efficient

differentiation methods to interrogate the size change of nucleus during lineage differentiation. To

differentiate NPCs from hESCs, we used two molecular compounds, SB431542 and LDN193189,

to inhibit the TGF-β (transforming growth factor beta) and BMP (bone morphogenetic protein)

pathways, which are also called Dual SMAD inhibitors (19). In addition, as we previously used,

overexpression of Ngn2 (neural gene neurogenin-2) could converts hESCs into iN cells (induced

neuronal cells) with nearly 100% yield in less than two weeks. The hESCs-derived iN cells grow

long axons that conduct electrical impulses away from the neuron's cell body (20). We utilized

DAPI to stain hESCs for 5 min, then examined the dynamic change of the nuclear size via

measuring the area of nucleus under a fluorescence microscope (Fig. 1*A*). Notably, it was shown

that there is a reduced size of approximately 40% from hESCs to NPCs, and 24% decrease from

NPCs to iN cells (Fig. 1*B*). Consistently as reported, the nuclear size of hESCs is bigger than NPCs,

and the nuclear size of NPCs is bigger than iN cells. In addition, it was shown that the nuclear size

of hESCs and NPCs are in a wider range than iN cells, which indicates that there is a potential of

stem cells in nuclear dynamics. The different sizes of nuclei may correspond with hESCs, NPCs,

and neural cells characterized with distinct unique chromatin architecture. During the

reprogramming of cell identity and cell fate determination, heterochromatin plays a critical role.

H3K9me3 (Tri-methylation of histone H3 lysine 9) is a histone mark associated with

heterochromatin, and has emerged as a key player in repressing lineage-inappropriate genes, and

shielding them from activation by transcription factors. We investigated the H3K9me3 expression

in hESCs and hESC-derived NPCs by immunofluorescence staining. It was shown that there are

high levels of expression of H3K9me3 in the nucleus (Fig. 1*C*). To directly address the role of

chromatin in nuclear morphology, we employed a range of commercially available chemical

compound inhibitors of the enzymes that modulate histone modification state to manipulate

chromatin alterations to treat hESCs (Fig. 1*D* and *E*). These available inhibitors of chromatin-

modifying enzymes that were represented the various functions of chromatin regulation, which

helped us to investigate broadly chromatin-modifying enzymes associated with nuclear alteration.

We purchased these inhibitors from tocris as the method shown. As reported, GSK-J1 is a highly

potent inhibitor of the H3K27 histone demethylases JMJD3 (KDM6B) and UTX (KDM6A),

likewise inhibits KDM5B, KDM5C, and KDM5A to increase H3K27me3. Histone demethylases

inhibitor increased heterochromatin levels and chromatin-based nuclear stiffness. Heterochromatin

is highly condensed DNA that leads to gene silencing (21). R2HG is oncometabolite produced by

mutant isocitrate dehydrogenase 1 (IDH1) and IDH2, inhibits KDM4A in murine embryonic

fibroblasts as well as increases H3K9me3, H3K36me3. 5-Aza is a DNA methyltransferase

inhibitor that induces demethylation. GSK-LSD1 is a potent and selective lysine specific

demethylase 1 (LSD1) inhibitor, increasing H3K4me3, H3K9me3, and H3K27me3. IOX1 is a

histone demethylase JMJD inhibitor, increasing H3K36me3, H3K9me3, and H3K27me3. JIB04

is a pan Jumonji histone demethylase inhibitor that inhibits H3K4 demethylases, decreases

H3K36me3, H3K9me3. NSC636819 is a KDM4A/KDM4B inhibitor, increasing H3K9me3. PBIT

is a JARID1 (Jumonji AT-Rich Interactive Domain 1) inhibitor that increases H3K4me3. C646 is

a inhibit histone acetyltransferase, which corresponds predominantly to decompacted euchromatin,

allowing transcription factors to bind to regulatory sites on DNA, in order to cause transcriptional

activation (22). Through the treatment with compounds function as chromatin-modifying enzyme

inhibitors, we found that 5-Aza treatment enlarges the nuclear size of hESCs remarkably (Fig. 1*D*),

also some other inhibitors. The great increase in nuclear size of hESCs from 5-Aza treatment

indicated that modulation of chromatin-modifying enzymes is enabled to change nuclear size (Fig.

1E). It suggested that perturbation of chromatin-modifying enzyme activity alters nuclear size in

normal cells.

In addition, we noticed that the treatment of high concentrations of IOX1 revealed the

cytotoxic effects on hESCs. Therefore, we tested a series of concentrations of compounds for

hESCs treatment. The high concentrations of inhibitors caused cell death of hESCs, such as 100

mM GSK-J1, 100 mM IOX1, 1 mM JIB04, 10 mM PBIT, 25 mM C646, especially, 10 μM, 1 mM,

10 mM 5-Aza induced numerous of hESCs death. The high concentrations of inhibitors have

cytotoxic effects on cells and impact cells alive. However, hESCs grow well in 60 μM, 1 mM, 10

mM GSK-J1, 100 mM R2HG, 10 mM and 100 mM GSK-LSD1, 1 mM and 10 mM IOX1, 10 μM

and 100 μM JIB04, 10 μM and 10 mM NSC636819, 100 μM and 1 mM PBIT, 250 μM and 2.5

mM C646 treatment. It indicated that compounds function as chromatin-modifying enzyme

inhibitors which effect nuclear size alteration, possibly through modulating histone dynamics. It

had been shown that utilizing histone methyltransferase inhibitors to decrease heterochromatin

results in a softer nucleus and nuclear blebbing, without perturbing lamins. Histone demethylase

inhibitors increase heterochromatin and chromatin nuclear rigidity (23). Given that measurements

in these reports demonstrate that there is evidence that nuclear size abnormalities can be caused

by alterations to chromatin regulatory proteins.

Inhibition of chromatin-modifying enzyme effects nuclear size and cell identification of NPCs

Not only hESCs, but we also found that the treatment of 5-Aza, C646, and JIB04 enhanced the

nuclear size of hESC-derived NPCs. In contrast, the treatment of R2HG, GSK-LSD1, NSC636819,

IOX1 reduced the nuclear size of hESC-derived NPCs (Fig. 2A). 5-Aza treatment slightly reduced

the nuclear size of hESC-derived iN cells (Fig. 2B). To gain further insights into whether these

inhibitors not only impact nuclear size and cell death but also effect additional features, such as

cell fate decisions, we treated the hESC-derived NPCs with DMSO and chromatin-modifying

enzyme inhibitors. The concentration we used was tested and the same concentration was used as

hESCs treatment, then imaged hESC-derived NPCs after staining for Sox1 (green), PI (red), and

DAPI (blue) (Fig. 2C). As known, Sox1 is a neural stem cell marker and is highly expressed in

NPCs. It was observed that the expression of Sox1 was significantly reduced in a large number of

NPCs in 5-Aza, JIB04, PBIT, GSK-J1, GSK-LSD1, c646, and IOX1 treatment (Fig. 2D). To

eliminate the possibility of Sox1 expression decrease caused by cell death, we performed

propidium iodide (PI) staining to NPCs after inhibitors treatment. As known, PI is not permanent

to live cells but is used to detect dead cells in a population. It was shown that Sox1 silence was not

caused by cell death due to PI staining being negative. Except IOX1, JIB04 and GSK-LSD1

increased PI staining in NPCs which suggested that it may induces cytotoxic responses of NPCs

occur as a consequence of cell death. Therefore, it suggested that the compound inhibitors of

chromatin-modifying enzymes perturb cell identification. Collectively, inhibition of chromatin-

modifying enzymes activity enabled to impact nuclear size in both hESCs and NPCs which

consequently leads cell fate to aberrant states.

Upregulated GASC1 impairs differentiation from hESCs to iN cells

As we previously discussed, the nuclear size dramatically reduced from hESCs to NPCs, and

to iN cells during lineage differentiation. Additionally, we want to validate the specificity of the

targeted chromatin-modifying enzyme instead of using chemical compounds. So, we selected three

represented histone enzymes such as GASC1, DNMT3A, and KDM4A, based on the screening of

inhibitors of chromatin-modifying enzyme activity, to investigate the impact of chromatin-

modifying enzymes in lineage differentiation. GASC1 (lysine-specific demethylase 4C, KDM4C)

is a histone demethylase as well as KDM4A (lysine-specific demethylase 4A), they both belong to

the families of lysine demethylases (KDM). DNMT3A (DNA methyltransferase 3A) catalyzes 5-

methylcytosine methylation. 5-Aza inhibits DNA methyltransferase (DNMT). We used the

doxycycline (dox)-inducible lentiviral vector system to transduce empty vector plasmid as control

and GASC1, DNMT3A, KDM4A into the hESCs H1 cell line, separately. As mentioned above,

we used the same approach to induce neuronal cells from differentiating hESCs by Ngn2

overexpression (20). It was shown that there is a significantly high efficiency of wt hESCs

differentiating to iN cells as nearly 100%. The iN cells display a mature neural morphology with

axon extension and packaged smaller cell nucleus (Fig. 3*A*). We stained hESC-iN cells with a

neuronal marker Tuj1 (tubulin beta-III) and a pluripotent marker Sox2. It demonstrated that iN

cells differentiated from wt hESCs showed Tuj1 positive expression and long axon extension, no

Sox2 indicated hESCs left, which also occurred in KDM4A overexpressed hESC-iN cells. In

contrast, hESC-iN cells with induced overexpression of GASC1 and DNMT3A that grew few

extend axons and showed big nuclei. In addition, a large number of cells were Sox2 positive during

differentiation (Fig. 3A). After measurement analysis of neurite length at 14 days of iN cell

differentiation, we found that hESC-iN cells with GASC1 and DNMT3A overexpression grew

remarkably shorter neurite compared to wt and KDM4A overexpressed iN cells (Fig. 3B).

Perturbation of chromatin-modifying enzymes activity alters the molecular and cellular identity of

stem cells. Overall, these data suggested that chromatin-modifying enzymes can drive lineage

specification as well as assist nuclear reconstruction, in order to facilitate a smoother transition

from stem cells to differentiated cells.

GASC1 promotes aberrant nuclear structure and cell cycle arrest of leukemia cells

As above discussed, chromatin-modifying enzyme activity is highly associated with nuclear

size alteration in normal stem cells during lineage differentiation. As reported, cancer cells are

known to develop larger nuclei as they become more malignant and increase the nuclear-to-

cytoplasmic ratio (24). The mechanisms that underlie the increase in nuclear size in malignant

cells remain largely unknown. Abnormal nuclear morphology has been used for cancer diagnoses

for nearly a century in tests such as the Pap smear (25). We measured the nuclear size of K562

leukemia cells by DAPI staining, in order to investigate the distinctive features of chromatin-

modifying enzyme activity in cancer cells compared to normal stem cells (Fig. 4A and B). We

tested a series of concentrations of compound inhibitors for K562 cells treatment. Consistent with

hESCs, high concentrations of inhibitors caused cell death of K562 cells. Especially, high doses

and long-time of 5-Aza and PBIT treatment induce cell death. In addition, we photographed the

DAPI stained nucleus of K562 cells treated with DMSO as control and 5-Aza, IOX1, GSK-LSD1,

and JIB04. It was shown that K562 cells treated with inhibitors have a bigger size of nucleus and

twisted shape as clover, especially in 5-Aza, IOX1, GSK-LSD1, and JIB04 treatment (Fig. 4*A*). It

demonstrated that the irregularity of nuclear contour occurred in inhibitors treatment from the

visible fluorescent images. The most studied types of nuclear morphological abnormalities are

protrusions from the normally ellipsoidal nucleus, termed "blebs." K562 cells showed a "blebs"

shape and seem to prefer to have a twisted nuclear shape but not die as quickly as hESCs and NPCs

did. It all suggested that not only nuclear size change but also nucleus misshapen are indicated the

pathological change of K562 leukemia cells induced by inhibition of chromatin-modifying enzyme

activity. Furthermore, the size of cancer cells was more similar to hESCs but bigger than NPCs.

After inhibitors treatment for 7 days (10 μM GSK-J1, 100 μM R2HG, 1 μM 5-Aza, 10 μM GSK-

LSD1, 10 μM IOX1, 100 μM JIB04, 10 μM NSC636819, 1 μM PBIT, 2.5 μM C646), compared

to DMSO control, the increased percentage of nuclear size of K562 cells is 5% ~ 44%. The

treatment of 5-Aza and IOX1 markedly lead to the enlargement of the nuclear size of K562

leukemia cells (Fig. 4*B*). Except for lamina perturbation affecting nuclear abnormalities (26), it

also suggested that enlarged and twisted nuclear size effect the survival of K562 cells which is

associated with chromatin-modifying enzyme activity.

As mentioned above, these chromatin-modifying enzymes effect lineage differentiation of

hESCs. These hESC-iN cells with GASC1 and DNMT3A aberrant expression retain their proliferation capability, fail to differentiate, and acquire resistance to cell death (Fig. 3). As previously reported, IDH1 mutated cells generate R(2)-2-hydroxyglutarate (2HG) (27), which causes GASC1 mediated hypermethylation (15, 28). IDH1 gene mutations occur frequently in AML (acute lymphoblastic leukemia). Accordingly, we investigated the role of GASC1 in K562 leukemia cells differentiation. Basically, healthy K562 cells are spherical in shape without any structural distortions, and grow in suspension (Fig. 4C). After being differentiated by Ngn2 overexpression for 14 days, it was shown that a portion of K562 cells grew spike, and Tuj1 was not abundantly expressed (Fig. 4C). K562 cells are hard to differentiate. When adding dox in cell culture for inducing GASC1 overexpression, the majority of K562 leukemia cells began to change their morphology from smooth surfaces of spherical shape to twisted cellular shape compared with no dox-induced K562 cells (Fig. 4D and E). The abnormal nuclear morphology and protrusions termed "blebs" are both associated with chromatin alterations (23). To validate the cell shape change corresponding to nucleus change, we stain K562 cells with DAPI. Under microscope examination of GASC1 overexpressed K562 cells with or without dox, we found that it is consistent with cell shape change in dox-induced overexpressed K562 cells, that showed nucleus enlarges and shapes as clover (Fig. 4F and G). To investigate whether the twisted shape consequently provides benefits of growth of the K562 cells or not, we employed the BrdU cell proliferation assay to interrogate the cell cycle of K562 cells growth. The dox-induced K562-GASC1 cells significantly proliferate slower than the no dox-induced K562-GASC1 cells (Fig. 4H

and *I*). Meanwhile, we count the cell number of K562 cells with or without dox-induced GASC1

overexpression, it reveals that there is a significant reduction in the growth of K562 cells with

GASC1 overexpression (Fig. 4*J*). To explore the mechanism of K562 cells with GASC1

expression was able to stop proliferation at the genetic level, we selected a range of key regular

genes function as cell cycle effectors, such as Rbl2, PRDM1, p21, p53, E2F4, lin9, and CDK2 (29,

30, 31). Based on the qPCR results, we found that there is a remarkably increased transcription

level of Rbl2, PRDM1, p21, E2F4, lin9, and CDK2 in the dox-induced GASC1 expression K562

cells than in no dox cells (Fig. 4*K*). p53 mRNA level was reduced markedly in the dox-induced

GASC1 expression K562 cells. As known, depletion of p53 in cells causes G1 arrest and apoptosis.

Taken together, modulation of GASC1 expression in leukemia cells leads to cell cycle arrest in the

G0/G1 phase and shows misshapen nuclei. Through regulating the activity of chromatin-

modifying enzymes, that lead to nuclear rupture and blebs, we sought to address a novel strategy

to regulate cancer cell growth.

Discussion

The objective of this study was to investigate the manner in which both cellular and nuclear

size change during differentiation and how such changes might relevant to chromatin-modifying

enzymes. However, to date, there have not been many investigations of chromatin contribution to

nuclear morphology, although altered histone modifications and chromatin decompaction are

consistently present in both nuclei with abnormal morphology and nuclear blebs.

261   Lamins are fibrous proteins as well as nuclear structural proteins. Mouse ES cells express high

262  levels of A-type lamin during cell differentiation (32, 33). Forced expression of the progerin mutant

263  isoform of lamin A/C (LMAC) increases osteogenic differentiation and reduces adipogenic

264  differentiation (33). Additionally, deregulation of lamin expression can lead to various cancer and

265  be used as diagnostic biomarkers. Lamin is a genetic reason for nuclear misshapen, however

266  epigenetic mechanism of nuclear shape change in cell fate determination is still obscure. Here, we

267  aim to understand whether activity alterations of chromatin-modifying enzymes affect nuclear

268  contour, and what mechanisms contribute to cellular identification of cell fate. We employed

269  histone-modifying enzyme inhibitors to get some clues of whether certain types of enzymes or all

270  histone-modifying enzymes are able to effect nuclear size alteration. We used stem cells such as

271  hESCs and NPCs to investigate the impact of histone-modifying enzyme inhibitors at the stage of

272  pluripotency and multipotency maintenance, which may have some distinctions. Also, we

273  investigated the impact of histone-modifying enzyme inhibitors on cancer cells such as leukemia

274  cells. We observed that there is enlarged nuclear size and misshapen nucleus via chromatin-

275  modifying enzyme inhibition. As known, heterochromatin appears larger, fewer, and decondensed

276  in ES cells, numerous and condensed in ES cell-derived neuronal progenitor cells, and appear

277  fewer but more compact in terminally differentiated neurons (34, 35). Additionally, cancer cells

278  have global alterations in chromatin organization, mostly a reduction in heterochromatin (36).

279  Therefore, it suggested that the original histone states identify the specific type of cell, and impact

280  the outcome of histone-modifying enzymes inhibitors treatment.

In addition, when treated with histone-modifying enzyme inhibitors, the nuclear size was able

to alter, and the nuclear shape becomes aberrant. It demonstrated that chromatin state is a major

factor determining the morphology of the nucleus. The increase of nuclear blebbing in nucleus

which indicated that a consequence of increasing euchromatin or decreasing heterochromatin, and

decreasing nuclear rigidity. In addition, during dysregulation of histone by treatment of histone-

modifying enzyme inhibitors, some cells were unable to endure the stimulation and die, but some

cells with a misshapen nucleus survived. Is that the reconstructive chromatin provides certain

benefits for cell survival during chromatin-modifying enzyme dysregulation? After treatment of

histone-modifying enzyme inhibitors in NPCs, it was shown that the decreased expression of genes

who play a role in stem cell identification. We also observed the nuclear size become smaller and

smaller when cells were differentiated from hESCs to NPCs, then to iN cells. Simultaneously,

chromatin appears decondensed in ES cells, then appears more compact in differentiated neurons.

Collectively, it implies that the major function of chromatin-modifying enzyme is to determine cell

fate, nuclear size alteration is a phenomenon accompanied by the occurrence of cell lineage

commitment in normal cells.

As known, histone modifications correlate with ESC differentiation. Hundreds of promoters

of pluripotent genes gain their epigenetic marks, whereas many others lose it, during the transition

from ESC to NPCs, and from NPCs to differentiated neurons (37, 38). Through modulating

chromatin-modifying enzymes that are able to control a huge number of epigenetic marks on the

promoters of stem cell genes. Therefore, we ask for is that possible to modulate chromatin-

modifying enzymes to alter cancer cell fate. We selected three representative chromatin-modifying

enzymes. Unlike KDM4A-B, KDM4C (GASC1) is associated with chromatin during mitosis, and

triggers the demethylation of H3K9me3 mark (10). Ectopic expression of GASC1 or other JMJD2

members markedly decreases H3K9me3/me2 levels, increases H3K9me1 levels, delocalizes HP1

(heterochromatin protein 1), and reduces heterochromatin in vivo (15). GASC1 is an Oct4 target

in mouse ES cells, and acts as a positive regulator of Nanog, also is preferentially expressed in

undifferentiated ES cells (8, 39). Depletion of GASC1 in ES cells induced differentiation with a

global increase in H3K9me3, confirming that the histone demethylase activity of GASC1 is linked

to the maintenance of pluripotency. The expression levels of GASC1 were also significantly related

to the grade of basal-like breast cancers which is the most aggressive type (14). It suggested that

histone-modifying enzymes impact the cell fate of normal cells as well as cancer cells. To

investigate the impact of GASC1 in cancer cells, we manipulated GASC1 expression in a

commonly used leukemia cell line K562. We found that overexpression GASC1 impairs the cell

cycle process, deprives K562 cells proliferation capacity, and shows nuclear abnormalities. It

indicated that chromatin-modifying enzymes are associated with nuclear size determination and

cell fate decision of normal cells as well as cancer cells. Through modulating chromatin-modifying

enzymes activity which enables to deprive the aggressive proliferation capacity of cancer cells,

arrest cells in the cell cycle to die, and as a consequence, show nucleus misshapen.

Methods

Cell culture and media.

All cell types were maintained at 37°C in a humidified 5% (vol/vol) CO2 environment during

culture. H1 ES cells were obtained from WiCell Research Resources (WiCell, WI). ES cells were

maintained as feeder-free cells in mTeSR™1 medium (Stem Cell Technologies). Cells were

allowed to spread overnight prior to transduction. We used a doxycycline (dox)-inducible lentiviral

vector system to transduce empty vector plasmid as a control and GASC1, DNMT3A, KDM4A

into the H1 ES cells, separately. Selection was carried out for 7 days in mTeSR™1 medium with

50 μg/ml hygromycin.

Generation of iN cells from human ES cells

H1 human ESCs were dissociated using accutase and plated as single cells in mTeSR™1 medium

supplemented with ROCK inhibitor Y-27632 (10 μM, StemCell Technologies, Inc.) on matrigel to

a final density of 20,000-30,000 cells per cm^2. On the second day, attached cells were infected with

lentivirus for Ngn2-puro and rtTA expression. After 24 hr (day 0), transgene expression was

induced by replacing the medium with N2/B27 medium (DMEM/F12, N2 supplement (Life

Technologies), B27 supplement (Life Technologies), 10 μg/ml insulin (Life Technologies)), and

2 μg/ml Doxycycline. Selection was carried out for 14 days (day 1 to day 14) in N2/B27 medium

with 2 μg/ml puromycin. From day 2, induced neuronal cells were maintained in N2/B27 medium

and changed medium every 2 days.

hESCs differentiate to neural progenitor cells

Growing hESCs in mTeSR media on matrigel coated dish until confluent, then enzymatically

passage with accutase, resuspend in Neurobasal-A medium, DMEM/F12, 1% N2, 1% B27 and

insulin media, with Y-27632, a specific inhibitor of Rho kinase activity, cultured in the same size

of the dish without matrigel coated. Overnight, colonies will form floating spherical clusters

termed "embryoid bodies" (EBs). Wait until the EB size grows to 200～400 μm, gently move it to

matrigel coated dish. The next day, it will attach to the bottom. Changing medium with N2/B27

media supplemented with dual SMAD inhibitors (LDN193189 and SB431542). Feed EBs every

second day with N2/B27 media supplemented with 0.1 μM LDN193189 and 10 μM SB431542.

After several days, the neural rosettes begin to appear, characterized as round clusters of

neuroepithelial cells with apicobasal polarity. Staining Sox1 to indicate the neural progenitor cells.

Virus generation

Lentiviruses were produced as described (40) in HEK293T cells (ATCC, VA) by co-transfection

with three helper plasmids (pRSV-REV, pMDLg/pRRE and vesicular stomatitis virus G protein

expression vector) (12 μg of lentiviral vector DNA and 6 μg of each of the helper plasmid DNA

per 75 cm^2 culture area) using calcium phosphate (41). Lentiviruses were harvested with the

medium 24 and 46 hr after transfection. Viral supernatant was filtered through a 0.45 μm filter,

pelleted by centrifugation (21,000 × g for 2 hours), resuspended in DMEM, aliquoted, and snap-

frozen in liquid N$_2$. Only virus preparations with >90% infection efficiency as assessed by EGFP

expression or puromycin resistance were used for experiments.

Lentiviral infections

Complementary DNAs for candidate genes were cloned into doxycycline-inducible lentiviral

vectors, as described previously (42). Lentiviral production and cell infection were performed as

described previously (43). Cells were infected with concentrated lentivirus and treated with

doxycycline (2 mg/ul) 16-24 h later.

Immunofluorescence

Cultured iN cells were fixed in 4% paraformaldehyde in PBS for 15 min at room temperature,

washed three times with PBS, and incubated in 0.2% Triton X-100 in PBS for 10 min at room

temperature. Cells were blocked in PBS containing 10% BSA for 3 hours at 4℃. Primary

antibodies were applied overnight. Cells were washed in PBS three times. Secondary antibodies

were applied for 3 hours, washed. 300 nM DAPI staining for indication of nucleus. Mount.

Antibody

The following primary antibodies with indicated dilution in blocking buffer were used: Rabbit

anti-Tuj1 (Covance, BioLegend, MRB-435P, 1:1,000), Mouse anti-Tuj1 (Covance, BioLegend,

MMS-435P, 1:1,000), Rabbit anti-human Sox1 (Millipore, AB15766, 1:200), Goat anti-Sox2

(Santa Cruz, 1:200). After staining with corresponding secondary antibodies in PBS plus 1% BSA,

coverslips were washed four times with PBS, each for 10 min, mounted with the mounting medium

onto glass slides, and examined under Olympus FluoView FV1000.

Inhibitors of chromatin-modifying enzymes

Purchase from tocris. GSK J1 (Cat. No. 4593), (R)-2-Hydroxyglutaric acid disodium salt (R2HG,

Cat. No. 6124), GSK-LSD 1 dihydrochloride (Cat. No. 5361), IOX 1 (Cat. No. 4464/10), JIB04

(Cat. No. 4972/10), NSC636819 (Cat. No. 5287/10), PBIT (Cat. No. 5930), C646 (Cat. No.

4200/10), 5-azacytidine (5-Aza, Cat. No. 3842).

qRT-PCR

RNA was isolated through standard methods following manufacturer's instructions (QIAGEN),

and reverse transcribed with Superscript III (Life Technology). qRT-PCR was performed with Fast

Start universal SYBR green master mix (Roche) along with gene specific primers on a 7900HT

real-time PCR machine (Applied Biosystems). Statistical significance was determined by Students

t test based on triplicated experiments.

Statistical Tests

Microsoft Excel was used for data analysis. Cumulative plots were generated from parameters

collected from individual cells under similar experimental conditions. Average data are presented

as bar graphs indicating means $\pm$ SD. Statistical comparisons between bar graphs were made using

the unpaired, two-tail, Student t test (*$P < 0.05$, **$P < 0.01$, and ***$P < 0.001$, all versus control).

Acknowledgements

I would like to thank all our laboratory members and collaborators for their comments and

dedicated work that have greatly contributed to the ideas presented here.

Author Contributions

L.M. conceived and designed the study; L.M. planned and performed the experiments; L.M. wrote

the manuscript.

Declaration of Interests

The authors declare no conflicts of interest in the presented study.

FIGURE LEGENDS

Figure 1

Fig. 1. Nuclear size is associated with cell fate decision and chromatin-modifying enzyme

activity.

(*A*) Immunofluorescence staining cell nucleus of hESCs, hESC-derived NPCs, hESC-derived iN

cells by DAPI (blue). White scale bars= 50 μm.

(*B*) Measurements of cell nucleus size of hESCs, hESC-derived NPCs, hESC-derived iN cells by

DAPI staining. ***$P < 0.001$.

(*C*) Brightfield and fluorescent images of hESCs and hESC-derived NPCs immunofluorescence

staining for H3K9me3 (red) and DAPI (blue).

(*D*) Measurements of nuclear size after hESCs treated with histone demethylase inhibitors (GSK

J1, (R)-2-Hydroxyglutaric acid disodium salt, GSK-LSD1, NSC636819, PBIT, IOX1, JIB04),

DNA methyltransferase inhibitor (5-Azacytidine), and histone acetyltransferase inhibitor (C646)

by DAPI staining. Statistically significant differences are marked with asterisks. *P < 0.05, ***P

< 0.001.

(*E*) Representative immunocytochemistry images of hESCs stained with DAPI. hESCs treated

with DMSO as control and 5-Aza. It was shown that the bigger nuclei occurred at 5-Aza treatment.

The images were taken at 40× on a confocal microscope. White scale bars= 20 µm.

Figure 2

Fig. 2. Inhibition of chromatin-modifying enzyme effects nuclear size and cell identification of NPCs.

(*A*) Measurements of nuclear size by DAPI staining after hESC-derived NPCs treated with DMSO as control and chromatin-modifying enzymes inhibitors.

(*B*) Measurements of nuclear size by DAPI staining after hESC-derived iN cells treated with DMSO as control and 5-Aza.

428   **(*C*)** Fluorescent images of hESC-derived NPCs immunofluorescence staining for Sox1 (green), PI

429   (red), and DAPI (blue) after treatment with DMSO as control and inhibitors of chromatin-

430   modifying enzymes. White scale bars= 50 μm.

431   **(*D*)** Percentage of cells expressing Sox1 (green) over DAPI (blue) total cells. $*P < 0.05$, $**P <$

432   0.01, $***P < 0.001$.

Figure 3

B

Neurite length

Fig. 3. Upregulated GASC1 impairs differentiation from hESCs to iN cells.

(*A*) Brightfield and fluorescent images of hESC-derived iN cells with wt, dox-induced overexpression of GASC1, DNMT3A, and KDM4A stained for the pluripotent marker Sox2 (green), the neural cells marker Tuj1 (red), and nucleus staining DAPI (blue) after 14 days of iN cells differentiation.

439   (**B**) Neurite length of hESC-derived iN cells was measured at 14 days of iN cells differentiation.

440   p-values were calculated by unpaired t-test, error bars represent standard deviation, *P < 0.05, ***

441   P< 0.001.

Figure 4

Fig. 4. GASC1 promotes aberrant nuclear structure and cell cycle arrest of leukemia cells.

(*A*) Representative images of DAPI staining for K562 cells treated with DMSO as control and 5-Aza, IOX1, GSK-LSD1, and JIB04. It was shown that the bigger size and twisted shape of nuclei in inhibitors treatment. The images were taken at 40× on a confocal microscope. White scale bars= 25 μm.

(**B**) Measurements of the nuclear size using DAPI staining under fluorescence microscopy after

K562 cells treated with inhibitors of chromatin-modifying enzymes. Two-tailed t-test, Mean $\pm$ SD.

**P $<$ 0.01, ***P $<$ 0.001.

(**C**) Brightfield and fluorescent images show that K562 leukemia cells differentiate to iN cells with

Ngn2 overexpression after 14 days differentiation and stain with the neuronal marker Tuj1 (red),

compared to healthy and as cultured K562 cells without any treatment. The yellow arrows indicate

the cells of morphological changes.

(**D and E**) Brightfield images of K562 leukemia cells with induced expression of GASC1 with or

without doxycycline (dox). The no dox-induced GASC1 overexpression K562 cells show spherical

morphology, same as normal cells. The dox-induced GASC1 overexpression K562 cells show

morphology change from spherical to twisted shape and cell blebs. The red arrow indicates the

cells with twisted shapes and blebs. White scale bars= 25 μm.

(**F and G**) Fluorescent images by DAPI staining demonstrate that the dox-induced GASC1

overexpression K562 cells show the twisted nuclear shape and bigger nuclei compared with the no

dox-induced GASC1 overexpression K562 cells. The red arrow indicates the enlarged and twisted

nucleus. White scale bars= 25 μm.

(**H and I**) Flow cytometric analysis of K562 cells were double-stained with BrdU and PI. The dox-

induced GASC1 overexpression K562 cells proliferate significantly slower than the no dox-

induced GASC1 overexpression K562 cells by BrdU assay.

(**J**) Cell number counting after induction by dox for 7 days. The cell number is decreased in dox-

induced GASC1 overexpression K562 cells than no dox-induced GASC1 overexpression K562

cells. ***P < 0.001. n=3.

(**K**) The genes play a role in cell-cycle arrest detected by qPCR in the dox-induced GASC1

overexpression K562 cells and no dox-induced GASC1 overexpression K562 cells. Unpaired two-

tailed Student t-test. Data are mean ± SD. P < 0.001.

REFERENCES

1. A. C. Tamashunas *et al.*, High-throughput gene screen reveals modulators of nuclear shape. *Mol Biol Cell* **31**, 1392-1402 (2020).

2. R. McBeath, D. M. Pirone, C. M. Nelson, K. Bhadriraju, C. S. Chen, Cell shape, cytoskeletal tension, and RhoA regulate stem cell lineage commitment. *Dev Cell* **6**, 483-495 (2004).

3. E. Bartova, J. Krejci, A. Harnicarova, S. Kozubek, Differentiation of human embryonic stem cells induces condensation of chromosome territories and formation of heterochromatin protein 1 foci. *Differentiation* **76**, 24-32 (2008).

4. J. D. Pajerowski, K. N. Dahl, F. L. Zhong, P. J. Sammak, D. E. Discher, Physical plasticity of the nucleus in stem cell differentiation. *Proc Natl Acad Sci U S A* **104**, 15619-15624 (2007).

5. H. Shen, W. Xu, F. Lan, Histone lysine demethylases in mammalian embryonic development. *Exp Mol Med* **49**, e325 (2017).

6. J. He *et al.*, Kdm2b maintains murine embryonic stem cell status by recruiting PRC1 complex to CpG islands of developmental genes. *Nat Cell Biol* **15**, 373-384 (2013).

7. J. D. Pereira *et al.*, Ezh2, the histone methyltransferase of PRC2, regulates the balance between self-renewal and differentiation in the cerebral cortex. *Proc Natl Acad Sci U S A* **107**, 15957-15962 (2010).

8. Y. H. Loh, W. Zhang, X. Chen, J. George, H. H. Ng, Jmjd1a and Jmjd2c histone H3 Lys 9 demethylases regulate self-renewal in embryonic stem cells. *Genes Dev* **21**, 2545-2557 (2007).

9. F. Kottakis *et al.*, NDY1/KDM2B functions as a master regulator of polycomb complexes and controls self-renewal of breast cancer stem cells. *Cancer Res* **74**, 3935-3946 (2014).

10. I. Kupershmit, H. Khoury-Haddad, S. W. Awwad, N. Guttmann-Raviv, N. Ayoub, KDM4C (GASC1) lysine demethylase is associated with mitotic chromatin and regulates chromosome segregation during mitosis. *Nucleic Acids Res* **42**, 6168-6182 (2014).

11. Z. Q. Yang *et al.*, Identification of a novel gene, GASC1, within an amplicon at 9p23-24 frequently detected in esophageal cancer cell lines. *Cancer Res* **60**, 4735-4739 (2000).

12. P. A. Northcott *et al.*, Multiple recurrent genetic events converge on control of histone lysine methylation in medulloblastoma. *Nat Genet* **41**, 465-472 (2009).

13. Z. Xiao *et al.*, GASC1 promotes glioma progression by enhancing NOTCH1 signaling. *Mol Med Rep* **23**,   (2021).

14. G. Liu *et al.*, Genomic amplification and oncogenic properties of the GASC1 histone demethylase gene in breast cancer. *Oncogene* **28**, 4491-4500 (2009).

15. P. A. Cloos *et al.*, The putative oncogene GASC1 demethylates tri- and dimethylated lysine 9 on histone H3. *Nature* **442**, 307-311 (2006).

16. L. Wu *et al.*, Histone Demethylases KDM4A and KDM4C Regulate Differentiation of

Embryonic Stem Cells to Endothelial Cells. *Stem Cell Reports* **5**, 10-21 (2015).

17.   H. Richly, L. Aloia, L. Di Croce, Roles of the Polycomb group proteins in stem cells and
cancer. *Cell Death Dis* **2**, e204 (2011).

18.   S. J. Heo *et al.*, Differentiation alters stem cell nuclear architecture, mechanics, and
mechano-sensitivity. *Elife* **5**,   (2016).

19.   S. M. Chambers *et al.*, Highly efficient neural conversion of human ES and iPS cells by
dual inhibition of SMAD signaling. *Nat Biotechnol* **27**, 275-280 (2009).

20.   Y. Zhang *et al.*, Rapid single-step induction of functional neurons from human pluripotent
stem cells. *Neuron* **78**, 785-798 (2013).

21.   T. A. Volpe *et al.*, Regulation of heterochromatic silencing and histone H3 lysine-9
methylation by RNAi. *Science* **297**, 1833-1837 (2002).

22.   K. K. Lee, J. L. Workman, Histone acetyltransferase complexes: one size doesn't fit all.
Nat Rev Mol Cell Biol **8**, 284-295 (2007).

23.   A. D. Stephens *et al.*, Chromatin histone modifications and rigidity affect nuclear
morphology independent of lamins. *Mol Biol Cell* **29**, 220-233 (2018).

24.   E. G. Fischer, Nuclear Morphology and the Biology of Cancer Cells. *Acta Cytol* **64**, 511-
519 (2020).

25.   G. N. Papanicolaou, H. F. Traut, The diagnostic value of vaginal smears in carcinoma of
the uterus. 1941. *Arch Pathol Lab Med* **121**, 211-224 (1997).

26.   T. Sullivan *et al.*, Loss of A-type lamin expression compromises nuclear envelope integrity
leading to muscular dystrophy. *J Cell Biol* **147**, 913-920 (1999).

27.   L. Dang *et al.*, Cancer-associated IDH1 mutations produce 2-hydroxyglutarate. *Nature* **462**,
739-744 (2009).

28.   C. Lu *et al.*, IDH mutation impairs histone demethylation and results in a block to cell
differentiation. *Nature* **483**, 474-478 (2012).

29.   M. Koivomagi, M. P. Swaffer, J. J. Turner, G. Marinov, J. M. Skotheim, G1 cyclin-Cdk
promotes cell cycle entry through localized phosphorylation of RNA polymerase II.
Science **374**, 347-351 (2021).

30.   J. A. Coffman, Cell cycle development. *Dev Cell* **6**, 321-327 (2004).

31.   L. Litovchick *et al.*, Evolutionarily conserved multisubunit RBL2/p130 and E2F4 protein
complex represses human cell cycle-dependent genes in quiescence. *Mol Cell* **26**, 539-551
(2007).

32.   G. Galiova, E. Bartova, I. Raska, J. Krejci, S. Kozubek, Chromatin changes induced by
lamin A/C deficiency and the histone deacetylase inhibitor trichostatin A. *Eur J Cell Biol*
87, 291-303 (2008).

33.   P. Scaffidi, T. Misteli, Lamin A-dependent misregulation of adult stem cells associated with
accelerated ageing. *Nat Cell Biol* **10**, 452-459 (2008).

34.   E. Meshorer *et al.*, Hyperdynamic plasticity of chromatin proteins in pluripotent embryonic
stem cells. *Dev Cell* **10**, 105-116 (2006).

35. E. Meshorer, T. Misteli, Chromatin in pluripotent embryonic stem cells and differentiation.
Nat Rev Mol Cell Biol **7**, 540-546 (2006).
36. C. Gurrion, M. Uriostegui, M. Zurita, Heterochromatin Reduction Correlates with the
Increase of the KDM4B and KDM6A Demethylases and the Expression of Pericentromeric
DNA during the Acquisition of a Transformed Phenotype. *J Cancer* **8**, 2866-2875 (2017).
37. B. E. Bernstein *et al.*, A bivalent chromatin structure marks key developmental genes in
embryonic stem cells. *Cell* **125**, 315-326 (2006).
38. T. S. Mikkelsen *et al.*, Genome-wide maps of chromatin state in pluripotent and lineage-
committed cells. *Nature* **448**, 553-560 (2007).
39. Y. Katoh, M. Katoh, Comparative integromics on JMJD2A, JMJD2B and JMJD2C:
preferential expression of JMJD2C in undifferentiated ES cells. *Int J Mol Med* **20**, 269-273
(2007).
40. T. Brambrink *et al.*, Sequential expression of pluripotency markers during direct
reprogramming of mouse somatic cells. *Cell Stem Cell* **2**, 151-159 (2008).
41. Z. P. Pang, P. Cao, W. Xu, T. C. Sudhof, Calmodulin controls synaptic strength via
presynaptic activation of calmodulin kinase II. *J Neurosci* **30**, 4132-4142 (2010).
42. M. Stadtfeld, N. Maherali, D. T. Breault, K. Hochedlinger, Defining molecular
cornerstones during fibroblast to iPS cell reprogramming in mouse. *Cell Stem Cell* **2**, 230-
240 (2008).
43. C. Chen, H. Okayama, High-efficiency transformation of mammalian cells by plasmid
DNA. *Mol Cell Biol* **7**, 2745-2752 (1987).