Title: The neurodevelopmental genes *alan shepard* and *Neuroglian* contribute to female mate

preference in African *Drosophila melanogaster*

Abstract:

Mate choice is a key trait that determines fitness for most sexually reproducing organisms, with

females often being the choosy sex. Female preference often results in strong selection on male

traits that can drive rapid divergence of traits and preferences between lineages, leading to

reproductive isolation. Despite this fundamental property of female mate choice, very few loci

have been identified that contribute to mate choice and reproductive isolation. We used a

combination of population genetics, quantitative complementation tests, and behavioral assays to

demonstrate that *alan shepard* and *Neuroglian* contribute to female mate choice, and could

contribute to partial reproductive isolation between populations of *Drosophila melanogaster*.

Our study is among the first to identify genes that contribute to female mate preference in this

historically important system, where female preference is an active premating barrier to

reproduction. The identification of loci that are primarily known for their roles in

neurodevelopment provides intriguing questions of how female mate preference evolves in

populations via changes in sensory system and higher learning brain centers.

Introduction

For sexually reproducing organisms mate choice is critical to balance maximizing fitness

with avoiding costly heterospecific mating events (Mendelson and Shaw, 2012; Gray, 2022).

Strong selection on male courtship traits and female preferences can drive rapid divergence

between closely related lineages (Fisher, 1958; Andersson, 2019), providing a link between

sexual selection and the evolution of reproductive isolation. A large body of work has focused on

documenting and describing female preference and identifying genes controlling male courtship

traits (Chenoweth and Blows, 2006; Yamamoto and Koganezawa, 2013; Neelon et al., 2019;

Munson et al., 2020). However, there have been fewer studies that have identified specific genes

that contribute to female mate preference, with most instead focusing on quantitative trait loci

(Bay et al., 2017; Svensson et al., 2017; Blankers et al., 2019; Xu and Shaw, 2019; Chowdhury

et al., 2020) and some identifying specific genes (Chowdhury et al., 2020). Identifying and

characterizing loci contributing to female preference is a critical first step to understand how

courtship signal detection, signal processing, and higher brain functions (i.e. learning and

memory) contribute to sexual selection and female mate preference evolution.

One challenge in identifying the genetic basis of female choice lies in the complexity of

preference-based behaviors (Andersson, 2019). During mate choice females are receiving many,

potentially multimodal, signals that then require processing before a decision to accept or reject a

mate is made (Gray, 2022; Munson et al., 2020). Progress has been made to understand male

courtship and how female preference exerts divergent selection between populations (Xu and

Shaw 2019a; Xu and Shaw 2021), yet even in systems where male courtship traits, and

presumably male signals, are well understood, identifying female loci can be difficult (Xu and

Shaw 2019b). The complexity of female preference implies that this trait is polygenic, but it is

unclear how many loci we would expect to contribute to this trait. Specifically, it remains

unclear whether multiple loci interact in an emergent fashion, contributing to female preference

as a whole, or if individual loci contribute to a specific facet (perception of a specific signal/cue)

of female preference (Boake et al., 2002; Chenoweth and Blows, 2006). Additionally, while

premating isolation evolves rapidly and is often thought to evolve before other barriers to

reproduction (Coyne and Orr, 1989; Turissini et al., 2018), it can be difficult to determine if the

behaviors that currently act as a mating barrier contributed to the speciation event or evolved

after species divergence (Coyne and Orr, 2004; Safran et al., 2013; Kopp et al., 2018). Focusing

on populations that vary in the degree of mating isolation can overcome some of these challenges

and can be used as a model to understand the early stages of divergence and identify the loci that

contribute to speciation and reproductive isolation.

Drosophila melanogaster originated in Southern Africa but is now a cosmopolitan

species with a worldwide distribution. This broad range has resulted in structured and cryptic

populations, even within Africa (Coughlan et al 2021), and strong premating isolation between

some specific populations (Wu et al., 1995). For example, females from strains collected in

Southern Africa strongly reject males from strains that are collected from non-African localities

in both choice and no-choice mating experiments (Wu et al., 1995; Hollocher et al., 1997;

Coughlan et al., 2021). These strong premating isolation behaviors are also observed in other

parts of the *D. melanogaster* range including the Southeast United States and the Caribbean

(Yukilevich and True, 2008). These mating preferences and behaviors that occur outside of

Africa are potentially a product of African ancestry in these populations and admixture with non-

African populations (Kao et al 2015; Bergland et al 2016). Combined, these observations of

strong and variable female preference and population structure provide an ideal system to

explore the evolution of female mate preference.

In this study, we leverage these *D. melanogaster* populations by combining population

genomics and behavioral assays to identify genes that contribute to female mate preference and

partial reproductive isolation. First, we use previously published genomic data from Kao et al.

(2015) to identify loci that show a clinal pattern of allele frequency in an admixed secondary

contact zone, suggesting a role in reproductive isolation (Harrison and Larson, 2014). We

compare these candidates to outlier loci from other studies allowing us to narrow our list and

focus on two loci, *alan Shephard* (*shep*) and *Neuroglian* (*Nrg*).We then examine patterns of

differentiation in these loci between African and non-African genomes and further provide

genetic evidence that these two loci contribute to African female mate preference using

quantitative complementation tests.

Methods

Genomic cline analysis

To understand how alleles potentially involved in reproductive isolation are structured we

compared allele frequencies across the Southeastern United States and Caribbean using

previously published data (Kao et al 2015). We focused on these populations because this clinal

structure allowed us to make explicit expectations for a gene that contributes to reproductive

isolation compared to the rest of the genome (Harrison and Larson 2014). Specifically, because

previous studies have demonstrated a correlation between latitude and African ancestry, with

more African ancestry in southern populations, we predict that genes involved in behavioral

isolation will have a steeper slope (ie larger regression coefficient) compared to the rest of the

genome. Using the genotype matrix (genotype calls for all variable positions they identified)

from Kao et al. (2015), we fit linear regression models between SNP allele frequency and

latitude for the populations.

<u>GO Enrichment Analysis:</u> We retained the top 1% outliers for use in a gene ontology enrichment

analysis. The list of genes used for the enrichment analysis are found in Supplemental Table 1.

This list included genes that contained outlier SNPs. SNPs that were not within the known

boundaries of a gene were not included in this analysis. That is, we did not assign SNPs to the

nearest gene. We used the web application FlyEnrichr (Chen et al. 2016; Kuleshov et al. 2016),

to look for enrichment in "biological processes". These biological processes are defined by

Flybase (The Gene Ontology Consortium 2017) and FlyEnrichr uses definitions made in the

2018 version of Flybase.

<u>Comparison to other datasets: We compared our list of outlier genes to two other datasets that</u>

<u>looked for differences in behavior between African and non-African flies using either DNA</u>

<u>sequence data or RNAseq transcriptomic data. In the first comparison we compared our set of</u>

<u>outliers to Coughlan et al. (2021). In this study the authors compared allele frequency between</u>

<u>strains that showed strong female choice and strains that did not show strong choice and</u>

<u>identified outlier loci using the PBE test statistic. In the second comparison we compared our list</u>

<u>of outliers to Bailey et al. (2011) who looked at transcriptomic data in female brains after they</u>

<u>interacted with "preferred" and "non-preferred" males in both African and non-African strains.</u>

Choice of candidate loci

After calculating regression coefficients for each SNP, we found that two genes previously

identified as important for female behavior *alan shepard* (*shep*; Chen et al., 2014) and

Neuroglian (*Nrg;* Carhan et al., 2005) were genomic outliers (See Results). These genes were

outliers in at least one of the additional data sets we compared our lists with (see Results) and

appeared as key loci contributing to significant GO terms in the enrichment analysis (See

Results). Previous data for these genes demonstrated that null mutant females rejected were

mating deficient, and we used these mutants to test for a specific role in African female behavior.

Population genetics of shep and Nrg across Drosophila populations

Several studies have examined the population genetics of *Drosophila* populations from

worldwide distributions (Lack et al. 2016; Kapopoulou et al. 2018; Sprengelmeyer et al. 2020),

but our focus was to specifically look at variation in *shep* and *Nrg*. The SNPs identified in Kao et

al (2015), the source of our cline analysis, likely represent a subset of the total variation observed

in these two loci, since the focused on SNPs that they could define as coming from either the

African or European source population. To understand the variation in these alleles and

population differentiation we used the PopFly application (Hervas et al. 2017) to provide

summary statistics for nucleotide diversity (π) and differentiation (F_{st}) for the two populations

with the largest number of sequences strains, Raleigh (RAL) and Zimbabwie (ZI) (Lack et al.

2016). We also exported FASTA sequences from PopFly to examine differentiation using

principal components analysis. This allowed to visualize and estimate the variation of

alleles/haplotypes within and between populations.

Quantitative complementation test

To determine if *Nrg* and *shep* alleles contribute to the female mate preference of African strains,

we used a quantitative complementation test combined with a binary mate choice assay

(Supplemental Figure 1). Quantitative complementation tests have previously been used for no-

choice tests (Chowdhury et al., 2020) and binary choice tests (Comeault et al., 2017). In a typical

complementation test, the starting point is two parental strains that have divergent phenotypes for

a continuous trait and are often the minimum and maximum trait values in a population

(reviewed in Mackay 2001). The trait of interest may be polygenic and hybrids have an

intermediate phenotype. A third strain is used that carries a genetic deficiency, a large deletion,

for a gene that is suspected to contribute to the trait of interest. The deficiency may contain many

genes, or just the gene of interest. In *Drosophila* these deficiencies are "balanced" by a

chromosome, known as a balancer chromosome, that has several complex inversions that

prevents recombination with the homologous chromosome (Muller 1918; Miller et al. 2019).

Balancer chromosomes also carry a dominant visible mutation to track the presence/absence of

the balancer chromosome. When both wild-type parental strains are crossed independently to the

deficiency strain (that also carries the balancer chromosome) four genotypes are produced and

can be phenotyped (Supplemental Figure 1). Two of the genotypes are heterozygous at the gene

of interest. They carry either parental allele and the allele from the balancer chromosome. The

other two genotypes are hemizygous. They only carry one the the parental alelles. The

homologous chromosome either lacks the gene of interest or has a non-functional allele. In

Drosophila balancer chromosomes have a marker with a dominant visible phenotype. This

causes the heterozygote individuals, but not hemizygote individuals, to have a phenotype that

could affect the trait of interest and must be taken into account during analysis (Turner et al.,

2011). Additional details about complementation tests using preference behavior and

considerations when using quantiative complementation tests are included in Supplemental

Information.

Using a binary choice test, we introduced a single female of a given genotype (one of the

four genotypes produced from the complementation tests crosses) to two males from

representative strains: Z53 representing an African male and DGRP882 representing a non-

African. Trios were observed for 60 minutes or until copulation occurred. When copulation

occurred between a pair of individuals, the non-copulating male was removed and the strain

identity was determined (see behavioral assay conditions, below). After the pair was finished

copulating we also examined the sucessful male to verify which strain the female chose. Our *a*

priori expectation was that the females hemizygous for the African allele will prefer African

males, whereas females hemizygous for the non-African allele will mate indiscriminately.

Drosophila strains and crosses

To determine whether *Nrg* and *shep* alleles contribute to female preference behavior

between populations of *D. melanogaster* we needed to identify wild-type strains that show strong

female mate preference and complementary strains that show no preference. Strains from

Southern Africa were once considered to be a single lineage from the ancestral range and were

referred to as Z-type. All other strains that migrated from the ancestral range (non-African and

some Northern African strains) were considered a single cosmopolitan lineage and referred to as

M-type. The M-type behavior contrasted with Z-type behavior in that M-type females often mate

indiscriminately, showing no preference, whereas Z-type females strongly reject M-type males

(Wu et al., 1995). Since this initial description, the demographic and evolutionary relationships

between *D. melanogaster* populations have been shown to be more complex (Coughlan et al.,

2021), and it is unlikely that all Z-type strains are a homogenous lineage.

We were able to choose strains based on geographic location and evaluated the female

preference behavior for wild type females prior to using these strains in our crosses. For our

representative African strain, we chose Z53 because its behavior profile is well documented and

has been consistent across studies (Wu et al., 1995; Hollocher et al., 1997; Moran, 2006; Jin et

al., 2022). For our representative non-African strain, we chose DGRP882 which we had

previously used and had expected baseline behavior (Jin et al., 2022). The African admixture has

been estimated for individuals of the DGRP panel (Pool, 2015) and DGRP882 has among the

least amount of African ancestry making it most likely to be differentiated from African strains.

In addition to these representative lines we also chose two additional African strains: Z30 and a

more recently collected strain CH11. Canton-S was used as an additional non-African strain

because it is commonly used in behavioral studies (Tompkins et al., 1982; Lasbleiz et al., 2006;

Ng and Kopp, 2008; Kohlmeier et al., 2021).

To create heterozygote and hemizygote genotypes for our complementation tests, we

chose to use loss of function mutant lines for both *shep* and *Nrg* that had been previously

characterized as null alleles (*shep* Chen et al., 2014; *Nrg* Bieber et al., 1989; Enniking et al.,

2013). Wild type females that have not mated (called virgins) readily mate with males. *shep* and

Nrg mutant females that are virgin do not mate and instead show rejection behaviors when

courted by males (*shep* Chen et al., 2014; *Nrg* Kerr et al. 1997; Carhan et al. 2005).We used the

shep^{BG00836} null mutation (BDSC #12513), hereafter *shep$^-$*. Homozygous null mutant females for

this allele are viable and mating deficient. For our complementation test we wanted to use a

strain that carried *shep$^-$* on one chromosome while the homologous chromosome was a balancer

chromosome. We combined the *shep$^-$* allele with the TM6B balancer chromosome. This

balancer chromosome carries the dominant visible Humoral (*Hu)* mutation. We inbred progeny

from this original cross for three generations before using this strain in our behavioral assays.

The mutant line and the line that contributed the TM6B chromosome were not the same genetic

background so residual heterozygosity could be present in this strain. Any effects of this

heterozygosity on the X chromosome and 2^{nd} chromosome would occur in all of the

crosses/treatments that we examined.

207   We used the Nrg^{17} allele, which is synonymous with Nrg^2 and hereafter Nrg^- (BDSC

208   #5595). This specific Nrg^- allele is homozygous lethal, so the stock is maintained with one X

209   chromosome carrying Nrg^- and the other X chromosome is the balancer FM7 chromosome. Fm7

210   carries the dominant visible marker *Bar*.

211   Importantly both the *shep$^-$* and *Nrg$^-$* strains are non-African and carry non-African alleles

212   of *shep* or *Nrg* on the respective balancer chromosomes. These alleles are not identical to

213   DGRP882 or Canton S alleles, however, all non-African alleles are phenotypically more similar

214   than they are to are to the African alleles, because they result in random mating.

216   *Experimental Crosses*

217   We crossed the *shep*/TM6B line to both DGRP882 and Z53 males to generate four F1

females. Non-African heterozygotes (N/N) have two non-African *shep* alleles, one from the

TM6B chromosome and one from the DGRP882 chromosome. Non-African hemizyogtes (N/N-

shep$^-$) only have the non-African *shep* allele. We designate the *shep$^-$* chromosome as N-*shep$^-$* to

highlight that the rest of the chromosome carries non-African (N) alleles. African heterozygotes

(A/N) have an African allele from the Z53 chromosome and a non-African allele from the TM6B

chromosome. African hemizyogotes (A/N-*shep$^-$*) only carry the African allele of *shep*.

We crossed the Nrg^-/FM7 line to DGRP882 and Z53 similar to the design for *shep.* For

experiments with *Nrg,* we tested two additional African and one non-African strains (described

above) for two reasons. First, we were not sure how or if the *Bar* locus, which affects eye

morphology, would affect female preference. This is particularly important because the impact of

vision for female mating behavior could vary between strains. We tested the effect of *Bar*

explicitly by including Canton S as an additional strain that should not exhibit strong mating

preference. Second, even though our cline analysis identified *Nrg* as a strong outlier, this locus

was only a marginal outlier in and independent study that identified putative female preference

loci (Coughlan *et al.*, 2021). This could indicate that the affect of *Nrg* differs between strains and

populations, or multiple alleles of *Nrg* segregate in African populations with only a subset

contributing to female preference phenotypes. Either scenario could result in a situation where

there is not a single high frequency allele contributing to female preference in African strains

motivating us to test multiple African strains for this locus.

Behavioral assay conditions

All stocks and virgins were kept on standard cornmeal molasses media in an incubator on

a 12:12 light cycle held at a constant 20 degrees Celsius and 50% humidity. Virgin males and

females were collected within the first 0-4 hours post eclosion to ensure they had not mated.

Females were phenotyped based on the presence of the dominant visible marker and housed in

groups of 5-10. Males were housed individually because group housing can potentially change

courtship vigor (Dixon *et al.* 2003). All individuals were allowed to age 7-10 days before

experiments because this is the timeframe that produces the maximum number of matings for the

African strains (Jin et al 2022). All behavioral assays were performed in a room held at a

constant 20C degrees Celsius and 50% humidity within two hours of the incubator lights turning

on to maximize the number of copulations we could observe.

The strains used in the experiment are phenotypically indistinguishable, so to be able to

tell males apart we placed them on food containing blue food coloring 48 hours prior to the

behavioral assay. This is a non-invasive, robust identification method (Wu et al., 1995;

Hollocher et al., 1997). In these studies, all males received food dye treatment, but in our

experiment, we only marked one male per mating with blue food dye. We found that we could

more easily distinguish between a dyed and non-dyed male rather than distinguishing between

males fed two different colors. We tested the effect of this treatment on female preference and

found no effect (Supplementary Information).

We used males from the representative strains Z53 and DGRP882 for all binary choice

tests. We collected all genotypes for a given locus in each block of our trials in equal numbers,

but due to differences in the available genotypes after aging, our final number of females that

were included differed for each genotype, but all genotypes were represented in each block.

While the rearing conditions and behavior rooms conditions were very well controlled, our

behavioral protocol also included completing a set of DGRP882 females in parallel to our

experimental crosses. For each block of trials, 20 DGRP882 females were used as a behavioral

control and we consistently had 8-12 (representing 40-60%) choosing Z males consistent with

the random mating for this genotype, allowing us to conclude that the environment was not

affecting our matings and pool replicates over blocks.

Statistical analysis of behavioral data

To analyze data from a quantitative complementation test using continuous traits, a linear model

of the following structure is used (Equation 1; Pasyukova and Mackay 2000; Mackay 2001). The

goal is to isolate the effects of the gene of interest on the phenotype while taking into account the

difference in parental strains and any effects the balancer chromosome might have on the

phenotype.

$$y = \mu + \beta_1 x_1 + \beta_2 x_2 + \beta_3 x_1 x_2 \qquad \text{Equation 1}$$

A full description of this analysis framework is included in the Supplemental Information.

For our behavioral preference experiments the data can be represented as binary choices.

An individual female replicate chooses either the DGRP882 or the Z53 male, which can be

represented as 0 and 1. Our statistical philosophy is to construct a model based on the properties

of the data (Warton et al. 2016) and would lead us to using binomial regression. One benefit of

the generalized linear models is that the same model structure can be used as Equation 1, except

now the scale is in log-odds due to the logit transformation of the data. The logit link function is

the most common link function for binomial regression (Equation 2).

$$logit(p) = \mu + \beta_1 x_1 + \beta_2 x_2 + \beta_3 x_1 x_2 \qquad \text{Equation 2}$$

While this approach matches the rich literature of complementation tests, there are some barriers

to using this model, specifically in interpreting the interaction term for binomial regression and

the lack of correspondence between interaction on the probability and logit scales. In the

binomial regression the interaction is modelled as a multiplicative increase in odds (Hosmer et

al., 2013) and because of the logit link the interaction is non-linear. As a result, there are well

documented examples where even when a significant interaction effect for the logit model exists

there will not always be a corresponding significant interaction on the probability scale (Hosmer

et al., 2013). For a behavioral preference experiment, we are ultimately interested in the

probability scale and effects at this level since this would represent the preference of genotypes.

The lack of correspondence between scales can be observed in the reverse comparison as well.

When there is a multiplicative increase in the predicted probabilities on the probability scale

(which would be of interest in our analysis) it is possible that there will not be a multiplicative

increase in the odds, and no significant interaction term in the logistic model (Hosmer et al.,

2013).

When a significant interaction occurs in the binomial regression neither the value nor the

sign of the coefficients gives us clear information about the nature of the interaction at the

probability scale (Ai and Norton 2003; Chen 2003). When the affects at the probability scale are

of interest it is recommended to further probe the interaction by evaluating simple slopes models

(Jaccard, 2001). This is also a solution to scenarios where there is discordance between the

presence/absence of interaction on the probability vs log odds scale and alternative models can

generated to explore the interaction effect (see below). This is important because in behavior we

are interested in the probability scale since this matches our estimate of female preference. We

discuss aspects of our data and compare them to other behavioral complementation tests to

demonstrate the need for alternative models in the Supplemental Information.

In the alternative model, instead of the interaction being coded as x1x2 we can capture

the effects of that genotype as its own variable x3 (Equation 3).

310                 $$logit(p) = \mu + \beta_1 x_1 + \beta_2 x_2 + \beta_3 x_3$$ Equation 3

While we lack the ability to test for an interaction explicitly, we estimate the same number of

coefficients as in as the interaction model. Since we do not expect strain effects ($\beta_1=\beta_2=0$) we

can now model the trait mean of each genotype independently. Using a Wald's test tested for

significant difference between β_2 and β_3 which would indicate a significant shift in preference

for the African hemizygote genotype equivalent to what we would observe if we combined our

coefficients from the interaction model to estimate the effect of this genotype.

In the results we report both the interaction models and the proposed alternative models.

For interaction models we report coefficients and their corresponding p-values to remain

consistent with other complementation test studies. For the alternative model we calculated 95%

confidence intervals and determined whether any interval contained 0. We also provide results

for the Wald's test specifically testing for differences in African genotypes. In the supplemental

material we provide results for non-interaction models. The reason for these models is to

compare them with interaction models and make observations about the stability, or lack thereof,

of coefficients. This could suggest a poor fit of this group of models in estimating coefficients

for our specific data and experimental design.

Results

Female behavior loci are genomic outliers

To generate candidate genes for functional validation, we leveraged genomic data from

populations of *D. melanogaster* that exhibit a cline in African ancestry (Kao et al., 2015;

Bergland et al., 2016) and African-like female preference behavior (Yukilevich and True, 2008).

Our goal was to identify outlier genes that showed a steeper cline in African ancestry compared

to the genome wide average. We used these genes to understand patterns of gene ontology

enrichment and compared our list of candidates to independent analyses that used populations

from Africa to identify female preference alleles (Coughlan et al., 2021) and changes in gene

expression in female brains between these populations (Bailey et al. 2011).

We retained the top 1% of outliers for loci that had the steepest clines with African alleles

being more frequent at lower latitudes. We determined whether SNPs occurred in annotated

genes, and retained these genes for a gene ontology (GO) enrichment analysis (Supplemental

Table 1). We focused on biological function and observed a pattern where axon and neuron

maintenance and development were enriched in our top 10 categories (Table 1). Courtship

behavior also showed significant enrichment, and within that list were two genes, *alan shepard*

(*shep*) and *Neuroglian* (*Nrg*) that contribute specifically to female behavior. Interestingly *Nrg*

shows up as a gene contributing to many of the GO terms that show significant enrichment.

344   We next compared our list of outliers to those outliers identified in Coughlan et al (2021).

Coughlan et al (2021) conducted a GO enrichment analysis, with a different focus, that returned

similar terms including behavior, mushroom body formation (neurogenesis), and olfaction as

enriched terms (Fig 4 of Coughlan et al 2021). When comparing the list of outliers highlighted in

that study (Supplemental Table 2) to our list of outliers we found five genes in common. These

included *shep*, *Rdl*, *rad*, *RunxA*, and *Dop2R*. *Nrg* was in the top 5% of outliers from the

Coughlan et al 2021 study. In our final comparison we compared our list of outliers to genes that

showed significantly different transcription in African and non-African female brains after

exposure to both types of males (Bailey et al. 2011). In total there were 36 genes shared between

these lists (Supplemental Table 3). The only gene that overlapped in all three studies was *shep*.

354   Given the patterns from GO enrichment analyses and comparisons with other data sets,

we focused on *shep* and *Nrg* for functional tests. Previous behavioral analysis of homozygous

viable mutant strains for these loci found evidence that these genes were involved female

behavior (Carhan et al., 2005; Chen et al., 2014). Our goal was to use natural variation in *shep*

and *Nrg* alleles to test for a role specifically in African female mate preference behavior.

Population genetics and differentiation of shep and Nrg

Using previously published genomic datasets that sampled a large number of strains, we

looked at patterns of nucleotide diversity (π) within two populations, patterns of differentiation

(Fst) between these populations, and structure for *Nrg* and *shep*. π and Fst were summarized for

1 kb windows and compared to genome wide averages found in Lack et al (2016). For both *shep*

and *Nrg* there was variability across the gene region for π and Fst (Table 2). While we would not

necessarily expect the entire locus to be differentiated, *Nrg* had an average Fst that was greater

than the genome wide average. The position of this locus on the X-chromosome could contribute

to this pattern (reviewed in Meisel and Connallon 2013). Within both *shep* and *Nrg* there were

windows with elevated Fst compared to the genome average. This differentiation was also

captured by looking at the clustering of strains/genotypes in principal component space (Figure

2). The strains largely clustered by population for both loci. For *shep* there was also structuring

within the ZI population, likely due to the presence of the segregating In(3L)P inversion in this

region of the genome (Pool et al 2012; Corbet-Dettig and Hartl 2012).

Shep contributes to female mate preference in an African strain

We used quantitative complementation tests to determine if *alan shepard* (*shep*) contributes to

female mate preference. We were able to validate that *shep* contributes to female mate

preference by specifically demonstrating that females that carry only the African *shep* allele

(African hemizygotes: A/N-*shep⁻*) prefer African males to non-African males (Figure 3A). When

these data were analyzed using a binomial regression that included an interaction, the model did

not contain any significant coefficients (Table 3). This is not unexpected given the specifics of

our data, the phenotypes of F1s from our crosses used, and the potential discordance between

interactions on the probability vs log-odds scales (See Methods and Supplemental Information

for a full discussion). Given the clear trend for the African hemizygote females preference, we

followed up with an alternative model that was chosen specifically to overcome these limitations

(see Supplemental Information). This analysis indicated that the African hemizygote females

(A/N-*shep*⁻) were the only genotype that had a significant mating preference (Table 4). All other

genotypes had 95% confidence intervals that contained zero indicating that they were consistent

with random mating. We compared preference of the African hemizygote (A/N-*shep*⁻) directly to

the preference of African heterozygote (A/N) using a Wald's-test. This test indicated a

significant difference in preference between these two genotypes (χ^2_2 = 13.0, P = 0.0015). Both

genotypes are primarily heterozygous across the entire genome for non-African and African

alleles since they are hybrids. The main difference between these genotypes is at the *shep* locus.

Nrg contributes female mate preference in African strains

We also used quantitative complementation tests to validate the role of *Nrg* alleles on female

mate preference in African strains. Our design and logic were identical for what we used to

analyze *shep*. For this gene we included more genotypes to determine if the visible marker on the

FM7 balancer chromsome, *Bar*, which affects eye phenotypes was in turn affecting mating

behavior. We also tested whether different African *Nrg* alleles might show different affects on

female mate preference.

To test for the affects of the *Bar* allele specifically on female behavior we tested the

behavior of another non-African strain, Canton S, and compared it to the DGRP882 strain. We

used Canton S because we had a clear expectation: that both female genotypes from a cross

between Canton S and the *Nrg*⁻ strain should have the same behavior as the DGRP882

genotypes, and they should mate randomly and not show preference for either male genotype.

We in fact observed that these genotypes mated indiscriminately, and we did not detect any

significant difference in behavior comparing DGRP882 genotypes and Canton S genotypes

(Supplemental Figure 2; Supplemental Table 5).

We then tested *Nrg* alleles from three African strains for effects on female mate

preference. The first strain CH11 provided inconclusive results because females from both the

heterozygote and hemizygote genotypes showed strong preference for Z53 males and had

confidence intervals that were greater than 0 (Supplemental Figure 2; Supplemental Table 5).

The dominant female preference from this strain precluded our ability to test the effect of the

CH11 *Nrg* allele specifically.

The remaining two African strains had a pattern consistent with *Nrg* contributing to

female mate preference (Figure 3B; Table 4). While binomial interaction models did not indicate

a significant interaction affect (Table 3) alternative models suggested increased preference for

the African hemizygotes (A/N-*Nrg*⁻) exclusively as these genotypes were the only ones with 95%

confidence intervals that did not contain 0 (Table 4). Additionally a Walds-test indicated a

significant difference between African hemizygotes (A/N-*Nrg*⁻) and African heterozygotes (A/N)

for both strains (Z53 $\chi^2_2 = 8.5$, *P*= 0.015 ; Z30 $\chi^2_2 = 17.9$, *P*= 0.00013).

Discussion

Identifying genes that contribute to female mate preference is an important first step in

understanding how this important behavior evolves and contributes to the evolution of

reproductive isolation and speciation. In this study, we demonstrate that *shep* and *Nrg* play a role

in female mate preference in some African strains. Our current study is one of very few studies

that has identified loci with direct genetic evidence for female mate preference (Chowdhury et

al., 2020) and importantly in a system where premating isolation is the strongest and most

relevant barrier to reproduction. These African populations of *D. melanogaster* have been a

historically important system in the study of speciation (Wu et al., 1995; Coyne et al., 1999;

Greenberg et al., 2003), and we have now identified, in part, genes that could contribute to

premating isolation in this system. Since the genetic basis of behavior is complex, and we

observed variation in behavior among strains (see below), it will be important to test the roles of

shep and *Nrg* in more strains from different locations within Africa. Regardless, these remain

promising candidates that we can leverage to understand the evolution of behavior. For example,

when this *D. melanogaster* system was initially described, mapping studies were only able to

identify large chromosomal regions on chromosome 3, the largest contributor to premating

isolation, and the X chromosome (Hollocher et al., 1997; Takahashi and Ting, 2004). Combining

population genetics with complementation tests, we identified a gene on chromosome 3, *shep* is

on 3L, and the X chromosome, *Nrg*, that contribute to female preference behavior. Getting gene

scale resolution will allow us to further understand how female mate preference has evolved.

Since both *Nrg* and *shep* are essential for neurodevelopment and female mating behavior

(Carhan et al., 2005; Chen et al., 2014), the identification of their role in female preference in

African strains can potentially hint at sensory pathways that are important for mating decisions

and partial reproductive isolation.

Female preference can evolve when structures/genes that are involved in sensory

perception evolve, when brain centers important for learning/decision making evolve, or when

both evolve in tandem (Stevens, 2013; Schaefer and Ruxton, 2015; Munson et al., 2020). Many

studies have documented the role that female preference plays in reproductive isolation

(Laturney and Moehring 2012), and further demonstrating how sexual selection can shape

evolution of female preference and the genetic architecture of female preference (Xu and Shaw

2021). Both *shep* and *Nrg* regulate neurological development, specifically neural remodeling

during metamorphosis, contributing heavily to the formation of higher learning centers,

specifically those responsible for processing olfactory signals (Carhan et al., 2005; Chen et al.,

2014). Our gene ontology results highlight a role for genes involved in neurogenesis and

behavior that complements other gene ontology analyses that suggest outlier genes are involved

in sensory perception and neurological development.

Previous work in African populations of *D. melanogaster* has suggested a large role for

olfaction in female mate preference (Grillet et al., 2012; Moran, 2006; Jin et al., 2022) and it is

possible that *Nrg* and *shep* may contribute to these observed patterns. *Nrg* controls, in part, the

development of the mushroom body (Goosens et al 2011;) which is a structure in *Drosophila*

that processes olfactory information (Li et al 2020). It is possible that differences in *Nrg* across

populations could result in differences in the mushroom body, either at the gross anatomical

level, or at the circuit level and how it receives input from olfactory receptors (Akalal et al.

2006). In closely related species of *Drosophila* differences in olfactory preference can be

explained by the connections made between olfactory receptors and the mushroom body (Ellis et

al. 2023). *shep* is an RNA/DNA binding gene that regulates alternative splicing (Chen et al.

2018; Olesnicky et al. 2018). Given the evidence for changes in *shep* expression in African and

non-African strains under different mating conditions (Bailey et al. 2011) *shep* might regulate

targets in specific neurons that contribute to mating behaviors . It has been demonstrated that

shep has specific targets during neuronal remodeling (Chen et al 2018) and might interact with

this or other targets in the adult brain (Olesniky 2018). Polymorphisms in the coding regions of

either gene could alter interactions with other genes/targets during neuronal development.

Changes in regulatory regions and/or introns could change expression patterns. More work is

needed to establish functional differences between African and non-African alleles. Specifically

whether functional changes in the neural networks of females are correlated with the olfactory

cues used in mate preference. Regardless, while we observed large effects of these single loci,

the genetic basis of female preference is likely complex, with additional loci contributing to

isolation in these strains.

One interesting observation that could suggest that more than these two loci contribute to

female preference is the variation in dominance that we observed for female mating preference in

hybrids between non-African and African strains. In our experiment, we used three African

strains that show strong preference for African males and never choose non-African males in

choice tests. We observed that female mate preference was recessive in F1 hybrids for two of the

strains and dominant in F1 hybrids the third strain when crossed to the same non-African strain

(see Results). The dominance in the CH11 strain precluded us from testing the effects of *Nrg* on

female preference for that strain. When assessing how important *shep* and *Nrg* are in

reproductive isolation across strains different mutations (i.e. CRISPR knockouts) can be used to

circumvent this issue. Nevertheless, testing both the effects of *shep* and *Nrg* and the dominance

of female behavior in more strains will be informative in and of itself and could highlight how

general this phenomenon is. This will be important for two reasons. First, if more strains show

effects of *shep* and *Nrg* on female preference this could suggest these genes are critical for

reproductive isolation in the system (see below). However, it should be cautioned that at early

stages of divergence we do not necessarily expect fixed differences between populations (Cutter,

2012; Castillo and Barbash, 2017). This contrasts with reproductive isolation between species

where we would assume genetic differences are fixed (Laturney and Moehring 2012). Second, in

terms of dominance, looking for general patterns will be important because the dominance of

reproductive barriers can have strong impacts on the outcome of speciation (Thompson et al.,

2021) and might indicate how particular mating barriers evolve.

Differences in dominance for female mate preference could reflect differences in the

selective environment caused by local mating dynamics. This scenario would be analogous to

parallel vs non-parallel evolution of ecological traits in different environments (Oke et al., 2017;

Bolnick et al., 2018). If different populations experience different selective pressures, that is

there is variation in male courtship behavior and female preference across populations, then

different alleles of *shep* and *Nrg* could segregate in these populations. African lineages are quite

diverse in terms of courtship behaviors (Colgrave et al 2000; Yukilevich and True 2008; Jin et

al. 2022) and while strong premating isolation is found in many populations it is likely that

female preferences and associated alleles could be segregating within Africa (Yukilevich and

True, 2008; Coughlan et al 2021) Another possibility, that is not mutually exclusive, is that

different genes contribute to female mate preference in different populations. This would

generate a pattern where genes show effects on behavior for some hybrid genotypes, but not

others. Given the limited data on the genetic basis of female preference it is difficult to infer the

likelihood of this process, but polymorphic incompatibilities are common in other systems

(Cutter, 2012; Castillo and Barbash, 2017).

One example in *Drosophila* suggests that the genetic architecture of premating isolation

can be variable. When different strains of *D. simulans* were crossed and tested for mate rejection

of *D. melanogaster,* different numbers, locations, and dominance of quantitative trait loci were

identified that were strain specific (Uenoyama and Inoue, 1995; Carracedo et al., 1998a;

Carracedo et al., 1998b; Carracedo et al., 2000). The variation in dominance and the number of

loci contributing to mate preference in this *D. simulans* example might be consistent with this

barrier evolving after the speciation event that separated these lineages (Laturney and Moehring,

2012). Moving forward it will be important to determine how common it is for variable loci to

contribute to premating isolation or uniform genetic architecture for reproductive isolation

between populations. Either way, the difference in dominance suggests there may be standing

variation for female mate preference that selection can act on, which is known to facilitate the

evolution of premating isolation through sexual selection (Mendelson et al., 2014; Castillo and

Delph, 2016).

530   Overall, our results provide evidence for two neurodevelopmental genes, *Nrg* and *shep,*

contributing to female mate preference in African strains, suggesting that these genes could

contribute to mating preferences between populations of *D. melanogaster*. Our results connect

population genetics from a behavioral cline, to allelic diversity in African and non-African

populations, to functional validation of the affect of these genes on female African preference

behavior. The differences in alleles for both loci in non-African vs African populations should be

further explored and the testing of more strains are needed to determine whether these genes

could contribute to reproductive isolation. Understanding the functional consequences of

different alleles on neural circuits and sensory perception for specific alleles could provide

insight into how female preference evolved in this system, and provide a framework for

exploring these type of data in other systems. Ultimately, this could lead to understand common

patterns for the evolution of female mate preference and the genetic and neural level.

Literature Cited

Ai, C.R., Norton, E.C., 2003. Interaction terms in logit and probit models. Economic Letters 80,
123-129.
Akalal, D.B., Wilson, C.F., Zong, L., Tanaka, N.K., Ito, K., Davis, R.L. 2006. Roles for
Drosophila mushroom body neurons in olfactory learning and memory. 13:659-68.
Andersson, M., 2019. Sexual selection. Princeton University Press.
Bailey, R.R., Innocenti, P., Morrow, E.H., Friberg, U., Qvarnström, A. 2011. Female *Drosophila*
melanogaster gene expression and mate choice: the X chromosome harbours candidate
genes underlying sexual isolation. Plos One 6: e17358.
Bay, R.A., Arnegard, M.E., Conte, G.L., Best, J., Bedford, N.L., McCann, S.R., Dubin, M.E.,
Chan, Y.F., Jones, F.C., Kingsley, D.M., Schluter, D., Peichel, C.L., 2017. Genetic
Coupling of Female Mate Choice with Polygenic Ecological Divergence Facilitates
Stickleback Speciation. Current Biology 27, 3344-3349.e4.
Bergland, A.O., Tobler, R., González, J., Schmidt, P., Petrov, D., 2016. Secondary contact and
local adaptation contribute to genome□wide patterns of clinal variation in *Drosophila*
melanogaster. Mol Ecol 25, 1157–1174.
Bieber, A.J., Snow, P.M., Hortsch, M., Patel, N.H., Jacobse, J.R., Traquina, Z.R., Schilling, J.,
Goodman C.S., 1989. *Drosophila* neuroglian: a member of the immunoglobulin
superfamily with extensive homology to the vertebrate neural adhesion molecule L1. Cell
59, 447-460.
Blankers, T., Berdan, E.L., Hennig, R.M., Mayer, F., 2019. Physical linkage and mate preference
generate linkage disequilibrium for behavioral isolation in two parapatric crickets.
Evolution 73, 777–791.
Boake, C.R., Arnold, S.J., Breden, F., Meffert, L.M., Ritchie, M.G., Taylor, B.J., Wolf, J.B.,
Moore, A.J., 2002. Genetic tools for studying adaptation and the evolution of behavior.
the american naturalist 160, S143–S159.
Bolnick, D.I., Barrett, R.D.H., Oke, K.B., Rennison, D.J., Stuart, Y.E., 2018. (Non)Parallel
Evolution. Annu. Rev. Ecol. Evol. Syst. 49, 303–330.
Carhan, A., Allen, F., Armstrong, J.D., Goodwin, S.F., O'Dell, K.M.C., 2005. Female receptivity
phenotype of icebox mutants caused by a mutation in the L1-type cell adhesion molecule
neuroglian. Genes Brain Behav 4, 449–465.
Carracedo, M., Asenjo, A., Casares, P., 1998a. Brief communication. Inheritance mode of
Drosophila simulans female mating propensity with *D. melanogaster* males. Journal of
Heredity 89, 101–104.
Carracedo, M.C., Suarez, A., Asenjo, A., Casares, P., 1998b. Genetics of hybridization between
Drosophila simulans females and *D. melanogaster males*. Heredity 80, 17–24.
Carracedo, M.C., Suarez, C., Casares, P., 2000. Sexual isolation between *Drosphila*
melanogaster, *D. simulans* and *D. mauritiana*: sex and species specific discrimination.
Genetica 108, 155–162.
Castillo, D.M., Barbash, D.A., 2017. Moving Speciation Genetics Forward: Modern Techniques
Build on Foundational Studies in *Drosophila*. Genetics 207, 825–842.
Castillo, D.M., Delph, L.F., 2016. Male–female genotype interactions maintain variation in traits
important for sexual interactions and reproductive isolation. Evolution 70, 1667–1673.
Chen, D., Qu, C., Bjorum, S.M., Beckingham, K.M., Hewes, R.S., 2014. Neuronal Remodeling
During Metamorphosis Is Regulated by the *alan shepard* (*shep*) Gene in *Drosophila*
melanogaster. Genetics 197, 1267–1283.
Chen, D., Dale, R.K., Lei, E.P. 2018. *shep* regulates Drosophila neuronal remodeling by

controlling transcription of its chromatin targets. Development 145: dev154047.

Chen, E.Y., Tan, C.M., Kou, Y., Duan, Q., Wang, Z., Meirelles G.V., Clark, N.R., Ma'ayan A. 2013. Enrichr: interactive and collaborative HTML gene list enrichment analysis tool. BMC Bioinformatics, 128.

Chen, J.J., 2003. Communication complex information: the interpretation of statistical interaction in multiple logistic regression analysis. American Journal of Public Health 93, 1376-1377.

Chenoweth, S.F., Blows, M.W., 2006. Dissecting the complex genetic basis of mate choice. Nature Reviews Genetics 7, 681–692.

Chowdhury, T., Calhoun, R.M., Bruch, K., Moehring, A.J., 2020. The *fruitless* gene affects female receptivity and species isolation. Proc. R. Soc. B. 287, 20192765.

Colegrave, N., Hollocher, H., Hinton, K., Ritchie, M.G. 2000. The courtship song of African *Drosophila melanogaster*. J. Evol. Biol 13:143-150.

Comeault, A.A., Serrato-Capuchina, A., Turissini, D.A., McLaughlin, P.J., David, J.R., Matute, D.R., 2017. A nonrandom subset of olfactory genes is associated with host preference in the fruit fly *Drosophila orena*: Behavioral Specialization In *D. Orena*. Evolution Letters 1, 73–85.

Corbett-Detig, R.B., Hartl, D.L. 2012 Population genomics of inversion polymorphisms in *Drosophila melanogaster*. PLoS Genet. 12:e1003056.

Coughlan, J.M., Dagilis, A.J., Serrato-Capuchina, A., Elias, H., Peede, D., Isbell, K., Castillo, D.M., Cooper, B.S., Matute, D.R., 2021. Patterns of and processes shaping population structure and introgression among recently differentiated *Drosophila melanogaster* populations. bioRxiv 2021.06.25.449842.

Coyne, J.A., Orr, H.A., 2004. Speciation. Sinauer associates Sunderland, MA.

Coyne, J.A., Orr, H.A., 1989. Patterns of speciation in *Drosophila*. Evolution 43, 362–381.

Coyne, J.A., Wicker-Thomas, C., Jallon, J.-M., 1999. A gene responsible for a cuticular hydrocarbon polymorphism in *Drosophila melanogaster*. Genet. Res. 73, 189–203.

Cutter, A.D., 2012. The polymorphic prelude to Bateson–Dobzhansky–Muller incompatibilities. Trends in Ecology & Evolution 27, 209–218.

Ellis, K.E., Bervoets, S., Smihula, H., Ganguly, I., Vigato, E., Auer, To.O., Benton, R., Litwin-Kumar, A., Caron, S.J.C. 2023. Evoltuion of connectivity architechture in the Drosophila mushroom body. bioRxiv 2023.02.10.528036.

Enneking, E.M., Kudumala, S.R., Moreno, E., Stephan, R., Boerner, J., Godenshwege, T.A., Pielage, J., 2013. Transsynaptic coordination of synaptic growth, function, and stability by the L1-type CAM *Neuroglian*. Plos Biology 11, e1001537.

Fisher, R.A., 1930. The genetical theory of natural selection. Clarendon Press

Gray, D.A., 2022. Sexual selection and 'species recognition' revisited: serial processing and order-of-operations in mate choice. Proc. R. Soc. B. 289, 20212687.

Greenberg, A.J., Moran, J.R., Coyne, J.A., Wu, C.-I., 2003. Ecological Adaptation During Incipient Speciation Revealed by Precise Gene Replacement. Science 302, 1754–1757.

Grillet, M., Everaerts, C., Houot, B., Ritchie, M.G., Cobb, M., Ferveur, J.-F., 2012. Incipient speciation in *Drosophila melanogaster* involves chemical signals. Sci Rep 2, 224.

Goossens, T., Kang, Y.Y., Wuytens, G., Zimmermann, P., Callaerts-Végh, Z., Pollarolo, G., Islam, R., Hortsch, M., Callaerts, P. 2011. The Drosophila L1CAM homolog *Neuroglian* signals through distinct pathways to control different aspects of mushroom body axon development. Development 138:1595-605.

Harrison, R.G., Larson, E.L., 2014. Hybridization, Introgression, and the Nature of Species
Boundaries. Journal of Heredity 105, 795–809.
Hervas, S., Sanz, E., Casillas, S., Pool, J.E., Barbadilla, A. 2017. PopFly: the Drosophila
population genomics browser. Bioinformatics 33: 2779-2780;
Hollocher, H., Ting, C.-T., Wu, M.-L., Wu, C.-I., 1997c. Incipient Speciation by Sexual Isolation
in *Drosophila melanogaster:* Extensive Genetic Divergence Without Reinforcement.
Genetics 147, 1191–1201.
Hosmer, D.W., Lemeshow, S., Sturdivant, R.X., Applied Logistic Regression 3 ed. John Wiley
& Sons, Inc., Hoboken, New Jersey.
Jaccard, J. 2001. Interaction effects in logistic regression (Vol. 135). Sage, Thousand Oaks, CA.
Jin, B., Barbash, D.A., Castillo, D.M., 2022. Divergent selection on behavioural and chemical
traits between reproductively isolated populations of *Drosophila melanogaster*. J of
Evolutionary Biology jeb.14007.
Kao, J.Y., Zubair, A. Salomon, M.P., Nuzhdin, S.V., Campo, D., 2015. Population genomic
analysis uncovers African and European admizture in *Drosophila melanogaster*
populations from the south-eastern United States and Caribbean Islands. Molecular
Ecology 24, 1499-1509.
Kapopoulou, A., Pfeifer, S.P., Jensen, J.D., Laurent, S. 2018. The demographic history of
African *Drosophila melanogaster.* Genome Bio. Evol. 10:2338-2342.
Kerr, C., Ringo, J., Dowse, H., Johnson, E. 1997. Icebox, a recessive X-linked mutation in
Drosophila causing low sexual receptivity. J. Neurogenet. 11: 213--229.
Kopp, M., Servedio, M.R., Mendelson, T.C., Safran, R.J., Rodríguez, R.L., Hauber, M.E.,
Scordato, E.C., Symes, L.B., Balakrishnan, C.N., Zonana, D.M., van Doorn, G.S., 2018.
Mechanisms of Assortative Mating in Speciation with Gene Flow: Connecting Theory
and Empirical Research. The American Naturalist 191, 1–20.
Kuleshov, M.V., Jones, M.R, Rouillard, E.D., Fernandez N.F., Duan Q., Wang Z., Koplev S.,
Jenkins S.L., Jagodnik K.M., Lachmann A., McDermott M.G., Monteiro C.D.,
Gundersen G.W., Ma'ayan A. 2016. Enrichr: a comprehensive gene set enrichment
analysis web server 2016 update. Nucleic Acids Research, gkw377.
Lack, J.B., Lange J.D., Tang A.D., Corbett-Detig R.B., Pool J.E. 2016. A thousand fly genomes:
an expanded Drosophila genome nexus. Mol Biol Evol. 33:3308-3313.
Laturney, M., Moehring, A.J., 2012. The Genetic Basis of Female Mate Preference and Species
Isolation in *Drosophila*. International Journal of Evolutionary Biology 2012, 1–13.
Li, F., Lindsey, J.W., Marin, E C., Otto, N., Dreher, M., Dempsey, G., et al. 2020. The
connectome of the adult Drosophila mushroom body provides insights into function. Elife
9:e62576.
Mackay, T.F.C., 2001. Quantitative trait loci in *Drosophila*. Nat Rev Genet 2, 11–20.
Meisel R. P., Connallon T., 2013. The faster-X effect: integrating theory and data. Trends Genet.
29: 537–544.
Mendelson, T.C., Martin, M.D., Flaxman, S.M., 2014. Mutation-order divergence by sexual
selection: diversification of sexual signals in similar environments as a first step in
speciation. Ecol Lett 17, 1053–1066.
Mendelson, T.C., Shaw, K.L., 2012. The (mis) concept of species recognition. Trends in ecology
& evolution 27, 421–427.
Miller, D.E, Cook, K.R., Hawley, R.S. 2019. The joy of balancers. PLoS Genet 7:e1008421.

Moran, J.R., 2006. The Genetics of Pre-mating Isolation at the Incipient Stage of Speciation in
Drosophila Melanogaster. University of Chicago, Committee on Genetics.
Muller, H.1918. Genetic variability, twin hybrids and constant hybrids, in a case of balanced
lethal factors. Genetics 3: 422 499.
Munson, A.A., Jones, C., Schraft, H., Sih, A., 2020. You're Just My Type: Mate Choice and
Behavioral Types. Trends in Ecology & Evolution 35, 823–833.
Neelon, D.P., Rodríguez, R.L., Höbel, G., 2019. On the architecture of mate choice decisions:
preference functions and choosiness are distinct traits. Proc. R. Soc. B. 286, 20182830.
Oke, K.B., Rolshausen, G., LeBlond, C., Hendry, A.P., 2017. How Parallel Is Parallel Evolution?
A Comparative Analysis in Fishes. The American Naturalist 190, 1–16.
Olesnicky, E.C., Antonacci, S., Popitsch, N., Lybecker, M.C., Titus, M.B., Valadez, R., Derkach,
P.G., Marean, A., Miller, K., Mathai, S.K., Killian, D.J. 2018. *shep* interacts with
posttranscriptional regulators to control dendrite morphogenesis in sensory neurons.
Dev. Biol. 444: 116-128.
Pool, J.E., 2015. The mosaic ancestry of the *Drosophila* genetic reference panel and the D.
melanogaster reference genome reveals a network of epistatic fitness interactions.
Molecular biology and evolution 32, 3236–3251.
Safran, R.J., Scordato, E.S.C., Symes, L.B., Rodríguez, R.L., Mendelson, T.C., 2013.
Contributions of natural and sexual selection to the evolution of premating reproductive
isolation: a research agenda. Trends in Ecology & Evolution 28, 643–650.
Schaefer, H.M., Ruxton, G.D., 2015. Signal Diversity, Sexual Selection, and Speciation. Annu.
Rev. Ecol. Evol. Syst. 46, 573–592.
Serrato☐Capuchina, A., Schwochert, T.D., Zhang, S., Roy, B., Peede, D., Koppelman, C.,
Matute, D.R., 2021. Pure species discriminate against hybrids in the Drosophila
melanogaster species subgroup. Evolution 75, 1753–1774.
Sprengelmeyer, Q.D., Mansourian S., Lange J.D., Matute D.R., Cooper B.S., Jirle E.V.,
Stensmyr M.C., Pool J.E. 2020. Recurrent collection of *Drosophila melanogaster* from
wild African environments and genomic insights into species history. Mol Biol Evol.
2020 37:627-638.
Stevens, M., 2013. Sensory ecology, behaviour, and evolution. Oxford University Press.
Svensson, O., Woodhouse, K., van Oosterhout, C., Smith, A., Turner, G.F., Seehausen, O., 2017.
The genetics of mate preferences in hybrids between two young and sympatric Lake
Victoria cichlid species. Proc. R. Soc. B. 284, 20162332.
Takahashi, A., Ting, C.-T., 2004. Genetic Basis of Sexual Isolation in *Drosophila melanogaster*.
Genetica 120, 273–284.
The Gene Ontology Consortium. 2017. Expansion of the Gene Ontology knowledgebase and
resources. Nuclei Acids Res. 45: D331-D338.
Thompson, K.A., Urquhart-Cronish, M., Whitney, K.D., Rieseberg, L.H., Schluter, D., 2021.
Patterns, Predictors, and Consequences of Dominance in Hybrids. The American
Naturalist 197, E72–E88.
Turissini, D.A., McGirr, J.A., Patel, S.S., David, J.R., Matute, D.R., 2018. The rate of evolution
of postmating-prezygotic reproductive isolation in Drosophila. Molecular Biology and
Evolution 35, 312–334.
Turner, T.L., Stewart, A.D., Fields, A.T., Rice, W.R., Tarone, A.M., 2011. Population-Based
Resequencing of Experimentally Evolved Populations Reveals the Genetic Basis of Body
Size Variation in Drosophila melanogaster. PLoS Genet 7, e1001336.

Uenoyama, T., Inoue, Y., 1995. Genetic studies on premating isolation in *Drosophila simulans*. I. A *D. simulans* line highly crossable with *D. melanogaster*. The Japanese Journal of Genetics 70, 365–371.

Warton, D.I., Lyons, M., Stoklosa, J., Ives, A.R. 2016. Three points to consider when choosing a LM or GLM test for count data. Methods in Ecol and Evol 7:882-890.

Wu, C.-I., Hollocher, H., Begun, D.J., Aquadro, C.F., Xu, Y., Wu, M.-L., 1995. Sexual isolation in *Drosophila melanogaster*: a possible case of incipient speciation. Proceedings of the National academy of sciences 92, 2519–2523.

Xu, M., Shaw, K.L., 2019a. Genetic coupling of signal and preference facilitates sexual isolation during rapid speciation. Proc. R. Soc. B. 286, 20191607.

Xu, M., Shaw, K.L., 2019b. The genetics of mating song evolution underlying rapid speciation: linking quantitative variation to candidate genes for behavioral isolation. Genetics 211:1089-1104.

Xu, M., Shaw, K.L., 2021. Extensive linkage and genetic coupling of song and preference loci underlying rapid speciation in *Laupala* crickets. J. Heredity 112:204-213.

Yamamoto, D., Koganezawa, M., 2013. Genes and circuits of courtship behaviour in *Drosophila* males. Nat Rev Neurosci 14, 681–692.

Yukilevich, R., True, J.R., 2008. African Morphology, Behavior and Phermones Underlie Incipient Sexual Isolation between US and Caribbean *Drosophila melanogaster*. Evolution 62, 2807–2828.

749 **Table 1.** GO enrichment analysis using biological function highlights genes involved in neuronal
750 development and behavior. Included are the top 10 GO terms with their representation in the data
751 set and adjusted p-values after correcting for multiple testing.
752

GO Term	Overlap	Z-score	Adjusted P-value	Genes
axon guidance	27/242	-1.06	1.2E-09	**Nrg**;Ptp61F;jeb;egh;msn;beat-Vc;tup;tok;robo2;CadN2;Wnt4;grn;Trim9;side;ko;SCAR;beat-Ia;NT1;beat-IIa;trol;sli;CadN;RhoGEF64C;beat-Ib;Lar;cher;mud
axonogenesis	24/212	-1.08	8.2E-09	spri;**Nrg**;Trim9;Ptp61F;jeb;SCAR;egh;beat-Ia;msn;NT1;sick;tup;sli;tok;Gclc;robo2;fz;Sh;CadN;dnc;RhoGEF64C;Lar;mud;tai
motor neuron axon guidance	15/75	-1.56	1.1E-08	**Nrg**;grn;side;ko;beat-Ia;NT1;beat-IIa;beat-Vc;trol;tup;tok;beat-Ib;Lar;cher;Wnt4
cell-cell adhesion	13/64	-1.80	1.4E-07	spri;**Nrg**;fz;Sh;SCAR;CadN;CadN2;dnc;Lar;tai
axon extension	10/36	-1.96	4.6E-07	Con;Cad88C;klg;beat-Ia;beat-IIa;beat-Vc;kirre;rst;fz;CadN;CadN2;beat-Ib;CG34353
courtship behavior	15/109	-1.17	1.3E-06	**Nrg**;cac;orb2;egh;fru;Gr58b;Gr77a;eloF;5-HT7;Sh;**shep**;TfAP-2;dnc;per;dlg1
homophillic cell adhesion	8/28	-2.45	1.1E-05	kirre;rst;Con;fz;CadN;klg;CadN2;CG34353
wing morphogenesis	20/240	-1.13	2.0E-05	disco-r;Spn88Ea;d;bi;fru;dpy;inv;Lpt;bs;tok;Gclc;Fkbp14;fz;EcR;Raf;cv-2;sfl;GEFmeso;hth;Dys
neuron projection extension	8/31	-2.37	2.0E-05	spri;**Nrg**;fz;Sh;CadN;dnc;Lar;tai
anterograde trans-synaptic signaling	12/89	-1.45	3.3E-05	Cep89;Shab;CDase;Dop1R1;5-HT7;cac;X11Lbeta;dnc;nAChRalpha5;Sap47;Rdl;dlg1

753
754 **Table 2.** The summary of population genetic statistics between the Raleigh (RAL) population,
755 representing non-African populations, and Zambia (ZI), representing African populations, for
756 *shep and Nrg*. Average values are provided for reference and come from Lack et al. (2016). The
757 mean across windows is provided with the (min, max) in parentheses.
758

Gene	RAL pi	ZI pi	Fst
shep	0.0078 (0.0011,0.0202)	0.0115 (0.0030,0.0253)	0.1575 (0.0306, 0.4478)
Nrg	0.0047 (0.0002,0.0140)	0.0104 (0.0037,0.0262)	0.2728 (0.0911,0.5788)
Averages	**RAL average pi**	**ZI average pi**	**Average Fst**
	0.00569	**0.00843**	**0.187**

760 **Table 3.** The analysis of female mating behavior using binomial regression containing
761 interaction terms for both *shep* and *Nrg*. The African strain used in the cross is given next to the
762 name of each locus. Coefficients represents the log-odds ratio for the specific effect/term. A
763 discussion of the difference in these models and models without interaction terms is found in the
764 Supplemental Information.
765

shep – **Z53**		
Effect	Coefficient	*P-value*
μ - baseline	0.1001	0.752
β_1 – deficiency	-0.4748	0.259
β_2 – strain	0.4520	0.291
β_3 - interaction	0.9905	0.109
Nrg – **Z53**		
Effect	β coefficient	*P-value*
μ - baseline	-0.3483	0.356
β_1 – deficiency	0.1858	0.711
β_2 – strain	0.2877	0.575
β_3 - interaction	0.9395	0.187
Nrg – **Z30**		
Effect	β coefficient	*P-value*
μ - baseline	-0.3483	0.356
β_1 – deficiency	0.185	0.711
β_2 – strain	0.936	0.086
β_3 - interaction	0.864	0.255

766
767

768   **Table 4.** The behavioral differences in female mate preference for both *shep* and *Nrg* using
769   alternative models that compare each genotype to the random mating expectation. The
770   coefficients represent log odds ratios, estimated from logistic regression, describing the increase
771   or decrease in a genotypes propensity to mate with Z53 males. For this analysis we estimated
772   95% confidence interals, and those that do not contain 0 are in bold.
773

shep		
Strain-Genotype	Lower 95% CI	Upper 95% CI
DGRP882 heterozygote	-0.5222	0.7289
DGRP882 hemizygote	-0.9300	0.1623
Z53 heterozygote	-0.0018	1.1347
Z53 hemizygote	**0.4141**	**1.7988**
Nrg		
Strain-Genotype	Lower 95% CI	Upper 95% CI
DGRP882 heterozygote	-1.1117	0.3833
DGRP882 hemizygote	-0.8204	0.4840
Z30 heterozygote	-0.1662	1.3996
Z30 hemizygote	**0.8901**	**2.5349**
Z53 heterozygote	-0.7523	0.6263
Z53 hemizygote	**0.3800**	**1.8343**

775
776

Figure 1. A). The genes *alan shepard* (*shep*) and *Neuroglian* (*Nrg*) are genomic outliers with a
stronger relationship in African allele frequency and latitude in an analysis using genomes from
populations collected in the Southeastern United States and the Bahamas (data previously
collected in Kao et al 2015). The dashed line corresponds to the value for *shep* and the solid line
corresponds to the value for *Nrg*. The change in allele frequency over latitude for B) *Nrg* and C)
shep. The regression coefficient and adjusted p-value are provided for each gene.

Figure 2. Population differentiation for both A) *shep* and B) *Nrg* are seen in principal components for allelic variation comparing populations from Raleigh NC (RAL, red) and Zimbabwe (ZI, blue).

A

B

Figure 3. A) *alan shepard* (*shep*) and B) *Neuroglian* (*Nrg*) contribute to African female mate
preference. Female flies that are hemizygous and carry only the African allele of *shep* and *Nrg*
show increased preference for African males compared to heterozygote genotypes. This increase
is in the direction of the wild-type African females. * represents a significant difference of the
hemizygote genotype from the corresponding heterozygote genotype using a Walds test.
Confidence intervals for each genotype are Wilson rank intervals estimated from the data.