The nerodevelopmental genes *alan shepard* and *Neuroglian* contribute to female mate preference

in African *Drosophila melanogaster*

Paula R. Roy *[1] and Dean M. Castillo *[2]

* School of Biological Sciences, University of Utah, Salt Lake City UT

Current affiliation: Department of Neurobiology, University of Utah, Salt Lake City UT

Corresponding Author, Current affiliation: Institute of Agriculture and Natural Resources, University of Nebraska-Lincoln, Lincoln NE, dmc79@cornell.edu

Abstract:

Mate choice is a key trait that determines fitness for most sexually reproducing organisms, with

females often being the choosy sex. Female preference often results in strong selection on male

traits that can drive rapid divergence of traits and preferences between lineages, leading to

reproductive isolation. Despite this fundamental property of female mate choice, very few loci

have been identified that contribute to mate choice and reproductive isolation. We used a

combination of population genetics, quantitative complementation tests, and behavioral assays to

demonstrate that *alan shepard* and *Neuroglian* contribute to female mate choice, and could

contribute to partial reproductive isolation between populations of *Drosophila melanogaster*.

Our study is among the first to identify genes that contribute to female mate preference in this

historically important system, where female preference is an active premating barrier to

reproduction. The identification of loci that are primarily known for their roles in

neurodevelopment provides intriguing questions of how female mate preference evolves in

populations via changes in sensory system and higher learning brain centers.

Introduction

For sexually reproducing organisms mate choice is critical to balance maximizing fitness with avoiding costly heterospecific mating events (Mendelson and Shaw, 2012; Gray, 2022). Strong selection on male courtship traits and female preferences can drive rapid divergence between closely related lineages (Fisher, 1930; Andersson, 2019), providing a link between sexual selection and the evolution of reproductive isolation. A large body of work has focused on documenting and describing female preference and identifying genes controlling male courtship traits (Chenoweth and Blows, 2006; Yamamoto and Koganezawa, 2013; Neelon et al., 2019; Munson et al., 2020). However, there have been fewer studies that have identified specific genes that contribute to female mate preference, with most instead focusing on quantitative trait loci (Bay et al., 2017; Svensson et al., 2017; Blankers et al., 2019; Xu and Shaw, 2019; Chowdhury et al., 2020) and some identifying specific genes (Chowdhury et al., 2020). Identifying and characterizing loci contributing to female preference is a critical first step to understand how courtship signal detection, signal processing, and higher brain functions (i.e., learning and memory) contribute to sexual selection and female mate preference evolution.

One challenge in identifying the genetic basis of female choice lies in the complexity of preference-based behaviors (Andersson, 2019). During mate choice females are receiving many, potentially multimodal, signals that then require processing before a decision to accept or reject a mate is made (Gray, 2022; Munson et al., 2020). The complexity of female preference implies that this trait is polygenic, but it is unclear how many loci we would expect to contribute to this trait. Specifically, it remains unclear whether multiple loci interact in an emergent fashion, contributing to female preference as a whole, or if individual loci contribute to a specific facet (perception of a specific signal/cue) of female preference (Boake et al., 2002; Chenoweth and

Blows, 2006). Additionally, while premating isolation evolves rapidly and is often thought to

evolve before other barriers to reproduction (Coyne and Orr, 1989; Turissini et al., 2018), it can

be difficult to determine if the behaviors that currently act as a mating barrier contributed to the

speciation event or evolved after species divergence (Coyne and Orr, 2004; Safran et al., 2013;

Kopp et al., 2018). Focusing on populations that vary in the degree of mating isolation can

overcome some of these challenges and can be used as a model to understand the early stages of

divergence and identify the loci that contribute to speciation and reproductive isolation.

Drosophila melanogaster originated in Southern Africa but is now a cosmopolitan

species with a worldwide distribution. This broad range has resulted in structured and cryptic

populations, even within Africa (Coughlan et al 2021), and strong premating isolation between

some specific populations (Wu et al., 1995). For example, females from strains collected in

Southern Africa strongly reject males from strains that are collected from non-African localities

in both choice and no-choice mating experiments (Wu et al., 1995; Hollocher et al., 1997;

Coughlan et al., 2021). These strong premating isolation behaviors are also observed in other

parts of the *D. melanogaster* range including the Southeast United States and the Caribbean

(Yukilevich and True, 2008). These mating preferences and behaviors that occur outside of

Africa are potentially a product of African ancestry in these populations and admixture with non-

African populations (Kao et al 2015; Bergland et al 2016). Combined, these observations of

strong and variable female preference and population structure provide an ideal system to

explore the evolution of female mate preference.

In this study, we leverage these *D. melanogaster* populations by combining population

genomics and behavioral assays to identify genes that contribute to female mate preference and

partial reproductive isolation. We focus on two loci, *alan Shephard* (*shep*) and *Neuroglian* (*Nrg*)

and demonstrate a role for both genes in female mate preference. First, we use previously

published genomic data from Kao et al. (2015) to demonstrate that these loci show a clinal

pattern of allele frequency in an admixed secondary contact zone, suggesting a role in

reproductive isolation (Harrison and Larson, 2014). We then provide genetic evidence that these

two loci contribute to female mate preference using quantitative complementation tests. Our

study is one of a small number to identify genes important for female mate preference, and in

this case it involves populations where premating isolation is an active barrier to gene flow.

Methods

Genomic cline analysis

To understand how alleles potentially involved in reproductive isolation are structured we

compared allele frequencies across the Southeastern United States and Caribbean using

previously published data (Kao et al 2015). We focused on these populations because this clinal

structure allowed us to make explicit expectations for a gene that contributes to reproductive

isolation compared to the rest of the genome (Harrison and Larson 2014). Specifically, because

previous studies have demonstrated a correlation between latitude and African ancestry, with

more African ancestry in southern populations, we predict that genes involved in behavioral

isolation will have a steeper slope (i.e., larger regression coefficient) compared to the rest of the

genome. Using the genotype matrix (genotype calls for all variable positions they identified)

from Kao et al. (2015), we fit linear regression models between SNP allele frequency and

latitude for the populations. After calculating regression coefficients for each SNP, we found that

two genes previously identified as important for female behavior (*alan shepard* (*shep*) Chen et

al., 2014 and *Neuroglian* (*Nrg)* Carhan et al., 2005) had steep clines. Previous data for these

genes demonstrated that null mutant females rejected wild-type male courtship and were mating

deficient. We determined how the cline regression coefficients for *Nrg* and *shep* compared to the

genomic distribution and the number of SNPs that were outliers and the SNP locations within

these loci.

Quantitative complementation test

To determine if *Nrg* and *shep* alleles contribute to the female mate preference of African strains,

we used a quantitative complementation test combined with a binary mate choice assay.

Quantitative complementation tests have previously been used for no-choice tests (Chowdhury et

al., 2020) and binary choice tests (Comeault et al., 2017). The basic premise of the quantitative

complementation test is to cross wild type strains with a strain containing a mutant allele of the

gene of interest to isolate the effects of different wild type alleles (Mackay, 2001). This is

achieved by comparing progeny that are hemizygous for the locus of interest to individuals,

ideally siblings, that are heterozygous at that locus. The complication of this test is that in

Drosophila these mutants and deletions/deficiencies are often maintained over a balancer

chromosome that has a marker with a dominant visible phenotype. This causes the heterozygote

individuals, but not hemizygote individuals, to have a phenotype that could affect the trait of

interest and must be taken into account during analysis (Turner et al., 2011).

Using a binary choice test, we introduced a single female of a given genotype to two

males from two representative strains: Z53 representing an African male and DGRP882

representing a non-African. Trios were observed for 60 minutes or until copulation occurred.

Upon copulation of a pair of individuals, the non-copulating male was removed, and the strain

identity was determined (see behavioral assay conditions, below). After the pair was finished

copulating, we also examined this male to verify which strain the female chose. Our *a priori*

expectation was that the females hemizygous for the African allele will prefer African males,

whereas females hemizygous for the non-African allele will mate indiscriminately.

Drosophila strains and crosses

To determine whether *Nrg* and *shep* alleles contribute to female preference behavior

between populations of *D. melanogaster* we needed to identify wild-type strains that show strong

female mate preference and complementary strains that show no preference. Strains from

Southern Africa were once considered to be a single lineage from the ancestral range and were

referred to as Z-type. All other strains that migrated from the ancestral range (non-African and

some Northern African strains) were considered a single cosmopolitan lineage and referred to as

M-type. The M-type behavior contrasted with Z-type behavior in that M-type females often mate

indiscriminately, showing no preference, whereas Z-type females strongly reject M-type males

(Wu et al., 1995). Since this initial description, the demographic and evolutionary relationships

between *D. melanogaster* populations have been shown to be more complex (Coughlan et al.,

2021), and it is unlikely that all Z-type strains are homogenous lineage.

We were able to choose strains based on geographic location and evaluated the female

preference behavior for wild type females prior to using these strains in our crosses. For our

representative African strain, we chose Z53 because its behavior profile is well documented and

has been consistent across studies (Wu et al., 1995; Hollocher et al., 1997; Moran, 2006; Jin et

al., 2022). For our representative non-African strain, we chose DGRP882 which we had

previously used and had expected baseline behavior (Jin et al., 2022). The African admixture has

been estimated for individuals of the DGRP panel (Pool, 2015) and DGRP882 has among the

least amount of African ancestry making it most likely to be differentiated African strains. In

addition to these representative lines, we also chose two additional African strains: Z30 and a

more recently collected strain CH11. Canton-S was used as an additional non-African strain

because it is commonly used in behavioral studies (Tompkins et al., 1982; Lasbleiz et al., 2006;

Ng and Kopp, 2008; Kohlmeier et al., 2021).

To create heterozygote and hemizygote genotypes we used loss of function mutant lines

for both *shep* and *Nrg* that had been previously characterized as null alleles (*shep* Chen et al.,

2014; *Nrg* Bieber et al., 1989; Enniking et al., 2013). We used the *shep^{BG00836}* null mutation

(BDSC #12513), hereafter *shep^{-}*. Homozygous null mutant females for this allele are viable but

have mating deficiencies where they do not accept matings as virgins (Chen et al., 2014). We

balanced this mutation over the TM6B chromosome. This balancer chromosome carries the

dominant visible *Hu* marker. We inbred progeny from this original cross for three generations

before using this strain in our behavioral assays. The mutant line and the line that contributed the

TM6B chromosome were not the same genetic background so residual heterozygosity could be

present in this strain. Any effects of this heterozygosity on the X chromosome and 2nd

chromosome would occur in all of the crosses/treatments that we examined. A *shep* deficiency

line (BDSC #7583), already balanced over the TM6B, was tested in preliminary experiments, but

female mating behavior in control crosses with DGRP882 produced highly variable results.

Additional large deficiencies are available but would have precluded us from testing the role of

shep directly. For this reason, we chose to use the balanced strain we had created. We crossed the

shep/TM6B line to both DGRP882 and Z53 males to generate F1 females that were either

heterozygous (had humoral phenotype) or hemizygous (had wild-type phenotype) for *shep*.

We used the Nrg^{17} allele, which is synonymous with Nrg^2 and hereafter Nrg^-, balanced

over the FM7 chromosome, which carries the *Bar* dominant visible marker (BDSC #5595). The

Nrg^{17}requires balancing because this mutation is homozygous lethal. We crossed the Nrg^-/FM7

line to DGRP882 and Z53 similar to the design for *shep.* For experiments with *Nrg,* we tested

two additional African and one non-African strains (described above) for two reasons. First, we

were not sure how or if the *Bar* locus, which affects eye morphology, would affect female

preference. This is particularly important because the impact of vision for female mating

behavior could vary between strains. We tested the effect of *Bar* explicitly by including Canton S

as an additional strain that should not exhibit strong mating preference. Second, even though our

cline analysis identified *Nrg* as a strong outlier, this locus was only a marginal outlier in and

independent study that identified putative female preference loci (Coughlan *et al.*, 2021). This

could indicate that the effect of *Nrg* differs between strains and populations, or multiple alleles of

Nrg segregate in African populations with only a subset contributing to female preference

phenotypes. Either scenario could result in a situation where there is not a single high frequency

allele contributing to female preference in African strains motivating us to test multiple African

strains for this locus.

Behavioral assay conditions

All stocks and virgins were kept on standard cornmeal molasses media in an incubator on

a 12:12 light cycle held at a constant 20 degrees Celsius and 50% humidity. Virgin males and

females were collected within the first 0-4 hours post eclosion to ensure they had not mated.

Females were phenotyped based on the presence of the dominant visible marker and housed in

groups of 5-10. Males were housed individually because group housing can potentially change

courtship vigor (Dixon et al., 2003). All individuals were allowed to age 7-10 days, before

experiments because this is the timeframe that produces the maximum number of matings for the

African strains (Jin et al 2022). All behavioral assays were performed in a room held at a

constant 20C degrees Celsius and 50% humidity within two hours of the incubator lights turning

on to maximize the number of copulations we could observe.

The strains used in the experiment are phenotypically indistinguishable, so to be able to

tell males apart we placed them on food containing blue food coloring 48 hours prior to the

behavioral assay. This is a non-invasive, robust identification method (Wu et al., 1995;

Hollocher et al., 1997; Serrato-Capuchina et al., 2021). In these studies, all males received food

dye treatment, but in our experiment, we only marked one male per mating with blue food dye.

We found that we could more easily distinguish between a dyed and non-dyed male rather than

distinguishing between males fed two different colors. We tested the effect of this treatment on

female preference and found no effect (Supplementary Information).

We used males from the representative strains Z53 and DGRP882 for all of our binary

choice tests. We collected all genotypes for a given locus in each block of our trials in equal

numbers, but due to differences in viability and the number of females that made choices, our

final number of females that were included differed for each genotype (Table 1), but all

genotypes were represented in each block. While the rearing conditions and behavior rooms

conditions were very well controlled, our behavioral protocol also included completing a set of

DGRP882 females in parallel to our experimental crosses. For each block of trials, 20 DGRP882

females were used as a behavioral control and we consistently had 8-12 (representing 40-60%)

choosing Z males consistent with the random mating for this genotype, allowing us to conclude

that the environment was not affecting our matings and pool replicates over blocks.

Statistical analysis of behavioral data

The analysis typically conducted for quantitative complementation tests compares the

difference in phenotype for heterozygote genotypes to the difference in hemizygous genotypes,

using two strains that maximize phenotype divergence (Pasyukova and Mackay 2000; Mackay

2001). Statistically, this has been accomplished by using a model that includes a strain effect, an

effect for carrying the balancer chromosome (which also captures hemizygosity), and the

interaction of these terms. This model works nicely for quantitative traits, but the interpretation

of interaction effects is not straightforward for non-linear models or models based on logistic

regression (Ai and Norton 2003; Chen 2003). For behavioral studies that have used

complementation tests the preference is to use logistic regression as this best captures the binary

nature of choice data, but not all studies have tested the interaction term explicitly. One study

compared hemizygotes to heterozygotes for an interspecies cross using the wildtype strain of

interest (Comeault et al., 2017), but this framework does not control for either hemizygosity

effects or the effects of dominant mutations carried on balancer chromosomes. This would have

required a within species control cross, whereas the study instead focused on interspecies

hybrids. Another study included multiple strains in their statistical models to evaluate

hemizygosity and balancer effects but still focused on the significance between hemizygote and

heterozygote genotypes that were generated from the interspecific crosses (Chowdhury et al.,

2020). This study modeled the interaction effect, but only reported the significance of this effect,

which is not sufficient to understand the sign of the interaction effect without recalculating the

odds ratios for the comparisons of interest (Ai and Norton 2003; Chen 2003). The interpretation

of interaction effects in logistic regression cannot be made based on this single coefficient

because the interaction is conditional on other effects (Ai and Norton 2003). Since this is not an

issue in statistical models using quantitative traits (i.e., linear regression) authors must be

cautious when applying interaction models to logistic data.

For our analysis, we used a statistical framework that can be readily interpreted and

parallels the intent of the original analysis outlined for quantitative complementation tests

(Pasyukova and Mackay 2000; Mackay 2001), but avoids modelling the interaction effect, and

the difficulty in the interpretation of this coefficient. By comparing each genotype in a set of

crosses to a single baseline genotype, we can test for a strain effect, a balancer effect, and the

effect of the hemizygote of interest. If a strain effect is detected a post-hoc test can be used to

determine if there is an interaction between strain and hemizygosity in a similar fashion of what

would be required to properly interpret a statistical interaction effect for a logistic regression.

We used logistic regression where the response variable was 0 if the female mated with

the DGRP882 male and 1 if they mated with the Z53 male. Female genotype was included as a

categorical variable in the model. We first determined whether any female genotypes (for a given

locus) were significantly different that the DGRP882 heterozygote genotype that we used as

baseline in our model. Comparing the heterozygote genotypes to the baseline allowed us to test

for any strain effects. If a given African strain heterozygote was not significantly different from

baseline, we could infer there was no strain difference. These heterozygotes would have one

common haploid genome, and the other complement coming from the strain used in the cross.

Testing the balancer effect, which also tests for hemizygosity effects, was completed by

comparing the DGRP882 heterozygote that carries a balancer chromosome, to the DGRP882

hemizygote. In the absence of balancer and heterozygote effects we would expect these

genotypes to show the same, random mating preference. For all crosses the DGRP882

hemizygote was never significantly different from the DGRP882 heterozygote, allowing us to

focus on whether any of the African hemizyogtes were significantly different from baseline. If

the African strain hemizygote was significantly different from the baseline, and DGRP882

hemizygote but the African strain heterozygote was not different from baseline we could infer

that this allele contributes to female preference behavior. would fulfill the criteria of the

difference in hemizygotes being greater than the difference in heterozygotes (Mackay, 2001). If

both the specific African strain heterozygote and African strain hemizygote were significantly

different from baseline we proceeded to the second analysis step. This involved computing a

linear contrast for the difference in these genotypes from their representative DGRP882

counterparts. This second step was only necessary for the African strain CH11 using in the *Nrg*

experiment.

Results

Female behavior loci are genomic outliers

To generate candidate genes for functional validation, we leveraged genomic data from

populations of *D. melanogaster* that exhibit a cline in African ancestry (Kao et al., 2015;

Bergland et al., 2016) and African-like female preference behavior (Yukilevich and True, 2008).

Our goal was to identify outlier genes that showed a steeper cline in African ancestry compared

to the genome wide average. We also compared our list of candidates to independent analyses

that used populations from Africa to identify female preference alleles (Coughlan et al., 2021).

We retained the top 1% of outliers for loci that had the steepest clines with African alleles

being more frequent at lower latitudes. Many of these SNPs did not occur in annotated genes, but

two interesting candidates emerged, *alan shepard* (*shep*) and *Neuroglian* (*Nrg*). There was a

single SNP in *shep* that was above the top 1% threshold and this SNP was in an intron of *shep*.

There were three additional SNPs close to the 1% threshold that were also in introns. For *Nrg*

there were two intronic SNPs that were in the top 1% and these SNPs were within 2 kb of each

other. Interestingly these loci were also outliers in the study by Coughlan et al (2021) with shep

in the top 1% of outliers and *Nrg* in the top 5% of outliers. Previous behavioral analysis of

homozygous viable mutant strains for these loci found evidence that these genes were involved

female behavior, where virgin females rejected male courtship to remain unmated (Carhan et al.,

2005; Chen et al., 2014). Taken together, our observation of a strong cline in allele frequency

and previous studies suggesting a role in behavior led us to functionally test these loci for a role

specifically in African female mate preference behavior.

Shep contributes to female mate preference in an African strain

We used quantitative complementation tests to determine if *alan shepard* (*shep*) contributes to

female mate preference. We were able to validate that *shep* contributes to female mate

preference by specifically demonstrating that females that are hemizygous for the Z53 African

allele have significantly different behavior than the other control genotypes. These African

hemizygote females show a significantly higher preference for African males over non-African

males (Fig. 2). The rest of the genotypes were not significantly different from our baseline

genotype, the DGRP882 heterozygote which carried the DGRP882 *shep* allele and the *shep* allele

on the TM6B chromosome (Table 1). The non-African hemizygote, that only carried the

DGRP882 allele of *shep* was not significantly different from the baseline (β=-0.374, *P*=0.1761),

neither was the Z53 heterozygote genotype (β=0.522, *P*=0.0552). The only genotype that was

significantly different was the Z53 hemizygote (β=1.068, *P*=0.0022). This genotype preferred

Z53 males over DGRP882 males in a similar direction, but to a lesser degree, as the wild-type

Z53 parental strain indicating that this allele of *shep* contributes to female mate preference.

Nrg contributes female mate preference in African strains

We also used quantitative complementation tests to validate the role of *Nrg* alleles on female

mate preference in African strains. Our design and logic were identical for what we used to

analyze *shep* and for this gene we included more genotypes to determine if the visible marker,

Bar, on the balancer chromosome was affecting behavior, and whether different *Nrg* alleles

might show different effects on female mate preference. We used the non-African genotype

DGRP882 heterozygote which carries the *Nrg* allele from DGRP882 and the *Nrg* allele from the

FM7 chromosome as our baseline.

To test for the effects of the *Bar* allele specifically on female behavior we tested the

behavior of another non-African strain, Canton S, and compared it to the DGRP882 strain. We

used Canton S because we had a clear expectation: that both female genotypes from a cross

between Canton S and the Nrg^-/FM7 strain should have the same behavior as the DGRP882

genotypes, and they should mate randomly and not show preference for either male genotype.

We in fact observed that these genotypes mated indiscriminately, and we did not detect any

significant difference in behavior comparing DGRP882 genotypes and Canton S genotypes (Fig

2; Table 1).

We then tested *Nrg* alleles from three African strains for effects on female mate

preference. The first strain CH11 provided inconclusive results because females from both the

heterozygote and hemizygote genotypes showed strong preference for Z53 males compared to

the DGRP882 baseline (Fig 2; Table 1). The strength of this preference resulted in both

genotypes being significantly different from the baseline, and indistinguishable from each other.

The dominant female preference from this strain precluded our ability to test the effect of the

CH11 *Nrg* allele specifically.

The remaining two African strains had a pattern consistent with *Nrg* contributing to

female mate preference (Fig 3; Supplemental Table 1). Both the heterozygous female genotypes

for the Z30 and Z53 strain were not significantly different from the baseline (Z30 β=0.936,

P=0.0862; Z53 β=0.288; *P*=0.5752). The hemizygous female genotypes for both strains were

significantly different from the baseline (Z30 β=1.986, *P*=0.0004; Z53 β=1.413; *P*=0.0072)

indicating that when females only have these African strain alleles they exhibit significant

preference for African males similar to the wild-type African parental strains.

Discussion

Identifying genes that contribute to female mate preference is an important first step in

understanding how this important behavior evolves and contributes to the evolution of

reproductive isolation and speciation. In this study, we demonstrate that *shep* and *Nrg* play a role

in female mate preference in some African strains. Our current study is one of very few studies

that has identified loci with direct genetic evidence for female mate preference (Chowdhury et

al., 2020) and importantly in a system where premating isolation is the strongest and most

relevant barrier to reproduction. These African populations of *D. melanogaster* have been a

historically important system in the study of speciation (Wu et al., 1995; Coyne et al., 1999;

Greenberg et al., 2003), and we have now identified, in part, genes that could contribute to

premating isolation in this system. Since the genetic basis of behavior is complex, and we

observed variation in behavior among strains (see below), it will be important to test the roles of

shep and *Nrg* in more strains from different locations within Africa. Regardless, these remain

promising candidates that we can leverage to understand the evolution of behavior. For example,

when this *D. melanogaster* system was initially described, mapping studies were only able to

identify large chromosomal regions on chromosome 3, the largest contributor to premating

isolation, and the X chromosome (Hollocher et al., 1997; Takahashi and Ting, 2004). Combining

population genetics with complementation tests, we identified a gene on chromosome 3, *shep* is

on 3L, and the X chromosome, *Nrg*, that contribute to female preference behavior. Getting gene

scale resolution will allow us to further understand how female mate preference has evolved.

Since both *Nrg* and *shep* are essential for neurodevelopment and female mating behavior

(Carhan et al., 2005; Chen et al., 2014), the identification of their role in female preference in

African strains can potentially hint at sensory pathways that are important for mating decisions

and partial reproductive isolation.

353   Female preference can evolve when structures/genes that are involved in sensory

perception evolve, when brain centers important for learning/decision making evolve, or when

both evolve in tandem (Stevens, 2013; Schaefer and Ruxton, 2015; Munson et al., 2020). Both

shep and *Nrg* regulate neurological development, specifically neural remodeling during

metamorphosis, contributing heavily to the formation of higher learning centers, specifically

those responsible for processing olfactory signals (Carhan et al., 2005; Chen et al., 2014). Our

results complement a gene ontology analysis of outliers for female mate preference by Coughlan

et al. (2021) that found significant outliers involved in sensory perception, including three

olfactory receptors, as well as genes important for learning and memory, cognition, and

neurological development. Both analyses are consistent with previous work in these populations

of *D. melanogaster* has suggested a large role for olfaction in female mate preference (Grillet et

al., 2012; Moran, 2006; Jin et al., 2022). Given these complementary observations it will be

important to understand how variation *Nrg* and *shep* alleles are structured in strains and

populations that show high or low mate discrimination preference. Identifying different alleles

could provide insight into the structural and/or functional changes in the neural networks of

females and whether this is correlated with the olfactory cues used in mate preference.

Regardless, while we observed large effects of these single loci, the genetic basis of female

preference is likely complex, with additional loci contributing to isolation in these strains.

One interesting observation that could suggest that more than these two loci contribute to

female preference is the variation in dominance that we observed for female mating preference in

hybrids between non-African and African strains. In our experiment, we used three African

strains that show strong preference for African males and never choose non-African males in

choice tests. We observed that female mate preference was recessive in F1 hybrids for two of the

strains and dominant in F1 hybrids the third strain when crossed to the same non-African strain

(see Results). The dominance in the CH11 strain precluded us from testing the effects of *Nrg* on

female preference for that strain. When assessing how important *shep* and *Nrg* are in

reproductive isolation across strains different experimental designs can be used to circumvent

this issue. Nevertheless, testing the dominance of female behavior in more strains will be

informative in and of itself and could highlight how general this phenomenon is which will be

important because the dominance of reproductive barriers can have strong impacts on the

outcome of speciation (Thompson et al., 2021) and might indicate how particular mating barriers

evolve.

Differences in dominance for female mate preference could reflect differences in the

selective environment caused by local mating dynamics. This scenario would be analogous to

parallel vs non-parallel evolution of ecological traits in different environments (Oke et al., 2017;

Bolnick et al., 2018). If different populations experience different selective pressures, then

different alleles of *shep* and *Nrg* could segregate in these populations. African lineages are quite

diverse and while strong premating isolation is found in many populations it is likely that female

preferences and associated alleles could be segregating within Africa (Yukilevich and True,

2008; Coughlan et al 2021) Another possibility, that is not mutually exclusive, is that the genetic

architecture of female preference in hybrids is genotype-dependent because different genes

contribute to female mate preference in these different populations. Given the limited data on the

genetic basis of female preference it is difficult to infer the likelihood of this process, but

polymorphic incompatibilities are common in other systems (Cutter, 2012; Castillo and Barbash,

2017).

One example in *Drosophila* suggests that the genetic architecture of premating isolation

can be variable. When different strains of *D. simulans* were crossed and tested for mate rejection

of *D. melanogaster* different numbers, locations, and dominance of quantitative trait loci (QTL)

were identified that were strain specific (Uenoyama and Inoue, 1995; Carracedo et al., 1998a;

Carracedo et al., 1998b; Carracedo et al., 2000). The variation in dominance and the number of

loci contributing to mate preference in this *D. simulans* example might be consistent with this

barrier evolving after the speciation event that separated these lineages (Laturney and Moehring,

2012). Moving forward it will be important to determine how common it is for variable loci to

contribute to premating isolation or uniform genetic architecture for reproductive isolation

between populations. Either way, the difference in dominance suggests there may be standing

variation for female mate preference that selection can act on, which is known to facilitate the

evolution of premating isolation through sexual selection (Mendelson et al., 2014; Castillo and

Delph, 2016).

Overall, our results provide evidence for two neurodevelopmental genes, *Nrg* and *shep,*

contributing to female mate preference in African strains, and could contribute to partial

reproductive isolation between populations of *D. melanogaster*. Our results connect population

genetics from a system that exhibits a cline in behavior and allele frequency, to functional

validation of the effect of these genes on behavior. The differences in alleles for both loci in non-

African vs African populations should be further explored and the testing of more strains are

needed to determine whether these could be considered speciation genes. Understanding allelic

diversity will help develop experiments that can test the functional consequences on neural

circuits and sensory perception for specific alleles. Neurobiological experiments could provide

insight into how female preference evolved in this system and provide a framework for exploring

these types of data in other systems. Ultimately, this could lead to understand common patterns

for the evolution of female mate preference and the genetic and neural level.

Literature Cited

Ai, C.R., Norton, E.C., 2003. Interaction terms in logit and probit models. Economic Letters 80, 123-129.

Andersson, M., 2019. Sexual selection. Princeton University Press.

Bay, R.A., Arnegard, M.E., Conte, G.L., Best, J., Bedford, N.L., McCann, S.R., Dubin, M.E., Chan, Y.F., Jones, F.C., Kingsley, D.M., Schluter, D., Peichel, C.L., 2017. Genetic Coupling of Female Mate Choice with Polygenic Ecological Divergence Facilitates Stickleback Speciation. Current Biology 27, 3344-3349.e4.

Bergland, A.O., Tobler, R., González, J., Schmidt, P., Petrov, D., 2016. Secondary contact and local adaptation contribute to genome-wide patterns of clinal variation in *Drosophila melanogaster*. Mol Ecol 25, 1157–1174.

Bieber, A.J., Snow, P.M., Hortsch, M., Patel, N.H., Jacobse, J.R., Traquina, Z.R., Schilling, J., Goodman C.S., 1989. *Drosophila* neuroglian: a member of the immunoglobulin superfamily with extensive homology to the vertebrate neural adhesion molecule L1. Cell 59, 447-460.

Blankers, T., Berdan, E.L., Hennig, R.M., Mayer, F., 2019. Physical linkage and mate preference
   generate linkage disequilibrium for behavioral isolation in two parapatric crickets.
   Evolution 73, 777–791.

Boake, C.R., Arnold, S.J., Breden, F., Meffert, L.M., Ritchie, M.G., Taylor, B.J., Wolf, J.B.,
   Moore, A.J., 2002. Genetic tools for studying adaptation and the evolution of behavior.
   The American Naturalist 160, S143–S159.

Bolnick, D.I., Barrett, R.D.H., Oke, K.B., Rennison, D.J., Stuart, Y.E., 2018. (Non)Parallel
   Evolution. Annu. Rev. Ecol. Evol. Syst. 49, 303–330.

Carhan, A., Allen, F., Armstrong, J.D., Goodwin, S.F., O'Dell, K.M.C., 2005. Female receptivity
   phenotype of icebox mutants caused by a mutation in the L1-type cell adhesion molecule
   neuroglian. Genes Brain Behav 4, 449–465.

Carracedo, M., Asenjo, A., Casares, P., 1998a. Brief communication. Inheritance mode of
   Drosophila simulans female mating propensity with *D. melanogaster* males. Journal of
   Heredity 89, 101–104.

Carracedo, M.C., Suarez, A., Asenjo, A., Casares, P., 1998b. Genetics of hybridization between
   Drosophila simulans females and *D. melanogaster males*. Heredity 80, 17–24.

Carracedo, M.C., Suarez, C., Casares, P., 2000. Sexual isolation between *Drosphila
   melanogaster*, *D. simulans* and *D. mauritiana*: sex and species-specific discrimination.
   Genetica 108, 155–162.

Castillo, D.M., Barbash, D.A., 2017. Moving Speciation Genetics Forward: Modern Techniques
   Build on Foundational Studies in *Drosophila*. Genetics 207, 825–842.

Castillo, D.M., Delph, L.F., 2016. Male–female genotype interactions maintain variation in traits
   important for sexual interactions and reproductive isolation. Evolution 70, 1667–1673.

Chen, D., Qu, C., Bjorum, S.M., Beckingham, K.M., Hewes, R.S., 2014b. Neuronal Remodeling
   During Metamorphosis Is Regulated by the *alan shepard* (shep) Gene in *Drosophila
   melanogaster*. Genetics 197, 1267–1283.

Chen, J.J., 2003. Communication complex information: the interpretation of statistical
   interaction in multiple logistic regression analysis. American Journal of Public Health 93,
   1376-1377.

Chenoweth, S.F., Blows, M.W., 2006. Dissecting the complex genetic basis of mate choice.
   Nature Reviews Genetics 7, 681–692.

Chowdhury, T., Calhoun, R.M., Bruch, K., Moehring, A.J., 2020. The *fruitless* gene affects
   female receptivity and species isolation. Proc. R. Soc. B. 287, 20192765.

Comeault, A.A., Serrato-Capuchina, A., Turissini, D.A., McLaughlin, P.J., David, J.R., Matute,
   D.R., 2017. A nonrandom subset of olfactory genes is associated with host preference in
   the fruit fly *Drosophila orena*: Behavioral Specialization In *D. Orena*. Evolution Letters
   1, 73–85.

Coughlan, J.M., Dagilis, A.J., Serrato-Capuchina, A., Elias, H., Peede, D., Isbell, K., Castillo,
   D.M., Cooper, B.S., Matute, D.R., 2021. Patterns of and processes shaping population
   structure and introgression among recently differentiated *Drosophila melanogaster*
   populations (preprint). Evolutionary Biology.

Coyne, J.A., Orr, H.A., 2004. Speciation. Sinauer associates Sunderland, MA.

Coyne, J.A., Orr, H.A., 1989. Patterns of speciation in *Drosophila*. Evolution 43, 362–381.

Coyne, J.A., Wicker-Thomas, C., Jallon, J.-M., 1999. A gene responsible for a cuticular
   hydrocarbon polymorphism in *Drosophila melanogaster*. Genet. Res. 73, 189–203.

Cutter, A.D., 2012. The polymorphic prelude to Bateson–Dobzhansky–Muller incompatibilities.
Trends in Ecology & Evolution 27, 209–218.
Dixon, S.M., Coyne, J.A., Noor, M.A.F., 2003. The evolution of conspecific sperm precedence
in Drosophila. Molecular Ecology 12:1179-1184.
Enneking, E.M., Kudumala, S.R., Moreno, E., Stephan, R., Boerner, J., Godenshwege, T.A.,
Pielage, J., 2013. Transsynaptic coordination of synaptic growth, function, and stability
by the L1-type CAM *Neuroglian*. Plos Biology 11, e1001537.
Fisher, R.A., 1930. The genetical theory of natural selection. Clarendon Press, Oxford.
Gray, D.A., 2022. Sexual selection and 'species recognition' revisited: serial processing and
order-of-operations in mate choice. Proc. R. Soc. B. 289, 20212687.
Greenberg, A.J., Moran, J.R., Coyne, J.A., Wu, C.-I., 2003. Ecological Adaptation During
Incipient Speciation Revealed by Precise Gene Replacement. Science 302, 1754–1757.
Grillet, M., Everaerts, C., Houot, B., Ritchie, M.G., Cobb, M., Ferveur, J.-F., 2012. Incipient
speciation in *Drosophila melanogaster* involves chemical signals. Sci Rep 2, 224.
Harrison, R.G., Larson, E.L., 2014. Hybridization, Introgression, and the Nature of Species
Boundaries. Journal of Heredity 105, 795–809.
Hollocher, H., Ting, C.-T., Wu, M.-L., Wu, C.-I., 1997c. Incipient Speciation by Sexual Isolation
in *Drosophila melanogaster:* Extensive Genetic Divergence Without Reinforcement.
Genetics 147, 1191–1201.
Jin, B., Barbash, D.A., Castillo, D.M., 2022. Divergent selection on behavioural and chemical
traits between reproductively isolated populations of *Drosophila melanogaster*. J of
Evolutionary Biology jeb.14007.
Kao, J.Y., Zubair, A. Salomon, M.P., Nuzhdin, S.V., Campo, D., 2015. Population genomic
analysis uncovers African and European admixture in *Drosophila melanogaster*
populations from the south-eastern United States and Caribbean Islands. Molecular
Ecology 24, 1499-1509.
Kohlmeier, P., Zhang, Y., Gorter, J.A., Su, C.Y., Billeter, J.C., 2021. Mating increases
Drosophila melanogaster females' choosiness by reducing olfactory sensitivity to a male
pheromone. Nature Ecology and Evolution 5:1165-1173.
Kopp, M., Servedio, M.R., Mendelson, T.C., Safran, R.J., Rodríguez, R.L., Hauber, M.E.,
Scordato, E.C., Symes, L.B., Balakrishnan, C.N., Zonana, D.M., van Doorn, G.S., 2018.
Mechanisms of Assortative Mating in Speciation with Gene Flow: Connecting Theory
and Empirical Research. The American Naturalist 191, 1–20.
Lasbleiz, C., Ferveur, J.F., Everaerts, C., 2006. Courtship behavior of *Drosophila melanogaster*
revisited. Animal Behavior 72:1001-1012.
Laturney, M., Moehring, A.J., 2012. The Genetic Basis of Female Mate Preference and Species
Isolation in *Drosophila*. International Journal of Evolutionary Biology 2012, 1–13.
Mackay, T.F.C., 2001. Quantitative trait loci in *Drosophila*. Nat Rev Genet 2, 11–20.
Mendelson, T.C., Martin, M.D., Flaxman, S.M., 2014. Mutation-order divergence by sexual
selection: diversification of sexual signals in similar environments as a first step in
speciation. Ecol Lett 17, 1053–1066.
Mendelson, T.C., Shaw, K.L., 2012. The (mis) concept of species recognition. Trends in ecology
& evolution 27, 421–427.
Moran, J.R., 2006. The Genetics of Pre-mating Isolation at the Incipient Stage of Speciation in
Drosophila Melanogaster. University of Chicago, Committee on Genetics.

Munson, A.A., Jones, C., Schraft, H., Sih, A., 2020. You're Just My Type: Mate Choice and Behavioral Types. Trends in Ecology & Evolution 35, 823–833.

Neelon, D.P., Rodríguez, R.L., Höbel, G., 2019. On the architecture of mate choice decisions: preference functions and choosiness are distinct traits. Proc. R. Soc. B. 286, 20182830.

Ng, C.S., Kopp, A., 2008. Se combs are important for male mating success in *Drosophila melanogaster*. Behavior Genetics 38:195-201.

Oke, K.B., Rolshausen, G., LeBlond, C., Hendry, A.P., 2017. How Parallel Is Parallel Evolution? A Comparative Analysis in Fishes. The American Naturalist 190, 1–16.

Pasyukova, E.G., Vieira, C., Mackay, T.F., 2000. Deficiency mapping of quantitative train loci affecting longevity in *Drosophila melanogaster*. Genetics 156:1129-1146.

Pool, J.E., 2015. The mosaic ancestry of the *Drosophila* genetic reference panel and the D. melanogaster reference genome reveals a network of epistatic fitness interactions. Molecular biology and evolution 32, 3236–3251.

Safran, R.J., Scordato, E.S.C., Symes, L.B., Rodríguez, R.L., Mendelson, T.C., 2013. Contributions of natural and sexual selection to the evolution of premating reproductive isolation: a research agenda. Trends in Ecology & Evolution 28, 643–650.

Schaefer, H.M., Ruxton, G.D., 2015. Signal Diversity, Sexual Selection, and Speciation. Annu. Rev. Ecol. Evol. Syst. 46, 573–592.

Serrato-Capuchina, A., Schwochert, T.D., Zhang, S., Roy, B., Peede, D., Koppelman, C., Matute, D.R., 2021. Pure species discriminate against hybrids in the Drosophila melanogaster species subgroup. Evolution 75, 1753–1774.

Stevens, M., 2013. Sensory ecology, behaviour, and evolution. Oxford University Press.

Svensson, O., Woodhouse, K., van Oosterhout, C., Smith, A., Turner, G.F., Seehausen, O., 2017. The genetics of mate preferences in hybrids between two young and sympatric Lake Victoria cichlid species. Proc. R. Soc. B. 284, 20162332.

Takahashi, A., Ting, C.-T., 2004. Genetic Basis of Sexual Isolation in *Drosophila melanogaster*. Genetica 120, 273–284.

Thompson, K.A., Urquhart-Cronish, M., Whitney, K.D., Rieseberg, L.H., Schluter, D., 2021. Patterns, Predictors, and Consequences of Dominance in Hybrids. The American Naturalist 197, E72–E88.

Turissini, D.A., McGirr, J.A., Patel, S.S., David, J.R., Matute, D.R., 2018. The rate of evolution of postmating-prezygotic reproductive isolation in Drosophila. Molecular Biology and Evolution 35, 312–334.

Turner, T.L., Stewart, A.D., Fields, A.T., Rice, W.R., Tarone, A.M., 2011. Population-Based Resequencing of Experimentally Evolved Populations Reveals the Genetic Basis of Body Size Variation in Drosophila melanogaster. PLoS Genet 7, e1001336.

Uenoyama, T., Inoue, Y., 1995. Genetic studies on premating isolation in *Drosophila simulans*. I. A *D. simulans* line highly crossable with *D. melanogaster*. The Japanese Journal of Genetics 70, 365–371.

Wu, C.-I., Hollocher, H., Begun, D.J., Aquadro, C.F., Xu, Y., Wu, M.-L., 1995. Sexual isolation in *Drosophila melanogaster*: a possible case of incipient speciation. Proceedings of the National academy of sciences 92, 2519–2523.

Xu, M., Shaw, K.L., 2019. Genetic coupling of signal and preference facilitates sexual isolation during rapid speciation. Proc. R. Soc. B. 286, 20191607.

Yamamoto, D., Koganezawa, M., 2013. Genes and circuits of courtship behaviour in *Drosophila* males. Nat Rev Neurosci 14, 681–692.

575   Yukilevich, R., True, J.R., 2008. African Morphology, Behavior and Phermones Underlie
576   Incipient Sexual Isolation between US and Caribbean *Drosophila melanogaster*.
577   Evolution 62, 2807–2828.
578

Table 1. The behavioral differences in female mate preference from the baseline genotype for
both *shep* and *Nrg*. The baseline genotype in each case was the DGRP882 heterozygote which
carried the DGRP882 allele for the specific locus and the allele from the corresponding balancer
chromosome. The NA represents that this genotype was used as the baseline. The remaining
coefficients represent log odds ratios, estimated from logistic regression, describing the increase
or decrease in a genotypes propensity to mate with Z53 males. The significance of each
coefficient was tested independently against the baseline and ** represents *P*<0.001.

shep				
Strain-Genotype	Females tested	Proportion choose	β coefficient	*P-value*
DGRP882 homozygote	40	0.48	NA	NA
Z53 heterozygote	53	0.64	0.552	0.0552
DGRP882 hemizygote	54	0.41	-0.3747	0.1761
Z53 hemizygote	43	0.74	1.068	0.0022**
Nrg				
Strain-Genotype	Females tested	Proportion choose Z	β coefficient	*P-value*
DGRP882 homozygote	29	0.58	NA	NA
Canton S homozygote	24	0.54	0.515	0.35464
Z30 heterozygote	28	0.64	0.936	0.0862
Z53 heterozygote	33	0.48	0.288	0.5752
CH11 heterozygote	22	0.82	1.852	0.0056**
DGRP882 hemizygote	37	0.46	0.186	0.7107
Canton S hemizygote	29	0.55	0.556	0.2947
Z30 hemizygote	43	0.84	1.986	0.0004**
Z53 hemizygote	39	0.74	1.413	0.0072**
CH 11 hemizygote	30	0.80	1.735	0.0034**

588
589

Figure 1. The genes *alan shepard* (*shep*) and *Neuroglian* (*Nrg*) are genomic outliers with a

stronger relationship in African allele frequency and latitude in an analysis using genomes from

populations collected in the Southeastern United States and the Bahamas (data previously

collected in Kao et al 2015). The dashed line corresponds to the value for *shep* and the solid line

corresponds to the value for *Nrg*. These regressions are for individual SNPs within these loci that

are in the top 1% for slope in this analysis.

Fig 2. *alan shepard* (*shep*) contributes to African female mate preference. Female flies that are

hemizygous and carry only the African allele of *shep* show increased preference for African

males, similar to wild-type African females. The horizontal dashed lines are the 95% confidence

interval for the baseline genotype, DGRP882 heterozygote. Each remaining genotype was

compared to the baseline in a logistic regression where * represents a significant difference from

the baseline at *P*<0.05. Confidence intervals are Wilson rank intervals from the binomial

distribution.

Fig 3. The dominant visible marker, *Bar*, on FM7 does not affect female mate preference for

either non-African or African strains. Canton S was included as a control because we expected it

to behave similarly to the DGRP882 strain. The CH11 strain was included to test multiple

African *Nrg* alleles. Because this strain exhibited a dominant female preference phenotype for

both heterozygous and hemizygous females these genotypes were indistinguishable. The

horizontal dashed lines are the 95% confidence interval for the baseline DGRP882

heterozygotes. Each remaining genotype was compared to this baseline in a logistic regression

where * represents a significant difference from the baseline at *P*<0.05. Confidence intervals are

Wilson rank intervals from the binomial distribution.

Fig 4. *Neruoglian* (*Nrg*) contributes to African female mate preference. Female flies that are

hemizygous and carry only the African allele of *Nrg* show increased preference for African

males, similar to wild-type African females. The horizontal dashed lines are the 95% confidence

interval for the baseline DGRP882 heterozygote. Each remaining genotype was compared to this

baseline with a logistic regression where * represents a significant difference from the baseline at

P<0.05. Confidence intervals are Wilson rank intervals from the binomial distribution.

626