The lncRNA *Firre* functions as a transcriptional activator from a distance

Christian Much[1], Erika L. Lasda[1], Jordan P. Lewandowski[2], Michael J. Smallegan[1,3*], John L. Rinn[1,4*]

[1]BioFrontiers Institute, University of Colorado Boulder, Boulder, CO 80303, USA

[2]Department of Stem Cell and Regenerative Biology, Harvard University, Cambridge, MA 02138, USA

[3]Department of Molecular, Cellular and Developmental Biology, University of Colorado Boulder, Boulder, CO 80302, USA

[4]Department of Biochemistry, University of Colorado Boulder, Boulder, CO 80303, USA

*Correspondence: michael.smallegan@colorado.edu**,** john.rinn@colorado.edu

Abstract

There are currently over 16,000 long-noncoding RNAs (lncRNAs) annotated in mammalian genomes, a number on par with protein-coding genes. Numerous studies have demonstrated that lncRNAs can influence gene expression programs but lack the temporal resolution to identify the immediate regulatory events. Thus, even for the most well-studied, the early and primary targets of lncRNAs remain elusive. Here we describe a multifaceted effort to determine the primary targets of a lncRNA and the underlying temporal dynamics. Our approach combines several genetically defined loss-of and gain-of-function, rescue, and inducible lncRNA models. We further resolved gene regulatory events temporally (30 min to four days). Applying this approach, we find that the lncRNA *Firre* is an RNA-based transcriptional activator. These transcriptional activation events are not mediated by the epigenetic regulators PRC2, WDR5, or G9A, nor are they due to the proximal binding of the *Firre* lncRNA. Instead, *Firre* activates transcription by increasing chromatin accessibility (approximately 30 min after induction) and activating primary target genes (about an hour later). Overall, this study suggests *Firre* functions as an RNA-based transcriptional activator.

Introduction

Mammalian genomes are replete with novel regulatory elements, including DNA enhancers, newly discovered small proteins, and long noncoding RNAs (lncRNAs) (Abascal et al. 2020; Consortium 2012; Derrien et al. 2012). These findings have inspired efforts to define the targets of these newfound noncoding regulatory elements. For example, there are more than 16,000 lncRNAs in the human genome (Derrien et al. 2012; Cabili et al. 2011; Consortium 2012; Abascal et al. 2020), a number on par with protein-coding genes – yet their primary regulatory sites remain poorly understood. Numerous studies have demonstrated that lncRNAs can influence gene expression programs (Rinn and Chang 2012, 2020; Wang and Chang 2011; Morris and Mattick 2014). However, only a few lncRNAs have been identified to play important roles *in vivo*, and even fewer have been mechanistically characterized (Andergassen and Rinn 2021).

A major limitation in understanding lncRNAs' mechanistic function has been the lack of knowing the direct regulatory targets. Determining primary targets for lncRNAs that function *in cis* has been facile and is usually due to a DNA-based enhancer encoded in the locus or a low abundance lncRNA that facilitates promoter activation *in cis* (Tong and Yin 2021; Tan et al. 2017; Groff et al. 2016; Maamar et al. 2013; Dimitrova et al. 2014; Paralkar et al. 2016; Guttman and Rinn 2012; Maass et al. 2012; Elling et al. 2018; Miyagawa et al. 2012; Engreitz et al. 2016; Rinn and Chang 2020; Carpenter et al. 2013). In contrast, *trans*-acting RNA-based gene regulatory events are less well understood (Guttman and Rinn 2012). This raises the question: Do lncRNAs act directly to regulate transcription, akin to transcription factors (TFs), or are they affecting other cellular processes resulting in gene regulation from a distance?

Transcription factors bind, in *trans*, to DNA and act by recruiting transcriptional machinery (Lee and Young 2000; Mitchell and Tjian 1989). TF binding in proximity to the transcription start site (TSS) is an orthogonal piece of information that permits the inference of direct action independent of the timing of gene expression changes (Lammers et al. 2020; Wang and Deng 2022;

Charoensawan et al. 2015; Stewart-Ornstein et al. 2013; Weikum et al. 2017). We currently lack a parallel mechanistic understanding of lncRNA-based gene regulation.

The time scale at which lncRNAs affect gene expression is also relatively unexplored. Most loss- or gain-of-function studies of lncRNAs measure the transcriptome at a single time point on the order of multiple hours or days (Guttman et al. 2011; Huarte et al. 2010; Loewer et al. 2010; Guttman and Rinn 2012). Thus, the earliest or primary effects of the lncRNA are obscured by layers of gene regulatory cascades and compensation mechanisms. In contrast, TFs can regulate primary gene targets within minutes. For example, p53 and NFKB *trans*-activate their primary target sites within minutes (Allen et al. 2014; Mahat et al. 2016; Sasse et al. 2019). For understanding TF regulation, the temporal dynamics of gene expression changes became essential to disentangling primary (and likely direct) effects from secondary and tertiary effects.

Here we set out to identify the primary targets and underlying temporal dynamics of lncRNA-based gene regulation *in trans*. This effort required choosing a lncRNA that is known to be functional and that functions *in trans*. To this end we chose to first focus on the lncRNA *Firre* (functional intergenic repeating RNA element), based on the following properties: (i) *Firre* is a *trans*-acting RNA that does not regulate gene expression *in cis*; (ii) gain- and loss-of *Firre* RNA function *in vivo* phenotypes range from lethality under LPS exposure to defects in hematopoiesis, respectively; (iii) *Firre* is syntenically conserved across vertebrates; (iv) *Firre* is genetically associated with human diseases (Lewandowski et al. 2019; Abe et al. 2014; Hacisuleyman et al. 2016, 2014; Barutcu et al. 2018). Although many studies have attempted to understand the mechanism of *Firre*, these efforts have resulted in several conflicting models; a conundrum that the knowledge of the primary regulatory sites of *Firre* would help to sort out.

To determine the primary regulatory sites of *Firre*, we used a combination of genetically defined mouse embryonic stem cell (mESC) lines. Briefly, we derived wild-type (WT) and *Firre* knockout (*Firre^{KO}*) lines with an inducible *Firre* transgene. Importantly, we have a control cell line in each

genetic background to account for the effect of doxycycline (dox) – thus controlling for non-specific dox effects.

We monitored gene regulation events by RNA-seq across 16 time points (from 30 min to four days) to identify the temporal order of *Firre*-based regulatory events. We also performed a time course of ATAC-seq to decipher the earliest changes in chromatin accessibility that lead to changes in target gene expression. Finally, we combined this temporal information across genotypes to identify *Firre* targets that occur in all conditions tested and are not due to dox effects. We identified 30 genes commonly regulated in WT and *Firre*^{KO} genetic backgrounds upon induction of *Firre*.

Interestingly, all but one of these genes are upregulated, suggesting that *Firre* has transcriptional activation activity. Using loss-of-function studies, we determined that this gene activation is not due to key epigenetic factors (WDR5, G9a, or PRC2), as previously suggested. We further determined where *Firre* was localized relative to these primary gene-regulation events using RNA affinity purification followed by DNA sequencing (RAP-DNA-seq) and single molecule RNA FISH (smRNA FISH). We found that *Firre* RNA does not appear to localize in proximity to these sites, despite the rapid induction. Instead, *Firre* functions to increase chromatin accessibility as measured by ATAC-seq in less than 30 minutes – that in turn leads to mature transcriptional output approximately an hour later. Collectively, our study identifies *Firre* as an RNA-based transcriptional activator that functions via increasing chromatin accessibility from a distance.

Results

Firre's gene regulatory effects across static WT and KO samples

We initially set out to pinpoint possible regulatory targets of *Firre* using a *Firre^{KO}* mouse model. We reasoned that the possible regulatory targets of *Firre* would be commonly deregulated across multiple tissues. To this end, we analyzed previously-generated RNA-seq experiments comparing nine WT and *Firre^{KO}* primary tissues (Fig. 1A) (Andergassen et al. 2019; Lewandowski et al. 2019). We observed a total of 842 genes that were differentially expressed (WT vs *Firre^{KO}* for a given tissue; P < 0.05, |fold change| >= 1.5) in at least one tissue (Fig. 1B). However, there were very few differentially expressed genes in common in multiple tissue contexts (69 genes in two or more tissues and four genes in three or more tissues). The only gene commonly differentially expressed in all nine tissues was *Firre*. Together, these results suggest either that *Firre* regulation may be highly tissue-specific or that tissue heterogeneity confounds identifying regulatory targets of *Firre*.

To control for cellular heterogeneity in tissues, we derived mESCs from WT and *Firre^{KO}* mice. We performed RNA-seq on WT and *Firre^{KO}* mESCs and found 921 genes that were differentially expressed (314 downregulated and 607 upregulated, P < 0.05, |fold change| >= 1.5; Fig. 1C). We note that this number of possible regulatory targets far exceeds the number of approximately 30 *Firre* molecules per cell in mESCs (Much et al. 2022). Due to compensatory mechanisms and the stoichiometry of *Firre* to differentially regulated genes, these expression changes are not likely to be due to the direct action of *Firre*.

Figure 1: Gene expression changes upon *Firre* knockout *in vivo* and in mESCs. (A) Schematic of tissues examined by RNA-seq in a *Firre* knockout mouse model. **(B)** Bar plot showing the number of genes differentially regulated in WT vs *Firre^{KO}* tissues classified by the number of tissues in which those genes are commonly deregulated. **(C)** Volcano plot showing a total of 921 genes significantly differentially expressed in *Firre^{KO}* relative to WT mESCs. *Firre* not shown: log$_2$ fold change -6.9, -log$_{10}$(p-value) = 196.6.

Temporal gene regulation by *Firre* in a rescue model

To dispense with the confounding effects of compensation, we need to be able to control the expression of *Firre* and measure the dynamics of gene expression changes before the cell returns to homeostasis. However, the rate at which lncRNAs mediate gene expression changes is unknown. Therefore, we started by measuring the transcriptome at widely spaced time intervals up to four days after the induction of *Firre*.

To temporally resolve gene regulatory effects by *Firre*, we generated *Firre^{KO}* mESCs that also contain a dox-inducible *Firre* transgene; we term this line *Firre^{RESCUE}* (Fig. 2A; Supplemental Fig. S1A). Since dox itself causes gene expression changes (Ahler et al. 2013), we derived a dox

control line that we term $Firre^{KO\text{-}CTL}$ (Fig. 2A). This cell line is genetically identical to the $Firre^{RESCUE}$ line but the reverse tetracycline-controlled transactivator (rtTA) element is not expressed (Supplemental Fig. S1B).

We first aimed to determine *Firre* transgene expression in the $Firre^{RESCUE}$ and $Firre^{KO\text{-}CTL}$ lines at various time points post induction. Specifically, we treated both cell lines with dox for 0, 12, 24, 48, and 96 hours and performed quantitative reverse-transcription PCR (RT-qPCR) using probes detecting endogenous and transgene *Firre*, respectively. In the $Firre^{RESCUE}$ line, *Firre* transgene expression was induced to WT levels by 12 hours and remained expressed through four days, while endogenous *Firre* remained undetectable (Fig. 2B). As expected, we did not observe any expression of transgenic *Firre* in the $Firre^{KO\text{-}CTL}$ line.

To identify temporally regulated genes upon the induction of *Firre*, we generated a transcriptome-wide time course in the $Firre^{RESCUE}$ and $Firre^{KO\text{-}CTL}$ cell lines. We treated cells with dox for 12, 24, 48, and 96 hours and performed RNA-seq with three replicates at each time point. We observed that *Firre*'s expression measured by RNA-seq followed the induction pattern detected by RT-qPCR (Fig. 2C; Supplemental Fig. S2A).

As dox is known to induce gene expression changes (Ahler et al. 2013), we next characterized the dox gene-regulatory effect in the dox control line $Firre^{KO\text{-}CTL}$. We determined differentially expressed genes due to dox by comparing each gene's expression at a given time point to the zero-time point. Surprisingly, treatment with dox dynamically regulated 1,978 genes ($P < 0.05$; |fold change| >= 1.5; Fig. 2D), indicating that it is essential to control for the gene-regulatory effects of dox in the forthcoming experiments.

To identify genes regulated by *Firre* and independent of the effect of dox, we used a likelihood ratio test between two regression models (Fig. 2E). The first model contains terms for time (t), the baseline difference between cell lines (F_i), and the interaction of the two ($t*F_i$). The second model lacks the interaction term. Genes regulated similarly in the $Firre^{RESCUE}$ and $Firre^{KO\text{-}CTL}$ lines fit well in both models. However, if the interaction term has an explanatory value, the likelihood ratio test

will have a small p-value. Using this approach, we found 414 genes to be significantly regulated

by *Firre* across time (P < 0.05; |fold change| >= 1.5; Fig. 2F; Supplemental Fig. S2B-C).

Figure 2: Temporal gene regulation in a *Firre* transgene rescue model. (A) Schematic showing the mouse blastocyst-derived mESC line *Firre^RESCUE* containing a dox-inducible *Firre* transgene in a *Firre* KO background, and the *Firre^KO-CTL* cell line that does not induce *Firre* expression and serves as a dox control. **(B)** Relative expression of transgenic *Firre* after dox administration in *Firre^RESCUE* mESCs as determined by qRT-PCR. The dashed line indicates *Firre* WT expression levels. **(C)** Expression levels of *Firre* in *Firre^RESCUE* (red) and *Firre^KO-CTL* mESCs (black) after the addition of dox as measured by RNA-seq. The dashed line indicates *Firre* WT expression levels. **(D)** $\log_2$ fold change of significantly differentially expressed genes that are affected by dox treatment in the *Firre^KO-CTL* control cell line. Fold changes are relative to the zero-time point. **(E)** Linear model to derive genes that are significantly differentially regulated upon *Firre* transgene induction and not due to dox treatment (likelihood ratio test). **(F)** Heatmap of the expression fold change of significantly differentially expressed genes changing over time upon *Firre* induction independent of dox effects. Shades of red and blue represent $\log_2$ fold changes relative to the zero-time point of up- and downregulation, respectively, upon *Firre* transgene induction. Black tick marks represent genes whose expression was "rescued", i.e., that were downregulated in the knockout and upregulated upon transgene induction, or vice versa. **(G)** Venn diagram of the number of significantly differentially expressed genes (overlap) in *Firre^KO-CTL* vs WT and *Firre^KO-CTL* vs *Firre^RESCUE* cells. **(H)** Heatmap of the expression of 68 rescued genes across each time point. Top and bottom heatmap represent the *Firre^KO* and WT expression levels, respectively. The heatmaps in between represent the temporal timing of when genes are rescued. **(I)** Principal component analysis of the rescued gene expression levels in WT, *Firre^KO* and *Firre^RESCUE* cells at each time point. The primary component (PC1 that represents 75% of the variation in the data) is plotted.

To assess the extent to which the induction of *Firre* in the *Firre^KO* background restored the WT expression pattern, we identified genes that were significantly differentially expressed in response

to *Firre*'s induction and differentially expressed in the static *Firre*[KO] vs WT analysis. The genes responding to *Firre*'s induction significantly overlapped the differentially expressed genes in the *Firre*[KO] mESCs (102 genes; hypergeometric $P = 5.79 \times 10^{-70}$; Fig. 2G). The overall pattern of expression changes indicated a reciprocal relationship (Pearson correlation = -0.81; $P < 2.2 \times 10^{-16}$; Supplemental Fig. S2D). We considered genes reciprocally regulated to within 20% of the *Firre*[KO] vs. WT fold change to be rescued (68 genes; Fig. 2H; Supplemental Fig. S2E-F). We next examined the temporal dynamics underlying how gene-expression programs return to WT levels (Fig. 2H). Notably, there appears to be a transition between 12 and 24 hours. To test this, we performed a principal component analysis on the expression values of the rescued genes. The principal component values showed a high similarity in cell lines not exposed to *Firre* and the 12-hour time point. We observed a shift towards rescued gene-expression levels at 24 hours. Interestingly, after this larger shift at 24 hours, the remaining changes are more gradual towards the WT state (Fig. 2H-I). In summary, we established a *Firre*-inducible expression system and model that can be harnessed to monitor gene expression changes over time.

Gene regulation by *Firre* before twelve hours of induction in the rescue model

The time course above established that gene expression changes occur by twelve hours; however, expression changes in a shorter time window would give more confidence that these constitute direct effects of *Firre*. As a first test, we measured the levels of *Firre* induction by RT-qPCR at 0, 3, 6, and 12 hours. Interestingly, *Firre* reached maximal expression at three hours (Fig. 3A). Based on the early induction of *Firre* at three hours we performed RNA-seq at a much more fine-grained temporal resolution prior to twelve hours. Specifically, we collected RNA every 30 minutes until 5.5 hours post addition of dox and performed RNA-seq (Fig. 3B). We observed induction of *Firre* expression already within 30 minutes (Fig. 3C), with a maximal induction around two hours (18.5-fold above WT levels; Supplemental Fig. S3A). We also observed that mESCs have minimal expression changes resulting from dox exposure in this short time window relative

to the response at twelve hours (twelve differentially expressed genes; P < 0.05; |fold change| >= 1.5; Fig. 3D).

We next analyzed this short time course to home in on early *Firre* regulatory events. We used the same likelihood ratio test approach (Fig. 2E) that determines genes significantly regulated by *Firre* induction across time while controlling for dox-responsive genes. We identified 18 genes (twelve upregulated and six downregulated) that respond to *Firre* (Fig. 3E; Supplemental Fig. S3B-C). We note that over the 5.5-hour time course, the upregulated genes continue their trajectory to higher expression levels, while the downregulated genes transiently decrease expression and then return to their original expression levels (Fig. 3E; Supplemental Fig. S3C-D). Together these results suggest that *Firre*'s rapid induction of expression mediates robust upregulation of some genes and transient downregulation of others.

Figure 3: Early temporal regulation in a *Firre* transgene rescue model. (A) Relative expression of the *Firre* transgene over time after dox administration in *Firre^RESCUE* mESCs as measured by RT-qPCR. **(B)** Schematic showing sampling of cells that were treated with dox for 0 to 5.5 hours at intervals of 30 min for RNA-seq. **(C)** Expression levels of *Firre* in *Firre^RESCUE* (red) and *Firre^KO-CTL* (black) cell lines over time as measured by RNA-seq. Bottom, zoom in of *Firre* expression at early time points. **(D)** $\log_2$ fold change of significantly differentially expressed genes that are affected by dox treatment in the *Firre^KO-CTL* control cell line. Fold changes are relative to the zero-time point. **(E)** Heatmap of the expression fold change of genes differentially expressed due to *Firre* transgene induction and not dox treatment at each early time point.

Gene regulation by *Firre* in an overexpression model

We hypothesized that the robust primary targets of *Firre* would be affected similarly in cells with a *Firre* transgene in a WT background. To test this, we generated a *Firre* WT mESC line harboring the same dox-inducible *Firre* transgene (*Firre* overexpression line / *Firre^OE*). WT cells served as a dox control (Fig. 4A). We first used RT-qPCR to characterize the induction of the *Firre* transgene in the WT and *Firre^OE* cells. In the *Firre^OE* line, we observed a rapid induction of the *Firre* transgene with a maximum expression at three hours and taper around six hours, while endogenous *Firre* levels remained unchanged (Fig. 4B). To mirror our previous experiments in the rescue model, we generated a long (96-hour; three replicates) and a short (5.5-hour; two replicates) time course that we sampled for RNA-seq. Similar to the kinetics in the *Firre^RESCUE* line, we observed a maximal induction of *Firre* in the *Firre^OE* line (46.2-fold WT levels) at two hours, and *Firre* expression levels returned to 4.5-fold WT levels by twelve hours (Fig. 4C-D; Supplemental Fig. S4A-C).

To discard false-positive *Firre*-responsive genes, we next analyzed our RNA-seq data for dox-related gene expression changes in the WT cell line. As we did previously, we characterized the effects of dox alone by comparing gene expression at different time points after dox administration

to the zero time-point. We observed 63 genes changing within the 5.5-hour window and 762 genes changing in the longer time course (Supplemental Fig. S4D-E). To identify *Firre*-responsive genes, we used the same regression models and likelihood ratio test described above (Fig. 2E), which identifies genes temporally regulated by *Firre* but not due to dox effects. In the 96-hour time course, we identified 176 differentially expressed genes (P < 0.05; |fold change| > 1.5; Fig. 4E; Supplemental Fig. S4F-G) that exhibit coherent regulation by *Firre* across time. We note that this is a smaller number of regulated genes than observed in the time course with the *Firre^{RESCUE}* line. Nonetheless, a significant fraction of affected genes overlap in the *Firre^{OE}* and *Firre^{RESCUE}* line (hypergeometric P = 2.2x10^{-107}; Fig. 4F). We next used the same analyses on our early time course to monitor gene expression changes every 30 minutes for 5.5 hours. We identified 26 (25 upregulated and one downregulated) potential early *Firre*-regulatory targets (Fig. 4G; Supplemental Fig. S4H-J). We also observed a significant (hypergeometric P = 4.21x10^{-42}; Fig. 4H) overlap between immediate *Firre*-responding genes in *Firre^{OE}* and *Firre^{RESCUE}* lines. Collectively, we found good correspondence in genes that are regulated in both the shorter and longer time course across overexpression and rescue models.

Figure 4: Temporal regulation of *Firre* transgene in a WT background. (A) Schematic showing the *Firre^OE* cell line containing a dox-inducible *Firre* transgene in a WT background, and a WT cell line that serves as a dox control. **(B)** Relative expression of endogenous and transgenic *Firre* after dox administration in WT and *Firre^OE* mESCs as determined by qRT-PCR. **(C, D)** RNA-seq measurement of *Firre* expression in WT (black) and *Firre^OE* (red) over the long (C) and short (D) time course. **(E, G)** Heatmap of expression fold changes of genes that are significantly

differentially expressed upon induction of *Firre* and independent of dox treatment over the long (E) and short (G) time course. **(F, H)** Venn diagram representing the overlap of *Firre*-responsive genes in *Firre^{OE}* and *Firre^{RESCUE}* models in the long (F) and short (H) time course.

Firre-regulatory targets across time and genotype

To generate a high-confidence list of *Firre* targets, we next wanted to determine the genes that respond rapidly to *Firre* but not dox in both the rescue and the overexpression model. To this end, we combined the data from the short time course: 96 RNA-seq experiments across 12 time points and four distinct genotypes (Fig. 5A). We used the same likelihood ratio test as described previously and added a term describing the genetic background (F_{KO}) (Fig. 5B).

Using this model, we identified 29 significantly differentially expressed (P < 0.05, |fold change| > 1.5) early *Firre*-responsive genes (Fig. 5C; Supplemental Fig. S5A-B). All but one (*Prdm14*) of these genes were upregulated. These genes started increasing expression in 1.5-4.5 hours, and 21 out of the 28 upregulated genes remained at elevated levels through 96 hours (Fig. 5C-D; Supplemental Fig. S5A-B). Since these 29 genes are rapidly regulated concordantly in two genetic backgrounds in response to *Firre*, they represent high-confidence candidates to be *Firre* target genes.

We next wanted to independently verify an early *Firre*-responsive gene by smRNA FISH. We chose *Shf* due to its higher basal expression levels and robust change in expression over time, and designed a complementary probe set to monitor its abundance and localization across time. We tested 0, 2, 4, and 6-hour time points after *Firre* induction with dox in both *Firre^{RESCUE}* and *Firre^{OE}* lines. In both cell lines, we observed a continuous increase in the expression of *Shf* after the induction of *Firre*, verifying our RNA-seq results (Supplemental Fig. S6A). In summary, we identified a cohort of high-confidence *Firre* target genes that are predominantly upregulated upon *Firre* induction.

Figure 5: *Firre* targets across genotype and time. (A) Schematic of all *Firre* mESC lines tested. **(B)** Linear model to determine genes that are regulated by *Firre* transgene independent of dox and across WT and KO backgrounds. **(C)** Heatmap of gene expression changes of Firre-responsive genes over time as determined by the linear model in (B). **(D)** Gene expression changes of representative genes due to *Firre* in *Firre*-inducing (red) but not in dox control (black) cells.

Epigenetic and proximal influence of *Firre* trans-activation

Although many lncRNAs have been identified to bind to epigenetic regulatory factors, how these interactions affect gene expression remains poorly understood. One hypothesis is that *Firre* could bind to epigenetic complexes that regulate histone modifications and alter these modifications proximal to regulatory sites. To test this hypothesis, we designed a strategy to examine the influence of the Polycomb repressive complex 2 (PRC2), *G9a* (EHMT2), and WDR5 that control histone H3 lysine 27 trimethylation (H3K27me3), H3 lysine 9 trimethylation (H3K9me3), and H3 lysine 4 trimethylation (H3K4me3) states, respectively.

Specifically, we used GapmeRs to knockdown the PRC2 components *Ezh2*, *Suz12*, *Jarid2*, and *Aebp2* as well as *G9a* and *Wdr5* in *Firre*^{OE} cells. After 24 hours of GapmeR exposure, the *Firre* transgene was induced by dox for six hours (Fig. 6A). We reasoned that if any of the three epigenetic complexes were required for *Firre*-mediated gene expression regulation, their knockdown would abolish the upregulation of the *Firre*-responsive genes *Adgrg1* and *Shf* (Fig. 6B).

We validated the effective depletion of PRC2, G9a, and WDR5 complexes at the RNA and protein level (Supplemental Fig. S6B). Using a non-targeting control (NTC) GapmeR, we confirmed the induction of *Firre* and the concomitant upregulation of the *Firre* responders *Adgrg1* and *Shf* after the addition of dox, while the non-*Firre* target *Tug1* remained unchanged (Fig. 6C and Supplemental Fig. S6C). Surprisingly, we found that the depletion of any of these three epigenetic regulatory complexes did not decrease the upregulation of the tested *Firre*-responsive genes (Fig. 6C). Collectively, these results demonstrate that *Firre*-mediated activation of target genes is not dependent on the epigenetic complexes PRC2, G9a, and WDR5.

To probe for another possible mechanism of how *Firre* regulates its targets, we next determined if the *Firre* RNA is in proximity to its target genes. We first leveraged previously published RNA affinity purification (RAP) data that identified 34 *Firre*-bound genomic locations in mESCs (Hacisuleyman et al. 2014) where *Firre* was bound and searched for overlaps with *Firre* responder

genes. The 34 RAP peaks did not overlap within 50 kb of the gene bodies of any of the *Firre* responder genes identified in any experiment (520 genes). Thus, under homeostatic conditions, *Firre* does not appear to be in proximity to genes with early or late expression changes.

Figure 6: Probing epigenetic and proximal mechanisms of *Firre*-mediated gene regulation.

(A) Timeline of GapmeR-mediated gene knockdown and dox-induced *Firre* transgene expression.

(B) Depiction of possible outcomes for *Firre*-responsive gene expression (red line) after knockdown of a regulatory factor (blue line) and induction of *Firre* transgene expression. **(C)** Relative gene expression of *Firre* and its targets *Adgrg1* and *Shf* in *Firre^OE* cells transfected with non-targeting control- (NTC), *Ezh2*-, *Suz12, Jarid2*-, *Aebp2*-, *Wdr5*-, or *G9a*-targeting GapmeRs, and treated without (zero hours) or with (six hours) dox as determined by RT-qPCR. **(D** and **E)** smRNA FISH-based quantification of the distance between *Shf* intron and *Firre* transgene (D) and *Shf* intron and the five closest *Firre* exons (distance averaged) (E) in *Firre^OE* and *Firre^RESCUE* cells after 0, 2, 4, and 6 hours of dox treatment. The images below depict the rationale of the distance measurements.

As a second approach, we aimed at determining the spatial relationship between *Firre* and its target genes by smRNA FISH. To this end, we designed differentially labeled probe sets complementary to *Firre* exons as well as the first intron of the early response gene *Shf* and performed smRNA FISH at 0, 2, 4, and 6 hours post-*Firre* induction in the *Firre^OE* and *Firre^RESCUE* cell lines. We then measured the distance of the site of transcription of *Shf*, as indicated by its intron signal, from the site of transcription of *Firre*, as approximated by the typical *Firre* exon-stained "expression cloud" amongst the many individual *Firre* molecules in the nucleoplasm. We found that the *Firre* transgene was not in close proximity to the *Shf* gene nor increased in proximity upon induction of *Firre*, indicating no direct interaction of the *Firre* and *Shf* loci (Fig. 6D). We next determined if individual *Firre* RNA molecules were in proximity to the *Shf* locus. We found that *Firre* exons on average were more than one micron away from the *Shf* locus and remained so despite increased expression of *Firre* and *Shf* (Fig. 6E, Supplemental Fig. S6A). Thus, both RAP and smRNA FISH demonstrate that *Firre* does not form direct contacts with its most immediate and primary gene targets.

Firre regulates chromatin accessibility proximal to regulatory targets

Another possibility for how _Firre_ could be activating genes is by increasing chromatin accessibility in _cis_-regulatory regions of the target genes. To this end, we performed ATAC-seq across time in the four different mESC lines. We focused on a short time window and generated ATAC-seq libraries from cells harvested every 30 min for 2.5 hours to identify the earliest changes in chromatin accessibility.

Using MACS2 to call peaks separately in each sequenced sample, we merged the peak sets to create a consensus peak set with 145,143 peaks. Peak signals were quantified by counting the number of reads in each peak across all samples. To detect peaks changing accessibility due to the effect of dox, we compared the peak signals of each time point to those of the zero-time point in the control cell lines. We found no peaks significantly changing in time due to dox (zero dox associated peak changes; $P < 0.05$). We further determined peaks that changed in both rescue and overexpression models. Briefly, we tested for differential accessibility by comparing all conditions in which _Firre_ is induced (time point > 0 in _Firre^{RESCUE}_ and _Firre^{OE}_ lines) to conditions in which the _Firre_ transgene was not induced (WT and _Firre^{KO-CTL}_ lines; zero-time point in _Firre^{RESCUE}_ and _Firre^{OE}_ lines). We identified a total of 55 ATAC peaks that changed ($P < 0.05$) in response to _Firre_'s induction (Fig. 7A; Supplemental Fig. S7A). Of these 55 peaks, the vast majority (51 peaks) increased in accessibility, while only four decreased in accessibility over the time course (Fig. 7A; Supplemental Fig. S7A).

Next, we wanted to determine how many of the differentially accessible ATAC peaks were in proximity to genes changing in expression upon _Firre_ induction. We identified ten differentially accessible ATAC peaks within +/- 50 kb of the TSS of early _Firre_-responsive genes. One of these peaks was in proximity to two neighboring genes which both increased in expression (_Duox1_ and _Shf_). This revealed eleven genes, including _Firre_ itself, that showed proximal increased chromatin accessibility and concordant temporal regulation by _Firre_ (Fig. 7B). Analysis of the ATAC-seq

peak center location relative to the TSS for the ten non-*Firre* genes revealed that all but one ATAC-peaks were downstream of the TSS (Fig. 7C). The gene *Rapgef4*, for example, rapidly gained ATAC peak signals within its first intron at a proximal enhancer element (Fig. 7D; Supplemental Fig. S7B).

We next wanted to define the timing of chromatin accessibility and gene expression changes upon *Firre* induction. To this end, we compared the temporal dynamics of ATAC- and RNA-seq changes side by side. We found that for all eleven genes, chromatin accessibility was increasing already within 30 min and before most gene expression changes occurred (Fig. 7E).

Figure 7: Changes in chromatin accessibility upon induction of *Firre* transgene. (A) Heatmap showing chromatin accessibility $\log_2$ fold changes across time. Increased chromatin accessibility relative to the zero-time point is depicted in shades of red, decreased chromatin accessibility in blue. Black tick marks indicate differentially expressed genes. **(B)** Venn diagram of *Firre*-responsive genes and chromatin accessibility sites. **(C)** TSS metaplot of the density of ATAC-seq peak centers relative to *Firre*-responsive gene TSSs. Tick marks indicate the peak centers. **(D)** Genome browser tracks of chromatin accessibility changes across time within *Rapgef4* in *Firre^RESCUE* cells. **(E)** Heatmap of chromatin accessibility and gene expression changes at each time point after induction of *Firre.* Time points where chromatin accessibility was not measured are indicated in grey. **(F)** Plot of chromatin accessibility (red) and gene expression (black) signal relative to the maximum measured over time. The dashed line represents the time difference in half-maximum values.

To formalize this observation, we created a meta-profile of ATAC-seq signal changes and RNA-seq signal changes for these genes (Fig. 7F). We observe that, on average, ATAC peaks and RNA abundance reach half-maximal at 47 minutes and 104 minutes, respectively. Collectively, these data provide evidence that *Firre* functions as an RNA-based transcriptional activator by increasing chromatin accessibility that in turn activates increased RNA transcription.

Discussion

The mammalian genome encodes thousands of recently discovered noncoding regulatory elements – ranging from newly identified enhancers to lncRNAs (Consortium 2012; Abascal et al. 2020; Birney et al. 2007; Consortium 2004). Many recent efforts have focused on identifying where, when, and how these sites regulate transcription. This has led to many new insights into how enhancers and lncRNAs work *in cis* to activate or repress target genes (Anderson et al. 2016; Fanucchi et al. 2018; Krawczyk and Emerson 2014; Gil and Ulitsky 2019). However, our understanding of *trans*-acting elements has remained poorly understood. Arguably the least is known about how lncRNAs work to regulate gene expression.

We found that *Firre*-mediated regulatory events occurred rapidly and rescued loss-of-function defects over much longer time scales. Specifically, we found that induction of *Firre* increased chromatin accessibility within 30 minutes, initial gene expression changes occurred in a few hours, and reached homeostasis (in a KO background) after 24 hours. These findings demonstrate the importance of combining very short and longer time frames of lncRNA-mediated gene regulation. Short time courses provide insight into the primary targets, and longer time courses identify stable changes to the gene expression program that are more likely to potentiate disease states associated with *Firre*.

We further investigated the possible mechanisms of the observed gene activation by *Firre*. We started with testing a common theme that lncRNAs bind and influence epigenetic processes (e.g., histone modifications). We found that the loss-of-function for three key epigenetic pathways (PRC2, WDR5, and G9a) did not influence gene activation by *Firre*. We observed similar *Firre*-mediated gene activation rates under all conditions tested – further underscoring the robustness of these regulatory events. Thus, these early gene regulatory events are unlikely due to a commonly proposed mechanistic model of recruitment or eviction of epigenetic complexes to derepress or activate *Firre* target genes.

We also tested if the *Firre* RNA localized to these sites by two independent methods: RAP and smRNA FISH. In neither case did we observe that *Firre* binds in proximity to its regulatory sites. Using smRNA FISH we homed in on time points that corresponded to near and after activation events and did not find colocalization of *Firre* exons at sites of gene activation. A possible caveat to both approaches is the targeted nature requiring probes to hybridize to *Firre in situ*. It is possible, although unlikely, that *Firre* cannot be targeted by either approach if it was bound by chromatin complexes making it inaccessible to probes. In contrast, we observed a swift increase in chromatin accessibility within minutes of inducing *Firre*. These findings are surprising in that *Firre* does not need to be in proximity to corresponding sites of gene regulation despite such rapid activation of these genes.

Together these results raise the question of how does a lncRNA regulate gene expression so rapidly? Based on our observation that chromatin accessibility changes precede any gene expression changes, it is unlikely that these effects are the result of changes in the abundance of a protein. That would require a gene to be activated, spliced, translated, and shuttled back to the nucleus, which seems unlikely to occur within 30 minutes. Moreover, it is unlikely that *Firre* directly binds to these sites and activates gene expression akin to a transcription factor. Finally, since *Firre* is not in proximity to these sites it is unlikely able to evict or recruit epigenetic complexes to derepress or activate chromatin.

Yet, the chromatin accessibility sites hold a clue of how *Firre* could activate transcription. Based on this information about what is unlikely we can envision a few possible mechanistic insights. First, most of the earliest regulatory events occur downstream of promoter regions, suggesting that *Firre* does not affect transcription at the promoter. Rather these data suggest *Firre* is activating promoter-proximal enhancers to induce target gene expression. Second, since *Firre* does not appear to localize to these sites it suggests that *Firre* must be functioning in the nucleoplasm (as *Firre* localization is almost exclusively nuclear). It is likely that this nucleoplasmic

function involves a ribonucleic-based molecular reaction that rapidly leads to either repressing a repressor or activating an activator of transcription, respectively (Brivanlou and Darnell 2002). This would also hold true for interactions with chromatin-remodeling complexes prior to their localization at these regulatory sites.

We can now test these, and other possible RNA-based gene activation models owing to the identification of the early and primary regulatory events. These sites can be assayed to determine what molecular interactions occur upon induction of *Firre*. Thus, we argue the best first step in determining the mechanism of a lncRNA is knowing the primary targets. It is much more difficult to determine the mechanism before knowing the primary regulatory sites. Collectively, these results represent a way forward in dissecting the mechanism by which a model lncRNA can activate gene expression from a distance.

Supplemental Figures

Supplemental Figure S1: Characterization of the mESC lines used. (A) Diagram showing the

Firre transgene construct and its induction by dox through the rtTA protein. **(B)** Genome browser

alignment tracks of RNA-seq reads over the rtTA3 element, which is expressed in *Firre*^{RESCUE} and

Firre^{OE} cell lines, but not in the WT and *Firre*$^{KO-CTL}$ dox controls.

Supplemental Figure S2: Temporal gene regulation in the *Firre^RESCUE* line. (A) *Firre* expression measured by RNA-seq in the *Firre^RESCUE* line (red) relative to WT line (black). **(B)** Time course plot of the shrunken $\log_2$ expression fold changes relative to the zero-time point of *Firre*-responsive genes with the numbers of up- and downregulated genes indicated. **(C)** Bar plot showing the number of genes with a |fold change| > 1.5 at each time point. Red bars indicate the number of genes upregulated while black bars indicate the number of genes downregulated. **(D)** Scatter plot of the expression fold changes of the significantly differentially expressed genes overlapping in the *Firre^KO* vs WT and in the *Firre^RESCUE* time course. The time course fold change (y-axis) is represented by the maximum fold change in any one time point. A linear regression is plotted in blue. **(E)** Scatter plot of the expression fold changes of the significantly differentially expressed genes overlapping in the *Firre^KO* vs WT and in the *Firre^RESCUE* time course. The dashed lines indicate the 20% reciprocal fold change cutoff for genes to be considered rescued. Rescued genes are plotted in red and genes that did not make the cut-off are plotted in grey. **(F)** Venn

diagram showing the number of rescued genes compared to the total number of *Firre^KO* vs WT

overlapping differentially expressed genes.

Supplemental Figure S3: Dynamic gene regulation in the *Firre^RESCUE* short time course. **(A)**

Firre expression in the *Firre^RESCUE* line (red) relative to the WT line (black) as determined by RNA-

seq. **(B)** Time course plot of the shrunken $\log_2$ fold changes for the genes significantly differentially

expressed. **(C)** Bar plot showing the number of genes with a |fold change| > 1.5 at each time

point. Red bars indicate the number of genes upregulated while black bars indicate the number

of genes downregulated. **(D)** Individual profile plots of the expression $\log_2$ fold change compared

to the zero-time point for the significantly differentially expressed genes.

Supplemental Figure S4: Gene regulation dynamics in the *Firre^OE* time course. **(A)** *Firre* expression profile measured by RNA-seq across all time points in the *Firre^OE* line (red) and WT line (black). **(B-C)** *Firre* expression in the *Firre^OE* line (red) relative to the WT line (black) in the long (B) and short time course (C) as determined by RNA-seq. **(D-E)** Time course plot of the shrunken $\log_2$ expression fold changes for the significantly differentially expressed genes in the WT dox control line in the short (D) and long (E) time course. **(F)** The number of differentially expressed genes with |fold change| > 1.5 at each time point in the *Firre^OE* line. **(G-H)** Time course plot of the shrunken $\log_2$ expression fold changes for the differentially expressed genes in the *Firre^OE* in the long (G) and short (H) time course. **(I)** The number of genes at each time point with |fold change| > 1.5 for the *Firre^OE* short time course. **(J)** Individual gene expression profile plots showing the fold change vs the zero-time point for the significantly differentially expressed genes in the *Firre^OE* short time course.

Supplemental Figure S5: Gene expression profile plots for genes detected as differentially expressed in the combined model. (A) Time course plot of the shrunken $\log_2$ expression fold change for the genes significantly differentially expressed in the short time course combined model ($Firre^{OE}$ and $Firre^{RESCUE}$). **(B)** Individual gene plots showing the $\log_2$ expression fold change in the $Firre$-inducing cell lines compared to the zero-time point for all measured time points. Genes with nearby differentially accessible ATAC-seq peaks are colored in red.

Supplemental Figure S6: Chromatin modifier knockdown validation. **(A)** Maximum intensity projections of smRNA FISH images for *Shf* exon and *Firre* exon (left panel) and *Shf* intron and *Firre* exon (right panel) in *Firre^OE* and *Firre^RESCUE* mESC colonies after 0, 2, 4, and 6 hours of dox treatment. Nuclei are stained with Hoechst (blue), *Firre* exon in grey, and *Shf* exon and intron in magenta. Scale bar is 5 μm. **(B)** Relative expression of the genes targeted for GapmeR knockdown (white bars) or the non-targeting control (NTC, black bars) as measured by RT-qPCR and western blot zero and six hours after dox treatment in *Firre^OE* cells. **(C)** Relative expression

of the control non-*Firre* target gene *Tug1* in *Firre*[OE] cells transfected with non-targeting control (NTC) or targeting GapmeRs, and treated without (zero hours) or with (six hours) dox as determined by RT-qPCR.

Supplemental Figure S7: Differentially accessible ATAC-seq peaks. (A) Volcano plot showing ATAC peak signal changes between *Firre*-inducing and dox control cells. Significantly differentially accessible peaks are plotted in red with a cutoff of padj < 0.05. **(B)** Genome browser tracks of ATAC-seq peak signal at the *Rapgef4* locus in the *Firre*KO-CTL and *Firre*RESCUE line through time after dox administration. Predicted promoter (prom) and enhancer elements (enhP and enhD) are depicted.

Methods

Derivation of genetically modified mESCs

mESC derivations were performed with the assistance of Harvard's Genome Modification Facility and performed as described in (Behringer et al. 2014). Briefly, 2-5 IU of pregnant mare serum gonadotropin (PMSG) and 2-5 IU of human chorionic gonadotropin (HCG) were administered 48 hours apart by intraperitoneal injection to female *Firre*-deficient (*Firre*$^{-/-}$) mice. PMSG/HCG-treated mice were setup for timed matings (1-2 per cage) with either male mice deficient for *Firre* (*Firre*$^{-/y}$) or with male mice that contain an inducible *Firre* transgene in the *Firre*-deficient background (*Firre*$^{-/y}$; tg-*Firre*; *CAGs-rtTA3*) (generation of mice previously described (Lewandowski et al. 2019)). Female mice were checked for copulation plugs the following morning. Approximately 72 hours later, blastocysts were flushed out of the uterine horns from female mice with successful copulation plugs. Blastocysts with a detected cavity were selected for culture and expansion. Individual clones were screened by PCR genotyping for *Firre* WT, KO, transgene, and *CAGs-rtTA3* alleles as described (Lewandowski et al. 2019).

mESCs were maintained on 0.1% gelatin in KnockOut DMEM (Thermo Fisher, 10829018) supplemented with 12.5% FCS (MilliporeSigma, ES-009-B), 1X GlutaMAX supplement (Thermo Fisher, 35050061), 1X non-essential amino acids (Thermo Fisher, 11140050), 100 U/mL penicillin/streptomycin (Thermo Fisher, 15140122), 100 µM 2-mercaptoethanol (Thermo Fisher, 31350010), 1 µM PD0325901 (MilliporeSigma, PZ0162-5MG), 3 µM CHIR99021 (MilliporeSigma, SML1046-5MG), and 100 U/mL LIF (MilliporeSigma, ESG1107) at 37°C and 7.5% CO_2. Cells were routinely tested for mycoplasma contamination by PCR.

Cells were treated with 1 ng/µL dox (MilliporeSigma, D9891-5G) the day after they were seeded and one hour after the medium was changed to reduce cell stress.

RNA isolation, quantification, and quality control

Total RNA was extracted using TRIzol lysis reagent (Thermo Fisher) and purified using the RNeasy Plus Mini Kit (Qiagen) according to the manufacturer's instructions. RNA concentration was measured with a Qubit fluorometer (Invitrogen), and RNA quality was determined using a Bioanalyzer or TapeStation (Agilent Technologies).

RT-qPCR

Total RNA was reverse-transcribed using the SuperScript IV First-Strand Synthesis System with random hexamers (Thermo Fisher) according to the manufacturer's instructions. qRT-PCR samples were prepared using TaqMan probes and the TaqMan Fast Advanced Master Mix (Thermo Fisher) according to the manufacturer's instructions and run in technical triplicates on a Bio-Rad C1000 Touch Thermal Cycler. C_t values were normalized against the internal controls *Gapdh* and *ActB*. Fold differences in expression levels were calculated according to the $2^{-\Delta\Delta Ct}$ method (Livak and Schmittgen, 2001).

RNA-Seq

RNA-seq libraries were prepared from 1 µg of total RNA using the TruSeq RNA Library Prep Kit v2 (Illumina) with poly(A) selection according to the manufacturer's instructions. Multiples of twelve randomized samples were each multiplexed, pooled, and run in one HiSeq lane.

ATAC-seq

ATAC-seq libraries were prepared from 25,000 cells following the Omni-ATAC protocol (Corces et al., 2017) and using the Tagment DNA Enzyme and Buffer Kit (Illumina). Samples were pooled and run in one HiSeq lane.

Knockdowns

For knockdown assays, triplicates of 3×10^5 mESCs were seeded per well of a 6-well plate. The next day, cells were reverse transfected with 25 nM (non-targeting control, *Jarid2*, *Aebp2*, *Suz12*, *Ezh2*) or 50 nM (*Wdr5*, *G9a*) GapmeRs using Lipofectamine RNAiMAX Transfection Reagent (Thermo Fisher) according to manufacturer's instructions. After 24 hours, either no dox (0-hour time point) or 1 ng/µL dox (6-hour time point) was added without performing a media change, and cells were harvested after six hours for RNA and protein extraction.

Protein extraction and western blotting

Cells were collected by trypsinization, washed with cold PBS, and incubated in lysis buffer (50 mM Tris pH 7.5, 150 mM NaCl, 1% NP-40, 3 mM MgCl2, 5% glycerol, 1X protease inhibitors) for 20 min on ice. Protein extracts were collected as supernatant after centrifugation at 14,000xg for 10 min at 4°C, resolved on an Any kD Mini PROTEAN TGX Precast Protein Gel (Bio-Rad) and transferred to a nitrocellulose membrane (Millipore) via wet transfer in the cold room overnight. Membranes were blocked in 5% milk in TBS-T (TBS with 0.1% Tween 20) and probed with anti-JARID2 (D6M9X), anti-AEBP2 (D7C6X), anti-EZH2 (D2C9), anti-SUZ12 (D39F6), or anti-GAPDH antibody (14C10, all Cell Signaling Technology) at a dilution of 1:1000 in blocking buffer overnight at 4°C or for one hour at room temperature. The membrane was washed in TBS-T, incubated with a rabbit HRP-coupled secondary antibody (Agilent) in TBS-T for one hour, and washed again in TBS-T. Proteins were detected with the SuperSignal West Pico PLUS Chemiluminescent Substrate (Thermo Fisher) and acquired using an ImageQuant LAS 4000 (GE Healthcare).

smRNA FISH

Oligonucleotides tiling *Firre* exons (including its RRDs), *Shf* exons, and *Shf* intron 1 were designed with the Stellaris RNA FISH probe designer (LGC Biosearch Technologies, version 4.2),

labeled with Quasar 570 or 670, and produced by LGC Biosearch Technologies. Probe sequences are provided in Supplemental Table S1.

Cells were seeded on cover glasses coated with 0.1% gelatin. Cover glasses were washed twice with PBS, fixed in 3.7% formaldehyde in PBS for 10 min at room temperature, and washed again twice with PBS. Cover glasses were incubated in 70% ethanol at 4°C for at least one hour and then washed with 1 mL of wash buffer A (LGC Biosearch Technologies) at room temperature for 5 min. Cells were hybridized with 100 µL of hybridization buffer (LGC Biosearch Technologies) containing the FISH probes at a 1:100 dilution in a humid chamber at 37 °C overnight up to 16 hours. The next day, cells were washed with 1 mL of wash buffer A at 37 °C for 30 min and stained with wash buffer A containing 10 µg/mL Hoechst 33342 (Thermo Fisher) at 37 °C for 30 min. Cover glasses were washed with 1 mL of wash buffer B (LGC Biosearch Technologies) at room temperature for 5 min, mounted with ProLong Gold (Thermo Fisher) on a glass slide, and left to curate at 4°C overnight before proceeding to image acquisition.

Image acquisition was performed using a DeltaVision Elite widefield microscope with an Olympus UPlanSApo 100×/1.40-numerical aperture oil objective lens and a PCO Edge sCMOS camera. Z-stacks of 200 nm step size capturing the entire cell colony were acquired. Images were deconvolved with the built-in DeltaVision SoftWoRx Imaging software and maximum intensity projections were created using ImageJ/Fiji.

Imaris image analysis software (Bitplane) was used to quantify the distances between FISH spots. Images were prepared and FISH spots detected as previously described (Gil and Ulitsky 2019; Khong and Parker 2018). Coordinates of each *Shf* intron spot and the closest '*Firre* cloud' or the five closest *Firre* exon spots within the same cell were recorded and used to calculate their Euclidean distance from each other.

Tissue and static *Firre^{KO}* vs WT analysis

FASTQs for *Firre^{KO}* and WT mouse tissue RNA-seq experiments were retrieved from GEO GSE83631 and GSE127554 corresponding to Lewandowski et al. (Lewandowski et al. 2019) and Andergassen et al. (Andergassen et al. 2019). The static *Firre^{KO}* vs WT comparison was carried out using the *Firre* WT and KO mESCs before adding dox (n = 12). Quality control, read mapping, and quantification was performed using nf-core/rnaseq v1.4.2 (Ewels et al. 2020). Reads were mapped to mm10 and quantified with Salmon v1.5.2 (Patro et al. 2017) using the Gencode M25 annotation. Significantly differentially expressed genes were calculated using DESeq2 (Love et al. 2014) (comparing *Firre^{KO}* vs WT) with an adjusted p-value cutoff of 0.05 and a fold change cutoff of 1.5 on the shrunken $\log_2$ fold change values from DESeq2.

Firre induction time course quantification

Quality control, read mapping, and quantification were performed using nf-core/rnaseq v1.4.2. Reads were mapped to mm10 and quantified with Salmon using the Gencode M25 annotation.

Doxycycline control RNA-seq

Differential expression was calculated using DEseq2 and the Wald test to compare each time point back to the zero-time point. The rescue and overexpression dox control lines and the corresponding short and long-time courses were analyzed separately. Cutoffs to call differentially expressed genes were made using an adjusted p-value cutoff of 0.05 and whether a gene achieved a fold change greater than 1.5 at any time point (calculated using the shrunken $\log_2$ fold change).

Firre induction time course analysis

Differential expression was calculated with DEseq2 using the likelihood ratio test to compare a full model which contains the *Firre* induction/time point interaction term to a reduced model lacking that term. Calling differentially expressed genes proceeded in two steps: 1. A cutoff was made on the adjusted p-values from the likelihood ratio test ($p < 0.05$), filtering to genes that changed significantly in time as compared to the dox control. 2. A fold change cutoff was made by comparing the expression through time back to the zero-time point (without any comparison to the dox control line). A gene was called differentially expressed (responding to *Firre*) if it achieved a fold change greater than 1.5 in any time point as measured by the shrunken $\log_2$ fold change from DEseq2.

The combined model using all four cell lines was analyzed in the same manner with the same thresholding strategy, but the full and reduced models contained an additional indicator variable for the *Firre* genotype (WT or KO).

The $\log_2$ fold changes shown in the heatmaps are the shrunken $\log_2$ fold changes reported by DEseq2.

Identification of rescued genes

A gene was considered rescued, if: 1. It was significantly differentially expressed in the static *Firre*[KO] vs WT comparison and in the *Firre*[RESCUE] induction experiment, and 2. If the fold change was reciprocal to within 20% of the fold change value in the *Firre*[KO] vs WT.

ATAC-seq time course analysis

Quality control, read mapping, and peak calling were carried out using nf-core/atacseq v1.2.0. Reads were mapped to mm10 and the mm10 v2 blacklist from ENCODE was used. Within the nf-core/atacseq pipeline MACS2 v2.2.7.1 (Feng et al. 2012; Zhang et al. 2008) was used to call peaks in each experiment separately. Consensus peaks were called using the peak regions

present in any single experiment. Reads within peaks were calculated using featureCounts. Differentially accessible peaks were calculated using DEseq2 to compare the read counts in the samples in which *Firre* was present (>0 time point in the *Firre^{RESCUE}* and *Firre^{OE}* lines) and the samples in which *Firre* was absent (WT and *Firre^{KO-CTL}* lines, and zero-time points). A cutoff was made using the adjusted p-value ($p < 0.05$).

Data access

All sequencing files have been deposited at Gene Expression Omnibus under the accession number GSE202406.

Competing interest statement

The authors declare that they have no conflict of interest.

Acknowledgements

We would like to thank Gabrijela Dumbović and Aaron Nguyen for their help with smRNA FISH, Nydia Chang for assistance in deriving the mouse ES cells, and Anthony Khong for help with the Imaris software. We are grateful to Roy Parker and Carolyn Decker for access to, and training on, the DeltaVision Elite microscope. We acknowledge the support of Theresa Nahreini and Nicole Kethley in the Biochemistry Cell Culture Facility, Joe Dragavon in the BioFrontiers Advanced Light Microscopy Core, and Amber L. Scott in the BioFrontiers Sequencing Core. J.L.R. is an HHMI Faculty Scholar. C.M. was supported by the Deutsche Forschungsgemeinschaft (DFG) (MU 4462/1-1). This research was supported by NIH/NIGMS grant P01 GM099117 to J.L.R.

References

Abascal F, Acosta R, Addleman NJ, Adrian J, Afzal V, Aken B, Akiyama JA, Jammal OA, Amrhein H, Anderson SM, et al. 2020. Expanded encyclopaedias of DNA elements in the human and mouse genomes. *Nature* 583: 699–710.

Abe Y, Kikuchi A, Kobayashi S, Wakusawa K, Tanaka S, Inui T, Kunishima S, Kure S, Haginoya K. 2014. Xq26.1-26.2 gain identified on array comparative genomic hybridization in bilateral periventricular nodular heterotopia with overlying polymicrogyria. *Developmental Medicine & Child Neurology* n/a-n/a.

Ahler E, Sullivan WJ, Cass A, Braas D, York AG, Bensinger SJ, Graeber TG, Christofk HR. 2013. Doxycycline Alters Metabolism and Proliferation of Human Cell Lines. Plos One 8: e64561.

Allen MA, Andrysik Z, Dengler VL, Mellert HS, Guarnieri A, Freeman JA, Sullivan KD, Galbraith MD, Luo X, Kraus WL, et al. 2014. Global analysis of p53-regulated transcription identifies its direct targets and unexpected regulatory mechanisms. *Elife* 3: e02200.

Andergassen D, Rinn JL. 2021. From genotype to phenotype: genetics of mammalian long non-coding RNAs in vivo. *Nat Rev Genet* 23: 229–243.

Andergassen D, Smith ZD, Lewandowski JP, Gerhardinger C, Meissner A, Rinn JL. 2019. In vivo Firre and Dxz4 deletion elucidates roles for autosomal gene regulation. *Elife* 8: e47214.

Anderson KM, Anderson DM, McAnally JR, Shelton JM, Bassel-Duby R, Olson EN. 2016. Transcription of the non-coding RNA upperhand controls Hand2 expression and heart development. *Nature* 539: 433–436.

Barutcu AR, Maass PG, Lewandowski JP, Weiner CL, Rinn JL. 2018. A TAD boundary is preserved upon deletion of the CTCF-rich Firre locus. *Nature communications* 9: 1444.

Behringer R, Gertsenstein M, Nagy KV, Nagy A. 2014. Manipulating the Mouse Embryo: A Laboratory Manual, 4th ed. Cold Spring Harbor Press, New York.

Birney E, Stamatoyannopoulos JA, Dutta A, Guigó R, Gingeras TR, Margulies EH, Weng Z, Snyder M, Dermitzakis ET, Stamatoyannopoulos JA, et al. 2007. Identification and analysis of functional elements in 1% of the human genome by the ENCODE pilot project. *Nature* 447: 799–816.

Brivanlou AH, Darnell JE. 2002. Signal Transduction and the Control of Gene Expression. *Science* 295: 813–818.

Cabili MN, Trapnell C, Goff L, Koziol M, Tazon-Vega B, Regev A, Rinn JL. 2011. Integrative annotation of human large intergenic noncoding RNAs reveals global properties and specific subclasses. *Genes & Development* 25: 1915–1927.

Carpenter S, Aiello D, Atianand MK, Ricci EP, Gandhi P, Hall LL, Byron M, Monks B, Henry-Bezy M, Lawrence JB, et al. 2013. A Long Noncoding RNA Mediates Both Activation and Repression of Immune Response Genes. *Science* 341: 789–792.

Charoensawan V, Martinho C, Wigge PA. 2015. "Hit-and-run": Transcription factors get caught in the act. *Bioessays* 37: 748–754.

Consortium EP. 2012. An integrated encyclopedia of DNA elements in the human genome. *Nature Publishing Group* 489: 57–74.

Consortium TEP. 2004. The ENCODE (ENCyclopedia Of DNA Elements) Project. *Science* 306: 636–640.

Derrien T, Johnson R, Bussotti G, Tanzer A, Djebali S, Tilgner H, Guernec G, Martin D, Merkel A, Knowles DG, et al. 2012. The GENCODE v7 catalog of human long noncoding RNAs: analysis of their gene structure, evolution, and expression. *Genome Research* 22: 1775–1789.

Dimitrova N, Zamudio JR, Jong RM, Soukup D, Resnick R, Sarma K, Ward AJ, Raj A, Lee JT, Sharp PA, et al. 2014. LincRNA-p21 Activates p21 In cis to Promote Polycomb Target Gene Expression and to Enforce the G1/S Checkpoint. *Molecular Cell* 1–14.

Elling R, Robinson EK, Shapleigh B, Liapis SC, Covarrubias S, Katzman S, Groff AF, Jiang Z, Agarwal S, Motwani M, et al. 2018. Genetic Models Reveal cis and trans Immune-Regulatory Activities for lincRNA-Cox2. *CellReports* 25: 1511-1524.e6.

Engreitz JM, Haines JE, Perez EM, Munson G, Chen J, Kane M, McDonel PE, Guttman M, Lander ES. 2016. Local regulation of gene expression by lncRNA promoters, transcription and splicing. *Nature Publishing Group* 539: 452–455.

Ewels PA, Peltzer A, Fillinger S, Patel H, Alneberg J, Wilm A, Garcia MU, Tommaso PD, Nahnsen S. 2020. The nf-core framework for community-curated bioinformatics pipelines. *Nat Biotechnol* 38: 276–278.

Fanucchi S, Fok ET, Dalla E, Shibayama Y, Börner K, Chang EY, Stoychev S, Imakaev M, Grimm D, Wang KC, et al. 2018. Immune genes are primed for robust transcription by proximal long noncoding RNAs located in nuclear compartments. *Nat Genet* 51: 138–150.

Feng J, Liu T, Qin B, Zhang Y, Liu XS. 2012. Identifying ChIP-seq enrichment using MACS. *Nat Protoc* 7: 1728–1740.

Gil N, Ulitsky I. 2019. Regulation of gene expression by cis-acting long non-coding RNAs. *Nat Rev Genet* 21: 102–117.

Groff AF, Sanchez-Gomez DB, Soruco MML, Gerhardinger C, Barutcu AR, Li E, Elcavage L, Plana O, Sanchez LV, Lee JC, et al. 2016. In Vivo Characterization of Linc-p21 Reveals Functional cis-Regulatory DNA Elements. *CellReports* 16: 2178–2186.

Guttman M, Donaghey J, Carey BW, Garber M, Grenier JK, Munson G, Young G, Lucas AB, Ach R, Bruhn L, et al. 2011. lincRNAs act in the circuitry controlling pluripotency and differentiation. *Nature* 477: 295–300.

Guttman M, Rinn JL. 2012. Modular regulatory principles of large non-coding RNAs. *Nature* 482: 339–346.

Hacisuleyman E, Goff LA, Trapnell C, Williams A, Henao-Mejia J, Sun L, McClanahan P, Hendrickson DG, Sauvageau M, Kelley DR, et al. 2014. Topological organization of multichromosomal regions by the long intergenic noncoding RNA Firre. *Nature Structural & Molecular Biology* 21: 198–206.

Hacisuleyman E, Shukla CJ, Weiner CL, Rinn JL. 2016. Function and evolution of local repeats in the Firre locus. *Nature communications* 7: 11021.

Huarte M, Guttman M, Feldser D, Garber M, Koziol MJ, Kenzelmann-Broz D, Khalil AM, Zuk O, Amit I, Rabani M, et al. 2010. A large intergenic noncoding RNA induced by p53 mediates global gene repression in the p53 response. *Cell* 142: 409–419.

Khong A, Parker R. 2018. mRNP architecture in translating and stress conditions reveals an ordered pathway of mRNP compaction. *J Cell Biol* 217: 4124–4140.

Krawczyk M, Emerson BM. 2014. p50-associated COX-2 extragenic RNA (PACER) activates COX-2 gene expression by occluding repressive NF-κB complexes. *Elife* 3: e01776.

Lammers NC, Kim YJ, Zhao J, Garcia HG. 2020. A matter of time: Using dynamics and theory to uncover mechanisms of transcriptional bursting. *Curr Opin Cell Biol* 67: 147–157.

Lee TI, Young RA. 2000. TRANSCRIPTION OF EUKARYOTIC PROTEIN-CODING GENES. *Genetics* 34: 77–137.

Lewandowski JP, Lee JC, Hwang T, Sunwoo H, Goldstein JM, Groff AF, Chang NP, Mallard W, Williams A, Henao-Meija J, et al. 2019. The Firre locus produces a trans-acting RNA molecule that functions in hematopoiesis. *Nat Commun* 10: 5137.

Loewer S, Cabili MN, Guttman M, Loh Y-H, Thomas K, Park IH, Garber M, Curran M, Onder T, Agarwal S, et al. 2010. Large intergenic non-coding RNA-RoR modulates reprogramming of human induced pluripotent stem cells. *Nature Genetics* 42: 1113–1117.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* 15: 550.

Maamar H, Cabili MN, Rinn J, Raj A. 2013. linc-HOXA1 is a noncoding RNA that represses Hoxa1 transcription in cis. *Genes & Development* 27: 1260–1271.

Maass PG, Rump A, Schulz H, Stricker S, Schulze L, Platzer K, Aydin A, Tinschert S, Goldring MB, Luft FC, et al. 2012. A misplaced lncRNA causes brachydactyly in humans. *The Journal of clinical investigation* 122: 3990–4002.

Mahat DB, Salamanca HH, Duarte FM, Danko CG, Lis JT. 2016. Mammalian Heat Shock Response and Mechanisms Underlying Its Genome-wide Transcriptional Regulation. *Mol Cell* 62: 63–78.

Mitchell PJ, Tjian R. 1989. Transcriptional Regulation in Mammalian Cells by Sequence-Specific DNA Binding Proteins. *Science* 245: 371–378.

Miyagawa R, Tano K, Mizuno R, Nakamura Y, Ijiri K, Rakwal R, Shibato J, Masuo Y, Mayeda A, Hirose T, et al. 2012. Identification of cis- and trans-acting factors involved in the localization of MALAT-1 noncoding RNA to nuclear speckles. *RNA* 18: 738–751.

Morris KV, Mattick JS. 2014. The rise of regulatory RNA. *Nature Reviews Genetics* 15: 423–437.

Much C, Smallegan MJ, Hwang T, Hanson SD, Dumbovic G, Rinn JL. 2022. Evolutionary divergence of Firre localization and expression. *Rna* rna.079070.121.

Paralkar VR, Taborda CC, Huang P, Yao Y, Kossenkov AV, Prasad R, Luan J, Davies JOJ, Hughes JR, Hardison RC, et al. 2016. Unlinking an lncRNA from Its Associated cis Element. *Molecular Cell* 62: 104–110.

Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. 2017. Salmon provides fast and bias-aware quantification of transcript expression. *Nat Methods* 14: 417–419.

Rinn JL, Chang HY. 2012. Genome regulation by long noncoding RNAs. *Annual review of biochemistry* 81: 145–166.

Rinn JL, Chang HY. 2020. Long Noncoding RNAs: Molecular Modalities to Organismal Functions. *Annu Rev Biochem* 89: 283–308.

Sasse SK, Gruca M, Allen MA, Kadiyala V, Song T, Gally F, Gupta A, Pufall MA, Dowell RD, Gerber AN. 2019. Nascent transcript analysis of glucocorticoid crosstalk with TNF defines primary and cooperative inflammatory repression. *Genome Res* 29: 1753–1765.

Stewart-Ornstein J, Nelson C, DeRisi J, Weissman JS, El-Samad H. 2013. Msn2 Coordinates a Stoichiometric Gene Expression Program. *Curr Biol* 23: 2336–2345.

Tan JY, Smith AAT, Silva MF da, Matthey-Doret C, Rueedi R, Sönmez R, Ding D, Kutalik Z, Bergmann S, Marques AC. 2017. cis-Acting Complex-Trait-Associated lincRNA Expression Correlates with Modulation of Chromosomal Architecture. *CellReports* 18: 2280–2288.

Tong C, Yin Y. 2021. Localization of RNAs in the nucleus: cis- and trans- regulation. *Rna Biol* 18: 1–14.

Wang KC, Chang HY. 2011. Molecular mechanisms of long noncoding RNAs. *Molecular Cell* 43: 904–914.

Wang Z, Deng W. 2022. Dynamic transcription regulation at the single-molecule level. *Dev Biol* 482: 67–81.

Weikum ER, Knuesel MT, Ortlund EA, Yamamoto KR. 2017. Glucocorticoid receptor control of transcription: precision and plasticity via allostery. *Nat Rev Mol Cell Bio* 18: 159–174.

Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nusbaum C, Myers RM, Brown M, Li W, et al. 2008. Model-based analysis of ChIP-Seq (MACS). *Genome Biol* 9: R137.