Bacterial sterol methylation confounds eukaryotic biomarker interpretations

Malory O. Brown[a] and Babatunde O. Olagunju[b], José-Luis Giner[b], and Paula V.
Welander[a]

[a]Department of Earth System Science, Stanford University, Stanford, CA 94305

[b]Department of Chemistry, State University of New York-Environmental Science and
Forestry, Syracuse, NY 13210

Sterol lipids are required by most eukaryotes and are readily preserved as sterane

molecular fossils. These geologic steranes are broadly interpreted as biomarkers for

ancient eukaryotes[1,2] although diverse bacteria also produce sterols[3]. Steranes with

side-chain methylations can act as more specific biomarkers[4] if their sterol precursors

are limited to particular extant eukaryotes and are absent in bacteria. An abundance

of one such sterane, 24-isopropylcholestane, in late Neoproterozoic rocks has been

attributed to marine demosponges and potentially represents the earliest evidence for

animals on Earth[5]. However, debates over this interpretation[6–14] continue given the

potential for alternative sources of 24-isopropylcholestane and the lack of

experimental evidence demonstrating the function of enzymes that methylate sterols

to give the 24-isopropyl side-chain. Here we show that sterol methyltransferases from

both sponges and bacteria are functional and identify three bacterial

methyltransferases each capable of sequential methylations resulting in the 24-

isopropyl sterol side-chain. We identified two of these propylating enzymes in a

demosponge metagenome suggesting bacterial symbionts contribute to 24-isopropyl

sterol biosynthesis in demosponges. Our results demonstrate yet-uncultured bacteria

have the genomic capacity to synthesize side-chain alkylated sterols and should

therefore be considered when interpreting side-chain alkylated sterane biomarkers

in the rock record.

Main text

According to the sponge biomarker hypothesis, two side-chain alkylated sterane

biomarkers, 24-isopropylcholestane (24-ipc) and 26-methylstigmastane (26-mes), indicate

marine sponges of the Demospongiae class[5,15] (Fig. 1**a**). The co-occurrence of 24-ipc and

26-mes in the late Neoproterozoic (660-540 Ma)[5,15], preceding the earliest sponge

macrofossils by ~100 million years[7], may represent the first evidence of animals in the

geologic record. However, debate over this interpretation continues for several reasons[6-14]

including the potential for alternative sources. Demosponges are the only extant organisms

known to contain sterols with the 24-ipc and 26-mes side-chain structures as their major

sterols[15-21], but pelagophytes, which produce 24-*n*-propylcholesterol as their major sterol,

also contain 24-isopropyl sterols[22-25]. Trace amounts of 24-ipc and 26-mes were also

reported in hydrogenated lipid extracts from Rhizaria[10], although attempts to replicate these

results were not successful[11]. In laboratory pyrolysis experiments under conditions that

may mimic diagenesis, 24-ipc and 26-mes form from certain C_{29} sterols suggesting

Neoproterozoic 24-ipc and 26-mes could have formed abiotically from algal sterols[13,14].

Further, microbial symbionts, including bacteria from diverse phyla, can constitute up to

40% of sponge biomass[26]. Bacterial symbionts may therefore contribute to 24-isopropyl

and 26-methyl sterol biosynthesis in sponges, although bacterial enzymes that methylate

the sterol side-chain have not been previously identified.

One approach to constrain uncertainties in the sponge biomarker hypothesis is to

identify and characterize the enzymes responsible for these distinct alkylations at C-24 and

C-26. Sterol 24-C-methyltransferases (SMTs) alkylate the C-24 position at an isolated

double bond in an *S*-adenosylmethionine (SAM) dependent mechanism[27] and have been

characterized extensively in fungi and plants. Bioinformatic analyses addressing this

reaction in sponges prompted the hypotheses that at least two SMT copies in a genome are

required to produce 24-propyl sterols, and that sponge SMTs can perform multiple

methylations at C-24[28]. However, no experimental evidence verifying this activity by any

sponge-derived SMT currently exists, and SMTs that produce 24-propyl sterols have not

been identified in any extant organism.

Sponge SMTs are functional. Our first goal was to express putative sponge SMTs

and verify methylation activity in vitro. We chose eight SMT homologs identified in

genomes and transcriptomes from eight sponges of the Demospongiae,

Homoscleromorpha, and Calcarea classes, each of which have been shown to contain 24-

ethyl sterols at the species[29–32] or genus[32–34] level. We tested each SMT with two potential

substrates, desmosterol or 24-methylenecholesterol, and analyzed sterol products using gas

chromatography-mass spectrometry (GC-MS). All eight of the sponge SMTs methylated

desmosterol to produce 24-methyl sterols including 24-methylenecholesterol and

(epi)codisterol (Table 1). Six SMTs also methylated 24-methylenecholesterol to 24-ethyl

sterols including fucosterol, isofucosterol, and (epi)clerosterol (Fig. 1**d**). None of the

sponge SMTs produced 24-propyl sterols.

Bacterial SMTs produce 24-isopropyl sterols. Our analyses of sponge symbiont

metagenomes revealed several SMT homologs, many of which occurred in gene clusters

with homologs of other known sterol biosynthesis genes including the bacterial C-4

demethylases *sdmA* and *sdmB*[35] (Fig. 2). We therefore hypothesized bacterial symbionts

may contribute to side-chain alkylation in sponges. We chose 10 putative bacterial SMTs

identified in metagenomes from four demosponge species for analysis. Eight symbiont

SMTs methylated desmosterol to 24-methyl sterols including (epi)codisterol, 24-

methylenecholesterol, and 24-methyldesmosterol (Table 1, Fig. 1**c**). Three of these SMTs

also methylated 24-methylenecholesterol to 24-ethyl sterols including fucosterol,

isofucosterol, and (epi)clerosterol (Fig. 1**d**). Two SMTs identified in a metagenome from

Aplysina aerophoba, a demosponge known to contain trace 24-isopropyl sterols[21], also

methylated 24-methylenecholesterol to two distinct C_{30} sterols. The first symbiont SMT

produced 24*S*-24-isopropylcholest-5,25-dienol while the second produced 24-

isopropylcholest-5,24-dienol. Both propylating SMTs were capable of all three

methylation steps required to synthesize the 24-ipc side-chain via a single protein. One

propylating SMT was identified in a sterol biosynthesis gene cluster suggesting a yet-

unknown bacterial sponge symbiont has the genomic capacity to synthesize 24-isopropyl

sterols de novo (Fig. 2).

Identification of bacterial symbiont SMTs capable of side-chain alkylation led us

to question if bacteria from environments outside a sponge host could also produce side-

chain alkylated sterols. Through additional searches, we identified SMT homologs in a

variety of environmental metagenomes and metagenome-assembled genomes (MAGs).

We tested 14 of these SMT homologs from marine, freshwater, thermal spring, and coral

reef environments. Twelve environmental SMTs alkylated the sterol side-chain to produce

24-methyl and/or 24-ethyl sterols (Table 1). One SMT, identified in a drinking water filter

MAG assigned to the phylum Chlamydiae, produced both 24*S*-24-isopropylcholest-5,25-

dienol and 24-isopropylcholest-5,24-dienol (Fig. 1**a**).

The identification of functional metagenomic SMTs likely derived from bacteria,

including three that produce 24-isopropyl sterols, is significant given that no bacterial

proteins had previously been shown to alkylate the sterol side-chain. To confirm that these

SMTs are truly of bacterial origin, we performed phylogenetic analyses of the 24

metagenomic SMTs we tested. In our maximum-likelihood tree of SMT proteins (Fig. 3**a**),

the majority of the metagenomic SMTs are distinct and more closely related to each other

than to the SMTs of eukaryotes known to produce side-chain alkylated sterols. Further, 23

of the 24 putative bacterial SMTs, including those from sponge metagenomes, form two

clades separate from the sponge SMT clade. It is therefore unlikely that these metagenomic

SMTs were acquired from sponges via horizontal gene transfer. However, the observed

bootstrap values were not robust enough to draw any additional evolutionary conclusions,

nor to eliminate any remaining uncertainty of the bacterial origin of these SMTs. We

therefore used the recently developed classifier Whokaryote, which predicts whether a

metagenomic contig is likely eukaryotic or prokaryotic based on intergenic distance, gene

density, gene length, and k-mer frequencies. Whokaryote predicts all 19 of the contigs

containing the metagenomic SMTs tested here to be of prokaryotic origin (Extended Data

Table 2).

The genomic context of the SMTs we tested provides additional evidence of their

bacterial origin. Of the 24 metagenomic SMTs, nine were in gene clusters or MAGs with

a homolog of oxidosqualene cyclase (OSC), a key enzyme in sterol biosynthesis

responsible for cyclizing oxidosqualene to lanosterol[3]. OSCs cluster into robust

phylogenetic clades with high bootstrap support[36]. Thus, phylogenetic analyses of OSCs

associated with the SMTs we tested could confirm their bacterial origin. In our maximum-

likelihood tree of OSC proteins (Fig. 3**b**), we see that seven of the OSCs associated with

the SMTs analysed here, including four from demosponge metagenomes, fall into the

Group 1 clade which contains only bacterial OSCs. The remaining 2 OSCs, from soil and

hot spring sediment metagenomes, fall into bacterial Group 2. The position of these OSCs

provides additional evidence that the metagenomic SMTs we tested originate from a

bacterial source.

Side-chain propylation mechanism. It was surprising to observe that one SMT

was capable of propylating the sterol side-chain on its own as it has been hypothesized that

at least two SMTs would be required for this reaction[28]. To better understand this

unexpected biosynthetic pathway, we performed additional in vitro reactions with two of

the bacterial propylating SMTs using [13]C-labeled sterol substrates and analyzed the

resulting products by nuclear magnetic resonance (NMR; Extended Data Table 3). In

experiments with [28-[13]C] 24-methylenecholesterol as the substrate of the *A. aerophoba*

symbiont SMT, the major C_{29} products were confirmed as isofucosterol and fucosterol

present in a 7:1 ratio. Trace epiclerosterol, but not clerosterol, was also detected. The

product of triple side-chain methylation was confirmed as *24S*-24-isopropylcholesta-5,25-

dienol. The label was found exclusively at C-25, and the *24R* isomer was absent (Extended

Data Fig. 2**a**). Fucosterol was not a substrate for the *A. aerophoba* symbiont SMT, but [29-

[13]C] isofucosterol yielded *24S*-24-isopropylcholesta-5,25-dienol with the label distributed

in a 4:1 label ratio between C-27 and C-26 (Extended Data Fig. 2**b**). This outcome shows

interconversion of the C-28 and C-25 cations by hydride-shifts is not involved in the

biosynthesis of this compound, and deprotonation of the initially formed cation is

unspecific.

The same set of reactions with the propylating Chlamydiae MAG SMT gave similar

results. Fucosterol was not a substrate and epiclerosterol, but not clerosterol, was detected.

However, minor differences were measured between these two bacterial SMTs. In

experiments with [29-^{13}C] isofucosterol and the Chlamydiae MAG SMT, the ratio of

isofucosterol and fucosterol was 30:1, and the ratio of label at C-27/C-26 of *24S*-24-

isopropylcholesta-5,25-dienol was 5:1. Additionally, the C_{30} Δ^{23} and Δ^{24} isomers, 24-

isopropylcholesta-5,23-dienol and 24-isopropylcholesta-5,24-dienol, were also detected.

The label was detected in the Z-vinylic methyl of 24-isopropylcholesta-5,24-dienol and in

one of the two methyls of the Z-isopropyl group of 24-isopropylcholesta-5,23-dienol.

These results provide evidence for one hydride-shift to generate the C-24 cation. The ratio

of *24S*-24-isopropylcholesta-5,25-dienol and its Δ^{24} and Δ^{23} isomers was estimated to be

20:4:1, respectively. Though minor mechanistic differences between these two propylating

SMTs exist, the predominant pathway to 24-isopropyl sterols observed here proceeds from

desmosterol to 24-methylenecholesterol to isofucosterol to 24*S*-24-isopropylcholest-5,25-

dienol (Fig. 1**b**).

Discussion. The results presented here experimentally verify that sponges do

indeed encode functional SMTs capable of sequentially methylating desmosterol to 24-

methylenecholesterol to 24-ethyl sterols, similarly to SMTs recently identified in annelid

worms[37]. However, as in Annelida[38], this activity was inconsistent among Porifera as the

SMTs from *Oscarella carmela* (Homoscleromorpha) and *Sycon ciliatum* (Calcarea) did

not sequentially methylate sterols, contrary to previous hypotheses[28]. Sponge-derived

SMTs that produce 24-ethyl sterols may therefore be restricted to certain species, although

we recognize these SMTs may require a different substrate than we provided in our assays.

Substrate specificity could also explain why no sponge SMT produced 24-isopropyl sterols

in our experiments, although this result was unsurprising given that we did not test SMTs

from sponges known to contain propylated sterols due to the lack of available genomic

data. However, we did show that bacterial symbionts have the genomic capacity to

synthesize these propylated lipids. Previous studies have provided evidence of de novo

sterol biosynthesis in demosponges[39–41], but the experiments were performed prior to the

development of 16S community sequencing using cell-free sponge extracts or whole

sponge tissue from which the presence of bacterial symbiont-derived enzymes cannot be

excluded. Additionally, transcripts for several key sterol biosynthesis genes in the

demosponge *Vaceletia* sp. were derived from bacteria[42]. Both sponge and bacterial

symbiont proteins may therefore be required to produce 24-isopropyl sterols in vivo.

However, the metagenomic gene clusters containing functional SMTs identified here,

along with steroid gene clusters containing SMT homologs previously identified in

bacterial MAGs associated with demosponges of the *Ircinia* genus[43], suggest bacterial

symbionts may produce 24-isopropyl sterols de novo without demosponge input. Our work

highlights the need for robust sequencing of both the sponge host and its microbiome

coupled to sterol analysis in demosponge species that contain 24-isopropyls as their major

sterols. This may reveal sponge-derived SMTs sufficient for the biosynthesis of these lipids

and confirm sponges and/or their bacterial symbionts as potential sources of 24-ipc

biomarkers.

The identification of three bacterial SMTs each capable of producing the 24-

isopropyl side-chain via a single protein is also significant. Molecular clock analyses

suggest demosponges were the likely source of Cryogenian 24-ipc biomarkers given that

demosponges acquired a second SMT copy prior to their deposition, while pelagophytes

did not acquire a third SMT copy until the Phanerozoic[28]. Additionally, researchers have

argued against Rhizaria as a source of 24-isopropyl sterols given that *Bigelowiella natans*

encodes only two SMT copies and is thus unlikely to synthesize 24-propyl sterols[11]. These

interpretations rely on the hypothesis that multiple SMT copies are required to produce 24-

isopropyl sterols. However, our results demonstrating one SMT is sufficient for side-chain

propylation make SMT copy number an unreliable predictor of an organism's ability to

produce 24-propyl sterols. Now that propylating SMTs have been identified, site-directed

mutagenesis as performed with yeast[44,45] and plant[46] SMTs may reveal specific amino acid

changes that allow an SMT protein to perform a third methylation. If such key residues are

identified, they could act as a more reliable predictor of 24-isopropyl sterol biosynthesis

directly from sequencing data than SMT copy number alone.

In addition to its use as a demosponge biomarker, 24-ipc has also been used to

indicate marine input into terrestrial systems[47] given that 24-ipc precursor sterols are only

found in marine demosponges and algae. However, the production of 24-isopropyl sterols

by an SMT identified in a freshwater Chlamydiae MAG suggests these sterols may occur

in freshwater environments. Chlamydiae have not been shown to produce sterols, and

homologs of other sterol biosynthesis genes were absent in the Chlamydiae MAG. It is

therefore unlikely that this bacterium synthesizes 24-isopropyl sterols de novo. However,

some endosymbiotic Chlamydiae recruit sterols from their eukaryotic host into their

membranes[48]. Certain freshwater Chlamydiae may therefore methylate host-acquired

sterols to give the 24-isopropyl side-chain, which could explain the presence of 22-

dehydro-24-isopropylcholestanol throughout a freshwater sediment core from Motte Lake,

France[49]. We would therefore exercise caution when interpreting the presence of 24-ipc as

evidence for seawater incursion events.

Finally, although we identified SMTs that produce 24-isopropyl sterols, proteins

that methylate sterols to give other geologically important side-chain structures remain

undiscovered. Identification of enzymes that methylate sterols at the C-26 position to give

the 26-mes side-chain structure remains especially important when considering the

biomarker evidence for Precambrian sponges. SMTs from sponges and bacteria produced

(epi)codisterol suggesting both groups have the genomic capacity to produce substrates for

hypothetical 26-SMTs[39,50]. However, it remains to be seen if sponges and bacteria, sponge-

associated and otherwise, encode SMTs capable of methylation at C-26. Once identified,

in-depth kinetics studies with these enzymes and multiple substrates may reveal if SMTs

from sponges and/or bacteria are likely to produce 24-ipc and 26-mes sterol precursors in

ratios corresponding to the C_{30} sterane distributions observed in the Cryogenian. Verified

side-chain alkylated sterol biosynthesis in a bacterial isolate along with identification of

any unique isotopic signatures resulting from this reaction could also confirm bacteria as a

source these biomarkers. Until then, the co-occurrence of 24-ipc and 26-mes in high

relative abundance continues to provide compelling evidence for derived sponges in the

Cryogenian. However, sterol side-chain methylation by symbiotic and free-living bacteria

should now be considered when interpreting the presence of ergostane, stigmastane, and

24-isopropylcholestane in the geologic record.

Main references

1. Summons, R. E., Bradley, A. S., Jahnke, L. L. & Waldbauer, J. R. Steroids,

triterpenoids and molecular oxygen. *Philos Trans R Soc Lond B Biol Sci* **361**, 951–

968 (2006).

2. Bobrovskiy, I. *et al.* Ancient steroids establish the Ediacaran fossil *Dickinsonia* as
one of the earliest animals. *Science* **361**, 1246–1249 (2018).

3. Wei, J. H., Yin, X. & Welander, P. V. Sterol synthesis in diverse bacteria. *Front*
Microbiol **7**, 990 (2016).

4. Brocks, J. J. *et al.* The rise of algae in Cryogenian oceans and the emergence of
animals. *Nature* **548**, 578–581 (2017).

5. Love, G. D. *et al.* Fossil steroids record the appearance of Demospongiae during the
Cryogenian period. *Nature* **457**, 718–721 (2009).

6. Antcliffe, J. B. Questioning the evidence of organic compounds called sponge
biomarkers. *Palaeontology* **56**, 917–925 (2013).

7. Antcliffe, J. B., Callow, R. H. T. & Brasier, M. D. Giving the early fossil record of
sponges a squeeze: the early fossil record of sponges. *Biol Rev* **89**, 972–1004 (2014).

8. Love, G. D. & Summons, R. E. The molecular record of Cryogenian sponges - a
response to Antcliffe (2013). *Palaeontology* **58**, 1131–1136 (2015).

9. Antcliffe, J. B. The oldest compelling evidence for sponges is still early Cambrian in
age - reply to Love and Summons (2015). *Palaeontology* **58**, 1137–1139 (2015).

10. Nettersheim, B. J. *et al.* Putative sponge biomarkers in unicellular Rhizaria question
an early rise of animals. *Nat Ecol Evol* **3**, 577–581 (2019).

11. Love, G. D. *et al.* Sources of C30 steroid biomarkers in Neoproterozoic–Cambrian
rocks and oils. *Nat Ecol Evol* **4**, 34–36 (2020).

12. Hallmann, C. *et al.* Reply to: Sources of C30 steroid biomarkers in Neoproterozoic–
Cambrian rocks and oils. *Nat Ecol Evol* **4**, 37–39 (2020).

13. Bobrovskiy, I. *et al.* Algal origin of sponge sterane biomarkers negates the oldest

evidence for animals in the rock record. *Nat Ecol Evol* **5**, 165–168 (2021).

14. van Maldegem, L. M. *et al.* Geological alteration of Precambrian steroids mimics

early animal signatures. *Nat Ecol Evol* **5**, 169–173 (2021).

15. Zumberge, J. A. *et al.* Demosponge steroid biomarker 26-methylstigmastane provides

evidence for Neoproterozoic animals. *Nat Ecol Evol* **2**, 1709–1714 (2018).

16. Hofheinz, W. & Oesterhelt, G. 24-isopropylcholesterol and 22-dehydro-24-

isopropylcholesterol, novel sterols from a sponge. *Helv Chim Acta* **62**, 1307–1309

(1979).

17. Do, H. Q. *et al.* Unusual sterolic mixture, and 24-isopropylcholesterol, from the

sponge *Ciocalypta* sp. reduce cholesterol uptake and basolateral secretion in Caco-2

cells. *J Cell Biochem* **106**, 659–665 (2009).

18. Bortolotto, M., Braekman, J. C., Daloze, D. & Tursch, B. Chemical studies of marine

invertebrates. XXXVI. Strongylosterol, a novel C-30 sterol from the sponge

Strongylophora durissima Dendy. *Bull Soc Chim Belges* **87**, 539–543 (1978).

19. Barnathan, G. *et al.* Unusual sterol composition and classification of three marine

sponge families. *Boll Mus Ist Biol Univ Genova* **68**, 201–208 (2004).

20. Giner, J.-L. Biosynthesis of marine sterol side chains. *Chem Rev* **93**, 1735–1752

(1993).

21. Zumberge, J. A. A lipid biomarker investigation tracking the evolution of the

Neoproterozoic marine biosphere and the rise of eukaryotes. (University of

California, Riverside, 2019).

22. Kokke, W. C. M. C., Shoolery, J. N., Fenical, W. & Djerassi, C. Biosynthetic studies

of marine lipids. 4. Mechanism of side chain alkylation in E-24-

propylidenecholesterol by a Chrysophyte alga. *J Org Chem* **49**, 3742–3752 (1984).

23. Giner, J.-L. & Djerassi, C. Biosynthetic studies of marine lipids. 31. Evidence for a

protonated cyclopropyl intermediate in the biosynthesis of 24-propylidenecholesterol.

J Am Chem Soc **113**, 1386–1393 (1991).

24. Giner, J.-L., Li, X. & Boyer, G. L. Sterol composition of *Aureoumbra lagunensis*, the

Texas brown tide alga. *Phytochemistry* **57**, 787–789 (2001).

25. Giner, J.-L., Zhao, H., Boyer, G. L., Satchwell, M. F. & Andersen, R. A. Sterol

chemotaxonomy of marine pelagophyte algae. *C&B* **6**, 1111–1130 (2009).

26. Taylor, M. W., Radax, R., Steger, D. & Wagner, M. Sponge-associated

microorganisms: evolution, ecology, and biotechnological potential. *MMBR* **71**, 295–

347 (2007).

27. Nes, W. D. Enzyme mechanisms for sterol C -methylations. *Phytochemistry* **64**, 75–

95 (2003).

28. Gold, D. A. *et al.* Sterol and genomic analyses validate the sponge biomarker

hypothesis. *Proc Natl Acad Sci U S A* **113**, 2684–2689 (2016).

29. Gold, D. A. *et al.* Lipidomics of the sea sponge *Amphimedon queenslandica* and

implication for biomarker geochemistry. *Geobiology* **15**, 836–843 (2017).

30. Bergmann, W. & McTigue, F. H. Contributions to the study of marine products. XXI.

Chondrillasterol. *J Org Chem* **13**, 738–741 (1948).

31. Venkateswarlu, Y., Reddy, M. V. R. & Rao, M. R. A new epoxy sterol from the

sponge *Ircinia fasciculata*. *J Nat Prod* **59**, 876–877 (1996).

32. Fromont, J., Kerr, S., Kerr, R., Riddle, M. & Murphy, P. Chemotaxonomic

relationships within, and comparisons between, the orders Haplosclerida and

Petrosida (Porifera: Demospongiae) using sterol complements. *Biochem Syst Ecol* **22**,

735–752 (1994).

33. Hagemann, A., Voigt, O., Wörheide, G. & Thiel, V. The sterols of calcareous

sponges (Calcarea, Porifera). *Chemistry and Physics of Lipids* **156**, 26–32 (2008).

34. Aiello, A., Fattorusso, E., Magno, S. & Menna, M. Isolation of five new 5α-hydroxy-

6-keto-Δ7 sterols from the marine sponge *Oscarella lobularis*. *Steroids* **56**, 337–340

(1991).

35. Lee, A. K. *et al.* C-4 sterol demethylation enzymes distinguish bacterial and

eukaryotic sterol synthesis. *Proc Natl Acad Sci USA* **115**, 5884–5889 (2018).

36. Santana-Molina, C., Rivas-Marin, E., Rojas, A. M. & Devos, D. P. Origin and

evolution of polycyclic triterpene synthesis. *Mol Biol Evol* **37**, 1925–1941 (2020).

37. Michellod, D. *et al.* De novo *phytosterol synthesis in animals*.

http://biorxiv.org/lookup/doi/10.1101/2022.04.22.489198 (2022)

doi:10.1101/2022.04.22.489198.

38. Gold, D. *et al. Sterol methyltransferases in annelid worms rewrite the molecular*

fossil record. https://www.researchsquare.com/article/rs-1686449/v1 (2022)

doi:10.21203/rs.3.rs-1686449/v1.

39. Giner, J.-L. & Djerassi, C. Biosynthetic studies of marine lipids-XXVIII. Use of

sponge cell-free extracts in the study of marine sterol biosynthesis. *Tetrahedron Lett*

31, 5421–5424 (1990).

40. Kerr, R. G., Stoilov, I. L., Thompson, J. E. & Djerassi, C. Biosynthetic studies of

marine lipids 16. De novo sterol biosynthesis in sponges. Incorporation and

transformation of cycloartenol and lanosterol into unconventional sterols of marine

and freshwater sponges. *Tetrahedron* **45**, 1893–1904 (1989).

41. Silva, C. J., Wünsche, L. & Djerassi, C. Biosynthetic studies of marine lipids 35. The

demonstration of de novo sterol biosynthesis in sponges using radiolabeled

isoprenoid precursors. *Comp Biochem Physiol B* **99B**, 763–773 (1991).

42. Germer, J., Cerveau, N. & Jackson, D. J. The holo-transcriptome of a calcified early

branching metazoan. *Front Mar Sci* **4**, 81 (2017).

43. Kelly, J. B., Carlson, D., Low, J. S. & Thacker, R. W. *Comparative metagenomics*

evidence distinct trends of genome evolution between sponge-dwelling bacteria and

their pelagic counterparts. http://biorxiv.org/lookup/doi/10.1101/2020.08.31.276493

(2020) doi:10.1101/2020.08.31.276493.

44. Ganapathy, K. *et al.* Molecular probing of the *Saccharomyces cerevisiae* sterol 24-C

methyltransferase reveals multiple amino acid residues involved with C2-transfer

activity. *Biochim Biophys Acta Mol Cell Biol Lipids* **1781**, 344–351 (2008).

45. Nes, W. D. *et al.* Site-directed mutagenesis of the sterol methyl transferase active site

from *Saccharomyces cerevisiae* results in formation of novel 24-ethyl sterols. *J Org*

Chem **64**, 1535–1542 (1999).

46. Nes, W. D. *et al.* Probing the sterol binding site of soybean sterol methyltransferase

by site-directed mutagenesis: functional analysis of conserved aromatic amino acids

in region 1. *Arch Biochem Biophys* **448**, 23–30 (2006).

47. Moldowan, J. M. C30-steranes, novel markers for marine petroleums and

sedimentary rocks. *Geochim Cosmochim Acta* **48**, 2767–2768 (1984).

48. Elwell, C. A. & Engel, J. N. Lipid acquisition by intracellular Chlamydiae. *Cell*

Microbiol **14**, 1010–1018 (2012).

49. Wünsche, L., Gülaçar, F. O. & Buchs, A. Several unexpected marine sterols in a

freshwater sediment. *Org Geochem* **11**, 215–219 (1987).

50. Stoilov, I. L., Thompson, J. E., Cho, J. Ho. & Djerassi, Carl. Biosynthetic studies of

marine lipids. 9. Stereochemical aspects and hydrogen migrations in the biosynthesis

of the triply alkylated side chain of the sponge sterol strongylosterol. *J Am Chem Soc*

108, 8235–8241 (1986).

Methods

Bioinformatics analysis. SMT homologs were identified in the Joint Genome

Institute Integrated Microbial Genomes & Microbiomes (JGI IMG), GenBank, and

Compagen (http://www.compagen.org) databases using the *Saccharomyces cerevisiae*

S299C SMT amino acid sequence as the BLASTP search query (locus tag: YML008C,

maximum e-value: $1e^{-50}$, minimum percent identity: 30%). Additional demosponge SMT

sequences from *C. nucula*, *S. fasciculatas*, and *P. ficiformis* transcriptomes were provided

by Prof. David A. Gold[28]. Identifying sequence information is available in Extended Data

Table 1. Conserved sterol methyltransferase C-terminal and methyltransferase 11 domains

were identified using the HmmerWeb tool (version 2.41.1).

Oxidosqualene cyclase homologs were identified through BLASTP searches of the

metagenome scaffolds and MAGs containing the SMTs analyzed in this study and all

bacterial and eukaryotic genomes available in JGI IMG as of October 2019 using the

Methylococcus capsulatus Bath OSC protein sequence (locus tag: MCA2873, maximum

e-value: $1e^{-50}$, minimum percent identity: 20%, minimum sequence length: 350 aa).

Redundancy was decreased through manual removal of OSC homologs with 100%

similarity at the amino acid level. The N-termini of fused SMO/OSC homologs were

trimmed out of the dataset to begin the sequences at the squalene-hopene cyclase N-

terminal domain as identified using HmmerWeb.

All protein sequence datasets were aligned via MUSCLE. Conserved SMT domains

were aligned separately, then concatenated using Geneious. Maximum-likelihood trees

were generated using IQ-Tree on XSEDE with 1,000 ultra-fast bootstrap replicates using

the best amino acid substitution models, gamma shape parameters, and invariable sites

proportions under the Bayesian Information Criterion and Akaike Information Criterion

using ModelTest-NG on XSEDE (Extended Data Table 4). Resulting phylogenetic trees

were visualized in iTOL[51].

All contigs containing metagenomic SMTs were retrieved from JGI IMG or

GenBank and analyzed with Whokaryote[52] using the default inputs with the minimum

contig size decreased to 2,000 base pairs. Calculated features and predictions are shown in

Extended Data Table 2.

Gene synthesis and molecular cloning. SMT DNA sequences were codon-

optimized for expression in *E. coli* and artificially synthesized by GeneArt (ThermoFisher

Scientific) or through the Department of Energy Joint Genome Institute (DOE JGI) DNA

Synthesis Science Program. SMT sequences from DOE JGI were obtained in the IPTG-

inducible plasmid pSRKGm-*lac*UV5-rbs5[53] in *E. coli* TOP10. SMT sequences from

GeneArt were subcloned into pSRKGm-*lac*UV5-rbs5 by sequence and ligase independent

cloning (SLIC)[54] and transformed into electrocompetent *E. coli* DH10B (Invitrogen) using

a MicroPulser Electroporator (BioRad). Oligonucleotides were purchased from Integrated

DNA Technologies (Coralville, IA). PCR was performed according the to the

manufacturer's protocol using Phusion DNA Polymerase (New England Biolabs). DNA

fragments were purified using the GeneJET Gel Extraction Kit (ThermoFisher Scientific).

Plasmid DNA was isolated using the GeneJET Plasmid Miniprep Kit (ThermoFisher

Scientific). DNA was sequenced by ELIM Biopharm (Hayward, CA).

Bacterial culture and heterologous expression. *E. coli* expression strains were

cultured in 50 mL terrific broth (TB) supplemented with gentamycin (15 μg/mL) at 37 °C

while shaking at 225 rpm. Cultures were induced with 500 μM isopropyl *β*-D-1-

thiogalactopyranoside (IPTG) at an OD_{600} of ~0.6, then incubated an additional 4 hours at

30 °C while shaking at 225 rpm. Cells were harvested by centrifugation at $4,500 \times g$ for 10

minutes at 4 °C. Cell pellets were stored at -80 °C until sonication.

Sterol methyltransferase assay. Cells pellets were resuspended in 5 mL buffer

containing the following: 50 mM Tris-HCl, 2 mM $MgCl_2$, 20% glycerol (v/v), and 0.1%

β-mercaptoethanol (v/v), pH 7.5. Cells were then lysed on ice via a Qsonica Sonicator

Q500 equipped with a 3.2 mm probe at 30% amplitude pulsing at 5 seconds on, 15 seconds

off for 8 minutes of total on time. Lysates were then partially clarified by centrifugation at

$4,500 \times g$ for 20 minutes at 4 °C. Total protein concentration in the resulting supernatant

was quantified using a Coomassie (Bradford) Protein Assay Kit (Thermo Scientific)

according to the manufacturer's Standard Microplate Protocol and a BioTek Synergy HT

Microplate Reader. SMT assays were then immediately prepared using enough supernatant

to give 4500 μg total protein. Other reaction components included 100 μM desmosterol

(Sigma-Aldrich), 24-methylenecholesterol (Avanti Polar Lipids, Inc.), [28-[13]C] 24-

methylenecholesterol (Avanti Polar Lipids, Inc.), or synthesized [29-[13]C] fucosterol or

isofucosterol, 100 μM *S*-(5′-sdenosyl)-*L*-methionine chloride dihydrochloride (Sigma-

Aldrich), and 0.1% Tween-80 (v/v) to a total reaction volume of 400 μL. Reactions were

held at 30 °C for 20 hours and then stored at -20 °C until lipid extraction.

Lipid extraction. Lipids were extracted using a modified Bligh-Dyer method[55,56]

with the completed in vitro reactions as the water phase. Reactions were sonicated in 10:5:4

(vol:vol:vol) methanol:dichloromethane:water for 1 hour. The organic phase was then

separated with twice the volume of 1:1 (vol:vol) dichloromethane:water followed by

storage at -20 °C for >1 hr. Following centrifugation at 2,800 x *g* for 10 min at 4 °C, the

organic phase was transferred and evaporated under N_2 to give total lipid extracts (TLEs).

TLEs were derivatized to trimethylsilyl ethers in 1:1 (vol:vol)

pyridine:Bis(trimethylsilyl)trifluoroacetamide for 1 hr at 70 °C prior to GC-MS analysis.

GC-MS analysis. Lipids were separated with an Agilent 7890B Series GC

equipped with two Agilent DB-17HT columns (30 m x 0.25 mm i.d. x 0.15 μm film

thickness) in tandem with helium as the carrier gas at a constant flow of 1.1 ml/min and

programmed as follows: 100°C for 2 min, then 12°C/min to 250°C and held for 10 min,

then 10°C/min to 330°C and held for 17.5 min. 2 uL of each sample was injected in splitless

mode at 250°C. The GC was coupled to an Agilent 5977A Series MSD with the ion source

at 230°C and operated at 70 eV in EI mode scanning from 50 to 850 Da in 0.5 s. Sterols

were identified based on retention time and comparison to previously published spectra

and laboratory standards.

NMR analysis. In addition to GC-MS analysis, structures of sterol products formed by the metagenomic sterol methyltransferases referred to as "Aplysina aerophoba JGIcombinedJ30088_10000835 1 of 2" and "Chlamydiae sp., drinking water 1 of 2" were confirmed by NMR. ^{13}C-labeling experiments were used to assist the identification of the biosynthetic products by 2D-NMR (heteronuclear single quantum coherence (HSQC) and heteronuclear multiple bond correlation (HMBC)) by increasing the amount of ^{13}C at a specific position by two orders of magnitude compared to natural abundance. The two terminal isopropyl groups of 24-isopropyl sterols come from biosynthetically disparate origins, the isoprenoid pathway and SAM methylation. Because of the possibilities of hydride-shifts and different positions of deprotonation, each of the four terminal methyl groups can have multiple different origins within the product[20]. In addition to enhanced sensitivity, ^{13}C-labeling provided mechanistic details of a pathway which, because of the intrinsic symmetry of 24-isopropyl sterols, would be otherwise hidden.

Total lipid extracts from pooled in vitro reactions using [28-^{13}C] 24-methylenecholesterol or [29-^{13}C] isofucosterol as the substrate for both SMTs were fractionated using small silica pipet columns (1 mL) eluting with 5 mL each of hexanes:ethyl acetate 9:1, 2:1, and 1:1 (vol:vol). Steroidal constituents were found in the 2:1 solvent fraction and concentrated with a stream of N$_2$ prior to further purification by preparative thin-layer chromatography (TLC; 4:1 hexanes:ethyl acetate). Sterols were separated by high-performance liquid chromatography (HPLC) and the fractions were analyzed by 800 MHz NMR[35]. Sterols were identified by comparison of the HSQC-DEPT and HMBC spectra with reference compounds.

In experiments with [28-^{13}C] 24-methylenecholesterol, the presence of

isofucosterol was confirmed by a ^{1}H-^{13}C cross peak corresponding to C-28 at 5.108 and

116.48 ppm. The presence of fucosterol was shown by an HSQC cross peak at 5.183 and

115.58 ppm. Ratios of isofucosterol to fucosterol were determined by comparison of the

integral ratios of the cross peaks[35] for these two isomers. The presence of epiclerosterol

was confirmed by cross peaks for C-28 at 1.290/1.380 and 26.01 ppm. Structures of the

C$_{29}$ sterol products were confirmed by long-range HMBC correlations from the 29-methyl

group. Isofucosterol showed an HMBC cross peak for H-29 and C-28 at 1.59 and 115.58

ppm; fucosterol and epiclerosterol showed HMBC cross peaks at 1.57 and 116.48 ppm and

0.799 and 26.01 ppm, respectively. The presence of *24S*-24-isopropylcholesta-5,25-dienol

labeled at the C-25 position was confirmed by HMBC from the 27-methyl group (cross

peak between 1.570 and 147.38 ppm). The presence of 24-isopropylcholesta-5,24-dienol

was confirmed by HMBC showing two adjacent cross peaks correlating the C-26 and C-

27 methyl protons (1.633/1.651 ppm) with C-25 (123.1 ppm).

In experiments with [29-^{13}C] isofucosterol, the label distribution between the C-27

and C-26 positions of *24S*-24-isopropylcholesta-5,25-dienol was shown by integration of

the HSQC signals for C-27 (1.570 and 18.95 ppm) and C-30 (4.602/4.740 and 111.85). The

label on the Z-vinylic methyl of 24-isopropylcholesta-5,24-dienol was detected by HSQC

(1.651 and 19.73 ppm). 24-Isopropylcholesta-5,23-dienol was detected by label at 20.96

ppm exclusively, which is one of the two methyls of the Z-isopropyl group. The ratio of

24S-24-isopropylcholesta-5,25-dienol, 24-isopropylcholesta-5,24-dienol, and 24-

isopropylcholesta-5,23-dienol was estimated by integration of the HSQC.

Preparation of [29-^{13}C] fucosterol and isofucosterol. Specifically labeled C_{29}

sterol precursors were prepared by modification of the procedure reported in[57]. The

precursor 24-formylcholesterol i-methyl ether was obtained either through hydroboration

of i-Me 24-methylenecholesterol followed by Dess-Martin periodinane oxidation or

ozonolysis of i-Me 24-vinylcholesterol. The latter reaction was carried out in 90:9:1

(vol:vol:vol) dichloromethane/methanol/pyridine and yielded the desired 28-

formylcholesterol *i*-methyl ether and its methyl hemiacetal in a 1:5 ratio. The hemiacetal

was isolated as a mixture of four isomers. ^{1}H-NMR (600MHz): 8.06-8.03 (1H, 28-OH),

4.68-4.63 (1H, 28-H), 3.56-3.52 (3H, 28-OMe) 3.332 (s, 3H, 6-OMe), 2.767 (s, 1H, 6-H),

1.108 (s, 3H, 19-H), 0.710 (s, 3H, 18-H). This mixture was reacted with ^{13}C-

methylmagnesium iodide to give four isomers of [29-^{13}C] i-Me 28-hydroxysitosterol:

24R,28R (H-28: 3.938 ppm; C-29: 21.97 ppm), 24S,28S (H-28: 3.938 ppm; C-29: 22.20

ppm), 24R,28S (H-28: 3.820 ppm; C-29: 21.05 ppm), and 24S,28R (H-28: 3.803 ppm; C-

29: 21.01 ppm). Preparative TLC (4:1 hexanes:ethyl acetate) gave two bands

corresponding to a mixture of the RR and SS isomers (higher Rf) and a mixture of the RS

and SR isomers (lower Rf) in ca. 2:1 ratio. These two mixtures were separately dehydrated

with phosphorus oxychloride to obtain after deprotection [29 ^{13}C] fucosterol (C-29: 13.16

ppm) and [29 ^{13}C] isofucosterol (C-29: 12.76 ppm), respectively.

Data availability

The data that support the findings of this study are available within this paper and its supplementary information files. Additional (raw) data are available from the corresponding author upon reasonable request.

Methods references

51. Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display and annotation. *Nucleic Acids Res* **49**, W293–W296 (2021).

52. Pronk, L. J. U. & Medema, M. H. Whokaryote: distinguishing eukaryotic and prokaryotic contigs in metagenomes based on gene structure. *Microbial Genomics* **8**, (2022).

53. Banta, A. B., Wei, J. H., Gill, C. C. C., Giner, J.-L. & Welander, P. V. Synthesis of arborane triterpenols by a bacterial oxidosqualene cyclase. *Proc Natl Acad Sci USA* **114**, 245–250 (2017).

54. Li, M. Z. & Elledge, S. J. SLIC: a method for sequence- and ligation-independent cloning. *Methods Mol Biol* **852**, 51–59 (2012).

55. Welander, P. V. *et al.* Identification and characterization of *Rhodopseudomonas palustris* TIE-1 hopanoid biosynthesis mutants. *Geobiology* **10**, 163–177 (2012).

56. Bligh, E. G. & Dyer, W. J. A rapid method of total lipid extraction and purification. *Can J Biochem Physiol* **37**, 911–917 (1959).

57. Giner, J. L., Zimmerman, M. P. & Djerassi, C. Synthesis of (24R,28R)- and

(24S,28S)-24,28-methylene-5-stigmasten-3.beta.-ol and biosynthetic implications of

cyclopropyl cleavage to 24-substituted cholesterols. *J. Org. Chem.* **53**, 5895–5902

(1988).

Acknowledgments. We thank Prof. David A. Gold for providing demosponge SMT

protein sequences, Jeremy H. Wei for technical assistance, and Hanon McShea for helpful

discussions. We also thank the Department of Energy Joint Genome Institute DNA

Synthesis Science Program (Project 503267) for synthesis and cloning of the majority of

the SMTs analyzed in this study. Funding for this study was provided by National Science

Foundation Grant EAR-1752564 (to P.V.W).

Author contributions. M.O.B. wrote the manuscript with input from all co-authors.
M.O.B. performed the molecular biology experiments, lipid extractions, and GC-MS
analyses. M.O.B. and P.V.W. performed the bioinformatic analyses. B.O.O. and J.-L.G.
synthesized the ^{13}C labeled sterols and performed the NMR analyses. M.O.B analyzed and
interpreted the bioinformatic and GC-MS data. B.O.O. and J.-L.G. analyzed and
interpreted the NMR data. P.V.W. and M.O.B. conceived the project with contribution
from J.-L.G.

Competing interest declaration. The authors declare no conflict of interest.

Corresponding author: welander@stanford.edu

535 **Table 1. Sterols identified as products of sponge and bacterial sterol**

536 **methyltransferases.**

	C28			C29			C30	
	(epi)codisterol (II)	24-methylenecholesterol (III)	24-methyldesmosterol (V)	(epi)clerosterol (IV)*	fucosterol (VI)	isofucosterol (VIII)	24S-24-isopropylcholest-5,25-dienol (VII)	24-isopropylcholest-5,24-dienol (IX)
DEMOSPONGIAE								
Amphimedon queenslandica		+			+			
Chondrilla nucula	+	+		+				
Sarcotragus fasciculatus	+	+			+			
Haliclona amboinensis	+	+		+	+	+		
Haliclona tubifera	+	+		+				
HOMOSCLEROMORPHA								
Oscarella pearsei	+	+		+	+	+		
Oscarella carmela	+	+						
CALCAREA								
Sycon ciliatum	+	+						
DEMOSPONGE METAGENOME								
Aplysina aerophoba JGIcombinedJ30088_10000835 1 of 2	+	+		+	+	+	+	
Aplysina aerophoba JGIcombinedJ30088_10000835 2 of 2								
Aplysina aerophoba Ga0209021_1000146	+	+	+		+	+		+
Petrosia ficiformis LXNJ01014139		+						
Petrosia ficiformis LXNJ01003202		+						
Petrosia ficiformis LXNJ01000389	+	+	+		+			
Petrosia ficiformis LXNJ01004423								
Sarcotragus foetidus		+						
DEMOSPONGE METAGENOME-ASSEMBLED GENOME								
Theonella swinhoei associated Chromatiales sp. 1 of 2		+						
Theonella swinhoei associated Chromatiales sp. 2 of 2								
OTHER METAGENOME								
Marine sediment, Gulf of Thailand		+						
Marine sediment, Helgoland, North Sea		+						
Western Arctic Ocean	+	+		+	+	+		
Freshwater, Lake Fryxell, Antarctica	+	+			+	+		
Lake sediment, Walker Lake, Nevada		+						
Hotspring, Beatty, Nevada								
Hotspring sediment, Dewer Creek, BC		+						
OTHER METAGENOME-ASSEMBLED GENOME								
Chlamydiae sp., drinking water 1 of 2**	+	+	+	+	+	+	+	+
Chlamydiae sp., drinking water 2 of 2	+		+					
Nitrospira sp., coral reef 1 of 2	+	+		+				
Nitrospira sp., coral reef 2 of 2								
Sandaracinus sp., marine 1 of 2	+	+						
Sandaracinus sp., marine 2 of 2	+	+		+	+	+		
Spirochaetes sp., soil		+						

537

538 *NMR confirmed epiclerosterol and not clerosterol as products of the "*Aplysina aerophoba*

539 JGIcombinedJ30088_10000835 1 of 2" and "Chlamydiae sp., drinking water 1 of 2" sterol

540 methyltransferases.

541 **24-isopropylcholesta-5,23-dienol was also detected as a product of the "Chlamydiae sp.,

542 drinking water 1 of 2" sterol methyltransferase by NMR, but was not detected by GC-MS.

Figure 1. Sterol methyltransferases from sponges and bacteria perform multiple side-chain methylations. **a**, Structures of the sponge biomarkers 26-methylstigmastane (26-mes) and 24-isopropylcholestane (24-ipc). **b**, Predominant pathway to 24-isopropyl sterols by bacteria sterol methyltransferases (SMTs). **c**, Representative extracted ion chromatograms (*m/z* 456, 470, 484, 498) of total lipid extracts from in vitro reactions performed with *E. coli* lysates containing an empty plasmid (**upper**), a sterol methyltransferase (SMT) identified in a freshwater metagenome-assembled genome assigned to the phylum Chlamydiae (**middle**), and an SMT identified in a metagenome from the microbiome of the demosponge *Aplysina aerophoba* (**lower**) with desmosterol as the substrate. **d**, Representative extracted ion chromatograms (*m/z* 456, 470, 484, 498) of total lipid extracts from in vitro reactions performed with *E. coli* lysates containing an empty plasmid (**upper**), an SMT identified in a transcriptome of the demosponge *Chondrilla nucula* (**middle**), and the same SMT identified in the *A. aerophoba* metagenome as in **a** (**lower**) with 24-methylenecholesterol as the substrate. **e**, Side-chain structures of sterols identified in **a** and **b**. All lipids were derivatized to trimethylsilyls prior to GC-MS analysis. Mass spectra of identified sterols are shown in Extended Data Fig. 1.

Figure 2. Sterol gene clusters in demosponge metagenomes suggest de novo biosynthesis by bacterial symbionts. a, Putative sterol biosynthesis gene clusters identified in metagenomes from the microbiomes of the demosponge species *Aplysina aerophoba, Petrosia ficiformis*, and *Theonella swinhoei*, each containing a functional sterol methyltransferase. **b**, Schematic of the bacterial sterol biosynthesis pathway from squalene to 4-methylcholesta-8,24,-dien-3-ol. Arrow colors at each step correspond to the homologs identifiied in **a**. Detailed information on the source of each metagenomic locus is available in Extended Data Table 1.

Figure 3. Metagenomic SMTs originate from bacteria. a, Maximum likelihood tree generated from a concatenated alignment of the conserved methyltransferase and C-terminal domains of SMT proteins. Metagenomic SMTs are in blue, sponge SMTs are in red, and other eukaryotic SMTs are in black. The numbers in parentheses indicate the number of proteins in collapsed clades. Bolded labels indicate SMTs tested in this study. **b**, Maximum likelihood tree of oxidosqualene cyclases (OSCs). OSC groups are assigned as in[36]. Bolded labels indicate OSCs in metagenomic gene clusters or metagenome-assembled genomes with SMTs tested in this study.

575 **Extended Data Figure 1**. **Mass spectra of sterols identified in the main text.** Total lipid

576 extracts were derivatized to trimethylsilyl ethers. Detailed GC-MS methods can be found

577 in the Methods section of the main text.

Extended Data Figure 2. ^{13}C-labeling experiments with propylating SMTs from an *Aplysina aerophoba* **symbiont and a Chlamydiae sp.** Positions of the ^{13}C labels are shown for the predominant propylation pathway proceeding from 24-methylenecholesterol (III) to isofucosterol (VI) to 24S-24-isopropylcholest-5,25,dienol (VIII). **a**, Experiments with [28-^{13}C] 24-methylenecholsterol as the substrate. **b**, Experiments with [29-^{13}C] isofucosterol as the substrate.

 Extended Data Table 1. Sources of sterol methyltransferases analyzed in this study.

SMT	Source	Genome ID/ BioProject	Locus Tag/Locus
DEMOSPONGIAE			
Amphimedon queenslandica	JGI IMG	2507525020	anonymous.512
Chondrilla nucula	(28)		
Sarcotragus fasciculatus	(28)		
Haliclona amboinensis	Compagen		
Haliclona tubifera	Compagen		
HOMOSCLEROMORPHA			
Oscarella pearsei	Compagen		
Oscarella carmela	Compagen		
CALCAREA			
Sycon ciliatum	Compagen		
DEMOSPONGE METAGENOME			
Aplysina aerophoba JGIcombinedJ30088_10000835 1 of 2	JGI IMG	3300002448	JGIcombinedJ30088_1000083511
Aplysina aerophoba JGIcombinedJ30088_10000835 2 of 2	JGI IMG	3300002448	JGIcombinedJ30088_100008359
Aplysina aerophoba Ga0209021_1000146	JGI IMG	3300027386	Ga0209021_100014628
Petrosia ficiformis LXNJ01014139	GenBank	PRJNA318959	LXNJ01014139.1
Petrosia ficiformis LXNJ01003202	GenBank	PRJNA318959	LXNJ01003202.1
Petrosia ficiformis LXNJ01000389	GenBank	PRJNA318959	LXNJ01000389.1
Petrosia ficiformis LXNJ01004423	GenBank	PRJNA318959	LXNJ01004423.1
Sarcotragus foetidus	GenBank	PRJNA318959	LXNI01004480.1
DEMOSPONGE METAGENOME-ASSEMBLED GENOME			
Theonella swinhoei associated Chromatiales sp. 1 of 2	JGI IMG	2690315631	Ga0136241_103361
Theonella swinhoei associated Chromatiales sp. 2 of 2	JGI IMG	2690315631	Ga0136241_10845
OTHER METAGENOME			
Marine sediment, Gulf of Thailand	JGI IMG	3300010430	Ga0118733_1000126028
Marine sediment, Helgoland, North Sea	JGI IMG	3300004097	Ga0055584_1001717352
Western Arctic Ocean	JGI IMG	3300010883	Ga0133547_104202862
Freshwater, Lake Fryxell, Antarctica	JGI IMG	3300009032	Ga0105048_100346915
Lake sediment, Walker Lake, Nevada	JGI IMG	3300009504	Ga0114946_100084975
Hot spring, Beatty, Nevada	JGI IMG	3300009444	Ga0114945_100210811
Hot spring sediment, Dewer Creek, BC	JGI IMG	3300007072	Ga0073932_10040971
OTHER METAGENOME-ASSEMBLED GENOME			
Chlamydiae sp., drinking water 1 of 2	JGI IMG	2619618861	Ga0074140_11584
Chlamydiae sp., drinking water 2 of 2	JGI IMG	2619618861	Ga0074140_11583
Nitrospira sp., coral reef 1 of 2	JGI IMG	2681812944	Ga0131816_12726
Nitrospira sp., coral reef 2 of 2	JGI IMG	2681812944	Ga0131816_12722
Sandaracinus sp., marine 1 of 2	JGI IMG	2775506781	Ga0248530_10706
Sandaracinus sp., marine 2 of 2	JGI IMG	2775506781	Ga0248530_10707
Spirochaetes sp., soil	JGI IMG	2708742746	Ga0154758_10818

587 **Extended Data Table 2.** Whokaryote predicts all metagenomic contigs containing sterol

588 methyltransferases analyzed in this study as prokaryotic.

SMT	contig	contig length	nr genes	ratio same orientation	ID general avg	ID general std	gene density	gene length	ID Q1	ID Q3	rbs ratio	tiara pred	predicted
Aplysina aerophoba JGIcombinedJ30088_10000835	JGIcombinedJ30088_10000835	32339	23	0.68	128.14	183.70	0.91	1285.09	3.75	181.00	0.96	1.0	prokaryote
Aplysina aerophoba Ga0209021_1000146	Ga0209021_1000146	97163	103	0.68	115.34	117.55	0.88	829.83	31.00	173.75	0.90	1.0	prokaryote
Petrosia ficiformis LXNJ01014139	LXNJ01014139.1	3313	4	0.00	100.67	62.53	0.91	752.00	56.50	124.50	0.25	1.0	prokaryote
Petrosia ficiformis LXNJ01003202	LXNJ01003202.1	8597	8	0.57	115.57	167.57	0.91	975.50	0.00	131.00	0.88	1.0	prokaryote
Petrosia ficiformis LXNJ01000389	LXNJ01000389.1	31336	22	0.67	79.90	78.14	0.94	1341.91	4.00	147.00	0.77	1.0	prokaryote
Petrosia ficiformis LXNJ01004423	LXNJ01004423.1	7072	6	0.60	102.20	58.00	0.93	1093.00	63.00	126.00	0.67	1.0	prokaryote
Sarcotragus foetidus	LXNI01004480.1	7225	5	0.50	105.25	156.88	0.94	1359.20	6.75	121.00	0.80	1.0	prokaryote
Theonella swinhoei associated Chromatiales sp. 1 of 2	2690316284	444506	427	0.74	74.66	122.32	0.93	968.13	1.00	104.25	0.68	1.0	prokaryote
Theonella swinhoei associated Chromatiales sp. 2 of 2	2690316289	117771	97	0.83	74.25	116.07	0.94	1142.96	0.00	97.50	0.65	1.0	prokaryote
Marine sediment, Gulf of Thailand	Ga0118733_100012602	21160	19	0.89	72.06	133.71	0.94	1042.53	0.00	65.50	0.84	1.0	prokaryote
Marine sediment, Helgoland, North Sea	Ga0055584_100171735	2192	3	0.50	80.50	75.50	0.93	676.00	42.75	118.25	0.00	1.0	prokaryote
Western Arctic Ocean	Ga0133547_10420286	2727	3	0.00	147.00	126.00	0.89	810.00	84.00	210.00	0.33	1.0	prokaryote
Freshwater, Lake Fryxell, Antarctica	Ga0105048_10034691	7428	9	0.63	86.00	80.31	0.88	722.33	13.50	104.75	0.44	1.0	prokaryote
Lake sediment, Walker Lake, Nevada	Ga0114946_10008497	6500	5	0.50	77.25	71.98	0.93	1212.80	22.50	103.75	0.40	1.0	prokaryote
Hotspring, Beatty, Nevada	Ga0114945_10021081	3480	3	0.00	63.00	7.00	0.96	1108.00	59.50	66.50	0.33	1.0	prokaryote
Hotspring sediment, Dewer Creek, BC	Ga0073932_1004097	18020	12	0.82	164.73	132.22	0.89	1342.75	67.00	269.00	0.92	1.0	prokaryote
Chlamydiae sp., drinking water 1 and 2	2619626156	1146800	1105	0.68	85.18	128.28	0.92	956.10	0.00	118.00	0.42	1.0	prokaryote
Nitrospira sp., coral reef 1 and 2	2681831735	70077	63	0.74	147.08	145.66	0.87	962.71	27.00	196.00	0.89	1.0	prokaryote
Sandaracinus sp., marine 1 and 2	2775515747	24518	17	0.88	112.25	106.29	0.89	1282.47	22.50	163.50	0.76	1.0	prokaryote
Spirochaetes sp., soil	2708772188	70818	55	0.54	151.69	167.86	0.89	1142.60	6.25	242.50	0.24	1.0	prokaryote

Extended Data Table 3. 800 MHz ^{1}H and 201 MHz ^{13}C NMR assignments of selected sterol side-chain positions at 30 °C.

	C-24	C-25	C-26	C-27	C-28	C-29	C-30
Fucosterol	**145.6**	2.199 **34.7**			5.183 **115.58**	1.574 **13.16**	
Isofucosterol	**147.1**	2.829 **28.6**			5.108 **116.48**	1.591 **12.76**	
Clerosterol*	1.838 **49.54**	**147.61**	4.730, 4.642 **111.33**	1.570 **17.85**	1.305, 1.341 **26.53**	0.805 **12.04**	
Epiclerosterol	1.81 **49.75**	**147.98**	4.725, 4.636 **111.06**	1.576 **18.13**	1.290, 1.380 **26.01**	0.800 **11.96**	
24*R*-24-Isopropylcholest-5,25-dienol*	1.56 **55.03**	**147.29**	4.613, 4.738 **112.02**	1.567 **18.63**	1.49 **30.25**	0.911 **20.85**	0.803 **21.60**
24*S*-24-Isopropylcholest-5,25-dienol	1.52 **55.50**	**147.38**	4.602 4.740 **111.85**	1.570 **18.95**	1.51 **30.24**	0.913 **20.85**	0.807 **21.49**
24-Isopropylcholest-5,23-dienol			1.013 **24.77**	1.013 **24.79**		0.980 **20.96**	0.980 **21.18**
24-Isopropylcholest-5,24-dienol	**137.0**	**123.1**	1.633 **20.81**	1.651 **19.73**		0.938 **21.34**	0.938 **21.41**

^{13}C data are in boldface. The assignments were determined by HSQC-DEPT, HMBC, and COSY 2D-NMR experiments. ^{1}H and ^{13}C chemical shifts are given to three and two decimal places, respectively, if directly observed, two and one, respectively, if determined from 2D spectra. Diastereotopic signals are not assigned. *Clerosterol and 24*R*-24-isopropylcholest-5,25-dienol were not detected as products of the *A. aerophoba* symbiont and Chlamydiae SMTs but are shown for comparative purposes.

597 **Extended Data Table 4.** Best amino acid substitution models, gamma shape parameters,

598 and invariable sites proportions under the Bayesian Information Criterion and Akaike

599 Information Criterion using ModelTest-NG on XSEDE for generation of sterol

600 methyltransferase and oxidosqualene cyclase maximum-likelihood trees.

Protein	Substitution Model	Gamma Shape Parameter	Invariable Sites Proportion
sterol methyltransferase	LG+I+G	1.260	0.020
oxidosqualene cyclase	LG+I+G	1.212	0.022