1   **Composition can buffer protein dynamics within liquid-like condensates.**

2   Stela Jelenic[1,2], Janos Bindics[1], Philipp Czermak[1], Balashankar R Pillai[1,2], Martine Ruer[3],
3   Carsten Hoege[3], Alex S Holehouse[4], Shambaditya Saha[1,5]
4
5   [1]Institute of Molecular Biotechnology of the Austrian Academy of Sciences (IMBA), Vienna
6   BioCenter (VBC), Dr. Bohr-Gasse 3, 1030 Vienna, Austria.
7
8   [2]Vienna BioCenter PhD Program, Doctoral School of the University of Vienna and Medical
9   University of Vienna, 1030 Vienna, Austria.
10
11 [3]Max Planck Institute of Molecular Cell Biology and Genetics, 01307 Dresden, Germany.
12
13 [4]Department of Biochemistry and Molecular Biophysics, Washington University School of Medicine,
14 St. Louis, MO 63110, USA.
15
[5]Correspondence: shambaditya.saha@imba.oeaw.ac.at

Summary

Most non-membrane-bound compartments in cells that form via phase separation have
complex composition. While phase separation of individual proteins that form these
compartments is well-documented, the mechanisms that modulate dynamics of individual
proteins in multicomponent systems remain unclear. Here, we used in vitro reconstitution
and in vivo experiments to investigate how the dynamics of a scaffold protein PGL-3 is
regulated within the liquid-like 'P granule' compartment in *C. elegans*. Using mutational and
biophysical perturbations, we generated PGL-3 constructs that form condensates in vitro
with widely varying dynamics. Using these PGL-3 constructs, we show that introducing
other P granule components buffers against change of dynamics within liquid-like
condensates. This dynamics-buffering effect is mediated by weak interactions among two
or more components. Such dynamics-buffering may contribute to robust functional output of
cellular liquid-like compartments.

Introduction

To organize biochemistry, cells assemble compartments with distinct compositions. Many of these compartments are not enclosed within lipid membranes and assemble via liquid-liquid phase separation (LLPS) of proteins and RNAs from the surrounding cytoplasm or nucleoplasm (Banani *et al*, 2017; Brangwynne *et al*, 2009; Lyon *et al*, 2021). Studies over the past decade have focused on identifying the molecular mechanisms of LLPS within cells. The current model is that LLPS is driven by 'scaffold' proteins and/or RNAs which participate in homotypic or heterotypic interactions with each other to assemble cellular non-membrane-bound compartments (Van Treeck & Parker, 2018; Banani *et al*, 2016, 2017; Lyon *et al*, 2021; Riback *et al*, 2020; Sanders *et al*, 2020; Saha *et al*, 2016). Other macromolecules called 'clients' do not contribute to compartment assembly but concentrate within those compartments and modulate biological function. It is now appreciated that LLPS can be driven by interactions of different types i.e. electrostatic, polar or hydrophobic interactions involving both folded domains and intrinsically disordered regions (Banani *et al*, 2017; Lyon *et al*, 2021). The mechanisms that regulate dynamics of components within these compartments, however, remain unclear.

Macromolecules can phase separate into condensates with different material properties, ranging from liquid-like phases with fast dynamics to dynamically-arrested gels as well as glasses (Banani *et al*, 2017; Jawerth *et al*, 2020). Many liquid-like condensates *in vitro* have also been shown to mature over time, starting as liquid-like droplets that transition into non-dynamic assemblies (Banani *et al*, 2017). The rates of biochemical reactions within biomolecular condensates must depend on the diffusion times of the components. Recent *in vitro* investigations of LLPS for simple two-component systems have shown that the behavior of one component can modulate the dynamics of another within condensates (Boeynaems *et al*, 2019; Rhine *et al*, 2020; Zhang *et al*, 2015; Alshareedah *et al*, 2021; Boyko *et al*, 2020). Cellular condensates contain hundreds of different proteins and RNAs (e.g. P granules, P bodies and stress granules (Youn *et al*, 2019; Phillips & Updike, 2022)). How such complex composition affects dynamics of macromolecules in liquid-like compartments remains poorly understood.

P granules are a canonical liquid-like compartment found in the germline of *C. elegans*

(Strome & Wood, 1982; Brangwynne *et al*, 2009). mRNA and over 70 proteins are known to

concentrate in P granules (Phillips & Updike, 2022). Mutations in P granule proteins are

associated with loss of fertility and fecundity in *C. elegans* (Phillips & Updike, 2022; Updike

& Strome, 2010). Eight proteins – PGL-1, PGL-3, GLH-1, GLH-2, GLH-3, GLH-4, DEPS-1

and MIP-1 are currently known to be constitutive P granule components i.e., these proteins

concentrate within P granules at all stages of germline development (Kawasaki *et al*, 1998,

2004; Spike *et al*, 2008a, 2008b; Price *et al*, 2021; Cipriani *et al*, 2021; Phillips & Updike,

2022). Proteins of the PGL family are specific to the *Caenorhabditis* genus and contain

RNA-binding motifs. The GLH- proteins belong to the highly conserved family of DEAD-box

RNA helicases. Based on current estimates of the concentrations of these proteins in vivo,

the PGL- and GLH- proteins are among the most abundant constitutive components in P

granules (Saha *et al*, 2016). In vitro reconstitution experiments have shown that, among the

constitutive P granule proteins, only the protein PGL-3 can phase separate into liquid-like

condensates in vitro (Saha *et al*, 2016, 201; Hanazawa *et al*, 2011). Two other proteins that

concentrate in P granules – MEG-3 and LAF-1 can also phase separate into liquid-like

condensates in vitro (Smith *et al*, 2016; Elbaum-Garfinkle *et al*, 2015). This suggests that

phase separation of these proteins also contribute to the assembly of P granules at some

stages of germline development. The interactions that support phase separation of the

PGL-3 protein remain unclear. Structural studies on the orthologous PGL-1 protein revealed

two dimerization domains (Aoki *et al*, 2016, 2021). This raises the possibility that PGL-3

also contains similar dimerization domains, and phase separation depends on interactions

involving these domains. The PGL-3 protein can phase separate in vitro only at high 'non-

physiological' protein concentrations (Saha *et al*, 2016). At physiological concentrations

(approx. 1 μM), binding to mRNA facilitates phase separation of the PGL-3 protein (Saha *et*

al, 2016). It remains unclear if interactions among the dimerization domains cooperate with

mRNA binding to support phase separation of the PGL-3 protein.

Here, we use in vitro reconstitution and in vivo experiments in adult *C. elegans* gonads to

investigate how complex composition within the P granule compartment contributes to the

dynamics of one of its scaffold proteins PGL-3 (Saha *et al*, 2016; Hanazawa *et al*, 2011).

We begin with structure-function analysis for insights into the mechanism of phase

separation of the PGL-3 protein. Using a combination of mutational analysis and

biophysical perturbations, we generated PGL-3 constructs that phase separate in vitro into

condensates with widely varying dynamics. Investigating the effect of other P granule

components on phase separation of PGL-3 constructs, we uncovered a role of complex

composition in buffering against large change of dynamics within cellular liquid-like

compartments. This "dynamics-buffering" effect is distinct from active processes that

depend on ATP hydrolysis, but is mediated by weak interactions among components in

liquid-like compartments.

Results

Homotypic and heterotypic interactions drive phase separation of PGL-3 and mRNA

at physiological concentrations.

We began by investigating the role of different domains in the phase separation properties

of the protein PGL-3 in vitro (Supplementary Fig. 1). Together with other proteins and

mRNA, PGL-3 drives the assembly of the canonical liquid-like P granules in the germline of

C. elegans. Bioinformatic analysis of the 693 residue-long PGL-3 protein predicted that the

N-terminus accounting for ~65% of PGL-3 is predominantly α-helical, while the C-terminus

accounting for ~35% is mostly disordered (Fig. 1a, Supplementary Fig. 2a, 2b). Consistent

with the bioinformatics predictions, circular dichroism (CD) spectroscopy under conditions

that do not support phase separation showed that the PGL-3 N-terminal fragment 1-452 is

predominantly α-helical, while the C-terminal fragment 370-693 is disordered (Fig. 1f).

Intrinsically disordered polypeptides have in many cases been shown to be necessary and

sufficient to drive phase separation. We therefore tested if a recombinantly purified C-

terminal fragment of PGL-3 (370-693), can drive phase separation in vitro. To our surprise,

we did not observe any phase separation even at high protein concentrations up to ~70 μM

(Fig. 1b, c). This greatly exceeds the physiological concentration of the PGL-3 protein in *C.*

elegans (estimated to be ~1 μM) (Saha *et al*, 2016). However, we found that a

recombinantly purified N-terminal fragment of PGL-3 (1-452) phase separates into

condensates at close to physiological concentrations of PGL-3 (Fig. 1b, c, Supplementary

figure 2e). Similar to full-length PGL-3, condensates of the N

Fig. 1 α-helical N
terminus drives phase
separation of PGL-3.
131
a Cartoon representing
PGL-3 and mutants.
b Bright field micrographs
of PGL-3 (23 µM), PGL-3
(1-452) (22 µM), PGL-3
(370-693) (67 µM) and
PGL-3 (K126E K129E)
(108 µM) in 25 mM HEPES
pH 7.5, 100 mM KCl, 1 mM
DTT in absence of
crowding agents. Scale bar
20 µm.
c Phase diagrams of PGL-3
and mutants at different
protein and salt (KCl)
concentrations. Blue circles
represent conditions where
phase separation was
observed. Black circles
represent conditions where
phase separation was
absent. Top row: Phase
diagrams of PGL-3, PGL-3
(370-693), and PGL-3
(K126E K129E) in 25 mM
HEPES pH 7.5, 1 mM DTT
in absence of crowding
agents. 5% of the pool of
protein was tagged with
mEGFP in each case.
Bottom row: Phase
diagrams of PGL-3, and
PGL-3 (1-452) in 25 mM
HEPES pH 7.5, 1 mM DTT,
1% PEG 35000. 5% of the
pool of protein was tagged
with mEGFP in each case.
d Time-lapse fluorescence
micrographs show fusion of
two PGL-3 (1-452)
condensates (5% tagged

with mEGFP in 25 mM HEPES pH 7.5, 100 mM KCl, 1 mM DTT, 1% PEG 35000). Scale bar 10 µm.
e Time-lapse fluorescence micrographs show fluorescence recovery after photobleaching in a condensate of PGL-3 (1-
452) (5% tagged with mEGFP in 25 mM HEPES pH 7.5, 100 mM KCl, 1 mM DTT, 1% PEG 35000). Scale bar 5 µm.
Shown below is a plot of average normalized fluorescence in bleached regions (1.5 µm x 1.5 µm) of the PGL-3 (1-452)
condensates (n = 24) over time. Error bars represent 1 SD.
f Plot of mean residue ellipticity of PGL-3 and mutants vs. wavelength assayed using circular dichroism (CD)
spectroscopy. Black circles: PGL-3 (3 µM, 10 mM Potassium phosphate pH 7.5, 300 mM KCl, 1 mM TCEP), Grey circles:
PGL-3 (1-452) (3 µM, 10 mM Potassium phosphate pH 7.5, 300 mM KCl, 1 mM TCEP), Blue circles: PGL-3 (370-693) (10
µM, 10 mM Potassium phosphate pH 7.5, 150 mM KCl, 1 mM TCEP). Error bars represent 1 SD among six CD scans.

terminal PGL-3 fragment (1-452) are liquid-like in nature. First, condensates are spherical in

shape (Fig. 1d, e, Supplementary Fig. 2d). Second, condensates fuse over time to generate

a larger condensate (Fig. 1d). Third, fluorescence recovery after photobleaching (FRAP)

experiments showed that PGL-3 (1-452) molecules intermix rapidly ($t_{1/2}$ = 4 s) within a

condensate (Fig. 1e). We conclude that phase separation of the protein PGL-3 is driven by

an α-helical region, while the disordered region may stabilize condensates by modulating

the driving forces for phase separation.

We next investigated how RNA impacts and alters the phase behaviour of PGL-3. It has

been previously shown that the PGL-3 protein cannot phase separate independently at

physiological concentration (Saha *et al*, 2016). At this concentration, sufficiently long mRNA

can trigger phase separation via binding to six arginine-glycine-glycine (RGG) repeats at

the C-terminal disordered region of PGL-3 (Saha *et al*, 2016). We therefore expected that

when the C-terminal disordered region was mixed with RNA, it would promote phase

separation. Unexpectedly, we found that in contrast to full-length PGL-3, mRNA does not

support phase separation of the C-terminal disordered region of PGL-3 (Fig. 2a, b). This

suggests that the N-terminal α-helical region is essential for phase separation. However, we

found that mRNA does not enhance phase separation of the N-terminal α-helical region of

PGL-3 either (Fig. 2a, b).

To investigate the underlying mechanism further, we began by testing if the N-terminal α-

helical region of PGL-3 can self-associate. Our analysis using size exclusion

chromatography followed by multi-angle light scattering (SEC-MALS) showed that this PGL-

3 fragment 1-452 forms a dimer (Supplementary Fig. 2f). Mutation of two residues (K126E

K129E) have been shown to interfere with interactions among the N-termini of PGL-3

molecules (Aoki *et al*, 2021). We mutated these two residues within the full-length PGL-3

protein (K126E K129E) (Fig. 1a) and found that this mutant PGL-3 (K126E K129E) protein

cannot phase separate even at high protein concentrations up to ~130 µM (Fig. 1b, c).

Addition of mRNA does not trigger phase separation of this protein at physiological

concentrations either (Fig. 2a, b). Taken together, our data is consistent with a model where

association among folded N-termini of PGL-3 molecules is essential for phase separation.

Fig. 2 Homotypic interaction among PGL-3 molecules is essential for mRNA-dependent phase separation.

a Fluorescence micrographs of PGL-3 and mutants (0.5 µM) in presence or absence of total mRNA (10 ng/µL). 5% of the pool of protein was tagged with mEGFP in each case. Buffer: 25 mM HEPES pH 7.5, 150 mM KCl, 1 mM DTT. 1% PEG 35,000 was added to PGL-3 (1-452). Scale bar 10 µm.
b Plot of the number of condensates with PGL-3 and mutants in presence or absence of total mRNA. For each condition, number of condensates was scored in nine different fluorescence micrographs (116 µm x 117 µm). Error bars represent 1 SD. t- test was performed to compare the number of condensates of each construct with or without the addition of total mRNA (p values: PGL-3: p = 1.04E-6; PGL-3 (1-452): p = 0.172; PGL-3 (370-693): p = 0.940; PGL-3 (K126E K129E): p = 0.828). The significance score was determined by the following criteria: ns: p > 0.05; ****: p ≤ 0.0001.
c Cartoon showing that at physiological concentrations, mRNA facilitates PGL-3 condensate formation. Association among folded N-termini of PGL-3 molecules is necessary for phase separation.

At physiological concentrations, mRNA facilitates this process by binding to the disordered C-terminus of PGL-3 (Fig. 2c).

α-helicity of PGL-3 correlates with dynamics in condensates in vitro.

P granules have a complex composition, containing mRNA and over forty proteins (Updike & Strome, 2010). It is unclear how dynamics within the P granule phase depends on interactions among those components. Two constitutive P granule proteins, PGL-1 and GLH-1, have been reported to exchange rapidly between the cytoplasm and liquid-like P granules in *C. elegans* gonads (FRAP half-time ($t_{1/2}$) = 7 – 28 s) (Brangwynne *et al*, 2009). Interestingly, in spite of their vastly reduced compositional complexity, we found that condensates generated in vitro with the PGL-3 protein alone show similar dynamics ($t_{1/2}$ = 7 s) (Supplementary Fig. 2c). To investigate the role of other P granule components on dynamics of PGL-3, we added mRNA and recombinantly purified PGL-1 and GLH-1 to condensates of PGL-3 in vitro at physiologically relevant proportions compared to PGL-3 (Saha *et al*, 2016). We found that PGL-1, GLH-1 and mRNA concentrate within

condensates of PGL-3 in vitro (Supplementary Fig. 4d-f), but have a modest effect on the

dynamics of PGL-3 ($t_{1/2}$ = 10 s) (Supplementary Fig. 2c). This suggests that the dynamics of

full length PGL-3 are intrinsic to the protein, and that the influence of complex composition

on dynamics is modest.

In order to address the interactions that drive dynamics within condensates, we began by

probing the dynamics of several PGL-3 constructs within phase-separated condensates in

vitro. The rate at which molecules of two PGL-3 constructs 1-452 and 1-577, that either lack

the disordered C-terminal region entirely or in part, rearrange within condensates are close

to full-length PGL-3 (Fig. 1e, Supplementary Fig. 2g, 3a, b). This suggests that interactions

involving the disordered C-terminal region of PGL-3 are not essential for the fast dynamics

within condensates. Therefore, we addressed the role of the N-terminal α-helical region (1-

452) in driving dynamics. In order to avoid engineering mutations that result in significant

misfolding of PGL-3 and concomitant loss of its ability to phase separate, we focused our

mutational analysis close to the junction of the folded N-terminus and the disordered C-

terminus of PGL-3. Surprisingly, we found that a full-length PGL-3 construct (Δ425-452) that

lacks only 27 residues phase separates into aggregates (Fig. 3a, c). Sequence analysis of

the PGL-3 protein predicts that this region 425-452 spans two α-helices (one complete helix

and fraction of a second helix) (Supplementary Fig. 3d). We generated a PGL-3 construct

(hereafter called 'S1') (Fig. 3a) in which the sequence in the region, 425-452, is shuffled

while keeping the overall amino acid composition unchanged. We found that S1 phase

separates into condensates that are 20- fold less dynamic than with wild-type PGL-3 (Fig.

3d, Supplementary Fig. 3c). Slow dynamics within condensates is not a result of

aggregation of S1 molecules, since condensates of S1 dissolve without detectable trace of

aggregates on addition of excess salt (Supplementary Figure 3j). Following assembly via

phase separation, the extremely slow dynamics in condensates of S1 does not change

significantly over 24 h (Supplementary Fig. 3f). Taken together, our data suggests that

interactions involving the N-terminal α-helical region of PGL-3 are required for fast

dynamics within condensates.

For more insights into the role of the N-terminal α-helical region in regulating dynamics

within condensates of PGL-3, we investigated the contribution of the region 425-452 further.

CD spectroscopy showed that α-helicity is reduced in the mutant S1 compared to wild-type

PGL-3 (Supplementary Fig. 3g). One possibility is that sequence-shuffling of residues 425-

452 changes the tertiary structure of PGL-3. Interestingly, we found that the mutant S1 can

phase separate at concentrations lower than wild-type PGL-3 (Supplementary Fig. 3e).

Further, mutation of two residues (K126E K129E), that interfere with interactions among the

N-termini of wild-type PGL-3 molecules (Aoki *et al*, 2021), also inhibit phase separation of

the S1 mutant even at high protein concentrations up to ~50 μM (Fig. 3a, Supplementary

Fig. 3c, e). This suggests that, regardless of change in tertiary structure, interactions among

the N-termini of S1 molecules are important for phase separation. In order to address the

consequence of sequence-shuffling of residues 425-452 further, we compared the

structural stability of S1 with wild-type PGL-3. We tested this through thermal unfolding

which monitor α-helical content between 20 °C and 90 °C, using the CD signal at 222 nm

as a measure of α-helical content. We found that wild-type PGL-3 unfolds in two phases as

temperature is increased (T_m = 47 °C and 61 °C) (Fig. 3b). On the other hand, S1 also

unfolds in two phases but sequence-shuffling results in lower thermostability (T_m = 41 °C

and 61 °C) (Fig. 3b). Taken together, our observations on the phase separation behavior of

the S1 mutant raises the possibility that loss of α-helicity of PGL-3 may correlate with

slower dynamics in condensates.

In order to address if dynamics in condensates correlate with structural stability of PGL-3,

we compared the dynamics of condensates formed via phase separation of folded and

unfolded PGL-3 constructs. Heat-induced unfolding of both PGL-3 and S1 is irreversible

(Supplementary Fig. 3h, i). These constructs do not recover lost α-helicity once the

temperature is lowered to 21 °C following unfolding at higher temperatures. Therefore, we

heated mEGFP-tagged PGL-3 and S1 proteins at different temperatures for 10 min followed

by cooling to room temperature and triggering phase separation (by diluting salt (KCl)

concentration to 150 mM). Wild-type PGL-3 does not lose α-helicity when heated to 43.5 °C

(Supplementary Fig. 3h). We observed that PGL-3-mEGFP preheated at 43.5 °C generates

condensates that are liquid-like (Fig. 3e). On the other hand, heating S1 to 43.5 °C results

in partial unfolding (Supplementary Fig. 3i), and phase separation of this partially unfolded

S1-mEGFP protein generates solid-like condensates with no observable FRAP recovery

(Fig. 3f). Similarly, heating wild-type PGL-3 to a higher temperature (50 °C) results in partial

unfolding (Supplementary Fig. 3h), and phase separation of this partially unfolded PGL-3-

mEGFP protein generates solid-like condensates with no observable FRAP recovery (Fig.

3e). Taken together, our data suggests that reduced α-helicity of PGL-3 correlates with

slower dynamics in condensates.

Fig. 3 α-helicity of PGL-3
correlates with dynamics in
condensates.
325
a Cartoon representing PGL-3, S1,
S1(K126E K129E) and PGL-3
(Δ425-452). Residues 425-452 of
wild-type PGL-3 are shuffled in
mutants S1 and S1(K126E K129E) .
b Plot of normalized circular
dichroism (CD) spectroscopy signal
at 222 nm with PGL-3 (black circle)
and S1 (blue circle) vs. temperature,
assayed at 5 µM protein
concentration in 25 mM HEPES pH
7.5, 1 M KCl, 1 mM DTT. Error bars
represent 1 SD among three
independent measurements. Arrows
highlight 43.5 °C and 50 °C –
temperatures where partial unfolding
of PGL-3 begins and ends
respectively.
c-d Time-lapse fluorescence
micrographs show fluorescence
recovery after photobleaching in
condensates of **c** PGL-3 (Δ425-452)-
mEGFP or **d** S1 (5% of the pool of
protein was tagged with mEGFP).
Arrowheads show photobleached
area in condensates of PGL-3
(Δ425-452)-mEGFP. Scale bars 5
µm. Shown below the time-lapse
micrographs are plots of average
normalized fluorescence over time **c**
in bleached regions (1.4 µm x 1.4
µm) of PGL-3 (Δ425-452)-mEGFP
condensates (n = 8) or **d** in bleached
regions (1.5 µm x 1.5 µm) of S1 (5%
tagged with mEGFP) condensates (n
= 18). In both plots, normalized
fluorescence over time in bleached
regions (1.5 µm x 1.5 µm) of wild-
type PGL-3 (5% of the pool of
protein tagged with mEGFP)
condensates (n = 18) are also shown
as reference. Error bars represent 1
SD.

e Time-lapse fluorescence micrographs show fluorescence recovery after photobleaching at room temperature in
condensates of PGL-3-mEGFP (preheated at 43.5 °C or 50 °C). Scale bar 5 µm. Shown next to the micrographs is a plot
of average normalized fluorescence over time in bleached regions (1.4 µm x 1.4 µm) within the condensates of PGL-3-
mEGFP preheated at 43.5 °C (black circles, n =8) or 50 °C (grey circles, n = 8). Error bars represent 1 SD.
f Time-lapse fluorescence micrographs show fluorescence recovery after photobleaching at room temperature in
condensates of S1-mEGFP (preheated at 43.5 °C). Scale bar 5 µm. Shown next to the micrographs is a plot of average
normalized fluorescence over time in bleached regions (1.4 µm x 1.4 µm) within the condensates of S1-mEGFP preheated
at 43.5 °C (n =8). Error bars represent 1 SD.

P granule components buffer against reduced dynamics in condensates.

We wanted to see if similar changes in dynamics are seen in P granules in *C. elegans*

worms. To address this, we injected mEGFP-tagged wild-type PGL-3 or S1 proteins into

gonads of adult *C. elegans* and investigated their dynamics. Both PGL-3-mEGFP and S1-

mEGFP target to perinuclear P granules (Fig. 4a). Fluorescence recovery after

photobleaching experiments showed that S1 exchanges between P granules and

cytoplasm at rates similar to wild-type PGL-3 (Fig. 4b, Supplementary Fig 4a). Similarly,

when partially unfolded PGL-3-mEGFP (preheated at 50 °C) is injected into gonads of adult

C. elegans, this protein concentrates in perinuclear P granules (Fig. 4a) and exchanges

between P granules and cytoplasm at rates similar to wild-type PGL-3 (Fig. 4b,

Supplementary Fig 4a). Experiments with microinjected mutant PGL-3 proteins most likely

assay the effect of the P granule phase on high non-physiological dose of partially

denatured PGL-3 proteins. To test the effect on a physiologically expressed mutant PGL-3

protein, we used CRISPR/Cas9 technology to generate a transgenic strain that expresses

the construct PGL-3 (Δ425-452)-mEGFP from the native *pgl-3* locus. We found that PGL-3

(Δ425-452)-mEGFP concentrates in perinuclear puncta (Fig. 4c) and shows dynamics

similar to wild-type PGL-3-mEGFP (Fig. 4e, Supplementary Fig. 4b, Supplementary Movie

1 and 2). Further, we found that fecundity of the strain expressing PGL-3 (Δ425-452)-

mEGFP is similar to the strain expressing PGL-3-mEGFP (Fig. 4d). This suggests that the

P granule phase can prevent slowdown of dynamics and associated loss-of-function

resulting from the expression of the aggregation-prone PGL-3 (Δ425-452)-mEGFP

construct. Taken together, our results suggest that the multicomponent P granule phase

can prevent slowdown of dynamics associated with reduced α-helicity of PGL-3.

To reconstitute the multicomponent nature of the P granule, we purified and added back

other P granule proteins to the PGL-3 phase formed in vitro. Interestingly, addition of

merely a few P granule components (mRNA and recombinantly purified PGL-1 and GLH-1)

to condensates of S1 in vitro enhances the dynamics of S1 almost 20-fold to match the

dynamics observed in condensates of wild-type PGL-3 (Fig. 4f, Supplementary Fig. 4c-e,

5f). Further, we found that dynamics within condensates of S1 can be significantly

enhanced simply by adding wild-type PGL-3 protein (Supplementary Fig. 5a). For better

insight into how the PGL-3 protein supports fast dynamics of S1, we added three different

PGL-3 constructs (the N-terminal α-helical fragment 1-452, the C-terminal fragment 370-

693, and the phase separation-incompetent mutant PGL-3 (K126E K129E)) at different

concentrations to condensates of S1. We found that all three PGL-3 constructs concentrate

within condensates of S1 to different degrees (Supplementary Figure 5c) and enhance the

dynamics of S1 (Supplementary Fig. 5a). Further, addition of the phase separation-

incompetent mutant S1 (K126E K129E) to condensates of S1 also enhanced the dynamics

of S1 within condensates (Supplementary Fig. 5a and 5c). Taken together, our data

suggests that enhanced S1 dynamics in condensates can be supported by a wide variety of

interactions involving other P granule components (PGL-1, GLH-1, mRNA) or different parts

of the PGL-3 protein.

Fig. 4 P granule components promote fast dynamics of PGL-3 constructs with reduced α-helicity.

a Fluorescence micrographs of gonads (pachytene region) in *C. elegans* expressing PGL-3-mCherry (shown in magenta) injected with recombinantly-purified PGL-3-mEGFP, S1-mEGFP, or PGL-3-mEGFP preheated at 50 °C (shown in green). A single confocal section is shown. Scale bar 5 μm.

b Plot showing average normalized fluorescence signal over time in bleached peri-nuclear regions (1.5 μm x 1.5 μm) in wild-type *C. elegans* gonads (pachytene region) injected with recombinantly purified PGL-3-mEGFP (black circles, n = 46), S1-mEGFP (blue triangles, n = 36), or PGL-3-mEGFP preheated at 50 °C (grey circles, n = 37). Error bars represent 1 SD. For each condition, data from 2 different animals injected with a recombinantly purified protein were analyzed. Statistical analysis of the estimated half times of fluorescence recovery is shown in Supplementary Fig. 4a.

c Fluorescence micrograph of gonad (pachytene region) in *C. elegans* expressing PGL-3 (Δ425-452)-mEGFP (shown in green). A single confocal section is shown. Scale bar 5 μm.

d Plot showing total number of progeny of worms expressing PGL-3-mEGFP (n=19) or PGL-3 (Δ425-452)-mEGFP (n=17) at 20 °C. Each black circle represents the number of progenies from a single worm. The mean and 1 SD are also indicated. For statistical analysis, T test was performed (p = 0.858). "ns" indicates no significant difference (p > 0.05) between the number of progenies of the two genotype.

e Plot showing average normalized fluorescence signal over time in bleached peri-nuclear regions in *C. elegans* gonads (pachytene region) expressing either PGL-3-mEGFP (black circles, n = 43) or PGL-3 (Δ425-452)-mEGFP (grey squares, n = 32). Error bars represent 1 SD. For each condition, data from 6 different animals were analyzed. Statistical analysis of the estimated half times of fluorescence recovery is shown in Supplementary Fig. 4b.

f Time-lapse fluorescence micrographs show fluorescence recovery after photobleaching over time in condensates of PGL-3 (5 μM) or S1 (5 μM) (5% of the pool of protein was tagged with mEGFP) in presence of PGL-1 (6.7 μM), GLH-1 (1 μM), and total mRNA (14 ng/μL). Scale bar 5 μm. Shown next to the micrographs is a plot of average normalized fluorescence over time in bleached regions (1.5 μm x 1.5 μm) within the condensates of PGL-3 (black circles, n = 20) or

We next wondered if addition of other P granule components is able to support or rescue a

dynamic state of partially unfolded PGL-3 within condensates. In contrast to S1, which

generates condensates that are liquid-like, partially unfolded PGL-3-mEGFP (preheated at

50 °C) phase separates into non-dynamic solid-like condensates in vitro (Fig. 3e). We

tested the ability of different components to rescue dynamics of partially unfolded PGL-3-

mEGFP. When P granule components (mRNA and recombinantly purified PGL-1 and GLH-

1) are added to partially unfolded PGL-3-mEGFP at concentrations similar to those used

with S1 (5 µM), the PGL-3 protein phase separates predominantly into non-dynamic

condensates (Fig. 4g, Supplementary Fig. 4f, g). However, when a lower concentration of

the partially unfolded PGL-3 protein is used (1 µM), addition of these P granule components

supports its phase separation into liquid-like condensates (Fig. 4g). These results revealed

a concentration-dependence to the extent with which unfolded PGL-3 impacts condensate

dynamics.

We wondered if, similar to partially unfolded PGL-3-mEGFP, the dynamics of S1 also

depends on its concentration. In the assays in which S1 is mixed with different additional

components, 5% of the total pool of S1 is tagged with mEGFP. This allowed us to measure

the dynamics of S1 within condensates using FRAP. Following photobleaching,

fluorescence recovered in bleached region within condensates in two phases – a fast phase

driven predominantly by internal rearrangement of molecules within the condensate, and a

slower phase driven by the exchange with the pool of unbleached molecules outside the

condensate (Supplementary Fig. 6a-f). We focused on the fast phase of recovery and used

single-exponential fits to estimate half-time of internal rearrangement of S1 molecules

within condensates of different composition. On addition of other P granule components

(PGL-1, GLH-1, mRNA) to different concentrations of S1, we found that the half-time of

FRAP recovery of S1 within condensates remains relatively unchanged in spite of almost 5-

fold change in mEGFP-fluorescence signal per unit area within the condensates (Fig. 5a).

Assuming that mEGFP-fluorescence reports on the concentration of S1, this observation

suggests that P granule components can buffer the dynamics of S1 close to rates observed

in condensates of wild-type PGL-3 alone (Fig. 5a, d). However, the dynamics of S1 slows

down above a threshold concentration (9 μM in Fig. 5a), suggesting that P granule

components can buffer the dynamics of S1 only within a range of concentrations in

condensates. We wondered if the observed dynamics buffering effect within condensates of

S1 is limited to the specific set of additives PGL-1, GLH-1 and mRNA. We assayed this by

adding to condensates of S1 the construct PGL-3 (K126E K129E). The mutations K126E

K129E are expected to interfere with interactions among N-termini of S1 and PGL-3 (K126E

K129E) molecules (Aoki *et al*, 2021). We found that addition of this phase separation-

incompetent mutant PGL-3 (K126E K129E) can also buffer dynamics within a 2- fold range

of S1 concentration within condensates (Fig. 5b, Supplementary Fig. 5g). However, the

buffering is different compared to when PGL-1, GLH-1 and mRNA are added. Specifically,

the range of concentrations of S1 in condensates where dynamics is buffered and the

associated half-time of FRAP recovery depend on the additives. We used condensates of

similar size for our analysis (average $\pm$ 1 SD of diameter of condensates are 6.4 $\pm$ 1.7 μm

(Fig. 5a) and 5.9 $\pm$ 0.4 μm (Fig. 5b)). Therefore, dynamics buffering here is likely to

represent similar diffusion rates of S1 within condensates.

For more insights into how addition of different components rescues slowdown of S1

dynamics within condensates, we added wild-type PGL-3, or different PGL-3 or S1

constructs to condensates of S1. These additives titrated the concentration of S1 within

condensates (tracked by S1-mEGFP fluorescence signal per unit area) over a range

spanning concentration observed with condensates of S1 alone and those in presence of

other P granule components (PGL-1, GLH-1, mRNA). Here, we found that the half-time of

FRAP recovery of S1 correlates with mEGFP-fluorescence signal per unit area within the

condensates (Fig. 5c). This suggests that, regardless of the condensate composition, fast

dynamics generally correlate with smaller concentrations of S1. In control assays, we found

that the dynamics of the wild-type PGL-3 protein does not correlate significantly with either

concentration of PGL-3 or the composition of the condensate (Fig. 5d, Supplementary Fig.

5b, d). While fast dynamics in condensates generally correlate with smaller concentrations

of S1 molecules (Fig. 5c), the influence of relative concentrations of macromolecules within

condensates on dynamics buffering remains unclear. To address this, we estimated the

ratio of intra-condensate concentrations of S1 and PGL-3 (K126E K129E) both within and

outside the 'buffering range' (see Fig. 5b). We found that dynamics of S1 molecules is

buffered when the intra-condensate concentration of the additive PGL-3 (K126E K129E) is

roughly equal to or exceeds the concentration of S1 within condensates (Supplementary

Fig. 5h).

Fig. 5 Dynamics of S1 in condensates of different composition.

Analysis of fluorescence recovery after photobleaching within 1.5 µm x 1.5 µm regions in condensates of different composition. Half-time of fluorescence recovery after photobleaching ($t_{1/2}$) in each case was estimated by fitting the data to a single-exponential curve (see methods section for details). **a** Plot of $t_{1/2}$ vs. mean S1-mEGFP fluorescence intensity per unit area (before photobleaching) within condensates. Each point on the plots represents data from a single photobleached region within a condensate of defined composition which includes 5 µM PGL-1, 0.75 µM GLH-1, 11 ng/ µM mRNA and different S1 concentrations indicated as follows. Grey plus: 1 µM S1,

solid hexagon: 2 µM S1, asterisk 3.5 µM S1, open circle: 5.5 µM S1, solid diamond: 7.5 µM S1, solid triangle: 8 µM S1, open square: 9 µM S1. Ten points are plotted for each composition (seven for 1 µM S1) and buffering range indicates a range of S1 concentrations where $t_{1/2}$ does not differ significantly between different conditions.

b Plot of t1/2 vs. mean S1-mEGFP fluorescence intensity per unit area (before photobleaching) within condensates. Each point on the plots represents data from a single photobleached region within a condensate of defined composition which includes 5 µM S1 and different concentrations of PGL-3 (K126E K129E) indicated as follows. Solid diamond: 20 µM PGL-3 (K126E K129E), open circle: 15 µM PGL-3 (K126E K129E), asterisk 10 µM PGL-3 (K126E K129E), solid triangle: 5 µM PGL-3 (K126E K129E), Ten points are plotted for each composition (nine for 20 µM PGL-3 (K126E K129E)) and buffering range indicates a range of S1 concentrations where t1/2 does not differ significantly between different conditions.

c Plot of $t_{1/2}$ vs. mean S1-mEGFP fluorescence intensity per unit area (before photobleaching) within condensates. Each point on the plots represents data from a single photobleached region within a condensate of defined composition as follows. Open circle: S1 alone (10 µM, 5% mEGFP-tagged), solid diamond: S1 (5 µM, 5% mEGFP-tagged) + PGL-3 (370-693) (20 µM), solid asterisk: S1 (5 µM, 5% mEGFP-tagged) + PGL-3 (K126E K129E) (5 µM), solid circle: S1 (5 µM, 5% mEGFP-tagged) + PGL-3 (1-452) (5 µM), solid triangle: S1 (5 µM, 5% mEGFP-tagged) + PGL-1 (6.7 µM) + GLH-1 (1 µM) + total mRNA (14 ng/µL), solid square: S1 (5 µM, 5% mEGFP-tagged) + PGL-3 (5 µM), open square: S1 (5 µM, 5% mEGFP-tagged) + PGL-3 (370-693) (5 µM), grey inverse triangle: S1 (5 µM, 5% mEGFP-tagged) + S1(K126E K129E) (5 µM). Ten points are plotted for each composition. All or part of the data were fit using linear regression (R^2 = 0.68, fit of all data; R^2 = 0.84, fit of data on all compositions excluding S1+ S1(K126E K129E)). Data points for S1+ S1(K126E K129E) are highlighted since S1(K126E K129E) is the only construct added to S1 which contains the shuffled 425-452 region.

d Plot of $t_{1/2}$ vs. mean PGL-3-mEGFP fluorescence intensity per unit area (before photobleaching) within condensates. Each point on the plots represents data from a single photobleached region (1.5 µm x 1.5 µm) within a condensate of defined composition as follows. Open hexagon: PGL-3 alone (10 µM, 5% mEGFP-tagged), open square: PGL-3 (5 µM, 5% mEGFP-tagged) + PGL-3 (370-693) (5 µM), solid asterisk: PGL-3 (5 µM, 5% mEGFP-tagged) + PGL-3 (K126E K129E) (5 µM), solid diamond: PGL-3 (5 µM, 5% mEGFP-tagged) + PGL-3 (370-693) (20 µM), solid triangle: PGL-3 (5 µM, 5% mEGFP-tagged) + PGL-1 (6.7 µM) + GLH-1 (1 µM) + total mRNA (14 ng/µL), solid circle: PGL-3 (5 µM, 5% mEGFP-tagged) + PGL-3 (1-452) (5 µM). Ten points are plotted for each composition and R^2 value for a linear regression fit among all compositions is indicated.

Discussion

To maintain homeostasis, cellular liquid-like compartments must be functionally robust against small changes in composition or denaturation of one or more of its components. The mechanisms that provide such robustness remain unclear. A recent study proposed that liquid-liquid phase separation may reduce variation in concentration of components within condensates (Klosin *et al*, 2020). It remains unclear if LLPS can similarly buffer diffusion rate of components within condensates. Recent studies with simple two-component LLPS systems showed that the behavior of one component can influence the dynamics of another within condensates (Boeynaems *et al*, 2019; Rhine *et al*, 2020; Zhang *et al*, 2015; Alshareedah *et al*, 2021). While components are expected to influence the dynamics of each other within multi-component condensates, it is unclear *a priori* how dynamics of biomolecular condensates with complex composition adapt to misfolding/denaturation of one or more component proteins. Here, we show that the liquid-like P granule phase can rescue slow-down of diffusion rate associated with denaturation of a component protein. Such dynamics-buffering may contribute to functional robustness of cellular liquid-like compartments.

In order to understand how composition supports dynamics-buffering in biomolecular condensates, we began by addressing why denaturation of the PGL-3 protein leads to slower dynamics in condensates. One possibility is that the conformational changes of PGL-3 associated with reduced α-helicity facilitate stronger inter-molecular interactions. For instance, loss of α-helicity could increase the valence of "stickers" leading to additional physical crosslinks among PGL-3 molecules (Choi *et al*, 2020). Stronger inter-molecular interactions could inhibit rapid rearrangement of interacting partners within condensates, resulting in slower dynamics. Alternatively, interactions involving unfolded or partially unfolded PGL-3 molecules could generate inter-polypeptide chain entanglements within condensates resulting in slower dynamics. With diffusion of PGL-3 molecules within condensates, more entanglements could develop over time resulting in further decrease in dynamics. Few observations suggest that slower dynamics within PGL-3 condensates result predominantly from stronger inter-molecular interactions in contrast to inter-chain entanglements. First, dynamics of the mutant S1 protein within condensates does not

change significantly over 24 h. Second, the mutations K126E K129E that interfere with

interactions among PGL-3 or S1 molecules (Aoki *et al*, 2021) help enhance dynamics of S1

molecules within condensates. For instance, when added to condensates of S1, both S1

(K126E K129E) and PGL-3 (K126E K129E) help enhance the dynamics of S1 within

condensates. Addition of S1 (K126E K129E) reduces concentration of S1 within

condensates more than addition of PGL-3 (K126E K129E). However, dynamics of S1 within

condensates is faster in spite of higher concentration with PGL-3 (K126E K129E). Taken

together, these observations support the model that slow dynamics of S1 within

condensates depend on interactions among S1 molecules and suggest that interaction

among S1 molecules is stronger than that between wild-type PGL-3 molecules. Stronger

interaction among S1 molecules is further supported by the observation that S1 molecules

can phase separate at lower concentrations compared to wild-type PGL-3 molecules

(Supplementary Fig. 3e).

Our data is consistent with the model that other regions of S1 molecules cooperate with

residues 425-452 (shuffled) to generate stronger inter-molecular interactions. For instance,

addition of the mutant S1 (K126E K129E) enhances dynamics of S1 within condensates in

contrast to maintaining the slower dynamics observed within condensates of S1 alone. This

suggests that the interactions disrupted by the mutations K126E and K129E also contribute

to slow S1 dynamics. One possibility is that interactions involving the residues K126 and

K129 favor S1 conformations that enhance 425-452 (shuffled)-dependent interactions.

Indeed, the mutations K126E K129E have been reported to interfere with interactions

among N-termini of PGL-3 molecules (Aoki *et al*, 2021). While two self-association domains

within the α-helical N-terminus of PGL-3 have been mapped (Aoki *et al*, 2021, 2016),

structural insights into those associations are limited. However, PGL-3 shares significant

sequence similarity with another protein PGL-1. Crystal structures are available for

fragments of the PGL-1 protein that show the two self-association domains at the N-

terminus are predominantly α-helical and globular in nature (Aoki *et al*, 2016, 2021).

Therefore, one possibility is that shuffling the sequence 425-452 of PGL-3 or heat-induced

unfolding of PGL-3 exposes hydrophobic residues that become available to participate in

inter-molecular interactions. This effect may be enhanced within phase separated

condensates if the local solvent environment favors more extended conformations of the

PGL-3 protein, allowing more inter-molecular interactions compared to within the

surrounding dilute phase. Indeed, recent studies reported that a 129- residue long

intrinsically disordered fragment of the human Tau protein or the prion-like low complexity

domain of hnRNPA1 preferentially sample more extended conformations within phase-

separated condensates compared to the surrounding dilute phase (Majumdar *et al*, 2019;

Bremer *et al*, 2021). Taken together, our data suggests that strength of inter-molecular

interactions regulate dynamics within condensates of PGL-3.

It is generally thought that chaperone proteins or RNA helicases counteract slow-down

of dynamics that may result from denaturation of components or inter-

polypeptide/RNA chain entanglements in biomolecular condensates. For instance, the

ATPase-dependent activity of chaperones is required to dissolve aberrant stress granules

that result from aggregation of misfolding-prone proteins (Ganassi *et al*, 2016; Mateju *et al*,

2017). In contrast, we found that weak interactions among two or more components can

buffer against large change of dynamics within the P granule phase. This effect is distinct

from active processes that depend on ATP hydrolysis. Specifically, we show that presence

of just three P granule components (PGL-1, GLH-1 and mRNA) is sufficient to support a

dynamic state of partially unfolded PGL-3-mEGFP within phase-separated condensates. In

absence of the P granule components, partially unfolded PGL-3-mEGFP phase separates

into non-dynamic condensates. We found that the ability of P granule components to

support a dynamic state of partially unfolded PGL-3-mEGFP depends on the relative

concentrations and hence stoichiometries of PGL-3 and the P granule components. This

suggests that P granule components support dynamics by participating in inter-molecular

interactions with PGL-3-mEGFP molecules. Similarly, different PGL-3 constructs – PGL-3

(K126E K129E), PGL-3 (1-452) and PGL-3 (370-693) enhanced dynamics of S1 molecules

within condensates to different degrees. This may correlate with the varying ability of these

constructs to interact with S1 molecules. We considered the specificity of interactions that

support fast dynamics within phase separated condensates. Our data showing that different

additives (P granule components, or wild-type PGL-3, or PGL-3 constructs that differ in

terms of their ability to fold or self-associate) can all rescue slowdown of dynamics of the

mutant S1 protein within condensates suggest that fast dynamics can be supported by

competing interactions among two or more components (Fig. 6). Future investigations will

reveal the individual contribution of different P granule proteins to dynamics buffering.

Similar to our data with P granules, interactions among components could be a potential

mechanism of storage of misfolding-prone proteins in non-aggregated state within the

liquid-like nucleolus under stress in vivo (Frottin *et al*, 2019). We speculate that these

interactions that support fast dynamics are weak in nature and allow fast dissociation rates.

Taken together, our study supports the possibility that weak interactions among two or

more components prevent aggregation or large change of dynamics within cellular liquid-

like compartments.

An interesting aspect of the dynamics buffering effect is that the dynamics of S1 remain

relatively unchanged regardless of the change of S1 concentration within condensates. The

buffering capacity depends on the additives i.e. the range of intra-condensate S1

concentrations where dynamics is buffered varies with additives. Once the buffering

capacity is exhausted i.e. beyond a threshold intra-condensate S1 concentration, dynamics

of S1 molecules slow down at higher S1 concentrations. More work is required to

understand the underlying mechanism. The "buffered" dynamics of S1 molecules within

condensates also vary with additives. With the additive PGL-3 (K126E K129E), we found

that dynamics of S1 molecules is buffered when the intra-condensate concentration of the

additive is roughly equal to or exceeds the concentration of S1 within condensates

(Supplementary Fig. 5h). This is consistent with the notion that weak heterotypic

interactions with PGL-3 (K126E K129E) molecules mediate dynamics buffering in

condensates of S1. Interestingly, we found that buffered dynamics of S1 molecules is faster

in presence of P granule components (PGL-1, GLH-1 and mRNA) compared to that with the

mutant PGL-3 (K126E K129E) (compare Fig. 5a and 5b). This difference in buffered

dynamics is unlikely simply due to difference in relative concentrations of S1 and additives

within condensates. We found that, with the additive PGL-3 (K126E K129E), the dynamics

of S1 within condensates remains buffered despite almost 3- fold variation in the relative

concentration of S1 and additive (Supplementary Fig. 5h). Future work will reveal if the

buffered dynamics depend on the number of interacting species within condensates (e.g.

four when PGL-1, GLH-1 and mRNA is added to condensates of S1; two when PGL-3

(K126E K129E) is added to S1 condensates) in addition to the strength/nature of inter-

molecular interactions.

Fig. 6 Weak interactions among two or more components promote fast dynamics in condensates.

Cartoon shows how dynamics in condensates change with composition. Each condensate contains one or more of three different macromolecules (depicted as thread, square and hexagon) at different concentrations. Homotypic interactions among threads are strong. Other homotypic and heterotypic interactions are weak. Dynamics generally correlate inversely with concentration of strong interactors in condensates. However, within a range of low concentrations of strong interactors, heterotypic interactions can buffer, at least in part, against significant change of dynamics.

Many cellular non-membrane-bound compartments have been characterized to contain two or more phases with distinct biophysical properties (Banani *et al*, 2017; Feric *et al*, 2016; Jain *et al*, 2016; Lyon *et al*, 2021; Wang *et al*, 2014). Further, within the same liquid-like phase, diffusion rates of two different components may differ from each other (Boeynaems *et al*, 2019; Mathieu *et al*, 2020; Zhang *et al*, 2015). In P granules of *C. elegans* early embryos, a gel-like phase of MEG proteins is known to associate with the liquid-like phase of PGL proteins (Wang *et al*, 2014; Smith *et al*, 2016; Putnam *et al*, 2019). The gel-like phase of MEG proteins is absent from P granules of adult *C. elegans* gonads. Two recent studies suggest that association of the gel-like MEG phase changes the surface tension of the liquid-like PGL phase without changing its viscosity or the diffusion rates of PGL proteins (Folkmann *et al*, 2021; Putnam *et al*, 2019). Taken together, one possibility is that the dynamics-buffering mechanism that we identified here adds stability to the steady-state dynamics of liquid-like phases.

An important part of our study is to show that interactions among the α-helical N-terminal domain of PGL-3 molecules can drive phase separation. Although the unstructured C-terminus is dispensable for phase separation of PGL-3, it fine-tunes the process. First, full-length PGL-3 can phase separate at concentrations lower than the α-helical N-terminus 1-452 alone (Fig. 1c), suggesting that interactions involving the C-terminus enhance the propensity of phase separation. Second, heterotypic interactions among the unstructured C-terminus of PGL-3 molecules and mRNA facilitate phase separation of PGL-3 (Saha *et al*, 2016). Our data is consistent with other studies which found that interactions involving both folded domains and intrinsically disordered regions of proteins drive liquid-liquid phase

separation to assemble many cellular non-membrane-bound compartments (Guillén-Boixet

et al, 2020; Mittag & Parker, 2018; Yang *et al*, 2020).

The concept of robustness is widely understood to be essential for the ability of genetic

circuits to function under a wide variety of perturbations (Félix & Barkoulas, 2015). We

propose that one route to achieving robustness is through condensates that contribute to

spatial and temporal regulation over biochemical reactions. The compositional

heterogeneity-dependent dynamics-buffering mechanism that we identified here could add

robustness to biochemical reactions within condensates against the deleterious

consequences of stresses, short-term effects of chemical modifications, intrinsic mutational

changes, and the problems that arise from accumulating high local concentrations of sticky

molecules. The ability to buffer against these changes to dynamics not only provides

robustness, but these networks of interactions could also provide a source of variation that

enable responsiveness when conditions change.

Materials and Methods

Protein expression and purification

All PGL-3, GLH-1 and PGL-1 constructs were purified recombinantly from insect cells using

a baculovirus infection system. Insect cells were harvested ~72 hours after viral infection

and lysed. Constructs were purified from lysates using a combination of Ni-NTA affinity,

anion exchange and/or size exclusion chromatography as described below.

PGL-3, PGL-1, PGL-3 (370-693) and PGL-3 (K126E K129E): The protocol for purification of

PGL-3-6xHis-mEGFP has been described earlier (Saha et al., 2016). PGL-3-6xHis-RFP,

PGL-1-6xHis-mEGFP, PGL-3 (370-693)-6xHis-mEGFP and PGL-3 (K126E K129E)-6xHis-

mEGFP were purified using the same protocol.

PGL-3 (1-452): Insect cells expressing PGL-3 (1-452)-6xHis-mEGFP were resuspended in

lysis buffer (25 mM HEPES pH 7.5, 300 mM KCl, 0.3 M L-Arginine, 10 mM Imidazole, 1 mM

DTT added with protease inhibitor cocktail Complete EDTA-free (Roche) and lysed using

sonicator. The lysates were centrifuged at 21,000 x g in a JS-13.1 rotor (Beckman-Coulter),

and Ni-NTA Agarose resin (Macherey-Nagel) was mixed with the supernatant to capture

PGL-3 protein molecules. The resin was then washed with 25 mM HEPES pH 7.5, 0.3 M

KCl, 20 mM Imidazole. Finally, the PGL-3 molecules were eluted into 25 mM HEPES pH

7.5, 300 mM KCl, 250 mM Imidazole.

In preparation for anion-exchange chromatography, the eluate from Ni-NTA column

was diluted 1:5 in 25 mM Tris pH 8.0, 1 mM DTT and 5 ml of Q Sepharose resin (GE

Healthcare) was mixed to capture PGL-3 protein molecules. After incubation for ~1 h, the

resin was washed with 25 mM Tris pH 8.0, 1 mM DTT buffers containing 350 mM, 500 mM

and 1 M KCl. The eluates in 25 mM Tris pH 8.0, 1 mM DTT buffers containing 500 mM KCl

and 1 M KCl were pooled together and further purified using size exclusion chromatography

with a TSKgel 5000PW column (Tosoh Bioscience) in 25 mM HEPES pH 7.5, 1 M KCl, 1

mM DTT.

PGL-3 (1-577): Insect cells expressing PGL-3 (1-577)-6xHis-mEGFP were resuspended in

lysis buffer (25 mM HEPES pH 7.25, 300 mM KCl, 10 mM Imidazole, 1 mM DTT added with

protease Inhibitor cocktail set III (Calbiochem) and lysed via passage through Avestin

Emulsiflex C5. The lysates were centrifuged at 20000 rpm in a Ti45 rotor (Beckman-

Coulter), and Ni-NTA Agarose resin (Qiagen) was mixed with the supernatant to capture

PGL-3 protein molecules. The resin was then washed with 25 mM HEPES pH 7.25, 300

mM KCl, 20 mM Imidazole, 1 mM DTT. Finally, the PGL-3 molecules were eluted into 25

mM HEPES pH 7.25, 300 mM KCl, 250 mM Imidazole.

In preparation for anion-exchange chromatography, the eluate from Ni-NTA column

was diluted 1:6 in 25 mM Tris pH 8.0, 1 mM DTT and 5-10 ml of Q Sepharose resin (GE

Healthcare) was mixed to capture PGL-3 protein molecules. After incubation for ~1 h, the

resin was washed with 25 mM Tris pH 8.0, 1 mM DTT buffers containing 100 mM, 200 mM,

350 mM, 500 mM and 1 M KCl. The PGL-3 molecules eluted into the 25 mM Tris pH 8.0, 1

mM DTT buffers containing 500 mM KCl and 1 M KCl.

The eluates were further purified using size exclusion chromatography with HiLoad

16/60 Superdex 200 column (GE Healthcare) in 25 mM HEPES pH 7.5, 0.3 M KCl, 1 mM

DTT. The fractions containing pure PGL-3 (1-577)-6xHis-mEGFP from both runs were

pooled together.

S1 and S1(K126E K129E): Insect cells expressing PGL-3 (S1)-6xHis-mEGFP or PGL-3

(S1(K126E K129E))-6xHis-mEGFP were resuspended in lysis buffer (25 mM HEPES pH

7.5, 300 mM KCl, 300 mM L-Arginine, 10 mM Imidazole, 1 mM DTT added with protease

inhibitor cocktail Complete EDTA-free (Roche) and lysed using sonicator. The lysates were

centrifuged at 21,000 x g in a JS-13.1 rotor (Beckman-Coulter), and Ni-NTA Agarose resin

(Macherey-Nagel) was mixed with the supernatant to capture PGL-3 protein molecules. The

resin was then washed with 25 mM HEPES pH 7.5, 1 M KCl, 20 mM Imidazole. Finally, the

PGL-3 molecules were eluted into 25 mM HEPES pH 7.5, 1 M KCl, 250 mM Imidazole.

Next, 500 units of Benzonase nuclease and 1 mM each of DTT and $MgCl_2$ were added to

the eluate and dialyzed overnight into 25 mM HEPES pH 7.5, 1 M KCl, 1 mM DTT, 1 mM

$MgCl_2$.

In preparation for anion-exchange chromatography, the dialysate was diluted 1:13 in

25 mM Tris pH 8.0, 1 mM DTT and 5-10 ml of Q Sepharose resin (GE Healthcare) was

mixed to capture PGL-3 protein molecules. After incubation for ~1 h, the resin was washed

with 25 mM Tris pH 8.0, 350 mM KCl, 1 mM DTT and finally the PGL-3 molecules were

eluted into 25 mM Tris pH 8.0, 1 M KCl, 1 mM DTT.

The eluates were further purified using size exclusion chromatography with a TSKgel

5000PW column (Tosoh Bioscience) in 25 mM HEPES pH 7.5, 1 M KCl, 1 mM DTT.

PGL-3 (Δ425-452): Insect cells expressing PGL-3 (Δ425-452)-6xHis-mEGFP were

resuspended in lysis buffer (25 mM HEPES pH 7.25, 300 mM KCl, 300 mM L-Arginine, 10

mM Imidazole, 1 mM DTT, 10 mM MgCl$_2$ added with protease Inhibitor cocktail set III

(Calbiochem) and Benzonase nuclease) and lysed via passage through Avestin Emulsiflex

C5. The lysates were centrifuged at 20000 rpm in a Ti45 rotor (Beckman-Coulter), and Ni-

NTA Agarose resin (Qiagen) was mixed with the supernatant to capture PGL-3 protein

molecules. The resin was then washed with 25 mM HEPES pH 7.25, 1 M KCl, 40 mM

Imidazole, 1 mM DTT. Finally, the PGL-3 molecules were eluted into 25 mM HEPES pH

7.25, 1 M KCl, 250 mM Imidazole, 1 mM DTT. Next, ~250 units of Benzonase nuclease and

10 mM MgCl$_2$ were added to the eluate and dialyzed overnight into 25 mM HEPES pH 7.25,

1 M KCl, 1 mM DTT, 10 mM MgCl$_2$.

In preparation for anion-exchange chromatography, the dialysate was diluted 1:12 in

25 mM Tris pH 8.0, 1 mM DTT and 5-10 ml of Q Sepharose resin (GE Healthcare) was

mixed to capture PGL-3 protein molecules. After incubation for ~1 h, the resin was washed

with 25 mM Tris pH 8.0, 320 mM KCl, 1 mM DTT and the PGL-3 (Δ425-452) molecules

were eluted into 25 mM Tris pH 8.0, 500 mM KCl, 1 mM DTT. Finally, the eluate containing

purified PGL-3 (Δ425-452)-6xHis-mEGFP was dialyzed into 25 mM HEPES pH 7.5, 1 M

KCl, 1 mM DTT.

GLH-1: Insect cells expressing GLH-1-6xHis-mEGFP were resuspended in lysis buffer (25

mM HEPES pH 7.5, 300 mM KCl, 10 mM Imidazole, 1 mM DTT added with protease

inhibitor cocktail Complete EDTA-free (Roche) and lysed using sonicator. The lysates were

centrifuged at 21,000 x g in a JS-13.1 rotor (Beckman-Coulter), and Ni-NTA Agarose resin

(Macherey-Nagel) was mixed with the supernatant to capture PGL-3 protein molecules. The

resin was then washed with 25 mM HEPES pH 7.5, 300 mM KCl, 20 mM Imidazole. Finally,

the GLH-1 molecules were eluted into 25 mM HEPES pH 7.5, 300 mM KCl, 250 mM

Imidazole.

In preparation for anion-exchange chromatography, the eluate was diluted 1:6 in 25

mM Tris pH 8.0, 1 mM DTT and 4 ml of Q Sepharose resin (GE Healthcare) was mixed to

capture GLH-1 protein molecules. After incubation for ~30 min, the resin was washed with

25 mM Tris pH 8.0, 1 mM DTT buffers containing 200 mM, 250 mM, 300 mM, 350 mM, 400

mM, 500 mM and 1 M KCl. The fractions with 250 to 350 mM KCl were pooled.

Pooled fractions were further purified using size exclusion chromatography with a

HiLoad 16/60 Superdex 200 column (GE Healthcare) in 25 mM HEPES pH 7.5, 300 mM

KCl, 1 mM DTT.

Untagged PGL-3, S1, PGL-3 (1-452), PGL-3 (370-693), PGL-3 (K126E K129E), S1 (K126E

K129E), PGL-1 and GLH-1 were purified by cleaving off 6xHis and mEGFP tags from the

C-terminus of the recombinant proteins using 6xHis-tagged TEV protease, followed by

incubation with Ni-NTA Agarose resin to remove 6xHis-tagged mEGFP and TEV protease.

Size exclusion chromatography and multi-angle light scattering

Molecular weight analysis of PGL-3 (1-452) was carried out using an OMNISEC multi-

detector instrument from Malvern Panalytical. All measurements were performed in 25 mM

HEPES, 300 mM KCl, 1 mM TCEP at 25°C. Prior to measurement, samples were

centrifuged at 10 000 g for 10 min. 90 µl of sample at 0.5 mg/ml was injected onto a

Superose 6 10/300 column (Cytiva). The eluate from the column was fed into a refractive

index detector followed by a right-angle light scattering detector (640 nm). Data were

analysed using the OMNISEC software (v11.10) using a refractive index increment (dn/dc)

of 0.185 ml/g. Bovine serum albumin (PierceTM) was used to calibrate the detectors.

Samples were measured in duplicate.

Labeling of proteins with fluorescent dye

In order to generate fluorescently labeled S1, PGL-1 and GLH-1, the proteins were first

dialyzed into buffer containing 1 mM TCEP and then incubated with AlexaFluor 594 C5-

maleimide (Invitrogen) in 25 mM HEPES pH 7.5, 300 mM KCl (1 M KCl for S1), 1 mM

TCEP for 2 h at room temperature. To remove unbound dye, repeated buffer exchange was

performed with 25 mM HEPES pH 7.5, 300 mM KCl (1 M KCl for S1), 1 mM TCEP until no

fluorescence was detected in the flow through.

Preparation of mRNA used in experiments

In order to track the localization of mRNA in our assays, fluorescently labeled mRNA was generated using in vitro transcription. Coding sequence of the act-4 gene from *C. elegans* was cloned to be transcribed from a T7 promoter. To generate in vitro transcribed act-4 mRNA, T7 RNA polymerase (New England BioLabs) was used and ribonucleotide solution mix was doped with 30% Aminoallyl-UTP-Cy5 (Jena Bioscience). The reaction was incubated for 1 h at 37 °C, then treated with DNase II and mRNA was finally purified using RNeasy Mini Kit (Qiagen). For all the other experiments, total mouse liver mRNA (Takara) was used.

Imaging and fluorescence recovery after photobleaching (FRAP) experiments

Phase separation properties of PGL-3 constructs under various conditions were imaged within chambers created by attaching cover glass to glass slide with imaging spacer between them (Grace Bio-Labs SecureSeal, Sigma Aldrich). 4 µL of solution was placed at the center of the imaging chamber, such that the spacer had no contact with the solution. PGL-3 (1-452) (Fig. 1b) was imaged directly on the cover glass (without glass slide), as condensates of PGL-3 (1-452) were not stable when imaged within the glass chamber. Addition of 1% crowding agent PEG-35000 stabilized condensates of this N-terminal fragment of PGL-3 allowing imaging over time (Fig. 1c-e, 2a, 2b, Supplementary Fig. 2d). The surfaces of glass slide and cover glass were passivated through PEG-silanization as described earlier (Alberti et al., 2018).

The images were acquired with an inverted microscope (Olympus IX83) equipped with a spinning-disk scan head (4,000 rpm, Yokogawa CSU-W1), a sCMOS camera (pixel size 11um, Prime 95B Teledyne Photometrics), and a 100x/1.4 oil-immersion objective (Olympus UPLSAPO). For imaging and FRAP experiments, we used Coherent OBIS lasers of wavelengths 405nm (FRAP), 488 nm, 561 nm, and 640 nm. Single confocal plane was imaged over time for all FRAP experiments.

Phase separation properties of the PGL-3 constructs PGL-3 (1-577), PGL-3 (Δ425-452) and FRAP experiments of preheated PGL-3-mEGFP and S1-mEGFP were imaged within flow chambers created by attaching cover glass to glass slide with two double-sided tapes (~60

μm thick) positioned parallel to each other. The surface of the cover glass was passivated

through PEG-silanization or coating with lipids as described earlier (Alberti et al., 2018). An

inverted spinning-disk confocal imaging system was used, fitted with diode-pumped solid-

state LASERs (wavelength 488 nm and 561 nm), Olympus UplanSApochromat 100x 1.4

NA oil-immersion objective, Yokogawa CSU-X1 (5000 rpm) spinning-disk scan head, and

Andor iXon DU-897 BV back illuminated EMCCD camera.

Image analysis

All image analysis was conducted using the Fiji image-processing package (http://fiji.sc/Fiji).

Images used for analysis of FRAP experiments were corrected for camera noise and

uneven illumination error. Intensities of the bleached areas were corrected for bleaching

and normalized to prebleached image. FRAP data was fitted to single exponential function

using Matlab (Supplementary figure 5e). All fits had R^2 value of 0.92 or higher.

FRAP data for the construct S1(K126E K129E) was acquired with the same spinning-disk

confocal imaging system as all other constructs, but a different FRAP filter was used. To

account for any difference in signal intensity resulting from the use of a different FRAP filter,

we compared the fluorescence intensities of S1-mEGFP, PGL-3-mEGFP and a uniformly

green fluorescent slide (Chroma Autofluorescent Plastic Slides) in images acquired using

the two different FRAP filters. This analysis using three different samples all gave similar

values of correction factor around 2.6. We used this correction factor to normalize the

intensities of S1(K126E K129E) and plotted the results in Fig. 5c.

Phase separation assays

For phase separation assays, different recombinantly-purified PGL-3 constructs were used,

and phase separation was triggered by diluting salt (KCl) concentration to either 150 mM or

100 mM in 25 mM HEPES, pH 7.5, 1 mM DTT. Following phase separation, condensates of

PGL-3 (1-452) dissolved rapidly when incubated within glass-based imaging chambers.

Addition of 1% crowding agent PEG-35000 stabilized condensates of this N-terminal

fragment of PGL-3 allowing imaging over time.

In some cases, PGL-3 constructs were preheated at higher temperatures (43.5 °C or 50 °C)

for 10 min, followed by incubation at room temperature for 10 mins and phase separation

via dilution of salt concentration. Incubation at higher temperatures was carried out in a

temperature-controlled water bath or PCR thermocycler. Heating untagged wild-type PGL-3

and the mutant S1 protein at 50 °C or 43.5 °C respectively resulted in aggregation. In

contrast, presence of mEGFP tag reduced the extent of heat-induced aggregation to a large

extent. Therefore, we used mEGFP-tagged PGL-3 constructs in these assays. Following

incubation at 43.5 °C or 50 °C, we centrifuged the samples (5 min at 20,000 g) to separate

the aggregates from the pool of soluble protein in the supernatant. To account for loss of

protein via aggregation, we remeasured the concentration of protein in the supernatant and

used this pool of mEGFP-tagged PGL-3 constructs for our assays.

At physiological concentrations (approx. 1 μM), the PGL-3 protein is unable to phase

separate into condensates in vitro (Saha *et al*, 2016). At these concentrations, mRNA

promotes phase separation of PGL-3 (Saha *et al*, 2016). To assay for mRNA-dependence

of condensate assembly, it is therefore essential to use physiological concentrations of wild-

type or mutant PGL-3 proteins (Figure 2). However, these condensates are generally too

small to assay rate of internal rearrangement of PGL-3 molecules within condensates using

fluorescence recovery after photobleaching experiments. Therefore, to generate large

condensates for measuring internal rearrangement of wild-type or mutant PGL-3 molecules,

we primarily used higher concentrations of these proteins in vitro (binding to RNA is not

essential for phase separation at these concentrations). However, to mimic the in vivo P

granule phase as closely as possible, we generally added constituent proteins in proportion

to their in vivo abundance (Saha *et al*, 2016) as described below.

1005

In assays investigating phase separation of two or more components (i.e. PGL-3, S1,

different PGL-3 constructs, PGL-1, GLH-1 or mRNA), the components were mixed with

each other before phase separation was triggered by diluting the salt (KCl) to final

concentration of 100 mM in 25 mM HEPES pH 7.5, 1 mM DTT. In experiments where we

studied the correlation between concentration of S1 or PGL-3 in condensates and their half-

time of FRAP recovery: 1) all tested mixtures contained the same concentration of S1 (5

μM) or PGL-3 (5 μM) and 5% of the protein pool of S1 or PGL-3 was mEGFP-tagged; 2)

Image acquisition settings were the same for all the experimental groups: 50 ms exposure;

488-laser power set to 100%; bleaching power set to 100%.

1015

We used fluorescently labeled PGL-3 constructs, PGL-1, GLH-1 or act-4 mRNA to check if

they colocalize with S1 or PGL-3 in condensates. All components were mixed with each

other before phase separation was triggered by diluting the salt (KCl) to final concentration

of 100 mM in 25 mM HEPES pH 7.5, 1 mM DTT. In these experiments, 5% of the pool of

PGL-3 or S1 proteins was fluorescently labeled. 100% of the other components in the mix

were labeled with a fluorescent probe. Colocalization of different PGL-3 constructs with

wild-type PGL-3 or S1 was tested by adding the different proteins in equal amounts (final

concentration of each component was 5 µM). In experiments where colocalization of

different P granule components with PGL-3 or S1 was assayed, concentration (or

concentration ratio) of the different components were the same as in FRAP experiments

(6.7 µM PGL-1, and 14 ng/ µL mRNA with 5 µM S1 or 5 µM PGL-3; 1 µM GLH-1 with 5 µM

PGL-3; 2 µM GLH-1 with 10 µM S1).

1028

We tested if phase separation of S1 is reversible. We triggered phase separation of S1 (20

µM) by diluting salt (KCl) concentration to 100 mM in 25 mM HEPES, pH 7.5, 1 mM DTT.

Fluorescence microscopy confirmed presence of condensates of S1. An aliquot containing

pre-formed condensates of S1 was mixed with buffer containing 1 M KCl. Here, we did not

detect condensates or aggregates of the S1 protein. Following addition of 1 M KCl

containing buffer, the final conditions were 6,7 µM S1 in 25 mM HEPES pH 7.5, 700 mM

KCl, 1 mM DTT.

1036

1037   ***C. elegans* strains, maintenance and transgenic strain construction**

1038   *C. elegans* strains were maintained on plates containing nematode growth media seeded

1039   with OP50 bacteria using standard methods (Brenner, 1974). The strains TH561, TH623

1040   and SSA130 were generated using CRISPR/Cas9-based technology (Table 1). Strains

1041   were verified by sequencing of the modified locus.

1042

1043

1044

Table 1. *C. elegans* strains used in this study.

C. elegans strain number	Genotype	Guide RNA sequence	Source
SSA27	N2 (wild-type)	-	CGC
TH561	*pgl-3(dd29[pgl-3::mEGFP]) V*	5'GAAGGCTTAGGAACCTCCAC3'	This study
TH623	*pgl-3(knu99[pgl-3::mCherry]) V*	5'GGATTCTTTGGTGGATCCCG3' 5'TTCTTTGGTGGATCCCGTGG3'	This study (generated by InVivo Biosystems)
SSA130	*pgl3(sds130[pgl-3(Δ425-452)::mEGFP]) V*	5'GATCACAGAGTTCTACGGAT3'	This study

1046

Fecundity assay

We assayed fecundity of the worms TH561 (*pgl-3::mEGFP*) and SS130 (*pgl-3(Δ425-452)::mEGFP*). These animals were grown and maintained at 20 °C throughout the experiment. L4 worms were singled for each strain. Each of these worms was transferred to a fresh plate every 24 h for five days. On the fourth day after transfer to a fresh plate, the number of progenies from each animal was scored. The total number of progenies for each worm was calculated and shown in Fig. 4d.

1054

Microinjection of recombinant proteins into *C. elegans* gonads

Recombinant PGL-3 constructs were microinjected into gonads of young adult *C. elegans* to study 1) their colocalization with P granules, and 2) the effect of the P granule phase on their dynamics. To study colocalization, mEGFP-tagged PGL-3 constructs were injected into gonads of worms expressing PGL-3-mCherry (strain TH623) from the native locus (to track P granules). Concentrations of injected proteins: PGL-3-mEGFP (10 μM), S1-mEGFP (6 μM) and PGL-3-mEGFP preheated at 50 °C (30 μM). To minimize the effect of fluorescent tag on dynamics, FRAP experiments were conducted on N2 worms microinjected with mEGFP-tagged PGL-3 constructs. Concentration of injected proteins: PGL-3-mEGFP (20 μM), S1-mEGFP (25 μM) and PGL-3-mEGFP preheated at 50 °C (30 μM). Worms were immobilized using 2 mM levamisole and imaged immediately following injections.

1067

Circular Dichroism (CD) spectroscopy

CD spectroscopy of PGL-3 constructs was carried out in a quartz cuvette (0.5 mm pathlength) using Chirascan Plus spectrometer (Applied Photophysics) equipped with TC1 temperature controller (Quantum Northwest). To measure the change in α- helical content of PGL-3 at different temperatures (data presented in Fig. 3b), the temperature of the cuvette was varied from 20 °C to 90 °C in 2 °C steps. At each temperature, the system was allowed to equilibrate for 180 s before measurement of CD signal between 200 and 260 nm. To test if unfolding of PGL-3 constructs at higher temperature is reversible once the temperature is lowered (data presented in Supplementary Fig. 3h and i), CD signal was measured initially at 21 °C, followed by measurements at a higher temperature (43.5 °C or 50 °C) and once again after the temperature was lowered back to 21 °C. At each temperature, the system was allowed to equilibrate for 15 min before measurement of CD signal between 200 and 260 nm.

Statistical analysis

Two-tailed t test was performed for comparing two groups of data (number of condensates with or without addition of mRNA in Fig 2b; number of progeny in Fig 4d; estimated half time of recovery of fluorescence signal for bleached P granules in TH561 (*pgl-3::mEGFP*) and SSA130 (*pgl-3(Δ425-452)::mEGFP*) in Supplementary Fig. 4b; estimated half time of recovery of fluorescence signal for bleached condensates of S1 and PGL-3 with addition of PGL-1, GLH-1 and mRNA in Supplementary Fig. 4c). One way ANOVA was performed for comparing three groups of data (estimated half time of recovery of fluorescence signal for bleached P granules for injected proteins: PGL-3, PGL-3 preheated at 50 °C and S1 in Supplementary Fig. 4a). p values are indicated in the figure legends. Statistical tests were performed using GraphPad Prism version 8.4.3 (GraphPad Software, San Diego, California USA). To estimate the half time of recovery for above mentioned statistical analysis, the data points for normalized fluorescence signal for individual P granule/condensate were fit using R (RStudio Version 1.2.1335) to the following equation: $F_t = (F_0 + F_\infty*(t/t_{1/2}))/(1 + t/t_{1/2})$ where F_t is fluorescence signal at time t, F_0 is fluorescence signal immediately after photobleaching (minimal fluorescence signal), F_∞ is maximal recovered fluorescence signal and $t_{1/2}$ is half time of fluorescence recovery (Yguerabide *et al*, 1982). Only high-confidence fits [i.e., goodness of fit ($p < 1E-11$)] were used for statistical tests.

1100

Acknowledgments

We thank A.A. Hyman, R.V. Pappu, A. Goloborodko for feedback on the manuscript and

helpful discussions. We thank A.A. Hyman and C. Hoege for the *C. elegans* strains TH623

and TH561. We thank Jana Sipkova, Anoop Kumar Yadav and Felicitas Arlt for help with

image analysis and technical support. S.S. initiated this study as a postdoctoral fellow in the

lab of A.A. Hyman. We thank the services and facilities at Vienna BioCenter for support -

IMBA BioOptics Facility for support with imaging and image analysis; IMBA Molecular

Biology Service for providing reagents for cloning and recombinant protein purification, and

sequencing of constructs; VBCF Protein Technologies for help with recombinant protein

expression and performing SEC-MALS experiment; IMBA Protein Chemistry Facility for

verification of recombinant proteins using mass spectrometry; VBCF Computational Biology

Training (A. Aszódi) for help with generating scripts for curve fitting. N2 strain was provided

by the CGC, which is funded by NIH Office of Research Infrastructure Programs (P40

OD010440). S.J., J.B., P.C., B.R.P., and S.S. are funded by the Austrian Academy of

Sciences. A.S.H. is scientific advisor at Dewpoint Therapeutics.

1116

1117 **Author contribution:** Conceptualization, S.J. and S.S.; Methodology, S.J., A.S.H. and

1118 S.S.; Investigation, S.J., J.B., P.C., B.R.P., M.R., C.H., S.S.; Writing – Original Draft, S.J.

1119 and S.S.; Writing – Review & Editing, S.J., A.S.H., and S.S.; Funding Acquisition, S.S.;

1120 Resources, A.S.H. and S.S.; Supervision, S.S.

1121

References

Alshareedah I, Thurston GM & Banerjee PR (2021) Quantifying viscosity and surface tension of multicomponent protein-nucleic acid condensates. *Biophysical Journal* 120: 1161–1169

Aoki ST, Kershner AM, Bingman CA, Wickens M & Kimble J (2016) PGL germ granule assembly protein is a base-specific, single-stranded RNase. *Proceedings of the National Academy of Sciences of the United States of America*

Aoki ST, Lynch TR, Crittenden SL, Bingman CA, Wickens M & Kimble J (2021) C. elegans germ granules require both assembly and localized regulators for mRNA repression. *Nat Commun* 12: 996

Banani SF, Lee HO, Hyman AA & Rosen MK (2017) Biomolecular condensates: organizers of cellular biochemistry. *Nature reviews Molecular cell biology* 18: 285–298

Banani SF, Rice AM, Peeples WB, Lin Y, Jain S, Parker R & Rosen MK (2016) Compositional Control of Phase-Separated Cellular Bodies. *Cell* 166: 651–663

Boeynaems S, Holehouse AS, Weinhardt V, Kovacs D, Van Lindt J, Larabell C, Van Den Bosch L, Das R, Tompa PS, Pappu RV, *et al* (2019) Spontaneous driving forces give rise to protein-RNA condensates with coexisting phases and complex material properties. *Proceedings of the National Academy of Sciences of the United States of America* 116: 7889–7898

Boyko S, Surewicz K, & Surewicz WK (2020) Regulatory mechanisms of tau protein fibrillation under the conditions of liquid-liquid phase separation. *Proceedings of the National Academy of Sciences of the United States of America* 117: 31882-31890

Brangwynne CP, Eckmann CR, Courson DS, Rybarska A, Hoege C, Gharakhani J, Jülicher F & Hyman AA (2009) Germline P granules are liquid droplets that localize by controlled dissolution/condensation. *Science* 324: 1729–1732

Bremer A, Farag M, Borcherds WM, Peran I, Martin EW, Pappu RV & Mittag T (2021) Deciphering how naturally occurring sequence features impact the phase behaviors of disordered prion-like domains Biophysics

Brenner S (1974) The genetics of Caenorhabditis elegans. *Genetics* 77: 71–94

Choi J-M, Holehouse AS & Pappu RV (2020) Physical Principles Underlying the Complex Biology of Intracellular Phase Transitions. *Annual review of biophysics* 49: 107–133

Cipriani PG, Bay O, Zinno J, Gutwein M, Gan HH, Mayya VK, Chung G, Chen J-X, Fahs H, Guan Y, *et al* (2021) Novel LOTUS-domain proteins are organizational hubs that recruit C. elegans Vasa to germ granules. *Elife* 10: e60833

Elbaum-Garfinkle S, Kim Y, Szczepaniak K, Chen CC-H, Eckmann CR, Myong S & Brangwynne CP (2015) The disordered P granule protein LAF-1 drives phase

separation into droplets with tunable viscosity and dynamics. *Proc Natl Acad Sci U S A* 112: 7189–7194

Félix M-A & Barkoulas M (2015) Pervasive robustness in biological systems. *Nature Reviews Genetics* 16: 483–496

Feric M, Vaidya N, Harmon TS, Mitrea DM, Zhu L, Richardson TM, Kriwacki RW, Pappu RV & Brangwynne CP (2016) Coexisting Liquid Phases Underlie Nucleolar Subcompartments. *Cell*: 1–13

Folkmann AW, Putnam A, Lee CF & Seydoux G (2021) Regulation of biomolecular condensates by interfacial protein clusters. *Science* 373: 1218–1224

Frottin F, Schueder F, Tiwary S, Gupta R, Körner R, Schlichthaerle T, Cox J, Jungmann R, Hartl FU & Hipp MS (2019) The nucleolus functions as a phase-separated protein quality control compartment. *Science* 365: 342–347

Ganassi M, Mateju D, Bigi I, Mediani L, Poser I, Lee HO, Seguin SJ, Morelli FF, Vinet J, Leo G, *et al* (2016) A Surveillance Function of the HSPB8-BAG3-HSP70 Chaperone Complex Ensures Stress Granule Integrity and Dynamism. *Molecular cell* 63: 796–810

Guillén-Boixet J, Kopach A, Holehouse AS, Wittmann S, Jahnel M, Schlüßler R, Kim K, Trussina IREA, Wang J, Mateju D, *et al* (2020) RNA-Induced Conformational Switching and Clustering of G3BP Drive Stress Granule Assembly by Condensation. *Cell* 181: 346-361.e17

Hanazawa M, Yonetani M & Sugimoto A (2011) PGL proteins self associate and bind RNPs to mediate germ granule assembly in C. elegans. *The Journal of Cell Biology* 192: 929–937

Jain S, Wheeler JR, Walters RW, Agrawal A, Barsic A & Parker R (2016) ATPase-Modulated Stress Granules Contain a Diverse Proteome and Substructure. *Cell* 164: 487–498

Jawerth L, Fischer-Friedrich E, Saha S, Wang J, Franzmann T, Zhang X, Sachweh J, Ruer M, Ijavi M, Saha S, *et al* (2020) Protein condensates as aging Maxwell fluids. *Science* 370: 1317–1323

Kawasaki I, Amiri A, Fan Y, Meyer N, Dunkelbarger S, Motohashi T, Karashima T, Bossinger O & Strome S (2004) The PGL family proteins associate with germ granules and function redundantly in Caenorhabditis elegans germline development. *Genetics* 167: 645–661

Kawasaki I, Shim YH, Kirchner J, Kaminker J, Wood WB & Strome S (1998) PGL-1, a predicted RNA-binding component of germ granules, is essential for fertility in C. elegans. *Cell* 94: 635–645

Klosin A, Oltsch F, Harmon T, Honigmann A, Jülicher F, Hyman AA & Zechner C (2020)
Phase separation provides a mechanism to reduce noise in cells. *Science* 367: 464–
468

Lyon AS, Peeples WB & Rosen MK (2021) A framework for understanding the functions of
biomolecular condensates across scales. *Nat Rev Mol Cell Biol* 22: 215–235

Majumdar A, Dogra P, Maity S & Mukhopadhyay S (2019) Liquid-Liquid Phase Separation
Is Driven by Large-Scale Conformational Unwinding and Fluctuations of Intrinsically
Disordered Protein Molecules. *The Journal of Physical Chemistry Letters* 10: 3929–
3936

Mateju D, Franzmann TM, Patel A, Kopach A, Boczek EE, Maharana S, Lee HO, Carra S,
Hyman AA & Alberti S (2017) An aberrant phase transition of stress granules
triggered by misfolded protein and prevented by chaperone function. *The EMBO
journal* 36: 1669–1687

Mathieu C, Pappu RV & Taylor JP (2020) Beyond aggregation: Pathological phase
transitions in neurodegenerative disease. *Science* 370: 56–60

Mittag T & Parker R (2018) Multiple Modes of Protein-Protein Interactions Promote RNP
Granule Assembly. *Journal of molecular biology* 430: 4636–4649

Phillips CM & Updike DL (2022) Germ granules and gene regulation in the Caenorhabditis
elegans germline. *Genetics* 220: iyab195

Price IF, Hertz HL, Pastore B, Wagner J & Tang W (2021) Proximity labeling identifies
LOTUS domain proteins that promote the formation of perinuclear germ granules in
C. elegans. *Elife* 10: e72276

Putnam A, Cassani M, Smith J & Seydoux G (2019) A gel phase promotes condensation of
liquid P granules in Caenorhabditis elegans embryos. *Nat Struct Mol Biol* 26: 220–
226

Rhine K, Makurath MA, Liu J, Skanchy S, Lopez C, Catalan KF, Ma Y, Ha T, Chemla YR &
Myong S (2020) ALS/FTLD-Linked Mutations in FUS Glycine Residues Cause
Accelerated Gelation and Reduced Interactions with Wild-Type FUS. *Molecular cell*

Riback JA, Zhu L, Ferrolino MC, Tolbert M, Mitrea DM, Sanders DW, Wei M-T, Kriwacki
RW & Brangwynne CP (2020) Composition-dependent thermodynamics of
intracellular phase separation. *Nature* 581: 209–214

Saha S, Weber CA, Nousch M, Adame-Arana O, Hoege C, Hein MY, Osborne Nishimura E,
Mahamid J, Jahnel M, Jawerth L, *et al* (2016) Polar Positioning of Phase-Separated
Liquid Compartments in Cells Regulated by an mRNA Competition Mechanism. *Cell*
166: 1572-1584.e16

Sanders DW, Kedersha N, Lee DSW, Strom AR, Drake V, Riback JA, Bracha D, Eeftens
JM, Iwanicki A, Wang A, *et al* (2020) Competing Protein-RNA Interaction Networks
Control Multiphase Intracellular Organization. *Cell* 181: 306-324.e28

Smith J, Calidas D, Schmidt H, Lu T, Rasoloson D & Seydoux G (2016) Spatial patterning
of P granules by RNA-induced phase separation of the intrinsically-disordered
protein MEG-3. *eLife* 5: e21337

Spike C, Meyer N, Racen E, Orsborn A, Kirchner J, Kuznicki K, Yee C, Bennett K & Strome
S (2008a) Genetic analysis of the Caenorhabditis elegans GLH family of P-granule
proteins. *Genetics* 178: 1973–1987

Spike CA, Bader J, Reinke V & Strome S (2008b) DEPS-1 promotes P-granule assembly
and RNA interference in C. elegans germ cells. *Development (Cambridge, England)*
135: 983–993

Strome S & Wood WB (1982) Immunofluorescence visualization of germ-line-specific
cytoplasmic granules in embryos, larvae, and adults of Caenorhabditis elegans.
Proceedings of the National Academy of Sciences of the United States of America
79: 1558–1562

Updike D & Strome S (2010) P granule assembly and function in Caenorhabditis elegans
germ cells. *Journal of andrology* 31: 53–60

Van Treeck B & Parker R (2018) Emerging Roles for Intermolecular RNA-RNA Interactions
in RNP Assemblies. *Cell* 174: 791–802

Wang JT, Smith J, Chen B-C, Schmidt H, Rasoloson D, Paix A, Lambrus BG, Calidas D,
Betzig E & Seydoux G (2014) Regulation of RNA granule dynamics by
phosphorylation of serine-rich, intrinsically disordered proteins in C. elegans. *eLife* 3:
449–23

Yang P, Mathieu C, Kolaitis R-M, Zhang P, Messing J, Yurtsever U, Yang Z, Wu J, Li Y,
Pan Q, *et al* (2020) G3BP1 Is a Tunable Switch that Triggers Phase Separation to
Assemble Stress Granules. *Cell* 181: 325-345.e28

Yguerabide J, Schmidt JA & Yguerabide EE (1982) Lateral mobility in membranes as
detected by fluorescence recovery after photobleaching. *Biophys J* 40: 69–75

Zhang H, Elbaum-Garfinkle S, Langdon EM, Taylor N, Occhipinti P, Bridges AA,
Brangwynne CP & Gladfelter AS (2015) RNA Controls PolyQ Protein Phase
Transitions. *Molecular cell* 60: 220–230

1262

1263

Supplementary Information

1264

1265

Supplementary Figure 1

1266
1267
SDS-PAGE of recombinantly-purified
proteins used in this study. Four SDS-
PAGE gels are shown (gel 1: lanes 1-13;
gel 2: lanes 14-20; gel 3: lanes 21-23, gel
4: lanes 24-26). Lane 21 is not adjacent
to lanes 22 and 23 in gel 3; lanes 24, 25
and 26 represent non-adjacent lanes in
gel 4.
1276

Supplementary Figure 2

a Sequence of PGL-3 with secondary structure highlighted in different colors as predicted by the web-based PSIPRED 4.0 server. Dark grey β strand, blue: α helix, light grey: coil. (http://bioinf.cs.ucl.ac.uk/psipred/)

b Plot of disorder probability across the length of PGL-3, as predicted by the web-based PrDOS (Protein DisOrder prediction System) server with false-positive rate of 5.0% (http://bioinf.cs.ucl.ac.uk/psipred/).

c Plot of average normalized fluorescence over time in bleached regions (1.5 μm x 1.5 μm) within condensates of PGL-3 alone (black circles, 10 μM, 5% mEGFP-tagged, n = 18), or PGL-3 (blue circles, 5 μM, 5% mEGFP-tagged, n = 21) in presence of PGL-1 (6.7 μM), GLH-1 (1 μM) and total mRNA (14 ng/μL). Error bars represent 1 SD.

d Time-lapse fluorescence micrographs show fluorescence recovery over time after a condensate of PGL-3 (1-452) (5 μM, 5% mEGFP-tagged) was photobleached. Scale bar 5 μm. Below is shown a plot of average normalized fluorescence over time in 11 condensates of PGL-3 (1-452) (21 μM, 5% mEGFP-tagged) that were photobleached. Error bars represent 1 SD.

e Bright field micrograph of PGL-3 (1-452) (5 μM) in 25 mM HEPES pH 7.5, 100 mM KCl, 1 mM DTT. Scale bar 20 μm.

f Representative plot of size exclusion chromatography and multi-angle light scattering (SEC-MALS) data for PGL-3 (1-452). Black: relative refractive index response (Y-axis, left), Grey: molecular weight in kDa (Y- axis, right). Measurements were performed in 25 mM HEPES, 300 mM KCl, 1 mM TCEP. Average molecular weight of PGL-3 (1-452) estimated from SEC-MALS is 109.5 (±1.4) kDa (n = 2). Expected molecular weight based on amino acid sequence is 50.5 kDa, suggesting that PGL-3 (1-452) exists as a dimer.

g Plot of average normalized fluorescence over time in bleached regions (1.5 μm x 1.5 μm) within condensates of PGL-3 (black circles, 5% mEGFP-tagged, n = 18), PGL-3 (1-452) (blue circles, 5% mEGFP-tagged, n = 24) and PGL-3 (1-577)-mEGFP (grey circles, bleached region 1.4 μm x 1.4 μm, n = 8). Error bars represent 1 SD.

a

b

c

d PGL-3 (1-452)

e PGL-3 (1-452)

g

f

Supplementary Figure 3

a Time-lapse fluorescence micrographs show fluorescence recovery over time after a 1.4 µm x 1.4 µm area within a condensate of PGL-3 (1-577)-mEGFP (20 µM) was photobleached. Scale bar 5 µm.
b Plot showing average normalized fluorescence of bleached region (1.4 µm x 1.4 µm) within PGL-3 (1-577)-mEGFP condensates (n=8) over time. Error bars represent 1 SD.
c Bright-field micrographs of S1 (8 µM) in 25 mM HEPES pH 7.5, 100 mM KCl, 1 mM DTT and S1 (K126E K129E) (50 µM) in 25 mM HEPES pH 7.5, 150 mM KCl, 1 mM DTT. Scale bar 20 µm.
d Sequence of PGL-3 (412-455) with secondary structure highlighted in different colors as predicted by the web-based PSIPRED 4.0 server. Light blue: α helix, grey: coil. (http://bioinf.cs.ucl.ac.uk/psipred/). Residues 425-452 (shuffled in S1) are highlighted within open black box.
e Phase diagrams of PGL-3, S1 and S1(K126E K129E) at different protein and salt (KCl) concentrations in 25 mM HEPES pH 7.5, 1 mM DTT. Blue circles represent conditions where phase separation was observed. Black circles represent conditions where phase separation was absent. 5% of the pool of protein was tagged with mEGFP in each case.
f Plot showing average normalized fluorescence of bleached regions (1.5 µm x 1.5 µm) within condensates of S1 (10 µM, 5% mEGFP-tagged) over time. Condensates were assayed at different times after assembly via phase separation and incubation at 20 °C (Black circles: immediately following assembly (n = 16), blue triangles: 2 h post-assembly (n = 14), grey squares: 24 h post-assembly (n = 11). Error bars represent 1 SD.
g Plot of mean residue ellipticity of PGL-3 and S1 vs. wavelength assayed using circular dichroism (CD) spectroscopy. Black circles: PGL-3 (3 µM, 10 mM Potassium phosphate pH 7.5, 300 mM KCl, 1 mM TCEP), Blue triangles: S1 (3 µM, 10 mM Potassium phosphate pH 7.5, 300 mM KCl, 1 mM TCEP). Error bars represent 1 SD among six CD scans.
h-i Plots of circular dichroism signal vs. wavelength of **h** PGL-3 (3 µM) and **i** S1 (3 µM) in 25 mM HEPES pH 7.5, 1 M KCl and 1 mM DTT at different temperatures. Error bars represent 1 SD among six CD scans. Black circles: 21 °C, blue squares: 43.5 °C, blue triangles: 21 °C (after preheating at 43.5 °C), grey squares: 50 °C, grey triangles: 21 °C (after preheating at 50 °C).

1383 **j** Fluorescence micrographs show increasing salt (KCl) concentration dissolve pre-formed condensates of the S1 protein.
1384 5% of total pool of the S1 protein was tagged with mEGFP. Condensates of S1 protein (20 µM) were prepared in 25 mM
1385 HEPES pH 7.5, 100 mM KCl, 1 mM DTT. After dilution with buffer containing 1 M KCl, the final concentration of S1 was
1386 6,7 µM in 25 mM HEPES pH 7.5, 700 mM KCl, 1 mM DTT. Scale bar 20 µm.
1387

Supplementary Figure 4

a-c Plots with estimated half times of recovery with statistical analysis for three experiments. Each black circle represents estimated half time (s) for one bleached P granule/condensate. The mean and 1 SD are also indicated. Half times were estimated by fitting the data points for normalized fluorescence signal for individual P granule/condensate to the equation: $F_t = (F_0 + F_\infty*(t/t_{1/2}))/(1 + t/t_{1/2})$. Only points with very confident fit (p<1E-11) were considered for the statistical analysis (described in more detail in Materials and Methods section). "ns" indicates no significant difference (p > 0.05) between the half times of recovery. Experiments were as following: **a** Bleaching of the P granules in live animals of N2 strain after PGL-3 (n = 46), preheated PGL-3 (n = 29) or S1 (n = 7) protein injection (Fig. 4b); One way ANOVA performed to compare the groups with following p values: p(PGL-3 vs. preheated PGL-3) = 0.241; p(PGL-3 vs. S1) = 0.926; p(preheated PGL-3 vs. S1) = 0.417 ; **b** Bleaching of the P granules in live animals in the strains expressing PGL-3 (Δ425-452)-mEGFP (n = 19) or PGL-3-mEGFP (n = 19) (Fig. 4b). T test performed to compare the two groups with resulting p value p = 0.205 **c** Bleaching of the condensates of PGL-3 (5 μM) (n=21) or S1 (5 μM) (n=20). 5% of the pool of protein was tagged with mEGFP) in presence of PGL-1 (6.7 μM), GLH-1 (1 μM), and total mRNA (14 ng/μL). T test performed to compare the two groups with resulting p value p = 0.104

d-g Fluorescence micrographs show concentration of PGL-1, GLH-1 and act-4 mRNA (in vitro transcribed) within condensates of PGL-3 **d**, S1 **e**, PGL-3-mGFP **f** and PGL-3-mEGFP preheated at 50 °C **g**. To track condensates of PGL-3 and S1 in **d-e**, 5% of the pool of PGL-3 or S1 proteins used were tagged with RFP, mEGFP or Alexa 594. Fluorophores used to track PGL-1 (mEGFP or Alexa 594), GLH-1 (mEGFP or Alexa 594), and mRNA (Cy5) are specified in each case. Scale bars **d-f** 20 μm, **g** 10 μm.

a Plot of average normalized S1-mEGFP fluorescence over time in bleached regions (1.5 µm x 1.5 µm) within the condensates of defined composition as follows. Black circles: S1 alone (10 µM, 5% mEGFP-tagged) (n = 18), light green triangle: S1 (5 µM, 5% mEGFP-tagged) + PGL-3 (370-693) (20 µM) (n = 24), blue square: S1 (5 µM, 5% mEGFP-tagged) + PGL-3 (1-452) (5 µM) (n = 16), light blue hexagon: S1 (5 µM, 5% mEGFP-tagged) + PGL-3 (5 µM) (n = 20), green triangle: S1 (5 µM, 5% mEGFP-tagged) + PGL-3 370-693) (5 µM) (n = 11), orange diamond: S1 (5 µM, 5% mEGFP-tagged) + PGL-3 (K126E K129E) (5 µM) (n = 17), magenta open circle: S1 (5 µM, 5% mEGFP-tagged) + S1 (K126E K129E) (5 µM) (n = 13). Error bars represent 1 SD.

b Plot of average normalized PGL-3-mEGFP fluorescence over time in bleached regions (1.5 µm x 1.5 µm) within the condensates of defined composition as follows. Black circles: PGL-3 alone (10 µM, 5% mEGFP-tagged) (n = 18), light green triangle: PGL-3 (5 µM, 5% mEGFP-tagged) + PGL-3 (370-693) (20 µM) (n = 12), blue square: PGL-3 (5 µM, 5% mEGFP-tagged) + PGL-3 (1-452) (5 µM) (n = 28), green triangle: PGL-3 (5 µM, 5% mEGFP-tagged) + PGL-3 (370-693) (5 µM) (n = 13), orange diamond: PGL-3 (5 µM, 5% mEGFP-tagged) + PGL-3 (K126E K129E) (5 µM) (n =12). Error bars represent 1 SD.

c-d Fluorescence micrographs show concentration of PGL-3-mEGFP, PGL-3 (1-452)-mEGFP, PGL-3 (370-693)-mEGFP, PGL-3 (K126E K129E)-mEGFP or S1(K126E K129E)-mEGFP within condensates of S1 **c** or PGL-3 **d**. To track condensates of S1 and PGL-3 in **c-d**, 5% of the pool of S1 or PGL-3 proteins used were tagged with Alexa 594 or RFP. Scale bar 10 µm.

e A representative plot of normalized PGL-3-mEGFP fluorescence over time in bleached regions (1.5 µm x 1.5 µm) within a condensate and corresponding single exponential fit. FRAP data was fitted to the single exponential function: y = a * (1 - exp(-b*x)). R^2 values were $\geq$ 0.92 for all fits.

f Plot of average normalized fluorescence over time in bleached regions (1.5 µm x 1.5 µm) within condensates of PGL-3
alone (black circles, 10 μM, 5% mEGFP-tagged, n = 18), PGL-3 in presence of PGL-1 (6.7 μM), GLH-1 (1 μM) and total
mRNA (14 ng/μL) (grey triangles, 5 μM, 5% mEGFP-tagged, n = 21), S1 alone (blue circle, 10 μM, 5% mEGFP-tagged, n
=18), or S1 in presence of PGL-1 (6.7 μM), GLH-1 (1 μM) and total mRNA (14 ng/μL) (light blue triangle, 5 μM, 5%
mEGFP-tagged, n = 21). Error bars represent 1 SD.
g Plot from Figure 5b shown here with additional data representing condensates with S1 alone. Plot of $t_{1/2}$ vs. mean S1-
mEGFP fluorescence intensity per unit area (before photobleaching) within condensates. Each point on the plot
represents data from a single photobleached region within a condensate of defined composition which includes 5 µM S1
alone (solid circle) or 5 µM S1 and different concentrations of PGL-3 (K126E K129E) indicated as follows. Solid diamond:
20 µM PGL-3 (K126E K129E), open circle: 15 µM PGL-3 (K126E K129E), asterisk 10 µM PGL-3 (K126E K129E), solid
triangle: 5 µM PGL-3 (K126E K129E). Ten points are plotted for each composition (nine for 20 µM PGL-3 (K126E
K129E)).
h Estimation of the ratio of intra-condensate concentrations of S1 and additives for conditions represented in Figure 5b.
Plot showing average mEGFP fluorescence intensity per unit area in condensates containing S1 and PGL-3 (K126E
K129E): from left to right in shades of gray of decreasing intensity represent condensates generated with 5 µM S1 and
PGL-3 (K126E K129E) (20 µM or 15 µM or 10 µM or 5 µM). Different sets of condensates were used to estimate the intra-
condensate concentration of S1 and PGL-3 (K126E K129E) separately, i.e. either 5% mEGFP-tagged S1 or 5% mEGFP-
tagged PGL-3 (K126E K129E) was used. 38 condensates were used to estimate S1 or PGL-3 (K126E K129E)
concentration in each case. Error bars represent 1 SD. Within the buffering range (highlighted using broken lines), $t_{1/2}$ of
S1 within condensates remains unchanged. Estimated ratio of intra-condensate concentrations of S1 and PGL-3 (K126E
K129E) is specified above the plot.
1516

Supplementary Figure 6

a Time-lapse fluorescence micrographs show fluorescence recovery after photobleaching over time in condensates of PGL-3 (10 µM) or S1 (10 µM) (5% of the pool of protein was tagged with mEGFP). Scale bars 5 µm.

b Cartoon defining position with respect to a condensate. This position scheme was used to generate fluorescence-profile plots shown in **c** and **d**.

c-d Profile plots mapping the fluorescence signal at different positions within a condensate of PGL-3 **c** or S1 **d**. Plots shows fluorescence signal in selected area (see **b** at different time points. Over time, fluorescence recovers within bleached regions through a combination of internal rearrangement of molecules within condensates and exchange of molecules between condensates and surroundings. Colors represent different recovery phases: blue for time before bleaching, grey for phase I and black for phase II. Phase I is a fast phase of fluorescence recovery that promotes homogenization of fluorescence signal within the condensate predominantly via internal rearrangement of molecules. Phase II is a slower phase of fluorescence recovery that depend predominantly on additional replenishment of unbleached molecules from the surroundings. Different time points are shown as follows. **c** Blue circle: -1 s (before bleaching), grey circle: 0 s (moment of bleaching), grey diamond: 10 s, grey triangle: 40 s, grey square: 80 s, black diamond: 260 s, black square: 620 s, black triangle: 920 s after bleaching. **d** Blue circle: -1 s (before bleaching), grey circle: 0 s (moment of bleaching), grey diamond: 150 s, grey triangle: 250 s, grey square: 395 s, grey reverse triangle: 876 s, black square: 1236 s, black triangle: 1536 s after bleaching.

e-f Plots of average normalized fluorescence over time in bleached regions (1.5 µm x 1.5 µm) within the condensates of PGL-3 (n = 9) **e** or S1 (n=9) **f** imaged as described in **a**. Error bars represent 1 SD.

Supplementary Movies

Supplementary Movie 1
Time-lapse fluorescence micrographs show fluorescence recovery after photobleaching of P granules in live *C. elegans* (pachytene region) expressing PGL-3-mEGFP. Time stamp and scale bar (3 µm) indicated (related to Fig. 4e).

Supplementary Movie 2
Time-lapse fluorescence micrographs show fluorescence recovery after photobleaching of P granules in live *C. elegans* (pachytene region) expressing PGL-3 (Δ425-452)-mEGFP. Time stamp and scale bar (3 µm) indicated (related to Fig. 4e).