Variation in the ACE2 receptor has limited utility for SARS-CoV-2 host prediction

Nardus Mollentze[1,2], Deborah Keen[1], Uuriintuya Munkhbayar[1], Roman Biek[1], Daniel G. Streicker[1,2]

1. Institute of Biodiversity, Animal Health, and Comparative Medicine, College of Medical, Veterinary, and Life Sciences, University of Glasgow, Glasgow, UK

2. Medical Research Council – University of Glasgow Centre for Virus Research, Glasgow, UK

Abstract

Transmission of SARS-CoV-2 from humans to other species threatens wildlife conservation and may create novel sources of viral diversity for future zoonotic transmission. A variety of computational heuristics have been developed to pre-emptively identify susceptible host species based on variation in the ACE2 receptor used for viral entry. However, the predictive performance of these heuristics remains unknown. Using a newly-compiled database of 96 species we show that, while variation in ACE2 can be used by machine learning models to accurately predict animal susceptibility to sarbecoviruses (accuracy = 80.2%, binomial confidence interval [CI]: 70.8 – 87.6%), the sites informing predictions have no known involvement in virus binding and instead recapitulate host phylogeny. Models trained on host phylogeny alone performed equally well (accuracy = 84.4%, CI: 75.5 – 91.0%) and at a level equivalent to retrospective assessments of accuracy for previously published models. These results suggest that the predictive power of ACE2-based models derives from strong correlations with host phylogeny rather than processes which can be mechanistically linked to infection biology. Further, biased availability of ACE2 sequences misleads projections of the number and geographic distribution of at-risk species. Models based on host phylogeny reduce this bias, but identify a very large number of susceptible species, implying that model predictions must be combined with local knowledge of exposure risk to practically guide surveillance. Identifying barriers to viral infection or onward transmission beyond receptor binding and incorporating data which are independent of host phylogeny will be necessary to manage the ongoing risk of establishment of novel animal reservoirs of SARS-CoV-2.

Introduction

The global spread and high incidence of severe acute respiratory syndrome-associated virus 2 (SARS-CoV-2) has enabled human-to-animal transmission (hereafter, "reverse zoonotic" transmission) in a variety of taxa and environmental contexts. For example, reports have described household transmission to domestic dogs and cats, transmission to lions and tigers in zoos, transmission to farmed mink, and transmission to free-living white-tailed deer (Barrs et al., 2020; Kuchipudi et al., 2022; McAloose et al., 2020; Sit et al., 2020). While the majority of such spillovers are likely to be dead-ends for the virus, repeated introductions followed by onward transmission have been reported in both European mink and North American deer (Kuchipudi et al., 2022; Oreshkova et al., 2020; Oude Munnink et al., 2021). The potential for reverse zoonotic transmission to both threaten wild and domestic animal health and to foster viral evolutionary diversification that could ultimately be re-introduced into humans (Hammer et al., 2021; Oude Munnink et al., 2021) has led to a surge of interest in pro-actively identifying potentially-susceptible species in which surveillance could more efficiently detect reverse zoonotic spillover or sustained transmission of SARS-CoV-2.

Evidence-based prioritisation of SARS-CoV-2 surveillance in animals has proven intractable through traditional approaches. Animal infection experiments have demonstrated that a broad range of species spanning multiple mammalian orders are susceptible (Freuling et al., 2020; Munster et al., 2020; Mykytyn et al., 2021; Schlottau et al., 2020; Shi et al., 2020; Zhao et al., 2020). Yet, infection experiments also suggest fine-scaled variation in susceptibility within mammalian orders. For example, Egyptian fruit bats (*Rousettus aegyptiacus*) can be infected experimentally with SARS-CoV-2, while attempts to establish infection in big brown bats (*Eptesicus fuscus*) have thus far failed (Hall et al., 2020; Schlottau et al., 2020). This combination of wide host range with variable susceptibility within groups creates logistical and financial challenges to establishing susceptibility using experimental infections, since large numbers of taxa would need to be investigated. Cell culture-based compatibility assays, an alternative to in vivo infections for evaluating potential host range, are more scalable, but how well they map to susceptibility remains unclear.

A large number of computational heuristics have been developed to address the challenge of predicting the host range of SARS-CoV-2 in the absence of comprehensive experimental evidence. Most utilise the amino acid sequence of angiotensin-converting enzyme 2 (ACE2, the host receptor involved in entry of SARS-CoV-2) and base predictions on measures of similarity to human ACE2, reasoning that changes relative to this sequence (particularly at amino acid positions known to interact with the SARS-CoV-2 spike protein) decrease the probability of successful infection (Ahmed et al., 2021; Alexander et al., 2020; Damas et al., 2020; Frank et al., 2020; Kumar et al., 2021; Luan et al., 2020; Melin et al., 2020; Qiu et al., 2020). A related structural approach has compared the predicted binding affinity between the modelled ACE2 protein structures of different species and the SARS-CoV-2 spike protein, reasoning that stronger binding supports susceptibility (Huang et al., 2020; Lam et al., 2020; Rodrigues et al., 2020). A final approach trained models to predict the binding affinity of species with known ACE2 sequences using ecological and physiological traits, reasoning that these traits might be correlated with ACE2 evolution and therefore allow projection of binding affinity – and by extension susceptibility – to species without ACE2 sequence information (Fischhoff et al., 2021). A limitation common to all efforts to date is that they have focused on untested surrogates of susceptibility, namely ACE2 similarity or binding affinity. This has two immediate drawbacks. First, how these surrogates map to susceptibility is unknown, leading to the development of largely arbitrary thresholds for deciding risk of infection. Second, since no model has been trained or validated using observed data on infection (the outcome of interest), standard metrics of predictive value (e.g., sensitivity, specificity, accuracy) have not been quantified. There are several reasons to doubt whether models based solely on ACE2 are likely to succeed. Virus replication depends on interactions with a wide range of host proteins, including during cell entry, so ignoring these additional molecular barriers may overestimate host range (Hoffmann et al., 2020; V'kovski et al., 2021). Next, SARS-CoV-2 has been circulating in humans for only a relatively short amount of time, meaning it is unlikely that human ACE2 is the optimal baseline for sequence- or binding affinity-based approaches. Indeed, models using similarity to human ACE2 tend to correctly predict other primates as susceptible to SARS-CoV-2, but frequently fail to predict susceptibility of the rhinolophid bats where this virus is thought to have originated (Boni et al., 2020; MacLean et al., 2021). Understanding the value of ACE2-based host range predictions for guiding surveillance therefore requires developing models based on the outcome of interest – susceptibility – and quantifying the accuracy of their predictions.

The recent accumulation of reports of SARS-CoV-2 spillover to various species, alongside ongoing efforts to characterise the host range of sarbecoviruses experimentally and via virus discovery, provides an exciting opportunity to develop verifiable models of sarbecovirus host range. Here, we assembled a database of compatibility across 96 species and trained sets of machine learning models

to distinguish susceptible from non-susceptible species. We sought to understand the relative predictive power of commonly employed sequence and structural binding-based representations of ACE2 and to retrospectively assess the accuracy of previously published prediction methods against our compatibility database. By incorporating new metrics into model training, we also tested whether the exclusive focus on ACE2 is warranted, and whether predictive performance could be improved by considering evolutionary relationships among hosts. Finally, we used the best-performing models to assess geographic and taxonomic biases in the distribution of predicted sarbecovirus hosts that may arise depending on input data available and explore whether sarbecovirus host range prediction can be used to generate actionable insights for ongoing SARS-CoV-2 surveillance efforts in non-human animals.

Results

Our final dataset contained records of interactions between sarbecoviruses and 96 mammalian and avian host species. Reports included natural infections (N=25) and the outcomes of experimental inoculations of captive animals (N=28), where in both cases subsequent detection of viral RNA in, or virus isolation from, nasal or rectal swabs or faeces was considered evidence of susceptibility. We also separately recorded whether species were shown to shed infectious virus (either through virus isolation or infection of other animals), but available data were limited to 23 species, of which only 17 shed infectious virus (Figure 1). To investigate the utility of in vitro data in the absence of natural infections or in vivo data, we also collected compatibility in cell culture experiments which involved either direct inoculation of cell cultures from each species (N=4) or inoculation of cells expressing heterologous ACE2 (N=39). We treated these different sources of information hierarchically, considering the best available evidence for compatibility or incompatibility in each host species (natural infection > experimental infection > cell culture > heterologous ACE2 cell culture experiments). Importantly, we included records from any strain of *Severe Acute Respiratory Syndrome Coronavirus*, a species which includes SARS-CoV, SARS-CoV-2 and several related strains. Therefore, a key assumption of our work is that susceptibility to different sarbecoviruses overlaps sufficiently to be modelled jointly. In support of this assumption, most susceptible species in our data (88%) have been linked to at least one sarbecovirus known to use the ACE2 receptor, and among the 10 included host species with records of natural SARS-CoV-2 infection, many were also reported to be susceptible to other sarbecoviruses (N=6), primarily SARS-CoV (N=5, supplementary table S1).

To explore the host range of sarbecoviruses and how susceptibility varies within animal groups, we first quantified how susceptible and non-susceptible species were distributed across both a general time-scaled phylogeny based on the accepted patterns of evolutionary divergence among hosts (derived from TimeTree (Kumar et al., 2017)) and a maximum likelihood phylogeny constructed from ACE2 orthologs. Sarbecovirus susceptibility was highly conserved within both host phylogenies (Pagel's λ > 0.686, likelihood ratio test p-values ≤ 0.003; figure 1 A–B). Clustering was particularly strong among non-susceptible species, which tended to be closer to other non-susceptible species than known susceptible host species, both in terms of divergence dates and ACE2 amino acid distances (Figure 1 C–F). This remained true when focusing on mammals only (Wilcoxon rank sum test p-value < 0.001), indicating that results were not driven by the large evolutionary divergence between birds (mostly non-susceptible) and mammals, which comprised the majority of susceptible records. Further, a phylogeny based on all available ACE2 ortholog sequences closely matched the topology and branch lengths of the time-scaled phylogeny (Baker's γ correlation coefficient = 0.934, permutation test p-value < 0.001; Spearman's ρ correlation of pairwise distances = 0.870, Supplementary figure S1). Together, these results show that variation in ACE2 largely mirrors host

evolutionary relationships, but it is still possible that current ACE2-based heuristics for predicting susceptibility to SARS-CoV-2 leverage areas of phylogenetic incongruence.

We therefore trained a series of machine learning models using metrics which captured the breadth of currently-used representations of the information embedded in ACE2 (sequence dissimilarity and binding affinity with SARS-CoV-2 spike). We also tested whether representing information about individual ACE2 amino acid positions could outperform these commonly used representations. Individual positions were represented by either the observed amino acids ("AA categorical"), the physicochemical properties of observed amino acids ("AA properties"), or as a distance to the most common amino acid observed among susceptible species at the same position ("AA consensus distance"). In terms of sensitivity, no ACE2 representation performed markedly better than a null model which randomly assigned species as "susceptible" or "non-susceptible" in proportion to the frequency of susceptible species in the training data (Figure 2). In contrast, specificity varied more dramatically among representations and all outperformed the null model, indicating that ACE2 can rule out non-susceptible species. However, since non-susceptible species were rare in our training data (N=21), only two models (using either the AA consensus distance or AA properties representations) clearly outperformed the null in terms of overall accuracy. The best-performing model (AA consensus distance) correctly predicted the susceptibility or non-susceptibility of 77 of 96 species (80.2%, 95% binomial confidence interval [CI]: 70.8 – 87.6%). Among other models, we found that encoding ACE2 amino acid distances relative to the closest susceptible species ("distance to susceptible") outperformed the widely-used metric of distance to human ACE2 ("distance to humans") in terms of specificity while maintaining high sensitivity. The AA categorical model performed considerably worse than the null across all performance metrics (Figure 2). Combining all representations of ACE2 had little effect on performance (74.0% accurate, CI: 64.0 – 82.4%), suggesting that divergence at key ACE2 amino acids from the susceptible consensus (and most other possible representations of ACE2 sequences) and structural models of binding affinity supply equivalent information. However, mis-classified species differed between individual models, meaning performance was improved by averaging predictions from any two ACE2-based models (ensemble models, Figure 2). This included an ensemble averaging predictions from two iterations of a model trained on the same ACE2 representation, suggesting that these minor improvements in performance stem purely from random variation between models (likely driven by the large numbers of features available for selection in any one representation) and not from unique information in different sequence representations (AA consensus distance / AA consensus distance ensemble, Figure 2).

Figure 1. Phylogenetic clustering of sarbecovirus host susceptibility and shedding data. (A) Species for which susceptibility to infection and shedding of infectious virus have been assessed. Colours indicate the best available evidence, while symbols show the viruses involved. Blank rows in the shedding panel indicate missing data. A composite phylogeny derived from TimeTree indicates evolutionary relationships and median estimated divergence times. (B) Measurements of phylogenetic clustering when considering increasingly relaxed evidence quality thresholds, based on the phylogeny in (A). Outlined circles show likelihood ratio p-values ≤ 0.003, indicating greater clustering of infection records than expected by chance, while the cross indicates a p-value of 0.186. (C–D) Pairwise cophenetic distances between each host and its closest infected (C) or non-infected (D) neighbour. (E–F) Total Grantham distance between the ACE2 sequence of each host and its closest infected (E) or non-infected (F) neighbour. P-values from a Wilcoxon rank sum test are indicated in (C-F), and overlapping values from different hosts are jittered horizontally. Note that while both p-values and boxplots are based on a single value for each host, overlapping symbols at the same position are used to indicate different viruses (for example, a species known to be susceptible to both SARS-CoV [▲] and SARS-CoV-2 [▼] would be indicated using a star formed of overlapping triangles).

High congruence between the overall host phylogeny and the phylogeny of ACE2 (Figure 1, Supplementary figure S1) raised the possibility that successful prediction of sarbecovirus hosts could arise through the correlation of variation in ACE2 with other evolutionarily-conserved features of host biology which define susceptibility, rather than reflecting the disproportionate mechanistic importance of ACE2. Exploiting this correlation would be practically desirable since it would produce models which could be applied to all species, rather than only those with publicly-available ACE2 sequences. We therefore tested models of sarbecovirus susceptibility which used phylogenetic eigenvectors to represent the time-scaled phylogeny for all amniotes (the taxonomic clade including both mammals and birds). The phylogeny-only model performed equivalently to the best ACE2-only models, correctly predicting susceptibility labels of 81 species (84.4%, CI: 75.5 – 91.0%) with a nearly identical sensitivity alongside improved specificity ('phylogenetic eigenvectors', Figure 2). Combining phylogeny with either the best-performing ACE2 representation or all ACE2 representations performed similarly to phylogeny alone (Figure 2), suggesting that ACE2 sequences and measures of binding affinity supply information that is redundant with host divergence patterns. Similar performance was also observed when averaging numeric predictions from individual ACE2-based and phylogeny-only models ("All ACE2 representations/phylogeny ensemble", Figure 2; 81.2% accurate, CI: 72.0 – 88.5%).

Figure 2. *Ability of models trained on different representations of either ACE2 sequences or a time-scaled amniote phylogeny to predict host susceptibility to sarbecovirus infection. Bars represent proportions from leave-one-out cross-validation, with error bars indicating binomial confidence intervals. Dashed vertical lines indicate the performance expected from a null model which randomly assigns susceptibility in proportion to its frequency in the training data (78.12% of available hosts are considered susceptible, N = 96). Dotted vertical lines highlight performance of the best model in each panel.*

Deeper analysis of the features used by our ACE2-only models to classify host susceptibility supported the hypothesis that much of their accuracy derives from correlation with host phylogeny. The model trained with all ACE2-representations (but not phylogeny) retained 25 features, of which just two were clearly linked to virus interaction (both representations of binding affinity, Supplementary figure S2A). The remaining 23 features described information about individual ACE2 positions, primarily in the form of amino acid properties (hydrophobicity, polarity, and Van der Waals

volume). Several of these features represented the same positions (e.g., hydrophobicity and Van der Waals volume of position 752), meaning the model used just 18 positions in the ACE2 alignment (Supplementary figure S2B). Notably, only one of these positions is known to interact with the SARS-CoV-2 spike protein (position 82), but included positions did contain more phylogenetic information than average (as measured by Shannon entropy, Figure 3; Wilcoxon rank sum test p-value < 0.001), suggesting they largely reconstructed host phylogeny. Similar results were observed in the best-performing ACE2-only model (the individual site-based "AA consensus distance" model), which retained no known spike-binding sites (Figure 3). This implies that the inclusion of binding affinity measurements cannot explain the "All ACE2 representation" model's selection of phylogenetically informative sites over known spike-binding sites. It is possible that the model-selected sites were randomly-selected proxies for the same information in correlated spike-binding sites. However, it is unclear why the same sites would have been repeatedly selected from among the large number of correlated alternatives (Figure 3), and why they would be consistently preferred over spike-interacting sites themselves. Further, a model based only on spike-binding sites performed considerably worse than other ACE2-based models (Supplementary figure S3).

Figure 3. Phylogenetic informativeness of all ACE2 amino acid positions available for selection in the model. Positions are stratified by whether or not they form part of any features retained by the combined ACE2-based model (i.e., the model trained with access to all ACE2 representations, left panel) or the best-performing ACE2-based model, which lacked binding affinity information (right panel). Colours indicate whether the site in question is known to interact with the SARS-CoV-2 spike protein (based on annotations in the human reference sequence, accession number NP_001358344.1) or belongs to a cluster of correlated features which also contains at least one such spike-interacting site. Clusters were obtained using affinity propagation clustering of pairwise Spearman correlations between sites. P-values obtained through a Wilcoxon rank sum test. Sequence positions indicated next to selected points refer to locations in the human ACE2 reference sequence. Overlapping points are jittered horizontally.

To ensure that our machine learning approach adequately captured earlier representations of ACE2 sequences, we next compared predictions from our ACE2- and host phylogeny-based models to those of previously published heuristics. Strong correlations between predictions from our combined

ACE2-based model and those from earlier studies confirmed that it was broadly representative of earlier approaches (Supplementary Text, Supplementary figure S4). Direct comparisons of model performance were difficult given the absence of performance metrics for previously published heuristics and wide variation in the number and identity of species for which prediction was attempted. However, performance appeared to be broadly similar across studies, with binding affinity-based predictions performing best (Huang et al., 2020), followed by distance-based and finally host trait-based approaches (Figure 4). Most distance-based approaches achieved their high accuracy by prioritising sensitivity (i.e., accurate prediction of susceptible species, which dominate current data), and had very low specificity (Figure 4B). The performance of our models based on fitted values (the most comparable approach to previous heuristics) was equivalent to that of the best previously published heuristic (Huang et al., 2020). Formal holdout-based measurements of performance for our ensemble and phylogeny-only models also overlapped with the best performing heuristics, further suggesting comparable performance, despite the more stringent evaluation. This identifies predictions based on host phylogeny as a viable alternative to ACE2 sequences for which availability remains limited.

Figure 4. *Performance of existing heuristics on our susceptibility data. (A) Overall accuracy, based on all species in our data for which predictions were available in each study. Accuracy measurements are arranged by increasing sample size, also indicated in colour. A dashed line indicates the performance expected from a null model which assigns labels in proportion to the frequency of susceptible species observed in the full training dataset. (B) Trade-offs between sensitivity and specificity. A dotted line indicates balanced sensitivity and specificity, colours as in (A). For clarity, only studies providing predictions for ≥ 30 species in our data are shown. In both panels, error bars represent binomial confidence intervals. Earlier studies provided no formal metrics of model performance and had a fixed cut-off that was decided post-hoc based on observed susceptibility in the predicted species, rather than from independent data. This makes it impossible to calculate performance metrics directly comparable to the formal holdout-based measures used in this study. To allow comparison, we therefore include the equivalent measures of performance based on training data for our models ('fitted values'), including the best-performing ACE2-based model (AA consensus distance / AA consensus distance ensemble) and the host phylogeny-based model. Note, however, that all such measures will overestimate*

performance on new taxa. Performance metrics based on 'holdout predictions' (Figure 2) provide a more generalizable view of future performance.

Finally, we investigated how including host phylogeny when predicting host range would influence recommendations for targeted surveillance of sarbecovirus spillovers. Our best-performing ACE2-based model, the AA consensus distance/AA consensus distance ensemble, predicted 87.6% of mammals to be susceptible (N=178). This is broadly in line with currently available data, where 82.2% of mammals are reported to be susceptible (N=90; Figure 1). Mapping the distribution of terrestrial wild mammal species predicted to be susceptible by this model revealed an apparent hot-spot in South-East Asia, as previously observed by Fischhoff et al. (2021) (Figure 5C). However, species in this region were not more likely to be susceptible to sarbecoviruses than expected by chance (Figure 5E). Instead, the observed concentration of putatively susceptible species appears to be shaped largely by the increased availability of ACE2 sequences for species from this region (Figure 5A). Indeed, the only regions in which more susceptible species were predicted than expected under a uniform spatial distribution were areas containing very few species with available ACE2 sequences (e.g., the arctic and the Sahara desert).

By allowing predictions on species without available ACE2 sequences, the phylogeny-only model leads to different conclusions on how surveillance might be targeted. This model reduces the proportion of susceptible mammals from 87.6% to 42.7% (N=4183), but this still implies the need for surveillance in 1785 mammal species (Supplementary figure S5). In contrast to the ACE2-based predictions, geographic risk was diffused across most of the tropics and subtropics with apparent hotspots limited to areas with very low species diversity (Figure 5D, F). Thus, this model – which performs comparably to or better than ACE2-based models – suggests that clearly defined geographic hotspots of host susceptibility suitable for targeted surveillance are unlikely to exist. Consistent with observed data, in most mammalian taxonomic orders that contained susceptible species, all included species were predicted to be susceptible (Supplementary figure S6). Interestingly however, primates, rodents, and bats deviated from this pattern, containing several families with no species predicted as susceptible (Supplementary figure S7 – S8). Thus, while the wide host range of sarbecoviruses makes it impractical to specifically target surveillance, it may be possible to exclude some groups from consideration.

Discussion

12
Frequent spillovers of SARS-CoV-2 from human to non-human hosts has led to a surge in the
development of heuristics that seek to predict which species are likely to be susceptible. However,
the predictive power of these heuristics or their utility for guiding research and surveillance
remained unclear. By cataloguing available data on sarbecovirus host range, we show that while a
variety of ACE2-based approaches produce relatively sensitive and specific predictions, these
predictions largely derive from strong correlations with host phylogeny, and limitations and biases in
their input data limit their actionability. Models based on host phylogeny alone perform equivalently,
enabling scalable prediction across nearly all mammals, but imply that in the absence of additional
metrics of inter-species exposures, vast numbers of species would need to be surveyed with limited
geographic or taxonomic focus.

ACE2 sequences showed surprising accuracy for predicting sarbecovirus host range, both in the
formally-trained models produced here and in earlier heuristics. However, various lines of evidence

suggest the predictive power of ACE2-based models derives primarily from phylogenetic correlation. First, susceptibility to sarbecovirus infection was highly conserved, clustering in patterns consistent with the evolutionary history of potential host species. Second, ACE2 is also evolutionarily conserved, and a phylogeny derived from ACE2 sequences was highly congruent with one reflecting broader evolutionary history. Third, commonly-used approaches for representing ACE2 sequence differences are largely equivalent in the amount of susceptibility information carried (Figure 2 & Supplementary figure S4), despite measuring very different aspects of ACE2 sequence variation and its interaction with sarbecovirus spike proteins. Fourth, models based on host evolutionary history performed similarly to the best ACE2-based models, while considering ACE2 in addition to evolutionary history provided no detectable improvements. Finally, in ACE2-based models with access to information about individual amino acid positions, most predictive power was derived from phylogenetically-informative sites rather than virus binding sites, even when measures of binding affinity and sequence similarity were removed. Taken together, these results question the exclusive focus on ACE2 as a predictor of sarbecovirus host range. Biologically, the apparent success of ACE2-based models in predicting sarbecovirus host range cannot be interpreted as evidence for a disproportionate mechanistic influence of ACE2 variation in restricting host range. Instead, sarbecovirus host range is likely to be shaped by a range of factors all showing some level of phylogenetic conservation (e.g., receptor distribution, innate immunity, behaviour). These results also mean caution is needed when interpreting the predictive power of specific sites in ACE2 sequences, as phylogenetic correlation means their biological significance cannot be adequately assessed from observational data.

A key drawback of ACE2-based models is that data are only available for a small fraction of potential host species (Figure 5A). This limits their applied use and may also introduce biases since many ACE2 sequences likely stem from existing sarbecovirus research which is geographically-biased. One previous effort to circumvent the limited availability of ACE2 sequences used host traits believed to follow similar patterns of phylogenetic conservation as ACE2 sequences (Fischhoff et al., 2021). However, this approach had reduced performance relative to models directly incorporating host phylogeny (Figure 4). Further, by conditioning one stage of model development on ACE2 binding, we speculate that this approach may have introduced the same biases plaguing ACE2 sequences, since it recovers similar geographic patterns to our ACE2-based models, which in turn mirror the geographic biases in ACE2 sequence availability. Our results show host phylogeny as a viable alternative, which performs comparably to the best ACE2-based models. However, beyond data availability, the wide host range of sarbecoviruses represents a more significant practical hurdle, since the scope of surveillance suggested by these models is massive. This issue does not stem from the models, which show high accuracy on current data, but rather are representative of a pattern of widespread susceptibility seen consistently across experimental infections (65.2% of mammals susceptible, CI: 42.7 – 83.6%), our wider training data (82.2%, CI: 72.7 – 89.5%), and predictions from both the ACE2-based ensemble and host phylogeny-based models (87.6% and 42.6%, respectively). In essence, the potential host range of sarbecoviruses may be genuinely too broad for accurate predictions to be actionable in the absence of information on the likelihood of exposure.

Given the broad predicted host range of sarbecoviruses, how should research and surveillance actions proceed? While it may be possible to improve predictive performance further by considering variation in additional host proteins linked to infection (e.g., TMPRSS2 (Hoffmann et al., 2020)), this is unlikely to be practically useful given expected correlation with phylogeny and the low numbers of species for which sequences of multiple relevant genes will be realistically attainable. Such an approach is also unlikely to significantly reduce the number of susceptible hosts predicted. Alternatively, surveillance might be better informed by targeting hosts in proportion to their

predicted risk. However, while conditioning our predictions on the top 30% of species captures the majority of currently-known susceptibles (73%), this still means at least 1255 mammals require surveillance while introducing a significant risk of missing key spillovers (Supplementary figure S5). In the interim, surveillance might therefore target species for which local knowledge indicates high human-animal contact, further refined by our model predictions (e.g., knowledge of likely non-susceptible groups, Supplementary figure S6 – S7). Looking further into the future, our data suggest that the majority of susceptible hosts are unlikely to transmit the virus effectively (Figure 1A), and indeed only a limited number of natural spillovers have thus far resulted in onward transmission (Kuchipudi et al., 2022; Oude Munnink et al., 2021). Predicting broader reservoir competence instead of susceptibility to infection may therefore be a path forward to optimise surveillance (Becker et al., 2020). However, the inability of current data to produce models capable of accurately predicting virus shedding (Supplementary figure S9) indicates a need for greater investment to gather sufficient data on the ability of hosts to transmit sarbecoviruses. Careful interrogation of observed spillover events to evaluate sustained transmission (Kuchipudi et al., 2022; Oude Munnink et al., 2021) and inclusion of naïve hosts in experimental infections to track transmission (Freuling et al., 2020; Shi et al., 2020) would be particularly valuable.

A clear limitation of our approach is the relatively small number of species for which sarbecovirus susceptibility data are available. Combining diverse data sources and expanding the scope to all sarbecoviruses allowed us to evaluate the rapidly growing number of untested heuristics predicting SARS-CoV-2 host range. Although our strategy to maximise sample size meant including data from cell culture-based experiments, we found no evidence to suggest that this skewed results (Supplementary text, Supplementary figure S10). Further, while previous approaches were not explicitly designed for non-SARS-CoV-2 sarbecoviruses, our results suggest the inclusion of additional sarbecoviruses did not have any adverse impact (Supplementary text, Supplementary figure S4, S11, S12). However, limited numbers of known non-susceptible species in particular make it difficult to distinguish between alternative approaches with confidence. Available susceptibility data may also be affected by many of the same taxonomic and geographic biases affecting ACE2 sequence availability, since directly-observed data will be biased to countries with capacity for surveillance, in vivo data are limited to animal models that can be maintained in captivity, and heterologous ACE2 experiments are limited by requiring prior knowledge of ACE2 sequences.

Our results imply major challenges for predicting the host range of generalist viruses such as SARS-CoV-2, even when models are reasonably accurate. Both current data and model predictions suggest a significant ongoing risk of reverse zoonosis to a wide range of species, implying a need for continued surveillance among species in contact with humans. Our results provide quantitatively-validated predictions of sarbecovirus susceptibility across all mammals, which together with complementary data on exposure risk could be used to narrow such surveillance. However, they also highlight a data gap which currently precludes models that predict not just susceptibility but competence for onward transmission. Such models will be vital to understand the risk that reverse zoonosis poses for establishing novel reservoirs of SARS-CoV-2 genetic diversity that may compromise global efforts to control the ongoing pandemic.

Methods

Data collection

Data on sarbecovirus host range were collected through literature searches and targeted follow-up of news reports, focusing on identifying reports of natural spillover and experimental infections. These were supplemented with records from cell culture infection assays and two studies which used functional assays in which cells expressing heterologous ACE2 from various species were inoculated with SARS-CoV-2 (Liu et al., 2021; Mykytyn et al., 2021). This allowed us to test the utility of adding in vitro data when information on susceptibility is limited at the organismal level. In all cases, data were treated in a hierarchical manner, preferentially recording susceptibility from the most reliable data type found (natural infection > experimental infection > cell culture > heterologous ACE2 experiment). Data were summarised to the host species level, allowing more reliable data sources to overrule observations from other sources (e.g., an observed natural infection was allowed to overrule negative experimental infections involving the same species, as such experiments may have failed for a variety of reasons such as insufficient dose). Where multiple equivalent studies provided conflicting evidence for the same species (e.g., one successful and one unsuccessful experimental infection), we recorded the species as susceptible, reflecting our desire to predict whether a species had ever been observed as susceptible to any sarbecovirus.

We also recorded whether or not species reported as susceptible were capable of shedding infectious virus, either through reports of onward transmission (e.g., to co-housed naïve individuals) or through isolation of virus from nasal or rectal swabs or from faeces. Species were recorded as negative for virus excretion only when either virus isolation or co-housing with naïve individuals was specifically attempted. These data were available for 23 species.

Data on ACE2 orthologs were obtained from the NCBI Gene database (GeneID: 59272; list of all orthologs downloaded 16 March 2022). This was supplemented by specific searches for host species with available susceptibility data to identify additional ACE2 sequences. For species with susceptibility data but no available ACE2 sequence, we used ACE2 sequences from their closest available relative (based on a time-scaled phylogeny from TimeTree (Kumar et al., 2017)). This replacement affected 21 species in the final dataset, and most (62%) involved replacement sequences from the same genus (Supplementary table S1).

Evaluating phylogenetic clustering

Amino acid sequences for all ACE2 orthologs were downloaded from the NCBI protein database and aligned using the E-INS-i option of MAFFT version 7.471 (Katoh and Standley, 2013). A maximum likelihood phylogeny was then generated using version 1.6.12 of IQ-TREE (Minh et al., 2020), with built-in model selection used to choose the substitution model minimising the Bayesian information criterion (Kalyaanamoorthy et al., 2017). The best model used the Jones, Taylor & Thornton (JTT) substitution matrix (Jones et al., 1992) with empirical amino acid frequencies and six categories of rate variation in the "FreeRate" model of Soubrier et al. (2012).

We also obtained a phylogeny reflecting the estimated divergence dates of all amniotes from TimeTree version 4 (Kumar et al., 2017). To allow comparison between phylogenies, the time-scaled phylogeny was trimmed to species with available ACE2 sequences, before calculating Baker's γ correlation coefficient (Baker, 1974) using version 1.15.1 of the dendextend library in R version 4.1.0 (Galili, 2015; R Core Team, 2021) along with version 2.1.0 of the phylogram library (Wilkinson and Davy, 2018). To calculate the expected null distribution of γ values given these particular phylogenies, species names on both phylogenies were randomly shuffled 1000 times.

To measure the amount of phylogenetic clustering among susceptible and non-susceptible species, we calculated Pagel's λ using version 0.7_80 of the phytools R library (Pagel, 1999; Revell, 2012). Pagel's λ was re-calculated while cumulatively adding data from species with susceptibility or non-susceptibility known from natural infection, experimental infection, cell culture, and finally heterologous ACE2 experiments. We also compared distances to the closest other susceptible or non-susceptible species, using both cophenetic distances and ACE2 amino acid distances. Cophenetic distances were calculated on the time-scaled (TimeTree) phylogeny using version 5.5 of the APE R library (Paradis et al., 2004), while amino acid distances were calculated as the total Grantham distance separating any two species across all sites in the ACE2 alignment (Grantham, 1974).

Sequence representations

Current heuristics for predicting susceptibility to SARS-CoV-2 infection focus on either some measure of ACE2 amino acid distance from its human counterpart or the predicted binding affinity between each ACE2 protein and the SARS-CoV-2 spike protein. To compare such metrics in a common framework, we calculated or obtained a range of amino acid sequence representations that broadly captured current approaches alongside novel alternatives. This included binding affinities from two recent studies (Fischhoff et al., 2021; Huang et al., 2020) ('binding affinity', represented as 2 separate feature columns). Neither study supplied binding affinity measures for all ACE2 sequences in our training data, but combined ensured at least one value for each included sequence. To represent the widely-used distance to human ACE2, we calculated the total amino acid distance between each species' ACE2 protein and that of humans (termed 'distance to humans' in all figures, and calculated as described above, 1 feature). Despite variation in which amino acid positions were included by previous distance-based studies, all measures were highly correlated (Supplementary figure S4), making the total amino acid distance (i.e., the distance calculated while considering all amino acid positions) to human ACE2 a reasonable proxy. Although this metric is commonly-used, whether human ACE2 is the appropriate baseline is unclear given the relatively recent association between SARS-CoV-2 and humans. We therefore also included a similar metric capturing the total amino acid distance to the ACE2 of the closest known susceptible species. This distance was calculated with reference to the relevant training dataset only, and separately expressed the distance to the closest susceptible known from either direct observations, experimental infections, heterologous ACE2 experiments, or across all data ('distance to susceptible', 4 features; data contained no susceptibles known only from cell culture experiments).

We also sought to include more nuanced information about specific ACE2 sites by including representations of individual positions in the ACE2 protein alignment. The amino acids observed at each alignment position showing variation was represented using one-hot encoding of all amino acids observed at a frequency > 10%, alongside an additional binary variable taking a value of 1 for all amino acids observed less frequently at a given position, and 0 otherwise ('AA categorical', 1657 features). We also summarised the amino acids at each variable alignment position using the physicochemical properties hydrophobicity, polarity, net charge, and Van der Waals volume ('AA properties', 1841 features), with values for individual amino acids obtained from the AAindex database (accessions JURD980101, ZIMJ680103, KLEP840101, FAUJ880103) (Fauchère et al., 1988; Juretić et al., 1998; Kawashima et al., 1999; Klein et al., 1984; Zimmerman et al., 1968). Finally, we included the Grantham distance between each observed amino acid and the most common amino acid observed at that site among susceptible species ('AA consensus distance', calculated with reference to the relevant training dataset only, 531 features).

Phylogenetic eigenvectors

The machine learning algorithms we used (described below) cannot directly represent phylogenetic information. Therefore, to include evolutionary relationships among possible hosts in our models, we used phylogenetic eigenvectors calculated from the time-scaled phylogeny described above and including all amniotes. Phylogenetic eigenvectors were calculated using version 0.3-6 of the MPSEM library in R, assuming Brownian motion (Guénard et al., 2013). The first 50 eigenvectors were retained and included in models ('phylogenetic eigenvectors', 50 features).

Machine learning

We trained a series of gradient-boosted classification tree models to predict the susceptibility of individual species using various combinations of ACE2 sequence representations or the phylogenetic eigenvectors. Models were constructed using XGBoost version 1.4.0 in R alongside version 0.1.3 of the tidymodels suite of helper libraries (Chen and Guestrin, 2016; Kuhn and Wickham, 2020). Given the relatively small amount of training data available, performance was evaluated using leave-one-out cross-validation. Throughout, we report proportions of species accurately predicted, alongside binomial confidence intervals calculated using the binom.test function in R (Clopper and Pearson, 1934).

Models were trained in a nested process to allow simultaneous tuning and evaluation of performance. For each species, data were split into training data and the single holdout species for cross-validation ('outer split'). The training data were then randomly split again, keeping 75% for model training with a given combination of tuning parameters, with performance evaluated on the remaining data ('inner split'). This inner split was repeated for each unique tuning parameter combination, testing 100 parameter combinations across a maximum entropy grid (Dupuy et al., 2015). After training an initial model on a given tuning parameter combination, a search was performed for the cutoff which best balanced sensitivity and specificity on the inner split's testing data when converting quantitative scores produced by the model into binary predictions. This was achieved by calculating the mean of sensitivity and specificity ('balanced accuracy') for each cutoff. The tuning parameter and cutoff combination which maximised balanced accuracy was used to train a final model on all training data (i.e., all data except for the single species held back in the outer split). This nested tuning and evaluation process was repeated to generate holdout predictions for all species, allowing calculation of the final reported performance statistics.

Ensemble models

In addition to the direct combination of different ACE2 sequence representations and phylogenetic eigenvectors in the same model, we also tested whether training separate models before combining predictions would improve performance. Models were combined by averaging the quantitative holdout predictions for each species. To convert these averages to binary predictions, we calculated the optimal cutoff balancing sensitivity and specificity as described above, using the averaged quantitative predictions for all species except a given focal species to evaluate potential cutoffs, before using the best cutoff to generate a binary prediction for the focal species. This procedure was repeated to generate predictions for all species.

Feature importance

The importance of individual features in shaping model predictions was measured using SHAP values (Lundberg and Lee, 2017), which measure the contribution of individual features to the final quantitative prediction for a given host species. Using a final model trained on all available data, we calculated feature importance as the mean of absolute SHAP values across all species.

Since many features represented individual ACE2 positions which are expected to co-vary, we also sought to characterise the correlation between positions. For each position in a given ACE2 sequence, we first calculated the Grantham distance between the observed amino acid and the most common amino acid observed at that position across all sequences used in model training. We then calculated Spearman correlations in this distance metric for all possible pairs of ACE2 positions, before using the correlations to perform affinity propagation clustering with version 1.4.8 of the apcluster library in R (Bodenhofer et al., 2011; Frey and Dueck, 2007). The resulting clusters were used to annotate positions in Figure 3, separating known spike binding sites ('S-binding'), sites occurring in the same cluster as a known spike-binding site ('correlated with S-binding site'), and all other sites ('other').

Comparison of other studies

To compare predictions across our models and to those from previously-published heuristics, we collected both binary predictions and the underlying quantitative scores / heuristics used to make these predictions from 12 studies. We compared predictions at the species level. In rare cases where previous studies provided predictions for multiple representatives of a given species (e.g., predictions for subspecies or for different ACE2 sequences of the same species), we used the mean of quantitative scores and the most common binary prediction. Scores from each study were re-scaled to lie between 0 and 1, allowing us to plot them on the same scale. Where needed, scores were reversed to ensure that higher values always reflected species considered more likely to be susceptible.

The similarity of predictions across our models and earlier heuristics was summarised by calculating pairwise spearman correlations, using all overlapping species in each pair of models/heuristics. These correlations were used for agglomerative hierarchical clustering, using the mean correlation across all points as the distance metric between clusters (i.e., the unweighted pair-group arithmetic average [UPGMA] method) in version 2.1.2 of the cluster R library (Kaufman and Rousseeuw, 1990).

For studies which supplied binary or categorical predictions, we also estimated sensitivity, specificity, and overall accuracy, all in light of the new susceptibility dataset generated here. Two studies supplied categorical rather than binary (susceptible / non-susceptible) predictions. In the case of Melin et al. (2020), we treated species predicted as having high susceptibility as susceptible, and those predicted to have low susceptibility as non-susceptible. Similarly, species predicted as having very high, high, or medium susceptibility by Damas et al. (2020) were considered susceptible, treating predictions of low and very low susceptibility as non-susceptible. Importantly, performance estimates for published heuristics relied on predictions as published. Since these heuristics did not include holdout species (i.e., data withheld as an independent test set), cutoffs chosen to judge susceptibility or non-susceptibility to SARS-CoV-2 would have been defined by all species to which the heuristic was applied. This is likely to artificially inflate their performance beyond what could be expected when applying these heuristics to host species that were not involved in the development of the heuristic. In contrast, our approach of withholding data during model development gives a more accurate representation of how performance will generalise to new potential host species.

Mapping predictions

To visualise the spatial distribution of species predicted as susceptible (and of available ACE2 sequences), we summarised IUCN range maps for terrestrial mammals into a grid with cells measuring 1/6 x 1/6 of a degree. Using predictions from models trained on all available data, we counted the number of susceptible species with ranges overlapping each grid cell. We also measured

whether the proportion of predicted susceptible species in any one grid cell was higher than expected by chance by dividing the proportion of species predicted as susceptible in each cell by the overall proportion of species predicted as susceptible across the entire dataset. This meant we were comparing the observed frequency of susceptible species in any one location to a baseline assumption of uniform geographic distribution of susceptible species.

Data availability

Data and code required to reproduce all analyses are available at DOI: 10.5281/zenodo.6552861.

Funding

NM and DGS were supported by a Wellcome Senior Research Fellowship (217221/Z/19/Z) and the Medical Research Council (MC_UU_12014/12). The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Acknowledgements

We thank members of the Streicker lab for feedback on earlier versions of this manuscript.

Author contributions

Conceptualization: NM, RB, DGS

Data curation: NM, DK, UM

Formal analysis: NM

Funding acquisition: DGS

Investigation: NM, DK, RB, DGS

Visualisation: NM

Supervision: RB, DGS

Writing – original draft: NM

Writing – review & editing: NM, RB, DS

References

Ahmed R, Hasan R, Siddiki AMAMZ, Islam MS. 2021. Host range projection of SARS-CoV-2: South Asia perspective. *Infect Genet Evol* **87**:104670. doi:10.1016/j.meegid.2020.104670

Alexander MR, Schoeder CT, Brown JA, Smart CD, Moth C, Wikswo JP, Capra JA, Meiler J, Chen W, Madhur MS. 2020. Predicting susceptibility to SARS-CoV-2 infection based on structural differences in ACE2 across species. *FASEB J* **34**:15946–15960. doi:10.1096/fj.202001808r

Baker FB. 1974. Stability of Two Hierarchical Grouping Techniques Case 1: Sensitivity to Data Errors. *J Am Stat Assoc* **69**:440–445. doi:10.2307/2285675

Barrs VR, Peiris M, Tam KWS, Law PYT, Brackman CJ, To EMW, Yu VYT, Chu DKW, Perera RAPM, Sit THC. 2020. SARS-CoV-2 in Quarantined Domestic Cats from COVID-19 Households or Close Contacts, Hong Kong, China. *Emerg Infect Dis* **26**. doi:10.3201/eid2612.202786

Becker DJ, Seifert SN, Carlson CJ. 2020. Beyond Infection: Integrating Competence into Reservoir Host Prediction. *Trends Ecol Evol* **35**:1062–1065. doi:10.1016/j.tree.2020.08.014

Bodenhofer U, Kothmeier A, Hochreiter S. 2011. APCluster: an R package for affinity propagation clustering. *Bioinformatics* **27**:2463–2464. doi:10.1093/bioinformatics/btr406

Boni MF, Lemey P, Jiang X, Lam TT-Y, Perry BW, Castoe TA, Rambaut A, Robertson DL. 2020.

Evolutionary origins of the SARS-CoV-2 sarbecovirus lineage responsible for the COVID-19 pandemic. *Nat Microbiol* 1–10. doi:10.1038/s41564-020-0771-4

Chen T, Guestrin C. 2016. XGBoost: A Scalable Tree Boosting SystemProceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining, KDD '16. San Francisco, California, USA: Association for Computing Machinery. pp. 785–794. doi:10.1145/2939672.2939785

Clopper CJ, Pearson ES. 1934. The Use of Confidence or Fiducial Limits Illustrated in the Case of the Binomial. *Biometrika* **26**:404–413. doi:10.2307/2331986

Damas J, Hughes GM, Keough KC, Painter CA, Persky NS, Corbo M, Hiller M, Koepfli K-P, Pfenning AR, Zhao H, Genereux DP, Swofford R, Pollard KS, Ryder OA, Nweeia MT, Lindblad-Toh K, Teeling EC, Karlsson EK, Lewin HA. 2020. Broad host range of SARS-CoV-2 predicted by comparative and structural analysis of ACE2 in vertebrates. *Proc Natl Acad Sci* **117**:22311–22322. doi:10.1073/pnas.2010146117

Dupuy D, Helbert C, Franco J. 2015. DiceDesign and DiceEval: Two R Packages for Design and Analysis of Computer Experiments. *J Stat Softw* **65**:1–38. doi:10.18637/jss.v065.i11

Fauchère J-L, Charton M, Kier LB, Verloop A, Pliska V. 1988. Amino acid side chain parameters for correlation studies in biology and pharmacology. *Int J Pept Protein Res* **32**:269–278. doi:10.1111/j.1399-3011.1988.tb01261.x

Fischhoff IR, Castellanos AA, Rodrigues JPGLM, Varsani A, Han BA. 2021. Predicting the zoonotic capacity of mammals to transmit SARS-CoV-2. *Proc R Soc B Biol Sci* **288**:20211651. doi:10.1098/rspb.2021.1651

Frank HK, Enard D, Boyd SD. 2020. Exceptional diversity and selection pressure on SARS-CoV and SARS-CoV-2 host receptor in bats compared to other mammals. *bioRxiv* 2020.04.20.051656. doi:10.1101/2020.04.20.051656

Freuling CM, Breithaupt A, Müller T, Sehl J, Balkema-Buschmann A, Rissmann M, Klein A, Wylezich C, Höper D, Wernike K, Aebischer A, Hoffmann D, Friedrichs V, Dorhoi A, Groschup MH, Beer M, Mettenleiter TC. 2020. Susceptibility of Raccoon Dogs for Experimental SARS-CoV-2 Infection. *Emerg Infect Dis* **26**. doi:10.3201/eid2612.203733

Frey BJ, Dueck D. 2007. Clustering by Passing Messages Between Data Points. *Science* **315**:972–976. doi:10.1126/science.1136800

Galili T. 2015. dendextend: an R package for visualizing, adjusting and comparing trees of hierarchical clustering. *Bioinformatics* **31**:3718–3720. doi:10.1093/bioinformatics/btv428

Grantham R. 1974. Amino Acid Difference Formula to Help Explain Protein Evolution. *Science* **185**:862–864. doi:10.1126/science.185.4154.862

Guénard G, Legendre P, Peres-Neto P. 2013. Phylogenetic eigenvector maps: a framework to model and predict species traits. *Methods Ecol Evol* **4**:1120–1131. doi:10.1111/2041-210X.12111

Hall JS, Knowles S, Nashold SW, Ip HS, Leon AE, Rocke T, Keller S, Carossino M, Balasuriya U, Hofmeister E. 2020. Experimental challenge of a North American bat species, big brown bat (Eptesicus fuscus), with SARS-CoV-2. *Transbound Emerg Dis* **68**:3443–3452. doi:https://doi.org/10.1111/tbed.13949

Hammer AS, Quaade ML, Rasmussen TB, Fonager J, Rasmussen M, Mundbjerg K, Lohse L, Strandbygaard B, Jørgensen CS, Alfaro-Núñez A, Rosenstierne MW, Boklund A, Halasa T, Fomsgaard A,

Belsham GJ, Bøtner A. 2021. SARS-CoV-2 Transmission between Mink (Neovison vison) and Humans, Denmark. *Emerg Infect Dis* **27**:547–551. doi:10.3201/eid2702.203794

Hoffmann M, Kleine-Weber H, Schroeder S, Krüger N, Herrler T, Erichsen S, Schiergens TS, Herrler G, Wu N-H, Nitsche A, Müller MA, Drosten C, Pöhlmann S. 2020. SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* **181**:271-280.e8. doi:10.1016/j.cell.2020.02.052

Huang X, Zhang C, Pearce R, Omenn GS, Zhang Y. 2020. Identifying the Zoonotic Origin of SARS-CoV-2 by Modeling the Binding Affinity between the Spike Receptor-Binding Domain and Host ACE2. *J Proteome Res* **19**:4844–4856. doi:10.1021/acs.jproteome.0c00717

Jones DT, Taylor WR, Thornton JM. 1992. The rapid generation of mutation data matrices from protein sequences. *Bioinformatics* **8**:275–282. doi:10.1093/bioinformatics/8.3.275

Juretić D, Lučić B, Zucić D, Trinajstić N. 1998. Protein transmembrane structure: recognition and prediction by using hydrophobicity scales through preference functions In: Párkányi C, editor. Theoretical and Computational Chemistry, Theoretical Organic Chemistry. Elsevier. pp. 405–445. doi:10.1016/S1380-7323(98)80015-0

Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017. ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods* **14**:587–589. doi:10.1038/nmeth.4285

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Mol Biol Evol* **30**:772–780. doi:10.1093/molbev/mst010

Kaufman L, Rousseeuw PJ. 1990. Agglomerative Nesting (Program AGNES)Finding Groups in Data. Hoboken, New Jersey: John Wiley & Sons, Ltd. pp. 199–252. doi:10.1002/9780470316801.ch5

Kawashima S, Ogata H, Kanehisa M. 1999. AAindex: Amino Acid Index Database. *Nucleic Acids Res* **27**:368–369. doi:10.1093/nar/27.1.368

Klein P, Kanehisa M, DeLisi C. 1984. Prediction of protein function from sequence properties: Discriminant analysis of a data base. *Biochim Biophys Acta BBA - Protein Struct Mol Enzymol* **787**:221–226. doi:10.1016/0167-4838(84)90312-1

Kuchipudi SV, Surendran-Nair M, Ruden RM, Yon M, Nissly RH, Vandegrift KJ, Nelli RK, Li L, Jayarao BM, Maranas CD, Levine N, Willgert K, Conlan AJK, Olsen RJ, Davis JJ, Musser JM, Hudson PJ, Kapur V. 2022. Multiple spillovers from humans and onward transmission of SARS-CoV-2 in white-tailed deer. *Proc Natl Acad Sci*. doi:10.1073/pnas.2121644119

Kuhn M, Wickham H. 2020. Tidymodels: a collection of packages for modeling and machine learning using tidyverse principles. (manual).

Kumar A, Pandey SN, Pareek V, Narayan RK, Faiq MA, Kumari C. 2021. Predicting susceptibility for SARS-CoV-2 infection in domestic and wildlife animals using ACE2 protein sequence homology. *Zoo Biol* **40**:79–85. doi:10.1002/zoo.21576

Kumar S, Stecher G, Suleski M, Hedges SB. 2017. TimeTree: A Resource for Timelines, Timetrees, and Divergence Times. *Mol Biol Evol* **34**:1812–1819. doi:10.1093/molbev/msx116

Lam SD, Bordin N, Waman VP, Scholes HM, Ashford P, Sen N, van Dorp L, Rauer C, Dawson NL, Pang CSM, Abbasian M, Sillitoe I, Edwards SJL, Fraternali F, Lees JG, Santini JM, Orengo CA. 2020. SARS-CoV-2 spike protein predicted to form complexes with host receptor protein orthologues from a broad range of mammals. *Sci Rep* **10**:16471. doi:10.1038/s41598-020-71936-5

Liu Y, Hu G, Wang Y, Zhao X, Ji F, Ren W, Gong M, Ju X, Zhu Y, Cai X, Wu J, Lan X, Xie Y, Wang X, Yuan Z, Zhang R, Ding Q. 2021. Functional and Genetic Analysis of Viral Receptor ACE2 Orthologs Reveals a Broad Potential Host Range of SARS-CoV-2. *bioRxiv* 2020.04.22.046565. doi:10.1101/2020.04.22.046565

Luan J, Lu Y, Jin X, Zhang L. 2020. Spike protein recognition of mammalian ACE2 predicts the host range and an optimized ACE2 for SARS-CoV-2 infection. *Biochem Biophys Res Commun* **526**:165–169. doi:10.1016/j.bbrc.2020.03.047

Lundberg SM, Lee S-I. 2017. A Unified Approach to Interpreting Model Predictions In: Guyon I, Luxburg UV, Bengio S, Wallach H, Fergus R, Vishwanathan S, Garnett R, editors. Proceedings of the 31st International Conference on Neural Information Processing Systems. Presented at the 31st Conference on Neural Information Processing Systems. Long Beach, CA, USA. pp. 4765–4774.

MacLean OA, Lytras S, Weaver S, Singer JB, Boni MF, Lemey P, Pond SLK, Robertson DL. 2021. Natural selection in the evolution of SARS-CoV-2 in bats created a generalist virus and highly capable human pathogen. *PLOS Biol* **19**:e3001115. doi:10.1371/journal.pbio.3001115

McAloose D, Laverack M, Wang L, Killian ML, Caserta LC, Yuan F, Mitchell PK, Queen K, Mauldin MR, Cronk BD, Bartlett SL, Sykes JM, Zec S, Stokol T, Ingerman K, Delaney MA, Fredrickson R, Ivančić M, Jenkins-Moore M, Mozingo K, Franzen K, Bergeson NH, Goodman L, Wang H, Fang Y, Olmstead C, McCann C, Thomas P, Goodrich E, Elvinger F, Smith DC, Tong S, Slavinski S, Calle PP, Terio K, Torchetti MK, Diel DG. 2020. From People to Panthera: Natural SARS-CoV-2 Infection in Tigers and Lions at the Bronx Zoo. *mBio* **11**. doi:10.1128/mBio.02220-20

Melin AD, Janiak MC, Marrone F, Arora PS, Higham JP. 2020. Comparative ACE2 variation and primate COVID-19 risk. *Commun Biol* **3**:1–9. doi:10.1038/s42003-020-01370-w

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, Lanfear R. 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. *Mol Biol Evol* **37**:1530–1534. doi:10.1093/molbev/msaa015

Munster VJ, Feldmann F, Williamson BN, van Doremalen N, Pérez-Pérez L, Schulz J, Meade-White K, Okumura A, Callison J, Brumbaugh B, Avanzato VA, Rosenke R, Hanley PW, Saturday G, Scott D, Fischer ER, de Wit E. 2020. Respiratory disease in rhesus macaques inoculated with SARS-CoV-2. *Nature* **585**:268–272. doi:10.1038/s41586-020-2324-7

Mykytyn AZ, Lamers MM, Okba NMA, Breugem TI, Schipper D, Doel PB van den, Run P van, Amerongen G van, Waal L de, Koopmans MPG, Stittelaar KJ, Brand JMA van den, Haagmans BL. 2021. Susceptibility of rabbits to SARS-CoV-2. *Emerg Microbes Infect* **10**:1–7. doi:10.1080/22221751.2020.1868951

Oreshkova N, Molenaar RJ, Vreman S, Harders F, Munnink BBO, Honing RWH der, Gerhards N, Tolsma P, Bouwstra R, Sikkema RS, Tacken MG, Rooij MM de, Weesendorp E, Engelsma MY, Bruschke CJ, Smit LA, Koopmans M, Poel WH van der, Stegeman A. 2020. SARS-CoV-2 infection in farmed minks, the Netherlands, April and May 2020. *Eurosurveillance* **25**:2001005. doi:10.2807/1560-7917.ES.2020.25.23.2001005

Oude Munnink BB, Sikkema RS, Nieuwenhuijse DF, Molenaar RJ, Munger E, Molenkamp R, van der Spek A, Tolsma P, Rietveld A, Brouwer M, Bouwmeester-Vincken N, Harders F, Hakze-van der Honing R, Wegdam-Blans MCA, Bouwstra RJ, GeurtsvanKessel C, van der Eijk AA, Velkers FC, Smit LAM, Stegeman A, van der Poel WHM, Koopmans MPG. 2021. Transmission of SARS-CoV-2 on mink farms between humans and mink and back to humans. *Science* **371**:172–177. doi:10.1126/science.abe5901

Pagel M. 1999. Inferring the historical patterns of biological evolution. *Nature* **401**:877–884.
doi:10.1038/44766

Paradis E, Claude J, Strimmer K. 2004. APE: Analyses of Phylogenetics and Evolution in R language.
Bioinformatics **20**:289–290. doi:10.1093/bioinformatics/btg412

Qiu Y, Zhao Y-B, Wang Q, Li J-Y, Zhou Z-J, Liao C-H, Ge X-Y. 2020. Predicting the angiotensin converting
enzyme 2 (ACE2) utilizing capability as the receptor of SARS-CoV-2. *Microbes Infect* **22**:221–225.
doi:10.1016/j.micinf.2020.03.003

R Core Team. 2021. R: A language and environment for statistical computing (manual). Vienna,
Austria.

Revell LJ. 2012. phytools: An R package for phylogenetic comparative biology (and other things).
Methods Ecol Evol **3**:217–223. doi:10.1111/j.2041-210X.2011.00169.x

Rodrigues JPGLM, Barrera-Vilarmau S, Teixeira JMC, Sorokina M, Seckel E, Kastritis PL, Levitt M. 2020.
Insights on cross-species transmission of SARS-CoV-2 from structural modeling. *PLOS Comput Biol*
16:e1008449. doi:10.1371/journal.pcbi.1008449

Schlottau K, Rissmann M, Graaf A, Schön J, Sehl J, Wylezich C, Höper D, Mettenleiter TC,
Balkema-Buschmann A, Harder T, Grund C, Hoffmann D, Breithaupt A, Beer M. 2020. SARS-CoV-2 in
fruit bats, ferrets, pigs, and chickens: an experimental transmission study. *Lancet Microbe*
1:e218–e225. doi:10.1016/S2666-5247(20)30089-6

Shi J, Wen Z, Zhong G, Yang H, Wang C, Huang B, Liu R, He X, Shuai L, Sun Z, Zhao Y, Liu P, Liang L, Cui
P, Wang J, Zhang X, Guan Y, Tan W, Wu G, Chen H, Bu Z. 2020. Susceptibility of ferrets, cats, dogs, and
other domesticated animals to SARS-coronavirus 2. *Science* **368**:1016–1020.
doi:10.1126/science.abb7015

Sit THC, Brackman CJ, Ip SM, Tam KWS, Law PYT, To EMW, Yu VYT, Sims LD, Tsang DNC, Chu DKW,
Perera RAPM, Poon LLM, Peiris M. 2020. Infection of dogs with SARS-CoV-2. *Nature* **586**:776–778.
doi:10.1038/s41586-020-2334-5

Soubrier J, Steel M, Lee MSY, Der Sarkissian C, Guindon S, Ho SYW, Cooper A. 2012. The influence of
rate heterogeneity among sites on the time dependence of molecular rates. *Mol Biol Evol*
29:3345–3358. doi:10.1093/molbev/mss140

V'kovski P, Kratzel A, Steiner S, Stalder H, Thiel V. 2021. Coronavirus biology and replication:
implications for SARS-CoV-2. *Nat Rev Microbiol* **19**:155–170. doi:10.1038/s41579-020-00468-6

Wilkinson SP, Davy SK. 2018. phylogram: an R package for phylogenetic analysis with nested lists. *J
Open Source Softw* **3**:790. doi:10.21105/joss.00790

Zhao Y, Wang J, Kuang D, Xu J, Yang M, Ma C, Zhao S, Li J, Long H, Ding K, Gao J, Liu J, Wang H, Li H,
Yang Y, Yu W, Yang J, Zheng Y, Wu D, Lu S, Liu H, Peng X. 2020. Susceptibility of tree shrew to
SARS-CoV-2 infection. *Sci Rep* **10**:16007. doi:10.1038/s41598-020-72563-w

Zimmerman JM, Eliezer N, Simha R. 1968. The characterization of amino acid sequences in proteins
by statistical methods. *J Theor Biol* **21**:170–201. doi:10.1016/0022-5193(68)90069-6