On the interplay between lipids and asymmetric dynamics of an NBS degenerate ABC transporter

Ágota Tóth, Angelika Janaszkiewicz, Veronica Crespi and Florent Di Meo*

INSERM U1248 Pharmacology & Transplantation, Univ. Limoges, 2 rue du Prof. Descottes, 87000 F-Limoges, France

Contact information

Dr. Florent Di Meo

INSERM U1248 Pharmacology & Transplantation, Univ. Limoges,

2 rue du Prof. Descottes,

87000 F-Limoges, France

Florent.di-meo@inserm.fr

Tel: +33(0)5 19 56 42 76

Abstract

Multidrug resistance-associated proteins are ABC C-family exporters. They are crucial in pharmacology as they transport various substrates across membranes. However, the role of the degenerate nucleotide-binding site (NBS) remains unclear likewise the interplay with the surrounding lipid environment. Here, we propose a dynamic and structural overview of MRP1 from *ca.* 110 μs molecular dynamics simulations. ATP binding to NBS1 is likely maintained along several transport cycles. Asymmetric NBD behaviour is ensured by lower signal transduction from NBD1 to the rest of the protein owing to the absence of ball-and-socket conformation between NBD1 and coupling helices. Even though surrounding lipids play an active role in the allosteric communication between the substrate-binding pocket and NBDs, our results suggest that lipid composition has a limited impact, mostly by affecting transport kinetics. We believe that our work can be extended to other degenerate NBS ABC proteins and provide hints for deciphering mechanistic differences among ABC transporters.

1. Introduction

ATP-binding cassette (ABC) transporters belong to one of the largest trans-kingdom protein superfamilies. The structural resolution of several ABC transporters has led to different conformations (namely, inward-facing – IF, outward-facing – OF, and occluded), which illustrate the alternating access as the most likely model to rationalize the substrate translocation along the transport cycle[1–3]. ABC transporter structures are made of at least two transmembrane domains (TMDs) consisting of six transmembrane helices (TMHs). TMDs are bound to two nucleotide-binding domains (NBDs) which are evolutionarily conserved over species. ABC transport cycle requires the binding of two ATP molecules and the energy released from the hydrolysis of at least one of them[3–7]. ATP molecules bind at the interface of the NBD dimer which adopts a non-covalent pseudo-symmetric head-to-tail arrangement; enabling the formation of two nucleotide-binding sites (NBSs). Both NBSs are formed with the conserved Walker A- and B-motifs, the A-, Q- and H-loops of one NBD and the ABC signature sequence and the X-loop of the other NBD[4,5,7,8].

Except for a few members (CFRT, ABCA4, ABCD4, SUR1/2)[2,3,9,10], eukaryotic ABC transporters are exporters, *i.e.*, they extrude substrates to the extracellular compartment. Eukaryotic ABC transporters used to be classified into type I (ABCB, ABCC, ABCD) and type II (ABCA, ABCG) families. Recently, the structural and functional diversities of ABC transporters have led to a new folding-based classification[2,3] in which the previous type I and type II exporters adopt the type IV and V folding, respectively.

Multidrug resistance-associated proteins (MRPs) are NBS degenerate ABC transporters[3–5,7,8]. In the non-canonical NBS1, the Walker-B catalytic glutamate, the A-loop tyrosine, and the first glycine residue of the ABC-signature motif are mutated into aspartate, tryptophan, and valine residues, respectively. These mutations were associated with significantly lower ATPase activity and higher ATP-binding affinity for the degenerate NBS1[4,7]. These observations have recently led to the development of a new asymmetric model for NBD function which may affect

the dynamics and function of the whole transporter[3,7]. *b*MRP1 function and kinetics were extensively and thoroughly investigated by combining structural information from cryo-EM experiments and single-molecule Förster Resonance Energy Transfer (smFRET)[5]. Despite the robust insights provided by the resolution of *b*MRP1 structure, unexplained differences between ABCB and ABCC exporters were observed, likely due to the non-native detergent-based environment used for *b*MRP1 experiments[7,11].

ABC transporters play a crucial role in pharmacology by transporting a tremendous variety of substrates, including xenobiotics and endogenous compounds, across cell membranes. For instance, their pharmacological role has been stressed by the International Consortium Transporter (ITC) which draws out a list of transporters of "emerging clinical importance" for which interactions with new xenobiotics have to be investigated in drug development[12]. Over the past decades, ABCC transporters, in which MRPs are included, have gained a growing interest owing to their role in pharmacology including patient inter-individual responses to treatments. For instance, investigations on ABCC2/MRP2 and ABCC4/MRP4 have been recommended by the ITC in order to retrospectively provide a mechanistic explanation of clinical observations regarding drug dispositions[13]. Given the role of MRPs in local pharmacokinetics and pharmacodynamics relationships (PK/PD) and therefore in local drug bioavailability[14], there is still a need to decipher *in situ* MRP transport cycle to provide a comprehensive overview of xenobiotic membrane crossing events. This is particularly relevant for MRPs located in proximal tubular kidney cells and liver hepatocytes since kidneys and liver are involved in the elimination of most worldwide -used xenobiotics[15].

Unfortunately, there is no experimentally resolved structure for human MRPs yet. An MD-refined protein threading MRP4 structure[16] has been proposed for which the computational resolution precludes functional investigations or thorough structural dynamics. However, several conformations of the bovine ABCC1/MRP1 (*b*MRP1) transporter have been resolved by cryogenic electron microscopy (cryo-EM)[4,5,8]. Given the high sequence similarity to the human ortholog *h*MRP1 (91%) as well as within other human MRPs (*ca.* 40-50%), the use of

*b*MRP1 structures appears relevant for investigating MRP transporter dynamics and functions[3]. MRP1 exporter adopts the type IV folding, however, it has an extra N-terminal TMD made of five TMHs. The so-called TMD_0 was shown not to play a role either in ABC ATPase activity or substrate transport[3,4,8]. Therefore, a TMD_0-less MRP1 model can be used as a prototype for ABCC exporters even for those which do not possess this domain (*e.g.*, MRP4). TMD_0 is connected to conventional ABC TMDs by a linker (L_0) which was shown to be mandatory for both trafficking and function[8,17]. The L_0 sequence is conserved in all members of the ABCC subfamily even in absence of TMD_0[8]. It is important to note that since the present model does not include TMD_0, the standard TMH labelling for ABC type IV will be used in the present manuscript, *i.e.*, TMH1 to TMH12.

The present work aims to map the ABC conformational space[1,18] of MRP1 considering different bound states (namely, apo, ATP and/or leukotriene C4 – LTX – bound states[8] for IF conformation and ATP bound state[4] for OF conformation) highlighting the importance of asymmetry in ABC domains. Furthermore, given the importance of surrounding lipids in the ABCC transport cycle[11,19], the interplay between the lipid bilayer membrane and protein dynamics was also investigated. This was achieved by using different computational symmetric membrane models made of (i) pure POPC (1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine), (ii) pure POPE (1-palmitoyl-2-oleoyl-sn-glycero-3-phosphoethanolamine), (iii) POPC:POPE (3:1), (iv) POPC:Chol (3:1) and (v) POPC:POPE:Chol (2:1:1); the last being the closest to mimic *in situ* MRP1 dynamics. All-atom unbiased microsecond-scaled molecular dynamics (MD) simulations were conducted to address the objectives.

2. Results

Projection of MRP1 onto the ABC conformational space

To examine the conformational space sampled during the simulations in POPC:POPE:Chol (2:1:1) accounting for bound states of *b*MRP1, different structural descriptors[1,11,18] were

considered. Namely, intracellular (IC) and extracellular (EC) angles were monitored for TMDs while NBD distance and NBD rocking-twist angle were used for NBDs (Fig. 1 and Supplementary Fig. 1-6). These structural descriptors were also measured on a large data set of experimentally resolved ABC proteins including *b*MRP1 cryo-EM structures[4,5,8] in order to reconstruct the ABC conformational space[1,18] (Fig. 1d and Supplementary Table 1). The known ABC conformations were efficiently featured, *i.e.*, IF open and occluded, OF as well as the recently resolved asymmetric unlock-returned (UR) turnover[1] conformations. *b*MRP1 MD simulations revealed the spontaneous closing of the intracellular cavity for all IF systems regardless of the bound state and the membrane composition (Fig. 1a-b, Supplementary Fig. 1-6 and Supplementary Tables 2-5). Average NBD distances and IC angles were significantly smaller than those calculated for *b*MRP1 cryo-EM structures. For instance, IF conformations converged toward similar IC angles and NBD distance values ranging from 26.0 ± 0.5 to 35.4 ± 1.8 and from 30.3 ± 1.9 to 55.0 ± 2.8 Å, respectively (Fig. 1b, Supplementary Fig. 1, 3-4, and Supplementary Tables 3-4). Interestingly, the spontaneous dimerization of NBDs was also observed in absence of ATP molecules. The difference between simulations and cryo-EM structure might be explained by the use of non-native detergent in experiments[7,11].

Figure 1. Overview of *b*MRP1 milestone structures along the transport cycle in POPC:POPE:Chol (2:1:1). *Inward-facing conformations:* apo-state (IF apo *b*MRP1), substrate-bound (*b*MRP1-LTX), ATP-bound (*b*MRP1-(ATP)₂), ATP/substrate-bound (*b*MRP1-LTX-(ATP)₂); *Outward-facing conformation:* ATP-bound (OF *b*MRP1-(ATP)₂). **a)** Representative snapshots of the different *b*MRP1 systems

investigated here at the beginning (left) and the end of MD simulations. **b)** Time-evolution of the IC and EC angles respectively for IF and OF structures. IC and EC angles were calculated according to the proposed ABC structural parameters defined in Methods section[1,11,18]. **c)** Time-evolution of key NBS distances defined by inter-NBD distances between Walker A Glycine and ABC signature serine residues[8]. **d)** Projection of bMRP1 structural parameters, each replica was averaged over the last 800 ns, onto the ABC conformational space obtained from multiple resolved ABC structures. PDB IDs of the resolved bMRP1 cryo-EM structures are explicitly mentioned. The first and second TMDs are respectively depicted in orange and blue, and NBD1 and NBD2 are respectively coloured yellow and cyan.

Indeed, resolved cryo-EM structures of bMRP1 exhibited larger IC angles and NBD distances than other ABC structures resolved in native environments (Fig. 1d). On the extracellular side of the lipid bilayer membrane, MD simulations of OF bMRP1-(ATP)$_2$ displayed minor openings of the EC gate as compared to bMRP1 OF cryo-EM structures, suggesting the existence of a slightly more open state. However, calculated EC angles remained small (lower than 20° in POPC:POPE:Chol (2:1:1), see Fig. 1a and Supplementary Fig. 1-2 and Supplementary Table 2), which precludes the substrate re-entry[1].

Even though trajectories performed with the pre-translocation state (*i.e.*, bMRP1-LTX-(ATP)$_2$) tend to populate the conformational subspace of OF conformation (Fig. 1d), substrate translocation was not observed. NBD twist values calculated for bMRP1-LTX-(ATP)$_2$ state are similar to OF ABC conformations. However, NBD distances remain larger than for resolved substrate-free OF ABC structures and OF bMRP1-(ATP)$_2$ simulations. Distances between Cα atoms of the ABC signature motif serine and a Walker A glycine were monitored (*i.e.*, Gly681-Ser1430 and Ser769-Gly1329, respectively denoted as d_{GS}^{NBS1} and d_{GS}^{NBS2}, Fig. 1c, Supplementary Fig. 7-8). Structural differences between the pre- and the post-translocation states (*i.e.*, bMRP1-LTX-(ATP)$_2$ and OF bMRP1-(ATP)$_2$, respectively) suggest that a conformational transition of NBD dimer is required prior to the substrate translocation event. Such transition from the so-called "non-competent" to "competent" NBD dimer conformations is likely to trigger TMD conformational transitions suggesting that it might be the limiting step for IF-to-OF transition.

Asymmetric dynamics of *b*MRP1 and modulation of its conformational landscape

Overall flexibilities were assessed by calculating root-mean-square fluctuations (RMSF, Supplementary Fig. 9). Per-residue RMSF confirms the asymmetric behaviour of NBDs[7], NBD2 being more flexible than NBD1. For each system, backbone-based principal component analyses (PCA) were conducted. Only the three first largest principal components were considered, revealing 85.6 % to 95.9 % of the overall structural variabilities depending on the system (Supplementary Fig. 10). For all IF simulations, the three first largest variabilities were systematically assigned to asymmetric NBD motions for which NBD2 contributed the most, from 27.0 to 59.6% of the motion (Supplementary Fig. 11). The first principal components were mostly assigned to NBD twist and rocking motions (Supplementary Movies 1-5). This is in agreement with the experimentally suggested higher flexibility of NBDs from smFRET experiments along the kinetic cycle of MRP1[5]. Regarding OF simulations, the two first principal components were associated with concerted NBD twist motion and opening of the extracellular side mediated mostly by TMH4, TMH5, TMH7 and TMH8. These TMHs have been suggested to behave as a single bundle in ABCB1/P-gp[18]. Furthermore, to a lesser extent than for IF conformations, NBD2 remained more involved in this shared motion than NBD1.

The asymmetric behaviour was also pictured by the supramolecular arrangement between NBD1 and NBD2 for which two main subpopulations were observed for apo, *b*MRP1-(ATP)$_2$ and bMRP1-LTX states (Fig. 2a). Interestingly, interactions within NBS1 across NBDs were maintained along our MD simulations, even in absence of ATP molecules. In contrast, NBS2 is more flexible leading to either open or closed NBD dimer arrangements. Surprisingly, both arrangements were also observed to a lesser extent in *b*MRP1-(ATP)$_2$ while ATP molecules were expected to maintain interactions at the NBD dimer interface. However, in presence of both substrate and ATP molecules, only the closed population was observed picturing the information transduction from the TMD substrate-binding pocket to NBDs in order to likely decrease the energy barrier for IF-to-OF transition in type IV folding ABC transporters[20,21].

In order to explain the asymmetric behaviour of NBDs, particular attention was paid to coupling helices (CH) which ensure the signal transduction from TMDs to NBDs[20,21]. Natively, ABC transporters exhibit four coupling helices linking intracellular domains of TMH2/TMH3, TMH4/TMH5, TMH8/TMH9, and TMH10/TMH11. The so-called CH_{2-3} and CH_{10-11} are in contact with NBD1 while CH_{4-5} and CH_{8-9} interact with NBD2 (Fig. 2a). Contacts between CHs and NBDs are not modified upon binding of ATP molecules or substrate (Supplementary Fig. 12). Interestingly, for a given NBD, one can consider a so-called "weak CH" (namely, CH_{2-3} and CH_{8-9} for NBD1 and NBD2, respectively) for which only a few contacts were observed. On the other hand, the so-called "strong CH" (namely, CH_{10-11} and CH_{4-5} for NBD1 and NBD2, respectively) exhibited more contact with the NBD and for which the "ball-and-socket" arrangement is significantly more pronounced than for the "weak CH". Interestingly, in mammalian MRP1, the NBD1 sequence exhibits a 13-aminoacid deletion which is associated with fewer contacts and significantly weaker interactions between CH_{2-3} and CH_{10-11} as compared to CH_{4-5} and CH_{8-9} for NBD2 (Supplementary Fig. 13). Furthermore, the few contacts observed between CH_{2-3} and CH_{10-11} are slightly decreased in presence of ATP molecules while the contact pattern for NBD2, *i.e.*, CH_{4-5} and CH_{8-9}, is well conserved regardless of the presence of ATP and/or leukotriene C4. This leads to the disruption of the so-called "ball-and-socket" arrangements of CH_{2-3} and CH_{10-11} in NBD1 which is expected (i) to be responsible for the aforementioned larger flexibility of NBS1 region and (ii) to preclude the information transduction between TMDs and NBD1[8].

IF conformations exhibited weaker NBD dimer interactions than OF conformation, explaining the overall larger flexibility. Likewise, MD simulations revealed slightly higher flexibility of IF apo state as compared to ATP- and/or LTX-bound systems. This is in agreement with previous observations suggesting that interactions between TMHs or between NBDs are modulated by the presence of substrate and/or ATP molecules[4,7,8,22]. Taking advantage of our extensive unbiased MD simulations, explored conformational subspaces were featured in terms of free energy using the InfleCS clustering methods[23] (Fig. 2b). Given the aforementioned non-

competent NBD dimer conformations, the focus was paid to NBD structural parameters, namely NBD twist and NBD distance. Larger variabilities for IF conformations were observed leading to multiple plausible minima for which interconversion is possible but slow. The expected to be competent NBD twist *versus* NBD distance subspace tends to be populated in presence of ATP molecule and/or substrate. Such finding highlights the central role of ATP- and substrate-binding events in the ABCC1 transport cycle.

To better decipher the NBD dimer dynamics, structural networks[24] were obtained by considering both contact maps (Supplementary Fig. 14) and dynamic cross-correlation matrices (Supplementary Fig. 15). Each NBD can be globally split into two subdomains regardless of the conformation or the bound states driven by ATP-binding (Supplementary Fig. 16). Indeed, it is worth mentioning that similar so-called communities were relatively well-conserved over the replicas and the systems investigated (Supplementary Table 6). The first community is defined by the Walker A and B as well as the A- and H-loops. On the other hand, the second community includes Q-and X-loops and the ABC signature sequence. Residues involved in these communities are highly correlated suggesting that local residue displacement upon ATP-binding will propagate. The two communities may be considered almost independent thus the binding of one ATP is not expected to strongly impact the motions of the second community. Meanwhile, given the asymmetric NBD dimer arrangement, ATP and magnesium are expected to link communities across NBDs. Interestingly, dynamic correlations in IF apo simulations exhibited larger variabilities since communities could be split into subcommunities (Supplementary Table 7), so that ATP-binding is expected to strengthen the structural cooperation within and between NBDs.

Figure 2. Asymmetric structural dynamics of _b_MRP1 systems in POPC:POPE:Chol (2:1:1) and their conformational landscape. a) Representative snapshots picturing the open and closed conformations of NBD dimers observed during MD simulations. IF apo _b_MRP1, _b_MRP1-(ATP)$_2$ and _b_MRP1-LTX revealed two main subpopulations for which black arrows highlight the motion; while the pre- and post-translocation conformations (namely, _b_MRP1-LTX-(ATP)$_2$ and OF _b_MRP1-(ATP)$_2$) exhibited only the closed NBD dimer conformation. NBD1, NBD2 and coupling helices are respectively depicted in cyan, yellow and pink. **b)** System-dependent local conformational landscapes obtained from the GMM-based approach developed in the InfleCS method[23,25] highlighting the influence of nucleotides and the substrate on _b_MRP1 structural dynamics. **c)** Averaged Coulomb (up) and van der Waals (down) potentials calculated between nucleotides and NBS1 and NBS2, separately. **d)** Calculated H-bond networks between ATP and NBS1 (left) or NBS2 (centre) and between LTX and substrate-binding pocket residues (right).

Conformation-dependent binding modes of ATP in degenerate and canonical NBSs

Particular attention was paid to the binding modes of ATPs by assessing van der Waals and Coulomb potentials as well as H-bond networks in the NBSs (Fig. 2c-d). It is important to note that such analyses should not be considered quantitatively owing to the compensation of errors between the two potentials, especially at short distances[26]. However, they can be used to provide qualitative hints in order to compare ATP-binding driving forces. Interestingly, both Coulomb and dispersive interactions exhibited less attractive energies for all IF conformations than for OF. Given that NBDs exhibit lower flexibility in OF conformation, the subsequent tighter NBD dimer interactions can be rationalized by the proper local arrangement of ATP molecules in NBSs. For instance, calculated lower π-stacking distances between ATP molecules and NBS conserved motifs were systematically larger in IF than in OF conformations leading to lower interaction energies between ATP molecules and MRP1 residues (Fig. 2c and Supplementary Table 8). Surprisingly, in IF conformations, NBS1 tends to exhibit lower dispersive interactions between ATP molecule and Trp653 A-loop residue as compared to Tyr1301 in NBS2 A-loop, while the spatial aromatic surface of tryptophane is larger than for tyrosine. On the other hand, in the OF conformation, dispersive contributions tend to be slightly larger for NBS1 than for NBS2. H-bond networks between ATP molecules and NBSs (Fig. 2d) suggest a similar network

between NBSs for IF conformations. The H-bond network is well conserved over IF and OF conformations regarding interactions with Walker A, however, ATP-bound IF systems do not exhibit the expected H-bond network with the signature motif as observed in OF simulations. Interestingly, MD simulations suggest an asymmetric behaviour between NBS regarding H-bonds with Q-loop glutamine residues, namely Gln713 and Gln1374, respectively for NBS1 and NBS2. Calculated distances suggested that ATP γ-phosphate binding to NBS1 Q-loop was weaker than for NBS2 in the OF conformation. The opposite behaviour was observed in MD simulations for IF conformations. Therefore, present simulations underline that proper ATP-binding modes in both NBSs are key in triggering conformational changes required for substrate translocation.

Towards deciphering the allosteric modulation between substrate- and nucleotide-binding sites

Particular attention was paid to the substrate-binding pocket and compared to the cryo-EM structure of *b*MRP1 bound to leukotriene C4[8]. In agreement with experiments, leukotriene C4 binding mode took place in the two so-called P- and H-pockets ("P" and "H" respectively standing for polar and hydrophobic). Given the amphiphilic feature of leukotriene C4, Coulomb and H-bond networks have been shown to be central for substrate-binding[8]. MD simulations highlighted the same key residues which were experimentally observed (Fig. 2d). For instance, strong salt-bridges were observed between arginine residues (Arg1248, Arg1196 and Arg593, Fig. 2d and Fig. 3a) maintaining at least two out of the three leukotriene carboxylate groups in the P-pockets. Interestingly, variabilities in terms of H-bond fractions or interaction energies (Fig. 2d and Supplementary Fig. 17) suggest a dynamic binding mode in agreement with its expected breaking along the IF-to-OF large-scale conformational changes.

Figure 3. Substrate-bMRP1 interactions and subsequent allosteric communications to nucleotide-binding sites. a) Substrate-binding pocket highlights important residues to leukotriene C4 binding. The structure of leukotriene is also shown for which amphiphilic features are stressed out. **b)** Definition of the allosteric pathway investigated in the present study for which NBS1 and NBS2 were treated separately. **c)** Calculated allosteric efficiencies of the information flow between substrate-binding

pocket and NBS1 (red) or NBS2 (blue) for the different systems embedded in POPC:POPE:Chol (2:1:1). Solid, dashed and dotted lines respectively depict efficiencies considering: Protein + lipids + nucleotides/substrate, Protein + nucleotides/substrate and standalone Protein. Protein **d)** and Lipid **e)** contributions to the information flow for allosteric communication from substrate-binding pocket to NBS1 (top) and NBS2 (bottom) show that NBD2 and its coupling helices (CH_{4-5} and CH_{8-9}) are systematically involved regardless of the sink region.

Even though differences in terms of interaction energies and H-bond networks between bMRP1-LTX-$(ATP)_2$ and bMRP1-LTX or bMRP1-$(ATP)_2$ remained low, they suggest a distant effect between NBSs and the substrate-binding pocket. Allosteric effect was assessed between the substrate-binding pocket and NBS1 and NBS2, independently (Fig. 3b). This was achieved by considering key residues for each binding site (Supplementary Table 9) for allosteric pathway network analyses[27,28]. Efficiencies (Fig. 3c) were calculated in presence or absence of substrate and ATP molecules. The impact of POPC:POPE:Chol (2:1:1) lipid bilayer was also considered. Natively, substrate-binding pocket and NBSs are allosterically connected through the protein as shown by calculated efficiencies without including nucleotides or substrates. As expected, the presence of substrate and/or ATP molecules substantially increased the allosteric communication from substrate-binding pocket to both NBSs. Interestingly, in spite of the aforementioned NBD asymmetric dynamics, present calculations did not exhibit a significant difference in terms of efficiencies between NBSs. Betweennesses were calculated to picture the residue and domain contributions to the allosteric pathway (Fig. 3d and Supplementary Fig. 18-20). Particular attention was paid to ATP-bound systems, *i.e.*, IF bMRP1-$(ATP)_2$, and bMRP1-LTX-$(ATP)_2$ as well as OF bMRP1-$(ATP)_2$ (Fig. 4d), other systems are reported in Supplementary Fig. 21. The main residues involved in the binding pocket-NBS allosteric pathways are mostly located in the intracellular part of TMHs. Interestingly, an asymmetric behaviour, for which TMH4, TMH5, TMH7 and TMH8 are significantly more involved than TMH1, TMH2, TMH10 and TMH11, was again observed. bMRP1 exhibits an asymmetric feature regarding the so-called "ball-and-socket" arrangements which was shown to be responsible for the structural cooperation between TMHs and NBDs

through coupling helices[8]. The thirteen amino acid deletion in NBD1 leads to lower cooperation between TMH1, TMH2, TMH10 and TMH11 with NBD1, which in turn decreases the allosteric communication between substrate-binding pocket and NBS1. Furthermore, it is worth mentioning that our calculations suggest that information mostly goes through NBD2 since the direct communication with NBD1 is significantly weakened by the absence of "ball-and-socket" conformation for CH_{10-11}[8]. Interestingly, POPC:POPE:Chol lipid bilayer was also shown to play a key role in the allosteric communication between substrate-binding pocket and NBSs (Fig. 3c). However, even though its impact is significant, lipid bilayer contributions appeared milder than in *e.g.*, Major Facilitator Superfamily membrane transporters[29].

On the interplay between the lipid bilayer and *b*MRP1 structures and dynamics

As nowadays more attention has been paid to the interplay between the surrounding lipid environment[11,19,22,27,29] and membrane proteins, lipid-dependent protein dynamics and lipid-protein interactions were investigated. This was achieved by carrying out MD simulations in different lipid bilayers, namely POPC, POPC:Chol (3:1), POPC:POPE:Chol (2:1:1). IF apo *b*MRP1 and OF *b*MRP1-$(ATP)_2$ systems were also considered in unrealistic POPE and POPC:POPE (3:1) lipid bilayers.

Projection of lipid-dependent structural parameters onto the ABC conformational space (Supplementary Fig. 1 and 22-23) revealed that most of the present simulations tend to explore similar ABC subspaces regardless of the lipid bilayer membrane composition. Figure 4a compares structural parameter averages according to lipid bilayer compositions for all MD simulations performed in the present study. For IF conformations, only intracellular structural parameters (*i.e.*, IC angle and NBD distance) exhibited slight deviations according to the lipid composition. Systems performed in pure POPC lipid bilayer exhibited slightly more open conformations. On the other hand, our calculations suggest that only EC angle is affected by lipid bilayer composition but only for OF conformations. Likewise, it is important to note that

calculated cavity radii (Supplementary Fig. 24) exhibited tiny differences while comparing lipid bilayer compositions. Even if these calculations underline a relatively limited overall impact of membrane composition on the bMRP1 structure, they did not sufficiently picture the dynamic variability over MD simulations and replicas (Supplementary Fig. 1-6 and 24).

Therefore, lipid-dependent conformational landscapes were calculated (Supplementary Fig. 25-26). For a given conformation and bound state, MD simulations preferentially populated similar regions regardless of lipid composition. However, structural variability was shown to be significantly affected by lipid composition. For instance, in pure POPC lipid bilayer, more open IF conformation subspaces, ranging from 30 to 55 Å NBD distance, were sampled. This effect was however reduced in presence of substrate which is expected to tend to maintain more contacts between TMHs and thus reduce the intracellular opening. MD simulations performed on POPC:POPE:Chol (2:1:1) exhibited a significantly smaller sampled region suggesting that the presence of PE lipids tends to close the intracellular gate of bMRP1 IF conformations, regardless of the bound state. On the other hand, regarding OF conformations and extracellular opening, global minima were interestingly observed in the same subspace of the conformational space as MD simulations performed in pure POPC and POPC:Chol (3:1). It suggests a limited impact of cholesterol on the opening of bMRP1 EC gate. However, the presence of PE lipids slightly shifted the calculated minima toward more opened OF structures, from 13.9 to 18.5°. Tilt angles between TMHs and lipid bilayer normal were measured in order to unravel slight differences between the distinct lipid bilayer compositions (Supplementary Fig. 27). Even though no clear conclusions can be drawn from these analyses, orientations of TMH3, TMH6 and TMH9 appeared more sensitive in absence of cholesterol. While considering that TMHs act as bundles as shown for ABCB1/P-gp[18], the impact of lipid bilayer membrane was more pronounced on the tilt orientations of given smaller bundles (Supplementary Fig. 28), namely Bundle C & D respectively consisting in TMH3/TMH6 and TMH9/TMH12. Interestingly, these bundles are expected to undergo larger conformational changes along the transport cycle[18]. This suggests that even though lipid composition seems to have a rather

limited impact on the overall structure of *b*MRP1, it may play a role in the kinetics of substrate transport by *b*MRP1.

Figure 4. Interplay between *b*MRP1 structural dynamics and lipid bilayer according to its composition. a) Average IC angle, EC angle and NBD distance for all *b*MRP1 systems embedded in

different lipid bilayers. **b)** Calculated C-atom lipid tail order parameters (S_{CD}) for palmitic (sn_1, left) and oleic tails (sn_2, right) for all systems embedded in POPC-based models. Solid, dashed and dotted lines respectively depict lipid order parameters obtained in POPC:POPE:Chol (2:1:1), POPC:Chol (3:1) and POPC lipid bilayer models. **c)** Calculated binding hotspots obtained from cholesterol and PE lipids defined by presence likelihood higher than 80% and 50% respectively for cholesterol and PE lipids. Zoom on Cryo-EM resolved cholesterol (purple) is shown to highlight the specific parallel orientation with respect to the lipid bilayer. **d)** Important lipid areas based on allosteric pathway analysis. The location of key cholesterol molecules involved in the allosteric communication between substrate-binding pocket and NBSs is shown. Both NBS is considered, highlighting the expected central roles of L0, pre-TMH1 and pre-TMH7 regions in specific lipid-protein interactions which might favour *b*MRP1 function.

To better understand the interplay between the lipid bilayer and *b*MRP1, particular attention was paid to the lipid bilayer membrane structure. The larger structural variability in pure POPC membrane was explained by a significantly more fluid lipid bilayer structure pictured by lower order parameters (S_{CD}) for palmitate and oleate tails (Fig. 4b). In line with the biophysics of pure lipid bilayers, the presence of cholesterol modulates the fluidity of POPC by increasing the lipid order which in turn led to lower flexibility of the lipid bilayer membrane[30]. To a lesser extent, the presence of PE lipid potentiated the structural effect of cholesterol, which is in agreement with the slightly lower structural variability of *b*MRP1 in POPC:POPE:Chol (2:1:1) as compared to other lipid bilayer membranes. Lipid order calculations also suggest a weak impact of protein dynamics on lipid bilayer structures. Indeed, IF apo and OF *b*MRP1-(ATP)$_2$ MD simulations exhibited slightly less ordered lipid tail profiles than IF *b*MRP1-LTX, *b*MRP1-(ATP)$_2$ and *b*MRP1-LTX-(ATP)$_2$. This may be explained by larger variability of intracellular and extracellular openings for IF and OF conformation, respectively, which in turn is likely to lead to more pronounced displacements of surrounding lipids.

The distribution of surrounding lipids revealed important cholesterol and PE lipid hotspots. For instance, PE lipids were shown to preferentially bind to pre-TMH7 elbow helix as well as close to the L0 domain, for more than 50% of the simulation (Fig. 4c). It is worth mentioning that electron density maps revealed three cholesterol molecules bound to the resolved OF *b*MRP1 structure. Interestingly, one by the pre-TMH7 was maintained near to its initial position with a

probability higher than 80% (Fig. 4c) for which cholesterol is oriented in line with pre-TMH7 elbow helix, *i.e.*, parallel to the lipid bilayer. Pre-TMH7 cholesterol hot spots were also observed for example in IF-apo POPC:POPE:Chol (2:1:1) simulations (Fig. 4c). Moreover, pseudo-symmetrically, a cholesterol hotspot was also observed near to pre-TMH1 elbow helix in simulations carried out with IF *b*MRP1 conformations (Fig. 4c). Interestingly, allosteric pathway analyses underlined the key role of cholesterol molecules close to pre-TMH1 and -TMH7 elbow helices in the information transduction from the substrate-binding pocket to NBSs as shown in Fig. 4d.

3. Discussion

*b*MRP1 is the only member of the ABC drug exporter C-family which has been resolved by cryo-EM so far, given that ABCC7/CFTR is a chloride channel[31] and sulfonylurea receptors (ABCC8 and ABCC9) are involved in the regulation of potassium channels[32]. Over the past decade, particular attention has been paid to MRP transporters including MRP1, MRP2 and MRP4 given their clinically and pharmacologically relevant roles in drug disposition as pointed out by the ITC[12,13]. In contrast to its cousin ABCB1/P-gp, knowledge about MRP1 dynamics and functions still remains fragmented although it has been resolved in multiple states, such as IF apo and substrate-bound states[8] and two OF states under pre-[4] and post-hydrolysis[5] conformations, respectively bound to either two ATP molecules or to ADP/ATP pair. In the present work, an extensive set of all-atom MD simulations were performed in order to capture conformational dynamics of *b*MRP1 in different states considering different mixtures of lipid bilayer models including PC and PE lipids as well as cholesterol. We propose an MD-based computational approach in order to (i) complete the experimental observations made in detergents and (ii) highlight structural patterns which might be extended to, at least, other ABC C-family transporters.

Global conformational dynamics in POPC:POPE:Chol (2:1:1) shows significant variations of IF structures with respect to cryo-EM structures. In IF states, spontaneous closing of NBDs was

systematically observed regardless of the presence of ATP molecules as pictured by IC angle and NBD distance values which all converged toward the same subspace. Regarding ABC conformational space, the presence of either ATP molecules in both NBS or substrate in the TMD binding pocket mostly shifts the dynamics of NBD dimerization by modulating the NBD twist value (Fig. 2b and Supplementary Fig. 5). Therefore, our MD simulations are in perfect agreement with recent observations[6,7] that wide-open IF structures are expected to be unlikely in membrane environments. Wide-open structures observed in cryo-EM experiments are thus believed to be due to artifacts owing to the use of non-physiological environments for structure resolution[7], in agreement with structural differences observed *e.g.*, for P-gp reconstituted either in detergents or in nanodiscs[11]. MRP1 and likely other MRPs are expected to adopt the same "ATP-switch" model feature as suggested for TM287/288[6]. ATP-bound to NBS1 is expected to stay over multiple transport cycles sterically acting as a pivot point in absence of substrate. This is *e.g.*, pictured by the existence of two subpopulations for IF apo, ATP- and LTX-bound states regarding NBD dimer (Fig. 2a), namely open and close for which NBS1 is systematically formed. The NBD1 13 amino-acid deletion is thus expected to play a central role in this pivot model. Indeed, the absence of the "ball-and-socket" structure for NBD1 might weaken the coupling between NBD1 and TMH10/TMH11. In absence of substrate, this is associated with more flexibility regarding NBD1 leading to the opening of NBD dimer. However, in presence of ATP molecules and substrate, only close NBD dimer conformations were observed (Fig. 2a) highlighting the allosteric communication between TMDs and both NBSs. Owing to the lower coupling between TMH10/TMH11 and NBD1, allostery between TMDs and NBDs is expected to be mostly mediated through NBD2. Interestingly, NBS1 was also formed even in absence of ATP molecules (Fig. 2a). For instance, this might also explain why IF-to-OF transitions were experimentally observed even in absence of ATP molecules[5]. Our results highlight the importance of NBD and NBS asymmetry as an evolution of ABC transporters which may result from energy saving while keeping transport function by requiring a single ATP hydrolysis.

MRP1 is expected to carry mostly anionic amphiphilic substrates contrary to P-gp which mostly transports hydrophobic substrates[2]. In agreement with structural observations, substrate access directly from either high- or low-density lipid tail regions of the membrane is very unlikely owing to the absence of an access channel in lipid bilayer contrary to P-gp[8]. However, amphiphilic substrates might partition in the high-density polar head region. Therefore, MRP1 substrate access is expected to occur either directly from the cytoplasm or from the high-density polar head region, for which access might be possible via TMH4 and TMH6 (Supplementary Fig. 29). On the other side of *b*MRP1, our MD simulations suggest that substrate access between TMH10 and TMH12 may be less likely for bulky substrates. However, such assumptions require further biochemical and structural investigations.

In the present work, we also investigated the interplay between lipid bilayer membranes and protein dynamics. Our results first underline the particular importance of the use of a reliable membrane environment to provide robust insights into transporter dynamics and functions. By playing with different lipid bilayer models such as cholesterol-free and PE-free membranes, *b*MRP1 dynamics is modulated. In line with previous biophysical studies, the presence of cholesterol in PC-based membrane increased membrane stiffness which in turn reduced conformational dynamics of MRP1[30]. The impact of PE lipids is expected to be rather limited for IF states. However, interestingly, EC opening under OF conformation appeared more favourable in presence of PE (Fig. 4a). Globally, our results also suggest that lipid bilayer composition is very unlikely to preclude overall transporter functions, but it is expected to mostly affect kinetics. This hypothesis should be considered carefully for other MRP transporters. Indeed, in contrast to *e.g.,* MRP2 and MRP4, MRP1 is a ubiquitous exporter which was observed in different cell types for which lipid bilayer composition might differ. We can thus infer that other MRP members might become more sensitive to lipid bilayer membrane compositions.

Out of the physical role of lipid bilayer on protein dynamics, our results support that lipid components also play an active role in transporter function. In line with computational

observations made on other membrane proteins[11,27,33,34], lipid components are involved in the allostery from TMDs to NBDs. More importantly, cholesterol bind to pre-TMH1 and -TMH7 elbow helices are strongly involved in these allosteric pathways. Such findings pave the way regarding the role of lasso pre-TMD motif (L_0) which was shown to be required for the MRP1 transport function[8,17]. Even if the role of lipids might sound limited on MRP1 as compared to other membrane receptors and channels, more attention should be paid to lipid-protein interactions, not only regarding the biophysical impact on the protein dynamics but also as transport modulator as recently proposed for P-gp[11].

We hope that the present work provides new insights into the function and the lipid-protein interplay of NBS degenerate ABC transporters for further investigations such as the role of MRPs in local pharmacokinetics, including *e.g.*, the impact of rare mutations/polymorphism as well as disease-based membrane lipid imbalance toward personalized medicine.

4. Methods

4.1. Construction of *b*MRP1 models embedded in lipid bilayer membranes

In order to have an overview of milestone structures along the transport cycle of *b*MRP1, different conformations and bound states were considered in the present study: IF apo *b*MRP1, IF *b*MRP1-(ATP)$_2$, IF *b*MRP1-LTX, IF *b*MRP1-LTX-(ATP)$_2$ and OF *b*MRP1-(ATP)$_2$. The cryo-EM structures were used as starting structures for IF (PDB ID: 5UJ9[8] and 5UJA[8]) and OF-conformations (PDB ID: 6BHU[4]). It is important to note that OF conformation was resolved using E1454Q mutant which was shown to lower the rate of ATP hydrolysis thus promoting OF structure determination[4]. This mutation was reverted manually for the present study. The so-called "TMD$_0$" was not included in the present models as it has already been shown not to affect the substrate transport[3,4,8]. However, it was shown that the so-called pre-TMH1 lasso domain (L$_0$) is mandatory for MRP1 function while it was not resolved in any cryo-EM MRP1 structures[8,17]. Missing parts of L$_0$ domain was modelled using either I-Tasser (Iterative Threading ASSEmbly Refinement) server[35] and modeller v9.23[36] for IF and OF conformations, respectively. Likewise, the missing loop between TMH6 and NBD1 was also added in the present models.

IF conformations were built in different bound states, namely apo, ATP$_2$-, LTX- and LTX-(ATP)$_2$ states while OF conformation was solely constructed in the ATP$_2$-bound state. IF *b*MRP1-(ATP)$_2$ and IF *b*MRP1-LTX-(ATP)$_2$ were constructed by superimposing separately NBD of 5UJ9 and 5UJA, respectively onto the NBDs of OF conformation in which both ATP molecules and Mg^{2+} are bound to NBSs. All final models were shortly minimized in vacuum using the Amber18 package[37,38] to avoid unphysical steric clashes.

CHARMM-GUI input generator[39,40] was used to embed the different *b*MRP1 models into different lipid bilayers, namely pure POPC, POPC:Chol (3:1) and POPC:POPE:Chol (2:1:1) taking advantage of *b*MRP1 coordinates obtained from the OPM (Orientations of Proteins in Membranes) database[41]. IF apo *b*MRP1 and OF *b*MRP1-(ATP)$_2$ were also embedded into pure POPE and POPC:POPE (3:1) lipid bilayers in order to specifically investigate lipid-protein interactions with PE lipids. It is worth mentioning that the resolved cryo-EM structure of OF *b*MRP1-(ATP)$_2$ also includes three resolved cholesterol molecules which were kept during the simulations in order to investigate their importance. From these different lipid bilayer compositions, POPC:POPE:Chol (2:1:1) mixture appeared the most relevant to model cell membranes. The other types of membranes were considered to help understand the role of each lipid in protein dynamics. The original total size of every system was *ca.* 120 x 120 x 180Å^3 (see Supplementary Tables 10-11 for system descriptions). To mimic physiological conditions, 0.15 M NaCl was used, and the systems were solvated using TIP3P explicit water molecules[42–44]. The final systems are made of *ca.* 245 000 atoms (see details in Supplementary Table 12).

4.2.   Molecular dynamics simulations

CHARMM-GUI[39,40] outputs were converted to Amber format using AmberTools scripts[37,38] (namely, charmmlipid2amber.py and pdb4amber). Regarding ATP and LTX-bound systems, substrate, nucleotides and Mg^{2+} ions were added after building protein-lipid systems; therefore, neutrality was ensured by randomly removing the corresponding number of counterions. Amber FF14SB[45], Lipid17[46] and the modified DNA.OL15[47,48] force fields were used to respectively model protein residues, lipids and ATP molecules. Water molecules, Mg^{2+} ions and counterions were modelled using the TIP3P water model[42–44] as well as the corresponding monovalent and divalent ion parameters from Joung and Cheatham[49,50]. LTX-substrate (Leukotriene C4) parameters were derived from the Generalized Amber Force Field version 2 (GAFF2)[51] using the Antechamber software[52]. LTX partial atomic charges were derived from quantum mechanical based calculations

at the HF/6-31G* level of theory, using the R.E.D. server[53]. Each system was simulated with periodic boundary conditions. The cutoff for non-bonded interactions was 12 Å for both Coulomb and van der Waals potentials. Long-range electrostatic interactions were computed using the particle mesh Ewald method[54].

Minimization and thermalization of the systems and MD simulations were carried out with Amber18 and Amber20 packages[37,38] using CPU and GPU PMEMD versions. Minimization was carried out in four steps by sequentially minimizing: (i) water O-atoms (20000 steps); (ii) all bonds involving H-atoms (20000 steps); (iii) water molecules and counterions (50000 steps) and (iv) the whole system (50000 steps). Each system was then thermalized in two steps: (i) water molecules were thermalized to 100 K during 50 ps under (N,V,T) ensemble conditions using a 0.5 fs time integration; (ii) the whole system was then thermalized from 100 K to 310 K during 500 ps under (N,P,T) ensemble conditions with 2 fs timestep in semi-isotropic conditions. Then, each system was equilibrated during 5 ns under (N,P,T) ensemble conditions with 2 fs timestep in semi-isotropic conditions, using Berendsen barostat. Production runs were then carried out at the microsecond scale with 2 fs integration timestep under (N,P,T) ensemble conditions with semi-isotropic scaling. Temperature was maintained using the Langevin dynamics thermostat[55] with 1.0 ps^{-1} collision frequency. Constant pressure set at 1 bar was maintained with semi-isotropic pressure scaling using either Berendsen barostat[56] for IF apo bMRP1 and OF bMRP1-(ATP)$_2$ or Monte Carlo barostat for IF bMRP1-(ATP)$_2$, IF bMRP1-LTX and IF bMRP1-LTX-(ATP)$_2$. The latter was used to speed up computational time.

In order to ensure the ATP docking into NBS, it is worth mentioning that restraint-MD simulations were carried out using a similar approach as proposed by Wen *et al.*[58]. Shortly, a set of distance-based restraints were applied between the A-loop tryptophan/tyrosine residues (Trp653 and Tyr1301, respectively for NBD1 and NBD2) and corresponding ATP purine moiety. Mg^{2+}-ATP-NBD arrangement was maintained by applying restraints between Mg^{2+} ions and ATP phosphate groups as well as between Mg^{2+} ions and Walker A Serine and Q-loop glutamine residue (namely, Ser685 and Gln713 for NBD1 and Ser1333, Gln1374 for NBD2). Moreover, ATP phosphate moieties were

also restrained with surrounding Walker A residues. All distances were restrained using harmonic potentials for which minimal distances and force constants are reported in Supplementary Tables 13-14. Distance-based restraints were applied for thermalization and box equilibration steps. They were then smoothly removed along the first 10 ns of production runs. Restraints for Mg^{2+} ions were kept during the whole simulation.

Snapshots were saved every 100 ps. For each system, three replicas were performed to better sample the local conformational space. Each production run was carried out for 2.0-2.5 μs and 1.5-2.0 μs, respectively for IF and OF models (Supplementary Table 15). Indeed, simulations performed using OF model reached the equilibrium faster than IF conformations (see time-dependent RMSDs in Supplementary Fig. 30). In the present study, the total MD time is 112.4 μs.

4.3.   Analysis and visualization

Simulations were analysed using the CPPTRAJ[59] package, and in-house Python scripts taking advantage of MDAnalysis module[60,61]. Plots were obtained using the matplotlib v3.3.1 Python package[62]. Structure visualization and rendering were prepared using VMD software[63] (v1.9.3 and the alpha-v1.9.4). The so-called ABC structural parameters (*i.e.*, IC angle, EC angle, NDB distance, NBD twist and EC distance) were calculated using the same definition as proposed by Hofmann *et al.*[1]. Shortly, IC angle describes the IC opening of substrate entry and is defined by the angle between two vectors; both starting from the centre-of-mass of the whole extracellular region and directed toward either the IC region of TMH1, TMH2, TMH3, TMH6, TMH10 and TMH11 or the IC region of TMH4, TMH5, TMH7, TMH8, TMH9 and TMH12. Likewise, EC angle describes the EC opening for substrate release and is defined by the angle between two vectors starting from the centre-of-mass of both NBDs and directed toward either the EC region of TMH1, TMH2, TMH9, TMH10, TMH11, and TMH12 or the EC region of TMH3, TMH4, TMH5, TMH6, TMH7, and TMH8. EC distance was defined as the distance between the EC regions of TMH1, TMH2, TMH9, TMH10, TMH11, and TMH12 and the EC region of TMH3, TMH4, TMH5, TMH6, TMH7, and TMH8. NBD distance was
28

defined as the distance between the two NBD centres-of-mass. Since these definitions were applied to the present MD models of *b*MRP1 but also to available resolved structures of other ABC transporters, intracellular and extracellular regions were defined based on the membrane thickness proposed in the OPM database[41]. Residue selections for each system and each structure parameter are reported in Supplementary Table 1. For each system and each lipid bilayer, local free energy landscape was calculated using the InfleCS approach which takes advantage of Gaussian Mixture Models (GMM)[23,25]. Structural parameters (*i.e.*, IC and EC angles, NBD distance and twist) were taken to monitor the free energy landscape using a grid size set at 80, from 2 to 12 gaussian components for each GMM obtained by a maximum of 20 iterations.

H-bond analyses were performed using CPPTRAJ[59] in which distance and angle cutoffs were set at 3.5 Å and 120°, respectively. Substrate cavities were calculated using the Hole2.2 software[64] on a trajectory made of frames selected every 10 ns of the equilibrated part of each replica separately. Profiles were then averaged and PC P-atom z-density profiles were used to define the centre of the lipid bilayer membrane (z = 0). Regarding tilt angles, all systems were aligned to the OF *b*MRP-(ATP)$_2$ model embedded in POPC:POPE:Chol (2:1:1). Lipid distributions were obtained using the CPPTRAJ[59] package while membrane thicknesses were obtained using the MEMBPLUGIN for VMD[63,65].

Principal component analyses were also performed using the CPPTRAJ[59] package, focusing on the ABC core, defined by backbone atoms of TMH1 to TMH12, NBD1 and NBD2. System variabilities were investigated by carrying out independent PCA for each system for which each replica was aligned to an average structure of the system. Network analyses were performed using the VMD Network Analysis plugin[63,66]. Dynamic cross-correlation matrices were calculated separately for each replica on which Cα-atoms were selected as nodes and all the default restrictions (notSameResidue, notNeighboringCAlpha, notNeighboringPhosphate, notNeighboringResidue) were applied. Communities were then calculated using gncommunities[66]. Allostery network pathways were determined using the recent Allopath approach developed by Westerlund *et al.*[27,28]. Shortly, the

distant "communication efficiency" between two domains was obtained from the contact map and the mutual information matrix of protein residues as well as non-protein interactors such as surrounding lipid and bound molecules (*e.g.,* nucleotides and substrate). Such an approach also provides a betweenness profile which pictures the involvement of each component (*i.e.,* residue, lipid, substrate and nucleotide). For each lipid molecule, three nodes were defined corresponding to the polar hear group and the two lipid tails. ATP was also split into three nodes: purine and ribose moiety as well as triphosphate tail. LTX substrate was divided into three nodes: glutathione moiety, poly-unsaturated tail and hydroxypentanoic acid). Mg^{2+} ions and cholesterol molecules were considered as one node each. Atom selections per node per are reported in Supplemental Fig. 31. Allosteric pathways were calculated from substrate-binding pocket to each NBS, separately, as defined in Supplemental Table 9).

5. References

1. Hofmann, S. *et al.* Conformation space of a heterodimeric ABC exporter under turnover conditions. *Nature* **571**, 580–583 (2019).

2. Thomas, C. & Tampé, R. Structural and Mechanistic Principles of ABC Transporters. *Annu. Rev. Biochem.* **89**, 605–636 (2020).

3. Kroll, T., Prescher, M., Smits, S. H. J. & Schmitt, L. Structure and Function of Hepatobiliary ATP Binding Cassette Transporters. *Chem. Rev.* **121**, 5240–5288 (2021).

4. Johnson, Z. L. & Chen, J. ATP Binding Enables Substrate Release from Multidrug Resistance Protein 1. *Cell* **172**, 81-89.e10 (2018).

5. Wang, L. *et al.* Characterization of the kinetic cycle of an ABC transporter by single-molecule and cryo-EM analyses. *eLife* **9**, e56451 (2020).

6. Hutter, C. A. J. *et al.* The extracellular gate shapes the energy profile of an ABC exporter. *Nat. Commun.* **10**, 2260 (2019).

7. Stockner, T., Gradisch, R. & Schmitt, L. The role of the degenerate nucleotide binding site in type I ABC exporters. *FEBS Lett.* **594**, 3815–3838 (2020).

8. Johnson, Z. L. & Chen, J. Structural Basis of Substrate Recognition by the Multidrug Resistance Protein MRP1. *Cell* **168**, 1075-1085.e9 (2017).

9. Scortecci, J. F. *et al.* Cryo-EM structures of the ABCA4 importer reveal mechanisms underlying substrate binding and Stargardt disease. *Nat. Commun.* **12**, 5902 (2021).

10. Xu, D. *et al.* Cryo-EM structure of human lysosomal cobalamin exporter ABCD4. *Cell Res.* **29**, 1039–1041 (2019).

11. Kapoor, K., Pant, S. & Tajkhorshid, E. Active participation of membrane lipids in inhibition of multidrug transporter P-glycoprotein. *Chem. Sci.* **12**, 6293–6306 (2021).

12. Giacomini, K. M. *et al.* Membrane transporters in drug development. *Nat. Rev. Drug Discov.* **9**, 215–236 (2010).

13. Zamek-Gliszczynski, M. J. *et al.* Transporters in Drug Development: 2018 ITC Recommendations for Transporters of Emerging Clinical Importance. *Clin. Pharmacol. Ther.* **104**, 890–899 (2018).

14. Mateus, A. *et al.* Prediction of intracellular exposure bridges the gap between target- and cell-based drug discovery. *Proc. Natl. Acad. Sci.* **114**, (2017).

15. Nigam, S. K. What do drug transporters really do? *Nat. Rev. Drug Discov.* **14**, 29–44 (2015).

16. Chantemargue, B. *et al.* Structural patterns of the human ABCC4/MRP4 exporter in lipid bilayers rationalize clinically observed polymorphisms. *Pharmacol. Res.* **133**, 318–327 (2018).

17. Bickers, S. C., Benlekbir, S., Rubinstein, J. L. & Kanelis, V. Structure of Ycf1p reveals the transmembrane domain TMD0 and the regulatory region of ABCC transporters. *Proc. Natl. Acad. Sci.* **118**, e2025853118 (2021).

18. Moradi, M. & Tajkhorshid, E. Mechanistic picture for conformational transition of a membrane transporter at atomic resolution. *Proc. Natl. Acad. Sci.* **110**, 18916–18921 (2013).

19. Immadisetty, K., Hettige, J. & Moradi, M. Lipid-Dependent Alternating Access Mechanism of a Bacterial Multidrug ABC Exporter. *ACS Cent. Sci.* **5**, 43–56 (2019).

20. Srikant, S. & Gaudet, R. Mechanics and pharmacology of substrate selection and transport by eukaryotic ABC exporters. *Nat. Struct. Mol. Biol.* **26**, 792–801 (2019).

21. Verhalen, B. *et al.* Energy transduction and alternating access of the mammalian ABC transporter P-glycoprotein. *Nature* **543**, 738–741 (2017).

22. Olsen, J. A., Alam, A., Kowal, J., Stieger, B. & Locher, K. P. Structure of the human lipid exporter ABCB4 in a lipid environment. *Nat. Struct. Mol. Biol.* **27**, 62–70 (2020).

23. Westerlund, A. M. & Delemotte, L. InfleCS: Clustering Free Energy Landscapes with Gaussian Mixtures. *J. Chem. Theory Comput.* **15**, 6752–6759 (2019).

24. Sethi, A., Eargle, J., Black, A. A. & Luthey-Schulten, Z. Dynamical networks in tRNA:protein complexes. *Proc. Natl. Acad. Sci.* **106**, 6620–6625 (2009).

25. Westerlund, A. M., Harpole, T. J., Blau, C. & Delemotte, L. Inference of Calmodulin's Ca2+-Dependent Free Energy Landscapes via Gaussian Mixture Model Validation. *J. Chem. Theory Comput.* **14**, 63–71 (2018).

26. Zgarbová, M., Otyepka, M., Šponer, J., Hobza, P. & Jurečka, P. Large-scale compensation of errors in pairwise-additive empirical force fields: comparison of AMBER intermolecular terms with rigorous DFT-SAPT calculations. *Phys. Chem. Chem. Phys.* **12**, 10476–10493 (2010).

27. Westerlund, A. M., Fleetwood, O., Pérez-Conesa, S. & Delemotte, L. Network analysis reveals how lipids and other cofactors influence membrane protein allostery. *J. Chem. Phys.* **153**, 141103 (2020).

28. Kang, P. W. *et al.* Calmodulin acts as a state-dependent switch to control a cardiac potassium channel opening. *Sci. Adv.* **6**, eabd6798 (2020).

29. Janaszkiewicz, A. *et al.* Insights into the structure and function of the human organic anion transporter 1 in lipid bilayer membranes. *Sci. Rep.* **12**, 7057 (2022).

30. de Meyer, F. & Smit, B. Effect of cholesterol on the structure of a phospholipid bilayer. *Proc. Natl. Acad. Sci.* **106**, 3654–3658 (2009).

31. Liu, F., Zhang, Z., Csanády, L., Gadsby, D. C. & Chen, J. Molecular Structure of the Human CFTR Ion Channel. *Cell* **169**, 85-95.e8 (2017).

32. Lee, K. P. K., Chen, J. & MacKinnon, R. Molecular structure of human KATP in complex with ATP and ADP. *eLife* **6**, e32481 (2017).

33. Martens, C. *et al.* Lipids modulate the conformational dynamics of a secondary multidrug transporter. *Nat. Struct. Mol. Biol.* **23**, 744–751 (2016).

34. Martens, C. *et al.* Direct protein-lipid interactions shape the conformational landscape of secondary transporters. *Nat. Commun.* **9**, 4151 (2018).

35. Yang, J. *et al.* The I-TASSER Suite: protein structure and function prediction. *Nat. Methods* **12**, 7–8 (2015).

36. Webb, B. & Sali, A. Comparative Protein Structure Modeling Using MODELLER. *Curr. Protoc. Bioinforma.* **54**, (2016).

37. Salomon-Ferrer, R., Case, D. A. & Walker, R. C. An overview of the Amber biomolecular simulation package: Amber biomolecular simulation package. *Wiley Interdiscip. Rev. Comput. Mol. Sci.* **3**, 198–210 (2013).

38. Case, D. A. *et al. AMBER 2018*. (niversity of California, San Francisco).

39. Jo, S., Kim, T., Iyer, V. G. & Im, W. CHARMM-GUI: A web-based graphical user interface for CHARMM. *J. Comput. Chem.* **29**, 1859–1865 (2008).

40. Wu, E. L. *et al.* CHARMM-GUI *Membrane Builder* toward realistic biological membrane simulations. *J. Comput. Chem.* **35**, 1997–2004 (2014).

41. Lomize, M. A., Pogozheva, I. D., Joo, H., Mosberg, H. I. & Lomize, A. L. OPM database and PPM web server: resources for positioning of proteins in membranes. *Nucleic Acids Res.* **40**, D370–D376 (2012).

42. Jorgensen, W. L., Chandrasekhar, J., Madura, J. D., Impey, R. W. & Klein, M. L. Comparison of simple potential functions for simulating liquid water. *J. Chem. Phys.* **79**, 926–935 (1983).

43. MacKerell, A. D. *et al.* All-atom empirical potential for molecular modeling and dynamics studies of proteins. *J. Phys. Chem. B* **102**, 3586–3616 (1998).

44. Price, D. J. & Brooks, C. L. A modified TIP3P water potential for simulation with Ewald summation. *J. Chem. Phys.* **121**, 10096–10103 (2004).

45. Maier, J. A. *et al.* ff14SB: Improving the Accuracy of Protein Side Chain and Backbone Parameters from ff99SB. *J. Chem. Theory Comput.* **11**, 3696–3713 (2015).

46. Dickson, C. J. *et al.* Lipid14: The Amber Lipid Force Field. *J. Chem. Theory Comput.* **10**, 865–879 (2014).

47. Meagher, K. L., Redman, L. T. & Carlson, H. A. Development of polyphosphate parameters for use with the AMBER force field. *J. Comput. Chem.* **24**, 1016–1025 (2003).

48. Galindo-Murillo, R. *et al.* Assessing the Current State of Amber Force Field Modifications for DNA. *J. Chem. Theory Comput.* **12**, 4114–4127 (2016).

49. Joung, I. S. & Cheatham, T. E. Determination of Alkali and Halide Monovalent Ion Parameters for Use in Explicitly Solvated Biomolecular Simulations. *J. Phys. Chem. B* **112**, 9020–9041 (2008).

50. Joung, I. S. & Cheatham, T. E. Molecular Dynamics Simulations of the Dynamic and Energetic Properties of Alkali and Halide Ions Using Water-Model-Specific Ion Parameters. *J. Phys. Chem. B* **113**, 13279–13290 (2009).

51. Wang, J., Wolf, R. M., Caldwell, J. W., Kollman, P. A. & Case, D. A. Development and testing of a general amber force field. *J. Comput. Chem.* **25**, 1157–1174 (2004).

52. Wang, J., Wang, W., Kollman, P. A. & Case, D. A. Automatic atom type and bond type perception in molecular mechanical calculations. *J. Mol. Graph. Model.* **25**, 247–260 (2006).

53. Vanquelef, E. *et al.* R.E.D. Server: a web service for deriving RESP and ESP charges and building force field libraries for new molecules and molecular fragments. *Nucleic Acids Res.* **39**, W511-517 (2011).

54. Darden, T., York, D. & Pedersen, L. Particle mesh Ewald: An N·log(N) method for Ewald sums in large systems. *J. Chem. Phys.* **98**, 10089–10092 (1993).

55. Davidchack, R. L., Handel, R. & Tretyakov, M. V. Langevin thermostat for rigid body dynamics. *J. Chem. Phys.* **130**, 234101 (2009).

56. Berendsen, H. J. C., Postma, J. P. M., van Gunsteren, W. F., DiNola, A. & Haak, J. R. Molecular dynamics with coupling to an external bath. *J. Chem. Phys.* **81**, 3684–3690 (1984).

57. Åqvist, J., Wennerström, P., Nervall, M., Bjelic, S. & Brandsdal, B. O. Molecular dynamics simulations of water and biomolecules with a Monte Carlo constant pressure algorithm. *Chem. Phys. Lett.* **384**, 288–294 (2004).

58. Wen, P.-C., Verhalen, B., Wilkens, S., Mchaourab, H. S. & Tajkhorshid, E. On the Origin of Large Flexibility of P-glycoprotein in the Inward-facing State. *J. Biol. Chem.* **288**, 19211–19220 (2013).

59. Roe, D. R. & Cheatham, T. E. PTRAJ and CPPTRAJ: Software for Processing and Analysis of Molecular Dynamics Trajectory Data. *J. Chem. Theory Comput.* **9**, 3084–3095 (2013).

60. Michaud-Agrawal, N., Denning, E. J., Woolf, T. B. & Beckstein, O. MDAnalysis: A toolkit for the analysis of molecular dynamics simulations. *J. Comput. Chem.* **32**, 2319–2327 (2011).

61. Gowers, R. J. *et al.* MDAnalysis: A Python Package for the Rapid Analysis of Molecular Dynamics Simulations. *Proc. 15th Python Sci. Conf.* 98–105 (2016) doi:10.25080/Majora-629e541a-00e.

62. Hunter, J. D. Matplotlib: A 2D Graphics Environment. *Comput. Sci. Eng.* **9**, 90–95 (2007).

63. Humphrey, W., Dalke, A. & Schulten, K. VMD: visual molecular dynamics. *J. Mol. Graph.* **14**, 33–38, 27–28 (1996).

64. Smart, O. S., Neduvelil, J. G., Wang, X., Wallace, B. A. & Sansom, M. S. HOLE: a program for the analysis of the pore dimensions of ion channel structural models. *J. Mol. Graph.* **14**, 354–360, 376 (1996).

65. Guixà-González, R. *et al.* MEMBPLUGIN: studying membrane complexity in VMD. *Bioinforma. Oxf. Engl.* **30**, 1478–1480 (2014).

66. Eargle, J. & Luthey-Schulten, Z. NetworkView: 3D display and analysis of protein·RNA interaction networks. *Bioinformatics* **28**, 3000–3001 (2012).

6. Acknowledgements

We acknowledge Dr. Benjamin Chantemargue and Mehdi Benmameri for stimulating and fruitful scientific discussions. We thank Xavier Montagutelli from the IT department of Limoges University for technically supporting supercomputer facilities. Calculations were performed using CALI ("CAlcul en LImousin") and "Baba Yaga" supercomputers hosting in Limoges University, as well as on the "Jean-Zay" national supercomputer from IDRIS HPC resources under the allocations 2020-A0080711487 and 2021-A0100711487 made by GENCI. This work was supported by grants from the "Agence Nationale de la Recherche" (ANR-19-CE17-0020-01 IMOTEP), Région Nouvelle Aquitaine and "Institut National de la Santé et de la Recherche Médicale" (INSERM, AAP-NA-2019-VICTOR).

7. Author Contributions

Á.T and F.D.M conceived the study. Á.T and F.D.M. conducted all the simulations. Á.T, A.J., V.C and F.D.M analysed the simulations. Á.T. and F.D.M. interpretated the results and discussed them together with A.J. and V.C. Á.T. and F.D.M wrote the manuscript and it was edited, reviewed and approved by all authors.

8. Competing interests statement

The authors declare no competing interests.