Boosting chloroplast ribosome biogenesis by a plastidial DEAD-box RNA helicase is critical for high light acclimation

El Batoul Djouani-Tahri[a], Sreedhar Nellaepalli[b*], Pascaline Auroy[a], Emmanuelle Billon[a], Adrien Burlacot[a#], Frédéric Chaux-Jukic[a+], Stéphan Cuiné[a], Virginie Epting[a], Marie Huleux[a], Bart Ghysels[a♣], Miriam Schulz-Raffelt[a§], Isabelle Te[a], Sabine Brugière[c], Yohann Couté[c], Yuichiro Takahashi[b], Yonghua Li-Beisson[a], Gilles Peltier[a¶]

[a] Aix Marseille Univ, CEA, CNRS, Institut de Biosciences et Biotechnologies Aix-Marseille, CEA Cadarache, 13108 Saint Paul-lez-Durance, France

[b] Research Institute for Interdisciplinary Science, Okayama University, 3-1-1 Tsushima-naka, Kita-ku, Okayama, 700-8530, Japan

[c] Université Grenoble Alpes, INSERM, CEA, UMR BioSanté U1292, CNRS, CEA, FR2048, 38000 Grenoble, France

[*] Present address: Department of Plant Sciences, University of Oxford, United Kingdom

[#] Present address: Department of Plant Biology, The Carnegie Institution, Stanford, CA, 94305 USA

[+] Present address: Sorbonne Université, CNRS, UMR7238, Institut de Biologie Paris-Seine, Laboratory of Computational and Quantitative Biology, 75005 Paris, France

[♣] Present address: Laboratory of Bioenergetics, Institute of Plant Biology B22, University of Liège, Liège, Belgium

[§] Present address: Molekulare Biotechnologie & Systembiologie, TU Kaiserslautern, Paul-Ehrlich Straße 23, D-67663 Kaiserslautern, Germany

[¶]Address correspondence to gilles.peltier@cea.fr

ABSTRACT

Photosynthetic organisms have developed sophisticated strategies to fine-tune light energy conversion

to meet the metabolic demand, thereby optimizing growth in fluctuating light environments. Although

mechanisms such as energy dissipation, photosynthetic control, or the photosystem II (PSII) damage

and repair have been widely studied, little is known about the regulation of protein synthesis capacity

during light acclimation. By screening a *Chlamydomonas reinhardtii* insertional mutant library using

chlorophyll fluorescence imaging, we isolated a high chlorophyll fluorescence mutant (hf_0) defected in

a gene encoding a putative plastid targeted DEAD-box RNA helicase called CreRH22. CreRH22 is

rapidly induced upon illumination and belongs to the GreenCut, a set of proteins specific to

photosynthetic organisms. While photosynthesis is slightly affected in the mutant under low light

(LL), exposure to high light (HL) induces a marked decrease in both PSII and PSI, and a strong

alteration of the light-induced gene expression pattern. These effects are explained by the inability of

hf_0 to increase plastid ribosome amounts under HL. We conclude that CreRH22, by promoting

ribosomal RNA precursor maturation in a light-dependent manner, enables the assembly of extra-

ribosomes required to synthesize photosystem subunits at a higher rate, a critical step in the

acclimation of algae to HL.

INTRODUCTION

In natural environments, photosynthetic organisms face highly fluctuating light conditions, and developed a complex network of regulatory mechanisms to adjust the capture and transformation of light energy to the metabolic demand. Such mechanisms are highly desirable since any disequilibrium between the conversion and use of light energy may result in severe photo-damage due to the formation of harmful reactive oxygen species (Niyogi, 1999). A wide range of mechanisms dynamically regulate light-energy conversion and photosynthetic electron flow, and operate at different time scales (Erickson et al., 2015). The short-term response involves energy dissipation by photosystems through a pH-dependent de-excitation of chlorophylls (Peers et al., 2009). On longer time scales (minutes to hour), other acclimation mechanisms operate in addition to the pH dependent NPQ, including state transitions (Depege et al., 2003; Allorent et al., 2013), adjustment of chlorophyll antenna size and photosystem stoichiometry (Bonente et al., 2012; Erickson et al., 2015). The PSII damage and repair cycle, although initially considered as a wasteful process (called photoinhibition) and involving selective destruction of PSII core subunits followed by active protein synthesis (Ohad et al., 1984), is now viewed as part of a dynamic and sophisticated machinery in which PSII is degraded and further repaired (Murata et al., 2012; Adams et al., 2013; Erickson et al., 2015; Li et al., 2018). The PSII repair cycle would protect the photosynthetic electron transfer chain, particularly PSI (Tikkanen et al., 2014). During the PSII repair cycle, translation of plastid-encoded PSII subunits D1, D2 and CP43 is enhanced (Christopher and Mullet, 1994; Kettunen et al., 1997; Minai et al., 2006; Jarvi et al., 2015). However, despite extensive studies on PSII repair, mechanisms involved in the regulation of translation remain largely unknown (Sun and Zerges, 2015). Analysis of gene expression during a day-night cycle showed that nuclear genes encoding chloroplast ribosome proteins are rapidly and transiently expressed upon sudden HL exposure, indicating that ribosome biogenesis is critical in such conditions (Zones et al., 2015).

DEAD-box RNA helicases catalyze ATP-dependent unwinding of RNA structures, and are involved in many aspects of RNA metabolism, including RNA synthesis, cleavage, modification, or ribosome biogenesis (Silverman et al., 2003). *CrhR*, the unique DEAD-box RNA helicase present in the genome of the cyanobacterium *Synechocystis* sp. PCC 6803, participates in redox regulation of gene expression (Kujat and Owttrim, 2000), low temperature acclimation of photosynthesis (Rosana et al., 2012; Sireesha et al., 2012), and was recently shown to interact with transcripts associated with photosynthesis (Migur et al., 2021). In plant chloroplasts, DEAD-box RNA helicases participate in ribosome biogenesis (Asakura et al., 2012; Chi et al., 2012), chloroplast differentiation (Wang et al., 2000), and responses to diverse abiotic stresses (Nawaz and Kang, 2017), but a role of these enzymes in the light-response of eukaryotic photosynthesis has not been reported so far.

With the aim to unveil new regulatory and acclimation mechanisms of photosynthesis, a forward genetic approach was developed in the unicellular microalga *Chlamydomonas reinhardtii* by screening an insertional mutant library based on the analysis of chlorophyll fluorescence (Tolleter et

al., 2011). We report here on the isolation of a high chlorophyll fluorescence mutant (hf_0) defected in a

gene encoding a putative plastid DEAD-box RNA helicase (*CreRH22*), and show its involvement in

ribosomal RNA processing and ribosome biogenesis. *CreRH22* belongs to the GreeenCut2

(Karpowicz et al., 2011) and its expression is rapidly and transiently induced upon HL exposure.

Photosynthetic activity and photoautotrophic growth of the hf_0 mutant are affected at HL intensity. We

conclude that rapid induction of CreRH22 promotes ribosome biogenesis upon HL exposure, thus

increasing the translation capacity and enabling efficient turnover of plastid-encoded photosystems

subunits.

RESULTS

Isolation of high chlorophyll fluorescence *Chlamydomonas* mutant defected in a DEAD-box RNA

helicase. From the screening of an insertion mutant library based on the analysis of chlorophyll

fluorescence transients of *Chlamydomonas* colonies grown in photo-autotrophic conditions (Tolleter et

al., 2011), a mutant (hf_0) showing a high F_0 fluorescence level was isolated (Fig. 1A). The hf_0 mutant

harbors a single insertion of the *AphVIII* paromomycin resistance cassette (Fig. 1B and Supplemental

Fig. S1) located within the 7^{th} exon of a gene (Cre03.g166650) encoding a putative DEAD-box RNA

helicase named as CreRH22 (Fig. 1B). The Cre03.g166650 transcript was not detected by RT-PCR in

hf_0 (Fig.1C). CreRH22 shares common properties of DEAD-box RNA helicases, including the Asp-

Glu-Ala-Asp (DEAD) sequence, an RNA-binding motif and an ATP hydrolyzing domain (Fig. 1B).

Phylogenetic analysis showed that *CreRH2* belongs to a DEAD-box RNA helicase clade containing a

previously characterized member in Arabidopsis (AtRH22) (Chi et al., 2012) (Supplemental Fig. S2).

AtRH22 was reported to be targeted to the chloroplast and to be involved in ribosomal RNA

processing and chloroplast ribosome biogenesis (Chi et al., 2012). By using the prediction tool

Predalgo specifically developed for intracellular targeting of microalgae (Tardif et al., 2012),

CreRH22 was predicted to be chloroplast targeted, like six other putative DEAD-box RNA helicases

encoded by the *C. reinhardtii* genome (Supplemental Fig. S3). Moreover, the CreRH22 protein

belongs to the GreenCut2 (identified as Hel15), an inventory of proteins identified on a phylogenomic

basis, conserved in plants and green algae but not in non-photosynthetic organisms (Karpowicz et al.,

2011). Complementation was attempted by both nuclear and chloroplast transformation. For nuclear

complementation, a genomic DNA fragment corresponding to the full *CreRH22* gene was cloned into

a vector containing a hygromycin resistance cassette, and resistant clones were screened by

chlorophyll fluorescence imaging (Supplemental Fig. S4). Full complementation was observed in one

strain out of 400 hygromycin resistant clones. For plastid transformation, a synthetic gene with a

codon usage adapted to the *Chlamydomonas* chloroplast genome was used (Supplemental Fig. S5).

Two independent homoplasmic chloroplast transformants were analyzed, chlorophyll fluorescence

measurements showing a full recovery of the wild-type phenotype in both transformants

(Supplemental Fig. S4). We conclude that inactivation of *CreRH22* is responsible for the *hf₀*

chlorophyll fluorescence phenotype, complementation by plastid expression confirming chloroplast

targeting.

We observed that the chlorophyll fluorescence phenotype of the *hf₀* mutant vanishes when

batch cultures reached a high cell density. Since the photon flux received by each cell is lower in a

dense culture than in a diluted culture due to cell-shading, we hypothesized that the mutant phenotype

may depend on the light intensity. Indeed, when cells were grown under HL (240 µmol photons. m^{-2} s^{-1}

) chlorophyll fluorescence induction of the mutant was strongly affected, showing higher F_0 and Fs

levels and reduced PSII yields as compared to wild-type control and two complemented lines (Fig. 2A,

C). However, when cells were grown under LL (15 µmol photons m^{-2} s^{-1}) differences in the F_0 levels

and PSII yields between *hf₀* and control strains were much smaller (Fig. 2B, D).

Ribosomal RNA processing is affected in *hf₀*. We next investigated whether CreRH22 is involved in

chloroplast ribosomal RNA processing by analyzing this mechanism in *hf₀*. Chloroplast ribosomal

RNAs are organized in polycistronic transcription units, which need to be matured into a set of

overlapping RNAs through a series of processing steps. The non-mature polycistronic rRNA

undergoes endo-nucleolytic cleavages to produce mature 7S, 3S, 23S and 5S rRNAs (Holloway and

Herrin, 1998) (Fig. 3A). Accumulation and processing of chloroplast ribosomal RNAs were further

examined by Northern blot using specific probes for mature 3S, 7S, and 23S rRNAs. We found that

RNA precursor levels are increased in *hf₀* as compared to the wild-type, mature RNA levels being

reduced accordingly (Fig. 3B). Plastid transcript level encoding PSII and PSI subunits, analyzed by

RT-qPCR, showed a slightly higher level of transcripts in *hf₀* compared to the wild-type both under LL

(30 µmol photons m^{-2} s^{-1}) and HL (240 µmol photons m^{-2} s^{-1}) (Supplemental Fig. S6A, B). Analysis of

chloroplast transcripts encoding non-polycistronic ribosomal components (*rpl36* and *rps18)* showed

no obvious difference in the levels of transcripts (Supplemental Fig. S6C, D), the 16S transcript level

being lower in *hf₀* compared to the wild-type (Supplemental Fig. S6E). We conclude from these

experiments that, as previously reported for AtRH22 (Chi et al., 2012), CreRH22 is involved in

chloroplast ribosomal RNA processing, its defect resulting in an accumulation of unprocessed rRNA

forms and thereby a decrease in the amount of plastid ribosomes.

Accumulation and turnover of PSII and PSI subunits are affected in *hf₀*. To better characterize the

effect of CreRH22 on protein synthesis, quantitative immuno-analysis was carried out on a set of

chloroplast proteins, including plastid ribosomal proteins and components of the photosynthetic

electron transport chain (Fig. 4). Under LL, no differences in protein amounts were observed between

hf₀ and the wild-type, except for chloroplast ribosomal proteins L30 and S21, both present at lower

amounts in *hf₀*. Upon HL exposure, the amount of both ribosomal proteins increased in the wild-type

while remaining at a low level in *hf₀*. In wild-type cells, HL exposure induced a slight decrease in the

amounts of PSII subunits PsbA (D1), PsbC (CP43) and PsbD (D2), and PSI subunits PsaA, PSAD and

PSAF, but the effect was more pronounced in hf_0. The amounts of other chloroplast components,
including AtpB, a subunit of the ATP synthase (CF0-CF1), PetA a subunit of the cytochrome b_6f
complex, and LHCB2 a light harvesting subunit remained unchanged. [35]S pulse-labeling experiments
of thylakoid proteins were then carried out to determine to what extent such changes in protein
amounts reflect changes in protein synthesis activity. No difference in pulse labeling patterns was
observed under LL between wild-type, hf_0 and two complemented lines (Fig. 5). However, upon 6h of
HL exposure a lower D1 labeling was observed in hf_0, the effect being reversed in both complemented
lines. The turnover of D1 is known as the highest of plastid proteins due to the existence of an efficient
damage/repair system (Jarvi et al., 2015). Note that slight differences in labeling between wild-type
and hf_0 were also observed, although at a lower extent, on other proteins including LHCII and the PSI
subunit PsaC. On the other hand, the labeling of ATP synthase (CF0-CF1) subunits AtpA and AtpB
was unaffected in the mutant. We conclude from these experiments that the protein
translation/synthesizing capacity in hf_0 is lower than in the wild-type due to a lower ribosome content,
which becomes limiting under HL, and results in a decreased accumulation of plastid proteins actively
synthesized under HL, such as D1.

In order to confirm that PSII inhibition observed in hf_0 under HL results from a defect in
protein synthesis capacity, we used chloramphenicol, an inhibitor of chloroplast protein synthesis. In
the absence of chloramphenicol, $\Phi_{PSIImax}$ and operating PSII yield progressively decreased upon HL
exposure in hf_0 but not in wild-type or in complemented lines (Fig. 6 A, C). As previously reported
(Schuster et al., 1988), PSII was more strongly inhibited by HL in the presence of chloramphenicol,
similar inhibitions being observed in hf_0, wild-type and complemented lines after 7h to 24h exposure
(Fig. 6 B,D). Since the chloramphenicol treatment of wild-type somehow mimicked the CreRH22
mutation, we conclude that the HL sensitivity of hf_0 likely results from a decreased protein synthesis
capacity. Since both turnover and accumulation of some PSII and PSI subunits were affected in HL-
acclimated hf_0 cells, we measured how PSII and PSI activities are impacted (Supplemental Fig. S7).
While a 50% reduction in the maximal PSII yield was observed in hf_0 under HL, a smaller decrease in
the PSII/PSI ratio was measured, consistent with the existence of reduced PSI activity, although to a
lesser extent than PSII inhibition.

Chlorophyll protein complexes are affected in hf_0. In order to gain insight on changes in
photosynthesis machinery occurring in hf_0 upon HL exposure, we first performed low temperature
(77K) chlorophyll fluorescence measurements (Fig. 7). When grown under LL both wild-type and
mutant strains showed closely related patterns, with a slightly higher fluorescence emission peak at
712nm (E_{712nm}) in the mutant (Fig. 7A). Upon HL exposure, the E_{712nm} peak dramatically increased in
hf_0 (Fig. 7B). The chlorophyll emission ratio (E_{712nm}/E_{686nm}) and the F_0 level (measured as the F_0/Fm
ratio) both strongly increased in the mutant while remaining at nearly constant values in the wild-type
(Fig. 7C, D). In order to determine whether the E_{712nm} fluorescence peak increase is related to the

phenomenon of state transition, phosphorylation of thylakoid proteins was analyzed (Fig. 7E). During

state transitions, reduction of the PQ pool triggers the STT7 kinase, which phosphorylates LHCII and

PSII subunits. The phosphorylation status of thylakoid proteins of wild-type and hf_0 are similar under

LL, and slightly differed under HL (Fig. 7E). However, contrary to what would have been expected if

hf_0 was more in state 2 than the wild-type, a lower phosphorylation status was observed in the mutant

as compared to the wild-type upon HL exposure. Moreover, the blue shift of the 712nm peak observed

in HL acclimated hf_0 cells (Fig. 7B) is likely due to an increased fluorescence emission from the PSI

core or LHCI oligomers, suggesting that part of PSI-LHCI is dissociated into PSI core and LHCI

oligomer or that LHCI oligomers accumulate because of the absence of PSI core (Takahashi et al.,

2004). We conclude from these experiments that both E_{712nm} and F_0 increases do not result from a

transition to state 2, but likely reflect the presence of LHCI oligomers. We then performed blue native

gel electrophoresis of thylakoid protein complexes, and observed no major difference in the

distribution of chlorophyll protein complexes in hf_0 as compared to the wild-type under LL (Fig. 7F).

However, under HL exposure, the pattern of chlorophyll protein complexes of hf_0 differed from that of

the wild-type, two weak upper bands (PSII-LHCII supercomplexes) disappearing and a lower band

(LHCI oligomer) appearing in the mutant, a similar effect being observed when LL wild-type cells

were treated with chloramphenicol (Supplemental Fig. S8). Two-dimensional blue native-PAGE

showed that a group of polypeptides, associated in wild-type cells to a large complex, are found

associated to a smaller complex in hf_0 (Fig. 8A, B). Mass spectrometry-based proteomic analysis was

performed on a gel strip taken on the 2D gel in the ~20-35kDa molecular weight range (see red box on

Fig. 8B). Various polypeptides were identified, and notably several subunits of PSI (PsaA, PsaB,

PsaC, PSAD and PSAF), light-harvesting proteins of PSI (LHCA3, LHCA4, LHCA5, LHCA6,

LHCA7), several subunits of PSII (PsbA, PsbB, PsbC and PsbD), and one Light Harvesting Complex

II (LHCII) polypeptide (LHCBM8), a component of trimeric LHCII (Girolomoni et al., 2017)

(Supplemental Data Set S1). Polypeptides with molecular weights comprised between 20 and 35kDa,

which corresponds to the gel strip taken for the proteomic analysis, include the five LHCA

polypeptides, the two nuclear-encoded PSI subunits (PSAD and PSAF) and LHCBM8. The presence

of higher molecular weight polypeptides such as PsaA, PsaB, PsbA, PsbB, PsbD in this area of the gel

suggests that some degradation of PSI and PSII photosystems occurred in HL-acclimated hf_0 cells. The

presence of orphan LHCAs, normally associated with high MW complexes (PSI-LHCI) in a stable

manner (Takahashi et al., 2004; Nellaepalli et al., 2021), indicates a defect in PSI organization, in line

with 77K chlorophyll fluorescence data (Fig. 7B). It is inferred from this experiment that accumulation

of PSI-LHCI and PSII-LHCII supercomplexes, which normally occurs when the synthesis of subunits

encoded by nuclear gene and chloroplast genes is properly synchronized, is impaired in hf_0 under HL

conditions. As a consequence, PSII-LHCII and PSI-LHCI bands disappear, some PSI and PSII

reaction centers are degraded and LHCI oligomers accumulate at detectable levels. The increase of

LHCII monomer observed in hf_0 under HL (Fig. 7F) reflects the fact that in contrast to LHCI, only a

small fraction LHCII is normally associated to PSII-LHCII supercomplexes (Tokutsu et al., 2012).

***CreRH22* expression is transiently upregulated during light transients.** By searching data related

to *CreRH22* (Cre03.g166650) transcript levels in a previous study reporting the dynamics of the *C.*

reinhardtii transcriptome in cells synchronized by a night-day cycle (Zones et al., 2015), we found that

CreRH22 expression shows a sharp and transient increase at the beginning of the light period

(Supplemental Fig. S9A). In this study a gene cluster named "light stress cluster" characterized by a

similar gene expression profile as *CreRH22* was defined. This cluster contains several *HSP* genes

(such as *HSP70B* and *HSP90C*), as well as light stress genes such as *PSBS2* or *VTC2* (Zones et al.,

2015). Although *CreRH22* was not assigned in this work as a member of the "light stress cluster", its

expression pattern actually meets most criteria used to define this cluster during the light period, the

transient expression increase observed during the dark period (reaching 51% of the maximum

expression) preventing the classification in this cluster (criteria limit <40%). In order to determine

whether *CreRH22* expression is regulated by a light transient independent of circadian regulations, we

then performed a dark to light transient on non-synchronized cells, which showed that *CreRH22*, like

HSP70B and *HSP90C,* is induced by a light (Supplemental Fig. S9B).

Light-dependent nuclear gene expression is severely impaired in hf_0. RNAseq analysis was then

performed in order to determine how the light-induced transcriptional response is affected in the

absence of CreRH22. For this purpose, *C. reinhardtii* cells were grown under LL in PBR operated as

turbidostats thus allowing acclimation to constant growth conditions, and then switched to HL. Global

transcript profiles were analyzed by RNA sequencing after 24h LL acclimation and then after 2h and

24h of HL exposure (Supplemental Data Set S2; Fig. 9). In wild-type cells, 3020 differentially-

expressed genes (DEGs) were up-regulated, and 445 down-regulated upon 2h of HL exposure (log

2(Fold Change) > 1 and <-1, respectively) (Fig. 9A). The number of up-regulated DEGs doubled (up

to 6023) after 24h under HL, the number of down-regulated DEGs remaining at a low level (576). The

effect of HL on gene expression was strongly attenuated in hf_0 in comparison to the wild-type, since

only 1241 and 1291 genes were up-regulated after 2h and 24h of HL, respectively. In contrast, the

number of down-regulated genes doubled as compared to the wild-type. MapMan classification of up-

regulated DEGs showed that most represented gene categories are distributed in a similar manner in

both strains at different time points (Fig. 9 B,C).

Since CreRH22 is involved in chloroplast ribosomal RNA processing, we focused on the

expression pattern of ribosomal protein genes (RPGs) of translational machineries located in the

cytosol, chloroplast, and mitochondria (Fig. 10). A previous study has established that RPG expression

is highly coordinated in the three cellular compartments during light-dark cycles, the expression of

cytosolic RPGs peaking at the beginning of the dark period, while the expression of chloroplast RPGS

is maximal at the very beginning of the light period (Zones et al., 2015). When wild-type cells were

shifted from LL to HL, the maximal RPG expression of cytosolic components was observed under LL
and strongly decreased upon HL exposure. In hf_0 cells, cytosolic RPG expression was much less
affected upon HL exposure (Fig. 10 A,D). The effect of HL on RPG expression of chloroplasts
components was dramatically different. Indeed, while expression of chloroplast RPGs was essentially
not affected by the HL shift in wild-type cells, a strong and progressive increase in chloroplast RPGs
expression was observed in hf_0 2h and 24h after the HL shift (Fig. 10 B,E). On the other hand,
mitochondrial RPGs showed a similar expression pattern in the wild-type and hf_0 after the HL shift, a
similar increase of expression being observed in both strains (Fig. 10 C,F). The dramatic difference
between chloroplast RPGs expression observed in hf_0 and wild-type cells in response to the HL
sharply contrasts with changes in ribosomes abundances, since an increase in ribosomal protein
amounts was observed in the wild-type and no change in hf_0 (Fig. 4). The increase in chloroplast
ribosome abundance observed in wild-type cells in response to HL and the absence of changes in
chloroplast RPG expression strongly suggests that chloroplast ribosome abundance is mainly driven
by post-transcriptional regulations (*i.e.* increased translation of ribosome subunits). The strong
increase in chloroplast RPG expression observed in hf_0 upon HL exposure and the absence of changes
in ribosome abundance suggests that 23S rRNA processing by CreRH22 is a critical step in the
biogenesis of additional ribosomes in wild-type, which cannot be compensated by an increased
expression of plastid RPGs in hf_0.

Biomass productivity is decreased under HL in hf_0. We then aimed to assess the impact of the
CreRH22 defect on growth capacity and biomass productivity under different light regimes by
performing PBR experiments in a turbidostat mode (Fig. 11). Steady-state dilution rates of cultures
maintained at constant biomass concentration allowed to determine specific growth rate, which
increased as a function of light intensity in both wild-type and hf_0 cell cultures until a saturation
plateau, which reached a 40% lower value in hf_0 as compared to the wild-type (Fig. 11A).
Remarkably, the decrease in the cellular chlorophyll content observed upon HL acclimation of *C.*
reinhardtii cells (Bonente et al., 2012) was strongly diminished in hf_0 (Fig. 11B) indicating a strong
impairment of light acclimation in the mutant. Under stress conditions, such as nutrient depletion
(Siaut et al., 2011) or HL (Goold et al., 2016), *Chlamydomonas* cells accumulate starch and oil, which
is generally viewed as the result of an imbalance between photosynthesis and growth capacities. While
intracellular starch and oil reached high levels in the wild-type at HL, starch accumulation was much
lower in hf_0 (Fig. 11D), the difference in oil contents being less marked (Fig. 11E). We conclude from
this experiment that CreRH22 is critical for Chlamydomonas to acclimate to HL conditions and
achieve maximal biomass productivity.

DISCUSSION
We have shown here that CreRH22, a DEAD-box RNA helicase targeted to the chloroplast and

involved in ribosomal RNA processing and ribosome biogenesis, has a critical function during HL

acclimation in the green unicellular alga *C. reinhardtii*. CreRH22 belongs to the GreenCut2

(Karpowicz et al., 2011) and shows similar gene expression profile as a transiently expressed cluster

of light stress genes (Zones et al., 2015). We propose that rapid induction of CreRH22 upon HL

exposure promotes ribosome assembly, thus increasing the translation capacity and allowing synthesis

of plastid-encoded proteins of the photosynthesis machinery, and particularly the D1 protein of PSII to

avoid photoinhibition. In the absence of CreRH22, plastid-encoded PSII subunits (and to a lesser

extent the PSI subunit PsaA/B and PsaC) are not synthesized at a sufficient rate during HL exposure,

thus leading to an imbalance between the synthesis of nuclear- and plastid- encoded subunits of

photosystems, to the accumulation of lower levels of PSII-LHCII and PSI-LHCI complexes and to the

accumulation of oligomeric LHCI, thus explaining the increase in the F_0 chlorophyll fluorescence. As

a consequence, light-induced gene expression, photosynthesis and biomass production are strongly

affected under HL (Figs. 11 and 12).

Physiological functions of plastidial DEAD-box RNA helicases. DEAD-box RNA helicases are

involved in different aspects of RNA metabolism and participate to many cellular functions (Linder

and Jankowsky, 2011). In Arabidopsis, among 58 DEAD-box RNA helicases identified in the genome,

ten and eight are predicted to be respectively targeted to chloroplasts or mitochondria and involved in

plant response to abiotic stresses (Nawaz and Kang, 2017). Proteomic analysis revealed the presence

of six of the plastidial DEAD-box RNA helicases, including RH22, in high molecular mass stromal

complexes, thus suggesting an involvement of these proteins in ribosome biogenesis (Olinares et al.,

2010), which was confirmed for RH3 and RH22 by studies of Arabidopsis mutants (Asakura et al.,

2012; Chi et al., 2012). However, despite the presence of DEAD-box RNA helicases in the chloroplast

stroma, a possible function of these proteins in regulating photosynthesis has not been documented so

far. Indeed, loss of the Arabidopsis *RH22* gene is lethal, and knock-down mutants show a severe

growth phenotype (Chi et al., 2012). In contrast, disruption of the *C. reinhardtii* gene confers a light-

dependent phenotype, the mutant growing normally under LL and at a slower rate than the wild-type

under moderate to HL. Such a phenotype difference between plants and unicellular algae may be

related to the requirement of a high protein synthesis capacity during the process of plastid biogenesis

and/or differentiation occurring during plant greening (Sun and Zerges, 2015). In this context, any

disturbance in the translation machinery has a strong impact on plant chloroplast biogenesis. However,

unicellular organisms such as microalgae do not experience such a greening process since chloroplasts

divide as green organelles (Jarvis and Lopez-Juez, 2013). Both CreRH22 and AtRH22 are thus likely

needed when active protein synthesis is required, either during chloroplast biogenesis in plants, or

during the PSII repair cycle in HL conditions in algae and most probably also in plants. However, the

later function is difficult to assess due to the strong phenotype of the Arabidopsis RH22 mutant.

CreRH22 is part of a network of proteins involved in ribosomal biogenesis. Remarkably, five out of the seven DEAD-box RNA helicases predicted to be plastid targeted in Chlamydomonas, including CReRH22, belong to the gene cluster 1 (Supplemental Fig. S3), defined as genes showing a strong induction during the first hour of illumination followed by a sharp decrease (Zones et al., 2015). These genes are expressed before the maximal expression of most ribosome protein genes, which peaked at two hours of light exposure (Zones et al., 2015), in line with a function in ribosome biogenesis. Very recently, four of these DEAD-box RNA helicases, including CreRH22, were identified as interacting with chloroplast ribosome proteins (Westrich et al., 2021) (Supplemental Fig. S3). Chloroplast ribosomes are made of a small subunit (30S) containing the 16S ribosomal RNA and a large subunit (50S) containing 23S ribosomal RNA. While AtRH22 in Arabidopsis (Chi et al., 2012) and CreRH22 in Chlamydomonas (this work) are involved in 23S ribosomal RNA maturation, a bacterial-type ribosome binding factor (RbfA) was proposed in Arabidopsis to participate in the processing of 16S ribosomal RNA. RbfA also belongs to the GreenCut, and was shown in Arabidopsis to be essential for chloroplast development and photoautotrophic growth (Fristedt et al., 2014). As it is the case for CreRH22, the Chlamydomonas RBF1 homolog (Cre02.g145000) belongs to the gene cluster 1 (Zones et al., 2015), was found in high molecular mass stromal complexes (Olinares et al., 2010), and was enriched in 30S ribosomal fractions interactome (Westrich et al., 2021). Together, these studies suggest that chloroplast CreRH22, in association with other plastid-targeted DEAD-box RNA helicases, is involved in ribosome biogenesis in response to the light supply. CreRH22 would be part of a complex network of regulatory proteins including other plastid targeted DEAD-box RNA helicases and other factors such as RBF1, involved in a coordinated increase in the translation machinery capacity required when algal cells are exposed to HL.

Crosstalk between transcriptional and post-transcriptional regulations during HL acclimation. *Chlamydomonas* implements diverse regulation strategies and photoprotection mechanisms to avoid photo-oxidative damages. PSII photoinhibition occurs when the rate of PSII damage exceeds the rate of repair. The PSII damage and repair cycle is actually part of a dynamic machinery, acting on a principle similar to that of a circuit breaker, which in coordination with other regulatory processes operating on shorter time scales (such as photosynthetic control or qE), protects PSI from irreversible photo-damages (Krieger-Liszkay et al., 2000; Suorsa et al., 2012; Tikkanen et al., 2014; Chaux et al., 2015). Optimal functioning of photosynthesis under HL therefore relies on a fine poise between PSII degradation and synthesis, the latter depending on the translation capacity, which in bacterial systems greatly relies on the level of free ribosomes (Kim et al., 2020). In such a scenario, chloroplasts need to improve the translation capacity to increase the production of photosystem subunits, particularly D1, in response to HL to avoid severe photo-oxidative damages. CreRH22, together with other plastid-targeted DEAD-box RNA helicases and other factors such as RBF1, may contribute to finely tuning the translation capacity to the protein synthesis demand. Ribosomes represent important energy cost

and resource allocation for cells, and in bacteria, the number of ribosomes is the result of a trade-off in

the use of nutrients between protein synthesis capacity and production of metabolic enzymes (Kim et

al., 2020). In bacteria and yeasts, allocation of resources towards ribosome synthesis is tuned to

optimize growth (Chure and Cremer, 2022). Photoautotrophic organisms are likely to have similar

constrains, especially when it comes to tune ribosomal content devoted to photosynthetic proteins

when light is the sole source of energy given the strong turnover rate of photosynthetic proteins (Li et

al., 2018). Optimizing the number of chloroplast ribosomes to available light resources is most likely

to be critical to maintain photosynthesis functioning and generate the required energy to optimize

fitness during adverse conditions of light fluctuations, and the DEAD-box RNA helicase CreRH22

appears to play a prominent role in such a regulation (Fig. 12).

Chloroplasts are semi-autonomous organelles of prokaryotic origin housing photosynthesis in

plant and algal cells. Most of the genes initially present in the endosymbiotic ancestor of chloroplasts

have been relocated to the nuclear genome, leading to a situation in which multi-subunit

photosynthetic complexes and ribosomes are partly encoded by nuclear and plastid genes. Therefore, a

coordinated control of gene expression and of protein synthesis between these two compartments is

needed to ensure a well-orchestrated assembly of these complexes and an optimal functioning of

photosynthesis (Stern et al., 2010). The expression of plastid genes is tightly controlled by complex

regulatory mechanisms involving more than hundred nucleus-encoded proteins (Stern et al., 2010),

and plastid-targeted DEAD-box RNA helicase most likely contribute to such a control.

An intriguing question relates to the origin of the strong changes in transcriptional regulation

that occur during light acclimation in hf_0. The expression of nuclear genes depends on a retrograde

signaling originating from the chloroplast, the origin and nature of which being a matter of intense

debate. It has been alternatively proposed that the redox state of the PQ pool or the redox state of PSI

acceptor side may contribute (Escoubas et al., 1995; Allen and Pfannschmidt, 2000; Pfalz et al., 2012;

Dietzel et al., 2015). In Arabidopsis leaves, the redox state of compounds on the reducing side of PSI

was considered to be more important than PQ-mediated redox signals in controlling the expression of

nuclear genes (Piippo et al., 2006). The redox status of the PQ pool can be assessed from chlorophyll

fluorescence measurements in certain conditions through the use of the parameter 1-qL (Kramer et al.,

2004). However, the high chlorophyll fluorescence of the hf_0 mutant, which results from the presence

of non-connected antennae, is not consistent with the "lake model" hypothesis that considers fully

connected units (Kramer et al., 2004), thus preventing the use of this parameter in this context. Since

phosphorylation of LHCII antennae, which depends on the reduction state of the PQ pool (Depege et

al., 2003), is not increased in hf_0 (Fig. 7. E), it seems likely that the redox state of the PQ pool is not

increased in hf_0. In fact, a lower redox state of the PQ pool is likely to occur in hf_0 since PSII is more

strongly inhibited than PSI under HL (Supplemental Fig. S7B). In this context, we propose that the

strong impairment of the transcriptome response observed in hf_0 under HL would be rather due to

limited over-reduction of PSI acceptors under HL resulting from a lower synthesis of photosystems.

Among the 32 Chlamydomonas DEAD-box RNA helicases listed in the Supplemental Fig. S3,

11 belong to a gene cluster (cluster 1) containing genes characterized by a strong induction during the

first hour of illumination followed by a sharp decrease (Zones et al., 2015), and four out of these 11

gene products were recently identified as physically interacting with plastid ribosomes (Westrich et

al., 2021). It seems therefore likely that, in addition of CreRH22, several other plastid-targeted DEAD-

box RNA helicases may participate, via the nuclear control of the plastid protein synthesis, to the

coordination between nuclear and plastid compartments, which is critical for a correct assemblage of

chloroplast complexes, particularly in changing light conditions. Recently, many DEAD-box RNA

helicases have been identified and partially characterized in crop plant genomes, including rice (Lu et

al., 2020), wheat (Ru et al., 2021), *Medicago truncatula* (Cheng et al., 2021) and *Brassica rapa*

(Nawaz et al., 2021), and plastid-targeted DEAD-box RNA helicases have been proposed to play a

role in the response to abiotic stress such a drought, salinity or cold (Nawaz and Kang, 2017). As light

is as a key parameter that potentiates the effects of abiotic stress on photosynthesis (Schwenkert, in

press), plastid-targeted DEAD-box RNA helicases may represent promising targets towards improving

the productivity of crops in stress conditions.

MATERIALS AND METHODS

Strains and culture conditions. *C. reinhardtii* strains (CC124 called wild-type, the hf_0 mutant and

complemented strains) were grown photo-autotrophically on a minimal medium (Harris, 2009). Batch

cultures were grown on a rotary shaker in Erlenmeyer flasks (100 mL) placed in a thermo-regulated

(25°C) incubator (Multitron, Infors) under continuous illumination (40 or 240 µmol photons m^{-2} s^{-1}) in

the presence of $2\%CO_2$ enriched air. For continuous photoautotrophic growth experiment, cells were

cultivated in four autoclavable 1L PBRs (BIOSTAT® Aplus, Sartorius Stedim Biotech) equipped with

a biomass probe (Excell probe, Exner, measuring O.D at 880 nm with a 2 cm light path) and operated

as turbidostats (Dang et al., 2014). Light was supplied by eight fluorescent tubes (Osram Dulux L

18W) disposed radially around the PBR to reach light intensities (measured at the surface of the PBR).

Growth rates (d^{-1}) were calculated by dividing the daily dilution volume by the PBR volume (1L) and

biomass productivity from dry weight measurements.

Mutant screening, isolation and molecular characterization. The *C. reinhardtii* strain CC124 (mt^-

nit1 nit2) was used as a genetic background to generate an insertion mutant library as described

previously (Tolleter et al., 2011). The pSL-X plasmid harboring the paromomycin resistance cassette

AphVIII was linearized by *KpnI* and used for nuclear transformation by electroporation. Around

12,000 random insertion mutants were isolated on selective medium and mutant phenotypes were

analyzed by chlorophyll fluorescence using a homemade imaging device. The hf_0 mutants showed a

high F_0 chlorophyll fluorescence level. The insertion site of the paromomycin cassette was identified

on genomic DNA upon phenol/chloroform extraction (Tolleter et al., 2011). First, Genome walker

protocol was adapted as follows from Universal GenomeWalker[TM] kit (Clontech). 250ng of gDNA

were digested overnight by 80U of *Afe*I (New England Biolabs). The restriction enzyme was

eliminated by precipitation with 1/10 (v/v) of ammonium acetate 2.5M (pH 5.6) and 1V of cold EtOH.

Digested DNA was dephosphorylated for 1h by 10U of antartic phosphatase (New England Biolabs).

DNA adaptator was ligated overnight with 2,000 u of T4 DNA ligase (New England Biolabs). Two

PCRs were carried out by using nested primers designed on the adaptator and on the paromomycin

resistance gene using the Advantage® GC genomic LA polymerase mix (Clontech). PCR products

were subcloned for sequencing and blasted on the *C. reinhardtii* genome, thus leading to the

identification of the Cre03.g166650 gene (thereafter called *CreRH22*).

Southern blot analysis. Genomic DNA (8μg) was digested by *NcoI* (New England Biolabs) overnight

and loaded on 0.8% agarose gel. After migration, gel was depurinated in 0.25M HCl for 10 min,

denatured in 0.5M NaOH/1.5M NaCl, and neutralized for 30 min in 0.5M Tris-HCl (pH 7.5), 1.5M

NaCl. The DNA was transferred overnight onto a nylon membrane using a 10 x SSC buffer, and was

fixed onto the membrane by a UV treatment (1,200W, UV Crosslinker, UVP Laboratory Product,

UK). Paromomycin probe was labeled with dUTP-DIG during PCR reaction (PCR DIG probe

synthesis kit, Roche; see table primers list). Membrane was pre-hybridized 1h at 50°C in a DIG eazy

Hyb (Roche) and then hybridized overnight at 50°C in same buffer using a denatured probe. Two

washing steps were realized, first using a 0.2 x SSC buffer containing 0.1% SDS (5 min at RT), and

then a 1 x SSC buffer containing 0.1% SDS (20 min at 65°C). Detection was done by Western Blot

with anti-digoxigenin antibody, coupled with alkaline phosphatase. We followed Roche procedure: 5

min in washing buffer, 30 min in blocking buffer, hybridization during 1h (1/10e4 anti-DIG in

blocking solution), two washing step of 15 min. Membrane was then equilibrated with detection buffer

(pH 9.5) before to add chemiluminescent substrate CSPD® (disodium 3-(4-methoxyspiro {1,2-

dioxetane-3,2'-(5'-chloro)tricyclo $[3.3.1.1^{3,7}]$decan}-4-yl) phenyl phosphate) (Roche). The chemio-

luminescent signal was detected by using a G-Box (Syngene).

Mutant complementation. Complementation of the *hf$_0$* mutant was first attempted by nuclear

transformation by cloning the genomic *CreRH22*. The CreRH22 gene was amplified by Phusion®

High-Fidelity PCR Master Mix with CDS helicase primers (Supplemental Fig. S10), subcloned into

the topoXL vector, and transferred into the pSL vector carrying the hygromycin resistance cassette

using *EcoRV/SpeI* sites (Berthold et al., 2002), and the CreRH22 gene under the control of *psa*D

promoter and terminator. The vector was linearized by *KpnI* and transformation was performed as

previously described (Tolleter et al., 2011). Transformants were selected on hygromycin (7.5μg mL^{-1})

and further analyzed for chlorophyll fluorescence properties. Complementation was also attempted by

means of chloroplast transformation. A synthetic *CreRH22* gene optimized for the *Chlamydomonas*

chloroplast codon usage (Supplemental Fig. S5) was cloned into the pLM21 vector under the control

of the *psa*A promoter using *BglII* restriction sites (Michelet et al., 2011; Tibiletti et al., 2016). Cells

were grown at 25°C under continuous light (100 µmol photons m² s^{-1}) in liquid Tris-Acetate-
Phosphate (TAP) medium until a density of 4.10^6 cells mL^{-1} and then spread onto agar plates.
Bombardment was realized at 7 bars with 10µg DNA of the PLM21 vector bound to gold carrier
particles (Seashell Technology kit). Transformants were selected on a medium containing 100 µg mL^{-1}
spectinomycin, and replated on increasing spectinomycin concentrations (up to 500 µg mL^{-1}) until
homoplasmy was reached, as controlled by PCR (Baltz et al., 2014).

RNA extraction and reverse transcription. After harvesting cells by centrifugation, cell pellets were
frozen in liquid nitrogen and resuspended in 0.5 mL of RNA reagent (Life Technologies™) to isolate
total RNAs. Cells were broken (two times for 10s at 5500 rpm, 4°C) using ceramic beads in a 2 mL
tube using a Precellys24® homogenizer (Bertin technologies, France). 100µL of NaCl 3M and 300 µL
of chloroform were added sequentially. After centrifugation (13,000g, 4°C) the upper aqueous phase
was mixed with isopropanol in order to precipitate nucleic acids. After washing with 70% EtOH RNA
samples were air dried. A DNAse treatment was done with TURBO® DNAse (Ambion®) followed by
purification with NucleoSpin® RNA clean-up (Macherey Nagel). Reverse transcription was
performed on 1µg of the total RNA using the SuperScript® vilo™ cDNA synthesis kit (Life
Technologies™). Expression of *CreRH22* was monitored by PCR on cDNA using TaKaRa LA Taq®
DNA Polymerase. *CBLP2* (RACK1 Cre06.g278222.t1.1) was used as a control (Supplemental Fig.
S10).

Northern blot and real time quantitative PCR. For Norther blots, 5µg of total RNA were separated
on 1.4% agarose denatured gel (2% formaldehyde), transferred on nylon membrane and hybridized
with DIG-labelled probes obtained by PCR DIG Probe Synthesis Kit (Roche) with specific primers
(see table primers list). The membrane was pre-hybridized and hybridized at 50°C in DIG eazy Hyb
(Roche). Two double washing steps were realized using 0.1% SDS, 0.2x SSC, during 5 min at room
temperature and 20 min at 65°C. Detection was performed using an anti-digoxigenin antibody,
coupled with alkaline phosphatase. RT-qPCR reactions were realized on LightCycler® 480 (Roche)
with MESA FAST qPCR MasterMix Plus for SYBR® Assay No ROX (Eurogentec). The cycling
conditions used were as follows: 95°C for 10 min, then 45 cycles at 95°C for 10 s, 60°C for 15 s, and
72°C for 10 s. The relative transcript ratio were calculated based on the $2^{-\Delta\Delta CT}$ method (Livak and
Schmittgen, 2001), with average cycle threshold obtained based on triplicate measurements.
Ribosomal protein gene *rpl36* was used as chloroplast reference gene.

RNA-seq experiments. 10 mL of PBR turbidostat cultures of both wild-type and *hf$_0$* (with a cell
density of 3 x 10^6 cells mL^{-1}) were harvested and centrifuged (4000 rpm, 1 min, 4°C). Samples of two
replicate cultures (two independent PBR experiments) were taken after 24 h cultivation under 30 µmol
photons m^{-2} s^{-1} and then 2h and 24h after a switch to 240 µmol photons m^{-2} s^{-1}. Total RNAs of all
samples (12 samples in total) were extracted as described above. Each library was sequenced using an
Illumina Sequencer generating 23.3-25 10^6 50 nt single end reads for each sample. The sequencing

reactions and RNA-Seq data analysis were performed at Beijing Genome Institute, Hong Kong. For

transcriptomic analysis, reads were aligned onto the *C. reinhardtii* genome assembly version v5 using

SOAPaligner/SOAP2 (Li et al., 2008; Li et al., 2009). No more than two mismatches were allowed in

the alignment. For each sample 17.2-19.8 10^6 reads were mapped to single copy sequences on the *C.*

reinhardtii genome. NoiSeq (Tarazona et al., 2011) was applied to screen Differentially Expressed

Genes (DEGs) between two groups. We used a False Discovery Rate (FDR) ≤ 0.001 and the absolute

value of ≥ 1 as the threshold to judge the significance of gene expression difference. These DEGs were

submitted to functional analysis using the MapMan 3.5.1R2 (Thimm et al., 2004) software package to

attribute DEGs to biological pathways.

Starch and oil measurements. Starch extraction was performed using the method described by Klein

and Betz (1978). One ml of culture was centrifuged (14,000 rpm, 2 min), re-suspended in 1ml of

methanol and centrifuged again. Chlorophyll concentration was estimated directly on methanol

supernatants by absorbance at 663 and 645 nm. Pellets were heated for 15 min at 120°C for starch

solubilisation after the addition of 400μL water. 200μL of amyloglucosidase solution Starch Assay

Reagent (Sigma-Aldrich, Saint-Louis, MO, USA) were then added and incubated at 55°C for 1h.

Glucose was subsequently assayed using an automated sugar analyser (Ysi model 2700 select, Yellow

springs, OH, USA). Lipid extraction and TAG analysis were performed as previously described (Siaut

et al., 2011).

Immunoblot analysis. Upon harvesting, 20×10^6 cells were harvested and frozen from PBR cultures.

Totals proteins were extracted with SDS 0.2% and acetone 80%. The pellets were treated with a

denaturing buffer from Thermofisher (LDS Sample Buffer supplied with DTT) and heated during 20

min at 70°C. Gel electrophoresis was performed using a NuPAGE Novex System (10% Bis-Tris gel,

MES running buffer) and proteins transfer using iBlot® 2 Dry Blotting System. Depending on the

sensitivity of the antibody, detection was based on chemiluminescence using HRP conjugated

antibody, chemiluminescent substrate and a GBOX imaging system (Syngene), or fluorescent

detection using a Fluorescent Alexa Fluor 680 dye conjugated antibody and Odyssey imaging system

(LiCor). Antibodies against PsbA, PsbD, PsbC, PsaC, PSAD, LHCB2, AtpB, and PetA were

purchased from Agrisera and the phosphothreonin antibody was purchased from InvitroGen. PsaA and

PSAF antibodies were kindly supplied by Michel Goldschmidt-Clermont (University of Geneva,

Switzerland), and L30 and S21 antibodies by William Zerges (Concordia University, Montreal,

Canada). Immuno-quantification was performed in the linear range of each antibody response. Protein

amounts loaded for immunoblots were 8μg (PsaA, PsaC, PSAD, PSAF, PsbC), 4μg (PetA, L30),

1.6μg (AtpB, PsbA), and 0.8μg (PsbD). Densitometries were analyzed by using the GeneTools

analysis software (Syngene) for chemiluminescence measurements or the Odyssey analysis software

(LiCor) for fluorescence measurements. Quantifications were based on biological triplicates and

expressed as relative values of the average wild-type protein level at LL.

Blue native PAGE and two-dimensional analysis of thylakoid proteins. Thylakoid membranes were purified under dim light at 4°C by using a protocol adapted from (Chua and Bennoun, 1975). Cells were harvested during exponential growth (about 5×10^7 cells) by centrifugation (5,000 rpm, 5 min, 4°C). Cell pellets were resuspended in 5mL of buffer A containing 5 mM HEPES-KOH (pH 7.5), 0.3M sucrose, 10 mM EDTA, Protease Inhibitor Cocktail (SigmaAldrich). Cells were disrupted using a French Press (Aminco) and centrifuged (12,000g, 10 min, 4°C). Cell pellets were resuspended in buffer A containing 1.8 M sucrose, and blended with a potter homogenizer. The homogenate was covered with 4 mL of buffer A without EDTA and containing 1.3M sucrose and 4 mL of buffer A containing 0.5 M sucrose. Upon ultracentrifugation (274,000g for 1 h), membranes were collected in the 1.3M sucrose layer and at the interface with the 0.5 M sucrose layer. Purified membranes were then diluted in the HEPES buffer and centrifuged (20,000 rpm for 20 min). Thylakoid membranes were then resuspended in s sample buffer (NativePAGE, ThermoFischer) at 1mg chlorophyll. mL^{-1}. Solubilization was carried out in the dark (5 min at 4°C) by adding an equal volume of 2% dodecyl maltoside. Insoluble material was removed by centrifugation (12,000 g, 20 min, 4°C) and one-tenth volume of Native PAGE 5% G 250 Sample Additive was added. Blue native PAGE was performed by using 3-12% Bis-Tris gels (NativePAGE, ThermoFischer). For two-dimensional PAGE, strips were excised and conserved at -80°C until use. After solubilization in LDS Buffer (Thermofisher) containing 5.5 M urea, two-dimensional PAGE was performed in 15% (w/v) polyacrylamide gels containing 5.5 M urea, and proteins revealed by silver staining.

Mass spectrometry-based proteomic analysis. For proteomic analysis, 2D gels were stained by Coomassie brilliant blue and proteins were digested in-gel using trypsin (modified, sequencing purity, Promega), as previously described (Casabona et al., 2013). The resulting peptides were analyzed by online nanoliquid chromatography coupled to MS/MS (Ultimate 3000 and LTQ-Orbitrap Velos Pro, Thermo Fisher Scientific) using a 30 min gradient. For this purpose, the peptides were sampled on a precolumn (300 μm x 5 mm PepMap C18, Thermo Scientific) and separated in a 75 μm x 250 mm C18 column (PepMap C18, 3 μm, Thermo Fisher Scientific). The MS and MS/MS data were acquired by Xcalibur (Thermo Fisher Scientific). Peptides and proteins were identified by Mascot (version 2.8.0, Matrix Science) through concomitant searches against the Uniprot database (*Chlamydomonas reinhardtii* taxonomy, January 2022 download), and a homemade classical database containing the sequences of classical contaminant proteins found in proteomic analyses (human keratins, trypsin, etc.). Trypsin/P was chosen as the enzyme and two missed cleavages were allowed. Precursor and fragment mass error tolerances were set respectively at 10 ppm and 0.6 Da. Peptide modifications allowed during the search were: Carbamidomethyl (C, fixed), Acetyl (Protein N-term, variable) and Oxidation (M, variable). The Proline software (Bouyssie et al., 2020) was used for the compilation, grouping, and filtering of the results (conservation of rank 1 peptides, peptide length ≥ 6 amino acids, false discovery rate of peptide-spectrum-match identifications < 1% (Coute et al., 2020), and

minimum of one specific peptide per identified protein group). Proteins from the contaminant database

were discarded from the final list of identified proteins. Proline was then used to perform MS1-based

label free quantification of the identified protein groups based on razor and specific peptides. The

relative abundances of the different proteins were evaluated through calculation of their intensity-

based absolute quantification (iBAQ, (Schwanhausser et al., 2011)) values.

Radiolabeling of total cellular proteins. Total cellular proteins from *C. reinhardtii* were labeled with

^{35}S as described previously (Ozawa et al., 2010) with some minor modifications. Cells grown photo-

autotrophically in batch cultures under LL (20 µmol photons m^{-2} s^{-1}) in the presence of 2% CO_2

enriched air until reaching a cell density of 2 x10^6 cells mL^{-1}, then switched to HL (300 µmol photons

m^{-2} s^{-1}) for 6 h, and resuspended (at 25 µg chlorophyll. mL^{-1}) in a sulfur free HSM media to induce

sulfur starvation for 1 hour. During starvation light conditions were same as they were grown

previously. Total cellular proteins were labeled by adding 5 µCi mL^{-1} of [^{35}S] Na_2SO_4 (American

Radiolabeled Chemicals) for 10 min. Labeling was stopped by adding 10 mM cold Na_2SO_4,

chloramphenicol (100 µg mL^{-1}) and cycloheximide (10 µg mL^{-1}). Thylakoid membranes were purified

by discontinuous sucrose density gradient ultracentrifugation, and radiolabeled polypeptides were

separated by (12-25%) SDS-PAGE (Nellaepalli et al., 2018). After electrophoresis, the gel was dried

and exposed to an imaging plate for several days and labeling was detected by using a fluorescence

image analyzer (FLA7000, Fujifilm).

Chlorophyll fluorescence and ECS measurements. Chlorophyll fluorescence measurements were

carried out at room temperature on whole cells using a Dual Pam-100 (http://www.walz.com).

Samples were taken from batch or PBR cultures and placed into the PAM cuvette under constant

stirring. The maximal PSII yield (Φ_{PSII}) was measured as the Fv/Fm ratio at a single saturation flash

illumination (15,000 µmol photons m^{-2} s^{-1}, 200 ms duration) upon 15-30 min dark adaptation. For ETR

(Electron Transport Rate) measurement, actinic light was increased step-wise from 3 to 800 µmol

photons m^{-2} s^{-1}. After 2 min stabilization at each light intensity, a saturating flash (15,000 µmol

photons m^{-2} s^{-1}, 200 ms duration) was applied, the PSII operating yield measured as (Fm'-Fs)/Fm' and

apparent ETR calculated by multiplying the PSII yield by the actinic light intensity. 77K chlorophyll

fluorescence was measured using a SAFAS Xenius optical fiber fluorescence spectrophotometer as

described previously (Tolleter et al., 2011; Dang et al., 2014). PSII/PSI ratios were determined from

electrochromic shifts of carotenoids (ECS) absorbance changes measurements at 520 nm using a

JTS10 spectrophotometer by measuring signals induced by a saturating single turnover flash in the

absence or in the presence of PSII inhibitors hydroxylamine and DCMU at final concentrations of 1

mM and 10 µM, respectively (Tolleter et al., 2011; Dang et al., 2014).

Accession numbers. Sequence data from this article can be found at Phytozome under the following
accession number: *RH22* (also named *HEL15* or *CGLD3*) as Cre03.g166650. The RNA-seq data
generated in this study were deposited to the SRA database at the National Center for
Biotechnological Information under the accession number PRJNA834659.
599
600
601
602

SUPPLEMENTAL DATA

Supplemental Figure S1. Southern blot analysis of wild-type and hf_0 *Chlamydomonas* mutant strain
indicates a single insertion of the paromomycin resistance cassette.

Supplemental Figure S2. Phylogenetic analysis of DEAD-box RNA helicases proteins.

Supplemental Figure S3. DEAD-box RNA helicase identified in the *Chlamydomonas* genome.

Supplemental Figure S4. Complementation of the hf_0 mutant by nuclear and chloroplast
transformation.

Supplemental Figure S5. Nucleotide sequence of the codon optimized synthetic CrRH22 used for
chloroplast transformation

Supplemental Figure S6. Accumulation of chloroplast transcripts as analyzed by RT-qPCR in the
WT and in the mutant hf_0 at two light intensities.

Supplemental Figure S7. Maximum PSII yields and PSII/PSI ratios determined in WT and hf_0 cells
grown under LL or HL.

Supplemental Figure S8. Blue native PAGE of thylakoid complexes from WT and hf0 grown under
HL in the absence or presence of chloramphenicol.

Supplemental Figure S9. Light-induced expression of CreRH22, HSP70B and HSP90C genes.

Supplemental Figure S10. List of PCR primers.

Supplemental Data Set S1. MS-based proteomic characterization of thylakoid proteins from hf_0 cells
exposed to HL and present in the two-dimensional blue native-PAGE region of interest.

Supplemental Data Set S2. RNA-seq results from hf_0 and wild-type control cells cultivated in low
light (30 µmol photons m^{-2} s^{-1}) and then exposed to high light (240 µmol photons m^{-2} s^{-1}) for 2h and
24h.
625

ACKNOWLEDGEMENTS
This work was supported by the French ANR (Agence Nationale pour la Recherche) projects
ALGOH2, OTOLHYD and by the A*MIDEX project (n° ANR-11-IDEX-0001-02) funded by the «
Investissements d'Avenir » French Government program. Experimental support was provided by the
HélioBiotec platform (funded by the European Regional Development Fund, the Région Provence
Alpes Côte d'Azur, the French Ministry of Research, and the "Commissariat à l'Energie Atomique et
aux Energies Alternatives"). Dr. Mohamed Barakat (CNRS, CEA Cadarache) is acknowledged for
managing the deposit of RNAseq files. The proteomic experiments were partially supported by

638   **AUTHOR CONTRIBUTIONS**

639   E.B.D.T. and G.P. designed the study. E.B.D.T., S.N., P.A., A.B., F.C., S.C., V.E., M.H., B.G.,

640   M.S.R., I.T., performed the experiments. E.B. performed phylogenetic and transcriptomic analysis,

641   S.B. performed proteomic analysis. E.B.D.T., S.N., S.C., S.D., Y.C., Y.T. and G.P. analysed the data.

642   E.B.D.T., S.N., Y.L.B and G.P. wrote the manuscript.

LEGENDS OF FIGURES

Figure 1. Isolation and molecular characterization of *hf₀*, a *C. reinhardtii* high chlorophyll fluorescence mutant. (A) The *hf₀* mutant was isolated from the screening of an insertion library based on the analysis of chlorophyll fluorescence transients (after recording F_0 fluorescence in the dark, a saturating flash at 5s was followed by 240 µmol photons m^{-2} s^{-1} actinic red light at 14 s). **(B)** The paromomycin resistance cassette (*AphVIII* gene) is integrated in exon 7 of the Cre03.g166650 locus encoding a gene annotated as a DEAD-box RNA helicase, renamed here *CreRH22* based on a phylogenetic analysis (Supplemental Fig. 2). The *CreRH22* gene is composed of 9 exons (blue boxes) and 8 introns (black lines). Orange arrows show primers used for RT-qPCR shown in (C). The CreRH22 protein shows typical features of DEAD-box RNA helicases, including a Q motif, an ATP binding domain containing a DEAD motif, and an ATPase RNA unwinding helicase domain. **(C)** RT-qPCR showing the absence of the *CreRH22* gene transcript in *hf₀*. *CBLP* (alias RACK1 Cre06.g27822) is used as loading control.

Figure 2. Effect of light on photosynthetic properties of the *hf₀* mutant. Wild-type (WT), *hf₀* and two complemented lines (*hf₀::HF₀-c7* and *hf₀::HF₀-g6*) were grown photo-autotrophically in 3% CO_2-enriched air under HL (240 µmol photons m^{-2} s^{-1} **A, C**) or LL (15 µmol photons m^{-2} s^{-1} **B, D**). Representative chlorophyll fluorescence measurements during a dark/light (100 µmol photons m^{-2} s^{-1}) transient for HL **(A)** or LL **(B)** grown cells. Black and white boxes indicate dark and light periods, respectively. **(C, D)** PSII yields determined from chlorophyll fluorescence measurements as F_v/F_m in the dark after 15 min dark adaptation or $(F_{m'}-F_s)/F_{m'}$ in the light in HL **(C)** or LL **(D)** grown cells.

Figure 3. Ribosomal RNAs processing is altered in the *hf₀* mutant. (A) Scheme showing the organization of the *C. reinhardtii* chloroplast ribosomal RNA (rRNA) operon and location of the different probes (black bars) used for Northern blot analysis. **(B)** rRNAs levels analyzed by Northern blot in WT and *hf₀* cells grown photo-autotrophically under a light intensity of 60 µmol photons m^{-2} s^{-1}. 5µg of total RNA were loaded on agarose gels, blotted on nylon membrane and hybridized with DIG-labeled probes. For loading controls (lower blue boxes), nylon membranes were stained with methylene blue after hybridization.

Figure 4. Accumulation of major chloroplast proteins in WT and *hf₀* upon HL exposure determined by immuno-analysis. Cells were grown photo-autotrophically in photobioreactors in 3% CO_2-enriched air at LL (7.5 µmol photons m^{-2} s^{-1}) and exposed to LL (60 µmol photons m^{-2} s^{-1}), for 6h or 26h, respectively corresponding to light intensities of about 20 and 200 µmol photons m^{-2} s^{-1} in flasks (Dang et al., 2014). Protein quantification based on immuno-analysis performed in the linearity range of the antibody detection. Shown are mean values (+/-SD, n=3) normalized on the WT protein content measured at t_0.

Figure 5. Pulse labeling experiments of thylakoid proteins show a strong decrease of D1 protein

synthesis under HL in hf_0. Cells were grown photo-autotrophically at LL up to a cell density of 2.

10^6 cells mL^{-1} and then switched to HL for 6 h to the cell density of 6. 10^6 cells mL^{-1}, and then labeled

for 10 min with ^{35}S (Na$_2$SO$_4$). Upon extraction and SDS-PAGE separation of total cellular proteins,

labeled proteins were visualized by autoradiography. Autoradiograms show labeled thylakoid

membrane proteins of WT, hf_0 and two complemented lines, $hf_0::HF_0$-c7 (complemented line obtained

by chloroplast transformation) and $hf_0::HF_0$-g6 (complemented line obtained by nuclear

transformation). The gel was stained with Coomassie brilliant blue before exposing to the imaging

plate.

Figure 6. Effect of HL and chloramphenicol addition on photosynthetic activity of the hf_0

mutant. Wild-type (WT), hf_0 and two complemented lines ($hf_0::HF_0$-c7 and $hf_0::HF_0$-g6) were grown

in batch cultures under LL (15 µmol photons m^{-2} s^{-1}). At t_0 cell cultures were switched to HL (240

µmol photons m^{-2} s^{-1}) in the absence **(A, C)** or in the presence **(B, D)** of 50 mg. L^{-1} chloramphenicol

(chlΦ) added 1h before the HL switch. At different time points after the light switch, maximal PSII

yields were measured as the F_v/F_m ratio after 15 min dark adaptation **(A, B)** and PSII operating yields

were measured as $(F_{m'}-F_s)/F_{m'}$ following a 3 min light period (430 µmol photons m^{-2} s^{-1}). Shown are

means ± SD (n=3).

Figure 7. Low temperature chlorophyll fluorescence and chlorophyll-protein complexes are

affected in hf_0, but state transitions are not. WT and hf_0 cells were grown photo-autotrophically in

photobioreactors in 3% CO$_2$-enriched air at LL (7.5 µmol photons m^{-2} s^{-1}) and then switched to HL (60

µmol photons m^{-2} s^{-1}), respectively corresponding to light intensities of about 20 and 200 µmol

photons m^{-2} s^{-1} in flasks (Dang et al. 2015). **(A, B)** 77K chlorophyll fluorescence emission spectra of

cells grown under LL **(A)** or 26 h after HL switch **(B)**. **(C)** 77K chlorophyll fluorescence emission

ratio E_{712nm}/E_{686nm}, and **(D)** F_0/F_m chlorophyll fluorescence at different time points after HL switch. **(E)**

Immunodetection of phosphorylated LHCII using an anti-phospho-threonine antibody. Arrows

indicate protein bands with lower abundance in hf_0. **(F)** Blue-native PAGE of thylakoid complexes

from WT and hf_0 grown under LL or upon 26h of HL exposure. Two left arrows indicate green bands

with lower abundance in hf_0 and the right arrow indicates a green band of higher abundance in hf_0.

Figure 8. Two-dimensional PAGE of thylakoid proteins from WT and hf_0 cells upon HL

exposure. WT and hf_0 cells were grown photo-autotrophically in batch cultures under LL (40 µmol

photons m^{-2} s^{-1}) and then exposed to HL (240 µmol photons m^{-2} s^{-1}) for 26h. Thylakoid proteins were

solubilized with 1% dodecyl-maltoside, separated by blue native PAGE in a first dimension and by

SDS-PAGE in a second dimension. The red dotted ellipse highlights proteins associated to high MW

PSI-LHCI supercomplexes in the WT **(A)** that migrates as low MW complexes attributed to LHCI

oligomers in hf_0 (red arrow in **B**). The gel strip in the 20-35kDa MW range shown as a red box **(B)** was

taken for proteomic analysis by mass spectrometry (Supplemental Table 1).

Figure 9. RNAseq analysis of light-induced gene expression in wild-type and hf_0 C. reinhardtii

cells. Cells were cultivated in PBRs operated as turbidostats in 3% CO_2-enriched air under LL (7.5

µmol photons m^{-2} s^{-1}) for 24h and then exposed to HL (60 µmol photons m^{-2} s^{-1}). Samples for RNAseq

analysis were taken from two biological replicates under LL before the HL switch, and the 2h and 24h

after the switch. (A) Shown are numbers of up and down Differentially Expressed Genes (DEGs) with

a log 2(Fold Change) > 1 and <-1, using a False Discovery Rate (FDR) ≤ 0.001. (B, C) Functional

analysis was performed DEGs from WT (A) and hf_0 (B) using the MapMan 3.5.1R2 package (Thimm

et al., 2004).

Figure 10. Expression patterns of genes encoding chloroplast-, mitochondria-, and cytosol-

targeted ribosomal proteins. (A to C) Heat maps depicting relative expression levels of ribosomal

protein genes and translational regulators targeted to cytosol **(A)**, chloroplast **(B)**, and mitochondria

(C). The maximum expression level for each gene is set to 1. ***** Genes without significant differential

expression (|log2FC|<2 as defined in Methods). **(D to F)** Boxplot of normalized expression values of

ribosomal protein genes targeted to cytosol **(D)**, chloroplast **(E)**, and mitochondria **(F)**.

Figure 11. Biomass productivity and concentrations of intracellular starch and TAGs are

reduced in hf_0. Wild-type and hf_0 cells were grown in PBRs operated as turbidostats in 3% CO_2-

enriched air and maintained at a constant biomass concentration (≈ 1.5 x 10^6 cells mL^{-1}) by continuous

addition of fresh medium. **(A)** Specific growth rates were determined from dilution rates measured at

different light intensities (ranging from 7.5 up to 300 µmol photons m^{-2} s^{-1}) following at least 24h

stabilization. Shown are means ± SD (n=4 for 60 and 200 µmol photons m^{-2} s^{-1}, n=2 for 7.5, 11, and

300 µmol photons m^{-2} s^{-1}). **(B)** Chlorophyll content, **(C)** chlorophyll a/b ratio, **(D)** intracellular starch,

and **(E)** intracellular TAG contents were determined on cell samples following at least 24h

stabilization at the specified light intensity (shown are means ± SD, n=3).

Figure 12. Hypothetical scheme proposed to explain the hf_0 mutant phenotype. The hf_0 mutant is

defected in a plastidial DEAD-box RNA helicase (CreRH22) involved in the processing of plastidial

rRNA operon and ribosome biogenesis. **(A)** Under LL the turnover of chloroplast proteins is low and

the translation capacity is sufficient to maintain photosynthesis at similar levels in the wild-type (WT)

and hf_0. **(B, C)** Under HL exposure, the amount of chloroplast ribosomes increases in the WT but

remains constant in the mutant. In these conditions, a much higher rate of protein synthesis is needed

due to the permanent degradation and synthesis of the D1 protein and to the biosynthesis of additional

PSII subunits and to a lesser extent PSI subunits. In these conditions, the hf_0 mutant is unable to

synthesize photosynthetic proteins at a sufficient rate, thus resulting in an inhibition of photosynthesis.

LITERATURE CITED

Adams, W.W., III, Muller, O., Cohu, C.M., and Demmig-Adams, B. (2013). May photoinhibition
be a consequence, rather than a cause, of limited plant productivity? Photosynth. Res. **117,** 31-44.

Allen, J.F., and Pfannschmidt, T. (2000). Balancing the two photosystems: photosynthetic electron
transfer governs transcription of reaction centre genes in chloroplasts. Philos. Trans. R. Soc. Lond.
B Biol. Sci. **355,** 1351-1357.

Allorent, G., Tokutsu, R., Roach, T., Peers, G., Cardol, P., Girard-Bascou, J., Seigneurin-Berny,
D., Petroutsos, D., Kuntz, M., Breyton, C., Franck, F., Wollman, F.A., Niyogi, K.K., Krieger-
Liszkay, A., Minagawa, J., and Finazzi, G. (2013). A dual strategy to cope with high light in
Chlamydomonas reinhardtii. Plant Cell **25,** 545-557.

Asakura, Y., Galarneau, E., Watkins, K.P., Barkan, A., and van Wijk, K.J. (2012). Chloroplast
RH3 DEAD box RNA helicases in maize and *Arabidopsis* function in splicing of specific group II
introns and affect chloroplast ribosome biogenesis. Plant Physiol. **159,** 961-974.

Baltz, A., Dang, K.V., Beyly, A., Auroy, P., Richaud, P., Cournac, L., and Peltier, G. (2014).
Plastidial expression of type II NAD(P)H dehydrogenase increases the reducing state of
plastoquinones and hydrogen photoproduction rate by the indirect pathway in *Chlamydomonas*
reinhardtii. Plant Physiol. **165,** 1344-1352.

Berthold, P., Schmitt, R., and Mages, W. (2002). An engineered *Streptomyces hygroscopicus aph 7″*
gene mediates dominant resistance against hygromycin B in *Chlamydomonas reinhardtii*. Protist
153, 401-412.

Bonente, G., Pippa, S., Castellano, S., Bassi, R., and Ballottari, M. (2012). Acclimation of
Chlamydomonas reinhardtii to different growth irradiances. J. Biol. Chem. **287,** 5833-5847.

Bouyssie, D., Hesse, A.M., Mouton-Barbosa, E., Rompais, M., Macron, C., Carapito, C., de
Peredo, A.G., Coute, Y., Dupierris, V., Burel, A., Menetrey, J.P., Kalaitzakis, A., Poisat, J.,
Romdhani, A., Burlet-Schiltz, O., Cianferani, S., Garin, J., and Bruley, C. (2020). Proline: an
efficient and user-friendly software suite for large-scale proteomics. Bioinformatics **36,** 3148-3155.

Casabona, M.G., Vandenbrouck, Y., Attree, I., and Coute, Y. (2013). Proteomic characterization
of *Pseudomonas aeruginosa* PAO1 inner membrane. Proteomics **13,** 2419-2423.

Chaux, F., Peltier, G., and Johnson, X. (2015). A security network in PSI photoprotection:
regulation of photosynthetic control, NPQ and O-2 photoreduction by cyclic electron flow. Front.
Plant Sci. **6**.

Cheng, J., Zhou, S., Yang, K., Yu, H., Chen, R., Zeng, L., Li, H., Wang, Y., and Song, J. (2021).
Identification of RNA helicases in *Medicago truncatula* and their expression patterns under abiotic
stress. Physiol. Mol. Biol. Plants **27,** 2283-2296.

Chi, W., He, B., Mao, J., Li, Q., Ma, J., Ji, D., Zou, M., and Zhang, L. (2012). The function of
RH22, a DEAD RNA helicase, in the biogenesis of the 50S ribosomal subunits of *Arabidopsis*
chloroplasts. Plant Physiol. **158,** 693-707.

Christopher, D.A., and Mullet, J.E. (1994). Separate photosensory pathways co-regulate blue
light/ultraviolet-A-activated *psb*D-*psb*C transcription and light-induced D2 and CP43 degradation
in barley (*Hordeum vulgare*) chloroplasts. Plant Physiol. **104,** 1119-1129.

Chua, N.H., and Bennoun, P. (1975). Thylakoid membrane polypeptides of *Chlamydomonas*
reinhardtii wild-type and mutant strains deficient in Photosystem 2 reaction center. Proc. Natl.
Acad. Sci. USA **72,** 2175-2179.

Chure, G., and Cremer, J. (2022). An optimal regulation of fluxes dictates microbial growth in and
out of steady-state. bioRxiv**,** 2022.2001.2027.477569.

Coute, Y., Bruley, C., and Burger, T. (2020). Beyond target-decoy competition: Stable validation of
peptide and protein identifications in mass spectrometry-based discovery proteomics. Anal. Chem.
92, 14898-14906.

Dang, K.V., Plet, J., Tolleter, D., Jokel, M., Cuine, S., Carrier, P., Auroy, P., Richaud, P.,
Johnson, X., Alric, J., Allahverdiyeva, Y., and Peltier, G. (2014). Combined increases in
mitochondrial cooperation and oxygen photoreduction compensate for deficiency in cyclic electron
flow in *Chlamydomonas reinhardtii*. Plant Cell **26,** 3036-3050.

Depege, N., Bellafiore, S., and Rochaix, J.D. (2003). Role of chloroplast protein kinase Stt7 in
LHCII phosphorylation and state transition in *Chlamydomonas*. Science **299,** 1572-1575.

Dietzel, L., Glaesser, C., Liebers, M., Hiekel, S., Courtois, F., Czarnecki, O., Schlicke, H., Yan,
Z., Boerner, T., Mayer, K., Grimm, B., and Pfannschmidt, T. (2015). Identification of early
nuclear target genes of plastidial redox signals that trigger the long-term response of Arabidopsis to
light quality shifts. Mol. Plant **8,** 1237-1252.

Erickson, E., Wakao, S., and Niyogi, K.K. (2015). Light stress and photoprotection in
Chlamydomonas reinhardtii. Plant J. **82,** 449-465.

Escoubas, J.M., Lomas, M., Laroche, J., and Falkowski, P.G. (1995). Light intensity regulation of
cab gene transcription is signaled by the redox state of the plastoquinone pool. Proc. Natl. Acad.
Sci. USA **92,** 10237-10241.

Fristedt, R., Scharff, L.B., Clarke, C.A., Wang, Q., Lin, C., Merchant, S.S., and Bock, R. (2014).
RBF1, a plant homolog of the bacterial ribosome-binding factor RbfA, acts in processing of the
chloroplast 16S ribosomal RNA. Plant Physiol. **164,** 201-215.

Girolomoni, L., Ferrante, P., Berteotti, S., Giuliano, G., Bassi, R., and Ballottari, M. (2017). The
function of LHCBM4/6/8 antenna proteins in *Chlamydomonas reinhardtii*. J. Exp. Bot. **68,** 628-
642.

Goold, H.D., Cuine, S., Legeret, B., Liang, Y., Brugiere, S., Auroy, P., Javot, H., Tardif, M.,
Jones, B., Beisson, F., Peltier, G., and Li-Beisson, Y. (2016). Saturating light induces sustained
accumulation of oil in plastidal lipid droplets in *Chlamydomonas reinhardtii*. Plant Physiol. **171,**
2406-2417.

Harris, E.H. (2009). The *Chlamydomonas* sourcebook. Second edition. (Academic Press).

Holloway, S.P., and Herrin, D.L. (1998). Processing of a composite large subunit rRNA: Studies
with *Chlamydomonas* mutants deficient in maturation of the 23S-like rRNA. Plant Cell **10,** 1193-
1206.

Jarvi, S., Suorsa, M., and Aro, E.-M. (2015). Photosystem II repair in plant chloroplasts -
Regulation, assisting proteins and shared components with photosystem II biogenesis. Biochim.
Biophys. Acta **1847,** 900-909.

Jarvis, P., and Lopez-Juez, E. (2013). Biogenesis and homeostasis of chloroplasts and other plastids.
Nat. Rev. Mol. Cell Biol. **14,** 787-802.

Karpowicz, S.J., Prochnik, S.E., Grossman, A.R., and Merchant, S.S. (2011). The GreenCut2
resource, a phylogenomically derived inventory of proteins specific to the plant lineage. J. Biol.
Chem. **286,** 21427-21439.

Kettunen, R., Pursiheimo, S., Rintamaki, E., VanWijk, K.J., and Aro, E.M. (1997).
Transcriptional and translational adjustments of *psbA* gene expression in mature chloroplasts
during photoinhibition and subsequent repair of photosystem II. Eur. J. Biochem. **247,** 441-448.

Kim, J., Darlington, A., Salvador, M., Utrilla, J., and Jimenez, J.I. (2020). Trade-offs between
gene expression, growth and phenotypic diversity in microbial populations. Curr. Opin. Biotechnol.
62, 29-37.

Kramer, D.M., Johnson, G., Kiirats, O., and Edwards, G.E. (2004). New fluorescence parameters
for the determination of Q(A) redox state and excitation energy fluxes. Photosynth. Res. **79,** 209-
218.

Krieger-Liszkay, A., Kienzler, K., and Johnson, G.N. (2000). Inhibition of electron transport at the
cytochrome b(6)f complex protects photosystem II from photoinhibition. FEBS Lett. **486,** 191-194.

Kujat, S.L., and Owttrim, G.W. (2000). Redox-regulated RNA helicase expression. Plant Physiol.
124, 703-713.

Li, L., Aro, E.-M., and Millar, A.H. (2018). Mechanisms of photodamage and protein turnover in
photoinhibition. Trends Plant Sci. **23,** 667-676.

Li, R.Q., Li, Y.R., Kristiansen, K., and Wang, J. (2008). SOAP: short oligonucleotide alignment
program. Bioinformatics **24,** 713-714.

Li, R.Q., Yu, C., Li, Y.R., Lam, T.W., Yiu, S.M., Kristiansen, K., and Wang, J. (2009). SOAP2:
an improved ultrafast tool for short read alignment. Bioinformatics **25,** 1966-1967.

Linder, P., and Jankowsky, E. (2011). From unwinding to clamping - the DEAD box RNA helicase
family. Nat. Rev. Mol. Cell Biol. **12,** 505-516.

Livak, K.J., and Schmittgen, T.D. (2001). Analysis of relative gene expression data using real-time
quantitative PCR and the 2(T)(-Delta Delta C) method **25,** 402-408.

Lu, C.-A., Huang, C.-K., Huang, W.-S., Huang, T.-S., Liu, H.-Y., and Chen, Y.-F. (2020). DEAD-
Box RNA helicase 42 plays a critical role in pre-mRNA splicing under cold stress. Plant Physiol.
182, 255-271.

Michelet, L., Lefebvre-Legendre, L., Burr, S.E., Rochaix, J.-D., and Goldschmidt-Clermont, M.
(2011). Enhanced chloroplast transgene expression in a nuclear mutant of *Chlamydomonas*. Plant
Biotechnol. J. **9,** 565-574.

Migur, A., Heyl, F., Fuss, J., Srikumar, A., Huettel, B., Steglich, C., Prakash, J.S.S., Reinhardt,
R., Backofen, R., Owttrim, G.W., and Hess, W.R. (2021). The temperature-regulated DEAD-
box RNA helicase CrhR interactome: autoregulation and photosynthesis-related transcripts. J. Exp.
Bot. **72,** 7564-7579.

Minai, L., Wostrikoff, K., Wollman, F.A., and Choquet, Y. (2006). Chloroplast biogenesis of
photosystem II cores involves a series of assembly-controlled steps that regulate translation. Plant
Cell **18,** 159-175.

Murata, N., Allakhverdiev, S.I., and Nishiyama, Y. (2012). The mechanism of photoinhibition in
vivo: Re-evaluation of the roles of catalase, alpha-tocopherol, non-photochemical quenching, and
electron transport. Biochim. Biophys. Acta **1817,** 1127-1133.

Nawaz, G., and Kang, H. (2017). Chloroplast- or mitochondria-targeted DEAD-box RNA helicases
play essential roles in organellar RNA metabolism and abiotic stress responses. Front. Plant Sci. **8**.

Nawaz, G., Sai, T.Z.T., Lee, K., Park, S.J., Dinh, S.N., and Kang, H. (2021). BrRH37, a cabbage
(*Brassica rapa*) DEAD-Box RNA helicase, confers drought tolerance and ABA response in
transgenic Arabidopsis plants. J. Plant Biol. **64,** 327-336.

Nellaepalli, S., Ozawal, S.I., Kuroda, H., and Takahashi, Y. (2018). The photosystem I assembly
apparatus consisting of Ycf3-Y3IP1 and Ycf4 modules. Nat. Commun. **9**.

Nellaepalli, S., Kim, R.G., Grossman, A.R., and Takahashi, Y. (2021). Interplay of four auxiliary
factors is required for the assembly of photosystem I reaction center subcomplex. Plant J. **106,**
1075-1086.

Niyogi, K.K. (1999). Photoprotection revisited: Genetic and molecular approaches. Ann. Rev. Plant
Physiol. Plant Mol. Biol. **50,** 333-359.

Ohad, I., Kyle, D.J., and Arntzen, C.J. (1984). Membrane protein damage and repair: Removal and
replacement of inactivated 32-kilodalton polypeptides in chloroplasts membranes. J. Cell Biol. **99,**
481-485.

Olinares, P.D.B., Ponnala, L., and van Wijk, K.J. (2010). Megadalton complexes in the chloroplast
stroma of *Arabidopsis thaliana* characterized by size exclusion chromatography, mass
spectrometry, and hierarchical clustering. Mol. Cell. Proteomics **9,** 1594-1615.

Ozawa, S.-i., Onishi, T., and Takahashi, Y. (2010). Identification and characterization of an
assembly intermediate subcomplex of Photosystem I in the green alga *Chlamydomonas reinhardtii*.
J. Biol. Chem. **285,** 20072-20079.

Peers, G., Truong, T.B., Ostendorf, E., Busch, A., Elrad, D., Grossman, A.R., Hippler, M., and
Niyogi, K.K. (2009). An ancient light-harvesting protein is critical for the regulation of algal
photosynthesis. Nature **462,** 518-521.

Pfalz, J., Liebers, M., Hirth, M., Gruebler, B., Holtzegel, U., Schroeter, Y., Dietzel, L., and
Pfannschmidt, T. (2012). Environmental control of plant nuclear gene expression by chloroplast
redox signals. Front. Plant Sci. **3**.

Piippo, M., Allahverdiyeva, Y., Paakkarinen, V., Suoranta, U.M., Battchikova, N., and Aro,
E.M. (2006). Chloroplast-mediated regulation of nuclear genes in *Arabidopsis thaliana* in the
absence of light stress. Physiol. Genomics **25,** 142-152.

Rosana, A.R.R., Ventakesh, M., Chamot, D., Patterson-Fortin, L.M., Tarassova, O., Espie, G.S.,
and Owttrim, G.W. (2012). Inactivation of a low temperature-induced RNA helicase in
Synechocystis sp PCC 6803: physiological and morphological consequences. Plant Cell Physiol.
53, 646-658.

Ru, J.-N., Hou, Z.-H., Zheng, L., Zhao, Q., Wang, F.-Z., Chen, J., Zhou, Y.-B., Chen, M., Ma,
Y.-Z., Xi, Y.-J., and Xu, Z.-S. (2021). Genome-wide analysis of DEAD-box RNA helicase family
in wheat (*Triticum aestivum*) and functional identification of TaDEAD-box57 in abiotic stress
responses. Front. Plant Sci. **12**.

Schuster, G., Timberg, R., and Ohad, I. (1988). Turnover of thylakoid Photosystem II proteins
during photoinhibition of *Chlamydomonas reinhardtii*. Eur. J. Biochem. **177,** 403-410.

Schwanhausser, B., Busse, D., Li, N., Dittmar, G., Schuchhardt, J., Wolf, J., Chen, W., and
Selbach, M. (2011). Global quantification of mammalian gene expression control. Nature **473,**
337-342.

Schwenkert, S.F., Alisdair R.; Geigenberger, Peter; Leister, Dario; Möhlmann, Torsten;
Naranjo, Belen ; Neuhaus, H. Ekkehard. (in press). Chloroplasts are key players to cope with
light and temperature stress. Trends Plant Sci.

Siaut, M., Cuine, S., Cagnon, C., Fessler, B., Nguyen, M., Carrier, P., Beyly, A., Beisson, F.,
Triantaphylides, C., Li-Beisson, Y., and Peltier, G. (2011). Oil accumulation in the model green
alga *Chlamydomonas reinhardtii*: characterization, variability between common laboratory strains
and relationship with starch reserves. Bmc Biotechnol. **11**.

Silverman, E., Edwalds-Gilbert, G., and Lin, R.J. (2003). DExD/H-box proteins and their partners:
helping RNA helicases unwind. Gene **312,** 1-16.

Sireesha, K., Radharani, B., Krishna, P.S., Sreedhar, N., Subramanyam, R., Mohanty, P., and
Prakash, J.S.S. (2012). RNA helicase, CrhR is indispensable for the energy redistribution and the
regulation of photosystem stoichiometry at low temperature in *Synechocystis* sp PCC6803.
Biochim. Biophys. Acta **1817,** 1525-1536.

Stern, D.B., Goldschmidt-Clermont, M., and Hanson, M.R. (2010). Chloroplast RNA metabolism.
Ann. Rev. Plant Biol. **61,** 125-155.

Sun, Y., and Zerges, W. (2015). Translational regulation in chloroplasts for development and
homeostasis. Biochim. Biophys. Acta **1847,** 809-820.

Suorsa, M., Jarvi, S., Grieco, M., Nurmi, M., Pietrzykowska, M., Rantala, M., Kangasjarvi, S.,
Paakkarinen, V., Tikkanen, M., Jansson, S., and Aro, E.-M. (2012). PROTON GRADIENT
REGULATION5 Is essential for proper acclimation of Arabidopsis photosystem I to naturally and
artificially fluctuating light conditions. Plant Cell **24,** 2934-2948.

Takahashi, Y., Yasui, T., Stauber, E.J., and Hippler, M. (2004). Comparison of the subunit
compositions of the PSI-LHCI supercomplex and the LHCI in the green alga *Chlamydomonas*
reinhardtii. Biochemistry **43,** 7816-7823.

Tarazona, S., Garcia-Alcalde, F., Dopazo, J., Ferrer, A., and Conesa, A. (2011). Differential
expression in RNA-seq: A matter of depth. Genome Res. **21,** 2213-2223.

Tardif, M., Atteia, A., Specht, M., Cogne, G., Rolland, N., Brugiere, S., Hippler, M., Ferro, M.,
Bruley, C., Peltier, G., Vallon, O., and Cournac, L. (2012). PredAlgo: a new subcellular
localization prediction tool dedicated to green algae. Mol. Biol. Evol. **29,** 3625-3639.

Thimm, O., Blasing, O., Gibon, Y., Nagel, A., Meyer, S., Kruger, P., Selbig, J., Muller, L.A.,
Rhee, S.Y., and Stitt, M. (2004). MAPMAN: a user-driven tool to display genomics data sets onto
diagrams of metabolic pathways and other biological processes. Plant J. **37,** 914-939.

Tibiletti, T., Auroy, P., Peltier, G., and Caffarri, S. (2016). *Chlamydomonas reinhardtii* PsbS
protein is functional and accumulates rapidly and transiently under high light. Plant Physiol. **171,**
2717-2730.

Tikkanen, M., Mekala, N.R., and Aro, E.-M. (2014). Photosystem II photoinhibition-repair cycle
protects Photosystem I from irreversible damage. Biochim. Biophys. Acta **1837,** 210-215.

Tokutsu, R., Kato, N., Bui, K.H., Ishikawa, T., and Minagawa, J. (2012). Revisiting the
supramolecular organization of Photosystem II in *Chlamydomonas reinhardtii*. J. Biol. Chem. **287,**
31574-31581.

Tolleter, D., Ghysels, B., Alric, J., Petroutsos, D., Tolstygina, I., Krawietz, D., Happe, T., Auroy,
P., Adriano, J.M., Beyly, A., Cuine, S., Plet, J., Reiter, I.M., Genty, B., Cournac, L., Hippler,
M., and Peltier, G. (2011). Control of hydrogen photoproduction by the proton gradient generated
by cyclic electron flow in *Chlamydomonas reinhardtii*. Plant Cell **23,** 2619-2630.

Wang, Y.C., Duby, G., Purnelle, B., and Boutry, M. (2000). Tobacco VDL gene encodes a plastid
DEAD box RNA helicase and is involved in chloroplast differentiation and plant morphogenesis.
Plant Cell **12,** 2129-2142.

Westrich, L.D., Gotsmann, V.L., Herkt, C., Ries, F., Kazek, T., Troesch, R., Armbruster, L.,
Muehlenbeck, J.S., Ramundo, S., Nickelsen, J., Finkemeier, I., Wirtz, M., Storchova, Z.,
Raeschle, M., and Willmund, F. (2021). The versatile interactome of chloroplast ribosomes
revealed by affinity purification mass spectrometry. Nucleic Acids Res. **49,** 400-415.

Zones, J.M., Blaby, I.K., Merchant, S.S., and Umen, J.G. (2015). High-resolution profiling of a
synchronized diurnal transcriptome from *Chlamydomonas reinhardtii* reveals continuous cell and
metabolic differentiation. Plant Cell **27,** 2743-2769.

Figure 1. Isolation and molecular characterization of hf_0, a *C. reinhardtii* high chlorophyll fluorescence mutant. (A) The hf_0 mutant was isolated from the screening of an insertion library based on the analysis of chlorophyll fluorescence transients (after recording F_0 fluorescence in the dark, a saturating flash at 5s was followed by 240 μmol photons m^{-2} s^{-1} actinic red light at 14 s). (B) The paromomycin resistance cassette (*AphVIII* gene) is integrated in exon 7 of the Cre03.g166650 locus encoding a gene annotated as a DEAD box RNA helicase, renamed here *CreRH22* based on a phylogenetic analysis (Supplemental Fig. 2). The *CreRH22* gene is composed of 9 exons (black boxes) and 8 introns (black lines). Orange arrows show primers used for RT-qPCR shown in (C). The CreRH22 protein shows typical features of DEAD box RNA helicases, including a Q motif, an ATP binding domain containing a DEAD motif, and an ATPase RNA unwinding helicase domain. (C) RT-qPCR showing the absence of the *CreRH22* gene transcript in hf_0. *CBLP* (alias RACK1 Cre06.g27822) is used as loading control.

Figure 2. Effect of light on photosynthetic properties of the *hf₀* mutant. Wild-type (WT), *hf₀* and two complemented lines (*hf₀::HF₀-c7* and *hf₀::HF₀-g6*) were grown photoautotrophically in 3% CO_2-enriched air under high light (240 µmol photons m^{-2} s^{-1} A, C) or low light (15 µmol photons m^{-2} s^{-1} B, D). Representative chlorophyll fluorescence measurements during a dark/light (100 µmol photons m^{-2} s^{-1})/dark transient for high light (A) or low light (B) grown cells. Black and white boxes indicate dark and light periods, respectively. (C, D) PSII yields determined from chlorophyll fluorescence measurements as F_v/F_m in the dark after 15 min dark adaptation or $(F_{m'}-F_s)/F_{m'}$ in the light in high light (C) or low light (D) grown cells.

Figure 3. Ribosomal RNAs processing is altered in the *hf₀* mutant. (A) Scheme showing the organization of the *C. reinhardtii* chloroplast ribosomal RNA (rRNA) operon and location of the different probes (black bars) used for Northern-blot analysis. (B) rRNAs levels analyzed by Northern blot in WT and *hf₀* cells grown photoautotrophically under a light intensity of 60 µmol m^{-2} s^{-1}. 5µg of total RNA were loaded on agarose gels, blotted on nylon membrane and hybridized with DIG-labeled probes. For loading controls (lower blue boxes), nylon membranes were stained with methylene blue after hybridation.

Figure 4. Accumulation of major chloroplast proteins in WT and *hf₀* upon high light exposure determined by immuno -analysis. Cells were grown photo-autotrophically in photobioreactors at a low light intensity (7.5 µmol photons m^{-2} s^{-1}) and exposed to high light (60 µmol photons m^{-2} s^{-1}), for 6h or 26h, respectively corresponding to light intensities of about 20 and 200 µmol photons m^{-2} s^{-1} in flasks (Dang et al. 2015). Protein quantification based on immuno-analysis performed in the linearity range of the antibody detection. Shown are mean values (+/-SD, n=3) normalized on the WT protein content measured at t_0.

Figure 5. Pulse labeling experiments of thylakoid proteins show a strong decrease of D1 protein turn-over under high light in hf_0. Cells were grown photoautotrophically at LL up to a cell density of 2. 10^6 cells mL^{-1} and then switched to HL for 6 h to the cell density of 6. 10^6 cells mL^{-1}, and then labeled for 10 min with ^{35}S (Na_2SO_4). Upon extraction and SDS-PAGE separation of total cellular proteins, labeled proteins were visualized by autoradiography. Autoradiograms show labeled thylakoid membrane proteins of WT, hf_0 and two complemented lines, $hf_0::HF_0$-c7 (complemented line obtained by chloroplast transformation) and $hf_0::HF_0$-g6 (complemented line obtained by nuclear transformation). The gel was stained with coomassie brilliant blue (CBB) before exposing to the imaging plate.

Figure 6. Effect of high light and chloramphenicol addition on photosynthetic activity of the hf_0 mutant. Wild-type (WT), hf_0 and two complemented lines ($hf_0::HF_0$-$c7$ and $hf_0::HF_0$-$g6$) were grown in batch cultures under 15 µmol photons m^{-2} s^{-1}. At t_0 cell cultures were switched to 240 µmol photons m^{-2} s^{-1} in the absence **(A, C)** or in the presence **(B, D)** of 50 mg. L^{-1} chloramphenicol added 1h before the high light switch. At different time points after the light switch, maximal PSII yields were measured as the F_v/F_m ratio after 15 min dark adaptation **(A, B)** and PSII operating yields were measured as $(F_{m'}-F_s)/F_{m'}$ following a 3 min light period (430 µmol photons m^{-2} s^{-1}). Shown are means ± SD (n=3).

Figure 7. Low temperature chlorophyll fluorescence and chlorophyll-protein complexes are affected in hf_0, but state transitions are not. WT and hf_0 cells were grown photo-autrophically in photobioreactors under 7.5 μmol photons m^{-2} s^{-1} (LL) and then switched to 60 μmol photons m^{-2} s^{-1} (HL), respectively corresponding to light intensities of about 20 and 200 μmol photons m^{-2} s^{-1} in flasks (Dang et al. 2015). (A,B) 77K chlorophyll fluorescence emission spectra of cells grown under LL (A) or 26 after HL switch (B). (C) 77K chlorophyll fluorescence emission ratio E_{712nm}/E_{686nm} , and (D) F_0/F_m chlorophyll fluorescence at different time points after HL switch. (E) Immunodetection of phosphorylated LHCII using an anti-phospho-threonine antibody. Arrows indicate protein bands with lower abundance in hf_0. (F) Blue-native PAGE of thylakoid complexes from WT and hf_0 grown under LL or upon 26h of HL exposure. Middle arrows indicate green bands with lower abundance in hf_0 and the right arrow a green band of higher abundance in hf_0.

Figure 8. Two-dimensional PAGE of thylakoid proteins from WT and hf_0 cells upon HL exposure. WT and hf_0 cells were grown photo-autotrophically in batch cultures under LL (40 µmol photons m^{-2} s^{-1}) and then exposed to HL (240 µmol photons m^{-2} s^{-1}) for 26h. Thylakoid proteins were solubilized with 1% dodecyl-maltoside, separated by blue native PAGE in a first dimension and by SDS-PAGE in a second dimension. The red dotted ellipse highlights proteins associated to high MW PSI-LHCI supercomplexes in the WT (A) that migrates as low MW complexes attributed to LHCI oligomers in hf_0 (red arrow in B). The gel strip in the 20-35kDa MW range shown as a red box (B) was taken for proteomic analysis by mass spectrometry (Supplemental Table 1).

Figure 9. RNAseq analysis of light-induced gene expression in wild-type and *hf₀ C. reinhardtii* cells. Cells were cultivated in PBRs operated as turbidostats in 3% CO_2-enriched air under LL (7.5 µmol photons m^{-2} s^{-1}) for 24h and then exposed to HL (60 µmol photons m^{-2} s^{-1}). Samples for RNAseq analysis were taken from two biological replicates under LL before the HL switch, and the 2h and 24h after the switch. (A) Shown are numbers of up and down Differentially Expressed Genes (DEGs) with a log 2(Fold Change) > 1 and <-1, using a False Discovery Rate (FDR) ≤ 0.001. (B, C) Functional analysis was performed DEGs from WT (A) and *hf₀* (B) using the MapMan 3.5.1R2 package (Thimm et al., 2004).

Figure 10. Expression patterns of genes encoding chloroplast-, mitochondria-, and cytosol-targeted ribosomal proteins. (A to C) Heat maps depicting relative expression levels of ribosomal protein genes and translational regulators targeted to cytosol (**A**), chloroplast (**B**), and mitochondria (**C**). The maximum expression level for each gene is set to 1. * Genes without significant differential expression (|log2FC|<2 as defined in Methods). (**D to F**) Boxplot of the normalized expression values of ribosomal protein genes targeted to cytosol (**D**), chloroplast (**E**), and mitochondria (**F**).

Figure 11. Biomass productivity and concentrations of intracellular starch and TAGs are reduced in *hf₀*. WT and *hf₀* cells were grown in 1L photobioreactors operated as turbidostats and maintained at a constant biomass concentration (≈ 1.5 x 10⁶ cells mL⁻¹) by continuous addition of fresh medium. (A) Specific growth rates were determined from dilution rates measured at different light intensities (from 7.5 up to 300 μmol photons m⁻² s⁻¹) following at least 24h stabilization. Shown are means ± SD (n=4 for 60 and 200 μmol m⁻² s⁻¹, n=2 for 7.5, 11, and 300 μmol m⁻² s⁻¹). (B) Intracellular starch, and (C) intracellular TAG contents were determined on cell samples following at least 24h stabilization at the specified light intensity (shown are means ± SD, n=3).

Figure 12. Hypothetical scheme proposed to explain the *hf₀* mutant phenotype. The *hf₀* mutant is defected in a plastidial DEAD box RNA helicase (CreRH22) involved in the processing of plastidial rRNA operon and ribosome biogenesis. (A) Under LL the turnover of chloroplast proteins is low and the translation capacity is sufficient to maintain photosynthesis at similar levels in the wild-type and *hf₀*. (B,C) Under HL exposure, the amount of chloroplast ribosomes increases in the wild-type (B) but remains constant in the mutant (C). In these conditions, a much higher rate of protein synthesis is needed due to the permanent degradation and synthesis of the D1 protein and to the biosynthesis of additional PSII subunits (essentially D1) and to a lesser extent PSI subunits (PsaA/B/C, not shown in the figure). In these conditions, the *hf₀* mutant is unable to synthesize photosynthetic proteins at a sufficient rate, thus resulting in an inhibition of photosynthesis.

Parsed Citations

Adams, W.W., III, Muller, O., Cohu, C.M., and Demmig-Adams, B. (2013). May photoinhibition be a consequence, rather than a cause, of limited plant productivity? Photosynth. Res. 117, 31-44.
Google Scholar: Author Only Title Only Author and Title

Allen, J.F., and Pfannschmidt, T. (2000). Balancing the two photosystems: photosynthetic electron transfer governs transcription of reaction centre genes in chloroplasts. Philos. Trans. R. Soc. Lond. B Biol. Sci. 355, 1351-1357.
Google Scholar: Author Only Title Only Author and Title

Allorent, G., Tokutsu, R., Roach, T., Peers, G., Cardol, P., Girard-Bascou, J., Seigneurin-Berny, D., Petroutsos, D., Kuntz, M., Breyton, C., Franck, F., Wollman, F.A, Niyogi, K.K., Krieger-Liszkay, A, Minagawa, J., and Finazzi, G. (2013). A dual strategy to cope with high light in Chlamydomonas reinhardtii. Plant Cell 25, 545-557.
Google Scholar: Author Only Title Only Author and Title

Asakura, Y., Galarneau, E., Watkins, K.P., Barkan, A., and van Wijk, K.J. (2012). Chloroplast RH3 DEAD box RNA helicases in maize and Arabidopsis function in splicing of specific group II introns and affect chloroplast ribosome biogenesis. Plant Physiol. 159, 961-974.
Google Scholar: Author Only Title Only Author and Title

Baltz, A, Dang, K.V., Beyly, A, Auroy, P., Richaud, P., Cournac, L., and Peltier, G. (2014). Plastidial expression of type II NAD(P)H dehydrogenase increases the reducing state of plastoquinones and hydrogen photoproduction rate by the indirect pathway in Chlamydomonas reinhardtii. Plant Physiol. 165, 1344-1352.
Google Scholar: Author Only Title Only Author and Title

Berthold, P., Schmitt, R., and Mages, W. (2002). An engineered Streptomyces hygroscopicus aph 7" gene mediates dominant resistance against hygromycin B in Chlamydomonas reinhardtii. Protist 153, 401-412.
Google Scholar: Author Only Title Only Author and Title

Bonente, G., Pippa, S., Castellano, S., Bassi, R., and Ballottari, M. (2012). Acclimation of Chlamydomonas reinhardtii to different growth irradiances. J. Biol. Chem. 287, 5833-5847.
Google Scholar: Author Only Title Only Author and Title

Bouyssie, D., Hesse, A.M., Mouton-Barbosa, E., Rompais, M., Macron, C., Carapito, C., de Peredo, A.G., Coute, Y., Dupierris, V., Burel, A, Menetrey, J.P., Kalaitzakis, A, Poisat, J., Romdhani, A, Burlet-Schiltz, O., Cianferani, S., Garin, J., and Bruley, C. (2020). Proline: an efficient and user-friendly software suite for large-scale proteomics. Bioinformatics 36, 3148-3155.
Google Scholar: Author Only Title Only Author and Title

Casabona, M.G., Vandenbrouck, Y., Attree, I., and Coute, Y. (2013). Proteomic characterization of Pseudomonas aeruginosa PAO1 inner membrane. Proteomics 13, 2419-2423.
Google Scholar: Author Only Title Only Author and Title

Chaux, F., Peltier, G., and Johnson, X. (2015). A security network in PSI photoprotection: regulation of photosynthetic control, NPQ and O-2 photoreduction by cyclic electron flow. Front. Plant Sci. 6.
Google Scholar: Author Only Title Only Author and Title

Cheng, J., Zhou, S., Yang, K., Yu, H., Chen, R., Zeng, L., Li, H., Wang, Y., and Song, J. (2021). Identification of RNA helicases in Medicago truncatula and their expression patterns under abiotic stress. Physiol. Mol. Biol. Plants 27, 2283-2296.
Google Scholar: Author Only Title Only Author and Title

Chi, W., He, B., Mao, J., Li, Q., Ma, J., Ji, D., Zou, M., and Zhang, L. (2012). The function of RH22, a DEAD RNA helicase, in the biogenesis of the 50S ribosomal subunits of Arabidopsis chloroplasts. Plant Physiol. 158, 693-707.
Google Scholar: Author Only Title Only Author and Title

Christopher, D.A., and Mullet, J.E. (1994). Separate photosensory pathways co-regulate blue light/ultraviolet-A-activated psbD-psbC transcription and light-induced D2 and CP43 degradation in barley (Hordeum vulgare) chloroplasts. Plant Physiol. 104, 1119-1129.
Google Scholar: Author Only Title Only Author and Title

Chua, N.H., and Bennoun, P. (1975). Thylakoid membrane polypeptides of Chlamydomonas reinhardtii wild-type and mutant strains deficient in Photosystem 2 reaction center. Proc. Natl. Acad. Sci. USA 72, 2175-2179.
Google Scholar: Author Only Title Only Author and Title

Chure, G., and Cremer, J. (2022). An optimal regulation of fluxes dictates microbial growth in and out of steady-state. bioRxiv, 2022.2001.2027.477569.
Google Scholar: Author Only Title Only Author and Title

Coute, Y., Bruley, C., and Burger, T. (2020). Beyond target-decoy competition: Stable validation of peptide and protein identifications in mass spectrometry-based discovery proteomics. Anal. Chem. 92, 14898-14906.
Google Scholar: Author Only Title Only Author and Title

Dang, K.V., Plet, J., Tolleter, D., Jokel, M., Cuine, S., Carrier, P., Auroy, P., Richaud, P., Johnson, X., Alric, J., Allahverdiyeva, Y., and Peltier, G. (2014). Combined increases in mitochondrial cooperation and oxygen photoreduction compensate for deficiency in cyclic electron flow in Chlamydomonas reinhardtii. Plant Cell 26, 3036-3050.
Google Scholar: Author Only Title Only Author and Title

Depege, N., Bellafiore, S., and Rochaix, J.D. (2003). Role of chloroplast protein kinase Stt7 in LHCII phosphorylation and state transition in Chlamydomonas. Science 299, 1572-1575.
Google Scholar: Author Only Title Only Author and Title

Dietzel, L., Glaesser, C., Liebers, M., Hiekel, S., Courtois, F., Czarnecki, O., Schlicke, H., Yan, Z., Boerner, T., Mayer, K., Grimm, B., and Pfannschmidt, T. (2015). Identification of early nuclear target genes of plastidial redox signals that trigger the long-term response of Arabidopsis to light quality shifts. Mol. Plant 8, 1237-1252.
Google Scholar: Author Only Title Only Author and Title

Erickson, E., Wakao, S., and Niyogi, K.K. (2015). Light stress and photoprotection in Chlamydomonas reinhardtii. Plant J. 82, 449-465.
Google Scholar: Author Only Title Only Author and Title

Escoubas, J.M., Lomas, M., Laroche, J., and Falkowski, P.G. (1995). Light intensity regulation of cab gene transcription is signaled by the redox state of the plastoquinone pool. Proc. Natl. Acad. Sci. USA 92, 10237-10241.
Google Scholar: Author Only Title Only Author and Title

Fristedt, R., Scharff, L.B., Clarke, C.A, Wang, Q., Lin, C., Merchant, S.S., and Bock, R. (2014). RBF1, a plant homolog of the bacterial ribosome-binding factor RbfA, acts in processing of the chloroplast 16S ribosomal RNA. Plant Physiol. 164, 201-215.
Google Scholar: Author Only Title Only Author and Title

Girolomoni, L., Ferrante, P., Berteotti, S., Giuliano, G., Bassi, R., and Ballottari, M. (2017). The function of LHCBM4/6/8 antenna proteins in Chlamydomonas reinhardtii. J. Exp. Bot. 68, 628-642.
Google Scholar: Author Only Title Only Author and Title

Goold, H.D., Cuine, S., Legeret, B., Liang, Y., Brugiere, S., Auroy, P., Javot, H., Tardif, M., Jones, B., Beisson, F., Peltier, G., and Li-Beisson, Y. (2016). Saturating light induces sustained accumulation of oil in plastidal lipid droplets in Chlamydomonas reinhardtii. Plant Physiol. 171, 2406-2417.
Google Scholar: Author Only Title Only Author and Title

Harris, E.H. (2009). The Chlamydomonas sourcebook. Second edition. (Academic Press).
Google Scholar: Author Only Title Only Author and Title

Holloway, S.P., and Herrin, D.L. (1998). Processing of a composite large subunit rRNA: Studies with Chlamydomonas mutants deficient in maturation of the 23S-like rRNA. Plant Cell 10, 1193-1206.
Google Scholar: Author Only Title Only Author and Title

Jarvi, S., Suorsa, M., and Aro, E.-M. (2015). Photosystem II repair in plant chloroplasts - Regulation, assisting proteins and shared components with photosystem II biogenesis. Biochim. Biophys. Acta 1847, 900-909.
Google Scholar: Author Only Title Only Author and Title

Jarvis, P., and Lopez-Juez, E. (2013). Biogenesis and homeostasis of chloroplasts and other plastids. Nat. Rev. Mol. Cell Biol. 14, 787-802.
Google Scholar: Author Only Title Only Author and Title

Karpowicz, S.J., Prochnik, S.E., Grossman, A.R., and Merchant, S.S. (2011). The GreenCut2 resource, a phylogenomically derived inventory of proteins specific to the plant lineage. J. Biol. Chem. 286, 21427-21439.
Google Scholar: Author Only Title Only Author and Title

Kettunen, R., Pursiheimo, S., Rintamaki, E., VanWijk, K.J., and Aro, E.M. (1997). Transcriptional and translational adjustments of psbA gene expression in mature chloroplasts during photoinhibition and subsequent repair of photosystem II. Eur. J. Biochem. 247, 441-448.
Google Scholar: Author Only Title Only Author and Title

Kim, J., Darlington, A, Salvador, M., Utrilla, J., and Jimenez, J.I. (2020). Trade-offs between gene expression, growth and phenotypic diversity in microbial populations. Curr. Opin. Biotechnol. 62, 29-37.
Google Scholar: Author Only Title Only Author and Title

Kramer, D.M., Johnson, G., Kiirats, O., and Edwards, G.E. (2004). New fluorescence parameters for the determination of Q(A) redox state and excitation energy fluxes. Photosynth. Res. 79, 209-218.
Google Scholar: Author Only Title Only Author and Title

Krieger-Liszkay, A, Kienzler, K., and Johnson, G.N. (2000). Inhibition of electron transport at the cytochrome b(6)f complex protects photosystem II from photoinhibition. FEBS Lett. 486, 191-194.
Google Scholar: Author Only Title Only Author and Title

Kujat, S.L., and Owttrim, G.W. (2000). Redox-regulated RNA helicase expression. Plant Physiol. 124, 703-713.
Google Scholar: Author Only Title Only Author and Title

Li, L., Aro, E.-M., and Millar, A.H. (2018). Mechanisms of photodamage and protein turnover in photoinhibition. Trends Plant Sci. 23, 667-676.
Google Scholar: Author Only Title Only Author and Title

Li, R.Q., Li, Y.R., Kristiansen, K., and Wang, J. (2008). SOAP: short oligonucleotide alignment program. Bioinformatics 24, 713-714.
Google Scholar: Author Only Title Only Author and Title

Li, R.Q., Yu, C., Li, Y.R., Lam, T.W., Yiu, S.M., Kristiansen, K., and Wang, J. (2009). SOAP2: an improved ultrafast tool for short read alignment. Bioinformatics 25, 1966-1967.
Google Scholar: Author Only Title Only Author and Title

Linder, P., and Jankowsky, E. (2011). From unwinding to clamping - the DEAD box RNA helicase family. Nat. Rev. Mol. Cell Biol. 12, 505-516.
Google Scholar: Author Only Title Only Author and Title

Livak, K.J., and Schmittgen, T.D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2(T)(-Delta Delta C) method 25, 402-408.
Google Scholar: Author Only Title Only Author and Title

Lu, C.-A., Huang, C.-K., Huang, W.-S., Huang, T.-S., Liu, H.-Y., and Chen, Y.-F. (2020). DEAD-Box RNA helicase 42 plays a critical role in pre-mRNA splicing under cold stress. Plant Physiol. 182, 255-271.
Google Scholar: Author Only Title Only Author and Title

Michelet, L., Lefebvre-Legendre, L., Burr, S.E., Rochaix, J.-D., and Goldschmidt-Clermont, M. (2011). Enhanced chloroplast transgene expression in a nuclear mutant of Chlamydomonas. Plant Biotechnol. J. 9, 565-574.
Google Scholar: Author Only Title Only Author and Title

Migur, A., Heyl, F., Fuss, J., Srikumar, A., Huettel, B., Steglich, C., Prakash, J.S.S., Reinhardt, R., Backofen, R., Owttrim, G.W., and Hess, W.R. (2021). The temperature-regulated DEAD-box RNA helicase CrhR interactome: autoregulation and photosynthesis-related transcripts. J. Exp. Bot. 72, 7564-7579.
Google Scholar: Author Only Title Only Author and Title

Minai, L., Wostrikoff, K., Wollman, F.A., and Choquet, Y. (2006). Chloroplast biogenesis of photosystem II cores involves a series of assembly-controlled steps that regulate translation. Plant Cell 18, 159-175.
Google Scholar: Author Only Title Only Author and Title

Murata, N., Allakhverdiev, S.I., and Nishiyama, Y. (2012). The mechanism of photoinhibition in vivo: Re-evaluation of the roles of catalase, alpha-tocopherol, non-photochemical quenching, and electron transport. Biochim. Biophys. Acta 1817, 1127-1133.
Google Scholar: Author Only Title Only Author and Title

Nawaz, G., and Kang, H. (2017). Chloroplast- or mitochondria-targeted DEAD-box RNA helicases play essential roles in organellar RNA metabolism and abiotic stress responses. Front. Plant Sci. 8.
Google Scholar: Author Only Title Only Author and Title

Nawaz, G., Sai, T.Z.T., Lee, K., Park, S.J., Dinh, S.N., and Kang, H. (2021). BrRH37, a cabbage (Brassica rapa) DEAD-Box RNA helicase, confers drought tolerance and ABA response in transgenic Arabidopsis plants. J. Plant Biol. 64, 327-336.
Google Scholar: Author Only Title Only Author and Title

Nellaepalli, S., Ozawal, S.I., Kuroda, H., and Takahashi, Y. (2018). The photosystem I assembly apparatus consisting of Ycf3-Y3IP1 and Ycf4 modules. Nat. Commun. 9.
Google Scholar: Author Only Title Only Author and Title

Nellaepalli, S., Kim, R.G., Grossman, A.R., and Takahashi, Y. (2021). Interplay of four auxiliary factors is required for the assembly of photosystem I reaction center subcomplex. Plant J. 106, 1075-1086.
Google Scholar: Author Only Title Only Author and Title

Niyogi, K.K. (1999). Photoprotection revisited: Genetic and molecular approaches. Ann. Rev. Plant Physiol. Plant Mol. Biol. 50, 333-359.
Google Scholar: Author Only Title Only Author and Title

Ohad, I., Kyle, D.J., and Arntzen, C.J. (1984). Membrane protein damage and repair: Removal and replacement of inactivated 32-kilodalton polypeptides in chloroplasts membranes. J. Cell Biol. 99, 481-485.
Google Scholar: Author Only Title Only Author and Title

Olinares, P.D.B., Ponnala, L., and van Wijk, K.J. (2010). Megadalton complexes in the chloroplast stroma of Arabidopsis thaliana characterized by size exclusion chromatography, mass spectrometry, and hierarchical clustering. Mol. Cell. Proteomics 9, 1594-1615.

Google Scholar: Author Only Title Only Author and Title

Ozawa, S.-i., Onishi, T., and Takahashi, Y. (2010). Identification and characterization of an assembly intermediate subcomplex of Photosystem I in the green alga Chlamydomonas reinhardtii. J. Biol. Chem. 285, 20072-20079.
  Google Scholar: Author Only Title Only Author and Title

Peers, G., Truong, T.B., Ostendorf, E., Busch, A, Elrad, D., Grossman, A.R., Hippler, M., and Niyogi, K.K. (2009). An ancient light-harvesting protein is critical for the regulation of algal photosynthesis. Nature 462, 518-521.
  Google Scholar: Author Only Title Only Author and Title

Pfalz, J., Liebers, M., Hirth, M., Gruebler, B., Holtzegel, U., Schroeter, Y., Dietzel, L., and Pfannschmidt, T. (2012). Environmental control of plant nuclear gene expression by chloroplast redox signals. Front. Plant Sci. 3.
  Google Scholar: Author Only Title Only Author and Title

Piippo, M., Allahverdiyeva, Y., Paakkarinen, V., Suoranta, U.M., Battchikova, N., and Aro, E.M. (2006). Chloroplast-mediated regulation of nuclear genes in Arabidopsis thaliana in the absence of light stress. Physiol. Genomics 25, 142-152.
  Google Scholar: Author Only Title Only Author and Title

Rosana, A.R.R., Ventakesh, M., Chamot, D., Patterson-Fortin, L.M., Tarassova, O., Espie, G.S., and Owttrim, G.W. (2012). Inactivation of a low temperature-induced RNA helicase in Synechocystis sp PCC 6803: physiological and morphological consequences. Plant Cell Physiol. 53, 646-658.
  Google Scholar: Author Only Title Only Author and Title

Ru, J.-N., Hou, Z.-H., Zheng, L., Zhao, Q., Wang, F.-Z, Chen, J., Zhou, Y.-B., Chen, M., Ma, Y.-Z, Xi, Y.-J., and Xu, Z-S. (2021). Genome-wide analysis of DEAD-box RNA helicase family in wheat (Triticum aestivum) and functional identification of TaDEAD-box57 in abiotic stress responses. Front. Plant Sci. 12.
  Google Scholar: Author Only Title Only Author and Title

Schuster, G., Timberg, R., and Ohad, I. (1988). Turnover of thylakoid Photosystem II proteins during photoinhibition of Chlamydomonas reinhardtii. Eur. J. Biochem. 177, 403-410.
  Google Scholar: Author Only Title Only Author and Title

Schwanhausser, B., Busse, D., Li, N., Dittmar, G., Schuchhardt, J., Wolf, J., Chen, W., and Selbach, M. (2011). Global quantification of mammalian gene expression control. Nature 473, 337-342.
  Google Scholar: Author Only Title Only Author and Title

Schwenkert, S.F., Alisdair R.; Geigenberger, Peter; Leister, Dario; Möhlmann, Torsten; Naranjo, Belen ; Neuhaus, H. Ekkehard. (in press). Chloroplasts are key players to cope with light and temperature stress. Trends Plant Sci.

Siaut, M., Cuine, S., Cagnon, C., Fessler, B., Nguyen, M., Carrier, P., Beyly, A, Beisson, F., Triantaphylides, C., Li-Beisson, Y., and Peltier, G. (2011). Oil accumulation in the model green alga Chlamydomonas reinhardtii: characterization, variability between common laboratory strains and relationship with starch reserves. Bmc Biotechnol. 11.
  Google Scholar: Author Only Title Only Author and Title

Silverman, E., Edwalds-Gilbert, G., and Lin, R.J. (2003). DExD/H-box proteins and their partners: helping RNA helicases unwind. Gene 312, 1-16.
  Google Scholar: Author Only Title Only Author and Title

Sireesha, K., Radharani, B., Krishna, P.S., Sreedhar, N., Subramanyam, R., Mohanty, P., and Prakash, J.S.S. (2012). RNA helicase, CrhR is indispensable for the energy redistribution and the regulation of photosystem stoichiometry at low temperature in Synechocystis sp PCC6803. Biochim. Biophys. Acta 1817, 1525-1536.
  Google Scholar: Author Only Title Only Author and Title

Stern, D.B., Goldschmidt-Clermont, M., and Hanson, M.R. (2010). Chloroplast RNA metabolism. Ann. Rev. Plant Biol. 61, 125-155.
  Google Scholar: Author Only Title Only Author and Title

Sun, Y., and Zerges, W. (2015). Translational regulation in chloroplasts for development and homeostasis. Biochim. Biophys. Acta 1847, 809-820.
  Google Scholar: Author Only Title Only Author and Title

Suorsa, M., Jarvi, S., Grieco, M., Nurmi, M., Pietrzykowska, M., Rantala, M., Kangasjarvi, S., Paakkarinen, V., Tikkanen, M., Jansson, S., and Aro, E.-M. (2012). PROTON GRADIENT REGULATION5 Is essential for proper acclimation of Arabidopsis photosystem I to naturally and artificially fluctuating light conditions. Plant Cell 24, 2934-2948.
  Google Scholar: Author Only Title Only Author and Title

Takahashi, Y., Yasui, T., Stauber, E.J., and Hippler, M. (2004). Comparison of the subunit compositions of the PSI-LHCI supercomplex and the LHCI in the green alga Chlamydomonas reinhardtii. Biochemistry 43, 7816-7823.
  Google Scholar: Author Only Title Only Author and Title

Tarazona, S., Garcia-Alcalde, F., Dopazo, J., Ferrer, A, and Conesa, A (2011). Differential expression in RNA-seq: A matter of depth. Genome Res. 21, 2213-2223.

Google Scholar: Author Only Title Only Author and Title

Tardif, M., Atteia, A, Specht, M., Cogne, G., Rolland, N., Brugiere, S., Hippler, M., Ferro, M., Bruley, C., Peltier, G., Vallon, O., and Cournac, L. (2012). PredAlgo: a new subcellular localization prediction tool dedicated to green algae. Mol. Biol. Evol. 29, 3625-3639.
Google Scholar: Author Only Title Only Author and Title

Thimm, O., Blasing, O., Gibon, Y., Nagel, A, Meyer, S., Kruger, P., Selbig, J., Muller, L.A, Rhee, S.Y., and Stitt, M. (2004). MAPMAN: a user-driven tool to display genomics data sets onto diagrams of metabolic pathways and other biological processes. Plant J. 37, 914-939.
Google Scholar: Author Only Title Only Author and Title

Tibiletti, T., Auroy, P., Peltier, G., and Caffarri, S. (2016). Chlamydomonas reinhardtii PsbS protein is functional and accumulates rapidly and transiently under high light. Plant Physiol. 171, 2717-2730.
Google Scholar: Author Only Title Only Author and Title

Tikkanen, M., Mekala, N.R., and Aro, E.-M. (2014). Photosystem II photoinhibition-repair cycle protects Photosystem I from irreversible damage. Biochim. Biophys. Acta 1837, 210-215.
Google Scholar: Author Only Title Only Author and Title

Tokutsu, R., Kato, N., Bui, K.H., Ishikawa, T., and Minagawa, J. (2012). Revisiting the supramolecular organization of Photosystem II in Chlamydomonas reinhardtii. J. Biol. Chem. 287, 31574-31581.
Google Scholar: Author Only Title Only Author and Title

Tolleter, D., Ghysels, B., Alric, J., Petroutsos, D., Tolstygina, I., Krawietz, D., Happe, T., Auroy, P., Adriano, J.M., Beyly, A, Cuine, S., Plet, J., Reiter, I.M., Genty, B., Cournac, L., Hippler, M., and Peltier, G. (2011). Control of hydrogen photoproduction by the proton gradient generated by cyclic electron flow in Chlamydomonas reinhardtii. Plant Cell 23, 2619-2630.
Google Scholar: Author Only Title Only Author and Title

Wang, Y.C., Duby, G., Purnelle, B., and Boutry, M. (2000). Tobacco VDL gene encodes a plastid DEAD box RNA helicase and is involved in chloroplast differentiation and plant morphogenesis. Plant Cell 12, 2129-2142.
Google Scholar: Author Only Title Only Author and Title

Westrich, L.D., Gotsmann, V.L., Herkt, C., Ries, F., Kazek, T., Troesch, R., Armbruster, L., Muehlenbeck, J.S., Ramundo, S., Nickelsen, J., Finkemeier, I., Wirtz, M., Storchova, Z, Raeschle, M., and Willmund, F. (2021). The versatile interactome of chloroplast ribosomes revealed by affinity purification mass spectrometry. Nucleic Acids Res. 49, 400-415.
Google Scholar: Author Only Title Only Author and Title

Zones, J.M., Blaby, I.K., Merchant, S.S., and Umen, J.G. (2015). High-resolution profiling of a synchronized diurnal transcriptome from Chlamydomonas reinhardtii reveals continuous cell and metabolic differentiation. Plant Cell 27, 2743-2769.
Google Scholar: Author Only Title Only Author and Title