The gradient-based parameter optimization to determine membrane ionic current composition of

human induced pluripotent stem cell-derived cardiomyocytes

Hirohiko Kohjitani[1], Shigeya Koda[2], Yukiko Himeno[2], Takeru Makiyama[1], Yuta Yamamoto[1], Daisuke

Yoshinaga[3], Yimin Wuriyanghai[1], Asami Kashiwa[1], Futoshi Toyoda[4], Yixin Zhang[2], Akira Amano[2], Akinori

Noma[2], Takeshi Kimura[1]

[1]Department of Cardiovascular Medicine, Kyoto University Graduate School of Medicine, Kyoto, Japan

[2]Graduate School of Life Sciences, Ritsumeikan University, Kusatsu, Japan

[3]Department Pediatrics, Kyoto University Graduate School of Medicine, Kyoto, Japan

[4]Department of Physiology, Shiga University of Medical Science, Otsu, Japan

* Correspondence:

Akira Amano

Keywords: Parameter optimization method, hiPSC-CMs, Cardiac action potential, Mathematical

model, Computer simulation

Abbreviations

hiPSC-CMs; human induced pluripotent stem cell-derived cardiomyocytes

AP; action potential

MDP; the maximum diastolic potential

SDD; slow diastolic depolarization

I_m; membrane current

V_m; membrane voltage

orp; optimization of randomized model parameters

PS; pattern search method

BP; base point for searching minimum MSE in the Pattern Search

NP; searching point in reference to BP in the Pattern Search

MSE; mean square error between two different V_m records

stp; step size to move NP

x; subscript to represent membrane current such as I_{Na}, I_{CaL}, I_{K1}, I_{ha}, I_{Kr}, I_{Kur}, I_{Ks} and I_{bNSC}

1.   Abstract

Premature cardiac myocytes derived from human-induced pluripotent stem cells (hiPSC-CMs) show

heterogeneous action potentials (APs), most probably because of different expression patterns of membrane

ionic currents. We aim to develop a method of determining expression patterns of functional channels in terms

of the whole-cell ionic conductances (G_x) using individual AP configurations. However, it has been suggested

that apparently identical AP configurations were obtained by different sets of ionic currents in a mathematical

model of cardiac membrane excitation. If so, the inverse problem of G_x estimation might not be solved. We

computationally tested the feasibility of the gradient-based optimization method. For realistic examination,

conventional 'cell-specific models' were prepared by superimposing the model output on each experimental AP

record by the conventional manual adjustment of G_x of the baseline model. Then, G_xs of 4 ~6 major ionic

currents of the 'cell-specific model' were randomized within a range of ± 5 ~15% and were used as an initial

parameter set for the gradient-based automatic G_x recovery by decreasing the mean square error (MSE)

between the target and model output. When plotted all data points of MSE - G_x relation during the

optimization, we found that the randomized population of G_xs progressively converged to the original value of

the cell-specific model with decreasing MSE. To confirm the absence of any other local minimum in the global

search space, we mapped the MSE by randomizing G_xs over a range of 0.1 ~ 10 times the control. No

additional local minimum of MSE was obvious in the whole parameter space except the global minimum of

MSE at the default model parameter.

2.   Introduction

During more than a half-century, the biophysical characteristics of ion transporting molecules

(channels and ion exchangers) have been extensively analyzed, and biophysical models of each functional

component have been largely detailed [1–4] (for hiPSC-CMs see [5–7]). In addition, various composite cell

models, including the membrane excitation, cell contraction, and the homeostasis of the intracellular ionic

composition, have been developed by integrating mathematical models at molecular levels into the cardiac cell

models [8–11]. These models have already been quite useful in visualizing individual currents underlying the

action potential (AP) configuration under various experimental conditions in matured cardiac myocytes.

However, the utility of these mathematical cell models has been limited because of the lack of extensive

validation for the accuracy of the model output. This is the drawback of the subjective manual fitting used in

almost all mathematical cardiac cell models so far published. A new challenge of such mechanistic models of

cardiac membrane excitation might be an examination in a very different paradigm to assess if the large but

continuous variety of cardiac AP configurations, for example, those recorded in the hiPSC-CMs, can be

reconstructed by applying the automatic parameter optimization method to the human cardiac cell models.

The automatic parameter optimization technique has been used to determine parameters objectively in

a wide range of various biological models (in cardiac electrophysiology; [12–15], in the systems

pharmacology; [16–20]). Because of this utility, a large variety of improvements have been made in the area of

information technology [21,22]. However, in electrophysiology, it has been suggested that different

combinations of model parameters can produce APs, which are very similar[23–25] (see also [13])[13,23–25].

It has been considered that the determination of current density at high fidelity and accuracy requires

additional improvements to the optimization method in the cardiac cell model because of complex interactions

among ionic currents underlying the membrane excitation (see [26], for review; [23]).

The final goal of our study is to develop an objective and accurate method of determining the current

profile (that is, the expression level of functional ionic currents) underlying individual AP configurations. As a

case study, we select a large variety of AP configurations in the hiPSC-CMs, which are difficult to classify into

the conventional nodal-, atrial- or ventricular-types. Nevertheless, it has been clarified that the molecular

bases of the ion channels expressed in the hiPSC-CMs well correspond to those in the adult cardiac

myocytes (GSE154580 GEO Accession viewer (nih.gov)). Thus, we use the human ventricular cell model

(hVC model, [11]) for the baseline model. In the present study, we computationally examine the feasibility of

the basic gradient-based optimization method, pattern search algorithm (PS; [21,27,28]) in the model of

cardiac AP generation. We prepared a given AP configuration using the respective 'cell specific models', which

were prepared by the conventional manual fitting of experimental recordings. To assess the accuracy of the PS

method of parameter optimization, this AP waveform of the cell-specific model was used as a target of

optimization. Then, the initial set of parameters for the optimization was prepared by uniform randomization

centered around the model's default values. The PS algorithm should return the original parameters by

decreasing the error function (MSE) between the modified model output and target AP waveforms. The

accuracy of optimization was definitely judged by recovering the original current amplitude as the MSE

progressively decreased toward zero.

3.   Materials and Methods

3.1.   The baseline model of hiPSC-CM membrane excitation

The baseline model of hiPSC-CMs was essentially the same as the human ventricular cell model (hVC

model), which has been fully described in references [10,11] and shares many comparable characteristics with

other human models so far published [8,9]. The model structure of the hVC model consists of the cell

membrane with a number of ionic channel species and two ion exchangers, the sarcoplasmic reticulum

equipped with the Ca^{2+} pump (SERCA), and the refined Ca^{2+} releasing units coupled with the L-type Ca^{2+}

channels on the cell membrane at the nano-scale dyadic space, the contractile fibers, and the cytosolic three

Ca^{2+} diffusion spaces containing several Ca^{2+}-binding proteins.

The source code of the present hiPSC-CM model was written in VB.Net and is available from our e-

Heart website (http://www.eheartsim.com/en/downloads/).

The kinetics of the ionic currents in the baseline model were readjusted according to new

experimental measurements if available in the hiPSC-CMs [29]. In the present study, the net membrane current

(I_{tot_cell}) is calculated as the sum of nine ion channel currents and two ion transporters (I_{NaK} and I_{NCX}) (Eq 1).

$$I_{tot\_cell} = I_{Na} + I_{CaL} + I_{ha} + I_{K1} + I_{Kr} + I_{Ks} + I_{Kur} + I_{Kto} + I_{bNSC} + I_{NaK} + I_{NCX} \qquad Eq.1$$

The membrane excitation of the model is generated by charging and discharging the membrane

capacitance (C_m) by the net ionic current (I_{tot_cell}) across the cell membrane (Eq 2). The driving force for the

ionic current is given by the potential difference between V_m and the equilibrium potential (E_x) (Eq 3). The net

electrical conductance of the channel is changed by the dynamic changes in the open probability (pO) of the

channel, which is mostly V_m-dependent through the V_m-dependent rate constants (α, β) of the opening and

closing conformation changes of the channel (Eq 4 and Eq 5).

119
$$\frac{dV_m}{dt} = -\frac{I_{tot\_cell}}{C_m} = -\frac{\sum I_x}{C_m}$$
Eq 2

120
$$I_x = \bar{G}_x \cdot pO \cdot (V_m - E_x)$$
Eq 3

121
$$\frac{dpO}{dt} = \alpha \cdot (1 - pO) - \beta \cdot pO$$
Eq 4

122
$$[\alpha\beta]^T = f(V_m)$$
Eq 5

The exchange of 3Na$^+$ / 2K$^+$ by the Na/K pump and the 3Na$^+$ / 1Ca^{2+} exchange by the NCX also

generate sizeable fractions of membrane ionic current, I_{NaK}, and I_{NCX}, respectively. We excluded background

currents of much smaller amplitude, such as I_{KACh}, I_{KATP}, I_{LCCa} and I_{Cab}, from the parameter optimization and

adjusted only the non-selective background cation current (I_{bNSC}) of significant amplitude for the sake of

simplicity [30–32]. The I_{bNSC} is re-defined in the present study as a time-independent net current, which

remained after blocking all time-dependent currents.

3.2.   The computational parameter optimization

The whole cell conductance G_x of a given current system (x) is modified by multiplying the limiting

conductance $\bar{G}_x$ (Eq 3) of the baseline model by a scaling factor sf_x (Eq 6) and are used for the parameter

optimization.

134
$$G_x = \bar{G}_x \cdot sf_x$$
Eq 6

135   The mean square error (MSE) function (Eq 7) was used in the parameter optimization, where $V_{m,a}$

136   represents adaptive V_m (the model output) generated by adjusting the set of sf_x of the baseline model. The target

137    $V_{m,t}$ represents the AP of the intact baseline model.

138
$$MSE = \frac{\sum(V_{m,a} - V_{m,t})^2}{N}$$
Eq 7

The MSE was stabilized by obtaining a quasi-stable rhythm of spontaneous APs through continuous

numerical integration of the model, usually 30 ~ 100 spontaneous cycles were calculated for a new set of sf_xs.

The MSE was calculated within a time window. The width of this time window was adjusted according to the

AP phase of interest. N is the number of digitized V_m points with a time interval of 0.1 ms.

In the usual parameter optimization, the $V_{m,a}$ is generated by modifying the baseline model for

comparison with the experimental record ($V_{m,t} = V_{m,rec}$). However, to evaluate the identifiability of the

parameter optimization, a simple approach was taken in the present study. Namely, we used the manually

adjusted 'cell-specific' model for the target ($V_{m,t}$), which was nearly identical to $V_{m,rec}$. More importantly, the

'cell-specific' V_m is totally free from extra-fluctuations (noise), which were observed in almost all AP

recordings in hiPSC-CMs. In the optimization process, the initial value of each optimization parameter was

prepared by randomizing the set of sf_x of the cell-specific model by $\pm$5~15% at the beginning of each run of

PS ($V_{m,orp}$) in Eq 8 and the PS runs of several hundred were repeated. Thus, the error function is,

151
$$MSE = \frac{\Sigma(V_{m,orp} - V_{m,t})^2}{N}$$
                    Eq 8

We call this optimization method 'orp test' in the present study.

The advantage of using a manually adjusted cell model for the optimization target is that the accuracy

of parameter optimization is proved by recovering all $sf_x = 1$ independently from the randomized initial

parameter set. Note the same approach was used in [23] in evaluating the accuracy of the parameter

optimization by applying the genetic algorithm (GA) to the TNNP model of the human ventricular cell [33].

The optimization of using the randomized initial model parameters were repeated for more than 200

runs. Thus, the method might be classified in a 'multi-run optimization'. The distribution of the sf_x data points

obtained during all test runs was plotted in a single sf_x-MSE coordinate to examine the convergence of

individual sf_xs with the progress of the parameter optimization.

3.3.   The pattern search method for the optimization

For a system showing the relatively simple gradient of MSE along the parameter axis, the gradient-

based optimization methods are more efficient in general than the stochastic methods for this kind of objective

function. We used one of the basic gradient-based optimization methods, the PS program. The computer

program code of the pattern search [34] is simple (see Supplemental Materials) and does not require

derivatives of the objective function. We implemented the code into a homemade program for data analysis (in

VB) to improve the method for better resolution and to save computation time.

The primary PS method uses a base and new points [27]. In brief, sf_x is coded with symbols BP_x and

NP_x in the computer program, representing a base point (BP_x) and a new searching point (NP_x), respectively.

Namely, MSE is calculated on each movement of NP_x by adding or subtracting a given step size (*stp*) *to the*

BP_x, and the search direction is decided by the smaller MSE. Then, the whole mathematical model is

numerically integrated (Eq 2, 3, 4, and 5) using NP_x to reconstruct the time course of AP ($V_{m,a}$). This

adjustment is conducted sequentially for each of the 4~6 selected currents in a single cycle of optimization.

The cycle is repeated until no improvement in MSE is gained by a new set of NP_xs. Then, the BP_x set is

renewed by the new set of NP_x for the subsequent series of optimization. Simultaneously, the *stp* is reduced by

a given reduction factor (*redFct* of 1/4). The individual PS run is continued until the new *stp* becomes smaller

than the critical *stp* (*crtstp*), which is set to 2~10E-5 in the present study.

3.4.   Selection of ionic currents for the optimization

When we get a new experimental record of AP, we do not start the analysis with an automatic

optimization of G_x but first adjust the baseline model by conducting the conventional manual fitting. The nine

ionic currents in Eq 1 in the baseline model are adjusted bit by bit to superimpose the simulated AP on the

experimental one. During this step, it is important to pay attention to the influences of each sf_x adjustment on

the simulated AP configuration on the computer display. Thereby, one may find several key current

components which should be used in the automatic parameter optimization. Usually, currents showing a

relatively large magnitude of G_x were selected for the automatic optimization according to Eq 2, while those

which scarcely modified the simulated AP were left as default values in the baseline model.

Note, the parameters adjusted in the manual fitting are an approximation and are used for the final

automatic parameter optimization using the experimental AP recordings (not shown in the present study). In

the present study, the approximated model is used specifically to prove the feasibility of the algorithm in the

PS parameter optimization in determining the cell-specific computational model.

3.5.   Principal component analysis of the cell-specific models

195   When the orp method is conducted with p elements, it is possible to record the final point BP where

196   the MSE is improved in the p-dimensional space. Suppose we represent the matrix when n data points are

197   acquired as an n x p matrix X. In that case, we obtain a vector space based on the unit vector that maximizes

198   the variance (first principal component: PC1) and the p-dimensional unit vector orthogonal to it (loadings

199   vector $\boldsymbol{w}_{(k)} = (w_1, w_2 \cdots, w_p)$. It is possible to convert each row, $\boldsymbol{x}_{(i)}$ of the data matrix X into a vector of

200   principal component scores, $\boldsymbol{t}_{(i)}$. The transformation is defined by

$$t_{k(i)} = \boldsymbol{x}_{(i)} \cdot \boldsymbol{w}_{(k)} \quad for \quad i = 1,2,\cdots,n \quad k = 1,2,\cdots,p \qquad \text{Eq 9}$$

202   In order to maximize variance, the first weight vector w ((1)) corresponding to the first principal

203   component thus has to satisfy,

$$\boldsymbol{w}_{(1)} = \arg\max_{\boldsymbol{w}} \left\{ \frac{\boldsymbol{w}^T \boldsymbol{X}^T \boldsymbol{X} \boldsymbol{w}}{\boldsymbol{w}^T \boldsymbol{w}} \right\} \qquad \text{Eq 10}$$

205   The k-th component can be found by subtracting the first k-1-th principal components from X

$$\widehat{\boldsymbol{X}_k} = \boldsymbol{X} - \sum_{s=1}^{k-1} \boldsymbol{X} \boldsymbol{w}_{(s)} \boldsymbol{w}_{(s)}^T \qquad \text{Eq 11}$$

207   Then the weight vector is given as a vector such that the variance of the principal component scores is

208   maximized for the new data matrix.

$$\boldsymbol{w}_{(k)} = \arg\max_{\boldsymbol{w}} \left\{ \frac{\boldsymbol{w}^T \widehat{\boldsymbol{X}}_k^T \widehat{\boldsymbol{X}}_k \boldsymbol{w}}{\boldsymbol{w}^T \boldsymbol{w}} \right\} \qquad \text{Eq 12}$$

211   **3.6.   Membrane excitation and its cooperativity with intracellular ionic dynamics**

212   When any of G_xs is modified, the intracellular ion concentrations ($[\text{ion}]_i$) change, although the

variation is largely compensated for with time in intact cells through modification of the activities of both

$3Na^+/2K^+$ pump and $3Na^+/1Ca^{2+}$ exchange (NCX). In the present study, we imitated this long-term

physiological homeostasis of $[\text{ion}]_i$ by introducing empirical Eq 13 and 14. These equations induced 'negative

feedback' to the capacity ($\max I_{NaK}$ and $\max I_{NCX}$) of these ion exchangers. Namely, each correcting factor (crf_x)

was continuously scaled to modify the limiting activity of the transporters to keep the $[\text{Na}^+]_i$ or the total

amount of Ca within the cell ($[Ca_{tot}]_i$) equal to their pre-set level (std_{Nai}, std_{Catot}) with an appropriate delay

(coefficients 0.3 and 0.008 in Eq 13 and Eq 14, respectively).

For the control of $[Na^+]_i$,

221
$$\Delta crf_{NaK} = -crf_{NaK} \cdot (stdNa_i - Na_i) \times 0.3, \qquad stdNa_i = 6.1mM,$$
$$I_{NaK} = (crf_{NaK} \cdot maxI_{NaK}) \cdot vcyc_{NaK} \qquad\qquad Eq\ 13$$

222   For the control of Ca_{tot},

223
$$\Delta crf_{NCX} = -crf_{NCX} \cdot (stdCa_{tot} - Ca_{tot}) \times 0.008, \qquad stdCa_{tot} = 79amol,$$
$$I_{NCX} = (crf_{NCX} \cdot maxI_{NCX}) \cdot (k_1 \cdot E_{1Na} \cdot E_{1NCX} - k_2 \cdot E_{2Na} \cdot E_{2NaCa}) \qquad Eq\ 14$$

The Ca_{tot} is given by $[Ca]_i$ included in the cytosolic three Ca-spaces *jnc*, *iz*, and *blk,* and SR_{up} and SR_{rl}

in the sarcoplasmic reticulum in the free or bound form.

$Ca_{tot\_cell} = [Ca_{tot}]_{jnc} \cdot vol_{jnc} + [Ca_{tot}]_{iz} \cdot vol_{iz} + [Ca_{tot}]_{blk} \cdot vol_{blk} + [Ca_{tot}]_{SRup} \cdot vol_{SRup} + [Ca_{tot}]_{SRrl} \cdot vol_{SRrl}$ Eq 15

Here, the *vol* is the volume of the cellular Ca compartment (see more detail, [11]).

4.   Results

4.1.   Mapping the magnitude of MSE over the nine global parameter space

Parameter identifiability has been one of the central issues in the parameter optimization of biological

models [14,20]. For confirmation of the identifiability of a unique solution using the parameter optimization

method, mapping of the MSE distribution is required over an enlarged parameter space defined by the sf_x of

the nine ionic currents of the baseline model. The randomization of sf_x ranged from 1/10 to ~ 10 times the

default values, and the calculation was performed for ~5,000,000 sets, as shown in Fig 2, where magnitudes of

$log(MSE)$ were plotted against each sf_x on the abscissa.

The data points of MSE at a given sf_x include all variable combinations of the other eight sf_xs. The

algorithm of the PS method searches for a parameter set, which gives the minimum MSE at a given stp through

the process of optimization. Although drawing a clear envelope curve by connecting the minimum MSE at

each sf_x was difficult because of the insufficient number of data points in these graphs (Fig 1), an approximate

envelope of the minimum MSE may indicate a single global minimum of MSE located at the control sf_x equals

one, as typically exemplified by I_{Kr}- and I_{bNSC}-MSE relations. On both sides of the minimum, steep slopes of MSE/sf_x are evident in all graphs. Outside this limited sf_x-MSE area, the global envelope showed a gentle and monotonic upward slope toward the limit on the right side. No local minimum was observed in all of the sf_x-MSE diagrams except the central sharp depression. It was concluded that the theoretical model of cardiac membrane excitation (hVC model) has only a single central sharp depression corresponding to the control model parameter.

Fig 1. Distribution of MSE calculated between the target and the simulated APs modified by randomizing the sfx of 9 ionic
currents in the coordinates of MSE-sfx.
All MSE data points were plotted on the logarithmic ordinate against the linear sfx. A total of 5,141,382 points were calculated
in cell model No.86 over the range of 1/10 ~ 10. Since the configuration of Vm records was largely unrealistic at sfx > 3, MSE
points were cut out over sfx > 3.0. To demonstrate the sharp decrease in MSE near the default sfx, the data points were densely
populated.

4.2.   The prompt necessity for a method of parameter optimization as indicated by hiPSC-CM APs

Fig 2 illustrates records of spontaneous APs (red traces) obtained in 12 experiments in the sequence of

MDP. All experimental records were superimposed with the simulated AP traces (black traces) obtained by the

conventional manual fitting. In most cases, MSE of 1~6 mV^2 remained (Eq 13) at the end of the manual fitting.

This extra component of MSE might be largely attributed to slow fluctuations of V_m of unknown origin

because the non-specific random fluctuations were quite different from the exponential gating kinetics of ion

channels. This extra-noise seriously interfered with the assessment of the accuracy of the parameter

optimization of G_x in the present study. Thus, APs produced by the manual adjustment ('cell specific model')

was used as the target AP, which were completely free from the extra noise when examining the feasibility of

the parameter optimization algorithm.

272   A comparison of AP configurations between these hiPSC-CMs clearly indicated that the

classification of these APs into atrial-, ventricular- and nodal-types was virtually impractical, as described

in [7]. On the other hand, if provided with the individual models fitted by an objective parameter

optimizing tools using the baseline model (black trace), the results should be fairly straightforward not

only in estimating the functional expression level of ion channels but also in clarifying the role of each

current system or the ionic mechanisms in generating the AP configuration in a quantitative manner.

Thus, the objective parameter optimization of the mathematical model is a vital requirement in cardiac

electrophysiology.

Fig 2. The manual fitting of variable AP configurations in different hiPSC-CMs.
Each panel shows the experimental record (red) superimposed by the model output (black) of the baseline
model by the conventional manual fitting. At the top of each pair of AP records, the experimental cell number
is presented. The extra fluctuations are obvious during the AP plateau in Cells 78, 08 and 01, while in Cells 15
and 74 during SDD. The length of abscissa is markedly different to illustrate the interval between two
successive peaks of AP.

Table 1. AP metrics recorded from hiPSC-CMs

	CL (ms)	OS (mV)	MDP (mV)	APD(ms) at -20mV	MSE (mV²)
Cell 78	983.8	29.6	-87.4	271.7	5.8443
Cell 86	1326.0	29.7	-85.0	289.2	4.0554
Cell 01	887.4	31.2	-82.2	435.0	3.9330
Cell 91	1058.0	33.0	-80.4	308.6	7.2156
Cell 08	551.4	32.0	-79.5	287.6	1.4043
Cell 11	695.0	25.3	-77.6	243.5	2.6683
Cell 02	603.9	26.4	-74.9	173.4	1.0412
Cell 74	622.8	18.5	-74.8	157.0	2.2589
Cell 10	564.3	23.0	-73.7	220.9	3.2194

Cell 38	425.4	24.2	-66.8	123.4	3.6626
Cell 15	239.5	13.8	-66.1	57.1	2.8607
Cell 12	458.6	19.7	-61.5	119.0	1.3514

Table 1 indicates the AP metrics; the cycle length (CL), the peak potential of the plateau (OS), the maximum diastolic potential
(MDP), and the AP duration measured at -20 mV in addition to the MSE between individual experimental record and the model
output fitted by manual fitting. The CL, MDP and AP were very variable among different AP recordings of cells shown in Fig 1.
The cells were arranged by the sequence of MDP. The manual fitting of varying AP configurations in different hiPSC-CMs.

Table 1 indicates the AP metrics; the cycle length (CL), the peak potential of the plateau (OS), the

maximum diastolic potential (MDP), and the AP duration measured at -20 mV in addition to the MSE between

individual experimental record and the model output fitted by manual fitting. The CL, MDP and AP were very

variable among different AP recordings of cells shown in Fig 1. The cells were arranged by the sequence of

MDP.

4.3.   Feasibility of the PS algorithm for parameter optimization of membrane excitation models

The automatic parameter optimization was applied to the model of cardiac membrane excitation

in a limited number of studies (for review, see [23,26,35,36]) using various optimization

methods, such as genetic algorithms. To the best of our knowledge, the principle PS algorithm

has not been successfully applied to the detailed mathematic models of cardiac membrane

excitation composed of both ionic channel and ion transporters models, except for the pioneering

work in [12], which applied more general gradient-based optimization method to the simple

308   ventricular cell model of Beeler and Reuter (BR model)[37].

Fig 3 shows a typical successful run of the new PS method in a hiPSC-CM, which showed an MDP

of ~-85 mV. The PS parameter optimization was started after randomizing the sf_xs of the major six currents,

I_{Kr}, I_{CaL}, I_{Na}, I_{ha}, I_{K1} and $I_{bNSC,}$ in the manual fit model within a range of $\pm 15\%$ around the default values

(normalized magnitude of 1). Fig 3A compares the simulated $V_{m,rmp}$ (black trace) with the target $V_{m,t}$ (red) (Eq

8). The OS, APD as well as the CL of spontaneous AP were markedly different at the first cycle of AP

reconstruction (Fig 3A-1). These deviations were largely decreased at the PS cycle (Fig 3A-2, at N = 50), and

became invisible in the final result (Fig 3A-3, N = 1167). The final individual current flow of nine current

components are demonstrated in the lower panel of Fig 3A-3 (I_m).

Fig 3. Results of the successful optimization in a cell (Cell86).
(A-1) Target AP (red line) and AP generated by randomized initial sf$_x$s (black line). (A-2) AP generated after 50 cycles of
adjusting BP (black line). (A-3) AP generated by the final sf$_x$s. I_m: shows corresponding time courses of each current for the
finalized AP shown in A-3. (B-1) Changes in sf$_x$s vs. log(MSE) during the successful optimization process of PS. (B-2)
Log(MSE) of all BP points during the search process in PS. The initial values of sf$_x$ are plotted at the top of each BP-log(MSE)
graph.
325

The time course of decreasing $log(MSE)$ evoked by the multi-run PS optimization is plotted for each

sf$_x$ in Fig 3B-1 every time of resetting the set of base points. Fig B2 shows all of the $log(MSE)$ obtained at

every adjustment by stepping individual BP points. The movement of all sf$_x$ were synchronized to decrease

$log(MSE)$ from ~2.4 to 1 during the initial 180 cycles of decreasing $log(MSE)$, but the search directions of

BP were quite variable. The detailed adjustment of sf$_x$s below $log(MSE) < 0$ was driven by adjusting I_{Kr}, I_{CaL}

and I_{bNSC} in this cell. The values of sf$_{Kr}$, sf$_{CaL}$ and sf$_{Na}$ approached to correct value 1, while those for I_{ha}, I_{K1} and

I_{bNSC} remained deviated from the unit by less than 10%. The explanation for the deviation of these three sf$_x$s

from the unit will be examined in the next section of Results.

4.4. The six-parameter multi-run PS method successfully determined the parameters of membrane excitation models

In individual runs, the PS optimization was frequently interrupted at intermediate levels during the

progress of optimization and the probability of reaching $log(MSE)$, for example, below -2, rapidly decreased

with increasing extent of the randomization of the initial set of parameters. Moreover, the complementary

relations between several ionic currents in determining dV_m/dt might have hampered the parameter

optimization. These facts indicate the requirement of statistical measures to improve the accuracy of the PS

method. Fig 4 shows the results of multi-run orp tests, in which the optimization shown in Fig 3 was repeated

several hundred times, and all results were plotted in a common coordinate of $log(MSE)$ and individual sf_xs.

The population of sf_x correctly converged at a single peak point very close to 1 with increasing negativity of

$log(MSE)$ for sf_{Kr}, sf_{CaL}, and sf_{Na}, while sf_{ha}, $sf_{K1,}$ and sf_{bNSC} showed obvious variance. Nevertheless, they also

showed a clear trend toward convergence to 1 in the average.

Fig 4. Convergence of sf_x in the multi-run orp test for Cell86.
The ordinate is the *log(MSE)* and the abscissa is the normalized amplitude of sf_x; *x* stands for *Kr, CaL, Na, ha, K1*, and *bNSC*.
Black points were obtained in the progress of optimization, and red ones are the final points in 829 runs of PS pattern search.

Table 2 summarizes the mean of sf_x determined for the top 20 runs of the PS parameter optimization in

each of the 12 cells illustrated in Fig 2. The [Na⁺] as well as [Ca²⁺$_{tot}$]$_i$ was well controlled to the reference

levels (stdNai, and stdCatot in Eq 16 and Eq 17) of 6.1 mM and 79 atto-mole, respectively, at the end of the

parameter optimization to ensure the constant [Na⁺] as well as [Ca²⁺$_{tot}$]$_i$. The mean of final $log(MSE)$ =

-2.74 indicates that the MSE was reduced by five orders of magnitude from the initial level by the multi-run

PS optimization method, like in the successful example shown in Fig 3B. The mean of individual sf_xs were

very close to 1 with a minimum standard error (SE) of mean, which were less than 1% of the mean, even for

I_{K1}, I_{bNSC} and I_{ha}, which showed weak convergence of the $log(MSE)$. These results well validate the accuracy

of the parameter optimization using the multi-run PS method in all of 12 cell-specific models, which showed

the large variety of spontaneous AP recorded in the hiPSC-CMs.

Table 2. Measurements of sfxs (mean + S.E n = 20), [Na+]$_i$, mM and [Ca]$_{tot}$ in atto-mole in the 12 cells.

Cell No.	log(MSE)	sfKr	sfK1	sfCaL	sfbNSC	sfha	sfNa	sfKur	[Na$^+$]$_i$(mM)	Ca$_{tot}$(amol)
78	-2.48321	1.00005	1.00157	1.00037	0.99460	1.00060	1.00134		6.10550	78.99979
91	-2.42008	0.99952	1.00644	1.00063	1.00280	1.00470	1.00068		6.09977	79.00044
86	-2.80257	1.00166	1.01394	1.00142	1.02670	1.00253	1.00702		6.09466	79.00008
01	-2.79709	0.99871	1.00157	0.99756	0.99692	1.00054	0.99779		6.08973	78.99984
08	-3.07432	0.00094	0.99982	1.00088	1.00041	0.99968	0.99985		6.12201	79.00056
11	-2.67641	1.00186	1.00686	1.00129	0.99768	1.00253	1.01028		6.10385	78.99995
10	-1.70278	1.00322	1.01081	1.00424	1.00396		0.99883		6.10968	79.00018
02	-2.35441	1.00161	1.02038	1.00341	0.99815	1.01324	1.00954		6.10184	79.00004
74	-2.43399	1.00126	1.01838	1.00308	0.99898	1.00004	1.00435		6.10118	79.99979
38	-3.01883	1.00075		1.00106	1.00061		0.98866	1.00151	6.10530	78.99969
15	-3.85992	1.00003		0.99894	0.99996		1.00015	0.98653	6.09902	79.00022
12	-3.33037	0.99978		1.00030	0.99990	0.97587		1.00188	6.10012	79.00007
Ave	-2.74617	0.99992	1.00886	1.00110	1.001723	0.99997	1.001681	0.99664	6.10272	79.08339
SE	0.07065	0.00093	0.010000	0.00164	0.00430	0.00729	0.00544	0.00690	0.00017	0.000345

The top 20 results obtained in the multi-run orp method were analyzed in each cell. Gland average and SE are
listed at the bottom rows.

4.5.   Complementary relationship among I_{K1}, I_{ha} and I_{bNSC}

Fig 5A illustrates the distribution of sf_xs amplitude in the top 20 data points, where the final sf_xs in

individual runs were connected with lines for each run of PS in Cell 86 (Fig 2). The values of standard error

(SE) of mean were quite small in the sf_{Kr} and $sf_{CaL,}$ less than 1%. In contrast, sf_{ha}, sf_{K1} and sf_{bNSC} showed

evidently larger deviation. This finding is interesting since the former currents are mainly involved in

determining the AP configuration and the latter group mainly in driving the relatively long-lasting SDD of

approximately 1 sec in duration.

Thus, we analyzed the distribution of sf_{ha}, sf_{K1} and sf_{bNSC} within the top 20 MSE. Fig 5C and D show

the distribution of sf_x points in the space of the three sf_x dimensions. In Fig 5C, the 20 data points seemed to be

dispersed randomly in the parameter space, but when the space was rotated to a specific angle, a linear

distribution was observed as in Fig 5D, indicating that the points are distributed approximately on a plane

surface in the 3D space. Using the multiple regression analysis, we could obtain an equation that fits the 20

data points as follows (R^2=0.872);

$$0.762 \cdot sf_{ha} - 0.619 \cdot sf_{K1} + 0.191 \cdot sf_{bNSC} = 0.333554 \qquad\qquad Eq\ 16$$

By replotting the data points in the 2D space with the abscissa for the sum of two inward-going

currents (0.76 sf_{ha} + 0.19 sf_{bNSC}) and the ordinate for the outward current 0.62 sf_{K1}, we obtained a regression

line as shown in Fig 5B. The close correlations among the three sf_xs were indicated with a quite large R^2 of

0.941. This finding well confirms that the three currents have complementary relations with each other to give

virtually identical configurations of spontaneous AP. In other words, $log(MSE)$ remains nearly constant as far

as the composition of the currents satisfies the relationship given by Eq 16.

Fig 5. Distribution of sf_x within the top 20 sets of sf$_x$s obtained from the multi-run orp test in Cell86 in Fig 3.
Data points of normalized sf_x in each set were depicted in a different color. (A) amplitudes of each sf_x (indicated on the abscissa)
were plotted. (C) Three parameters, sf_{ha}, sf_{K1}, and sf_{bNSC} were plotted in the 3D plot. (D) A different solid angle view of the 3D
plot showed a linear correlation; see text for the plot in (B)
394

The complementary relationship was further examined by conducting the orp test after fixing one of

the two factors, sf_{K1} or (sf_{ha} + sf_{bNSC}), illustrated in Fig 5B. Fig 6A shows the $log(MSE)$ vs. sf_{K1} relation when

the (sf_{ha} + sf_{bNSC}) were fixed at the values obtained by the multi-run orp test. Indeed, the typical convergence of

the $sf_{K1}$was obtained. Alternatively, if the sf_{K1} was fixed, the convergence was obviously improved for both sf_{ha}

and sf_{bNSC} (Fig 6B-1, 2), but it was less sharp if compared to sf_{Kr}, sf_{CaL} and sf_{Na} (not shown, but refer to

corresponding results in Fig 4A). This finding was further explained by plotting the relationship between the

two inward currents, I_{ha} and $I_{bNSC,}$ as illustrated in Fig 6C. The regression line for the data points was fitted by

Eq 17 with R^2 = 0.86, supporting the complemental relationship between the two inward currents, I_{ha} and I_{bNSC}.

403                         $$0.8098 \cdot sf_{bNSC} + 0.1897 \cdot sf_{ha} = 1 \qquad\qquad Eq\ 17$$

405   The moderately high R^2 indicates that the SDD is determined not only by the major I_{ha} and I_{bNSC} *but*

406   also by other currents, such as I_{K1}, I_{Kr}, the delayed component of I_{Na} (I_{NaL}) and I_{CaL}, which were recorded

407   during the SDD as demonstrated in Fig 2.

408   Essentially the same results of complementary relationship among sf_{ha}, sf_{bNSC} and sf_{K1} were obtained in

409   Cell 91, which also showed the long-lasting SDD with the very negative MDP as in Cell 86, as shown in Fig 2

410   and Table 2. The regression relation for the data points was fitted by Eq 18 and Eq 19 with R^2 = 0.656 and

411   0.472, respectively.

412                    $$0.572 \cdot sf_{ha} - 0.132 \cdot sf_{K1} + 0.810 \cdot sf_{bNSC} = 1.25891 \qquad\qquad Eq\ 18$$

413                         $$0.5238 \cdot sf_{bNSC} + 0.4715 \cdot sf_{ha} = 1.3415 \qquad\qquad Eq\ 19$$

Fig 6. The complementary relations among sf_{K1}, sf_{ha} and sf_{bNSC}.
(A) and (B) results of the multi-run orp test. A; the perfect convergence of sf_{K1} when sf_{ha} and sf_{bNSC} were fixed. (B1) improved
convergence of sf_{ha} and (B2) sf_{bNSC} when sf_{K1} was fixed. In these two orp tests, sf_x of other currents showed quite comparable
convergence as in Fig 4A. (C) the correlation between sf_{ha} and sf_{bNSC}.

4.6.   Principal components in the hiPSC-CM model

The pattern search frequently got stuck during the progress of parameter optimization and failed to

find the global minimum in the present study (Fig 5 and Fig 6). The major cause of this interruption may most

probably be attributed to the fact that sf_xs were used directly as the search vector of the pattern search. In

principle, the algorithm of PS parameter optimization gives the best performance when the parameters search

is conducted in orthogonal dimensions where each dimension does not affect the adjustment of other sf_x [28].

To get deeper insights, we applied the principal component (PC) analysis to the set of 6 sf_xs selected in the

baseline model. We performed principal component analysis on the data points recorded in the vicinity of the

minima (using the top 20 data).

As illustrated in Fig 7, each of the 6 PCs was not composed of a single sf_x but mostly included

multiple sf_x sub-components, except the first PC, which is virtually dependent only on sf_{bNSC}. This finding

indicates the inter-parameter interactions during the process of parameter optimization. For example, any

alteration of sf_{K1} or sf_{ha} should affect PCs No. 2~5, while the changes in sf_{CaL} or sf_{Na} simultaneously affect

No.3, 4, 6 or 3, 4, 5 PCs, respectively. sf_{Kr} shows a broad range of interactions. It might be concluded that the

frequent interruptions of PS parameter optimization are most probably caused by the sporadic appearance of

the local minimum of MSE through interactions among sf_xs.

Fig 7. Principal components (PCx) to describe the *sf_x* distribution mathematically in the *sf_x-log*(MSE) relations in Fig 2.

Results were obtained in Cell 86. The magnitudes of PCx were normalized to give a unit magnitude in the total. Note each PC is composed of multiple components of ionic current, which are indicated in the Index with corresponding colors. Principal component analysis was performed on the data population of the top 50 runs of the PS analyses, which showed good optimization results.

5.   Discussion

New findings in the present study are listed below.

(1) Mapping the MSE distribution over the enlarged parameter space was conducted by randomizing

the G_x of the baseline model. It was confirmed that the baseline model has only a single sharp

depression of MSE at the default G_xs (Fig 1).

(2) The preliminary cell-specific models were firstly prepared by the conventional manual tuning of

G_x to superimpose the model output on each of twelve experimental AP recordings (Fig 2).

Thereby, the parameter search space was restricted to a relatively small space to facilitate

parameter optimization.

(3) The sf_xs of 4~6 G_x parameters were initially assigned random values from a uniform distribution

ranging between $\pm10\%$ of default values. The MSE was calculated between the randomized

model output and the intact model AP as the target of optimization (Fig 3).

(4) Plotting parameter sf_x in a common sf_x - MSE coordinates during each run of several hundred runs

of optimization (Fig 4), we found that the sf_x distribution of I_{Kr}, I_{CaL}, and I_{Na} converged sharply to

a single point with decreasing MSE, which exactly equal the default ones. On the other hand,

estimates of sf_{K1}, sf_{ha} and sf_{bNSC} deviated slightly within a limited range around the default values

in cells showing long-lasting SDD (Fig 4).

(5) For statistical evaluation, the mean $\pm$SE of sf_x in the top 20 estimates of MSE was calculated in

467   individual cells (Table 2). The results of the parameter optimization in the 12 cells definitely

468   indicated that the means of sf_xs were very close to 1.00, with the S.E less than 0.01 for all G_xs.

469   (6) A complementary relationship was found between sf_{K1}, sf_{ha} and sf_{bNSC} in determining the gentle

470   slope of long-lasting SDD in two representative cells (Fig 5). Supporting this view, the sf_{K1} clearly

471   focused on the unit provided that sf_{ha} and sf_{bNSC} were fixed and vice versa (Fig 6).

472   (7) The nine search vectors of sf_x of the presented model could be replaced by the same number of

473   theoretical PCs, and each PC was mostly composed of multiple sf_xs (Fig 7). This finding

474   definitely supports the view [12] that the complex interactions among I_xs might interrupt the

475   progress of the parameter optimization when sf_xs were used as the search vector instead of using

476   theoretical orthogonal ones.

The use of an initial randomized set of parameters was crucial in examining if an optimization method

can determine unique estimates independent from the initial set of parameters, as used in the GA-based method

for determining the G_xs of the mathematical cardiac cell model [23]. The findings listed above well confirmed

the feasibility of the PS method. Most probably, the PS method is applicable to variable mathematical models

of other cell functions as well. See [26] for a more systematic review of the parameter optimization in the

cardiac model development.

It has been suggested that different combinations of parameters may generate simple outputs that are

very similar [12,23–25]. In the present study, this notion may be explained at least in part by the

complementary relationship, for example, between the I_{K1}, I_{ha} and I_{bNSC} in determining dV_m/dt of SDD, which

is a function of the total current (Eq 2, Fig 5, and Fig 6). The gradient-based optimization method relies on the

precise variation in the time course of dV_m/dt induced by the time-dependent changes in individual sf_xs (Eq 2).

Therefore, the MSE was calculated over the whole time course of the spontaneous APs. Note, we did not use

the AP metrics, which reflect only indirectly the kinetic properties of individual currents. Even with this

measure of calculating the MSE, the time-dependent changes in pO (Eq 3) might be relatively small between

two major currents, I_{K1} and I_{ha}, in comparison to I_{bNSC}, which has no V_m-dependent gate during the SDD as

shown in the current profile Fig 3A-3. We assume that the gradient-based optimization method will be able to

determine different contributions of individual currents if the optimization is conducted only within a selected

time window of SDD. If MSE is calculated over multiple phases of the spontaneous AP, the influence of a

particular phase on the MSE should be diluted. In our preliminary parameter optimization, this problem was

partly solved by using a weighted sum for different phases of the spontaneous AP in summing up the MSE.

The small amplitude of a given current might be an additional factor in the weak convergence of sf_x

observed in the diagram of sf_x - MSE in the orp method of optimization. If the current amplitude was much

smaller in reference to the sum of all currents in determining dV_m/dt (Eq 2), the resolution of the PS method

would get lower. [24] demonstrated that the model output, for example, the AP plateau phase were almost

superimposable when the different ratio of G_{Kr} and G_{pK} were used in reconstructing the model output (Fig

1)[24]. They described that the AP metrics used for comparisons, such as APD, OS and APA seemed quite

similar. It should be noted, however, that the results were obtained by applying different combinations of sf_x to

the same TNNP model [33]. This means that the relative amplitudes of I_{Kr} and I_{pK} in the TNNP model were

much smaller than the major I_{CaL} during the AP plateau, even though I_{Kr} and I_{pK} have totally different gating

kinetics. Thus, the results of parameter optimization should be model-dependent. The same arguments will

also be applied to the use of FR guinea pig model [38] in [23].

The gradient-based parameter optimization method was applied to the cardiac model of membrane

excitation in [12], which analyzed the classic BR model[37]. The whole cell current in the BR model was

composed of a minimum number of ionic currents, a background I_{K1}, and three time-dependent currents; I_{Na}, I_s,

and I_{x1}, which were dissected from the voltage clamp experiments by applying the sucrose gap method to the

multicellular preparation of ventricular tissue. The gating of the latter three currents was formulated according

to the Hodgkin-Huxley type gating kinetics, which was quite simpler if compared with the recent detailed

description of the ionic currents. They described that the parameter optimization was difficult if the AP

configuration was used as the target of the parameter optimization, and they used the time course of the whole

cell current as of the target of parameter optimization. However, the number of parameters was quite large, 63

in total, including limiting conductances and the gating kinetics. They suggested the feasibility of the

parameter optimization method will be improved if provided with additional experimental data.

In the modern mathematical cardiac cell models, most ionic currents were identified by the whole-cell

voltage clamp and single channel recordings in dissociated single myocyte [39] using the patch clamp

technique [40] and by identifying their molecular basis of membrane protein. It has been clarified that the

molecular basis of the ion channels expressed in the hiPSC-CMs is mostly identical to in the adult cardiac

myocytes rather than in the fetal heart (GSE154580 GEO Accession viewer (nih.gov)). Moreover, the gating

kinetics have been much detailed to characterize the ionic currents within the cell model. In principle, the

detailed characterization of individual currents should facilitate the identifiability of the model parameter but

should not necessarily interfere with parameter optimization. We consider that the manual fitting of the model

parameters to the AP recording by using a priori knowledge of biophysical mechanisms should largely

facilitate the subsequent automatic parameter optimization. It might also be noted that the ionic currents left at

the default values work as a kind of constraint to improve the identifiability of the target parameters.

After validating the automatic parameter optimization method, the final goal of our study is to find the

principle of ionic mechanisms, which are applicable to the full range of variations of spontaneous AP records

in both hiPSC-CMs and matured cardiomyocytes. For this purpose, we will apply the multi-run PS method to

the experimental AP recordings using the initial parameter sets obtained by the conventional manual fit. The

protocol of measuring the G_xs will be the same as used in the present study except for the use of experimental

AP recordings in place of the output of the 'cell-specific model. In our preliminary analysis, the magnitude of

individual model parameters obtained by the manual tuning was corrected by less than ~15% by the objective

parameter optimization. Finally, the ionic mechanisms underlying the SDD of variable time courses will be

analyzed in a quantitative manner, for example, by using the lead potential analysis [41], which explains

changes in V_m in terms of G_x of individual currents.

Limitations

In general, obvious limitations of the mathematical models of cardiac membrane excitation so far

published are caused by a shortage of functional components inherent in intact cells. For example, the $[ATP]_i$

controlled by energy metabolism is a vital factor in maintaining the physiological function of ion channels as

well as the active transport Na^+/K^+ pump [42]. Moreover, the followings are still not implemented in most

models; the modulation of the ion channel activity through phosphorylation of the channel proteins, detailed

modulation of the channel by the intracellular $[Ca^{2+}]_i$, the alterations of ion channel activity by PIP_2 [43,44]

and by the tension of the cell membrane through the cell volume change [45–48]. The detailed Ca^{2+} dynamics

of the intracellular $[Ca^{2+}]_i$ are still not implemented in most of the cardiac cell models; such as the Ca^{2+} release

from SR activated through the coupling of a few L-type Ca^{2+} channels with a cluster of RyRs at the dyadic

junction [49], the Ca^{2+} diffusion influenced by the Ca^{2+}-binding proteins [50]. To simulate the Ca^{2+}-binding to

troponin during the development of the contraction, it is necessary to include a dynamic model of contracting

fibers [51–54]. These limitations should be thoroughly considered when pathophysiological phenomena, such

as arrhythmogenesis are concerned. The scope of the present study is limited to the AP configurations of

hiPSC-CMs, which were assumed to be 'healthy' with respect to the above concerns; for example, $[ATP]_i$,

$[Na^+]_i$ and $[Ca_{tot}]_i$ were kept constant, and the standard contraction model was implemented in the hVC model.

The parameter optimization presented in this study could be achieved in a practical way by limiting

the number of unknown parameters. We determined only G_xs based on the assumption that ion channel

kinetics are preserved as the same in the hiPSC-CMs as in the matured myocytes. Usually, 4~6 ionic currents

were selected for the optimization. We found that the multi-run orp method could be performed simultaneously

for all nine ionic currents described in Eq 1. However, the computation time was radically prolonged, and the

resolution was not as high as obtained by using the modest number of parameters. We consider that the

determination of the limited number of G_xs is quite relevant to solving physiological problems in terms of

detailed model equations for each current system.

Although I_{NCX} and I_{NaK} contribute sizeable fractions of the whole-cell outward and inward currents,

respectively (Fig 3A-3), we excluded the scaling factors, sf_{NaK} and sf_{NCX} from the parameter optimization for

the sake of simplicity. Instead, the possible drift of the intracellular ion concentrations was virtually fixed

during the repetitive adjustment of ionic fluxes by varying sf_x as shown in Table 2. The introduction of the

empirical equations (Eq 13 and Eq 14) was quite useful to adjust the $[Na^+]_i$ and $[Ca_{tot}]_i$ (Table 2) so that the

time course as well as magnitude of I_{NCX} remained almost constant during the parameter optimization. When

influences of varying $[Na^+]_i$ and/or $[Ca_{tot}]_i$ are examined under various experimental conditions in future, the

reference levels of $[Na^+]_i$ and/or $[Ca_{tot}]_i$ (*stdNai* and *stdCatot* in Eq 13 and 14) might be replaced by

experimental measurements.

The parameter optimization method was not applied to several ionic currents. For example, it was

difficult to determine the kinetics of T-type Ca^{2+} channel (Ca_V 3.1) and excluded in the present study. A very

fast opening and inactivation rates described in [55] suggest a complete inactivation of I_{CaT} over the voltage

range of SDD, while a sizeable magnitude of window current described in [56] suggests a much larger

contribution to SDD. The kinetics of I_{CaT} (Ca_V 3.1) still remain to be clarified in experimental examinations.

The sustained inward current, I_{st}, is recently attributed most probably to the Ca_V 1.3 [57], which is activated at

a more negative potential range than the activation of I_{CaL} (Ca_V 1.2) [58,59]. The I_{bNSC} was used to represent

net background conductance in the present study. However, several components of the background

conductance have been identified on the level of molecular basis in matured myocytes (see for review TRPM4,

[60]). Experimental measurements of the current magnitude for each component are also awaited.

Gábor and Banga indicated that the multi-run method had shown good performance in certain cases,

especially when high-quality first-order information is used and the parameter search space is restricted to a

relatively small domain [16] (see also [19]). Indeed, the manual fitting of the parameters (Fig 1) was required

to utilize the presented multi-run PS method over the restricted search space. One of the major difficulties in

the manual fitting of individual G_xs arose during the SDD, where I_{Kr}, I_{K1}, I_{bNSC}, and I_{ha}, in addition to I_{NaK} and

I_{NCX} constitute the whole-cell current (Fig 3A-3). Close inspection of the current components (Fig 3A-3),

however, suggests hints of how to do with the manual fit. The transient peak of I_{Kr} dominates the current

profile during the final repolarization phase from -20 to -60 mV in all 12 hiPSC-CMs [61], since I_{CaL} and I_{Ks}

rapidly deactivated before repolarizing to this voltage range. The I_{NaK} and I_{NCX} are well controlled by the

extrinsic regulation in Eq 6 and Eq 7. Thus, the manual fitting of sf_{Kr} is firstly applied to determine sf_{Kr}. The

MDP more negative than -70 mV is adjusted by the sum of time-dependent ($I_{Kr} + I_{K1}$) and the time-

independent I_{bNSC}. Then, I_{Kr} is deactivated when depolarization becomes obvious after the MDP, and gradual

activation of I_{ha} and the depolarization-dependent blocking of I_{K1} by the intracellular substances [62] take the

major role in promoting the initial linear phase of SDD. Thus, the amplitude of sf_{K1} and sf_{bNSC} might be

approximated during the initial half of SDD. The late half of SDD, including the foot of AP, namely the

exponential time course of depolarization toward the rapid rising phase of AP, is mainly determined by the

subthreshold V_m-dependent activation of I_{Na} (after MDP more negative than -70 mV) and/or I_{CaL} (after MDP

less negative than -65 mV). Thus, the sf_{Na} and sf_{CaL} are roughly determined by fitting the foot of AP and the

timing of the rapid rising phase of AP. The plateau time course of AP is determined by sf_{CaL} and the factor of

Ca^{2+}-mediated inactivation of I_{CaL} (the parameter KL, [4]). Since the kinetics of outward currents, I_{Kur}, I_{Kto}

(endo-type), and I_{Ks} are quite different from that of I_{Kr}, the plateau configuration is determined bit by bit by

adjusting these currents. We failed to observe the phase 1 rapid and transient repolarization in the hiPSC-CMs

(Fig 2), which is the typical sign of the absence of epicardial-type I_{Kto}.

In hiPSC-CMs showing less negative MDP than ~-65 mV, the contribution of I_{K1}, I_{Na} and I_{ha} should

be negligibly small because I_{K1} is nearly completely blocked by the intracellular Mg^{2+} and polyamine, I_{Na}

inactivated, and I_{ha} deactivated during SDD, even if any expressed.

Nevertheless, parameter optimization might be laborious and time-consuming for those who are not

familiar with the electrophysiology of the cardiac myocyte. This difficulty might be largely eased by

accumulating both AP configuration and the underlying current profile obtained in the parameter optimization

into a database in the future. If this database becomes available, the computer may search for several candidate

APs for the initial parameter set, which is used for automatic parameter optimization.

6.   Funding and financial conflicts of interest

The authors declare that the research was conducted without any commercial or financial relationships

that could be construed as a potential conflict of interest.

This work was supported by JSPS KAKENHI (Grant-in-Aid for Young Scientists) Grant Numbers

JP19K17560 for HK, 16K18996 for YH, and 21K06781 for FT.

7.   Author Contributions

625   HK, YH, DY, YW, AK, and TM performed the wet experiments and analyzed them. HK, SK, YH,

626   YZ, FT, AA, and AN developed the simulation model and the parameter optimization method. HK, YH, AA

627   and AN wrote the manuscript. All authors reviewed the manuscript. AA and TK organized the research team.

8.   Acknowledgments

630   The authors would like to thank our laboratory colleagues for their valuable comments and

631   discussions.

9.   References

1. Noble D, Garny A, Noble PJ. How the Hodgkin-Huxley equations inspired the Cardiac Physiome Project:
Hodgkin-Huxley equations and the cardiac Physiome Project. J Physiology. 2012;590: 2613–2628.
doi:10.1113/jphysiol.2011.224238

2. Noble D, Rudy Y. Models of cardiac ventricular action potentials: iterative interaction between experiment
and simulation. Philosophical Transactions Royal Soc Lond Ser Math Phys Eng Sci. 2001;359: 1127–1142.
doi:10.1098/rsta.2001.0820

3. Winslow RL, Cortassa S, O'Rourke B, Hashambhoy YL, Rice JJ, Greenstein JL. Integrative modeling of the
cardiac ventricular myocyte. Wiley Interdiscip Rev Syst Biology Medicine. 2011;3: 392–413.
doi:10.1002/wsbm.122

4. Hinch R, Greenstein JL, Tanskanen AJ, Xu L, Winslow RL. A Simplified Local Control Model of Calcium-
Induced Calcium Release in Cardiac Ventricular Myocytes. Biophys J. 2004;87: 3723–3736.
doi:10.1529/biophysj.104.049973

5. Paci M, Hyttinen J, Aalto-Setälä K, Severi S. Computational models of ventricular- and atrial-like human
induced pluripotent stem cell derived cardiomyocytes. Ann Biomed Eng. 2013;41: 2334–2348.
doi:10.1007/s10439-013-0833-3

6. Paci M, Hyttinen J, Rodriguez B, Severi S. Human induced pluripotent stem cell-derived versus adult
cardiomyocytes: an in silico electrophysiological study on effects of ionic current block. Brit J Pharmacol.
2015;172: 5147–5160. doi:10.1111/bph.13282

7. Lei CL, Wang K, Clerx M, Johnstone RH, Hortigon-Vinagre MP, Zamora V, et al. Tailoring Mathematical
Models to Stem-Cell Derived Cardiomyocyte Lines Can Improve Predictions of Drug-Induced Changes to
Their Electrophysiology. Front Physiol. 2017;8: 986. doi:10.3389/fphys.2017.00986

8. Grandi E, Pasqualini FS, Bers DM. A novel computational model of the human ventricular action potential
and Ca transient. Journal of molecular and cellular cardiology. 2010;48: 112–121.
doi:10.1016/j.yjmcc.2009.09.019

9. O'Hara T, Virág L, Varró A, Rudy Y. Simulation of the Undiseased Human Cardiac Ventricular Action
Potential: Model Formulation and Experimental Validation. McCulloch AD, editor. Plos Comput Biol. 2011;7:
e1002061-29. doi:10.1371/journal.pcbi.1002061

10. Asakura K, Cha CY, Yamaoka H, Horikawa Y, Memida H, Powell T, et al. EAD and DAD mechanisms
analyzed by developing a new human ventricular cell model. Prog Biophysics Mol Biology. 2014;116: 11–24.
doi:10.1016/j.pbiomolbio.2014.08.008

11. Himeno Y, Asakura K, Cha CY, Memida H, Powell T, Amano A, et al. A Human Ventricular Myocyte
Model with a Refined Representation of Excitation-Contraction Coupling. Biophys J. 2015;109: 415–427.
doi:10.1016/j.bpj.2015.06.017

12. Dokos S, Lovell NH. Parameter estimation in cardiac ionic models. Prog Biophysics Mol Biology.
2004;85: 407–431. doi:10.1016/j.pbiomolbio.2004.02.002

13. Dutta S, Chang KC, Beattie KA, Sheng J, Tran PN, Wu WW, et al. Optimization of an In silico Cardiac
Cell Model for Proarrhythmia Risk Assessment. Front Physiol. 2017;8: 616. doi:10.3389/fphys.2017.00616

14. Whittaker DG, Clerx M, Lei CL, Christini DJ, Mirams GR. Calibration of ionic and cellular cardiac
electrophysiology models. Wiley Interdiscip Rev Syst Biology Medicine. 2020;12: e1482.
doi:10.1002/wsbm.1482

15. Cairns DI, Fenton FH, Cherry EM. Efficient parameterization of cardiac action potential models using a
genetic algorithm. Chaos Interdiscip J Nonlinear Sci. 2017;27: 093922. doi:10.1063/1.5000354

16. Gábor A, Banga JR. Robust and efficient parameter estimation in dynamic models of biological systems.
Bmc Syst Biol. 2015;9: 74. doi:10.1186/s12918-015-0219-2

17. Degasperi A, Fey D, Kholodenko BN. Performance of objective functions and optimisation procedures for
parameter estimation in system biology models. Npj Syst Biology Appl. 2017;3: 20. doi:10.1038/s41540-017-
0023-2

18. Penas DR, González P, Egea JA, Doallo R, Banga JR. Parameter estimation in large-scale systems biology
models: a parallel and self-adaptive cooperative strategy. Bmc Bioinformatics. 2017;18: 52.
doi:10.1186/s12859-016-1452-4

19. Villaverde AF, Fröhlich F, Weindl D, Hasenauer J, Banga JR. Benchmarking optimization methods for
parameter estimation in large kinetic models. Bioinformatics. 2019;35: 830–838.
doi:10.1093/bioinformatics/bty736

20. Sher A, Niederer SA, Mirams GR, Kirpichnikova A, Allen R, Pathmanathan P, et al. A Quantitative
Systems Pharmacology Perspective on the Importance of Parameter Identifiability. B Math Biol. 2022;84: 39.
doi:10.1007/s11538-021-00982-5

21. Coope ID, Price CJ. A direct search conjugate directions algorithm for unconstrained minimization.
Anziam J. 2000;42: 478–498. doi:10.21914/anziamj.v42i0.609

22. Hough PD, Kolda TG, Torczon VJ. Asynchronous Parallel Pattern Search for Nonlinear Optimization.
Siam J Sci Comput. 2001;23: 134–156. doi:10.1137/s1064827599365823

23. Groenendaal W, Ortega FA, Kherlopian AR, Zygmunt AC, Krogh-Madsen T, Christini DJ. Cell-Specific
Cardiac Electrophysiology Models. Plos Comput Biol. 2015;11: e1004242. doi:10.1371/journal.pcbi.1004242

24. Sarkar AX, Sobie EA. Regression Analysis for Constraining Free Parameters in Electrophysiological
Models of Cardiac Cells. Plos Comput Biol. 2010;6: e1000914. doi:10.1371/journal.pcbi.1000914

25. Zaniboni M, Riva I, Cacciani F, Groppi M. How different two almost identical action potentials can be: A
model study on cardiac repolarization. Math Biosci. 2010;228: 56–70. doi:10.1016/j.mbs.2010.08.007

26. Krogh-Madsen T, Sobie EA, Christini DJ. Improving cardiomyocyte model fidelity and utility via dynamic
electrophysiology protocols and optimization algorithms. J Physiology. 2016;594: 2525–2536.
doi:10.1113/jp270618

27. Hooke R, Jeeves TA. `` Direct Search'' Solution of Numerical and Statistical Problems. J Acm Jacm.
1961;8: 212–229. doi:10.1145/321062.321069

28. Torczon V. On the Convergence of Pattern Search Algorithms. Siam J Optimiz. 1997;7: 1–25.
doi:10.1137/s1052623493250780

29. Ma J, Guo L, Fiene SJ, Anson BD, Thomson JA, Kamp TJ, et al. High purity human-induced pluripotent
stem cell-derived cardiomyocytes: electrophysiological properties of action potentials and ionic currents. 2011.

30. Hagiwara N, IRISAWA H, Kasanuki H, Hosoda S. Background current in sino-atrial node cells of the
rabbit heart. The Journal of physiology. 1992;448: 53–72. doi:10.1113/jphysiol.1992.sp019029

31. Kiyosue T, Spindler AJ, Noble SJ, NOBLE D. Background inward current in ventricular and atrial cells of
the guinea-pig. Proceedings Biological sciences. 1993;252: 65–74. doi:10.1098/rspb.1993.0047

32. Cheng H, Li J, James AF, Inada S, Choisy SCM, Orchard CH, et al. Characterization and influence of
cardiac background sodium current in the atrioventricular node. J Mol Cell Cardiol. 2016;97: 114–24.
doi:10.1016/j.yjmcc.2016.04.014

33. Tusscher KHWJ ten, Noble D, Noble PJ, Panfilov AV. A model for human ventricular tissue. Am J
Physiol-heart C. 2004;286: H1573–H1589. doi:10.1152/ajpheart.00794.2003

34. Ashford JR, Colquhoun D. Lectures on Biostatistics: An Introduction to Statistics with Applications in
Biology and Medicine. J Royal Statistical Soc Ser Gen. 1972;135: 606–606. doi:10.2307/2344687

35. Syed Z, Vigmond E, Nattel S, Leon LJ. Atrial cell action potential parameter fitting using genetic
algorithms. Medical Biological Eng Comput. 2005;43: 561–571. doi:10.1007/bf02351029

36. Guo T, Abed AA, Lovell NH, Dokos S. Optimisation of a Generic Ionic Model of Cardiac Myocyte
Electrical Activity. Comput Math Method M. 2013;2013: 706195. doi:10.1155/2013/706195

37. Beeler GW, Reuter H. Reconstruction of the action potential of ventricular myocardial fibres. J Physiology.
1977;268: 177–210. doi:10.1113/jphysiol.1977.sp011853

38. Faber GM, Rudy Y. Action Potential and Contractility Changes in [Na+]i Overloaded Cardiac Myocytes:
A Simulation Study. Biophys J. 2000;78: 2392–2404. doi:10.1016/s0006-3495(00)76783-x

39. Powell T, Twist VW. A rapid technique for the isolation and purification of adult cardiac muscle cells
having respiratory control and a tolerance to calcium. Biochem Bioph Res Co. 1976;72: 327–333.
doi:10.1016/0006-291x(76)90997-9

40. Sakmann B, Neher E. Patch Clamp Techniques for Studying Ionic Channels in Excitable Membranes.
Annu Rev Physiol. 1984;46: 455–472. doi:10.1146/annurev.ph.46.030184.002323

41. Cha CY, Himeno Y, Shimayoshi T, Amano A, Noma A. A novel method to quantify contribution of
channels and transporters to membrane potential dynamics. Biophys J. 2009;97: 3086–3094.
doi:10.1016/j.bpj.2009.08.060

42. Winslow RL, Walker MA, Greenstein JL. Modeling calcium regulation of contraction, energetics,
signaling, and transcription in the cardiac myocyte. Wiley Interdiscip Rev Syst Biology Medicine. 2016;8: 37–
67. doi:10.1002/wsbm.1322

43. Hilgemann DW, Feng S, Nasuhoglu C. The Complex and Intriguing Lives of PIP2 with Ion Channels and
Transporters. Sci Stke. 2001;2001: re19. doi:10.1126/stke.2001.111.re19

44. Suh B-C, Hille B. PIP2 Is a Necessary Cofactor for Ion Channel Function: How and Why? Biophysics.
2008;37: 175–195. doi:10.1146/annurev.biophys.37.032807.125859

45. Sasaki N, Mitsuiye T, Wang Z, Noma A. Increase of the delayed rectifier K+ and Na(+)-K+ pump currents
by hypotonic solutions in guinea pig cardiac myocytes. Circ Res. 2018;75: 887–895.
doi:10.1161/01.res.75.5.887

46. Hammami S, Willumsen NJ, Olsen HL, Morera FJ, Latorre R, Klaerke DA. Cell volume and membrane
stretch independently control K + channel activity: Cell volume, membrane stretch and K + channel activity. J
Physiology. 2009;587: 2225–2231. doi:10.1113/jphysiol.2008.163550

47. Peyronnet R, Nerbonne JM, Kohl P. Cardiac Mechano-Gated Ion Channels and Arrhythmias. Circ Res.
2016;118: 311–329. doi:10.1161/circresaha.115.305043

48. Gao J, Yun T, Xie X-L, Zhao J, Liu C, Sheng Y, et al. Losartan inhibits hyposmotic-induced increase of
IKs current and shortening of action potential duration in guinea pig atrial myocytes. Anatol J Cardiol.
2020;23: 35–40. doi:10.14744/anatoljcardiol.2019.75332

49. Cannell MB, Kong CHT. Local control in cardiac E–C coupling. J Mol Cell Cardiol. 2012;52: 298–303.
doi:10.1016/j.yjmcc.2011.04.014

50. Bers DM. Calcium Cycling and Signaling in Cardiac Myocytes. Annu Rev Physiol. 2008;70: 23–49.
doi:10.1146/annurev.physiol.70.113006.100455

51. Greenstein JL, Hinch R, Winslow RL. Mechanisms of Excitation-Contraction Coupling in an Integrative
Model of the Cardiac Ventricular Myocyte. Biophys J. 2006;90: 77–91. doi:10.1529/biophysj.105.065169

52. Negroni JA, Lascano EC. Simulation of steady state and transient cardiac muscle response experiments
with a Huxley-based contraction model. Journal of molecular and cellular cardiology. 2008;45: 300–312.
doi:10.1016/j.yjmcc.2008.04.012

53. Timmermann V, Edwards AG, Wall ST, Sundnes J, McCulloch AD. Arrhythmogenic Current Generation
by Myofilament-Triggered Ca2+ Release and Sarcomere Heterogeneity. Biophys J. 2019;117: 2471–2485.
doi:10.1016/j.bpj.2019.11.009

54. Niederer SA, Campbell KS, Campbell SG. A short history of the development of mathematical models of
cardiac mechanics. J Mol Cell Cardiol. 2019;127: 11–19. doi:10.1016/j.yjmcc.2018.11.015

55. Hagiwara N, Irisawa H, Kameyama M. Contribution of two types of calcium currents to the pacemaker
potentials of rabbit sino-atrial node cells. J Physiology. 1988;395: 233–253.
doi:10.1113/jphysiol.1988.sp016916

56. Zhou Z, Lipsius SL. T-Type Calcium Current in Latent Pacemaker Cells Isolated from Cat Right Atrium. J
Mol Cell Cardiol. 1994;26: 1211–1219. doi:10.1006/jmcc.1994.1139

57. Guo J, Ono K, Noma A. A sustained inward current activated at the diastolic potential range in rabbit sino-
atrial node cells. J Physiology. 1995;483: 1–13. doi:10.1113/jphysiol.1995.sp020563

58. Toyoda F, Mesirca P, Dubel S, Ding W-G, Striessnig J, Mangoni ME, et al. CaV1.3 L-type Ca2+ channel
contributes to the heartbeat by generating a dihydropyridine-sensitive persistent Na+ current. Sci Rep-uk.
2017;7: 7869. doi:10.1038/s41598-017-08191-8

59. Toyoda F, Wei-Guang D, Matsuura H. Heterogeneous functional expression of the sustained inward Na+
current in guinea pig sinoatrial node cells. Pflügers Archiv - European J Physiology. 2018;470: 481–490.
doi:10.1007/s00424-017-2091-y

60. Guinamard R, Bouvagnet P, Hof T, Liu H, Simard C, Sallé L. TRPM4 in cardiac electrical activity.
Cardiovasc Res. 2015;108: 21–30. doi:10.1093/cvr/cvv213

61. Doss MX, Diego JMD, Goodrow RJ, Wu Y, Cordeiro JM, Nesterenko VV, et al. Maximum diastolic
potential of human induced pluripotent stem cell-derived cardiomyocytes depends critically on I(Kr). Barbuti
A, editor. Plos One. 2012;7: e40288. doi:10.1371/journal.pone.0040288

62. Ishihara K, Yan D, Yamamoto S, Ehara T. Inward rectifier K+ current under physiological cytoplasmic
conditions in guinea-pig cardiac ventricular cells. J Physiology. 2002;540: 831–841.
doi:10.1113/jphysiol.2001.013470

63. Takahashi K, Yamanaka S. Induction of pluripotent stem cells from mouse embryonic and adult fibroblast
cultures by defined factors. Cell. 2006;126: 663–676. doi:10.1016/j.cell.2006.07.024

64. Nakagawa M, Koyanagi M, Tanabe K, Takahashi K, Ichisaka T, Aoi T, et al. Generation of induced
pluripotent stem cells without Myc from mouse and human fibroblasts. Nat Biotechnol. 2008;26: 101–106.
doi:10.1038/nbt1374

65. Yang L, Soonpaa MH, Adler ED, Roepke TK, Kattman SJ, Kennedy M, et al. Human cardiovascular
progenitor cells develop from a KDR+ embryonic-stem-cell-derived population. Nature. 2008;453: 524–528.
doi:10.1038/nature06894

10.   Appendix

10.1.   Preparation of dissociated hiPSC-CMs and recording of spontaneous APs

801   201B7 and 253G1 hiPSC lines generated from a healthy individual were used in this study [63,64].

Differentiation of hiPSCs into cardiomyocytes was promoted using an embryoid body (EB) differentiating

system [65]. hiPSC were incubated at 37 ºC in 5% CO_2, 5% O_2, and 90% N_2 for the first 12 days to promote

differentiation. The hiPSCs aggregated to form EBs and were cultured in suspension for 20 days. On the 20th

day of culture, EBs were treated with collagenase B (Roche, Basel, Switzerland) and trypsin EDTA (Nacalai

Tesque, Kyoto, Japan) and dispersed into single cells or small clusters, which were plated onto 0.1% gelatin-

coated dishes. hiPSC-CMs were then maintained in a conditioned medium. The experimental study using the

hiPSC-CMs was approved by the Kyoto University ethics review board (G259) and conformed to the

principles of the Declaration of Helsinki.

10.2.   Electrophysiological recordings of hiPSC-CM APs

811   For single-cell patch-clamp recordings, gelatin-coated glass coverslips were placed into each well of a

6-well plate, and 2 ml of DMEM/F12 containing 2% FBS and 80,000-120,000 CMs were added in each well.

Spontaneous APs were recorded from beating single CM using the perforated patch-clamp technique with

amphotericin B (Sigma-Aldrich) at 36 ± one ºC. Data were acquired at 20 kHz with the Multiclamp 700B

amplifier (Molecular Devices, Sunnyvale, CA, USA), Digidata 1440 digitizer hardware (Molecular Devices),

and pClamp 10.4 software (Molecular Devices). The glass pipettes had a resistance of 3-6 MΩ after filling

them with the intracellular solution. The external solution used for AP recordings was composed of the

following composition (in mM): NaCl 150, KCl 5.4, $CaCl_2$ 1.8, $MgCl_2$-$6H_2O$ 1, glucose 15, HEPES 15, and

Na-pyruvate 1; pH was adjusted to 7.4 by titrating NaOH. Intracellular solution contained (in mM): KCl 150,

NaCl 5, $CaCl_2$ 2, EGTA 5, MgATP 5, HEPES 10 (pH adjusted to 7.2 with KOH), and amphotericin B 300

µg/ml.