In-vivo **detection of cyclic-di-AMP in** *Staphylococcus aureus*

Running Title: RNA biosensor based detection of cyclic-di-AMP in *S. aureus*

Nagaraja Mukkayyan,[a,b,#] Raymond Poon,[a,b,#] Philipp N. Sander,[c] Li-Yin Lai,[a,b] Zahra

Zubair-Nizami,[a,b] Ming C. Hammond[c,d]* and Som S. Chatterjee[a,b,e]*

[a] Department of Microbial Pathogenesis, School of Dentistry, University of Maryland,

Baltimore, MD.

[b] Institute of Marine and Environmental Technology, Baltimore, MD.

[c] Department of Chemistry, University of California, Berkeley, CA.

[d] Department of Chemistry and Henry Eyring Center for Cell and Genome Sciences,

University of Utah, Salt Lake City, UT.

[e] University of Maryland Center for Environmental Science, Baltimore, MD.

[#] Equal contributors

* Correspondence: Ming C. Hammond (ming.hammond@utah.edu) and Som S.

Chatterjee (schatterjee@umaryland.edu)

ABSTRACT

Cyclic-di-AMP (CDA) is a signaling molecule that controls various cellular functions

including antibiotic tolerance and osmoregulation in *Staphylococcus aureus*. In this study,

we developed a novel biosensor (*bsuO* P6-4) for *in-vivo* detection of CDA in *S. aureus*.

Our study showed that *bsuO* P6-4 could detect a wide concentration range of CDA in

both laboratory and clinical strains making it suitable for use in both basic and clinical

research applications.

KEYWORDS: Cyclic-di-AMP, biosensors, *Staphylococcus aureus*

Main text

Cyclic-di-AMP (CDA) is a newly discovered second messenger, which is present in bacteria belonging to the phyla firmicutes and actinobacteria (1). Recent studies have demonstrated that CDA plays important roles in regulating vital biological processes such as DNA repair, ion homeostasis, and central carbon metabolism among others (2-5). In addition, CDA has also been implicated in controlling processes that are important for bacterial pathogenesis such as biofilm formation, antibiotic tolerance, and virulence (6-10). CDA mediates its function through binding to its cognate effectors (i.e. proteins and riboswitches) and thereby modifying their function through allosteric changes and/or through altered gene expression. Thus, maintaining the correct concentration of CDA in bacterial cells is critical not only to retaining cellular homeostasis but also to responding to changing environmental needs. This is attained by controlling CDA's synthesis or degradation machinery(s) that is present in bacterial cells.

In *Staphylococcus aureus*, CDA synthesis and degradation are mediated by DacA (diadenylate cyclase) and GdpP (the primary CDA phosphodiesterase) respectively (1). Recent studies have highlighted that increased CDA concentrations promote tolerance to β-lactam antibiotics and allow cell wall restructuring (8, 11). Furthermore, a growing number of contemporary clinical surveillance studies have identified mutations in *gdpP* among β-lactam resistant or non-susceptible natural isolates of *S. aureus* (12-15). Since many of these mutations have been either shown (8) or are predicted to attenuate the function of GdpP, causing increased CDA concentrations in cells, these findings underscored the clinical importance of CDA and highlighted the importance of accurate determination of its concentrations for both basic and clinical research settings.

Detection and quantification of CDA are typically carried out either through HPLC/MS,

indirect ELISA, or dye intercalation assay (16-18). These assays determine CDA's

abundance in a static manner, i.e. in samples containing bacterial cell lysates. Of these,

HPLC/MS despite being a gold standard in the quantification of small molecules such as

CDA requires expensive technical infrastructure and operational expertise. The operation

of ELISA is relatively easy but requires expertise in protein purification. In this study, we

present a novel RNA-based CDA biosensor (*bsuO* P6-4), which can determine its

concentration in live cells through flow cytometric analysis.

bsuO P6-4 is a second-generation CDA biosensor with improved CDA affinity and signal-

to-noise ratio than its predecessor, *yuaA* P1-4 (19). This improvement was achieved by

fusing the pro-fluorescent dye-binding RNA aptamer (Spinach) to the P6 stem instead of

the P1 stem (as in *yuaA* P1-4) of the natural CDA binding riboswitch sequence present in

the upstream region of *ydaO/yuaA* genes in *Bacillus subtilis* (20) (Fig.1A). This rational

design of *bsuO* P6-4 restored the pseudoknot interaction between the P1 and P8 stems,

which acts as a native stabilizer of the *ydaO/yuaA* riboswitch structure (21-23).

Additionally, a modified fluorescent dye binding module, coined as cpSpinach2 (24) that

was circularly permutized to accept a transducer stem was used for the construction of

bsuO P6-4 (Fig.1A). Thus, *bsuO* P6-4 consists of two components, the CDA-binding

*yda*O/*yua*A module, and the pro-fluorescent, DFHBI-binding cpSpinach2 module. Binding

of CDA to *bsuO* P6-4 enabled appropriate folding of the cpSpinach2 module, allowing

DFHBI binding and production of a fluorescence signal (Fig.1B) and thereby enabling

detection of CDA.

An *in-vitro* fluorescence assay testing the CDA biosensors revealed a >10-fold higher

affinity of *bsuO* P6-4 compared to its predecessor *yuaA* P1-4 (Fig.1C). In preparation for

in-vivo experiments, a tRNA scaffold was added to flank the 5' and 3' ends of the

biosensors for increased RNA half-life (Table S1) (25). The resultant biosensors were

cloned into a constitutive expression vector and transformed into a wild-type (Wt) *S.*

aureus and its isogenic Δ*gdpP* strain. While both the biosensors were able to report the

higher level of CDA that is characteristic of a Δ*gdpP* strain (8), *bsuO* P6-4 showed

significantly enhanced fluorescence compared to that of *yuaA* P1-4. More importantly,

bsuO P6-4 compared to *yuaA* P1-4 displayed a higher dynamic range of signal (3.73X vs

1.55X) between Wt and Δ*gdpP* strains making it amenable for detection of a wider

concentration range of CDA (Fig.1D). The reason why *yuaA* P1-4 exhibits higher

background fluorescence than *bsuO* P6-4 *in vivo* is unknown.

Next, we sought to determine whether *bsuO* P6-4 could report different concentrations of

CDA in the cells. For this purpose, isogenic *gdpP* point mutants were created that

displayed varying degrees of GdpP's loss of function that were identified in our previous

study (8). As shown in figure 2, *bsuO* P6-4 was able to detect differing CDA

concentrations in the isogenic strains (Fig.2A), which correlated well (R^2=0.9453) with the

results independently obtained through ELISA assay among the identical strains (Fig.2B

& C). In addition to the isogenic strains, *bsuO* P6-4 was also able to determine different

CDA concentrations in clinical isolates (Fig. S1), which suggested that it could be used in

both laboratory and clinical strains. Our results further showed that the Wt and Δ*gdpP*

strains with *bsuO* P6-4 were also suitable for fluorescent microscopic analysis (Fig. S2).

However, this method was not sensitive enough to differentiate the intermediate

concentrations of CDA of the other isogenic strains (data not shown).

In summary, we have developed a novel *bsuO* P6-4 biosensor that is effective in

determining different CDA concentrations in live *S. aureus* cells through flow cytometry.

The plasmid harboring *bsuO* P6-4 can be transformed into both laboratory and clinical *S.*

aureus isolates for reporting CDA concentration. This biosensor-based approach could

be used in flux detection of CDA concentrations in future studies.

ACKNOWLEDGMENTS

This work was supported by NIH grants 2R01AI100291 and startup funds provided by the

University of Maryland, Baltimore, and the University of Maryland Center for

Environmental Science to SSC and NIH grant R01 GM124589 to MCH. The authors

would also like to thank the Charles A. and Lois H. Miller Foundation for their generous

gift to purchase the flow cytometer used in this study.

REFERENCES

1.   Corrigan RM, Grundling A. 2013. Cyclic di-AMP: another second messenger

enters the fray. Nat Rev Microbiol 11:513-24.

2.   Corrigan RM, Campeotto I, Jeganathan T, Roelofs KG, Lee VT, Grundling A. 2013.

Systematic identification of conserved bacterial c-di-AMP receptor proteins. Proc

Natl Acad Sci U S A 110:9084-9.

3.   Oppenheimer-Shaanan Y, Wexselblatt E, Katzhendler J, Yavin E, Ben-Yehuda S.

2011. c-di-AMP reports DNA integrity during sporulation in Bacillus subtilis. EMBO

Rep 12:594-601.

4.   Sureka K, Choi PH, Precit M, Delince M, Pensinger DA, Huynh TN, Jurado AR,

Goo YA, Sadilek M, Iavarone AT, Sauer JD, Tong L, Woodward JJ. 2014. The

cyclic dinucleotide c-di-AMP is an allosteric regulator of metabolic enzyme

function. Cell 158:1389-1401.

5.   Witte G, Hartung S, Buttner K, Hopfner KP. 2008. Structural biochemistry of a

bacterial checkpoint protein reveals diadenylate cyclase activity regulated by DNA

recombination intermediates. Mol Cell 30:167-78.

6.   Fahmi T, Port GC, Cho KH. 2017. c-di-AMP: An Essential Molecule in the Signaling

Pathways that Regulate the Viability and Virulence of Gram-Positive Bacteria.

Genes (Basel) 8.

7.   Huynh TN, Luo S, Pensinger D, Sauer JD, Tong L, Woodward JJ. 2015. An HD-

domain phosphodiesterase mediates cooperative hydrolysis of c-di-AMP to affect

bacterial growth and virulence. Proc Natl Acad Sci U S A 112:E747-56.

8.   Poon R, Basuino L, Satishkumar N, Chatterjee A, Mukkayyan N, Buggeln E,

Huang L, Nair V, Argudin MA, Datta SK, Chambers HF, Chatterjee SS. 2022. Loss

of GdpP Function in Staphylococcus aureus Leads to beta-Lactam Tolerance and

Enhanced Evolution of beta-Lactam Resistance. Antimicrob Agents Chemother

66:e0143121.

9.   Witte CE, Whiteley AT, Burke TP, Sauer JD, Portnoy DA, Woodward JJ. 2013.

Cyclic di-AMP is critical for Listeria monocytogenes growth, cell wall homeostasis,

and establishment of infection. mBio 4:e00282-13.

10.   Woodward JJ, Iavarone AT, Portnoy DA. 2010. c-di-AMP secreted by intracellular

Listeria monocytogenes activates a host type I interferon response. Science

328:1703-5.

11.   Corrigan RM, Abbott JC, Burhenne H, Kaever V, Grundling A. 2011. c-di-AMP is

a new second messenger in Staphylococcus aureus with a role in controlling cell

size and envelope stress. PLoS Pathog 7:e1002217.

12.   Argudin MA, Dodemont M, Taguemount M, Roisin S, de Mendonca R, Deplano A,

Nonhoff C, Denis O. 2017. In vitro activity of ceftaroline against clinical

Staphylococcus aureus isolates collected during a national survey conducted in

Belgian hospitals. J Antimicrob Chemother 72:56-59.

13.   Argudin MA, Roisin S, Nienhaus L, Dodemont M, de Mendonca R, Nonhoff C,

Deplano A, Denis O. 2018. Genetic Diversity among Staphylococcus aureus

Isolates Showing Oxacillin and/or Cefoxitin Resistance Not Linked to the Presence

of mec Genes. Antimicrob Agents Chemother 62.

14.   Ba X, Kalmar L, Hadjirin NF, Kerschner H, Apfalter P, Morgan FJ, Paterson GK,

Girvan SL, Zhou R, Harrison EM, Holmes MA. 2019. Truncation of GdpP mediates

beta-lactam resistance in clinical isolates of Staphylococcus aureus. J Antimicrob

Chemother 74:1182-1191.

15.   Sommer A, Fuchs S, Layer F, Schaudinn C, Weber RE, Richard H, Erdmann MB,

Laue M, Schuster CF, Werner G, Strommenger B. 2021. Mutations in the gdpP

gene are a clinically relevant mechanism for beta-lactam resistance in meticillin-

resistant Staphylococcus aureus lacking mec determinants. Microb Genom 7.

16.   Burhenne H, Kaever V. 2013. Quantification of cyclic dinucleotides by reversed-

phase LC-MS/MS. Methods Mol Biol 1016:27-37.

17.   Underwood AJ, Zhang Y, Metzger DW, Bai G. 2014. Detection of cyclic di-AMP

using a competitive ELISA with a unique pneumococcal cyclic di-AMP binding

protein. J Microbiol Methods 107:58-62.

18.   Zhou J, Sayre DA, Zheng Y, Szmacinski H, Sintim HO. 2014. Unexpected complex

formation between coralyne and cyclic diadenosine monophosphate providing a

simple fluorescent turn-on assay to detect this bacterial second messenger. Anal

Chem 86:2412-20.

19.   Kellenberger CA, Chen C, Whiteley AT, Portnoy DA, Hammond MC. 2015. RNA-

Based Fluorescent Biosensors for Live Cell Imaging of Second Messenger Cyclic

di-AMP. J Am Chem Soc 137:6432-5.

20.   Nelson JW, Sudarsan N, Furukawa K, Weinberg Z, Wang JX, Breaker RR. 2013.

Riboswitches in eubacteria sense the second messenger c-di-AMP. Nat Chem Biol

9:834-9.

21.   Gao A, Serganov A. 2014. Structural insights into recognition of c-di-AMP by the

ydaO riboswitch. Nat Chem Biol 10:787-92.

22.   Jones CP, Ferre-D'Amare AR. 2014. Crystal structure of a c-di-AMP riboswitch

reveals an internally pseudo-dimeric RNA. EMBO J 33:2692-703.

23.   Ren A, Patel DJ. 2014. c-di-AMP binds the ydaO riboswitch in two pseudo-

symmetry-related pockets. Nat Chem Biol 10:780-6.

24.   Su Y, Hickey SF, Keyser SG, Hammond MC. 2016. In Vitro and In Vivo Enzyme

180   Activity Screening via RNA-Based Fluorescent Biosensors for S-Adenosyl-l-

181   homocysteine (SAH). J Am Chem Soc 138:7040-7.

182   25.   Ponchon L, Dardel F. 2007. Recombinant RNA technology: the tRNA scaffold. Nat

183   Methods 4:571-6.

FIGURE LEGENDS

Figure 1: Construction of the cyclic-di-AMP biosensor *bsuO* P6-4 and its ability to

detect cyclic-di-AMP. A) & B) Schematic representation of the creation and function of

bsuO P6-4. C) & D) Comparison of *in-vitro* and *in-vivo* detection of CDA respectively by

bsuO P6-4 and *yuaA* P1-4.

Figure 2: Ability of *bsuO* P6-4 in detecting varying concentrations of cyclic-di-AMP

in *S. aureus*. A) & B) Detection of CDA in wild-type and isogenic strains of *S. aureus*

which carried GdpP loss of function mutations using flow cytometry and ELISA assay

respectively. C) Correlation of signals obtained in A & B.

Figure 1: Construction of the cyclic-di-AMP biosensor *bsuO* P6-4 and its ability to detect cyclic-di-AMP. A) & B) Schematic representation of the creation and function of *bsuO* P6-4. C) & D) Comparison of *in-vitro* and *in-vivo* detection of CDA respectively by *bsuO* P6-4 and *yuaA* P1-4.

Figure 2: Ability of *bsuO* P6-4 in detecting varying concentrations of cyclic-di-AMP in *S. aureus*. A) & B) Detection of CDA in wild-type and isogenic strains of *S. aureus* which carried GdpP loss of function mutations using flow cytometry and ELISA assay respectively. C) Correlation of signals obtained in A & B.