Title: Evolution and diversification dynamics of butterflies

Authors:
Akito Y. Kawahara[1,2,3*], Caroline Storer[1], Ana Paula S. Carvalho[1], David M. Plotkin[1,2], Fabien
Condamine[4], Mariana P. Braga[5,6], Emily A. Ellis[1], Ryan A. St Laurent[1,3], Xuankun Li[1,11], Vijay
Barve[1], Liming Cai[7,8], Chandra Earl[1,10], Paul B. Frandsen[9], Hannah L. Owens[10,12], Wendy A.
Valencia-Montoya[7], Kwaku Aduse-Poku[13,14], Emmanuel F. A. Toussaint[1,15], Kelly M. Dexter[1],
Tenzing Doleck[13,16], Amanda Markee[1], Rebeccah Messcher[1], Y-Lan Nguyen[13], Jade Aster T.
Badon[17], Hugo A. Benítez[18,19], Michael F. Braby[20,21], Perry A. C. Buenavente[22], Wei-Ping
Chan[7], Steve C. Collins[23], Richard A. Rabideau Childers[7], Even Dankowicz[7], Rod Eastwood[7],
Zdenek F. Fric[24], Riley J. Gott[1,2], Jason P. W. Hall[25], Winnie Hallwachs[26], Nate B. Hardy[27],
Rachel L. Hawkins Sipe[7], Alan Heath[7,28], Jomar D. Hinolan[29], Nicholas T. Homziak[1,2], Yu-Feng
Hsu[30], Yutaka Inayoshi[31], Micael G.A. Itliong[13], Daniel H. Janzen[26], Ian J. Kitching[32],
Krushnamegh Kunte[33], Gerardo Lamas[34], Michael J. Landis[6], Elise A. Larsen[35], Torben B.
Larsen†[36], Jing V. Leong[13,24], Vladimir Lukhtanov[37], Crystal A. Maier[7], Jose I. Martinez[1,2], Dino
J. Martins[38], Kiyoshi Maruyama[39], Sarah C. Maunsell[7], Nicolás Oliveira Mega[40], Alexander
Monastyrskii[41,42], Ana B. B. Morais[43], Chris J. Müller[44], Mark Arcebal K. Naive[45,46,47], Gregory
Nielsen[48], Pablo Sebastián Padrón[1,49], Djunijanti Peggie[50], Helena Piccoli Romanowski[40],
Szabolcs Sáfián[51], Motoki Saito[52], Stefan Schröder[53], Vaughn Shirey[35], Doug Soltis[10], Pamela
Soltis[10], Andrei Sourakov[1], Gerard Talavera[7,54], Roger Vila[55], Petr Vlasanek[24,56], Houshuai
Wang[57], Andrew D. Warren[1], Keith R. Willmott[1], Masaya Yago[58], Walter Jetz[59], Marta A.
Jarzyna[59,60,61], Jesse W. Breinholt[62,63], Marianne Espeland[64], Leslie Ries[35], Robert P. Guralnick[10],
Naomi E. Pierce[7*], David J. Lohman[13,16,22*]

Author Affiliations:
[1]McGuire Center for Lepidoptera and Biodiversity, Florida Museum of Natural History, University of
Florida, Gainesville, FL, USA
[2]Entomology and Nematology Department, University of Florida, Gainesville, FL, USA
[3]Department of Biology, University of Florida, Gainesville, FL, USA
[4]CNRS, Institut des Sciences de l'Evolution de Montpellier (Université de Montpellier), Montpellier,
France
[5]Department of Ecology, Swedish University of Agricultural Sciences, Uppsala, Sweden
[6]Department of Biology, Washington University in St. Louis, St. Louis, MO, USA
[7]Department of Organismic and Evolutionary Biology and Museum of Comparative Zoology, Harvard
University, Cambridge, MA, USA
[8]Department of Botany and Plant Sciences, University of California, Riverside, Riverside, CA, USA
[9]Department of Plant and Wildlife Sciences, Brigham Young University, Provo, UT, USA
[10]Florida Museum of Natural History, University of Florida, Gainesville, FL, USA
[11]Center for Biodiversity Research, Department of Biological Sciences, University of Memphis,
Memphis, TN, USA
[12]Center for Macroecology, Evolution, and Climate, GLOBE Institute, University of Copenhagen,
Copenhagen, Denmark
[13]Biology Department, City College of New York, City University of New York, USA
[14]Department of Life and Earth Sciences, Perimeter College, Georgia State University, GA
[15]Department of Entomology, Natural History Museum of Geneva, Geneva, Switzerland
[16]PhD Program in Biology, Graduate Center, City University of New York, New York, NY, USA
[17]Rodolfo B. Gonzales Museum of Natural History, Biology Department, Silliman University, Philippines

[18]Laboratorio de Ecología y Morfometría Evolutiva, Centro de Investigación de Estudios Avanzados del Maule, Universidad Católica del Maule, Talca, Chile

[19]Centro de Investigación en Recursos Naturales y Sustentabilidad (CIRENYS), Universidad Bernardo O'Higgins, Avenida Viel 1497, Santiago, 8370993, Chile

[20]Division of Ecology and Evolution, Research School of Biology, The Australian National University, Acton, ACT, Australia

[21]Australian National Insect Collection, Canberra, ACT, Australia

[22]Entomology Section, National Museum of Natural History, Manila, Philippines

[23]African Butterfly Research Institute, Nairobi, Kenya

[24]Biology Centre CAS, České Budějovice, Czech Republic

[25]Department of Entomology, National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

[26]Department of Biology, University of Pennsylvania, Philadelphia, PA, USA

[27]Department of Entomology and Plant Pathology, Auburn University, Auburn, AL, USA

[28]Iziko South African Museum, Cape Town, South Africa

[29]Institute of Biology, College of Science, University of the Philippines Diliman, Quezon City 1101, Philippines

[30]College of Life Science, National Taiwan Normal University, Taipei, Taiwan

[31]Sritana Condominium 2, 96/173, Huay Kaeo Rd. T. Suthep, A. Muang, Chiang Mai, Thailand

[32]Natural History Museum, London, UK

[33]National Centre for Biological Sciences, Tata Institute of Fundamental Research, Bengaluru, India

[34]Museo de Historia Natural, Universidad Nacional Mayor de San Marcos, Lima, Peru

[35]Department of Biology, Georgetown University, Washington, DC, USA

[36]Deceased

[37]Department of Karyosystematics, Zoological Institute of Russian Academy of Sciences, St. Petersburg, Russia

[38]Mpala Research Center, Department of Ecology and Evolutionary Biology, Princeton University, Princeton, NJ, USA

[39]89-7-106 Minami-Ôsawa 5 chome, Hachiôji-shi, Japan

[40]Departamento de Zoologia, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS 91501970, Brazil

[41]Fauna & Flora International, Vietnam Programme, Hanoi, Vietnam

[42]Vietnam National Museum of Nature, Vietnam Academy of Science and Technology, 18 Hoang Quoc Viet, Cau Giay, Hanoi, Vietnam

[43]Universidade Federal de Santa Maria, Centro de Ciências Naturais e Exatas, Pós-Graduação em Biodiversidade Animal, Santa Maria, RS, Brazil

[44]Australian Museum, Sydney, NSW, Australia

[45]Center for Integrative Conservation, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Mengla, Yunnan 666303, China

[46]University of Chinese Academy of Sciences, Beijing 100049, China.

[47]College of Arts and Sciences, Jose Rizal Memorial State University, Tampilisan Campus, ZNAC, Tampilisan 7116, Zamboanga del Norte, Philippines

[48]Aquapro, Km 6 Villavicencio, Meta, Colombia

[49]Entomology Laboratory, Museo de Zoología, Universidad del Azuay, Cuenca, Ecuador

[50]Museum Zoologicum Bogoriense, Research Center for Biology-National Research and Innovation Agency (previously Indonesian Institute of Sciences), Cibinong-Bogor, Indonesia

[51]Institute of Silviculture and Forest Protection, University of West Hungary, Sopron, Hungary

[52]The Research Institute of Evolutionary Biology (Insect Study Division), Kami-youga, Setagaya-ku, Tokyo, Japan

[53]Auf dem Rosenhügel 15, D-50997 Köln, Germany

[54]Institut Botànic de Barcelona (IBB, CSIC-Ajuntament de Barcelona), Barcelona, Catalonia, Spain
[55]Institut de Biologia Evolutiva (CSIC-UPF), Barcelona, Catalonia, Spain
[56]T.G. Masaryk Water Research Institute, Prague, Czech Republic
[57]Department of Entomology, College of Plant Protection, South China Agricultural University,
Guangzhou, China
[58]The University Museum, The University of Tokyo, Tokyo, Japan
[59]Department of Ecology & Evolutionary Biology, Yale University, New Haven, CT, USA
[60]Department of Evolution, Ecology and Organismal Biology, The Ohio State University, Columbus, OH,
USA
[61]Translational Data Analytics Institute, The Ohio State University, Columbus, OH, USA
[62]Intermountain Healthcare, Intermountain Precision Genomics, Saint George, Utah, USA
[63]RAPiD Genomics, Gainesville, Florida, USA
[64]Leibniz Institute for the Analysis of Biodiversity Change, Zoological Research Museum Alexander
Koenig, Bonn, Germany
112
113
114   *Corresponding authors
115

Abstract

Butterflies are a diverse and charismatic insect group that are thought to have diversified via

coevolution with plants and in response to dispersals following key geological events. These

hypotheses have been poorly tested at the macroevolutionary scale because a comprehensive

phylogenetic framework and datasets on global distributions and larval hosts of butterflies are

lacking. We sequenced 391 genes from nearly 2,000 butterfly species to construct a new,

phylogenomic tree of butterflies representing 92% of all genera and aggregated global

distribution records and larval host datasets. We found that butterflies likely originated in what is

now the Americas, ~100 Ma, shortly before the Cretaceous Thermal Maximum, then crossed

Beringia and diversified in the Paleotropics. The ancestor of modern butterflies likely fed on

Fabaceae, and most extant families were present before the K/Pg extinction. The majority of

butterfly dispersals occurred from the tropics (especially the Neotropics) to temperate zones,

largely supporting a "cradle" pattern of diversification. Surprisingly, host breadth changes and

shifts to novel host plants had only modest impacts.

Background

Butterflies have captivated naturalists, scientists, and the general public for centuries, and they have played a central role in studies of speciation, community ecology, plant-insect interactions, mimicry, genetics, and conservation. Despite being the most studied group of insects, butterfly evolutionary history and drivers of their diversification are still poorly understood[1,2]. Due to over a century of efforts by amateur naturalists, butterflies are one of the few insect groups for which substantial trait data (e.g., geographic and host plant information) exist to test diversification hypotheses. Until now, these data have largely been scattered across the literature, museum collections, and local databases. Butterflies are thought to have diversified in relation to multiple abiotic and biotic factors, including adaptations to novel climates and species interactions, with geographic history and caterpillar-host interactions playing a major role[3]. However, these have not been studied because a synthesis of associated traits and a robust phylogenetic framework at the taxonomic scale needed to examine their evolution has not been available.

We sequenced 391 genes from nearly 2,000 butterfly species to construct a new, robust, phylogenomic tree of butterflies representing 92% of all genera, aggregated global distribution records, and assembled a comprehensive host association dataset. Using this tree, we infer the evolutionary timing, biogeographic history, and diversification patterns of butterflies. We address three long-standing questions related to butterfly evolution: 1) did butterflies originate in the northern (Laurasia) or southern (Gondwana) hemisphere?[4], 2) what plants did the ancestor of butterflies feed on?[5], and 3) were plants a major driver promoting butterfly diversification?[6]

Results and Discussion

To elucidate patterns of global butterfly diversification in space and time, we used targeted exon capture [7] to assemble a dataset of 391 gene regions (161,166 nucleotides and 53,722 amino acids) from 2,244 butterfly species. The majority (1,914 specimens) of butterflies sampled were newly sequenced for this study, and represented all families, tribes, and 92% of recognized genera, with most samples coming from museum collections across the world (see Supplementary Materials). Phylogenomic trees inferred with nucleotides and amino acids were highly congruent, with strong support for the monophyly of all families and nearly all subfamilies with traditional branch support metrics (Table S1), multispecies coalescence (MSC)

analyses (Table S1), and 4-cluster likelihood mapping (FcLM) (Table S2). Our results strongly

support the need for revision of the classification of at least 32 butterfly tribes (25% of total) as

currently circumscribed (Table S1).

We conducted 24 dating analyses using different fossil and secondary calibration

schemes along with sensitivity analyses to assess the impact of analytical and sampling bias.

Across analyses, our results revealed largely congruent timing of butterfly divergence events

(Table S3), indicating that butterflies originated from nocturnal, herbivorous moth ancestors

during the mid-Cretaceous (101.4 Ma, 102.5–100.0 Ma). This provides independent evidence for

the Cretaceous origin recovered in previous butterfly phylogenies[2,8]. We conducted

diversification rate analyses that used time-variable models and clade-heterogeneous models that

are less prone to sampling bias (Supplementary Methods). Both approaches recovered similar

patterns of diversification across butterflies, with increased rates in clades such as the diverse

skipper tribe Eudamini and lycaenid subfamily Poritiinae (Fig. 1, Fig. S1).

To determine the geographic origin of butterflies, we used our dated phylogeny (Fig. 1)

to conduct a global biogeographic analysis with 15,764 newly aggregated country-level

distribution records (Table S4). Modeling with three different area categorizations, models of

range evolution, and parameters (e.g., adjacency matrices, time slices, etc.) consistently

recovered butterflies as originating in the Americas, in what is present-day western North

America or Central America (Fig. 4, Table S5). All extant butterfly families excluding the

Neotropical Hedylidae diversified ~10–30 Ma after the Cretaceous Thermal Maximum (CTM),

~90 Ma, when the global climate cooled by nearly 5° C[9] (Figs. 1, 2). During the Cretaceous,

butterflies appeared to be dispersing out of the Neotropics at a much higher rate than all other

dispersal pathways (Fig. S2). As new butterfly lineages became established in other bioregions,

other inter-bioregion dispersal avenues became more frequently used, particularly out of

Indomalaya (Figs. S3, S4).

Beginning around 60 Ma, the Neotropics served as an evolutionary "cradle"[10] with high

in situ butterfly speciation (Fig. S5), and many dispersal events occurring out of the Neotropics

to other areas (Fig. S6). Relative rate of dispersal out of the Neotropics was still high during the

early Cenozoic, although not as high as it was during the Cretaceous (Figs. S2, S3). Over the

course of their evolution, butterflies experienced significantly higher speciation rates in the

tropics compared to temperate zones (Data S1), and also more dispersal events out of the tropics

(Fig. S6), as observed by high relative mean dispersal rates out of the tropics (e.g., tropical

Indomalaya to temperate East Palearctic, and the Neotropics to Nearctic; Fig. 3). This pattern

differs from that seen in mammals, which are thought to have dispersed primarily in the opposite

direction during the Pliocene[11–13]. Some butterflies such as swallowtails showed greater

immigration into the Neotropics, following a "museum" model of diversification (Fig. S7),

corroborating prior findings[14]. The majority of dispersal events between the Neotropics and the

Nearctic took place after the Eocene-Oligocene boundary, ~33.9 Ma (Fig. S4). Two lineages

dispersed from the East Palearctic around 17 Ma, and these appear to be the first colonizers of

Europe: ancestors of a Nymphalini subclade including *Aglais, Nymphalis,* and *Polygonia*, and a

clade of checkered skippers (Carcharodini; Table S6). Butterflies were present on what are now

all modern continental landmasses by the late Eocene (Table S7).

The K/Pg boundary (~66 Ma) marked a major global extinction event that dramatically

reduced vertebrate and marine diversity[15]. Less is known, however, about the extent to which

insects were affected by this event[16]. We used two analytical approaches to identify sudden

extinction events and estimate changes in diversification rates, both of which concluded that

butterflies, like amphibians and some mammals[17,18], did not experience a major extinction event

at the K/Pg boundary (Fig. S8). Plants experienced a vegetation turnover from gymnosperms to

angiosperms in the Late Cretaceous[19] and continued to increase in diversity up to the present

(Fig. 2). The survival of many plant lineages across the K/Pg boundary may have buffered the

effects of the mass extinction event of butterflies.

Two tests of extinction demonstrate that butterflies underwent two major extinctions after

the K/Pg boundary — one at the Eocene–Oligocene transition (EOT) which coincides with the

Eocene–Oligocene global extinction event (~33.9 Ma), and another during the mid–Miocene that

coincides with the mid-Miocene Climatic Optimum (~14 Ma; Fig. S8). These were global

cooling and warming periods, respectively, that led to the disappearance of many plant and

animal lineages, followed by sharp floral and faunal turnover in temperate and tropical

environments[15]. Speciation analyses with BioGeoBEARS[20] (see Methods), also supports these

findings; there was a dramatic drop in the number of speciation events around 34 Ma, especially

in the East Palearctic region (Fig. S5), which occurred around the EOT extinction event that

significantly reduced floral and faunal diversity of that region[21]. Soon thereafter, the families

Hesperiidae and Nymphalidae experienced dramatic increases in the number of speciation events

228 in the East Palearctic (Fig. S9). This marks a period when East Palaearctic habitats changed from

229 dense forest to forest-steppe, temperate grassland, and mixed forest biomes[21]. These serve as

230 primary habitats of the diverse modern butterfly groups – fritillaries, satyrs, and grassland

231 skippers, which dominate this region today (Argynnini, Satyrini, and Hesperiinae, respectively).

232   It is often argued that butterfly evolution is closely tied to the diversification of

233 angiosperms[5,6]. Yet, few studies have examined the timing and pattern of butterflies and their

234 hosts on a broad, macroevolutionary scale. To understand the temporal diversification of

235 butterflies and plants, we compiled 31,456 butterfly host records from 186 books and databases

236 (Table S8). We examined how butterfly lineages increased through time and compared these

237 results with four recent studies that examined plant diversification dynamics. We found that

238 butterfly diversification lagged far behind the origin of angiosperms (Fig. 2), corroborating prior

239 studies[7,22]. Ancestral state estimation provided support for Fabaceae as the host plant of the most

240 recent common ancestor of butterflies (Tables S9-S10, Fig. S10), a widely accepted hypothesis[5]

241 that has lacked empirical support. The crown age of the most recent common ancestor of

242 Fabaceae is thought to be ~98 Ma[23,24], largely coincident with the origin of butterflies.

243   Although the vast majority of butterfly larvae in our dataset are herbivores, a small

244 number also feed on detritus, lichens, or insects (Table S8). The oldest associations in the

245 entirely entomophagous Miletinae (Lycaenidae) appear to originate by 58.4 Ma (58.9–57.1 Ma),

246 an estimate that corresponds with an earlier estimation of the origin of this group[25] (Tables S3,

247 S11). The Lycaenidae, with caterpillars that are ancestrally symbiotic with ants[7,26], date back to

248 64.5 Ma (65.4–63.7 Ma) (Fig. S11), long after the origin of ants (139–158 Ma[27]). Together with

249 plants, ants appear to have provided a template for diversification of Lycaenidae and certain

250 members of its sister clade, Riodinidae.

251   To address the possibility that butterfly diversification may have been spurred by co-

252 diversifying plants, we evaluated eight paleoenvironment-dependent models relating butterfly

253 diversification to host-plant evolution[28]. We first estimated time-dependent variation of

254 speciation and extinction rates with time-continuous birth-death models and compared them with

255 constant-rate diversification models[29]. In the best-fitting model, speciation rates vary

256 exponentially through time without extinction (92% of the trees, Akaike $\omega = 0.549$). This time-

257 dependent model is far better supported than the angiosperm-dependent models, the best of

258 which was ranked fourth (Table S12). We also ran the HiSSE model[30] as implemented in the R

package hisse[31] to test for a potential impact of host-plant preference on diversification

dynamics. In total, we compared 18 different models of HiSSE and BiSSE-like implementations,

accounting for hidden states to alleviate recent concerns regarding the reliability of SSE models

and the high incidence of false positive results[32]. We found little or no indication that association

with a specific plant group is linked to butterfly diversification, largely rejecting the hypothesis

of hostplant driven diversification (Table S13, Data S2). These results indicate that

coevolutionary interactions with angiosperm host-plants were unlikely to have been a driver of

butterfly diversification at macroevolutionary scales. Even within families, hypotheses such as

the correlated diversification of grasses with grass-feeding skippers (Hesperiinae) and satyrs

(Satyrini)[33] were uncorroborated (Table S13). Our results are consistent with smaller studies on

particular butterfly groups such as skippers (e.g., Sahoo et al.[34]), that show that alternative

drivers (hidden states) explain best butterfly diversification when compared to hostplant-

dependent models.

Butterfly diversification may not have been driven by the availability of particular plants,

but instead by host specificity (specializing on a few plants [e.g., Janz, Nylin and Nyblom[35]]). It

has been proposed through the resource-use hypothesis[36] that specialists that feed on few hosts

have higher speciation and extinction rates. Numerous studies on vertebrates support this

hypothesis[37,38], but few studies have tested the hypothesis on insects. We examined host plant

specificity on the butterfly phylogeny (Fig. 1) and found that more than two-thirds of butterfly

species feed on a single plant family (67.7%), whereas less than one-third (32.3%) are generalists

feeding on two or more (Table S14), a pattern largely in agreement with ecological studies on

butterflies[39]. We also found that 94.2% of generalists feed on plant families that are significantly

closely related compared to a randomly sampled null distribution, suggesting that generalists,

although capable of feeding on different host families, still consume closely related plants. This

finding supports the pattern reported by Ehrlich and Raven[6] that related butterflies feed on

related plants. Finally, we used trait-dependent methods to examine whether diversification rates

changed following switches between generalist and specialist, and host plant shifts (Table S13).

Our results demonstrate that – at least in analyses of Papilionoidea and Hesperiidae, where

results were significant – specialists have lower rates of speciation (Fig. S12), contradicting the

resource-use hypothesis.

This study includes the most comprehensive taxonomic sampling of butterflies to date to understand their evolutionary history. We also compiled two comprehensive trait datasets (geography and host plants) that demonstrate the importance of assembling large trait databases to test key hypotheses. Our study overturns some long-held assumptions of butterfly evolution and provides a framework for future studies of this model insect lineage. The consistency of results obtained using different approaches for each of our analyses suggests that our conclusions are robust. Assembling species-level datasets will be an important future goal to address fine-scale trends within particular butterfly clades.

Our data support the hypothesis that butterflies originated in the Americas in the Late Cretaceous 100 million years after the origin of angiosperms, and that they first fed on legumes. They dispersed from the Americas to the East Palearctic likely across Beringia ~75 Ma before diversifying in the Paleotropics. They appear to have been minimally affected by the K/Pg mass extinction. While some evidence points to a Nearctic origin, the evidence for Nearctic versus Neotropical origin is not strong – we therefore only tentatively conclude that a Laurasian origin is possible. Diversification of some lineages has been impacted by host associations, but an escape-and-radiate model of coevolution with angiosperms does not appear to have been a powerful driver of diversification at a broad macroevolutionary scale. Butterfly evolution is better viewed as compatible with a model of diffuse coevolution[40] in which plants provided an ecological template on which butterflies diversified as opportunities arose.

References

1. Chazot, N. et al. Priors and posteriors in Bayesian timing of divergence analyses: the age of butterflies revisited. *Systematic Biology* **68**, 797–813 (2019).

2. Allio, R. et al. Whole genome shotgun phylogenomics resolves the pattern and timing of swallowtail butterfly evolution. *Systematic Biology* **69**, 38–60 (2020).

3. Boggs, C. L., Watt, W. B. & Ehrlich, P. R. *Butterflies: Ecology and Evolution Taking Flight*. (University Of Chicago Press, 2003).

4. Braby, M. F., Trueman, J. W. H. & Eastwood, R. When and where did troidine butterflies (Lepidoptera: Papilionidae) evolve? Phylogenetic and biogeographic evidence suggests an origin in remnant Gondwana in the Late Cretaceous. *Invertebrate Systematics* **19**, 113–143

320   (2005).

5. Janz, N. & Nylin, S. Butterflies and plants: a phylogenetic study. *Evolution* **52**, 486–502
    (1998).

6. Ehrlich, P. R. & Raven, P. H. Butterflies and plants: a study of coevolution. *Evolution* **18**,
    586–608 (1965).

7. Espeland, M. et al. A comprehensive and dated phylogenomic analysis of butterflies. *Current
    Biology* **28**, 770-778.e5 (2018).

8. Wahlberg, N., Wheat, C. W. & Peña, C. Timing and patterns in the taxonomic diversification
    of Lepidoptera (butterflies and moths). *PLOS ONE* **8**, e80875 (2013).

9. Linnert, C. et al. Evidence for global cooling in the Late Cretaceous. *Nature
    Communications* **5**, 4194 (2014).

10. Stebbins, G. L. *Flowering plants: Evolution above the species level*. (Belknap Press of
    Harvard Univ. Press, 1974).

11. Domingo, L., Tomassini, R. L., Montalvo, C. I., Sanz-Pérez, D. & Alberdi, M. T. The Great
    American Biotic Interchange revisited: a new perspective from the stable isotope record of
    Argentine Pampas fossil mammals. *Scientific Reports* **10**, 1608 (2020).

12. Carrillo, J. D. et al. Disproportionate extinction of South American mammals drove the
    asymmetry of the Great American Biotic Interchange. *Proceedings of the National Academy
    of Sciences* **117**, 26281–26287 (2020).

13. Rolland, J., Condamine, F. L., Beeravolu, C. R., Jiguet, F. & Morlon, H. Dispersal is a major
    driver of the latitudinal diversity gradient of Carnivora. *Global Ecology and Biogeography*
    24, 1059–1071 (2015).

14. Condamine, F. L., Silva-Brandão, K. L., Kergoat, G. J. & Sperling, F. A. H. Biogeographic
    and diversification patterns of Neotropical Troidini butterflies (Papilionidae) support a
    museum model of diversity dynamics for Amazonia. *BMC Evolutionary Biology* **12**, (2012).

15. Hallam, A. & Wignall, P. B. *Mass Extinctions and Their Aftermath*. (Oxford University
    Press, UK, 1997).

16. Schachat, S. R. & Labandeira, C. C. Are insects heading toward their first mass extinction?
    Distinguishing turnover from crises in their fossil record. *Annals of the Entomological
    Society of America* **114**, 99–118 (2021).

17. Grossnickle, D. M. & Newham, E. Therian mammals experience an ecomorphological

radiation during the Late Cretaceous and selective extinction at the K-Pg boundary.

Proceedings of the Royal Society B: Biological Sciences **283**, 20160256 (2016).

18. Feng, Y.-J. et al. Phylogenomics reveals rapid, simultaneous diversification of three major

clades of Gondwanan frogs at the Cretaceous-Paleogene boundary. *Proceedings of the*

National Academy of Sciences **114**, E5864–E5870 (2017).

19. Condamine, F. L., Silvestro, D., Koppelhus, E. B. & Antonelli, A. The rise of angiosperms

pushed conifers to decline during global cooling. *Proceedings of the National Academy of*

Sciences **117**, 28867 (2020).

20. Matzke, N. J. BioGeoBEARS: biogeography with Bayesian (and likelihood) evolutionary

analysis in R scripts. (2013).

21. Sun, J. et al. Synchronous turnover of flora, fauna and climate at the Eocene–Oligocene

Boundary in Asia. *Scientific Reports* **4**, 7463 (2014).

22. Heikkilä, M., Kaila, L., Mutanen, M., Peña, C. & Wahlberg, N. Cretaceous origin and

repeated tertiary diversification of the redefined butterflies. *Proceedings of the Royal Society*

B: Biological Sciences **279**, 1093–1099 (2011).

23. Magallón, S., Gómez-Acevedo, S., Sánchez-Reyes, L. L. & Hernández-Hernández, T. A

metacalibrated time-tree documents the early rise of flowering plant phylogenetic diversity.

New Phytologist **207**, 437–453 (2015).

24. Li, H.-T. et al. Origin of angiosperms and the puzzle of the Jurassic gap. *Nature Plants* **5**,

461–470 (2019).

25. Kaliszewska, Z. A. et al. When caterpillars attack: Biogeography and life history evolution

of the Miletinae (Lepidoptera: Lycaenidae). *Evolution* **69**, 571–588 (2015).

26. Pierce, N. E. et al. The ecology and evolution of ant association in the Lycaenidae

(Lepidoptera). *Annual Review of Entomology* **47**, 733–771 (2002).

27. Moreau, C. S. & Bell, C. D. Testing the museum versus cradle tropical biological diversity

hypothesis: phylogeny, diversification, and ancestral biogeographical range evolution of the

ants. *Evolution* **67**, 2240–2257 (2013).

28. Condamine, F. L., Rolland, J. & Morlon, H. Macroevolutionary perspectives to

environmental change. *Ecology Letters* **16**, 72–85 (2013).

29. Morlon, H., Parsons, T. L. & Plotkin, J. B. Reconciling molecular phylogenies with the fossil

record. *Proceedings of the National Academy of Sciences* **108**, 16327 (2011).

30. Beaulieu, J. M. & O'Meara, B. C. Detecting hidden diversification shifts in models of trait-
dependent speciation and extinction. *Systematic Biology* **65**, 583–601 (2016).

31. Beaulieu, J., O'Meara, B., Caetano, D., Boyko, J. & Vasconcelos, T. *Package 'hisse'*.
(2021).

32. Rabosky, D. L. & Goldberg, E. E. Model inadequacy and mistaken inferences of trait-
dependent speciation. *Systematic Biology* **64**, 340–355 (2015).

33. Peña, C. & Wahlberg, N. Prehistorical climate change increased diversification of a group of
butterflies. *Biology Letters* **4**, 274–278 (2008).

34. Sahoo, R. K., Warren, A. D., Collins, S. C. & Kodandaramaiah, U. Hostplant change and
paleoclimatic events explain diversification shifts in skipper butterflies (Family:
Hesperiidae). *BMC Evolutionary Biology* **17**, 174 (2017).

35. Janz, N., Nyblom, K. & Nylin, S. Evolutionary dynamics of host-plant specialization: a case
study of the tribe Nymphalini. *Evolution* **55**, 783–796 (2001).

36. Vrba, E. S. Ecology in relation to speciation rates: some case histories of Miocene-Recent
mammal clades. *Evolutionary Ecology* **1**, 283–300 (1987).

37. Fernández, M. H. et al. Macroevolution and climate changes: a global multi-family test
supports the resource-use hypothesis in terrestrial mammals. *null* 1–9 (2022)
doi:10.1080/08912963.2022.2042807.

38. Cantalapiedra, J. L., Hernández Fernández, M. & Morales, J. Biomic specialization and
speciation rates in Ruminants (Cetartiodactyla, Mammalia): A test of the resource-use
hypothesis at the global scale. *PLOS ONE* **6**, e28749 (2011).

39. Forister, M. L. et al. The global distribution of diet breadth in insect herbivores. *Proceedings
of the National Academy of Sciences* **112**, 442 (2015).

40. Levin, S. A., Segel, L. A. & Adler, F. R. Diffuse coevolution in plant-herbivore
communities. *Theoretical Population Biology* **37**, 171–191 (1990).

41. Lemmon, A. R., Emme, S. A. & Lemmon, E. M. Anchored hybrid enrichment for massively
high-throughput phylogenomics. *Systematic Biology* **61**, 727–744 (2012).

42. Breinholt, J. W. et al. Resolving relationships among the megadiverse butterflies and moths
with a novel pipeline for Anchored Phylogenomics. *Systematic Biology* **67**, 78–93 (2018).

43. Regier, J. C. et al. Arthropod relationships revealed by phylogenomic analysis of nuclear
protein-coding sequences. *Nature* **463**, 1079–1083 (2010).

44. Zwick, A. *Degeneracy Coding Web Service*. (2010).

45. Minh, B. Q., Nguyen, M. A. T. & von Haeseler, A. Ultrafast approximation for phylogenetic

Bootstrap. *Molecular Biology and Evolution* **30**, 1188–1195 (2013).

46. Hoang, D. T., Chernomor, O., von Haeseler, A., Minh, B. Q. & Vinh, L. S. UFBoot2:

Improving the Ultrafast Bootstrap approximation. *Molecular Biology and Evolution* **35**, 518–

522 (2018).

47. Guindon, S. et al. New algorithms and methods to estimate maximum-likelihood

phylogenies: assessing the performance of PhyML 3.0. *Systematic Biology* **59**, 307–321

(2010).

48. Strimmer, K. & von Haeseler, A. Likelihood-mapping: A simple method to visualize

phylogenetic content of a sequence☐alignment. *Proceedings of the National Academy of*

Sciences **94**, 6815 (1997).

49. Smith, S. A. & O'Meara, B. C. TreePL: divergence time estimation using penalized

likelihood for large phylogenies. *Bioinformatics* **28**, 2689–2690 (2012).

50. Kawahara, A. Y. et al. Phylogenomics reveals the evolutionary timing and pattern of

butterflies and moths. *Proceedings of the National Academy of Sciences of the United States*

of America **116**, 22657–22663 (2019).

51. Parham, J. F. et al. Best practices for justifying fossil calibrations. *Systematic Biology* **61**,

346–359 (2012).

52. de Jong, R. Fossil butterflies, calibration points and the molecular clock (Lepidoptera:

Papilionoidea). *Zootaxa* **4270**, 1–63 (2017).

53. Graur, D. & Martin, W. Reading the entrails of chickens: molecular timescales of evolution

and the illusion of precision. *TRENDS in Genetics* **20**, 80–86 (2004).

54. Smith, S. A. & Brown, J. W. Constructing a broadly inclusive seed plant phylogeny.

American Journal of Botany **105**, 302–314 (2018).

55. Foster, C. S. et al. Evaluating the impact of genomic data and priors on Bayesian estimates of

the angiosperm evolutionary timescale. *Systematic Biology* **66**, 338–351 (2017).

56. Shields, O. Fossil butterflies and the evolution of Lepidoptera. *Journal of Research on the*

Lepidoptera **15**, 132–143 (1976).

57. Wahlberg, N. That awkward age for butterflies: insights from the age of the butterfly

subfamily Nymphalinae (Lepidoptera: Nymphalidae). *Systematic Biology* **55**, 703–714

(2006).

58. Rabosky, D. L. Automatic detection of key innovations, rate shifts, and diversity-dependence

on phylogenetic trees. *PLOS ONE* **9**, e89543–e89543 (2014).

59. Rabosky, D. L. et al. BAMMtools: An R package for the analysis of evolutionary dynamics

on phylogenetic trees. *Methods in Ecology and Evolution* **5**, 701–707 (2014).

60. Maddison, W. P. & Maddison, D. R. Mesquite: a modular system for evolutionary analysis.

(2018).

61. Moore, B. R., Höhna, S., May, M. R., Rannala, B. & Huelsenbeck, J. P. Critically evaluating

the theory and performance of Bayesian analysis of macroevolutionary mixtures.

Proceedings of the National Academy of Sciences **113**, 9569–9574 (2016).

62. Rabosky, D. L., Mitchell, J. S. & Chang, J. Is BAMM flawed? Theoretical and practical

concerns in the analysis of multi-rate diversification models. *Systematic Biology* **66**, 477–498

(2017).

63. Höhna, S. et al. A Bayesian approach for estimating branch-specific speciation and

extinction rates. *bioRxiv* 555805 (2019) doi:10.1101/555805.

64. Höhna, S. et al. RevBayes: Bayesian phylogenetic inference using graphical models and an

interactive model-specification language. *Systematic Biology* **65**, 726–736 (2016).

65. Louca, S. & Doebeli, M. Efficient comparative phylogenetics on large trees. *Bioinformatics*

34, 1053–1055 (2018).

66. Stadler, T. Mammalian phylogeny reveals recent diversification rate shifts. *Proceedings of*

the National Academy of Sciences **108**, 6187 (2011).

67. Höhna, S., May, M. R. & Moore, B. R. TESS: an R package for efficiently simulating

phylogenetic trees and performing Bayesian inference of lineage diversification rates.

Bioinformatics **32**, 789–791 (2016).

68. Pinkert, S., Barve, V., Guralnick, R. P. & Jetz, W. Global geographical and latitudinal

variation in butterfly species richness captured through a comprehensive country☐level

occurrence database. *Global Ecology and Biogeography* (2022) doi:10.1111/geb.13475.

69. Ree, R. H. & Smith, S. A. Maximum likelihood inference of geographic range evolution by

dispersal, local extinction, and cladogenesis. *Systematic Biology* **57**, 4–14 (2008).

70. Smith, S. A. Taking into account phylogenetic and divergence-time uncertainty in a

parametric biogeographical analysis of the Northern Hemisphere plant clade Caprifolieae.

Journal of Biogeography **36**, 2324–2337 (2009).

71. Beeravolu, C. R. & Condamine, F. L. An extended Maximum Likelihood inference of

geographic range evolution by dispersal, local extinction and cladogenesis. *bioRxiv* 038695

(2016) doi:10.1101/038695.

72. Allio, R. et al. Genome-wide macroevolutionary signatures of key innovations in butterflies

colonizing new host plants. *Nature Communications* **12**, 354 (2021).

73. Udvardy, M. D. F. *A classification of the biogeographical provinces of the world. Morges*

(Switzerland): International Union of Conservation of Nature and Natural Resources. IUCN

Occasional Paper no. 18. (1975).

74. Ree, R. H. Detecting the historical signature of key innovations using stochastic models of

character evolution and cladogenesis. *Evolution* **59**, 257–265 (2005).

75. Ronquist, F. Dispersal-vicariance analysis: a new approach to the quantification of historical

biogeography. *Systematic Biology* **46**, 195–203 (1997).

76. Li, X. et al. A diversification relay race from Caribbean-Mesoamerica to the Andes:

historical biogeography of *Xylophanes* hawkmoths. *Proceedings of the Royal Society B:*

Biological Sciences **289**, 20212435 (2022).

77. Edger, P. P. et al. The butterfly plant arms-race escalated by gene and genome duplications.

Proceedings of the National Academy of Sciences **112**, 8362 (2015).

78. Braga, M. P., Landis, M. J., Nylin, S., Janz, N. & Ronquist, F. Bayesian inference of

ancestral host–parasite interactions under a phylogenetic model of host repertoire evolution.

Systematic Biology **69**, 1149–1162 (2020).

79. Kembel, S. W. et al. Picante: R tools for integrating phylogenies and ecology. *Bioinformatics*

26, 1463–1464 (2010).

80. Beckett, S. J. Improved community detection in weighted bipartite networks. *Royal Society*

Open Science **3**, 140536 (2016).

81. Dormann, C. F., Gruber, B. & Fründ, J. Introducing the bipartite package: analysing

ecological networks. *R News* **8**, 8–11 (2008).

82. R Core Team. R: A language and environment for statistical computing. (2009).

83. Revell, L. J. phytools: an R package for phylogenetic comparative biology (and other

things). *Methods in ecology and evolution* **3**, 217–223 (2012).

84. Morlon, H. et al. RPANDA: an R package for macroevolutionary analyses on phylogenetic

trees. *Methods in Ecology and Evolution* **7**, 589–597 (2016).

85. Scotese, C. R., Song, H., Mills, B. J. W. & van der Meer, D. G. Phanerozoic

paleotemperatures: The earth's changing climate during the last 540 million years. *Earth-*

Science Reviews **215**, 103503 (2021).

86. van Nieukerken, E. J. et al. Order Lepidoptera Linnaeus, 1758. In: Zhang, Z.-Q. (Ed.)

Animal biodiversity: An outline of higher-level classification and survey of taxonomic

richness. *Zootaxa* **3148**, 212 (2011).

87. Chazot, N. et al. Conserved ancestral tropical niche but different continental histories explain

the latitudinal diversity gradient in brush-footed butterflies. *Nature Communications* **12**,

5717 (2021).

88. Wahlberg, N. et al. Synergistic effects of combining morphological and molecular data in

resolving the phylogeny of butterflies and skippers. *Proceedings of the Royal Society B:*

Biological Sciences **272**, 1577–1586 (2005).

Methods

Taxon sampling and sequence acquisition.

A total of 2,248 butterfly specimens representing 2,244 species in 1,644 genera was

included for the molecular component of this study (Table S15). We obtained marker loci used

for phylogenetic analysis via Anchored Hybrid Enrichment (AHE) exon capture of DNA extracts

and subsequent Illumina sequencing[41] or by bioinformatically extracting these sequences from

published genomes and transcriptomes. We used the BUTTERFLY1.0 probe set[7], which can

capture up to 450 loci.

We extracted DNA from 1,915 specimens that were: 1) stored in ethanol and frozen; 2)

dried and stored in glassine envelopes under ambient conditions (papered); or 3) dried, spread

and pinned in a museum collection. Locus assembly and sequence clean-up followed the pipeline

of Breinholt et al.[42]. Published sequences comprised: 1) genome assemblies; 2) genomic reads;

and 3) paired or 4) single-end transcriptomes. Three sequence datasets were created for this

study: nt123 (a nucleotide dataset with all codon positions), degen (a nucleotide dataset that

excludes all synonymous changes, created using the perl script, Degen1 v.1.4[43,44]), and aa (an

amino acid dataset translated from the nt123 dataset) (Data S3).

Phylogenetic analysis and dating.

Maximum likelihood (ML) tree inference was conducted on all three datasets in IQ-

TREE 2.0 (nt123, degen, and aa), and parameter settings for each analysis can be found in Table

S16. Branch support was calculated with 1,000 ultrafast bootstrap replicates (UFBS; '-B 1000'

command)[45,46], and Shimodaira-Hasegawa approximate likelihood ratio tests (SH-aLRT; '-alrt

1000' command)[47]. Quartet sampling (see *Section 8. Sequence quality check*) was performed on

the most-likely degen359 and aa154 trees to examine the topology and provide more

comprehensive and specific information on branch support. Four-cluster likelihood mapping

(FcLM) analyses[48] were performed on the degen and aa datasets to assess the placement of

particular butterfly clades that have been the subject of previous phylogenetic studies. We

applied this approach in addition to standard branch support metrics because the latter can be

subject to inflated estimates[48].

We obtained divergence time estimates using a penalized-likelihood based approach

implemented in treePL[49]. We implemented three different methods for calibrating the trees to

assess similarity among results. Method 1: Dating with secondary calibrations only. We used the

95% credibility intervals of Lepidoptera ages from Figure S12 of Kawahara et al.[50] to assign

minimum and maximum ages to 27 ingroup and 6 outgroup nodes in our tree. Method 2: Dating

with fossils and one secondary root calibration. In this approach, we followed the guidelines of

Parham et al.[51] by calibrating nodes with 11 butterfly fossils that could be assigned to a butterfly

lineage's geological age with confidence as verified by de Jong[52]. None of the outgroup nodes

could be calibrated because the only reliable fossils associated with our non-butterfly

Lepidoptera were too young to influence the ages of the deeper nodes representing multi-

superfamily clades. Consequently, preliminary treePL analyses yielded highly dubious age

estimates for deep nodes on the tree, hundreds of millions of years older than expected based on

the literature. We therefore added a single secondary calibration to the root of the tree. Although

combining secondary and fossil calibrations in a single analysis can create redundancy that

negatively impacts the resulting age estimates[53], the limited fossil record of Lepidoptera made it

a necessity in order to obtain comparable results derived primarily from fossils. We ran two

versions of this method, each with a different root calibration. Method 2A used a maximum-age

estimate of 139.4 Ma, based on the angiosperm age estimate of Smith and Brown[54]. Method 2B

used a more conservative maximum-age estimate of 251 Ma, based on the older end of the

credibility interval for the age of angiosperms in Foster et al.[55]. Both of these calibrations were

used under the assumption that butterflies diverged from their moth ancestors after their most

frequent host-plants, angiosperms, were already present[56,57]. Method 3: Secondary calibrations

and six fossils. In this approach, we combined the 33 secondary calibrations from Method 1 with

six fossil calibrations, including some of the fossils used in Method 2. Fossils previously used to

time-calibrate trees of Kawahara et al.[50] were excluded from this analysis to avoid circularity and

redundancy with secondary calibrations. Whenever possible, redundant fossil calibrations from

Method 2 were replaced with calibrations from unrelated fossils that could be associated with a

different node in the same clade.

Diversification rate analyses.

579   We performed a Bayesian analysis of macroevolutionary mixtures using the program

BAMM v.1.10.4[58] to detect shifts in diversification rates between clades. The reversible-jump

markov chain Monte Carlo was run for 50 million generations and sampled every 50,000

generations. Priors were estimated with the R package BAMMtools v.2.1.6[59] using the command

'setBAMMpriors'. The tree was trimmed in Mesquite v.3.6[60] to remove all outgroups. Six

analyses were performed using different priors for expected numbers of shifts (5, 10, 20, 30, 40,

and 50 shifts).

Because BAMM has been criticized for incorrectly modeling rate shifts on extinct

lineages (i.e., extinct or non-sampled lineages inherit the ancestral diversification process and

cannot experience subsequent diversification-rate shifts[61,62]), we conducted a lineage-specific

birth-death shift analysis[63] in RevBayes[64]. We also utilized the R package castor v.1.6.9[65] to infer

deterministic lineage through time (dLTT), pulled speciation rate (PSR), and pulled

diversification rate (PDR) plots for Papilionoidea and six of its families, excluding Hedylidae.

Finally, we performed diversification analyses in TreePar v.3.3[66] and TESS v.2.1.0[67] to

determine relative likelihoods of models with different amounts of shifts in diversification and

turnover rates.

We conducted a series of diversification analyses to evaluate whether there is a

correlation between butterflies and plants. For this, we used HiSSE (Hidden State Speciation and

Extinction) and a BiSSE-like (Binary State Speciation and Extinction) implementation of

HiSSE[30] in the R package hisse[31]. We pruned outgroups from the tree (aa154 dated tree, Strategy

A) and compared 20 HiSSE models and BiSSE-like implementations of HiSSE. The BiSSE

equivalent of HiSSE tests whether there are different diversification rates associated with the two

host-plant use states. Other models were built in the HiSSE framework to test alternative

combinations of presence or absence of hidden state and host-plant use associations while also

considering different transition rate matrices, net turnover rates τ_i (speciation plus extinction: λ_i

+ μ_i), and extinction fractions ε_i (extinction divided by speciation: μ_i/λ_i) (Table S17). We tested

whether diversification rates were linked to feeding (A) as a larval specialist or generalist (Table

S18), (B) on Poales (Table S19) in Papilionoidea, Hesperiidae, and Nymphalidae, (C) on Fabales

(Table S20) in Papilionoidea and Nymphalidae, (D) on Brassicales (Table S21) in all butterflies

and Pieridae, (E) on Fagales (Table S22), (F) on Poaceae module (Table S23), (G) on Fabaceae

module (Table S24), and (H) on Fabaceae in Eudaminae (Tables S13, S24). We compared these

different models of HiSSE and BiSSE-like implementations to account for hidden states to

alleviate recent concerns regarding the reliability of SSE models and the high incidence of false

positive results[32].

Biogeographic analyses.

To assess the role of geography on diversification, we first aggregated data from multiple

sources to create a global butterfly checklist for each country. Primary data sources included: 1)

the *Lepidoptera and other life forms* database (http://ftp.funet.fi/index/Tree_of_life/insecta/

lepidoptera); 2) WikiSpecies (https://species.wikimedia.org); and 3) the type locality of each

species or subspecies in our list of valid butterfly names, which was obtained from 1, above. This

initial global checklist was vetted using published country checklists and the ButterflyNet Trait

Database (https://butterflytraits.org). Trait data from *ca*. 100 comprehensive and country-specific

field guides have been entered into this database, allowing us to generate species lists to cross-

validate checklists assembled[68].

We estimated the ancestral area of origin and geographic range evolution for butterflies

using two approaches: the ML approach of the DECX model[69] as implemented in the C++

version[70,71], available at https://github.com/champost/DECX), and with the program

BioGeoBEARS v.1.1.2[20]. We designated 14 biogeographic regions across the globe (Fig. S13,

Table S25), determined which of these regions were occupied by each species in our tree, and

developed a 14-state character matrix. DECX uses a time-calibrated tree, the modern distribution

of each species for a set of geographic areas, and a time-stratified geographic model that is

represented by connectivity matrices for specified time intervals spanning the evolutionary

history of clade of interest[72].

Because we could not estimate immigration and emigration rates in DECX, we also ran

BioGeoBEARS with 7 and 8 areas (Figs. S14, S15, Table S25). BioGeoBEARS analyses could

not be run with 14 states due to computational limitations due to the complexity of our dataset

(2,248 tree tips). The 7 and 8 bioregions largely correspond to the biogeographic realms defined

by Udvardy[73]. In BioGeoBEARS, we implemented both Dispersal Extinction Cladogenesis

(DEC) model[69,74] and the Likelihood equivalent of the Dispersal-Vicariance approach

(DIVALIKE)[75] models and different adjacency matrices (Data S4). Both approaches gave largely

consistent results, regardless of the model and parameters used (Tables S5, S26).

We performed biogeographic stochastic mapping to examine *in-situ* speciation,

immigration, and emigration between the 7-bioregions in BioGeoBEARS. We followed the

protocol of Li et al.[76] with 1,000 simulations with the DEC model. Relative mean dispersal rates

between all permutations of bioregions were calculated and presented in Figure 3 (see also Data

S5). These mean dispersal rates represent dispersal of butterfly lineages throughout the entire

evolutionary history of Papilionoidea, and thus cannot reveal changes in rates over time. In order

to look at historical biogeography of butterflies during different epochs, rates along all possible

inter-bioregion colonization rates were calculated at specific time intervals of 5 million years,

following Li et al.[76] (Table S27). These relative rates were then averaged to represent relevant

geological time periods and presented in Figures S2-S4.

Larval host plant analyses.

Larval host records were compiled from numerous sources (Table S8, Data S6). Given

the size of our host datasets and the scale of our analyses, we chose to examine relationships

between individual butterfly species and host families that are consumed by their larvae. Plant

families are commonly adopted as the taxonomic rank used for examining the evolution of host

use[77,78]. For each plant-feeding butterfly species in our tree, we quantified host-plant richness

and phylogenetic distance using six different metrics implemented in the package picante v.

1.8.2[79]. To calculate these metrics, we used the calibrated tree of seed plants from Smith and

Brown[54].

Because the number of host groups was too large for an ancestral state reconstruction

(nearly 50 host-plant orders, ~200 plant families plus insect associations), we first reduced the

number of host groups by using a network analysis. The Beckett algorithm[80], as implemented in

the function 'computeModules' from the package bipartite[81] in R v. 3.6.2[82], assigns plants and

butterflies to modules and computes the modularity index, Q. By maximizing Q, the algorithm

finds groups of butterflies and hosts that interact more with each other than with other taxa in the

network. Thus, host-plants that are assigned to the same module tend to be used by the same

butterflies. We found 13 modules for butterfly host associations in our module analysis (Table

S28, Table S17). We then conducted three larval host ancestral state reconstruction analyses

using stochastic character mapping with SIMMAP in phytools v.0.7-70[83] using the

'make.simmap' command. We reconstructed the ancestral state of (A) generalist versus specialist

feeding (2 states, Data S7), (B) plant, lichen, Hemiptera, or Hymenoptera as a food source (4

states, Data S8), and (C) plant module (13 states as described above, Data S9).

We examined the speciation rate of butterflies over time with maximum-likelihood birth-

death models to fit constant-rate (CST), time-dependent (TimeVar), and angiosperm-dependent

(AngioVar) models of diversification using the R package RPANDA v.1.9[84]. These models

jointly tested whether the diversification of angiosperms could have fostered the diversification

of butterflies in a single statistical framework. We conducted a series of diversification analyses

to evaluate whether there is a correlation between butterflies and plants. For this, we used HiSSE

(Hidden State Speciation and Extinction) and a BiSSE-like (Binary State Speciation and

Extinction) implementation of HiSSE in the R package hisse[30,31].

Acknowledgments

We thank M. Kuhn and E. Mavrodiev for assembling trait data. S. Epstein, T. Girard-Ang, P.

Pezzi, and L. Xiao assisted with lab work. J. Barber, M. Brownlee, S. Cinel, J. Daniels, R.

Godfrey, H. Gough, C. Hamilton, G. Hill, P. Houlihan, C. Huang, J. Miller, K. Miner, C. Mitter,

K. Mitter, A. Renevier-Faure, J. Rubin, M. Scallion, Y. Sondhi, and L. Wu provided specimens

or helped improve the manuscript. RAPiD Genomics (Gainesville, FL, USA) conducted

sequencing. K. Meusemann (1KITE) provided unpublished FcLM scripts. T. Barve, C. Couch,

H. Dansby, K. Casarella, R. Merritt, and X. Zheng helped create figures. High performance

clusters at Brigham Young University, Smithsonian Institution, University of Florida, and

Zoological Research Museum Alexander Koenig provided computational support.

Funding: Funding came from the US National Science Foundation (NSF) GoLife

"ButterflyNet" collaborative grant (DEB-1541500, 1541557, 1541560) to AK, RG, DL, and

NEP. Specimen collection and preservation was funded by NSF DBI-1349345, 1601369, DEB-

1557007 and IOS-1920895 (AYK), NSF DEB-1120380 (DJL), Grants 9285-13 and WW-227R-

17 from the National Geographic Society (DJL), NSF DBI-1256742 (AYK and KRW), NSF

DEB-0639861 (KRW) and NSF SES-0750480, DEB-0447244 and DEB-9615760 to NEP. ME

was supported by the Research Council of Norway (#204308), and the Hintelmann Scientific

Award for Zoological Systematics. FLC is supported by the European Research Council (ERC)

under the European Union's Horizon 2020 research and innovation programme (project GAIA,

#851188). RV was supported by the Spanish Ministry of Science and Innovation grant PID2019-

107078GB-I00/AEI/10.13039/501100011033. GT was supported by the Spanish Ministry of

Science and Innovation (grants PID2020-117739GA-I00/AEI/ 10.13039/501100011033 and

RYC2018-025335-I). VL was supported by the Russian Science Foundation grant 19-14-00202.

MY was supported by MEXT KAKENHI #19916010 and JSPS KAKENHI Grant #13010131,

23570111, 26440207, 17K07528 and 21H02215. ABBM, HPR and NOM were supported by

CNPQ grants Proc #563332/2010-7 and 304273/2014-7. We are thankful for continuous support

from Putnam Expedition grants from the Museum of Comparative Zoology. We acknowledge

Research Computing at the University of Florida and the Faculty of Arts and Sciences, Harvard

University, for providing computational resources (https://www.rc.ufl.edu,

https://www.rc.fas.harvard.edu).

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: All data in the main text, supplementary materials, or in

gzipped data archives will be made available at the Dryad Digital Repository

(www.datadryad.org.xxxx) upon acceptance.

Author contributions:

Analysis: APSC, AYK, CE, CS, DMP, EAE, EFAT, FLC, HLO, JWB, ME, MPB, PBF, RAS,
XL

Conceptualization: AYK, DJL, LR, JWB, ME, NEP, RG

Funding acquisition: AYK, DJL, MJZ, NEP, RG, WJ

Data assembly: CS, DJL, DMP, EAE, EAL, FLC, HW, JH, JWB, LR, ME, MI, MN, NEP, RG,
VB, VS, WAVM

Methodology: CW, EH, FTGS, HP, JLS, JRK, JWB

Project administration: AYK, DJL, NEP

Sampling: ABBM, APSC, AYK, DJL, EFAT, FH, GT, HPR, JIM, JWB, ME, MY, NEP, RAS,
RV, PV, ZFF

Sequence workflow: JWB, ME

Specimen identification and preparation: ADW, AM, APSC, AYK, DJL, HPR, JIM, JVL, KMD,
ME, MY, NOM, RAS, RM, SCM, TD, YLN, ZFF

Supervision: AYK, DJL, JWB, ME, NEP, RG

Taxonomy and curation: DJL, GL, RG, VB

Visualization: AM, APSC, AYK, DJL, FLC, XL

Writing – original draft: APSC, AYK, DJL, DMP

Writing – review & editing: All authors

741
Fig. 1. Evolutionary relationships and diversification patterns of butterflies. Time-calibrated tree
based on 2,244 species, 391 loci, and 150 amino acid partitions. Branches show distinct shift
configurations (circles) as estimated by clade-specific diversification models. Letters at nodes
refer to clades with significant rate shifts (See section 6 of Supplementary Text). Colored lines in
the outer ring beside tips indicate association with one of the 13 host modules (see section 17 of
Extended Online Methods). Black lines in the host-association ring are species without data. The
inset graph shows the posterior probability distribution, with alternative and expected number of
diversification rate shifts, converging around 30.
750

Fig. 2. Global butterfly diversification over time. Butterfly diversity increased well after the origin of flowering plants. Colored lines with dots show butterfly diversity compared to vascular plant diversity from three widely accepted, recent plant diversification studies. Mean global temperature and major geological events during the last 350 Ma were calculated from Scotese et al.[85]. The angiosperm stem to clade credibility interval is a consensus of seven studies (see section 14 of Extended Online Methods).

Fig. 3. Relative mean dispersal rates of butterflies between different bioregions. Numbers associated with each arrow are the average rates from 1000 simulations using biogeographical stochastic mapping in BioGeoBEARS, which were then divided by 100 for ease of comparison (raw values can be found in Data S5).

Fig. 4. Distribution of ancestral butterflies over time. Each map corresponds to a 15 Ma interval of butterfly evolution. Bioregion color indicates the number of lineages in the Papilionoidea phylogeny that are associated with that bioregion during that time period, as determined by the BioGeoBears ASR.