Long title: Nitrogenase resurrection and the evolution of a singular

enzymatic mechanism

Short title: Nitrogenase resurrection and evolution

Amanda K. Garcia[a], Derek F. Harris[b], Alex J. Rivier[a], Brooke M. Carruthers[a], Azul Pinochet-

Barros[a], Lance C. Seefeldt[b], Betül Kaçar[a]*

[a] Department of Bacteriology, University of Wisconsin–Madison, Madison, WI 53706

[b] Department of Chemistry and Biochemistry, Utah State University, Logan, UT 84322

*To whom correspondence should be addressed: Betül Kaçar, [602 263 3622], bkacar@wisc.edu

Keywords: ancestral sequence reconstruction, evolutionary biochemistry, metabolic

engineering, nitrogen fixation, nitrogenase

ABSTRACT

The planetary biosphere is powered by a suite of key metabolic innovations that emerged early in the history of life. However, it is unknown whether life has always followed the same set of strategies for performing these critical tasks. Today, microbes access atmospheric sources of bioessential nitrogen through the activities of just one family of enzymes, nitrogenases. Here, we show that the only dinitrogen reduction mechanism known to date is an ancient feature conserved from nitrogenase ancestors. We designed a paleomolecular engineering approach wherein ancestral nitrogenase genes were phylogenetically reconstructed and inserted into the genome of the diazotrophic bacterial model, *Azotobacter vinelandii,* enabling an integrated assessment of both *in vivo* functionality and purified nitrogenase biochemistry. Nitrogenase ancestors are active and robust to variable incorporation of one or more ancestral protein subunits. Further, we find that all ancestors exhibit the reversible enzymatic mechanism for dinitrogen reduction, specifically evidenced by hydrogen inhibition, that is also exhibited by extant *A. vinelandii* nitrogenase isozymes. Our results suggest that life may have been constrained in its sampling of protein sequence space to catalyze one of the most energetically challenging biochemical reactions in nature. The experimental framework established here is essential for probing how nitrogenase functionality has been shaped within a dynamic, cellular context to sustain a globally consequential metabolism.

IMPACT STATEMENT

The enzymatic mechanism for dinitrogen reduction is an ancient feature of nitrogenases that persisted over hundreds of millions of years.

INTRODUCTION

The evolutionary history of life on Earth has generated tremendous ecosystem diversity, the sum of which is orders of magnitude larger than that which exists at present [1]. Life's historical diversity provides a measure of its ability to solve adaptive problems within an integrated planetary system. Accessing these solutions requires a deeper understanding of the selective forces that have shaped the evolution of molecular-scale, metabolic innovations. However, the early histories of many of life's key metabolic pathways and the enzymes that catalyze them remain coarsely resolved.

Important efforts to advance understanding of early metabolic innovations have included phylogenetic inference [2, 3], systems-level network reconstructions [4], and the leveraging of extant or mutant biological models as proxies for their ancient counterparts [5-7]. However, methods to directly study these metabolic evolutionary histories across past environmental and cellular transitions remain underexplored. A unified, experimental strategy that integrates historical changes to enzymes, which serve as the primary interface between metabolism and environment, and clarifies their impact within specific cellular and physiochemical contexts is necessary. To address this, phylogenetic reconstructions of enzymes can be directly integrated within laboratory microbial model systems [8-10]. In this paleomolecular framework, predicted ancestral enzymes can be "resurrected" within a compatible host organism for functional characterization. These experimental systems can ultimately integrate multiple levels of historical analysis by interrogating critical features of ancient enzymes as well as dynamic interactions between enzymes, their broader metabolic networks, and the external environment.

The study of biological nitrogen fixation offers a promising testbed to thread these investigations of early metabolic evolution. Both phylogenetic and geological evidence [11-15] indicate that the origin of biological nitrogen fixation was a singular and ancient evolutionary event on which the modern biosphere has since been built. The only known nitrogen fixation pathway (compared to, for example, at least seven carbon-fixation pathways [16]) is catalyzed by an early evolved family of metalloenzymes called nitrogenases that reduce highly inert, atmospheric dinitrogen (N_2) to bioavailable ammonia (NH_3). The nitrogenase family comprises three isozymes that vary in their metal dependence (i.e., molybdenum, vanadium, and iron) and, in certain cases, all coexist within the same host organism [17]. Many diazotrophs depend on genetic strategies for coordinating the biosynthesis and expression of multiple nitrogenase isozymes and their respective

metalloclusters [18, 19], and, in oxic environments, protecting the oxygen-sensitive
metalloclusters from degradation [20]. Thus, nitrogenase enzymes are a central component of a
broader, co-evolving nitrogen fixation machinery. These features that create significant
experimental challenges for nitrogen fixation engineering [21-23] nevertheless also make this
metabolism an ideal candidate for systems-level, paleomolecular study.

How biological nitrogen fixation emerged and evolved under past environmental conditions is still
poorly constrained relative to its importance in Earth's planetary and biological history. Because
nitrogen has been a limiting nutrient over geological timescales [24, 25], nitrogenase has long
been a key constituent of the expanding Earth biosphere. The impact of nitrogen limitation is
underscored by human reliance on the industrial Haber-Bosch process, an energetically and
environmentally costly workaround for nitrogen fertilizer production [26] designed to supplement
a remarkable molecular innovation that biology has tinkered with for more than three billion years.
How the structural domains and regulatory network of nitrogenase were recruited [27, 28] and
under what selective pressures the metal dependence of nitrogenases were shaped [11, 29]
remain open questions. Importantly, it is not known how the enzymatic mechanism for dinitrogen
reduction has been tuned by both peptide and metallocluster to achieve one of the most difficult
reactions in nature [30-32]. At the enzyme level, previous insights into nitrogenase sequence-
function relationships have primarily derived from single or dual substitution studies. These have
often yielded diminished or abolished nitrogenase activity [30, 31], though in certain cases
improved reactivity toward alternate, industrially relevant substrates [30]. Despite illuminating key
features of extant nitrogenase mechanism in select model organisms, the combination of detailed
functional studies within an explicit evolutionary scheme has not previously been accomplished
for the nitrogen fixation system.

Here, we seek guidance from the Earth's evolutionary past to reconstruct the history of the key
metabolic enzyme, nitrogenase. We establish an evolutionary systems biology approach for the
cellular- and molecular-level characterization of ancestral nitrogenases resurrected within the
model diazotrophic bacterium, *Azotobacter vinelandii* (*A. vinelandii*). We find that variably
replacing different protein subunits of the nitrogenase complex with inferred ancestral
counterparts enables nitrogen fixation in *A. vinelandii*. Purified ancestral enzymes exhibit the
specific N_2 reduction mechanism retained by their studied, extant counterparts, and maintain the
same catalytic selectivity between N_2 and protons. Thus, the core strategy for biological nitrogen
fixation is conserved across the investigated timeline. Our paleomolecular approach opens a new

route to study the ancient functionality and evolution of nitrogenases both deeper in its ancestry
and within the broader context of its supporting, cellular machinery.

RESULTS

A resurrection strategy for ancestral nitrogenases

We designed an engineering pipeline for the resurrection and experimental characterization of
ancestral nitrogenases (**Fig. 1***A*). In this scheme, phylogenetically inferred, ancestral nitrogenase
genes are synthesized and engineered into the genome of a modern diazotrophic bacterium,
enabling the assessment of *in vivo* nitrogenase activity and expression in parallel with biochemical
analysis of purified enzyme. The engineering and functional assessment of ancestral
nitrogenases required a suitable diazotrophic microbial host, owing to challenges associated with
nitrogenase heterologous expression [26]. We selected the obligately aerobic
gammaproteobacterium, *A. vinelandii* (strain "DJ"), an ideal experimental model due to its genetic
tractability and the availability of detailed studies on the genetics and biochemistry of its nitrogen
fixation machinery [33]. We specifically targeted the extant *A. vinelandii* molybdenum-dependent
("Mo-") nitrogenase (hereafter referred to as wild-type, "WT"), which is the best studied isozyme
[30] relative to the *A. vinelandii* vanadium ("V-") and iron ("Fe-") dependent nitrogenases . The
WT Mo-nitrogenase complex comprises multiple subunits, NifH, NifD, and NifK (encoded by
nifHDK genes), which are arranged into two catalytic components: a NifH homodimer and a NifDK
heterotetramer (**Fig. 1***B*). During catalysis, both components transiently associate to transfer one
electron from NifH to NifDK and subsequently dissociate. Transferred electrons accumulate at the
active-site Mo-containing metallocluster ("FeMoco") housed within the NifD subunits for reduction
of N_2 substrate to NH_3.

To infer Mo-nitrogenase ancestors, we built a maximum-likelihood nitrogenase phylogeny from a
concatenated alignment of NifHDK amino acid sequences (**Fig. 2***A***; Fig. 2-Figure supplement**
1). The phylogeny contains 385 sets of homologs representative of known nitrogenase molecular
sequence diversity (including Mo-, V-, and Fe-nitrogenases), and is rooted by dark-operative
protochlorophyllide oxidoreductase proteins classified within the nitrogenase superfamily [34]. For
this study, we selected ancestors that fall within the direct evolutionary lineage of *A. vinelandii*
WT (**Fig. 2***B*), "Anc1" and "Anc2" (listed in order of increasing age), having ~90% and ~85% amino
acid sequence identity to WT across the full length of their concatenated NifHDK proteins,

respectively (**Fig. 2C; Supplementary File 1a**). A relatively conservative percentage identity

threshold was chosen based on prior studies benchmarking functional expression of ancestral

elongation factor proteins in *Escherichia coli* [35] and *Synechococcus elongatus* [10]. The high-

dimensional, nitrogenase protein sequence space occupied by both extant and ancestral

homologs is visualized in two-dimensions in **Fig. 2D** by machine-learning embeddings (see

Materials and methods). This analysis highlights the swath of sequence space targeted here,

as well as that made accessible by resurrection of nitrogenase ancestors more broadly. WT,

Anc1, and Anc2 lie within a Mo-nitrogenase clade (previously termed "Group I" [12]) that contains

homologs from diverse aerobic and facultatively anaerobic taxa, including proteobacteria and

cyanobacteria (**Fig. 2A**). A maximum age constraint of ~2.5 Ga for Group I nitrogenases (and

thus for both Anc1 and Anc2) can be reasoned based on timing of the Great Oxidation Event [36]

and downstream emergence of aerobic taxa represented nearly exclusively within this clade.

Residue-level differences between ancestral and WT nitrogenases ("ancestral substitutions") are

broadly distributed along the length of each ancestral sequence (**Fig. 1B; Fig. 2-Figure**

Supplement 2). An ancestral substitution proximal to the active-site FeMoco metallocluster lies

within a loop considered important for FeMoco insertion [37] (NifD I355V; residue numbering from

WT)) and is observed across all targeted NifD ancestors (**Fig. 1B**). Other ancestral substitutions

are notable for their location at relatively conserved residue sites (assessed by ConSurf [38]; **Fig.**

2-Figure Supplement 3; see **Materials and methods**) and/or within subunit interfaces, including

two at the NifD:NifK interface that are proximal to one another, F429Y (NifD) and R108K (NifK).

The Anc2 NifD protein contains five more ancestral substitutions at conserved sites than the

younger Anc1 NifD protein. In all studied ancestors, the C275 and H442 FeMoco ligands, as well

as other strictly conserved nitrogenase residues, are retained.

Phylogenetic analysis informs the compatibility of selected ancestors in extant microbial hosts.

Extant nitrogenases within the Group I nitrogenase clade (which include Anc1 and Anc2

descendants) are associated with numerous accessory genes likely recruited to optimize

synthesis and regulation of the oxygen-sensitive nitrogenase for aerobic or facultative

metabolisms [27]. For example, in addition to the structural *nifHDK* genes, *A. vinelandii* WT is

assembled and regulated with the help of >15 additional *nif* genes. Likewise, the extant

descendants of Anc1 and Anc2 are primarily aerobic or facultative proteobacteria and are thus

associated with higher complexity *nif* gene clusters (**Fig. 2B**). We hypothesized that the likely

oxygen-tolerant, ancient proteobacterium harboring these ancestral nitrogenases were similar in

nif cluster complexity to extant *A. vinelandii*. Thus, we predicted that the *nif* accessory genes

present in *A. vinelandii* would support the functional expression of resurrected nitrogenase

ancestors.

To gauge the degree of compatibility between ancestral and extant nitrogenase proteins and

maximize the chance of recovering functional nitrogenase variants, we executed two parallel

genome engineering strategies. First, we constructed *A. vinelandii* strains harboring only the

ancestral *nifD* gene from both targeted ancestral nodes ("Anc1A" and "Anc2"), thereby expressing

"hybrid ancestral-WT" nitrogenase complexes (**Fig. 1C**). This strategy is similar to *in vitro* "cross-

reaction" studies that have evaluated the compatibility of nitrogenase protein components from

differing host taxa [39]. Second, we constructed a strain harboring all Anc1 *nifHDK* genes,

expressing a fully ancestral nitrogenase complex ("Anc1B"; sequence reconstructed from a node

equivalent to Anc1A from an alternate phylogeny, see **Materials and methods**). *A. vinelandii*

strains were constructed by markerless genomic integration of ancestral nitrogenase genes, as

described in **Materials and methods**.

Ancestral nitrogenases enable diazotrophic microbial growth

All *A. vinelandii* constructs harboring ancestral genes enabled diazotrophic growth in

molybdenum-containing, nitrogen-free media. All strains had comparable doubling times to WT

during the exponential phase ($p > .05$; **Fig. 3A, 3B**). The only significant difference among strains

was a ~14-h increase in the lag phase of strain Anc2 relative to WT, harboring the oldest

nitrogenase ancestor ($p \approx$ 2e-7). We did not detect growth under the same conditions for a control

Δ*nifD* strain (DJ2278, see **Supplementary File 1c**). This result confirmed that the growth

observed for ancestral strains did not stem from leaky expression of the alternative, V- or Fe-

dependent nitrogen fixation genes in *A. vinelandii,* which were left intact.

An acetylene reduction assay was performed to measure cellular nitrogenase activity in

engineered strains. This assay quantifies the reduction rate of the non-physiological substrate

acetylene (C_2H_2) to ethylene (C_2H_4) [40], here normalized to total protein content. *A. vinelandii*

strains harboring only ancestral *nifD* (Anc1A, Anc2) exhibited mean C_2H_2 reduction rates of ~5 to

6 μmol C_2H_4/mg total protein/h, ~40-45% that of WT ($p \approx$ 6e-4 and $p \approx$ 3e-4, respectively) (**Fig.**

3C). Strain Anc1B, harboring ancestral *nifHDK,* exhibited a mean acetylene reduction rate of ~9

μmol C_2H_4/mg total protein/h, ~70% that of WT ($p \approx$ 3e-2).

The phenotypic variability we observed among engineered and WT *A. vinelandii* strains might result both from differences in nitrogenase expression and nitrogenase activity. To provide insights into these disparate effects, we quantified nitrogenase protein expression in engineered strains by immunodetection of ancestral and WT Strep-tagged NifD proteins (the latter from strain DJ2102, see **Supplementary File 1c**) and did not conclusively detect significant differences in protein quantity relative to WT ($p > .05$; **Fig. 3D**).

Purified ancestral nitrogenases conserve extant N_2 reduction mechanism and efficiency

Ancestral nitrogenase NifDK protein components were expressed and purified for biochemical characterization. All ancestral NifDK proteins were assayed with WT NifH protein (ancestral NifH proteins were not purified) for reduction of H^+, N_2, and C_2H_2. Ancestors were found to reduce all three substrates *in vitro*, supporting the cellular-level evidence of ancestral nitrogenase activity (**Fig. 4A**).

We investigated whether the ancestral nitrogenases studied here would exhibit the general mechanism for N_2 binding and reduction that has been observed for the studied, extant nitrogenase isozymes of *A. vinelandii* (Mo, V, and Fe) [32, 41]. This mechanism involves the accumulation of 4 electron/protons on the active-site cofactor as metal-bound hydrides, generating the $E_4(4H)$ state (**Fig. 4B**). Once generated, N_2 can bind to the $E_4(4H)$ state through a reversible reductive elimination/oxidative addition (*re/oa*) mechanism, which results in the release (*re*) of a single molecule of hydrogen gas (H_2). N_2 binding is reversible in the presence of sufficient H_2, which displaces bound N_2 and results in reformation of $E_4(4H)$ with two hydrides (*oa*). Thus, a classic test of the (*re/oa*) mechanism is the ability of H_2 to inhibit N_2 reduction. We observed that the reduction of N_2 to NH_3 for all nitrogenase ancestors was inhibited in the presence of H_2, indicating that the ancestors follow the same mechanism of N_2 binding determined for extant enzymes (**Fig. 4C**).

In the event the $E_4(4H)$ state fails to capture N_2, nitrogenases will simply produce H_2 from the $E_4(4H)$ state to generate the $E_2(2H)$ state. The ratio of H_2 formed to N_2 reduced (H_2/N_2) can be used as a measure of the efficiency of nitrogenases in using ATP and reducing equivalents for N_2 reduction. The stoichiometric minimum of the mechanism is $H_2/N_2 = 1$. Experimentally (under 1 atm N_2), a ratio of ~2 is seen for Mo-nitrogenase and ~5 and ~7 for V- and Fe-nitrogenase,

respectively [32]. H_2/N_2 values for all ancestors under 1 atm N_2 was ~2, similar to extant Mo-
nitrogenase (**Fig. 4D**).

DISCUSSION

In this study, we leverage a new approach to investigate ancient nitrogen fixation by the
resurrection and functional assessment of ancestral nitrogenase enzymes. We demonstrate that
engineered *A. vinelandii* cells can reduce N_2 and C_2H_2 and exhibit diazotrophic growth rates
comparable to WT, though we observe that the oldest ancestor, Anc2, has a significantly longer
lag phase. Purified nitrogenase ancestors are active for reduction of H^+, N_2, and C_2H_2, while
maintaining the catalytic efficiency (described by the H_2/N_2 ratio) of WT enzymes. Our results also
show that ancestral N_2 reduction is inhibited by H_2, indicating an early emergence of the reductive-
elimination N_2-reduction mechanism preserved by characterized, extant nitrogenases [32, 41].
These properties are maintained despite substantial residue-level changes to the peripheral
nitrogenase structure (including relatively conserved sites), as well as a handful within the active-
site or protein-interface regions within the enzyme complex.

It is important to consider that the nitrogenase ancestors resurrected here represent hypotheses
regarding the true ancestral state. Uncertainty underlying ancestral reconstructions might derive
from incomplete extant molecular sequence data, as well as incorrect assumptions associated
with the implemented evolutionary models [8]. For example, a complicating feature of nitrogenase
evolution is that it has been shaped significantly by horizontal gene transfer [12, 15], which in
certain cases has led to different evolutionary trajectories of individual nitrogenase structural
genes. Specifically, certain H-subunit genes of the V-nitrogenase appear to have different
evolutionary histories relative to their DK components [12]. However, since our study targets a
Mo-nitrogenase lineage, we do not expect horizontal transfer to be a significant source of
uncertainty in our reconstructions.

The N_2-reduction activity of nitrogenase ancestors suggests that the required protein-protein
interactions—both between subunits that comprise the nitrogenase complex as well as those
required for nitrogenase assembly in *A. vinelandii*—and metallocluster interactions are sufficiently
maintained for primary function. Still, our results reveal the degree to which the organism-level
phenotype of host strains can be perturbed by varying both the number and age of ancestral
subunits. Importantly, these changes appear to impact phenotypic properties in complex ways,

representative of the type of cellular constraints on nitrogenase evolution that would be
unobservable through an *in vitro* study alone. For example, we observed comparable growth
characteristics of strains harboring single (Anc1A, ancestral NifD) versus multiple (Anc1B,
ancestral NifHDK) ancestral subunits of equivalent age, whereas the lag phase of Anc2 hosting
a single, older subunit (ancestral NifD) was increased. Here, growth is sensitive to older, ancestral
substitutions in a single subunit while permissive of more recent ancestral substitutions across
one or more subunits within the nitrogenase complex. However, a different pattern is observed
across *in vivo* acetylene reduction rates. These are most negatively impacted relative to WT in
strains with a single NifD ancestor (Anc1A, Anc2), whereas rates are more modestly decreased
in a strain with a complete, ancestral NifHDK complex (Anc1A). These results suggest that
ancestral subunits of equivalent age have greater compatibility and yield greater *in vivo* activity
compared to subunits of disparate ages, perhaps owing to modified protein interactions within the
nitrogenase complexes The discrepancy between *in vivo* activity and growth characteristics may
also be attributable to impacted cellular processes external to the biochemical properties of the
nitrogenase complex itself, and yet nevertheless vital in determining the overall fitness of the host
organism. Finally, though we do not detect significant differences in ancestral protein expression
here, it is possible that phenotypic outcomes of future reconstructions might be impacted by
perturbed expression levels (e.g., [10, 42]). To what degree these expression levels are
representative of the ancestral state and impact the phenotypic property of interest should be
considered in future work.

That nitrogenase ancestors perform the reductive-elimination N_2-reduction mechanism—as the
distantly related [11], extant Mo-, V-, and Fe-nitrogenases of *A. vinelandii* do today [32]—likely
indicates that this enzymatic characteristic was set early in nitrogenase evolutionary history and
sustained through significant past environmental change [36, 43, 44] and ecological diversification
[27, 45]. It is possible that life's available strategies for achieving N_2 reduction may be
fundamentally limited, and that a defining constraint of nitrogenase evolution has been the
preservation of the same N_2 reduction mechanism across shifting selective pressures. For
example, in the acquisition of V- and Fe-dependence from Mo-dependent ancestors [11],
nitrogenases may have required substantial sequence and structural changes [46, 47] in order to
facilitate reductive-elimination given a different active-site metallocluster. It is also possible that
alternate strategies for biological nitrogen fixation evolved early in the history of life and were
subsequently outcompeted, leaving no trace of their existence in extant microbial genomes. Why
these alternate possibilities were evidently not explored by nature to the same degree remains an

open question, particularly given the several abiotic mechanisms for nitrogen fixation [48-50] and
the multiple biological pathways for another, globally significant metabolism, carbon fixation [16].
Because our paleomolecular approach is ultimately informed by extant sequence data, it cannot
directly evaluate extinct sequences that, for instance, due to contingency or entrenchment, did
not persist and become preserved in extant microbial genomes. Nevertheless, evolutionarily
informed studies of nitrogenase functionality that define the sequence-function space of this
enzyme family will provide a foundation for laboratory efforts aimed toward exploring alternate
scenarios. Future work that explores deeper into nitrogenase evolutionary history (and across
extant and ancestral nitrogenase sequence space, as charted here (**Fig. 2*D***)) will clarify the
degree of functional constraint exhibited by the nitrogenase family, both past and present.
323
324   **CONCLUSION**
325
Broadening the historical level of analysis beyond a single enzyme to the organism level is
necessary to generate comprehensive insights into the evolutionary history and engineering
potential of nitrogen fixation. Paleomolecular work that has expanded toward the systems-level
investigation of early evolved, crucial metabolic pathways remains in its infancy, despite the
potential for provocative connections between molecular-scale innovations and planetary history
[8, 10]. Our results highlight the evolutionary conservation of a critical metabolic pathway that has
shaped the biosphere over billions of years, as well as establish the tractability of leveraging
phylogenetic models to carry out extensive, empirical manipulations of challenging enzymatic
systems and their microbial hosts. Building on the empirical framework presented here will
illuminate the evolutionary design principles behind ancient metabolic systems more broadly as
well as leverage these histories to understand how key enzymes that allowed organisms to
access nitrogen from the atmosphere evolved.
338
339

340 **MATERIALS AND METHODS**

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
strain, strain background (*Azotobacter vinelandii*)	DJ	DOI: 10.1128/JB.00504-09	n/a	Dennis Dean, Virginia Tech; Wild-type (WT); Nif+
genetic reagent (*Azotobacter vinelandii*)	DJ2102	DOI: 10.1016/bs.mie.2018.10.007	n/a	Dennis Dean, Virginia Tech; Strep-tagged WT NifD; Nif+
genetic reagent (*Azotobacter vinelandii*)	DJ2278	Other	n/a	Dennis Dean, Virginia Tech; Δ*nifD*::KanR; Nif-
genetic reagent (*Azotobacter vinelandii*)	DJ884	Other	n/a	Dennis Dean, Virginia Tech; *nifD*R187I mutant; Nif+(slow); overexpresses NifH
genetic reagent (*Azotobacter vinelandii*)	AK022	This paper	n/a	Δ*nifHDK*::KanR; Nif-

genetic reagent (*Azotobacter vinelandii*)	AK013	This paper	n/a	"Anc1A"; Δ*nifD*::*nifD*^{Anc1A}; Nif+
genetic reagent (*Azotobacter vinelandii*)	AK023	This paper	n/a	"Anc1B"; Δ*nifHDK*::*nifHDK*^{Anc1B}; Nif+
genetic reagent (*Azotobacter vinelandii*)	AK014	This paper	n/a	"Anc2"; Δ*nifD*::*nifD*^{Anc2}; Nif+
antibody	StrepMAB-Classic (Mouse monoclonal)	IBA Lifesciences	Cat# 2-1507-001, RRID: AB_513133	WB (1:5000)
recombinant DNA reagent	pAG25	This paper	n/a	KanR cassette (APH(3')-I gene) + 400-bp *nifHDK* flanking homology regions, synthesized into XbaI/KpnI sites in pUC19; used to construct strain AK022 from DJ
recombinant DNA reagent	pAG13	This paper	n/a	*nifD*^{Anc1A} + 400-bp *nifD* flanking homology regions, synthesized into XbaI/KpnI sites in pUC19; used to construct strain Anc1A from AK022

recombinant DNA reagent	pAG19	This paper	n/a	*nifHDK*[Anc1B] + 400-bp *nifHDK* flanking homology regions, synthesized into XbaI/KpnI sites in pUC19; used to construct strain Anc1B from AK022
recombinant DNA reagent	pAG14	This paper	n/a	*nifD*[Anc2] + 400-bp *nifD* flanking homology regions, synthesized into XbaI/KpnI sites in pUC19; used to construct strain Anc2 from AK022
sequence-based reagent	306_nifH_F	This paper	PCR primers	GCCGAACGT TCAAGTGGAA A
sequence-based reagent	307_nifH_R	This paper	PCR primers	AGAGCCAATC TGCCCTGTC
sequence-based reagent	308_nifD_F	This paper	PCR primers	CACCCGTTAC CCGCATATGA

sequence-based reagent	309_nifD_R	This paper	PCR primers	ACTCATCTGTGAACGGCGTT
sequence-based reagent	310_nifK_F	This paper	PCR primers	GCTAACGCCGTTCACAGATG
sequence-based reagent	311_nifK_R	This paper	PCR primers	TCAGTTGGCCTTCGTCGTTG
software, algorithm	MAFFT	MAFFT	RRID:SCR_011811	
software, algorithm	trimAl	trimAl	RRID:SCR_017334	
software, algorithm	IQ-TREE	IQ-TREE	RRID:SCR_017254	
software, algorithm	RAxML	RAxML	RRID:SCR_006086	
software, algorithm	PAML	PAML	RRID:SCR_014932	

software, algorithm	MODELLER	MODELLER	RRID:SCR_0 08395	
software, algorithm	ChimeraX	ChimeraX	RRID:SCR_0 15872	
software, algorithm	Growthcurve r	Growthcurver	n/a	R package

Nitrogenase ancestral sequence reconstruction and selection

The nitrogenase protein sequence dataset was assembled by BLASTp [51] search of the NCBI non-redundant protein database (accessed August 2020) with *A. vinelandii* NifH (WP_012698831.1), NifD (WP_012698832.1), and NifK (WP_012698833.1) queries and a 1e-5 Expect value threshold (**Supplementary File 1b**). BLASTp hits were manually curated to remove partially sequenced, misannotated, and taxonomically overrepresented homologs. BLASTp hits included protein sequences from homologous Mo-, V-, and Fe-nitrogenase isozymes [11]. H-, D, and K-subunit sequences from these isozymes were individually aligned by MAFFT v7.450 [52] and concatenated along with outgroup dark-operative protochlorophyllide oxidoreductase sequences (Bch/ChlLNB). The final dataset included 385 nitrogenase sequences and 385 outgroup sequences. For sequences used to construct Anc1A and Anc2 (internal nodes #960 and #929, respectively), tree reconstruction (using a trimmed alignment generated by trimAl v1.2 [53]) and ancestral sequence inference (using the initial untrimmed alignment) were both performed by RAxML v8.2.10 [54] with the LG+G+F evolutionary model (model testing performed by the ModelFinder [55] in the IQ-TREE v.1.6.12 package [56]).

Due to concerns that RAxML v.8.2 does not implement full, marginal ancestral sequence reconstruction as described by Yang et al. [57], we performed a second phylogenetic analysis as follows. The extant sequence dataset described above was realigned by MAFFT (untrimmed) and

364 tree reconstruction was again performed by RAxML. Ancestral sequence reconstruction was
365 instead performed by PAML v4.9j [58] using the LG+G+F model. From this second reconstruction,
366 Anc1B (internal node #1312), equivalent to Anc1A, was selected for experimental analysis. Anc1B
367 and Anc1A have identical sets of descendent homologs, and their NifD proteins are 95% identical
368 (**Fig. 2*B***).
369
370 Only the ancestral sequences inferred with the most probable residue at each protein site were
371 considered for this study (mean posterior probabilities of targeted nitrogenase subunits range
372 from .95 to .99; see **Supplementary File 1a**). All ancestral sequences were reconstructed from
373 well-supported clades (SH-like aLRT = 99-100 [59]).
374
375 **Ancestral nitrogenase structural modeling and sequence analysis**
376
377 Structural homology models of ancestral sequences were generated by MODELLER v10.2 [60]
378 using PDB 1M34 as a template for all nitrogenase protein subunits and visualized by ChimeraX
379 v1.3 [61].
380
Extant and ancestral protein sequence space was visualized by machine-learning embeddings,
where each protein embedding represents protein features in a fixed-size, multidimensional
vector space. The analysis was conducted on concatenated (HDK) nitrogenase protein
sequences in our phylogenetic dataset. The embeddings were obtained using the pre-trained
language model ESM2 [62, 63], a transformer architecture trained to reproduce correlations at
the sequence level in a dataset containing hundreds of millions of protein sequences. Layer 33 of
this transformer was used, as recommended by the authors. The resulting 1,024 dimensions were
reduced by UMAP [64] for visualization in a two-dimensional space.
389
390 Protein site-wise conservation analysis was performed using the Consurf server [38]. An input
391 alignment containing only extant, Group I Mo-nitrogenases was submitted for analysis under
392 default parameters. Conserved sites were defined by a Consurf conservation score >7.
393
394 ***A. vinelandii* strain engineering**
395
396 Nucleotide sequences of targeted ancestral nitrogenase proteins were codon-optimized for *A.*
397 *vinelandii* by a semi-randomized strategy that maximized ancestral nucleotide sequence identity

398 to WT genes. Ancestral and WT protein sequences were compared using the alignment output of

399 ancestral sequence reconstruction (RAxML or PAML). For sites where the ancestral and WT

400 residues were identical, the WT codon was assigned. At sites where the residues were different,

401 the codon was assigned randomly, weighted by *A. vinelandii* codon frequencies (Codon Usage

402 Database, https://www.kazusa.or.jp/codon/). Nucleotide sequences were synthesized into

403 XbaI/KpnI sites of pUC19 vectors (unable to replicate in *A. vinelandii*) (Twist Bioscience;

404 GenScript). Inserts were designed with 400-base-pair flanking regions for homology-directed

405 recombination at the relevant *A. vinelandii nif* locus. An "ASWSHPQFEK" Strep-II-tag was

406 included at the N-terminus of each synthetic *nifD* gene for downstream NifD immunodetection

407 and NifDK affinity purification. See **Supplementary File 1c** for a list of strains and plasmids used

408 in this study.

410 Engineering of *A. vinelandii* strains used established methods, following Dos Santos [65]. *A.*

411 *vinelandii* WT ("DJ"), DJ2278 (Δ*nifD::*KanR), DJ2102 (Strep-II-tagged WT NifD), and DJ884 (NifH-

412 overexpression mutant) strains were generously provided by Dennis Dean (Virginia Tech)

413 (**Supplementary File 1c**). Strains Anc1A and Anc2 were constructed from the DJ2278 parent

414 strain via transformation with plasmids pAG13 and pAG19, respectively (**Supplementary File**

415 **1c**). For construction of strain Anc2, we first generated a Δ*nifHDK* strain, AK022, by transforming

416 the DJ strain with pAG25. Genetic competency was induced by subculturing relevant parent

417 strains in Mo- and Fe-free Burk's medium (see below) . Competent cells were transformed with

418 at least 1 µg of donor plasmid. Transformants were screened on solid Burk's medium for rescue

419 of the diazotrophic phenotype ("Nif+") and loss of kanamycin resistance, followed by Sanger

420 sequencing of the PCR-amplified *nifHDK* cluster (see **Supplementary File 1d** for a list of

421 primers). Transformants were passaged at least 3 times to ensure phenotypic stability prior to

422 storage at -80 °C in phosphate buffer containing 7% DMSO.

424 ***A. vinelandii* culturing and growth analysis**

426 *A. vinelandii* strains were grown diazotrophically in nitrogen-free Burk's medium (containing 1 µM

427 Na_2MoO_4) at 30 °C and agitated at 300 rpm. To induce genetic competency for transformation

428 experiments, Mo and Fe salts were excluded. For transformant screening, kanamycin antibiotic

429 was added to solid Burk's medium at a final concentration of 0.6 µg/mL. 50 mL seed cultures for

430 growth rate and acetylene reduction rate quantification were grown non-diazotrophically in flasks

431 with Burk's medium containing 13 mM ammonium acetate.

For growth rate quantification, seed cultures were inoculated into 100 mL nitrogen-free Burk's medium to an optical density of ~0.01 at 600 nm (OD600), after Carruthers et al. [66], and monitored for 72 h. Growth parameters were modeled using the R package Growthcurver [67].

Microbial acetylene reduction assays

A. vinelandii seed cultures representing independent biological replicates were prepared as described above and used to inoculate 100 mL of nitrogen-free Burk's medium to an OD600 ≈ 0.01. Cells were grown diazotrophically to an OD600 ≈ 0.5, at which point a rubber septum cap was affixed to the mouth of each flask. 25 mL of headspace was removed and replaced by injecting an equivalent volume of acetylene gas. The cultures were subsequently shaken at 30 °C and agitated at 300 rpm. Headspace samples were taken after 15, 30, 45, and 60 mins of incubation for ethylene quantification by a Nexis GC-2030 gas chromatograph (Shimadzu). After the 60 min incubation period, cells were pelleted at 4,700 rpm for 10 mins, washed once with 4 mL of phosphate buffer, and pelleted once more under the same conditions prior to storage at -80 °C. Total protein was quantified using the Quick Start^{TM} Bradford Protein Assay kit (Bio-Rad) according to manufacturer instructions and a CLARIOstar Plus plate reader (BMG Labtech). Acetylene reduction rates for each replicate were normalized to total protein.

Nitrogenase expression analysis

Strep-II-tagged NifD protein quantification was performed on all ancestral strains (Anc1A, Anc1B, Anc2) and DJ2102 (harboring Strep-II-tagged WT NifD). Diazotrophic *A. vinelandii* cultures (100 mL) representing three independent biological replicates were prepared as described above and harvested at an OD600 ≈ 1. Cell pellets were resuspended in TE lysis buffer (10 mM Tris, 1 mM EDTA, 1 mg/mL lysozyme) and heated at 95 °C for 10 min. Cell lysates were centrifuged at 5,000 rpm for 15 min. Total protein in the resulting supernatant was quantified using the Pierce^{TM} BCA Protein Assay kit (ThermoFisher) following manufacturer instructions. Normalized protein samples were diluted in 2 × Laemmli buffer at a 1:1 (v/v) ratio prior to SDS-PAGE analysis. Proteins were transferred to nitrocellulose membranes (ThermoFisher), stained with Revert 700 Total Protein Stain (LI-COR), and imaged on a Odyssey Fc Imager (LI-COR). Membranes were then destained with Revert Destaining Solution (LI-COR) and blocked with 5% non-fat milk in PBS solution (137 mM NaCl, 2.7 mM KCl, 10 mM Na_2HPO_4, 1.8 mM KH_2PO_4.) for 1 h at room

temperature. Membranes were rinsed once with PBS-T (PBS with 0.01% Tween-20) and
incubated with primary Strep-tag II antibody (Strep-MAB-Classic, IBA Lifesciences, Cat# 2-1507-
001, RRID: AB_513133; 1:5000 in 0.2% BSA) for 2 h at room temperature. Membranes were then
incubated in LI-COR blocking buffer containing 1:15000 IRDye 680RD Goat anti-Mouse (LI-COR)
for 2 h at room temperature and subsequently imaged with an Odyssey Fc Imager (LI-COR).
Densitometry analysis was performed with ImageJ [68], with Strep-II-tagged NifD signal intensity
normalized to that of the total protein stain.
473
474 **Nitrogenase expression, purification, and biochemical characterization**
475
Ancestral nitrogenase NifDK proteins were expressed from relevant *A. vinelandii* strains (Anc1A,
Anc1B, Anc2) and purified according to previously published methods [69] with the following
modifications: cells were grown diazotrophically in nitrogen-free Burk's medium and no
derepression step to a sufficient OD600 (~1.8) before harvesting. WT NifH was expressed in *A.*
vinelandii strain DJ884 and purified by previously published methods [70]. Protein purity was
assessed at ≥ 95% by SDS-PAGE gel with Coomassie blue staining (**Fig. 4-Figure Supplement**
1).
483
Assays were performed in 9.4 mL vials with a MgATP regeneration buffer (6.7 mM MgCl2, 30 mM
phosphocreatine, 5 mM ATP, 0.2 mg/mL creatine phosphokinase, 1.2 mg/mL BSA) and 10 mM
sodium dithionite in 100 mM MOPS buffer at pH 7.0. Reaction vials were made anaerobic and
relevant gases (N_2, C_2H_2, H_2) were added to appropriate concentrations with the headspace
balanced by argon. NifDK proteins (~240 kDa) were added to 0.42 µM, the vial vented to
atmospheric pressure, and the reaction initiated by addition of NifH (~60 kDa) protein to 8.4 µM.
Reactions were run, shaking, at 30° C for 8 minutes and stopped by the addition of 500 µL of 400
mM EDTA pH 8.0. NH_3 was quantified using a fluorescence protocol [71] with the following
modifications: an aliquot of the sample was added to a solution containing 200 mM potassium
phosphate pH 7.3, 20 mM o-phthalaldehyde, and 3.5 mM 2-mercaptoethanol and incubated for
30 minutes in the dark. Fluorescence was measured at $\lambda_{excitation}$ of 410 nm and $\lambda_{emission}$ of 472 nm
and NH_3 was quantified using a standard generated with NH_4Cl. H_2 and C_2H_4 were quantified by
gas chromatography by a thermal conductivity detector (GC-TCD) and gas chromatography with
flame ionization detector (GC-FID) respectively, according to published methods [72, 73].
498
499

Statistical analyses

Experimental data was statistically analyzed by one-way ANOVA with post-hoc Tukey HSD test.

MATERIALS AVAILABILITY

Physical materials including bacterial strains and plasmids are available to the scientific community upon request.

DATA AND CODE AVAILABILITY

Phylogenetic data, including sequence alignments and phylogenetic trees, and the script for ancestral gene codon-optimization are publicly available at https://github.com/kacarlab/garcia_nif2023. All other data are included as source data and supplementary files.

ACKNOWLEDGEMENTS

We thank Dennis Dean and Valerie Cash for providing *A. vinelandii* strains DJ, DJ2278, DJ2102, and DJ884 and for guidance in genomic manipulations; Jean-Michel Ané, April MacIntyre, and Junko Maeda for guidance and instrumentation support in performing the *in vivo* acetylene reduction assays; Bruno Cuevas for assistance with visualizing nitrogenase sequence space; and the members of the Metal Selection and Utilization Across Eons (MUSE) Consortium for helpful suggestions and discussions. This research was supported by the National Aeronautics and Space Administration (NASA) Interdisciplinary Consortium for Astrobiology Research: Metal Utilization and Selection Across Eons, MUSE (19- ICAR19_2-0007), the University of Wisconsin-Madison College of Agricultural and Life Sciences, the NASA Postdoctoral Program (A.K.G.), NASA Arizona Space Grant (B.M.C.), and the NASA Early Career Faculty Award (B.K.).

COMPETING INTERESTS

The authors declare no competing interests.

SUPPLEMENTARY FILES & LEGENDS

garcia_elife_2023_suppfile1.pdf: **Supplementary File 1**

Supplementary File 1a. Sequence characteristics of ancestral nitrogenase subunits.

Supplementary File 1b. Host taxa of nitrogenase and outgroup dark-operative
protochlorophyllide oxidoreductase homologs included for phylogenetic analysis.

Supplementary File 1c. Strains and plasmids used in this study.

Supplementary File 1d. Primers used in this study.

TABLES

n/a

Fig. 1. Engineering strategy for ancestral nitrogenase resurrection. (*A*) Experimental pipeline

for nitrogenase resurrection in *A. vinelandii* and subsequent characterization, as described in the

main text. (*B*) Structural overview of ancestral nitrogenases reconstructed in this study. Homology

models (template PDB 1M34) of Anc1B NifH and NifDK proteins are shown with ancestral

substitutions (relative to WT) highlighted in red. Select substitutions at relatively conserved sites

in proximity to FeMoco (NifD, I355V) and the NifD:NifK interface (NifD, F429Y; NifK, R108K) are

displayed in the insets. (*C*) Parallel genome engineering strategies executed in this study,

involving both ancestral replacement of only *nifD* (Anc1A and Anc2) and replacement of *nifHDK*

(Anc1B). "P*nifH*": *nifH* promoter, "KanR": kanamycin resistance cassette. *Anc1A and Anc1B were

each reconstructed from equivalent nodes of alternate phylogenies (see **Materials and**

methods).

Fig. 2. Phylogenetic and genomic context of resurrected ancestral nitrogenases. (*A*)
Maximum-likelihood phylogenetic tree from which ancestral nitrogenases were inferred. Extant
nodes are colored by microbial host taxonomic diversity. Red box highlights the clade targeted in
this study and depicted in (*B*). Tree shown was used to infer Anc1A and Anc2 sequences (an

alternate tree was used for Anc1B inference; see **Materials and methods**). (*B*) *nif* gene cluster

complexity within the targeted nitrogenase clade. Presence and absence of each *nif* gene are

indicated by grey and white colors, respectively. Because some homologs for phylogenetic

analysis were obtained from organisms lacking fully assembled genomes, the absence of

accessory *nif* genes may result from missing genomic information. (*C*) Amino acid sequence

identity matrix of nitrogenase protein subunits harbored by WT and engineered *A. vinelandii*

strains. (*D*) Extant and ancestral nitrogenase protein sequence space visualized by machine-

learning embeddings, with the resulting dimensionality reduced to two-dimensional space. UMAP

dimension axes are in arbitrary units. The field demarcated by dashed lines in the left plot is

expanded on the right plot.

589
590

Figure 2-Figure Supplement 1. Maximum-likelihood phylogenies built from nitrogenase
NifHDK homologs. Anc1A and Anc2 sequences were inferred from the left tree, and the Anc1B
sequence was inferred from the right tree (targeting an equivalent node to Anc1A). Both trees
were reconstructed from the same extant sequence dataset (see **Materials and methods** for a
description of phylogenetic reconstruction and ancestral sequence inference methods). Branch
length scale indicates amino acid substitutions per site and applies to both trees.

Fig. 3. Cellular-level characterization of ancestral nitrogenase activity and expression. (*A*) Diazotrophic growth curves of *A. vinelandii* strains measured by the optical density at 600 nm ("OD600"). A smoothed curve is shown alongside individual data points obtained from five biological replicates per strain. The non-diazotrophic DJ2278 (Δ*nifD*) strain was used as a negative control. (*B*) Mean doubling and midpoint times of *A. vinelandii* strains, calculated from data in (*A*). (*C*) *In vivo* acetylene (C_2H_2) reduction rates quantified by production of ethylene (C_2H_4). Bars represent the mean of three biological replicates per strain. (*D*) Immunodetection and protein quantification of Strep-II-tagged WT ("WT-Strep", strain DJ2102) and ancestral NifD. Top gel image shows Strep-II-tagged NifD proteins detected by anti-Strep antibody and bottom gel image shows total protein stain. Plot displays relative immunodetected NifD signal intensity normalized to total protein intensity and expressed relative to WT. Bars in plot represent the mean of three biological replicates per strain. (*B-D*) Error bars indicate ± 1 SD and asterisks indicate *p* < .01 (one-way ANOVA, post-hoc Tukey HSD) compared to WT or WT-Strep.

Fig. 4. *In vitro* analyses of ancestral nitrogenase activity profiles and mechanism. All measurements obtained from assays using purified NifDK assayed with WT NifH. (*A*) Specific activities measured for H^+, N_2, and C_2H_2 substrate. (*B*) Partial schematic of the reductive-elimination N_2-reduction mechanism of nitrogenase is shown above, centering on the N_2-binding $E_4(4H)$ state of FeMoco (see main text for discussion) [32]. (*C*) Inhibition of N_2 reduction by H_2, evidencing the mechanism illustrated in (*B*). (*D*) Catalytic efficiencies of ancestral nitrogenases,

629   described by the ratio of formed H_2 to reduced N_2 (H_2/N_2), mapped across the targeted

630   phylogenetic clade. NifD homology models (PDB 1M34 template) are displayed with ancestral

631   substitutions highlighted in red. (*A,C*) Bars represent the mean of two independent experiments

632   with individual data points shown as black circles.

REFERENCES

1. Jablonski D. Extinction: past and present. Nature. 2004;427(6975):589. doi: 10.1038/427589a.

2. Gold DA, Caron A, Fournier GP, Summons RE. Paleoproterozoic sterol biosynthesis and the rise of oxygen. Nature. 2017;543(7645):420-3. Epub 2017/03/07. doi: 10.1038/nature21412. PubMed PMID: 28264195.

3. Sanchez-Baracaldo P, Cardona T. On the origin of oxygenic photosynthesis and Cyanobacteria. New Phytol. 2020;225(4):1440-6. Epub 2019/10/11. doi: 10.1111/nph.16249. PubMed PMID: 31598981.

4. Goldford JE, Hartman H, Smith TF, Segre D. Remnants of an ancient metabolism without phosphate. Cell. 2017;168(6):1126-34. Epub 2017/03/07. doi: 10.1016/j.cell.2017.02.001. PubMed PMID: 28262353.

5. Zerkle AL, House CH, Cox RP, Canfield DE. Metal limitation of cyanobacterial N_2 fixation and implications for the Precambrian nitrogen cycle. Geobiology. 2006;4(4):285-97. doi: 10.1111/j.1472-4669.2006.00082.x.

6. Soboh B, Boyd ES, Zhao D, Peters JW, Rubio LM. Substrate specificity and evolutionary implications of a NifDK enzyme carrying NifB-co at its active site. FEBS Lett. 2010;584(8):1487-92. Epub 2010/03/12. doi: 10.1016/j.febslet.2010.02.064. PubMed PMID: 20219465.

7. Hurley SJ, Wing BA, Jasper CE, Hill NC, Cameron JC. Carbon isotope evidence for the global physiology of Proterozoic cyanobacteria. Sci Adv. 2021;7(2). doi: 10.1126/sciadv.abc8998.

8. Garcia AK, Kacar B. How to resurrect ancestral proteins as proxies for ancient biogeochemistry. Free Radic Biol Med. 2019;140:260-9. Epub 2019/04/06. doi: 10.1016/j.freeradbiomed.2019.03.033. PubMed PMID: 30951835.

9. Kacar B, Guy L, Smith E, Baross J. Resurrecting ancestral genes in bacteria to interpret ancient biosignatures. Philos Trans R Soc A. 2017;375(2109):20160352. Epub 2017/11/15. doi: 10.1098/rsta.2016.0352. PubMed PMID: 29133450; PubMed Central PMCID: PMCPMC5686408.

10. Kedzior M, Garcia AK, Li M, Taton A, Adam ZR, Young JN, et al. Resurrected Rubisco suggests uniform carbon isotope signatures over geologic time. Cell Rep. 2022;39(4):110726. Epub 2022/04/28. doi: 10.1016/j.celrep.2022.110726. PubMed PMID: 35476992.

11. Garcia AK, McShea H, Kolaczkowski B, Kacar B. Reconstructing the evolutionary history of nitrogenases: Evidence for ancestral molybdenum-cofactor utilization. Geobiology. 2020;18:394-411. Epub 2020/02/18. doi: 10.1111/gbi.12381. PubMed PMID: 32065506.

12.   Raymond J, Siefert JL, Staples CR, Blankenship RE. The natural history of nitrogen
fixation. Mol Biol Evol. 2004;21(3):541-54. Epub 2003/12/25. doi: 10.1093/molbev/msh047.
PubMed PMID: 14694078.

13.   Boyd ES, Anbar AD, Miller S, Hamilton TL, Lavin M, Peters JW. A late methanogen origin
for molybdenum-dependent nitrogenase. Geobiology. 2011;9(3):221-32. Epub 2011/04/21. doi:
10.1111/j.1472-4669.2011.00278.x. PubMed PMID: 21504537.

14.   Stueken EE, Buick R, Guy BM, Koehler MC. Isotopic evidence for biological nitrogen
fixation by molybdenum-nitrogenase from 3.2 Gyr. Nature. 2015;520(7549):666-9. Epub
2015/02/18. doi: 10.1038/nature14180. PubMed PMID: 25686600.

15.   Parsons C, Stüeken EE, Rosen CJ, Mateos K, Anderson RE. Radiation of nitrogen-
metabolizing enzymes across the tree of life tracks environmental transitions in Earth history.
Geobiology. 2020;19:18-34. doi: 10.1111/gbi.12419.

16.   Garcia AK, Cavanaugh CM, Kacar B. The curious consistency of carbon biosignatures
over billions of years of Earth-life coevolution. ISME J. 2021. Epub 2021/04/14. doi:
10.1038/s41396-021-00971-5. PubMed PMID: 33846565.

17.   Mus F, Alleman AB, Pence N, Seefeldt LC, Peters JW. Exploring the alternatives of
biological nitrogen fixation. Metallomics. 2018;10(4):523-38. Epub 2018/04/10. doi:
10.1039/c8mt00038g. PubMed PMID: 29629463.

18.   Perez-Gonzalez A, Jimenez-Vicente E, Gies-Elterlein J, Salinero-Lanzarote A, Yang ZY,
Einsle O, et al. Specificity of NifEN and VnfEN for the Assembly of Nitrogenase Active Site
Cofactors in Azotobacter vinelandii. mBio. 2021;12(4):e0156821. Epub 2021/07/21. doi:
10.1128/mBio.01568-21. PubMed PMID: 34281397; PubMed Central PMCID:
PMCPMC8406325.

19.   Buren S, Jimenez-Vicente E, Echavarri-Erasun C, Rubio LM. Biosynthesis of Nitrogenase
Cofactors. Chem Rev. 2020;120(12):4921-68. Epub 2020/01/25. doi:
10.1021/acs.chemrev.9b00489. PubMed PMID: 31975585; PubMed Central PMCID:
PMCPMC7318056.

20.   Gallon JR. Reconciling the Incompatible - N_2 Fixation and O_2. New Phytol.
1992;122(4):571-609. PubMed PMID: WOS:A1992KH15800001.

21.   Smanski MJ, Bhatia S, Zhao D, Park Y, L BAW, Giannoukos G, et al. Functional
optimization of gene clusters by combinatorial design and assembly. Nat Biotechnol.
2014;32(12):1241-9. Epub 2014/11/25. doi: 10.1038/nbt.3063. PubMed PMID: 25419741.

22.   Bennett EM, Murray JW, Isalan M. Engineering Nitrogenases for synthetic nitrogen
fixation: From pathway engineering to directed evolution. BioDesign Research. 2023. doi:
10.34133/bdr.0005.

23.   Buren S, Rubio LM. State of the art in eukaryotic nitrogenase engineering. FEMS Microbiol
Lett. 2018;365(2). Epub 2017/12/15. doi: 10.1093/femsle/fnx274. PubMed PMID: 29240940;
PubMed Central PMCID: PMCPMC5812491.

24.   Falkowski PG. Evolution of the nitrogen cycle and its influence on the biological
sequestration of CO_2 in the ocean. Nature. 1997;387(6630):272-5. doi: 10.1038/387272a0.
PubMed PMID: WOS:A1997WZ16700049.

25.   Allen JF, Thake B, Martin WF. Nitrogenase Inhibition Limited Oxygenation of Earth's
Proterozoic Atmosphere. Trends Plant Sci. 2019;24(11):1022-31. Epub 2019/08/27. doi:
10.1016/j.tplants.2019.07.007. PubMed PMID: 31447302.

26.   Vicente EJ, Dean DR. Keeping the nitrogen-fixation dream alive. Proc Natl Acad Sci USA.
2017;114(12):3009-11. Epub 2017/03/12. doi: 10.1073/pnas.1701560114. PubMed PMID:
28283657; PubMed Central PMCID: PMCPMC5373348.

27.   Boyd ES, Costas AM, Hamilton TL, Mus F, Peters JW. Evolution of molybdenum
nitrogenase during the transition from anaerobic to aerobic metabolism. J Bacteriol.
2015;197(9):1690-9. Epub 2015/03/04. doi: 10.1128/JB.02611-14. PubMed PMID: 25733617;
PubMed Central PMCID: PMCPMC4403663.

28.   Mus F, Colman DR, Peters JW, Boyd ES. Geobiological feedbacks, oxygen, and the
evolution of nitrogenase. Free Radic Biol Med. 2019. Epub 2019/02/09. doi:
10.1016/j.freeradbiomed.2019.01.050. PubMed PMID: 30735835.

29.   Boyd ES, Hamilton TL, Peters JW. An alternative path for the evolution of biological
nitrogen fixation. Front Microbiol. 2011;2:205. Epub 2011/11/09. doi: 10.3389/fmicb.2011.00205.
PubMed PMID: 22065963; PubMed Central PMCID: PMCPMC3207485.

30.   Seefeldt LC, Yang ZY, Lukoyanov DA, Harris DF, Dean DR, Raugei S, et al. Reduction of
Substrates by Nitrogenases. Chem Rev. 2020;120:5082-106. Epub 2020/03/17. doi:
10.1021/acs.chemrev.9b00556. PubMed PMID: 32176472.

31.   Stripp ST, Duffus BR, Fourmond V, Leger C, Leimkuhler S, Hirota S, et al. Second and
Outer Coordination Sphere Effects in Nitrogenase, Hydrogenase, Formate Dehydrogenase, and
CO Dehydrogenase. Chem Rev. 2022;122(14):11900-73. Epub 2022/07/19. doi:
10.1021/acs.chemrev.1c00914. PubMed PMID: 35849738.

32.   Harris DF, Lukoyanov DA, Kallas H, Trncik C, Yang ZY, Compton P, et al. Mo-, V-, and
Fe-Nitrogenases Use a Universal Eight-Electron Reductive-Elimination Mechanism To Achieve

N_2 Reduction. Biochemistry. 2019;58(30):3293-301. Epub 2019/07/10. doi: 10.1021/acs.biochem.9b00468. PubMed PMID: 31283201.

33.   Noar JD, Bruno-Barcena JM. *Azotobacter vinelandii*: the source of 100 years of discoveries and many more to come. Microbiology. 2018;164(4):421-36. Epub 2018/03/14. doi: 10.1099/mic.0.000643. PubMed PMID: 29533747.

34.   Ghebreamlak SM, Mansoorabadi SO. Divergent Members of the Nitrogenase Superfamily: Tetrapyrrole Biosynthesis and Beyond. ChemBioChem. 2020;21(12):1723-8. Epub 2020/03/18. doi: 10.1002/cbic.201900782. PubMed PMID: 32180329.

35.   Kacar B, Garmendia E, Tuncbag N, Andersson DI, Hughes D. Functional Constraints on Replacing an Essential Gene with Its Ancient and Modern Homologs. MBio. 2017;8(4):e01276-17. Epub 2017/08/31. doi: 10.1128/mBio.01276-17. PubMed PMID: 28851849; PubMed Central PMCID: PMCPMC5574714.

36.   Lyons TW, Reinhard CT, Planavsky NJ. The rise of oxygen in Earth's early ocean and atmosphere. Nature. 2014;506(7488):307-15. Epub 2014/02/21. doi: 10.1038/nature13068. PubMed PMID: 24553238.

37.   Dos Santos PC, Dean DR, Hu Y, Ribbe MW. Formation and insertion of the nitrogenase iron-molybdenum cofactor. Chem Rev. 2004;104(2):1159-73. Epub 2004/02/12. doi: 10.1021/cr020608l. PubMed PMID: 14871152.

38.   Ashkenazy H, Abadi S, Martz E, Chay O, Mayrose I, Pupko T, et al. ConSurf 2016: an improved methodology to estimate and visualize evolutionary conservation in macromolecules. Nucleic Acids Res. 2016;44(W1):W344-50. Epub 2016/05/12. doi: 10.1093/nar/gkw408. PubMed PMID: 27166375; PubMed Central PMCID: PMCPMC4987940.

39.   Smith BE, Thorneley RN, Eady RR, Mortenson LE. Nitrogenases from *Klebsiella pneumoniae* and *Clostridium pasteurianum*. Kinetic investigations of cross-reactions as a probe of the enzyme mechanism. Biochemical Journal. 1976;157(2):439-47. doi: 10.1042/bj1570439.

40.   Hardy RW, Holsten RD, Jackson EK, Burns RC. The acetylene-ethylene assay for N_2 fixation: laboratory and field evaluation. Plant Physiol. 1968;43(8):1185-207. Epub 1968/08/01. doi: 10.1104/pp.43.8.1185. PubMed PMID: 16656902; PubMed Central PMCID: PMCPMC1086994.

41.   Harris DF, Badalyan A, Seefeldt LC. Mechanistic Insights into Nitrogenase FeMo-Cofactor Catalysis through a Steady-State Kinetic Model. Biochemistry. 2022;61(19):2131-7. Epub 2022/09/15. doi: 10.1021/acs.biochem.2c00415. PubMed PMID: 36103672.

42.   Garcia AK, Kedzior M, Taton A, Li M, Young JN, Kaçar B. System-level effects of CO_2 and
RuBisCO concentration on carbon isotope fractionation. bioRxiv, submitted. 2022. doi:
10.1101/2021.04.20.440233.

43.   Som SM, Buick R, Hagadorn JW, Blake TS, Perreault JM, Harnmeijer Jelte P, et al. Earth's
air pressure 2.7 billion years ago constrained to less than half of modern levels. Nat Geosci.
2016;9(6):448-51. doi: 10.1038/ngeo2713.

44.   Catling DC, Zahnle KJ. The Archean atmosphere. Sci Adv. 2020;6(9):eaax1420. doi:
10.1126/sciadv.aax1420.

45.   Zehr JP, Jenkins BD, Short SM, Steward GF. Nitrogenase gene diversity and microbial
community structure: a cross-system comparison. Environmental Microbiology. 2003;5(7):539-
54. doi: 10.1046/j.1462-2920.2003.00451.x.

46.   Sippel D, Einsle O. The structure of vanadium nitrogenase reveals an unusual bridging
ligand. Nat Chem Biol. 2017;13(9):956-60. Epub 2017/07/12. doi: 10.1038/nchembio.2428.
PubMed PMID: 28692069; PubMed Central PMCID: PMCPMC5563456.

47.   Eady RR. Structure-function relationships of alternative nitrogenases. Chemical Reviews.
1996;96(7):3013-30. doi: 10.1021/cr950057h.

48.   Cherkasov N, Ibhadon AO, Fitzpatrick P. A review of the existing and alternative methods
for greener nitrogen fixation. Chemical Engineering and Processing: Process Intensification.
2015;90:24-33. doi: 10.1016/j.cep.2015.02.004.

49.   Dorr M, Kassbohrer J, Grunert R, Kreisel G, Brand WA, Werner RA, et al. A possible
prebiotic formation of ammonia from dinitrogen on iron sulfide surfaces. Angew Chem Int Ed Engl.
2003;42(13):1540-3. Epub 2003/04/17. doi: 10.1002/anie.200250371. PubMed PMID: 12698495.

50.   Yung YL, McElroy MB. Fixation of nitrogen in the prebiotic atmosphere. Science.
1979;203(4384):1002-4. Epub 1979/03/09. doi: 10.1126/science.203.4384.1002. PubMed PMID:
17811121.

51.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+:
architecture and applications. BMC Bioinformatics. 2009;10:421. Epub 2009/12/17. doi:
10.1186/1471-2105-10-421. PubMed PMID: 20003500; PubMed Central PMCID:
PMCPMC2803857.

52.   Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Mol Biol Evol. 2013;30(4):772-80. Epub 2013/01/19.
doi: 10.1093/molbev/mst010. PubMed PMID: 23329690; PubMed Central PMCID:
PMCPMC3603318.

53.   Capella-Gutierrez S, Silla-Martinez JM, Gabaldon T. trimAl: a tool for automated alignment
trimming in large-scale phylogenetic analyses. Bioinformatics. 2009;25(15):1972-3. Epub
2009/06/10. doi: 10.1093/bioinformatics/btp348. PubMed PMID: 19505945; PubMed Central
PMCID: PMCPMC2712344.

54.   Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies.   Bioinformatics.   2014;30(9):1312-3.   Epub   2014/01/24.   doi:
10.1093/bioinformatics/btu033.   PubMed   PMID: 24451623;   PubMed   Central   PMCID:
PMCPMC3998144.

55.   Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. ModelFinder: fast
model selection for accurate phylogenetic estimates. Nature Methods. 2017;14(6):587-9. doi:
10.1038/nmeth.4285.

56.   Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective
stochastic   algorithm   for   estimating   maximum-likelihood   phylogenies.   Mol   Biol   Evol.
2015;32(1):268-74. Epub 2014/11/06. doi: 10.1093/molbev/msu300. PubMed PMID: 25371430;
PubMed Central PMCID: PMCPMC4271533.

57.   Yang Z, Kumar S, Nei M. A new method of inference of ancestral nucleotide and amino
acid sequences. Genetics. 1995;141(4):1641-50. doi: 10.1093/genetics/141.4.1641.

58.   Yang Z. PAML 4: phylogenetic analysis by maximum likelihood. Mol Biol Evol.
2007;24(8):1586-91. Epub 2007/05/08. doi: 10.1093/molbev/msm088. PubMed PMID: 17483113.

59.   Anisimova M, Gascuel O. Approximate likelihood-ratio test for branches: A fast, accurate,
and   powerful   alternative.   Syst   Biol.   2006;55(4):539-52.   Epub   2006/06/21.   doi:
10.1080/10635150600755453. PubMed PMID: 16785212.

60.   Webb B, Sali A. Comparative Protein Structure Modeling Using MODELLER. Current
Protocols in Bioinformatics. 2016;54(1). doi: 10.1002/cpbi.3.

61.   Pettersen EF, Goddard TD, Huang CC, Meng EC, Couch GS, Croll TI, et al. UCSF
ChimeraX: Structure visualization for researchers, educators, and developers. Protein Sci.
2021;30(1):70-82. Epub 2020/09/04. doi: 10.1002/pro.3943. PubMed PMID: 32881101; PubMed
Central PMCID: PMCPMC7737788.

62.   Lin Z, Akin H, Rao R, Hie B, Zhu Z, Lu W, et al. Language models of protein sequences
at the scale of evolution enable accurate structure prediction. bioRxiv. 2022. doi:
10.1101/2022.07.20.500902.

63.   Rives A, Meier J, Sercu T, Goyal S, Lin Z, Liu J, et al. Biological structure and function
emerge from scaling unsupervised learning to 250 million protein sequences. Proc Natl Acad Sci
USA. 2021;118(15). doi: 10.1073/pnas.2016239118.

64. McInnes L, Healy J, Melville J. UMAP: Uniform Manifold Approximation and Projection for
Dimension Reduction. arXiv. 2020. doi: 10.48550/arXiv.1802.03426.

65. Dos Santos PC. Genomic Manipulations of the Diazotroph *Azotobacter vinelandii*.
Methods Mol Biol. 2019;1876:91-109. Epub 2018/10/15. doi: 10.1007/978-1-4939-8864-8_6.
PubMed PMID: 30317476.

66. Carruthers BM, Garcia AK, Rivier A, Kacar B. Automated Laboratory Growth Assessment
and Maintenance of *Azotobacter vinelandii*. Curr Protoc Microbiol. 2021;1(3):e57. Epub
2021/03/04. doi: 10.1002/cpz1.57. PubMed PMID: 33656286.

67. Sprouffske K, Wagner A. Growthcurver: an R package for obtaining interpretable metrics
from microbial growth curves. BMC Bioinformatics. 2016;17:172. Epub 2016/04/21. doi:
10.1186/s12859-016-1016-7. PubMed PMID: 27094401; PubMed Central PMCID:
PMCPMC4837600.

68. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image
analysis. Nat Methods. 2012;9(7):671-5. Epub 2012/08/30. doi: 10.1038/nmeth.2089. PubMed
PMID: 22930834; PubMed Central PMCID: PMCPMC5554542.

69. Jimenez-Vicente E, Martin Del Campo JS, Yang ZY, Cash VL, Dean DR, Seefeldt LC.
Application of affinity purification methods for analysis of the nitrogenase system from *Azotobacter
vinelandii*. Methods Enzymol. 2018;613:231-55. Epub 2018/12/05. doi:
10.1016/bs.mie.2018.10.007. PubMed PMID: 30509468.

70. Christiansen J, Goodwin PJ, Lanzilotta WN, Seefeldt LC, Dean DR. Catalytic and
biophysical properties of a nitrogenase Apo-MoFe protein produced by a *nifB*-deletion mutant of
Azotobacter vinelandii. Biochemistry. 1998;37(36):12611-23. Epub 1998/09/09. doi:
10.1021/bi981165b. PubMed PMID: 9730834.

71. Corbin JL. Liquid Chromatographic-Fluorescence Determination of Ammonia from
Nitrogenase Reactions: A 2-Min Assay. Appl Environ Microbiol. 1984;47(5):1027-30. doi:
10.1128/aem.47.5.1027-1030.1984.

72. Khadka N, Dean DR, Smith D, Hoffman BM, Raugei S, Seefeldt LC. CO_2 Reduction
Catalyzed by Nitrogenase: Pathways to Formate, Carbon Monoxide, and Methane. Inorg Chem.
2016;55(17):8321-30. Epub 2016/08/09. doi: 10.1021/acs.inorgchem.6b00388. PubMed PMID:
27500789; PubMed Central PMCID: PMCPMC5068488.

73. Yang ZY, Dean DR, Seefeldt LC. Molybdenum nitrogenase catalyzes the reduction and
coupling of CO to form hydrocarbons. J Biol Chem. 2011;286(22):19417-21. Epub 2011/04/02.
doi: 10.1074/jbc.M111.229344. PubMed PMID: 21454640; PubMed Central PMCID:
PMCPMC3103320.