Short-term monocular deprivation boosts neural responsiveness to audio-visual events for the undeprived eye

Federici A.[1], Bernardi G.[1], Senna I.[2], Fantoni M.[1], Ernst M.O.[2], Ricciardi E.[1], Bottari D.[1]

[1] MoMiLab, IMT School for Advanced Studies Lucca, 55100 Lucca, Italy

[2] Applied Cognitive Psychology, Ulm University, 89081 Ulm, Germany

Corresponding author: Alessandra Federici alessandra.federici@imtlucca.it

Abstract

A brief period of monocular deprivation (MD) induces short-term plasticity of the adult visual system. Whether MD elicits changes beyond visual processing is yet unclear. Here, we assessed the specific impact of MD on multisensory processes. Neural oscillations associated with visual and audio-visual processing were measured for both the deprived and the undeprived eye. Results revealed that MD changed neural activities associated with unimodal and multisensory processes in an eye-specific manner. Selectively for the deprived eye, alpha activity was reduced within the first 150 ms of visual processing. Conversely, gamma activity was enhanced in response to audio-visual events only for the undeprived eye within 100-300 ms after stimulus onset. The analysis of gamma responses to unimodal auditory events revealed that MD elicited a crossmodal upweighting for the undeprived eye. Distributed source modeling suggested that the right parietal cortex played a major role in all neural effects induced by MD. Finally, visual and audio-visual processing alterations emerged selectively for the induced (but not the evoked) component of the neural oscillations, indicating a major role of feedback connectivity. These findings support a model in which MD increases excitability to visual events for the deprived eye and to audio-visual and auditory events for the undeprived eye. On the one hand, these results reveal the causal impact of MD on both unisensory and multisensory processes but with distinct frequency-specific profiles. On the other hand, they highlight the feedback nature of short-term neural plasticity. Overall this study shed light on the high flexibility and interdependence of unimodal and multisensory functions.

Keywords: neural plasticity, multisensory, monocular deprivation, EEG, neural oscillations

Highlights:

- We unveiled the impact of temporary MD on visual and audio-visual processing
- MD enhanced visual excitability for the deprived eye
- MD boosted neural responses to audio-visual events for the undeprived eye
- Analyses of auditory processing revealed crossmodal effects following MD
- Short-term MD primarily affects induced, non-phase-locked, oscillatory activity

Introduction

While neuroplasticity of the visual system is maximal during development (Hubel and Wiesel, 1970; Hensch, 2005; Espinosa and Stryker, 2012; Levelt and ubener, 2012; Reh et al., 2020), evidence that a certain degree of plasticity is retained in the adult visual cortex has accumulated (Karmarkar and Dan, 2006; Spolidoro et al., 2009; Espinosa and Stryker, 2012; Hensch and Quinlan, 2018). Studies in adults employing psychophysics and neuroimaging methods showed that a brief period of MD, a pivotal model for testing the plasticity of the visual system, alters the ocular balance by strengthening visual processing of the deprived eye and weakening the undeprived eye (Castaldi et al., 2020). These functional changes reflect ocular dominance shifts in V1 in favor of the deprived eye (Lunghi et al., 2015a; Binda et al., 2018) and are supposedly driven by homeostatic plasticity (Lunghi et al., 2015b), the mechanism involved in preserving the cortical excitatory-inhibitory balance (Turrigiano, 2012). Besides this well-documented MD effect on visual processing, little is known about how MD affects multimodal processing.

Alterations induced by MD seem to impact beyond visual processing. At the behavioral level, short-term MD was found to affect the interplay between sensory modalities (Lo Verde et al., 2017; Opoku-Baah and Wallace, 2020). Overall, results were consistent with a decreased multisensory interplay for the deprived eye, in which the visual processing is generally strengthened following MD, and vice versa, an increased multisensory interplay for the undeprived eye, in which visual processing is typically weakened following MD. However, the neural correlates of MD effects on multisensory processes are still unknown. To fill this gap, we exploited the model of MD to elicit short-term plasticity and investigated specific changes in visual and audio-visual processes for the deprived and undeprived eye. Overall, this study aimed to address whether a brief period of monocular visual experience in adulthood can affect the neural processing associated with multisensory audio-visual events, and which are the underpinning neural mechanisms.

Here, we combined psychophysical and electrophysiological approaches. We employed the *sound-induced flash illusion* (Shams et al., 2000, Hirst et al., 2020), in which the number of perceived flashes can be biased by the number of concurring beeps. To investigate the impact of MD on visual processing, we measured neural oscillations in response to a single flash before and after a deprivation phase. To determine the existence of an audio-visual MD effect, the analysis focused on neural oscillations elicited by the *fission* illusion trials in which a single flash is coupled with two beeps, and two flashes can be perceived (Shams et al., 2000). This illusion reveals the impact of auditory input on visual processing in case of conflictual audio-visual information.

Neural correlates associated with visual and audio-visual processing are characterized by their spectral, temporal, and spatial properties. Reflecting high and low neuronal excitability cycles, neural oscillations reveal perceptual organization in both visual (Jensen et al., 2014; VanRullen, 2016) and multisensory analyses (Cooke et al., 2019; Lennert et al., 2021). We expected MD to induce changes in alpha activity, which is known to reflect levels of inhibition (Klimesch et al., 2007; Klimesch, 2012), and, thus, increase the visual system excitability for the deprived eye (Lunghi et al., 2015a). Moreover, we hypothesized MD to alter responses in the gamma range. Gamma activity has been as well associated with excitatory-inhibitory balance (Jensen et al., 2010; 2012), and its modulation was found to be involved in eliciting a perceptual fission illusion (Bhattacharya et al., 2002; Mishra et al., 2007; Lange et al., 2011, 2013; Balz et al., 2016).

Importantly, distinct components of neural oscillations characterize different types of processing according to the direction of information flow: while the evoked activity, which is phase-locked to the stimulation, has mainly been associated with feedforward processing (thalamo-cortical), the induced oscillatory activity, which is not phase-locked to the onset of the stimulus, has mainly been associated with feedback processing (cortico-cortical connectivity, Klimesch et al., 1998; Tallon-Baudry and Bertrand, 1999; Chen et al., 2012). Studies in humans and non-human animal models demonstrated that permanent sensory deprivation primarily affects induced oscillatory activity (Bottari et al., 2016; Yusuf et al., 2017; Bednaya et al., 2021). As homeostatic plasticity is an intrinsic feedback mechanism (Turrigiano and Nelson, 2004), we predicted to primarily observe changes in induced oscillatory activity following temporary MD for both visual and audio-visual processing. Finally, source modeling was employed to investigate the neural sources of short-term plasticity effects.

Materials and Methods

Participants

Since the effect of MD on multisensory processing was unknown, we estimated the minimum sample size needed to reach the expected effect of MD on visual processing as previously reported in the literature. We expected the MD effect on visual processing to be at occipito-parietal electrodes, in the alpha range [8-14 Hz] (Lunghi et al., 2015a), and within the first stages of visual processing [0-120 ms] (comprising the earliest visual evoked potential, C1 wave, known to be modulated by MD; Lunghi et al., 2015a). The power analysis was performed by simulating our planned analysis (t1-t0 Deprived eye vs. t1-t0 Undeprived eye) on the alpha frequency power using a cluster-based permutation test (Wang and Zhang, 2021). The analysis revealed an estimated

minimum sample size of 17 participants (for further details on sample size estimation see Supplementary Materials and Figure S1). Note that previous studies investigating the effect of MD using EEG analyzed up to 16 participants (Lunghi et al., 2015a; Zhou et al., 2015; Schwenk et al., 2020).

To determine individual suitability for the *main experiment* (EEG experiment), twenty-seven potential participants completed a *preliminary behavioral assessment* (see the section below) in order to assess whether they met the following inclusion criteria: (i) to perceive the *fission illusion* with the dominant eye (i.e., >20% illusory rate), and (ii) to not be completely biased by the sound in the illusory conditions (i.e., >95% illusory rate). Out of twenty-seven young adults tested (mean age 28.22 ± 2.41 SD, twelve males and fifteen females), six participants were excluded as they did not meet these inclusion criteria or could not comply with the experimental instructions (see Supplementary Materials). Out of the 21 participants who performed the *main experiment*, one further participant was excluded due to his poor behavioral performance (the number of errors was 3 SD above the group mean in the conditions in which only auditory stimuli were presented, i.e., the control conditions). The final sample included 20 young-adult participants (mean age 28.45 ± 2.67 SD, eight males and twelve females). They all had normal or corrected-to-normal vision (visual acuity ≥ 8/10) and did not report hearing deficits or a history of neurological conditions. Since one EEG and one behavioral dataset, from different participants, went lost due to technical issues during acquisitions, the analyzed data sample included 19 behavioral and 19 EEG datasets.

The study was approved by the local Ethical Committee (Comitato Etico di Area Vasta Nord Ovest Regione Toscana protocol n. 24579). Each participant signed a written informed consent before taking part in the experiment. The experimental protocol adhered to the principles of the Declaration of Helsinki (2013).

Stimuli and apparatus

The experiment was performed in a dimly lit and sound-attenuated chamber (BOXY, B-Beng s.r.l., Italy). Participants were comfortably sitting in front of the apparatus, with their eyes at a distance of 60 cm from the monitor. Visual stimuli were presented on an LCD monitor (60 Hz refresh rate; 24.5 inches; 1920 × 1080 screen resolution), and audio stimuli were delivered via a single speaker (Bose® Companion 2, series III multimedia) located below the screen and aligned with its center. Stimuli were flashes and beeps. Both visual and audio stimuli were created using Matlab (The Mathworks, Inc. - version 2017b). The audio stimulus was a 7 ms quadratic beep with a 3.5 kHz frequency and a sampling rate of 44.1 kHz, which was presented at about 75 dB. The visual stimulus was a 2° diameter grey dot displayed 5° below the center of the screen for 17 ms

(corresponding to 1 frame) on a black background. The contrast level of the grey dot was selected

individually via a staircase procedure to elicit the fission illusion in about 50% of trials (Pérez-

Bellido et al., 2015, see below *Preliminary behavioral assessment*). Stimuli were delivered using E-

Prime® software (version 2, Psychology Software Tools, Inc. www.pstnet.com). The Audio/Visual

(AV) Device (Electrical Geodesics, Inc.) was employed to ensure accurate optimal synchronization

between the presented stimuli and the recorded EEG traces.

Experimental Design

The whole procedure consisted of two main parts performed on two separate days: a *preliminary*

behavioral assessment and the *main experiment*, in which the dominant eye was deprived of

patterned visual input using a translucent eye patch. In both the *preliminary behavioral assessment*

and the *main experiment,* participants performed a *monocular visual discrimination task*.

Monocular visual discrimination task

Participants were asked to report the number of perceived flashes (0, 1, or 2) while task-irrelevant

beeps (0, 1, or 2) were presented. Responses were given by pressing one of three keypad buttons

using the right-hand fingers. In each trial, audio (A) and visual (V) stimuli could be presented

coupled or isolated constituting eight conditions: half the conditions were unisensory and comprised

single or couples of visual or auditory events (visual: V and VV; auditory: A and AA) and the other

half were multisensory (coherent audio-visual stimulation: AV and AVAV; illusory audio-visual

stimulation: AVA and VAV, for a schematic summary of all conditions see the table reported in

Figure 1a). Unisensory auditory trials (i.e., A and AA) represented control conditions. They were

employed to ensure that participants correctly performed the task (participants with a number of

errors 3 SD above the group mean were excluded from the analyses as described in the Participants

section). The presentation order of the conditions was randomized.

Since our main aim was to explore changes in the neural response to audio-visual events caused by

a short-term MD, the main analysis focused on the audio-visual condition inducing fission illusion

(AVA). Generating an unstable percept, this illusion allows investigating subtle changes in audio-

visual processing. As a control, we investigated changes in the unisensory visual condition (V).

Notably, the visual stimulus was the same single flash in AVA and V conditions.

All trials started with a grey fixation cross presented at the center of the screen on a black

background. After 717 ms, the stimulation was delivered. In the case of the fission illusion (i.e.,

AVA), each visual and auditory stimuli were separated by one frame (i.e., 17 ms; see an example

for AVA condition in Figure 1a and Supplementary Materials). Following the stimulation, a blank

screen appeared for 500 ms (response-free time window), then the fixation cross became white, and participants were asked to respond within 1 second (except for the staircase procedure, in which participants had infinite time to respond; see *Preliminary behavioral assessment*). As soon as the response was given, a blank screen was presented for 300 ms before the beginning of the subsequent trial. Participants were asked to maintain their gaze at the fixation cross throughout the duration of the trial.

Preliminary behavioral assessment

The ocular dominance via the Porta Test (see Supplementary Materials), the visual acuity via the eye chart, and the rate of the illusory percept during the *monocular visual discrimination task* were measured to verify if the participant could take part in the *main experiment*.

Once ocular dominance and visual acuity were tested, participants performed two short versions of the *monocular visual discrimination task*. The first short version comprised a staircase procedure (see Supplementary Materials for details) that was used to identify, at the individual level, the luminance contrast between the grey dot and the black background needed to elicit the fission illusion in about 50% of trials (Pérez-Bellido et al., 2015). Participants performed this test monocularly, with the dominant and non-dominant eye (the order was randomized). The contrast level for the dominant and non-dominant eye did not differ within-participant (t(26)= -1.466, p=0.155). The second short version of the *monocular visual discrimination task* (30 trials for each of the following conditions V, VV, AVA, VAV, A, and 6 for AA, AV, AVAV) was then performed with the dominant eye (with the contrast identified by the staircase procedure) to evaluate whether the participant was fulfilling the inclusion criteria to take part in the *main experiment* (see Participants section).

Previous evidence revealed that individual sensory preference (*Audio* or *Visual*) impacts on multisensory processing (Giard and Perronet, 1999). To assess whether individual sensory predisposition might affect multisensory short-term plasticity, participants monocularly performed a speeded object recognition task based on auditory, visual, and audio-visual information (see Supplementary Materials for details). Each participant's Sensory-Preference (*Audio* or *Visual*) was classified for each eye, and no significant difference was found between them (p>0.68). Sensory-Preference was an additional measure that we took into consideration: for this reason, the N within the *Audio* and *Visual* groups were not balanced.

Main experiment

Participants recruited in the *main experiment* came to the laboratory a second time on a different

day. Since the *main experiment* lasted about five hours, the data were always acquired in the

morning (approximately between 9 am and 2 pm) to avoid possible confounds associated with the

circadian cycle or related to (visual) activities performed before the experiment. Each participant

repeated the short version of the *monocular visual discrimination task* with the dominant eye

comprising the staircase procedure to ensure the test-retest reliability of the selected visual stimulus

contrast (no significant difference was found between the contrast levels measured in the two

assessments; $t(20)=0.637$, $p=0.532$).

A brief practice of 16 trials was run before the *main experiment*. Participants performed the

monocular visual discrimination task, with each eye, before (t0) and after (t1) a period of

monocular deprivation (MD) (see Figure 1b), while their EEG signal was recorded. Thus, each

participant performed a total of four sessions of the *monocular visual discrimination task* (i.e., at t0

and t1, both with the dominant and the non-dominant eye). Note that whether they started with the

dominant or the non-dominant eye was counterbalanced across participants. MD consisted of 150

minutes in which the dominant eye was occluded by a translucent eye patch, following a validated

procedure (Lunghi et al., 2011). From now on, we will refer to the dominant eye as the 'Deprived

eye' and to the non-dominant eye as the 'Undeprived eye'.

To prompt active and controlled multisensory interactions with the environment during the MD

phase, participants were engaged in predefined activities. Since the physical activity was

demonstrated to boost short-term homeostatic plasticity in the adult visual cortex (Lunghi and Sale,

2015), all participants were engaged in the following activities: table football, table hockey, ping

pong, and billiards, each of them lasting 15 minutes. The EEG cap was always kept on the scalp for

the entire experiment duration.

Each monocular session (i.e., t0 Deprived, t0 Undeprived, t1 Deprived, and t1 Undeprived)

comprised 100 trials for the conditions V, VV, AVA, VAV, A, and only 30 trials for the conditions

AA, AV, AVAV, and was divided into five blocks (118 trials each) lasting about 5 minutes each.

The number of trials was chosen to keep the duration of the monocular session within the estimated

length of the MD effect (which has been demonstrated to be present for up to 90 minutes, but

substantially decrease after 15 minutes; see Lunghi et al., 2011).

Figure 1. Schematic illustration of the experimental paradigm. (a) The top left table shows all the possible combinations of the stimuli (visual stimuli V and auditory stimuli A), representing all the eight possible conditions (conditions of interest are highlighted, and the number of trials in each condition is shown in brackets). On the right, an example of a single trial presentation in AVA condition is depicted. (b) Experimental procedure showing the four sessions performed with the Deprived eye (upper line, red contour) and the Undeprived eye (bottom line, blue contour). At t0 and t1, participants performed the visual monocular task with each eye (eye order was counterbalanced across participants). Between t0 and t1, participants wore a translucent eye patch on the dominant eye (Deprived eye) for 150 minutes (MD phase).

EEG recording and preprocessing

EEG data were collected continuously during the four monocular task sessions (i.e., t0 Deprived, t0 Undeprived, t1 Deprived, and t1 Undeprived), using Electrical Geodesics EEG system with 64-channels (EGI; 500 Hz sampling rate). Offline, the data of the four sessions were concatenated at the individual level to detect common stereotypical artifacts. Data were preprocessed by implementing a validated approach (Stropahl et al., 2018; Bottari et al., 2020). The continuous recordings were filtered (low-pass cut-off at 40 Hz, Hanning filter, order 500; high-pass cut-off at 1 Hz, Hanning filter, order 100) and downsampled to 250 Hz to reduce the computational time. The filtered and downsampled data were segmented into consecutive 1-second epochs and cleaned using joint probability criterion: segments displaying an activity with a joint probability across all channels larger than 3 SD were removed (pop_jointprob function of EEGLAB; Delorme et al., 2007). Independent Component Analysis (ICA) based on the extended Infomax (Bell and Sejnowski, 1995; Jung et al., 2000a, 2000b) was then performed. The resulting ICA weights were saved and applied to the raw continuous unfiltered data (Stropahl et al., 2018; Bottari et al., 2020). Components associated with stereotypical artifacts, such as eye blinks and eye movements, were identified and removed using a semiautomatic procedure (CORRMAP, Viola et al., 2009). The data were then low-pass and high-pass filtered (100 Hz, filter order 100; 0.1 Hz, filter order 500) with a Hanning filter. Noisy channels were identified based on visual inspection and then interpolated using spherical spline interpolation (mean interpolated electrodes per subject 2.32 ± 2.26 SD) and

re-referenced to the average. Finally, the residual power line fluctuations at 50 Hz were removed using the CleanLine EEGLAB plugin (https://github.com/sccn/cleanline). The EEG data were then split again into the original four sessions. Each session was then segmented into epochs of 2.2 seconds, from -1 to 1.2 seconds with respect to the onset of the stimulation. Noisy epochs were then rejected based on the joint probability across channels (Delorme et al., 2007) with a threshold of 3 SD (mean epochs rejected per subject in each session for V condition: t0 Deprived 12% ± 4 SD, t0 Undeprived 12% ± 6 SD, t1 Deprived 14% ± 5 SD, and t1 Undeprived 12% ± 5 SD, and for AVA condition: t0 Deprived 12% ± 6 SD, t0 Undeprived 12% ± 5 SD, t1 Deprived 12% ± 5 SD, and t1 Undeprived 14% ± 6 SD). All these steps were performed with EEGLAB software (Delorme and Makeig, 2004). Data were then imported into Fieldtrip (Oostenveld et al., 2011) to perform time-frequency decomposition and statistical analyses.

Time-frequency decomposition

Time-frequency decomposition of the EEG data was performed within each session and separately for the visual and audio-visual conditions. Within each condition and session, we first extracted the induced power at a single trial level after subtracting the evoked activity (that is, subtracting from each trial the ERP computed averaging across trials without low-pass filtering). Time-frequency decomposition of single-trials was computed at each channel, separately for low (2-30 Hz) and high (30-80 Hz) frequency ranges. The oscillations in low frequencies were estimated using a Hanning taper with a frequency-dependent window length (4 cycles per time window) in steps of 2 Hz. Oscillations with higher frequencies were estimated using a Multitapers method with Slepian sequence as tapers, in steps of 5 Hz with a fixed-length time window of 0.2 seconds and fixed spectral smoothing of ± 10 Hz. For both frequency ranges, the power was extracted over the entire epoch (from -1 to 1.2 seconds) in steps of 0.02 seconds. Then, the average across trials was computed at the individual level within each session (t0 Deprived, t0 Undeprived, t1 Deprived, t1 Undeprived), conditions (V and AVA), and frequency range (low and high). The resulting oscillatory activity was baseline-corrected to obtain the relative signal change with respect to the baseline interval. For the low-frequency range, the baseline was set between -0.7 and -0.3 seconds, while for the high-frequency range, it was between -0.2 and -0.1 seconds. The low-frequency range, having longer cycles, required a wide baseline for the appropriate estimation of slow oscillations and was kept temporally distant from the stimulus onset to avoid temporal leakage of post-stimulus activity into the baseline period. The same procedure, without ERP subtraction from single trials, was implemented to estimate the total power. The baseline-corrected evoked power was computed by subtracting the baseline-corrected induced power from the baseline-corrected total power.

Source reconstruction

To better characterize the neural alterations induced by MD, source estimation of the neural effects was performed using Brainstorm software (Tadel et al., 2011) on preprocessed EEG data. Sources were extracted by applying a dynamic statistical parametric mapping (dSPM; Dale et al., 2000), adopting minimum-norm inverse maps to estimate the locations of scalp electrical activities. For each dataset, we used single trials pre-stimulus baseline [– 0.1 to 0.002 s] to calculate single subject noise covariance matrices and to estimate individual noise standard deviations at each location (Hansen et al., 2010). The boundary element method (BEM) provided in OpenMEEG was adopted as a head model; the model was computed on the first dataset and then applied to all the others (default parameters in Brainstorm were selected). Source estimation was performed by selecting the option of constrained dipole orientations (Tadel et al., 2011). Time-frequency decomposition was computed for each participant on the estimated sources at the single-trial level using the same approach described for the time-frequency decomposition performed at the sensor level.

Statistical analysis

Behavioral data

For each participant, we computed the d-prime (d') as visual and audio-visual sensitivity indices: d' = $z(p$ hits) - $z(p$ false alarms), where z is the inverse cumulative normal function, and p is the proportion of hits and of false alarms out of signal and noise, respectively. Values equal to 0 or 1 were corrected as 1/n and (n-1)/n, respectively, with n being the number of signal or noise trials. To compute the visual d', we defined as hits, trials in which participants perceived one flash (V condition) and correctly responded 'one'. Consequently, false alarms were trials in which two flashes were presented (VV), and participants responded to having seen one flash. To compute the audio-visual d', we defined, coherently with the fission illusion literature, false alarms AVA trials in which participants reported two flashes (Watkins et al., 2006; Whittingham et al., 2014; Pérez-Bellido et al., 2015; Vanes et al., 2016; Keil, 2020). Thus, AVAV trials in which participants correctly responded 'two flashes' were considered hits. Note that the audio-visual sensitivity (d') is inversely related to the amount of fission illusion: smaller d' indicated a greater amount of illusory percepts and vice-versa (note that Response means of each condition are reported in Supplementary Materials Figure S2).

Two mixed-design ANOVA, with Eye (Deprived and Undeprived) and Time (t1 and t0) as within-subjects factors, and Sensory-Preference (*Audio* and *Visual*) in the Deprived eye and Sensory-

Preference (*Audio* and *Visual*) in the Undeprived eye as between-subjects factors, were performed

separately on visual and audio-visual d' values. The Sensory-preference between-subjects factors

were inserted in order to control their impact on the visual and audio-visual perception.

Neural oscillations

Oscillatory activity occurring after stimulus onset was analyzed separately for visual (V) and audio-

visual (AVA) trials to assess MD impact on visual and audio-visual processing. Time-frequency

analyses were separately performed for induced and evoked oscillatory activity for both low [4-30

Hz] and high [30-80 Hz] frequency ranges.

To assess the impact of MD, we subtracted the oscillatory activity recorded before MD from the

oscillatory activity recorded after MD (i.e., t1 minus t0). This difference was computed separately

for the Deprived and Undeprived eye. From now on, *PowChangeDeprived* represents relative

changes in power due to MD for the Deprived eye, and *PowChangeUndeprived* the relative changes

in power due to MD for the Undeprived eye.

To compare the impact of MD on Deprived and Undeprived eye, a series of non-parametric cluster-

based permutation tests were performed via a paired-sample t-test without a priori assumptions (i.e.,

across all electrodes, time-points, and frequencies) between *PowChangeDeprived* and

PowChangeUndeprived. We used the Monte Carlo method with 1000 random permutations;

cluster-level statistics were calculated taking the sum of the t-values within every cluster, with an

alpha level of 0.05 (two-tailed) and a minimum neighbor channel = 1. Identified clusters were

considered significant at $p<0.025$ (corresponding to a critical alpha level of 0.05 in a two-tailed

test). We focused on the post-stimulus activity, and thus, statistical tests were performed for the

entire response-free time window, that is, from 0 to 0.5 seconds. The time period after 0.5 seconds

from the stimulation onset was discarded to prevent including motor artifacts. If a significant

difference due to MD emerged in this test (*PowChangeDeprived* vs. *PowChangeUndeprived*), we

assessed whether differences between the two eyes emerged at t0 or t1. To this end, two planned

comparisons (i.e., t0 Deprived vs. t0 Undeprived and t1 Deprived vs. t1 Undeprived) were

performed using the same cluster-based permutation analysis approach and the same parameters

reported above. In case a significant difference between Deprived and Undeprived eye would

emerge only at t1 and not at t0, it would be indicative of a specific effect of the MD manipulation

and rule out possible differences between the two eyes at baseline.

Correlations between neural and behavioral changes

After assessing the normality of the data with Shapiro-Wilk tests, Pearson correlations were
employed to assess whether the neural changes related to short-term MD were associated with
behavioral changes. When multiple correlations were performed to the same dataset, their results
were compared employing a bootstrap method adapted for independent samples (see
https://github.com/GRousselet/blog/tree/master/comp2dcorr).

The datasets and code used in the present study are available from the corresponding author on
reasonable request.

Results
Behavioral data

Unisensory visual
The mixed-design ANOVA on visual d' with Eye (Deprived and Undeprived) and Time (t1 and t0)
as within-subjects factors and Sensory-preference in the Deprived or in the Undeprived eye as
between-subjects factor revealed a significant main effect of Time ($F_{(1,15)}=13.52$, $p=0.002$). No
main effects of Eye, Sensory-Preference in the Deprived or in the Undeprived eye, nor other
interaction effects were found (all ps>0.1). These findings suggest a general decrease of sensitivity
at t1 (mean d' ± SE for each session: t0 Deprived 1.44 ± 0.21; t1 Deprived 1.13 ± 0.22; t0
Undeprived 1.37 ± 0.26; t1 Undeprived 1.14 ± 0.23; see Figure 2a).

Audio-visual
The mixed-design ANOVA performed on audio-visual d' revealed a significant main effect of Eye
($F_{(1,15)}=7.53$, $p=0.015$), showing that the Undeprived eye was less susceptible to the fission
illusion (mean d' ± SE for each session: t0 Deprived 0.58 ± 0.16; t1 Deprived 0.53 ± 0.19; t0
Undeprived 0.84 ± 0.23; t1 Undeprived 0.66 ± 0.17). A tendency towards significance emerged for
the interaction between Eye, Sensory-Preference in the Deprived eye, and Sensory-Preference in the
Undeprived eye ($F_{(1,15)}=4.01$; $p=0.064$). This tendency might suggest that the participant's
Sensory-Preference had an eye-specific impact; being an *Audio*-subject or a *Visual*-subject might
affect the level of the fission illusion perceived with that eye (in Deprived eye mean d' ± SE A-
group: 0.52 ms ± 0.19; V-group: 0.69 ms ± 0.20; in the Undeprived eye A-group: 0.65 ms ± 0.20;
V-group: 0.96 ms ± 0.36). No significant main effects of Time, Sensory-Preference in the Deprived
or in the Undeprived eye, nor other interactions emerged (all ps>0.08). Notably, a strong fission
illusion was elicited in both eyes as highlighted by the small d' measured in each of the four
sessions (see Figure 2b).

Since no main effect of the Sensory-Preference emerged neither in visual nor in audio-visual

ANOVAs, the analyses of EEG activity were performed on the whole group.

Figure 2. Behavioral performance computed as sensitivity (d'). (a) Both in visual and (b) audio-visual conditions,

the group mean d' is reported for t0 and t1 within each eye (Deprived and Undeprived); superimposed dots represent

single subjects' data; significant differences are highlighted with *. Sensitivity in the visual condition decreased after

MD in both Deprived and Undeprived eye. Fission illusion in the audio-visual condition was greater for the Deprived

than Undeprived eye.

Neural oscillations

To specifically investigate whether MD primarily affected feedback and/or feedforward

connectivity, we assessed the impact of MD on induced and evoked neural oscillations associated

with the processing of visual and audio-visual stimuli.

Unisensory Visual

Induced power. The cluster-based permutation test performed on induced oscillatory activity within

the low-frequency range [4-30 Hz] revealed a significant difference for *PowChangeDeprived* vs.

PowChangeUndeprived (p<0.009) spanning from occipital to frontal regions. MD elicited a marked

decrease of induced activity in the alpha range [10-16 Hz] between 0 to 0.12 seconds selectively for

the Deprived eye (see Figure 3a, b). While no significant effects were found when comparing the

oscillatory activity between the two eyes at t0 (all ps>0.51), the comparison performed on induced

oscillatory activity measured at t1 revealed a significant difference (p<0.002) between Deprived and

Undeprived eye for the power in the alpha range [10-16 Hz]. These planned comparisons confirmed

a direct effect of MD, and excluded possible confounds due to differences between the eyes at

baseline. To further investigate the time-course of the MD effect in the alpha range, we tested the

difference between t1 and t0 for each eye (i.e., Deprived, Undeprived). For each session and
participant, we extracted the mean induced alpha power [10-16 Hz] measured across three occipital
electrodes (E36, E38, E40, which corresponded to the peak of the statistical effect in the
PowChangeDeprived vs. *PowChangeUndeprived* cluster-based permutation test). A series of paired
t-tests were performed between t0 and t1 for each eye, at each time-point within the whole-time
window of interest [0-0.5 s] (FDR corrected, q=0.05). A significant difference was found only for
the Deprived eye, showing a clear decrease in the alpha synchronization after MD (from 0 to 0.14 s;
for Undeprived eye all ps>0.97; see Figure 3c).

No significant differences between *PowChangeDeprived* and *PowChangeUndeprived* were found
in the high-frequency range [30-80 Hz] (all ps>0.67, see Figure S3).

Evoked power. Cluster-based permutation analyses on evoked oscillatory activity were performed
contrasting *PowChangeDeprived* vs. *PowChangeUndeprived* to test whether MD alters feedforward
visual processing. No significant differences emerged in either low or high frequencies (all *p*s>0.13,
see Figure S4).

In sum, a visual MD effect emerged selectively for the Deprived eye and in the induced oscillatory
activity within the alpha range.

Source analysis
We investigated the electrical sources of the visual MD effect. To this end, a permutation paired t-
test (1000 randomizations) was performed at the source level, on the power in the alpha range [10-
16 Hz] between t0 and t1 for the Deprived eye (time window [0-0.5 s]; FDR correction was applied
on the time dimension). Results revealed that the visual MD effect was mainly located within the
right hemisphere and comprised the superior parietal gyrus, the superior occipital gyrus,
intraparietal and subparietal sulcus, and extended to calcarine sulcus (corrected p-threshold: 0.003;
see Figure 3d).

442
443
Figure 3. The effect of Monocular Deprivation (MD) on visual processing, visual MD effect. (a) Oscillatory activity
calculated as the difference between t1 minus t0 at each eye: *PowChangeDeprived* (upper row) and
PowChangeUndeprived (bottom row) are plotted as a function of time [-0.25 - 0.5 s] and frequency [4-30 Hz]. The
plots show the average across occipital electrodes (E36, E38, E40); 0 s indicates stimulus onset. Topographies in the
alpha range [10-16 Hz] at a representative time window [0.02 - 0.06 s]; arrows indicate which eye was stimulated (here
depicted a participant with right-eye dominance); crosses represent the eye covered by a translucent patch. (b) Statistical
results. Time-frequency plot highlighting significant differences between *PowChangeDeprived* and
PowChangeUndeprived identified by the cluster-based permutation test (p<0.025, two-tailed) and the corresponding
topography for the alpha range [10-16 Hz] at a representative time window [0.02 - 0.06 s]; electrodes belonging to the
significant cluster are highlighted with white asterisks. (c) Time-course at the group level of the mean power in the
alpha range [10-16 Hz] at t0 and t1 (data are averaged across electrodes E36, E38, E40) separately displayed for the
Deprived and the Undeprived eye (upper and bottom rows); shaded areas represent the standard error of the mean; the
continuous horizontal grey line indicates the significant difference between t0 and t1 in the Deprived eye (from 0 to
0.14 s; p<0.05, FDR corrected). The dashed grey boxes represent the time window [0.02 - 0.06 s], comprising the alpha
peak, in which the power in the alpha range [10-16 Hz] was extracted for each subject (across channels E36, E38, E40)
and shown in the corresponding violin plots (right side; each dot represents individual data). (d) Source analysis
performed to localize the visual MD effect; the image shows, at 40 ms after stimulus onset, the area in which the power
in the alpha range [10 - 16 Hz] significantly decreased at t1 with respect to t0.

Audio-visual

After we assessed the impact of MD on unisensory visual processing (visual MD effect), we
investigated whether MD can also affect audio-visual processing at the neural level. The induced
and evoked oscillatory activities were tested separately within low and high-frequency ranges.

Induced power. The cluster-based permutation performed on induced oscillatory activity within the low-frequency range [4-30 Hz] between *PowChangeDeprived* and *PowChangeUndeprived* showed no significant effects (all ps>0.45, see Figure S5). In contrast, the same analysis performed within the high-frequency range [30-80 Hz] revealed a significant effect in gamma activity (p<0.015). An increase in gamma activity [65-75 Hz] was found in the Undeprived eye between 0.16 and 0.26 s, mainly in posterior electrodes (see Figure 4a, b). The planned comparison showed no significant effect at t0 (all ps>0.07), while a significant difference between the two eyes emerged only at t1 (p<0.022), confirming that MD specifically guided the effect. To further investigate the time-course of the induced gamma effect, we tested the difference between t0 and t1 within each eye. For each session and participant, we extracted the mean induced gamma power [65-75 Hz] measured across two parieto-occipital electrodes (E40 and E42, which corresponded to the peak of the statistical effect in the *PowChangeDeprived* vs. *PowChangeUndeprived* cluster-based permutation test). A series of paired t-tests were performed between t0 and t1 for each eye, at each time-point within the whole-time window of interest [0-0.5 s] (FDR corrected, q=0.05). Only for the Undeprived eye, a significant effect was found (from 0.12 to 0.2 s; for Deprived eye all ps>0.5; see Figure 4c).

The gamma band increase during audio-visual processing in the Undeprived eye following MD might suggest an upweighting of the auditory modality. If this would be the case, a greater neural response in the gamma range to unimodal acoustic stimulation (A and AA condition) should emerge selectively for the Undeprived eye after MD. To this end, a hypothesis-driven (p<0.05, one tail) cluster-based permutation test was performed on the induced oscillatory activity in response to auditory stimuli (average across A and AA trials) in the high-frequency range [30-80 Hz] across all electrodes, frequencies, and time-points [0-0.5 s], between t0 and t1, separately in each eye. The analysis revealed a significant increase in gamma activity between 100 and 300 ms in response to auditory stimulation after MD (p<0.04), selectively for the Undeprived eye (see Figure 5). Conversely, no significant difference emerged for the Deprived eye (p>0.35). This significant effect emerged in parietal electrodes as for the audio-visual MD effect. These findings support our hypothesis that the increased induced gamma activity during audio-visual processing is due to an upweighting of auditory input.

Evoked power. When the difference between *PowChangeDeprived* and *PowChangeUndeprived* was tested in the evoked power, no significant difference was found within the low-frequency range (all ps>0.09), nor within the high-frequency range (p>0.05, see Figure S7).

To sum up, the audio-visual MD effect selectively emerged for the Undeprived eye in the induced

gamma power. This neurophysiological change seems to be driven by increased responsiveness to

auditory inputs.

Source analysis

We investigated the sources of the audio-visual MD effect. To this end, a permutation paired t-test

(1000 randomizations) was performed at the source level, on the power in the gamma range [65-75

Hz] between t0 and t1 for the Undeprived eye (time window [0-0.5 s]; FDR correction was applied

on the time dimension).

Results revealed that audio-visual MD effect was mainly located at the cortical level around the

right intraparietal sulcus (corrected p-threshold: 0.002; see Figure 4d).

Figure 4. The effect of Monocular Deprivation (MD) on audio-visual processing, audio-visual MD effect. (a)

Oscillatory activity calculated as the difference between t1 minus t0 at each eye: *PowChangeDeprived* (upper row) and

PowChangeUndeprived (middle row) are plotted as a function of time [-0.25 - 0.5 s] and frequency [30-80 Hz]. The

plots show the average across posterior electrodes (E40, E42); 0 s indicates the stimulus onset. Topographies in the

gamma range [65-75 Hz] at representative time window [0.16 - 0.2 s]; arrows indicate which eye was stimulated

(represented in a participant with a right-eye dominance), and crosses represent the eye covered by a translucent patch.

(b) Statistical results. Time-frequency plot highlighting significant differences between *PowChangeDeprived* and

PowChangeUndeprived identified by the cluster-based permutation test (p<0.025, two-tailed) and corresponding

topography for the gamma range [65-75 Hz] at a representative time window [0.16 - 0.2 s]; electrodes belonging to the

significant cluster are highlighted with white asterisks. (c) Time-course at the group level of the mean power in gamma

range [65-75 Hz] at t0 and t1 (data are averaged across electrodes E40, E42) separately displayed for the Deprived and the Undeprived eye (upper and bottom rows); shaded areas represent the standard error of the mean; the continuous horizontal grey line indicates a significant difference between t0 and t1 for the Undeprived eye (from 0.12 to 0.2 s after stimulus onset; p<0.05, FDR corrected). The dashed grey boxes represent the time window [0.16 - 0.2 s] in which the power in the gamma range [65-75 Hz] was extracted for each subject (across channels E40 and E42) and shown in the corresponding violin plots (right side). (d) Source analysis performed to localize the audio-visual MD effect; the image shows at 120 ms after stimulus onset the area in which the power in the gamma range [65-75 Hz] significantly increases at t1 with respect to t0.

Figure 5. Increased induced gamma activity after MD during unisensory auditory processing. (a) Oscillatory activity at t0 and t1 are plotted as a function of time [-0.25 - 0.5 s] and frequency [30-80 Hz] for the Deprived (upper row) and Underprived eye (bottom row). The plots show the average across central-posterior electrodes (E34, E40, E41, E42, E50); 0 s indicates the stimulus onset. (b) Results of the cluster-based permutation test are shown in the time-frequency plots and highlight significant differences between t0 and t1, which emerged only for the Undeprived eye. Topography shows the results of the cluster-based permutation test for the Undeprived eye in the gamma range [50-60 Hz] over a representative time window [0.24 - 0.32 s]; white asterisks highlight significant electrodes (E34, E40, E41, E42, E50). (c) Time-course at the group level of the mean power in gamma range [50-60 Hz] at t0 and t1 (data are averaged across electrodes E34, E40, E41, E42, E50), separately displayed for the Deprived and the Undeprived eye (upper and bottom rows); shaded areas represent the standard error of the mean; the continuous horizontal grey line indicates a significant difference between t0 and t1 for the Undeprived eye (from 0.16 to 0.34 s after stimulus onset; p<0.05, FDR corrected).

Association between neural and behavioral changes due to MD

For visual and audio-visual conditions, we investigated the degree of association between brain activity alterations and changes in behavioral performance. The average power at the frequencies of

interest was extracted within a time window and across channels that resulted significant in the

cluster-based permutation tests (visual and audio-visual MD effects). At the individual level, we

computed the normalized difference between t1 and t0 within each eye ([*PowChangeDeprived/t0*

Deprived] * *100;* [*PowChangeUndeprived/t0 Undeprived*] * *100*). Shapiro-Wilk tests confirmed

that data in each condition were normally distributed (all ps >0.05). Thus, within each condition

(i.e., visual and audio-visual) and for each eye (i.e., Deprived and Undeprived), the power change

was correlated with the corresponding change in sensitivity between t0 and t1 using Pearson's

correlation coefficient.

Unisensory Visual

We first tested whether observed changes of oscillatory activity in induced alpha power following

MD (visual MD effect) were associated with changes in behavioral performance. To this aim, we

extracted the average power in the 10-16 Hz range between 0.02 and 0.06 s from the three

significant electrodes (E36, E38, and E40) for each session. Then, the normalized t1-t0 difference of

PowChangeDeprived and *PowChangeUndeprived* was computed.

A significant positive correlation between normalized *PowChangeDeprived* and visual sensitivity

change in the Deprived eye (d' change) was found (r(18) = 0.492, p=0.038). Following MD, the

visual sensitivity in the Deprived eye decreased in parallel with a reduction in induced alpha power

in the same eye. When the correlation between induced alpha change and visual sensitivity change

was tested for the Undeprived eye, no significant effect was found (p>0.99).

Audio-visual

Next, we tested whether the increase in induced gamma after MD period (audio-visual MD effect)

was associated with illusory fission percept. Thus, within each session, the induced power between

65 and 75 Hz was extracted within the 0.16-0.2 s time window and across the two significant

posterior channels (E40 and E42). Then, the normalized t1-t0 difference of *PowChangeDeprived*

and of *PowChangeUndeprived* was computed. No significant correlation was found neither for the

Deprived eye nor for the Undeprived eye (all ps>0.43). However, at the behavioral level, a tendency

for an interaction between illusion perception and sensory preferences was found (F(1,15)=4.01,

p=0.06), indicating that *Visual* subjects tended to experience less illusion than *Audio* subjects,

especially in the Undeprived eye (see section Audio-visual in the Behavioral Results). Interestingly,

the perception of the multisensory input is known to partially depend on individual sensory

predisposition (e.g., Giard and Perronet, 1999; Hong et al., 2021). Therefore, despite this was not

the focus of the paper, we additionally explored whether the association between neural and

behavioral changes might be affected by individual sensory preference. Specifically, we calculated
the correlation between *PowChangeUndeprived* and audio-visual d' change in the Undeprived eye
within *Audio* and *Visual* groups, classified according to participants' Sensory-Preference (estimated
for the same eye). Pearson correlation revealed a significant negative correlation in the V-group
$(r(5) = -0.937; p=0.019)$, while a tendency toward a significant positive correlation was found in the
A-group $(r(13) = 0.513; p=0.073)$. The two correlations significantly differed (difference=1.45 CI
[0.17 1.80]). Since Pearson correlations were performed, the confidence interval was adjusted as
described in Wilcox (2009); see Supplementary Materials Figure S8). While these results are based
on an explorative analysis and small sub-samples, they seem to suggest that the audio-visual MD
effect might have a different impact on the illusory percept according to the participants' Sensory-
Preference: in V-group, the increase of induced gamma activity in the Undeprived eye was
positively associated with a fission illusion increase (smaller d' after MD), while in A-group the
increase of induced gamma activity seemed to be associated with a fission illusion decrease (larger
d' after MD).

598
Figure 6. Visual and audio-visual MD effects on induced oscillatory activity. The plots show the % normalized
power change due to MD in unisensory visual and multisensory audio-visual processing. On the left, the visual MD
effect: decreased induced alpha power selectively for the Deprived eye (red) during unisensory visual processing. On
the right, the audio-visual MD effect: enhanced induced gamma power selectively for the Undeprived eye (blu) during
audio-visual processing. Each dot represents the group mean and bars the SE. Significant differences are highlighted
with * (red indicates effect for the Deprived eye and blu for the Undeprived eye).

Discussion

In this study, adult neural plasticity of both unisensory visual and multisensory audio-visual
processes was investigated to assess whether the impact of MD extends beyond visual processing.

Induced and evoked oscillatory activity changes both in visual and audio-visual processing were
measured after 150 minutes of altered visual experience (brief MD). Induced alpha associated with
the early phase of visual processing (<150ms) decreased after MD selectively for the Deprived eye.
Conversely, induced gamma associated with audio-visual processing increased after MD only in the
Undeprived eye, within a later temporal window (~100-300ms). Notably, both visual and audio-
visual processing alterations were found selectively for the induced component of neural
oscillations (see Figure 6). Source modeling linked both the visual and audio-visual MD effects to
the right parieto-occipital cortex. Our data reveal the specific neural signatures of temporary MD
effects on visual and audio-visual processes and shed light on their shared feedback nature. We
demonstrated that a brief period of monocular visual experience in adulthood specifically changes
the neural response to multisensory audio-visual events because of plasticity in feedback
connectivity.

621
Spectro-temporal properties of the visual MD effect
The observed visual MD effect is in line with previous studies (Lunghi et al., 2015a, 2015b; Zhou et
al., 2015; Binda et al., 2018; Schwenk et al., 2020). The reduced alpha synchronization observed
when the task was performed with the Deprived eye after MD is consistent with increased
excitability to compensate for the absence of stimulation during the deprivation phase. Source
modeling suggested that the alpha reduction was mainly localized in the right hemisphere in
superior parieto-occipital areas, with some activities extending to primary visual areas (calcarine
sulcus). These results confirm the modulation of alpha rhythm induced by short-term plasticity,
previously shown in the frequency domain as a change in alpha peak amplitude (Lunghi et al.,
2015a), and characterize the spectro-temporal properties of this neural effect. Namely, a selective
decrease of alpha synchronization [10-16 Hz] occurs in the early stages of visual processing
(<150ms). Notably, only induced activity (i.e., not phase-locked to stimulus onset) was altered by
MD, revealing the feedback nature of visual short-term plasticity. Overall, these results are coherent
with spectroscopy data showing an increase of excitability in the early visual cortex as indicated by
reductions of inhibitory gamma-aminobutyric acid (GABA) concentration after short-term MD
(Lunghi et al., 2015b). Moreover, selectively for the Deprived eye, a significant correlation was
found between the decrease in induced alpha activity and the decrease in visual sensitivity after
MD, suggesting a potential link between this neural change and the ability to discriminate temporal
aspects of visual processing.

641
The impact of MD on audio-visual processing

Previous behavioral studies have shown that multisensory perception can be altered by MD (Lo
Verde et al., 2017; Opoku-Baah and Wallace, 2020). By measuring changes in neural oscillations,
we assessed neural mechanisms underpinning multisensory short-term plasticity and revealed the
specific enhancement of induced gamma activity [65-75 Hz] selectively for the Undeprived eye
when processing audio-visual input. The audio-visual MD effect involved the right intraparietal
sulcus, suggesting its central role in short-term plasticity induced by MD.

We hypothesized that the audio-visual MD effect could be due to a crossmodal upweighting of the
other modality (i.e., audition). To this end, we tested whether the neural response to unimodal
auditory stimuli was increased following MD in each eye. The observation that gamma response to
auditory input increased following MD selectively for the Underprived eye confirmed our
hypothesis. This result revealed that short-term plasticity following MD alters both visual and
auditory neural representations. Interestingly, alterations of neural excitability due to temporary
binocular deprivation were previously shown to increase heteromodal responses in the visual cortex
(Merabet et al., 2008).

From a neurophysiological perspective, it is important to remark that neural profiles of short-term
plasticity following MD seem to depend on the type of input at hand. While the visual stimulus was
the same in visual and audio-visual conditions, short-term plasticity was characterized by specific
oscillatory fingerprints (Siegel et al., 2012): alpha decreased during visual processing while gamma
increased during audio-visual and auditory processing.

To what extent does this neural alteration interact with individuals' sensory predisposition? In the
Undeprived eye, the correlation between enhanced gamma activity in audio-visual processing and
behavioral performance seems to indicate opposite MD impacts on illusory perception with respect
to participants' Sensory-Preference. Although preliminary, this result opens the possibility that the
upweighting of auditory information during audio-visual processing after MD affects visual
perception according to individual sensory preference. Coherently with the extreme flexibility and
adaptability of multisensory functions (Giard and Perronet, 1999; Fujisaki et al., 2004; Van
Atteveldt et al., 2014), the perception of the multisensory input after MD might change as a
function of individual sensory predisposition. This inference should be further verified with
psychophysics experiments (see Rohe et al., 2019), designed to directly estimate auditory modality's
weight changes during audio-visual processing after MD. Interestingly, a recent study investigating
cross-modal recalibration highlighted how individual variability in one sensory modality (visual
reliability) differently affects recalibration of the other modality (audition) (Hong et al., 2021).

Binocular input was shown to be critical for developing audio-visual perception: anomalies in
different audio-visual perceptual tasks were reported in cases of monocular enucleation (Moro and

Steeves, 2018a, 2018b), individuals affected by early monocular cataracts (Chen et al., 2017), and people suffering from amblyopia (Narinesingh et al., 2015, 2017; Richards et al., 2017). The present results, revealing that MD induces short-term plasticity of audio-visual processing, encourage possible treatments of audio-visual anomalies associated with MD. Increasing evidence in animal studies supports clinical treatment of adult amblyopia (Hensch and Quinlan, 2018), and crucially a recent study conducted with adult people affected by amblyopia has shown that MD combined with physical exercise could be a promising clinical intervention to promote visual recovery (Lunghi et al., 2019). Future studies might help understanding whether multisensory audio-visual processing could also benefit from this novel clinical treatment.

The pivotal role of induced cortical response in experience-dependent plasticity

Both visual and audio-visual MD effects were selectively found in induced neural oscillations, likely reflecting main alterations in feedback processing integrating sensory input and ongoing cortical activity (Galambos 1992; Klimesch et al., 1998; Tallon-Baudry and Bertrand, 1999; Chen et al., 2012; Keil et al., 2022). Instead, the evoked phase-locked activity (feedforward), related to the processing within the thalamo-cortical pathway (Galambos 1992; Keil et al., 2022), was not affected.

Visual MD effect was found in the induced alpha band, which was previously demonstrated to be drastically impaired by the transient absence of visual experience during development (Bottari et al., 2016). Animal studies involving congenital deaf cats demonstrated that induced and not evoked oscillatory activity in the auditory cortex is extremely reduced across a wide range of frequency bands (e.g., Yusuf et al., 2017). The authors hypothesized that the absence of sensory stimulation prevents the development of neural mechanisms allowing the integration of sensory signals and internal representations. Evidence of alterations of induced oscillatory activity following sensory deprivation was reported not only within modality, for visual (Bottari et al., 2016) and auditory systems (Yusuf et al., 2017), but also cross-modally. In humans, early-onset deafness selectively affects induced oscillatory activity associated with visual processing (Bednaya et al., 2021). The present audio-visual MD effect (increased induced gamma activity) provides evidence in the same direction also for multisensory processing. Taken together, this evidence suggests a substantial alteration of cortico-cortical feedback activities, in case of sensory input absence in both developmental and adult brain, for unisensory and multisensory functions. This is in line with previous evidence suggesting that the plasticity of feedback connectivity represents an extremely flexible mechanism to process sensory information according to changing demands (Polley et al., 2006).

Limitations of the study

Here, we focused on neural oscillations changes, and given the time constraints intrinsic in the MD

effect (which is maximum within the first 15 minutes following the deprivation, Lunghi et al.,

2011), we could not insert a direct measure of ocular dominance, which is usually assessed with

binocular rivalry (e.g., Lunghi et al., 2011). Thus, we relied on eye-specific oscillatory activity

alterations selectively at t1 (and their lack at t0) as indications of the change in interocular

excitability balance: the decreased alpha during visual processing is compatible with the

strengthening of the Deprived eye after MD. While a clear reduction of strength for the Undeprived

eye (i.e., increased alpha activity) did not emerge, this is in line with the literature: the strongest

impact of MD is known to be on the Deprived eye, while the opposite effect on the Undeprived eye

was found to be much smaller (Lunghi et al., 2015a; Binda et al., 2018) or even absent (Zhou et al.,

2015). Thus, it is possible that larger sample sizes are required to measure neural effects on the

Undeprived eye during visual processing. Further studies might help to assess whether depriving

the non-dominant eye will lead to the same MD effects. However, given the absence of difference at

t0, we can rule out the possibility that our effects are due to baseline differences between the

dominant and the non-dominant eye.

Noticeably, the visual MD effect emerged here for the Deprived eye, while the audio-visual MD

effect emerged only for the Undeprived eye. While these effects are in line with previous behavioral

studies (Lo Verde et al., 2017; Opoku-Baah and Wallace, 2020), they also support a crucial role of

MD in inducing flexible alterations of interocular excitability balance and, in turn, in audio-visual

processes. Moreover, although an interaction between MD and eye dominance cannot be excluded,

a recent study reported that the MD effect was the same regardless of whether the dominant or the

non-dominant eye was deprived (Schwenk et al., 2020).

Finally, while the interaction between sensory preference and gamma change in audio-visual

processing seems of interest, a larger sample with a balanced number of subjects with visual and

auditory sensory preferences is needed to confirm our intriguing preliminary result on how

multisensory plasticity can be affected by individual sensory predisposition.

Conclusions

741 These results demonstrated that a brief period of monocular visual experience in adulthood is able

742 to change the neural response not only to visual stimuli but also to multisensory events. The data

743 unveiled the spectral fingerprints of adult short-term plasticity induced by a brief period of MD for

visual and audio-visual processing. We found enhanced excitability (i.e., decreased induced alpha activity) for the Deprived eye during an early phase of visual processing, and we demonstrated the presence of neural alterations beyond the visual processing. Gamma activity associated with audio-visual processing was increased by MD at a later latency and only when the task was performed with the Undeprived eye. The analyses of responses to unimodal auditory input indicate an upweighting of sound input following MD, selectively for the Undeprived eye. Importantly, these distinct neural effects were selectively found in the induced neural oscillations, revealing that experience-dependent plasticity mainly involves alterations in feedback processing not only during development but also in adulthood. This observation is consistent with the existence of a general mechanism shared across sensory modalities and across the life cycle.

Acknowledgments

This work was funded by PRIN 2017 research grant (Prot. 20177894ZH) to Davide Bottari.

We would like to thank Cesare V. Parise for the helpful discussion and feedback regarding data analysis.

Declaration of Competing Interest

The authors declare no competing financial interests.

Data Availability

The datasets and code used in the present study are available from the corresponding author on reasonable request .

CRediT authorship contribution statement

Alessandra Federici: Conceptualization, Investigation, Data curation, Formal analysis, Writing - original draft, Writing - review & editing. Giulio Bernardi: Conceptualization, Writing - review & editing. Irene Senna: Formal analysis, Writing - review & editing. Marta Fantoni: Investigation. Marc Ernst: Formal analysis, Writing - review & editing. Emiliano Ricciardi: Resources, Writing - review & editing. Davide Bottari: Conceptualization, Investigation, Data curation, Formal analysis,

Writing - original draft, Writing - review & editing, Supervision, Project administration, Funding acquisition.

References

Balz J, Keil J, Roa Romero Y, Mekle R, Schubert F, Aydin S, Ittermann B, Gallinat J, Senkowski D (2016) GABA concentration in superior temporal sulcus predicts gamma power and perception in the sound-induced flash illusion. Neuroimage 125:724–730

Bednaya E, Pavani F, Ricciardi E, Pietrini P, Bottari D (2021) Oscillatory signatures of Repetition Suppression and Novelty Detection reveal altered induced visual responses in early deafness. Cortex.

Bell AJ, Sejnowski TJ (1995) An information-maximization approach to blind separation and blind deconvolution. Neural Comput 7:1129–1159.

Bhattacharya J, Shams L, Shimojo S (2002) Sound-induced illusory flash perception: Role of gamma band responses. Neuroreport 13:1727–1730.

Binda P, Kurzawski JW, Lunghi C, Biagi L, Tosetti M, Morrone MC (2018) Response to short-term deprivation of the human adult visual cortex measured with 7T bold. Elife 7:1–25.

Bottari D, Bednaya E, Dormal G, Villwock A, Dzhelyova M, Grin K, Pietrini P, Ricciardi E, Rossion B, Röder B (2020) EEG frequency-tagging demonstrates increased left hemispheric involvement and crossmodal plasticity for face processing in congenitally deaf signers. Neuroimage 223:117315.

Bottari D, Troje NF, Ley P, Hense M, Kekunnaya R, Röder B (2016) Sight restoration after congenital blindness does not reinstate alpha oscillatory activity in humans. Sci Rep 6.

Castaldi E, Lunghi C, Morrone MC (2020) Neuroplasticity in adult human visual cortex. Neurosci Biobehav Rev 112:542–552.

Chen CC, Kiebel SJ, Kilner JM, Ward NS, Stephan KE, Wang WJ, Friston KJ (2012) A dynamic causal model for evoked and induced responses. Neuroimage 59:340–348.

Chen YC, Lewis TL, Shore DI, Maurer D (2017) Early Binocular Input Is Critical for Development of Audiovisual but Not Visuotactile Simultaneity Perception. Curr Biol 27:583–589.

Cooke J, Poch C, Gillmeister H, Costantini M, Romei V (2019) Oscillatory properties of functional

connections between sensory areas mediate cross-modal illusory perception. J Neurosci 39:5711–5718.

Dale AM, Liu AK, Fischl BR, Buckner RL, Belliveau JW, Lewine JD, Halgren E (2000) Dynamic statistical parametric mapping: combining fMRI and MEG for high-resolution imaging of cortical activity. Neuron 26:55–67.

Delorme A, Makeig S (2004) EEGLAB: An open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. J Neurosci Methods 134:9–21.

Delorme A, Sejnowski T, Makeig S (2007) Enhanced detection of artifacts in EEG data using higher-order statistics and independent component analysis. Neuroimage 34:1443–1449.

Espinosa JS, Stryker MP (2012) Development and Plasticity of the Primary Visual Cortex. Neuron 75:230–249.

Farahani ED, Wouters J, van Wieringen A (2019). Contributions of non-primary cortical sources to auditory temporal processing. NeuroImage, 191: 303-314.

Froesel M, Cappe C, Ben Hamed S (2021) A multisensory perspective onto primate pulvinar functions. Neurosci Biobehav Rev 125:231–243.

Fujisaki W, Shimojo S, Kashino M, Nishida S (2004) Recalibration of audiovisual simultaneity. Nat Neurosci 2004 77 7:773–778.

Galambos, R. (1992). A comparison of certain gamma band (40-Hz) brain rhythms in cat and man. In Induced rhythms in the brain (pp. 201-216). Birkhäuser, Boston, MA.

Giard MH, Peronnet F (1999). Auditory-visual integration during multimodal object recognition in humans: a behavioral and electrophysiological study. Journal of cognitive neuroscience, 11:473-490.

Hansen P, Kringelbach M, Salmelin R (2010) MEG: An Introduction to Methods. Oxford University Press (OUP).

Hensch TK (2005) Critical period plasticity in local cortical circuits. Nat Rev Neurosci 6:877–888.

Hensch TK, Quinlan EM (2018) Critical periods in amblyopia. Vis Neurosci 35:E014.

Hirst RJ, McGovern DP, Setti A, Shams L, Newell FN (2020) What you see is what you hear: Twenty years of research using the Sound-Induced Flash Illusion. Neurosci Biobehav Rev 118:759–774.

Hong, F., Badde, S., & Landy, M. S. (2021). Causal inference regulates audiovisual spatial
recalibration via its influence on audiovisual perception. PLoS computational biology, 17(11),
e1008877.

Hubel DH, Wiesel TN (1970) The period of susceptibility to the physiological effects of unilateral
eye closure in kittens. J Physiol 206:419–436.

Jensen O, Bonnefond M, VanRullen R (2012) An oscillatory mechanism for prioritizing salient
unattended stimuli. Trends Cogn Sci 16:200–206.

Jensen O, Gips B, Bergmann TO, Bonnefond M (2014) Temporal coding organized by coupled
alpha and gamma oscillations prioritize visual processing. Trends Neurosci 37:357–369.

Jensen O, Mazaheri A (2010). Shaping functional architecture by oscillatory alpha activity: gating
by inhibition. Frontiers in human neuroscience, 186.

Jung T-P, Makeig S, Humphries C, Lee T-W, McKeown MJ, Iragui V, Sejnowski TJ (2000a)
Removing electroencephalographic artifacts by blind source separation. Psychophysiology
37:163–178.

Jung TP, Makeig S, Westerfield M, Townsend J, Courchesne E, Sejnowski TJ (2000b) Removal of
eye activity artifacts from visual event-related potentials in normal and clinical subjects. Clin
Neurophysiol 111:1745–1758.

Karmarkar UR, Dan Y (2006) Experience-dependent plasticity in adult visual cortex. Neuron
52:577–585.

Keil J (2020) Double Flash Illusions: Current Findings and Future Directions. Front Neurosci 14:1–
8.

Keil, A., Bernat, E. M., Cohen, M. X., Ding, M., Fabiani, M., Gratton, G., ... & Weisz, N. (2022).
Recommendations and publication guidelines for studies using frequency domain and
time‐frequency domain analyses of neural time series. Psychophysiology, 59(5), e14052.

Klimesch W (2012) Alpha-band oscillations, attention, and controlled access to stored information.
Trends Cogn Sci 16:606–617.

Klimesch W, Russegger H, Doppelmayr M, Pachinger T (1998) A method for the calculation of
induced band power: Implications for the significance of brain oscillations. Electroencephalogr
Clin Neurophysiol 108:123–130.

Klimesch W, Sauseng P, Hanslmayr S (2007) EEG alpha oscillations: The inhibition-timing

hypothesis. Brain Res Rev 53:63–88.

Lange J, Oostenveld R, Fries P (2011) Perception of the touch-induced visual double-flash illusion
correlates with changes of rhythmic neuronal activity in human visual and somatosensory
areas. Neuroimage 54:1395–1405.

Lange J, Oostenveld R, Fries P (2013) Reduced Occipital Alpha Power Indexes Enhanced
Excitability Rather than Improved Visual Perception. J Neurosci 33:3212–3220.

Lennert T, Samiee S, Baillet S (2021) Coupled oscillations enable rapid temporal recalibration to
audiovisual asynchrony. Commun Bio, 4:1-12.

Levelt CN, □ubener M (2012) Critical-period plasticity in the visual cortex. Annu Rev Neurosci
35:309–330.

Lo Verde L, Morrone MC, Lunghi C (2017) Early Cross-modal Plasticity in Adults. J Cogn
Neurosci 29:520–529.

Lunghi C, Berchicci M, Morrone MC, Di Russo F (2015a) Short-term monocular deprivation alters
early components of visual evoked potentials. J Physiol 593:4361–4372.

Lunghi C, Burr DC, Morrone C (2011) Brief periods of monocular deprivation disrupt ocular
balance in human adult visual cortex. Curr Biol 21:538–539.

Lunghi C, Emir UE, Morrone MC, Bridge H (2015b) Short-Term monocular deprivation alters
GABA in the adult human visual cortex. Curr Biol 25:1496–1501.

Lunghi C, Sale A (2015) A cycling lane for brain rewiring. Curr Biol 25:R1122–R1123.

Lunghi C, Sframeli AT, Lepri A, Lepri M, Lisi D, Sale A, Morrone MC (2019) A new
counterintuitive training for adult amblyopia. Ann Clin Transl Neurol 6:274–284.

Merabet LB, Hamilton R, Schlaug G, Swisher JD, Kiriakopoulos ET, Pitskel NB, Kauffman T,
Pascual-Leone A (2008). Rapid and reversible recruitment of early visual cortex for touch.
PLoS one, 3:e3046.

Mishra J, Martinez A, Sejnowski TJ, Hillyard SA (2007) Early cross-modal interactions in auditory
and visual cortex underlie a sound-induced visual illusion. J Neurosci 27:4120–4131.

Moro SS, Steeves JKE (2018a) Normal temporal binding window but no sound-induced flash
illusion in people with one eye. Exp Brain Res 236:1825–1834.

Moro SS, Steeves JKE (2018b) Audiovisual plasticity following early abnormal visual experience:

Reduced McGurk effect in people with one eye. Neurosci Lett 672:103–107.

Narinesingh C, Goltz HC, Raashid RA rha., Wong AMF (2015) Developmental Trajectory of

McGurk Effect Susceptibility in Children and Adults With Amblyopia. Invest Ophthalmol Vis

Sci 56:2107–2113.

Narinesingh C, Goltz HC, Wong AMF (2017) Temporal binding window of the sound-induced

flash illusion in amblyopia. Investig Ophthalmol Vis Sci 58:1442–1448.

Oostenveld R, Fries P, Maris E, Schoffelen JM (2011) FieldTrip: Open source software for

advanced analysis of MEG, EEG, and invasive electrophysiological data. Comput Intell

Neurosci 2011.

Opoku-Baah C, Wallace MT (2020) Brief period of monocular deprivation drives changes in

audiovisual temporal perception. J Vis 20:1–13.

Pérez-Bellido A, Ernst MO, Soto-Faraco S, López-Moliner J (2015) Visual limitations shape audio-

visual integration. J Vis 15:1–15.

Polley DB, Steinberg EE, Merzenich MM (2006). Perceptual learning directs auditory cortical map

reorganization through top-down influences. Journal of neuroscience, 26:4970-4982.

Reh RK, Dias BG, Nelson CA, Kaufer D, Werker JF, Kolbh B, Levine JD, Hensch TK (2020)

Critical period regulation acrossmultiple timescales. Proc Natl Acad Sci U S A 117:23242–

23251.

Richards MD, Goltz HC, Wong AMF (2017) Alterations in audiovisual simultaneity perception in

amblyopia. PLoS One 12:1–20.

Rohe T, Ehlis AC, Noppeney U. (2019). The neural dynamics of hierarchical Bayesian causal

inference in multisensory perception. Nature Communications, 10:1-17.

Schwenk JCB, VanRullen R, Bremmer F (2020) Dynamics of Visual Perceptual Echoes Following

Short-Term Visual Deprivation. Cereb Cortex Commun 1:1–11.

Shams L, Kamitani Y, Shimojo S (2000) What You See Is What You Hear. Nature 408:2000.

Siegel M, Donner TH, Engel AK (2012) Spectral fingerprints of large-scale neuronal interactions.

Nat Rev Neurosci 13:121–134.

Spolidoro M, Sale A, Berardi N, Maffei L (2009) Plasticity in the adult brain: Lessons from the

visual system. Exp Brain Res 192:335–341.

Stropahl M, Bauer AKR, Debener S, Bleichner MG (2018) Source-Modeling auditory processes of
EEG data using EEGLAB and brainstorm. Front Neurosci 12:1–11.

Tadel F, Baillet S, Mosher JC, Pantazis D, Leahy RM (2011) Brainstorm: A user-friendly
application for MEG/EEG analysis. Comput Intell Neurosci 2011.

Tallon-Baudry C, Bertrand O (1999) Oscillatory gamma activity in humans and its role in object
representation. Trends Cogn Sci 3:151–162.

Turrigiano G (2012) Homeostatic synaptic plasticity: Local and global mechanisms for stabilizing
neuronal function. Cold Spring Harb Perspect Biol 4.

Turrigiano GG, Nelson SB (2004). Homeostatic plasticity in the developing nervous system. Nature
reviews neuroscience, 5:97-107.

Tyll S, Budinger E, Noesselt T (2011) Thalamic influences on multisensory integration. Commun
Integr Biol 4:378–381.

Van Atteveldt N, Murray MM, Thut G, Schroeder CE (2014) Multisensory integration: Flexible use
of general operations. Neuron 81:1240–1253.

Vanes LD, White TP, Wigton RL, Joyce D, Collier T, Shergill SS (2016) Reduced susceptibility to
the sound-induced flash fusion illusion in schizophrenia. Psychiatry Res 245:58–65.

VanRullen R (2016) Perceptual Cycles. Trends Cogn Sci 20:723–735.

Viola FC, Thorne J, Edmonds B, Schneider T, Eichele T, Debener S (2009) Semi-automatic
identification of independent components representing EEG artifact. Clin Neurophysiol
120:868–877.

Watkins S, Shams L, Tanaka S, Haynes JD, Rees G (2006) Sound alters activity in human V1 in
association with illusory visual perception. Neuroimage 31:1247–1256.

Whittingham KM, McDonald JS, Clifford CWG (2014) Synesthetes show normal sound-induced
flash fission and fusion illusions. Vision Res 105:1–9.

Yusuf PA, Hubka P, Tillein J, Kral A (2017) Induced cortical responses require developmental
sensory experience. Brain 140:3153–3165.

Zhou J, Baker DH, Simard M, Saint-Amour D, Hess RF (2015) Short-term monocular patching
boosts the patched eye's response in visual cortex. Restor Neurol Neurosci 33:381–387.