Adaptation to ER-stress via serine glycine metabolism licences STING signalling and

CMV control in intestinal epithelial cells

Authors

Felix Wottawa[1*], Björn Becker[2*], Mohamed Bakr[1*], Lisa Mayr[3], Julia Kugler[1], Samuel J

Windross[4], Laura Neises[2], Silke van den Bossche[1], Nassim Kakavand[1], Florian Tran[1,13], Lina

Welz[1,13], Julian Schwärzler[3], Zoltán Bánki[5] Stephanie Stengel[1], Go Ito[6], Olivia Coleman[7],

Christian Jaeger[8], Eric Koncina[9], Elisabeth Letellier[9], Dirk Haller[7], Søren R Paludan[4], Richard

Blumberg[10], Arthur Kaser[11], Luka Cicin-Sain[12], Stefan Schreiber[1,13], Timon Adolph[3], Philip

Rosenstiel[1+] , Johannes Meiser[2+] and Konrad Aden[1,13+]

Affiliation

1 Institute of Clinical Molecular Biology, Christian-Albrechts-University and University

Hospital Schleswig-Holstein, Campus Kiel, 24105 Kiel, Germany

2 Luxembourg Institute of Health, Department of Cancer Research, Luxembourg

3 Department of Internal Medicine I, Gastroenterology, Hepatology, Metabolism &

Endocrinology, Medical University of Innsbruck, Innsbruck, Austria

4 Department of Biomedicine, Aarhus University, Aarhus, Denmark

5 Institute of Virology, Department of Hygiene, Microbiology and Public Health, Medical

University of Innsbruck, Innsbruck, Austria.

6 Department of Gastroenterology and Hepatology, Tokyo Medical and Dental University,

Tokyo, Japan;

7 Chair of Nutrition and Immunology, TUM School of Life Sciences, Technical University of

Munich

8 Luxembourg Centre for Systems Biomedicine, University of Luxembourg, Esch-sur-Alzette,

Luxembourg

9 Faculty of Science, Technology and Medicine, Department of Life Sciences and Medicine,

Université du Luxembourg

10 Gastroenterology Division, Department of Medicine, Brigham and Women's Hospital,

Harvard Medical School, Boston, MA.

12 Division of Gastroenterology and Hepatology, Department of Medicine, Addenbrooke's

Hospital, University of Cambridge, Cambridge, England, UK

12 Helmholtz Zentrum für Infektionsforschung, Braunschweig, Germany

13 Department of Internal Medicine I., Christian-Albrechts-University and University Hospital

Schleswig-Holstein, Campus Kiel, 24105 Kiel, Germany

*These authors contributed equally

[+] shared senior authorship

Correspondence to:

Konrad Aden, MD

Department of Internal Medicine I,

University Medical Center Schleswig Holstein

Rosalind Franklin Straße 11,

24105 Kiel, Germany

T: +49-431 500-22455,

F: +49 431-500-15168,

E-Mail: k.aden@ikmb.uni-kiel.de

Philip Rosenstiel, MD

Professor of Molecular Medicine

Director, Institute of Clinical Molecular Biology,

Christian-Albrechts-University Kiel

Rosalind Franklin Straße 11, 24105 Kiel, Germany

T: +49-431 500-15111,

F: +49 431-500-15168,

E-Mail: p.rosenstiel@mucosa.de

Johannes Meiser, PhD

Group Leader of Cancer Metabolism Group

Department of Cancer Research

Luxembourg Institute of Health

84, Val Fleuri,

L-1526 Luxembourg

Luxembourg

T: +352 26970-453

Email: Johannes.Meiser@lih.lu

Disclosures: The authors have no conflict of interest.

Author Contributions

PR, KA, JM, FW and BB designed the study.

FW, BB, MB, LM, JK, SJW, LN, SB, NK, FT, LW, SSt, GI, OC, CJ, EK, EL, DH, SRP, TA,

JM, PR and KA performed experiments and analyzed data.

LCS, RB, AK and SSc provided intellectual input and guidance.

PR, KA, JM, FW and BB planned the project and supervised the experiment.

PR, KA, JM and FW wrote the manuscript.

Highlights:

-    **Chronic ER stress abrogates gut epithelial cGAS/STING signalling and increases**
susceptibility to CMV infection
-    **Chronic ER stress increases serin-glycin metabolism to counterbalance reactive oxygen**
species overload in IECs
-    **Abrogation of serine-glycine metabolism exacerbates ROS generation and inhibits**
epithelial STING signalling.
-    **Pharmacologic ROS scavenging in ER stressed epithelium restores cGAS/STING**
activation and leads to improved control of CMV infection

Abstract:

Mucosal cytomegalovirus (CMV) infection represents a leading cause for complicated disease

behaviour and proctocolectomy in patients with inflammatory bowel disease (IBD). Using a

genetic loss-of-function mouse model of the hypomorphic IBD risk gene X-box-binding

protein 1 (*XBP1*), isotope-assisted metabolomics and pharmacologic approaches, we unravel

the molecular control of gut epithelial STING signalling by unresolved endoplasmic reticulum

(ER) stress. We demonstrate that unresolved epithelial ER stress, evoked by *Xbp1* deficiency,

leads to exhausted STING signalling and an impaired ability to control CMV infection, which

is driven by the generation of reactive oxygen species (ROS). ROS generation is controlled by

cellular glycine influx and de-novo serin synthesis enabling glutathione production, which can

be pharmacologically exploited to restrore STING signaling. Pharmacological scavenging of

ROS with N-acetly cysteine restores epithelial STING signalling and limits CMV infection in

IECs. Our findings unravel the serine-glycine dependent metabolic control by ER stress that

licences STING signaling and susceptibility to infection in gut epithelium.

[Words: 156/160]

Important abbreviations used in this paper:

2'3'-cGAMP: 2'3' linked cyclic guanosine monophosphate-adenosine monophosphate; *ATF6*:
Activating transcription factor 6; *BI*: BI-4924; *cGAS*: Cyclic GMP-AMP synthase; *CMV*:
cytomegalovirus; *CORE*: consumption and release; *CXCL10*: C-X-C Motif Chemokine Ligand
10; *DAMP*: danger-associated molecular pattern; *DMXAA*: dymethylxanthone acetic acid;
dsDNA: double-stranded DNA; *ER*: endoplasmic reticulum; *ER-stress*: endoplasmic reticulum
stress; *GGC*: dipeptide γ-glutamyl cysteine; *GPX*: glutathione peroxidases; *GS*: gluthathione
synthetase; *GSH*: gluthathione; *GSSG*: glutathione disulphide; *IBD*: inflammatory bowel
disease; *IECs*: intestinal epithelial cells; *IFNβ*: interferon beta; *IL*: interleukin; *IRF3*:
interferon-regulatory-factor 3; *ISG*: interferon-stimulated gene; *L. monocytogenes*: *Listeria
monocytogenes*; *LPO*: lipid peroxidation; *MCMV*: murine cytomegalovirus; *MEF*: mouse
embryonic fibroblasts; *MOI*: multiplicity of infection; *NAC*: N-acetly cysteine; *PAMP*:
pathogen-associated molecular pattern; *ROS*: reactive oxygen species; *RSL3*: RAS-selective
lethal 3; *S/G*: serine-glycine; *SNP*: single nucleotide polymorphism; *SSP*: serine *de novo*
synthesis pathway; *STING*: stimulator of interferon genes; *TBK1*: tank-binding kinase 1; *TM*:
Tunicamycin; *TNF*: tumor necrosis factor; *UC*: ulcerative colitis; *UPR*: unfolded protein
response; *XBP1*: X-box binding protein 1

Introduction:

Inflammatory bowel disease is a chronic relapsing inflammatory disorder of the gastrointestinal

tract. Although the introduction of targeted inhibition of pro-inflammatory pathways (e.g. anti-

TNF, anti-IL12/23) has significantly improved patient care, disease-associated complications

such as IBD-related hospitalizations and surgery still contribute to the major socioeconomic

burden of IBD (1). IBD-related cytomegalovirus (CMV) infections have seen a steep rise in

the past years and recurrent CMV-infections in IBD patients are associated with increased risk

of hospitalization and proctocolectomy (2). Although meta-analysis of retrospective clinical

studies indicate that disease extent and disease severity directly correlate with CMV colitis

risk, the underlying molecular mechanisms that drive susceptibility to CMV infection in IBD

remain enigmatic (3).

Endoplasmic reticulum stress (ER-stress) describes a state of disrupted cellular homeostasis,

which occurs if the balance between ER folding capacity and protein load is disturbed(4–9).

Cells with a high secretory demand such as intestinal epithelial cells (IECs, more specifically

Paneth and Goblet cells) are especially prone to accumulate unfolded proteins and thus highly

susceptible to ER-stress-associated pathologies (9,10). Ultimately, ER-stress triggers a

detrimental inflammatory reaction that drives mucosal tissue damage in human IBD (11,12).

The unfolded protein response (UPR) consists of three major pathways (specifically, the

IRE1/XBP1-, PERK/ATF4- and ATF6-pathway) and aims to reinforce ER homeostasis, thus

counteracting inflammation. Specific single-nucleotide-polymorphisms (SNPs) in the *X-box

binding protein 1* (*XBP1*) gene result in hypomorphic variant predisposing for ER-stress and

have been identified to mediate increased IBD susceptibility and mechanistically augment

epithelial-derived mucosal inflammation (13,14).

The cGAS/STING (cyclic GMP-AMP synthase/stimulator of interferon genes) pathway allows

nucleic acid sensing and pathogen defence, by directly sensing cytosolic double-stranded DNA

(dsDNA) derived from viruses (e.g., CMV) or cyclic dinucleotides derived from intracellular

bacteria (e.g., *Listeria monocytogenes*) (15–17). After binding of cytosolic dsDNA and

subsequent 2′3′-cGAMP (2'3' linked cyclic guanosine monophosphate-adenosine

monophosphate) production by cGAS, STING is activated. STING is an ER-resident adaptor

protein which mediates activation of TBK1 (Tank-binding-kinase 1) and IRF3 (Interferon-

regulatory-factor 3), controlling interferon beta (IFNβ) production and secretion. IFNβ belongs

to the family of type I IFNs and induces a variety of antiviral, immunomodulatory or

antiproliferative interferon-stimulated genes (ISGs), e.g. *CXCL10* (17). As such, cGAS/STING

signaling is critically involved in type I IFN induction during CMV infection, and CMV

actively antagonizes STING-mediated IFN induction, underscoring the potential role of

STING as a crucial first line of defense against CMV infection (18–20). Due to its localization

as an ER-resident protein, it is postulated that STING function is closely intertwined with ER

homeostasis. To which extent disturbance of the UPR directly affects cGAS/STING sensing

is not known. Several studies support the notion that acute ER stress is an important upstream

modulator of STING activation in response to PAMPs and DAMPs (21,22) and pro-

regenerative IL-22 signaling (23). We hypothesised that chronic unresolved ER stress directs

epithelial STING signaling in response to CMV infection. We find that chronic ER stress

instigates immune exhaustion of the STING pathway in the intestinal epithelium which renders

IECs susceptible to CMV infection. Using a genetic model of chronic ER stress, i.e. conditional

deletion of $Xbp1$ in intestinal epithelial cells ($Xbp1^{\Delta IEC}$), we show that immune exhaustion of

STING depends on accumulation of reactive oxygen species due to insufficient scavenging by

glutathione. We show that epithelial STING signaling relies on the metabolic control of serine-

glycine dependent glutathione synthesis during ER stress, which allows to cope with

environmental challenges, such as CMV infection.

Results & Discussion:

Chronic ER-stress suppresses STING signaling in intestinal epithelial cells

Chronic intestinal inflammation has been associated with risk for CMV infection in IBD. As

cGAS/STING signaling is involved in CMV control and cytomegalovirus induces the unfolded

the $Xbp1$ branch of the unfolded protein response(24), we tested a potential crosstalk between

chronic ER stress and cGAS/STING signaling in response to CMV infection.

We used a transgenic mouse model that allows intestinal epithelial deletion of the $Xbp1$ gene

in mice ($Xbp1^{\Delta IEC}$, $Xbp1^{fl/fl}$) or and knockdown in a non-transformed small intestinal epithelial

cell line by CRISPR Cas9 gene editing (Mode-K cells; *iCtrl* and *iXbp1*) (5,6). We first observed

that chronic epithelial ER stress impairs STING expression in *Xbp1*$^{\Delta IEC}$ mice when compared

to *Xbp1*$^{fl/fl}$ (WT) littermates, which was unrelated to STING transcription (Fig. 1 A, B, Fig.

S1A, B). In line with this, we observed reduced expression of *Cxcl10* and *Ifnb* in small

intestinal crypts from *Xbp1*$^{\Delta IEC}$ mice, indicative of reduced STING/IFN-I activity (Fig. 1C, Fig.

S1C). Next, we investigated whether chronic ER stress impairs cGAS/STING signalling. To

do so, we stimulated *iCtrl* and *iXbp1* IECs with double-stranded DNA (dsDNA) and observed

a nearly complete abrogation of STING activity (Fig. S1D) and *Cxcl10* expression (Fig. S1E)

in *Xbp1*-deficient IECs. This observation was phenocopied by chemical induction of chronic

ER stress with low-dose Tunicamycin 24h prior to dsDNA exposure (Fig. S1F, G). In contrast,

induction of acute ER stress potently activated the cGAS/STING pathway (Fig. S1H, I). These

findings suggested that unresolved ER stress promoted exhaustion of the STING pathway with

impaired type I IFN production in response to pathogen sensing. Indeed, the STING agonist

DMXAA activated STING and TBK1 phosphorylation in iCtrl but to a lesser extent in iXbp1

IECs (Fig. 1D), which was accompanied by impaired *Cxcl10* and *Ifnb* induction (Fig. 1E, F).

Moreover, we validated the impact of chronic ER stress on epithelial STING signalling using

an independent transgenic mouse model, i.e. *nAtf6*^{IEC} VilCreERT2 mice carrying a Tamoxifen-

inducible intestinal epithelial *Atf6* overexpression (leading to constitutive ER stress(25,26)).

Tamoxifen-induced overexpression of *Atf6* led to epithelial STING suppression (Fig. S1J, K).

Collectively, our data demonstrated that chronic ER stress suppresses STING signalling in the

intestinal epithelium.

ER stress mediated STING exhaustion increases susceptibility to CMV infection in IEC`s

To test if ER stress-induced perturbation of the STING signalling links to susceptibility towards

infection, we infected chronically stressed i*Xbp1* Mode-K IECs (and controls) with murine

CMV (MCMV). We stimulated iCtrl and i*Xbp1* cells with xy copies of MCMV for xy h and

assessed STING-dependent type I IFN production. Indeed, iCtrl IECs responded to MCMV

infection with STING activation, type I IFN induction and subsequent viral containment when

compared to i*Xbp1* IECs, which failed to induce type I IFNs (Fig. 1G, Fig. S1L) and showed

increased CMV loads (Fig. 1H). To verify the impact of epithelial STING signalling on CMV

infection *in vivo*, we challenged *Xbp1*^{fl/fl} and *Xbp1*^{ΔIEC} mice with MCMV intraperitoneally and

assessed the degree of intestinal inflammation five days after the initial infection, as described

previously (27,28). In line with previous reports, the total viral load within the intestinal

mucosa was not detectable at day of sacrifice. However, we observed increased severity of

enteritis by a semiquantitative histology score in *Xbp1*^{ΔIEC}, but not in *Xbp1*^{fl/fl} mice (Fig. 1I, J)

5 days after CMV exposure. This observation confirmed that MCMV infection controlled the

inflammatory tone in genetically susceptible mice that exhibit chronic epithelial ER stress. To

test whether the observed STING exhaustion is relevant to other pathogen infections, we

further infected IECs with the intracellular pathogen *Listeria monocytogenes*. As described

before (29), *L. monocytogenes* infection resulted in strong type I IFN induction in the iCtrl

cells, which was barely detectable in i*Xbp1* IECs (Fig. S1M, N) . These data indicate that

chronic ER stress results in a functional impairment of STING signalling which leads to

inefficient pathogen clearance in the intestinal epithelium.

ER stress induces ROS-dependent metabolic adaptation to maintain STING signalling

We next aimed at identifying the underlying molecular processes that links chronic ER stress

with impaired STING signalling in IECs. Chronic ER stress has been associated with

fundamental changes of amino acid metabolism in the context of carcinogenesis (30). We

therefore tested the hypothesis that ER stress mediated alterations in amino acid metabolism

drives STING exhaustion under conditions of chronic ER stress. We performed an unbiased

metabolic profiling by assessing consumption and release (CORE) rates from the cell culture

medium, steady-state levels of intracellular metabolites and relative flux distribution through central carbon metabolism using stable isotope-assisted metabolomics (Fig. S2A). Intracellular analysis of metabolite steady-state levels showed significant increases of various metabolites. Most consistently, we observed a strong increase of intracellular metabolites in i*Xbp1*, that are required for glutathione synthesis, such as glycine, glutamate and cystine (Fig. 2A). Glutathione (GSH) is the most abundant intracellular antioxidant, and GSH synthesis is a two-step process in which the glutamate-cysteine ligase catalyzes the rate limiting step to form the dipeptide γ-glutamyl cysteine (GGC) from cysteine and glutamate (31,32). Intracellular cysteine availability can be increased via the cystine/glutamate antiporter *Slc7a11* and is enhanced in response to ROS formation (Fig. 2B) (33). To test whether an atltered metabolic state towards increased GSH synthesis is seen upon chronic ER stress we assessed CORE rates in iCtrl and i*Xbp1* cells and observed increased glutamine consumption and a striking switch from glutamate consumption towards glutamate release (Fig. 2C), pointing toward increased cystine/glutamate exchange in i*Xbp1* cells. We further validated the observation of increased glutamate release rates upon chronic ER stress in intestinal organoids stimulated with Tunicamycin (Fig. S2F). Although ER stress induced a strong ROS signature based on the amino acid exchange rates, central carbon metabolism remained largely unchanged in i*Xbp1* cells. Following a metabolic flux analysis with [U-^{13}C]glucose and [U-^{13}C]glutamine no striking differences have been observed, suggesting that cells primarily repond to ROS stress by altered amino acid exchange rates. (FigS2 B-E).

Based on this observation we tested the hypothesis that ER-stress induces a metabolic shift towards increased intracellular availability of amino acids needed for GSH synthesis. Hence, we repeated the metabolic profiling in response to the chemical ER stress inducer Tunicamycin (TM). Indeed, TM treatment induced a similar, dose-dependent glutamate release in both iCtrl and i*Xbp*1 cells (Fig. 2D). Cystine uptake rates increased upon TM treatment along with a

switch from glycine release towards glycine uptake (Fig. 2E, F). Most likely, the glycine switch

can be explained by an increased glycine demand for GSH synthesis. Generally, cells satisfy

their glycine demand from serine catabolism and show net excretion of glycine. However, in

context of ER stress the serine derived glycine does not seem to be sufficient for IECs which

causes a glycine switch from net excretion to net consumption (Fig. S2G). To further test the

hypothesis that unresolved ER stress induces a metabolic switch aiming to mount sufficient

GSH levels, we also observed increased expression of the glutamate/cystine antiporter *Slc7a11*

and increased intracellular GSH levels in response to ER stress (Fig. 2G). The primordial

function of GSH is to counterbalance free radicals derived from ROS by being oxidized to

glutathione disulphide (GSSG) (34,35) (Fig. S2H). We therefore assessed GSH, GSSG levels

and the ratio of GSH/GSSG. Despite increased steady state levels of GSH, GSSG and the ratio

of GSH over GSSG was reduced in *iXbp*1 cells compared to *iCtrl* cells suggesting an

unbalanced intracellular ROS homeostasis (Fig. 2H-J).

The neutralization of reactive oxygen species is conducted via the reduction of hydroperoxides

via oxidation of reduced GSH into its disulfide form GSSG by glutathione peroxidases (GPX).

We therefore tested the expression of glutathione peroxidase (GPX) 1-4 expression and

observed that i*Xbp1* cells showed a significant reduction of GPX 1-4, indicating that reduced

GPX expression accounts for decreased GSH/GSSG ratio in *iXbp1* cells that ultimately would

result in increased accumulation of ROS in chronic ER-stressed cells (Fig. S2I). We therefore

assessed whether ER-stress results in increased ROS generation in IECs, as postulated

previously for e.g. mouse embryonic fibroblasts (MEF) (36). Indeed, we noted increased ROS

levels at baseline in i*Xbp1* cells compared to i*Ctrl* cells which was further augmented in

response to TM treatment (Fig. 2J). As GSH is also utilized to counterbalance ROS-dependent

lipid peroxidation (LPO) (34,35,37,38), we further assessed LPO levels, which were also

significantly increased in i*Xbp1* cells and further increased upon acute TM treatment (Fig. 2K).

Altogether, our metabolic profiling revealed that i) ER-stress induces a metabolic shift towards

increased GSH synthesis to counterbalance ROS and that ii) chronic ER stress renders cells

into a state of ROS accumulation. Based on our data we hypothesize that this metabolic

adaptation process represents a bottleneck for IECs to maintain healthy ROS homeostasis while

excessive ROS levels might impair proper STING signalling in response to CMV infection.

Serine and glutathione metabolism is a critical gatekeeper of STING function in intestinal

epithelial cells

Serine has multiple important roles to support cellular functionality (39,40). Besides protein

biosynthesis, serine represents a building block for sphingolipid synthesis and its catabolism

provides essential building blocks for nucleotides synthesis. Consequently, serine/glycine

(S/G) starvation has gained much attention as a dietary anticancer intervention (41). Alternative

to anabolic needs, catabolism of serine via the mitochondrial folate pathway can also give rise

to net production of ATP (42). Additionally, serine catabolism also generates glycine which

represents an essential building block for GSH synthesis. Finally, serine derived one-carbon

units can be oxidized via the one-carbon cycle to generate NADPH, an important co-factor for

the reduction of oxidised glutathione. Hence, we predicted that upon ER-stress and

concomitant elevated ROS levels, serine metabolism becomes polarized to counterbalance

ROS levels by providing glycine and NADPH. To validate the hypothesis that serine and

glutathione metabolism is a critical gatekeeper in response to ER-stress aiming to maintain

STING function, we used the following approach: we exposed iCtrl/i*Xbp1* to serine-glycine

free (S/G⁻) cell culture medium to deprive GSH precursors. A decrease of intracellular glycine

and serine levels as well as an increase in GGC confirmed functional consequences of S/G

starvation. (Fig. S3A-C). In context of S/G deprivation the cell is able to induce a metabolic

compensatory mechanism of the serine *de novo* synthesis pathway, aiming to generate serine

from glucose via phosphoglyceratedehydrogenase (*Phgdh)* (Fig. 3A).

We therefore applied [U-^{13}C]-glucose tracing to monitor the relative glucose flux through the

serine *de novo* synthesis pathway (SSP) which can act as a compensatory mechanism upon

serine starvation. Illustrated by the M+3 serine isotopologue, serine starvation strongly

increased SSP (Fig. S3D) upon S/G starvation. Notably, this increased SSP activity was not

sufficient to fully compensate S/G deprivation with regard to serine, glycine or GGC levels

(Fig. S3A-C). Of note, at standard conditions, the SSP activity of i*Xbp1* cells was not increased

compared to iCtrl cells (Fig. S3D).

We next tested the direct impact of S/G starvation on ROS and LPO formation in iCtrl and

i*Xbp1* cells. As S/G starvation led to a compensatory activation of the SSP, which might

directly feed into GSH synthesis we combined S/G starvation with the inhibition of PHGDH,

the rate limiting enzyme of the SSP pathway, by using the specific and competitive inhibitor

BI-4916 (BI, Fig. 3A) (43,44). As expected, BI treatment blocked SSP and dual perturbation

with BI and S/G starvation further reduced intracellular serine and GSH levels, confirming the

efficacy of our treatment (Fig. 3B, C, Fig. S3E). Interestingly, GSH depletion was more

pronounced in i*Xbp1* cells compared to iCtrl cells (Fig. 3C).

In order to assess whether the metabolic shift towards GSH synthesis functionally led to

increased accumulation of ROS levels and LPO, we assessed ROS and LPO levels in i*Xbp1*

and iCtrl cells. We found that the deprivation of S/G w/wo the inhibition of serine *de novo*

synthesis led to increased accumulation of LPO and ROS, which was again increased in i*Xbp1*

cells (Fig. 3D, E).

Finally, to formally show that deprivation of the compensatory serine metabolism disrupts

epithelial STING signalling, we performed a single (S/G deprivation) or dual (S/G deprivation

+ *de novo* synthesis inhibition) perturbation of serine metabolism of iCtrl cells and stimulated

with the STING agonist DMXAA. While S/G starvation alone had no significant impact on

Cxcl10 expression of DMXAA treated iCtrl cells, dual perturbation (S/G + BI) resulted in a

strong and significant reduction of *Cxcl10* expression. (Fig. 3F) and decreased phosphorylation

of STING and TBK (Fig. 3G). Altogether our data show that serine-glycine metabolism serves

as a critical gatekeeper to facilitate STING/IFN-I signalling in the stressed intestinal epithelial

cells.

Serine-glycine metabolism is increased in the intestinal epithelium of IBD patients

To validate whether serine-glycine metabolism also provides a functional metabolic adaptation

process to facilitate IFN-I signalling in the inflamed mucosa of IBD patients, we used available

transcriptome datasets from UC patients. Doing so we observed that the expression of genes

involved in compensatory serine-glycine metabolism (*PHGDH, SLC7A11, ALDH1L2,*

MTHFD1L, MTHFD2, SHMT2) significantly increased with endoscopic and histologic

inflammation (Fig. S4A, B). Notably the induction of genes involved in the serine-glycine

metabolism also significantly correlated with the induction of interferon-inducible genes (ISG

score), indicating that indeed serine-glycine metabolism is essential in licensing mucosal IFN-

I production (Fig. S4C, D).

To further show that induction of serine-glycine metabolism corresponds with disease activity

in IBD we assessed the induction of genes involved in serine-glycine metabolism in clinical

responder and non-responder IBD patients and observed that targeted treatment with biologic

therapies leads to significant downregulation of serine-glycine metabolism in therapy

responders (Fig. S4E). We further assessed single-cell RNA sequencing data obtained from the

SCP259 scRNA-Seq IBD dataset (colon mucosa of 18 UC patients and 12 healthy individuals

(45)) to understand whether the observed association of serine metabolism and IFN-I induction

is specifically present in IECs. We confirmed that genes involved in ER-stress, serine

metabolism and ISG production are expressed in intestinal epithelial cells and clusters with

high serine score were attributed to secretory cell types, such as Paneth cells and goblet cells

(Supplementary Figure F,G).

Restoration of glutathione function restores STING signalling and CMV control in

intestinal epithelial cells

Having shown that serine metabolism orchestrates STING signalling in IECs and that this

metabolic process is also present in IECs of IBD patients, we tested the hypothesis that

pharmacological ROS inhibition restores epithelial STING signalling and thereby reinforces

CMV protection. We therefore treated iCtrl and i*Xbp1* cells with the antioxidant N-acetyl

cysteine (NAC). NAC treatment successfully reverted the observed glutamate switch in i*Xbp1*

cells (Fig. 4A) and lowered intracellular ROS and LPO levels (Fig. 4B, C). In addition, NAC-

treatment completely restored DMXAA-induced pSTING and pTBK1 levels in i*Xbp1* cells

(Fig. 4D), indicating that indeed ER-stress impairs STING signalling via ROS generation. As

we had shown that *Xbp1*-deficient cells show decreased GPX4 expression and increased ROS

formation, we further confirmed that GPX4 inhibition, using the GPX4 inhibitor RSL3, led to

a nearly complete abrogation of STING signalling, which was completely rescued in the

presence of NAC (Fig. 4E-G). Along the same line the impact of combined S/G starvation and

PHGDH inhibition on STING signalling was fully reverted in the presence of NAC (Fig. 4H,

I). Finally, we wondered whether restoration of STING signalling via NAC directly improves

CMV control in IECs. To this purpose we co-treated MCMV infected iCtrl and i*Xbp1* cells

with NAC. In line with our hypothesis i*Xbp1* cells showed a robust restoration of type I IFN

signalling (Fig. 4J, K) which further resulted in reconstituted suppression of MCMV viral load

(Fig. 4L).

Discussion:

Mucosal CMV infection is a leading cause of complicated disease behaviour in IBD. However, the underlying molecular pathophysiology driving increased susceptibility towards CMV colitis in IBD patients is poorly understood. Our data provide a novel molecular nexus on how chronic ER-stress paves the susceptibility to CMV infection and thereby generate a molecular inside into a well-established and highly relevant disease complication in IBD. We show that i) chronic ER-stress generates a molecular susceptibility to CMV infection via impairment of STING signalling. We further provide evidence that ii) the functional impairment of epithelial STING signalling depends on intracellular ROS generation. In addition, we show that iii) ER-stressed IECs aim to counterbalance ROS and reinstall STING signalling by funnelling the cellular metabolic flux towards serine-glycine dependent glutathione synthesis. Our data indicate that serine-glycine metabolism is essential in counterbalancing ER-stress induced ROS and licences STING signalling in the intestinal epithelium. We postulate that this mechanism has a profound impact on host immunity and might provide a layer of susceptibility towards pathogen infection in the inflamed mucosa of IBD patients.

In this context it needs to be noted that recent data indicate that the intestinal epithelium indeed presents the primary sensor of mucosal CMV infection, as IECs have been shown to be permissive to CMV replication and elicit a proinflammatory mucosal immune signal in response to CMV signalling (46). Over thirty years ago is has been described that HCMV infection increases the glycolytic flux of host cell and that glycolysis crucially impacts productive infection (47). Here, we now show that the IEC response indeed fuels the serine pool via the glycolytic serine *de novo* synthesis pathways in order to sustain GSH synthesis and maintain STING signalling. The functional outcome of increasing *de novo* serine synthesis appears to be cell-type specific, as it has been previously shown that abrogation of *de novo* serine synthesis via PHGDH exacerbates IFN-I signalling in macrophages via a mechanism

involving the YAP lysosomal degradation (48). Importantly, our data indicate that *de novo*

serine synthesis in IECs is primarily needed for glutathione synthesis and the reduction of

intracellular ROS, resulting in maintenance of STING signalling.

These findings might point to an important cell type-specific utilization of serine metabolism,

which is needed to contain CMV infections in the intestinal epithelium as the first line of

defence. Bearing in mind that the mechanical breakdown of the epithelial lining is a critical

hallmark of IBD, our data strongly put forward the idea of an ER-stress driven immune

exhaustion of the STING pathway in the intestinal epithelium as a critical layer of susceptibility

for CMV infection in IBD.

<u>Figure legends</u>

Figure 1: Chronic ER-stress suppresses STING signalling in intestinal epithelial cells

(A) *Xbp1*$^{fl/fl}$ (n= 13) and *Xbp1*$^{\Delta IEC}$ (n= 10) mouse small intestinal sections were stained for

STING, representative images. (B) Corresponding quantification of STING-positive cells per

crypt. (C) *Xbp1* fl/fl (n= 5) and *Xbp1*$^{\Delta IEC}$ (n= 5) mouse small intestinal crypts were isolated and

Cxcl10 expression was assessed by q-RT-PCR. (D) iCtrl and i*Xbp1* Mode-K cells were

stimulated with DMXAA (100 µg/ml, 1 h) and protein expression assessed by Western Blot.

(E) *Cxcl10* and (F) *Ifnb* gene expression of Mode-K iCtrl and i*Xbp1* cells stimulated with

DMXAA (100 µg/ml, 3 h, n=3). (G) iCtrl and i*Xbp1* Mode-K cells were infected with MCMV

at a MOI of 10 and *Cxcl10* gene expression and (H) MCMV fold induction was assessed by

PCR (n= 4). (J) *Xbp1* fl/fl and *Xbp1*$^{\Delta IEC}$ mice were infected intraperitoneally with vehicle (n=11

for each group) or MCMV (n=12 (*Xbp1* fl/fl); n=14 (*Xbp1*$^{\Delta IEC}$)) and intestinal inflammation

was assessed. (K) representative images of HE stained small intestinal sections. Depicted data

representative of minimum of three independent experiments.

Figure 2: ER-stress induces a distinct ROS-dependent metabolic adaptation to sustain STING signalling

(A) In vivo uptake of amino acids after 24h. Absolute consumption and release rates (CORE) in fmol/PCV/h are displayed (n ≥ 3). (B) Relative intracellular metabolite level of iCtrl and i*Xbp1* cells (n ≥ 3). Peak areas are normalized to PCV and internal standards. (C) Schematic depicting of the GSH synthesis pathway. (GCL = glutamate-cysteine ligase; GS = glutathione synthetase). (D-F) iCtrl and i*Xbp1* cells were treated with TM and CORE of (D) glutamate, (E) cystine and (F) glycine was determined via GC-MS (n = 4, Two-way ANOVA). (G) Relative *Slc7a11* mRNA expression of iCtrl and i*Xbp1* cells treated with (+) or without (-) TM (n = 3, paired t-test). (H) Relative intracellular GSH level of iCtrl and i*Xbp1* cells (n = 9, paired t-test). (I) GSH/GSSG ratios of iCtrl and i*Xbp1* cells (n =3, paired t-test). (J) Relative intracellular ROS levels of iCtrl and i*Xbp1* cells treated with TM using DCF-DA (n ≥ 3, MFI ± SEM). (K) Relative percentage of FITC+ cells in iCtrl and i*Xbp1* cells treated with 0.1 µg/ml TM for 24h assessed by FACS (n =3). Depicted data representative of minimum of three independent experiments.

Figure 3: Serine and glutathione metabolism are critical gatekeepers of STING function in intestinal epithelial cells

(A) Schematic illustration of the different intervention strategies used to interfere with GSH synthesis and GPX4 activity. (B) Relative intracellular serine level of iCtrl and i*Xbp1* cells in medium lacking serine and glycine (-SG) in the presence (+) or absence (-) of 15 µM BI-4916 (n = 3) (C) Relative GSH level of iCtrl and i*Xbp1*cells in medium lacking serine and glycine (-SG) treated with (+) or without (-) of 15 µM BI-4916 (n = 2). (D) Relative intracellular ROS level of iCtrl and i*Xbp1* cells in medium lacking serine and glycine (-SG) treated with (+) or without (-) of 15 µM BI-4916 (n ≥3. (E) Relative percentage of FITC+ cells in iCtrl and i*Xbp1*

cells treated with 0.1 µg/ml TM (n ≥ 3). (F) Mode-K iCtrl and i*Xbp1* cells were starved of

serine/glycine and treated with 15 µM BI for 24 h before stimulation with DMXAA (100 µg/ml,

3 h) and *Cxcl10* gene expression was analyzed by q-RT-PCR (n=3). (G) Western Blot of both

Mode-K iCtrl and i*Xbp1* cells stimulated with DMXAA (100 µg/ml, 1 h) after 24 h of SG

starvation and treatment with BI-4916 (15 µM). Depicted data representative of minimum of

three independent experiments.

Figure 4: Restoration of glutathione function restores STING signalling and CMV

control in intestinal epithelial cells

(A) Absolute consumption and release rates of glutamate in iCtrl and i*Xbp1* cells after NAC

treatment for 24h (n ≥ 3). (B) Relative intracellular ROS level of iCtrl and i*Xbp1* cells after

NAC treatment for 24 h (n ≥4, paired t-test). (C) Relative percentage of FITC+ cells in iCtrl

and i*Xbp1* cells after NAC treatment for 24h (n = 3, paired t-test). (D) Western Blot of both

Mode-K iCtrl and i*Xbp1* cells stimulated with DMXAA (100 µg/ml, 1 h) after 24 h of NAC

pretreatment (10 mM). (E) Western Blot of Mode-K iCtrl cells stimulated with DMXAA (100

µg/ml, 1 h) after 24 h pretreatment with NAC (10 mM) and RSL3 (0.5 µM). (F) q-RT-PCR of

Cxcl10 (G) and *Ifnb1* expression from Mode-K iCtrl cells stimulated with DMXAA (100

µg/ml, 3 h) after 24 h of SG starvation and co-treatment with BI-4916 (15 µM) and NAC (10

mM) (n=3). (H) Western Blot of Mode-K iCtrl cells stimulated with DMXAA (100 µg/ml, 1

h) after 24 h of SG starvation and treatment with NAC (10 mM) and BI-4916 (15 µM). (I) q-

RT-PCR of *Cxcl10* expression from Mode-K iCtrl cells stimulated with DMXAA (100 µg/mL,

3 h) after 24 h of SG starvation and treatment with BI-4916 (15 µM) and NAC (10 mM) (n=4).

(J) Mode-K iCtrl and i*Xbp1* cells were stimulated with NAC (10mM) and infected MCMV. q-

RT-PCR of *Ifnb*, *Cxcl10* (K) and viral load as measured by viral DNA copy number (L) (n=4).

Depicted data representative of minimum of three independent experiments.

Supplementary Figure 1: Chronic ER-stress abrogates intestinal epithelial STING signalling

(A) *Xbp1* fl/fl (n=2), STING -/- (n=1) and *Xbp1*$^{\Delta IEC}$ (n=2) mouse small intestinal crypts were isolated and assessed for protein expression by Western Blot. (B) *Tmem173* and (C) *Ifnb1* gene expression of *Xbp1* fl/fl (n=5) and *Xbp1*$^{\Delta IEC}$ (n=5) as assessed by q-RT-PCR. (D) iCtrl and i*Xbp1* Mode-K cells were stimulated with dsDNA (1 µg/ml) for the indicated time and protein concentration was assessed by Western Blot. (E) *Cxcl10* gene expression of the same experiment as assessed by q-RT-PCR. (n=3). (F) Mode-K cells were subjected to either vehicle or low-dose Tunicamycin (0,1 µg/ml) pretreatment, followed by dsDNA (1 µg/ml) stimulation. Protein expression was assessed by Western Blot. (G) *Cxcl10* gene expression as assessed by q-RT-PCR of the same experiment (n=4). (H) Mode-K iCtrl cells were subjected to *Tmem173* knockdown by siRNA and subsequently stimulated with Tunicamycin (1 µg/ml) for the indicated time-course and protein expression was analysed by Western blot. (I) Corresponding gene expression analysis by q-RT-PCR (n=4). (J) *nATF6* VilCreERT2 tg or fl mice carrying a Tamoxifen-inducible intestinal epithelial overexpression of ATF6, resulting in ER-stress induction, were treated with Tamoxifen for 4d prior to sacrifice and STING expression was assessed by immunohistochemistry. Representative HE/STING-stained small intestinal sections. (K) Corresponding quantification of STING-positive cells per crypt (n=4 (fl) and n=6 (tg)). (L) *Ifnb* gene expression corresponding to Fig. 1G, H (n=4). (M) iCtrl or i*Xbp1* Mode-K cells were exposed to *Listeria monocytogenes* at a MOI of 50 for 1 h before medium was exchanged and Gentamicin (50 µg/ml) was added for 1 h, followed by Gentamicin at 5 µg/ml for 22 h. *Cxcl10* and (N) *Ifnb* gene expression was assessed by q-RT-PCR (n=4). Depicted data representative of three independent experiments.

Supplementary Figure 2: Metabolic adaptations in response to ER-stress to maintain

redox balance

(A) Schematic overview of the experimental workflow used to analyze metabolites from

medium and cell extracts. (B) Schematic illustration of the mass isotopomer distribution (MID)

of the TCA cycle from glucose. Filled red circle represent ^{13}C atoms derived from [U-^{13}C]

glucose. (C) Relative MIDs of the indicated TCA cycle intermediates in iCtrl or i*Xbp1* cells

cultured in [U-^{13}C] glucose medium for 24h. (D) Schematic illustration of the mass isotopomer

distribution (MID) of the TCA cycle from glutamine. Filled blue circle represent ^{13}C atoms

derived from [U-^{13}C] glutamine. (E) Relative MIDs of the indicated TCA cycle intermediates

in iCtrl or i*Xbp1* cells cultured in [U-^{13}C] glutamine medium for 24h. (F) Absolute consumption

and release rates of the indicated amino acids in organoids cultures derived from Xbp1 fl/fl

mouse small intestinal crypts after 24h. Representative experiment is shown. (G) Consumption

and release rates of serine in iCtrl and i*Xbp1* cells treated with TM for 24h (n ≥ 4, Two-way

ANOVA). (H) Simplified illustration of the cellular GSH cycle (GR = glutathione reductase,

GPx = glutathione peroxidase genes). (I) Relative mRNA expression of the indicated GPx

genes in iCtrl and i*Xbp1* cells (n = 4, paired t-test). Depicted data representative of three

independent experiments.

Supplementary Figure 3: Epithelial STING function relies on serine and gluthathione

metabolism

(A-C) Relative intracellular metabolite level of (A) glycine, (B) serine and (C) γ-glutamyl-

cysteine (GGC) in iCtrl and i*Xbp1* cells cultured in complete or serine/glycine-free (-S/G)

medium for 24h (n = 3). (D) Fractional enrichment of serine M+3 in iCtrl and i*Xbp1* cells

cultured in complete or serine/glycine-free (-S/G) medium for 24h (n =6). (E) Fractional

enrichment of serine M+3 in iCtrl and i*Xbp1* cells cultured in complete or serine/glycine-free

(-S/G) medium in the presence (+) and absence (-) of 15 mM BI-4916 for 24h (n =3, two-way

ANOVA). Depicted data representative of three independent experiments.

Supplementary Figure 4: Serine-glycine metabolism is increased in the intestinal

epithelium of IBD patients

(A, B) ISG score and expression of serine pathway related genes in UC patients in GSE73661

(A) and GSE109142 (B) at timepoint 0. Red dots and connecting lines show the median

expression. (C, D) Correlation of serine pathway related genes with the ISG score in GSE73661

(C) and GSE109142 (D) in all available samples. The Pearson's correlation coefficients *r* and

p-values are shown. (E) ISG score and serine pathway related genes expression in GSE73661

patients according to therapy response (the latest available timepoint was analysed for each

patient). (F) Heatmap of ISG or Serine scores and *ATF4* expression in the main cell-types of

the SCP259 scRNA-Seq IBD dataset (colon mucosa of 18 UC patients and 12 healthy

individuals (45)). Colour shows z-scores and the dot size the percent of expressing cells. (G)

Heatmap of ISG or Serine scores and *ATF4* expression in selected epithelial cell clusters of

SCP259 in healthy (H), NI (non-inflamed) and inflamed (I) tissue biopsies. *$p<0.05$, **$p<0.01$,

***$p<0.001$.

Methods

Resource availability

Lead contact

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the lead contacts, Konrad Aden, MD (k.aden@ikmb.uni-kiel.de), Philip Rosenstiel,

MD (p.rosenstiel@mucosa.de) and Johannes Meiser, PhD (Johannes.Meiser@lih.lu).

Materials availability

This study did not generate new unique reagents.

Data and code availability

All data reported in this paper will be shared by the lead contact upon request.

This paper does not report original code.

Any additional information required to reanalyze the data reported in this paper is available

from the lead contact upon request.

Experimental model and subject details

Mice

Xbp1 fl/fl mice were generated by targeted insertion of LoxP sites flanking the exon 2 of the

Xbp1 gene. VillinCre+/- mice (strain: B6.SJL-Tg (Vil-cre)997Gum/J) were purchased from

Jackson Laboratory. Conditional knockout of *Xbp1* in the intestinal epithelium was then

established by crossing VillinCre+/- mice with *Xbp1* fl/fl mice, resulting in specific deletion of

Xbp1 (*Xbp1*$^{\Delta IEC}$) in the intestinal epithelium (49). Tamoxifen-inducible nATF6-HA

overexpressing Villin-Cre n*Atf6*^{IEC} mice (n*Atf6* VilCreERT2 transgene (tg)) mice were

generated as described previously (26). Overexpression of nATF6-HA was induced using

Tamoxifen for four days before sacrifice. *Tmem173* -/- mice were purchased from Jackson

Laboratory (previously described by Jin et al., 2011, strain #025805) (50).

Mice were housed under specific pathogen - free (SPF) conditions in individually ventilated

cages (IVCs). Animals were held in a 12 h light-dark cycle at 21 °C ± 2 °C and 60 % ± 5 %

humidity. Food and water were available ad libitum. All animal experiments were approved by

federal authorities and performed in accordance to institutional guidelines.

MCMV infection *in vivo*

For *in vivo* MCMV infection 8 week old *Xbp1* fl/fl and *Xbp1*$^{\Delta IEC}$ mice were infected

intraperitoneally with either 10^5 PFU MCMV or with vehicle. During the infection mice were

fed a standard chow diet and had free access to food and water. Five days after infection mice

were sacrificed and the intestines were harvested.

Cell culture

In this study, the SV40 large T-antigen-immortalized murine small intestinal epithelial cell line

MODE-K (51) was employed as an in vitro model for the intestinal epithelium. MODE-K cells

harbouring an RNA interference (RNAi) based knockdown of *Xbp1* and a control RNA

interference (i*Xbp1*, iCtrl) were kindly gifted by Artur Kaser, University of Cambridge (6). In

brief, RNA interference was achieved by stable introduction of a small hairpin RNA specific

for *Xbp1* and a control small hairpin RNA identical to Lee et al. (52), except that SFGΔU3hygro

was used. Knockdown of *Xbp1* was confirmed by qPCR. The following chemical stimulants

were used (name, company, catalogue-number): DMXAA/Vadimezan (Hölzel, HY-10964), N

Acetyl-L-cystein (Th. Geyer/Sigma, A7250-25G), BI-4916 (Hoelzel, HY-126253-5mg), RSL3

(Selleck Chemicals, S8155). siRNA knockdown of *Tmem173* was performed using the

Viromer® Blue kit (Lipocalyx, Halle, Germany) according to the manufacturer's protocol.

Tmem173 siRNA was purchased from Qiagen (catalogue-number GS72512).

Organoid generation and handling

Mouse small intestinal organoids were cultivated as described previously (53,54). To generate

sustainable cultures of small intestinal organoids, mice were sacrificed and the small intestine

was removed and cut into pieces. PBS wit 10 nM EDTA was added to the pieces for 10 min

and the specimen was placed under mild agitation. Afterwards, the supernatant was removed,

fresh PBS/EDTA solution was added and incubated under mild agitation for 10 min again. This

was repeated four times. To remove debris, the suspension was filtered through a 100 µm

strainer and centrifuged at 1200 rpm for 10 min. Crypts were then either used for protein

respectively RNA isolation or resuspended in Matrigel and embedded in 24-well plates to

generate organoid cultures. Medium was changed twice per week. Organoids were stimulated

after 5 to 7 days of cultivation.

Method details

<u>Histology</u>

After sacrifice, the small intestine was removed and immediately fixed in 10 % (w/v) formalin

for 24 h at 4 °C. The tissue was then immersed in a series of ethanol solutions of increasing

concentrations until 100 %. Next, the ethanol was gradually replaced with xylene, which is

then replaced by paraffin. Paraffin embedded tissue was cut into 3.5-4.5 µm thin sections using

the RM2255 microtome. Paraffin embedded tissue sections were stained for 2-5 min in

haematoxylin. The cytoplasm was counterstained with 1 % (v/v) eosin solution for 2 min.

Stained slides were then dehydrated and embedded in Roti-Histo-Kit mounting medium and

examined with a Zeiss AxioImager.Z1 apotome microscope and the AxioVision Rel 4.9

software.

<u>STING immunohistochemistry and quantification</u>

To quantify STING expression in the intestinal epithelium, 3,3'-Diaminobenizidine staining

was performed. Slides were rehydrated and boiled for 20 min in citrate buffer (pH 6.0) for

antigen retrieval. Unspecific binding sites were blocked by incubating slides with 5 % (v/v)

goat serum in phosphate-buffered saline (PBS). Naturally occurring peroxidases were

inactivated by applying 3 % hydrogen peroxide. Antibody binding and 3,3'- diaminobenzidine

staining was performed according to the manufacturer's instruction (Vectastain Elite ABC Kit,

Vector Labs, Peterborough, United Kingdom). In brief, after primary antibody binding, the

signal was amplified by adding a horseradish peroxidase-conjugated streptavidin antibody.

Quantification was performed by counting average STING positive cells per crypt. For each

animal, 50 crypts were counted and averaged.

<u>Western blot</u>

Cells and tissues were lysed using 40-100 µl SDS-based DLB buffer + 1 % Halt™ Protease

inhibitor cocktail (ThermoFisher Scientific, Waltham, USA) before incubation for 5 minutes

at 95 °C followed by sonification for 2x 5 seconds. To remove cell remnants, lysates were

centrifuged at 16.000 x g for 15 minutes at 4°C and supernatants were transferred into a new

tube. For protein extraction of organoids, Matrigel™ was removed by incubation of organoids

in Corning™ Cell Recovery Solution (Corning, New York, USA) for 45-60 min at ice after

transferring organoids to tubes, centrifugation at 3000 g for 2 min and discarding supernatants.

Next, the lysis buffer as described above was added. The protein concentration was determined

applying the copper-based DC Protein Assay (Bio-Rad, Munich, Germany) according to the

manufacturer´s protocol. Proteins were separated by their molecular weight using precast

gradient polyacrylamide gels for improved separation of a wide range of proteins (NuPAGE®

4-12 % Bis-Tris gels, Life Technologies, Darmstadt, Germany). Protein lysates were prepared

by supplementation with the corresponding NuPAGE LDS Sample Buffer (Thermo Scientific,

Bremen, Germany), followed by incubation at 70 °C for 70 min. SDS-PAGE electrophoresis

was run at 160 V and a maximum current of 300 mA for 1 to 2 h in the XCell SureLock Mini

Cell system (Life Technologies, Darmstadt, Germany). NuPAGE MOPS SDS running buffer

was used to run the gels and Prestained PageRuler plus protein ladder 10-250 K (Thermo

Scientific, Bremen, Germany) served to estimate protein size. To detect proteins using

immunodetection by specific antibodies, separated proteins were transferred onto

polyvinylidene difluoride (PVDF) membranes. PVDF membranes (Bio-Rad, Munich,

Germany) were activated under mild agitation using methanol for 10 s, followed by 5 min of

washing in A. bidest. and then transferred for 5 min to anode buffer 1. Separated proteins were

transferred from the gel onto the membrane by semi-dry blotting using the Trans-Blot Turbo™

Transfer System (Bio-Rad, Munich, Germany) with a discontinuous buffer system consisting

of one cathode buffer and two anode buffers. Proteins were blotted onto the membrane at 0,3

A, 25 V for 60 min. After the transfer membranes were blocked with 5 % (w/v) blotting grade

blocker (non-fat dry milk) in Tris - buffered saline (TBS) supplemented with 0.1 % (v/v) Tween

20 (TBS - T) for 1 h. Membranes were probed with the appropriate primary and secondary

antibody. The primary antibody was applied in 5 % (w/v) blotting grade blocker (non-fat dry

milk) or bovine serum albumin (BSA) in Tris-buffered saline (TBS) supplemented with 0.1 %

(v/v) Tween 20 according to the manufacturer's instruction and incubated overnight at 4 °C

under mild shaking. Next, the membrane was washed three times with TBS - T for 15 min to

remove excess antibody. A horseradish peroxidase (HRP) conjugated secondary antibody was

diluted in 5 % (w/v) blotting grade blocker (non-fat dry milk) in TBS-T according to the

manufacturer's recommendation and incubated for 1 h at room temperature. Binding of the

secondary antibody was detected using a chemiluminescent substrate kit (Thermo Scientific)

and recorded with an automated developer machine (Agfa, Mortsel, Belgium). The following

antibodies were used: Actin β (Sigma-Aldrich, Munich, Germany, A-5441), β-Tubulin (MBL

International, Woburn, USA, M150-3), phospho-STING (Cell Signalling Technology, Leiden,

The Netherlands, 72971), phospho-TBK1 (Cell Signalling Technology, Leiden, The

Netherlands, 5483), STING (Cell Signalling Technology, Leiden, The Netherlands, 13647),

TBK1 (Cell Signalling Technology, Leiden, The Netherlands, 38066), mouse-HRP

(Amersham Biosciences, Glattbrugg, Switzerland, NA931V), rabbit-HRP (Amersham

Biosciences, Glattbrugg, Switzerland, NA934V).

q-RT-PCR

RNA extraction was performed using the RNeasy Mini Kit (Qiagen, Hilden, Germany)

according to the manufacturer's instruction. To generate cDNA, 100-1000 ng total RNA

(depending on the amount of previously extraced RNA) was transcribed into complementary

DNA (cDNA) by reverse transcription using the Maxima H Minus First Strand cDNA

Synthesis Kit (Thermo Scientific, Bremen, Germany). Reverse transcription was performed

following the manufacturer´s instruction. Quantitative real-time PCRs (q-RT-PCR) was used

to amplify and simultaneously quantify mRNA levels using target-specific oligonucleotides.

Predesigned TaqMan probes (Applied Biosystems, Carlsbad, USA were purchased and used.

The 7900HT Fast Real Time PCR System (Applied Biosystems, Darmstadt, Germany) was

used for qPCR experiments. Samples were run in duplicates on 384-well plates. For the PCR

reaction, 5-10 ng cDNA and 0,5 μl of the respective TaqMan gene expression assays were used.

The PCR program was carried out following the manufacturer's recommendation (TaqMan

Gene Expression Master Mix protocol, Applied Biosystems, Darmstadt, Germany).

Alternatively, RT-qPCR was performed from 10 ng cDNA per sample using Fast SYBR™

Green Master Mix (Lifetech) at 95°C for 20 s, 40 cycles of 95°C for 1 s and 60°C for 20 s on

the QuantStudio 5 Real-Time PCR System (ThermoFisher). Relative quantification of each

gene was done by QuantStudio Design&Analysis v1.5.1 software (ThermoFisher) and

comparative Ct method. All samples were analysed in technical duplicates and repeat as

independent, biological experiments (n $\geq$ 3). Moreover, all expression data were normalized to

housekeeping genes (*Tbp*, *Hrpt1* or *Actinb*). Taqman assay probes were purchased from Life

Technologies, Darmstadt, Germany (*Cxcl10*, 04331182; *Gapdh*, 99999915; Ifnb1, 04331182;

Tmem173, 04331182; *Xbp1*, 03464496). All non-TaqMan assay mouse primers used in this

study can be found in the table below:

Gene name	Forward primer sequence ((5' – 3')	Reverse primer sequence(5' – 3')

Slc7a11	GCTCGTAATACGCCCTGGAG	GGAAAATCTGGATCCGGGCA
Gpx1	ACCGTGTATGCCTTCTCCGC	CTCGGTGTAGTCCCGGATCG
Gpx2	AGTTCGGACATCAGGAGAACTG	GATGCTCGTTCTGCCCATTG
Gpx3	ACAATTGTCCCAGTGTGTGCAT	TGGACCATCCCTGGGTTTC
Gpx4	CATTGGTCGGCTGCGTGA	CGCGGGATGCACACATGG
Tbp	GTCATTTTCTCCGCAGTGCC	GCTGTTGTTCTGGTCCATGAT
Hrpt1	TGGATACAGGCCAGACTTTGTT	CAGATTCAACTTGCGCTCATC

Listeria monocytogenes infection

Listeria monocytogenes serotype 1/2a strain EGD was used to test type I IFN induction in MODE-K cells. Bacteria were plated out from glycerol stocks onto brain-heart infusion (BHI; BD Biosciences, Franklin Lakes, USA) agar plates overnight. *Listeria* colonies were then transferred into 1 ml BHI medium and incubated overnight at 37 °C to allow for bacterial replication. Next, bacteria were diluted and reincubated at 37 °C to induce a log - phase growth until OD600 = 0.6 was reached. After washing with PBS, bacteria were diluted in MODE-K medium and added to the cells according to multiciply of infection (MOI). As described previously (55), MODE-K cells were infected at a MOI of 50. After allowing bacterial invasion for 1 h, the bacteria-containing medium was removed, cells were washed once with PBS and then treated with gentamicin (50 µg/ml) for 1 h to kill extracellular bacteria. The gentamicin-containing medium was then replaced with medium containing a lower concentration of Gentamicin (5 µg/ml), and cells were lysed for protein and RNA harvesting after another 22 h.

MCMV infection and quantification of replication

MCMV was prepared as described previously (56) and infection of MODE-K cells was performed at a MOI of 10. Fold induction of MCMV was quantified by RT-PCR using the MCMV M54 gene sequence: ATCATCCGTTGCATCTCGTTG (forward), CGCCATCTGTATCCGTCCAT (reverse).

<u>Histology score CMV colitis</u>

After staining, an expert pathologist assessed inflammation with a semi quantitative scoring

system as previously reported (5). To assess the inflammation five subscores are summed up

for every sample and multiplied by the extent of the affected area (1, <10%; 2, 10%-25%; 3,

25%-50%; 4, >50% of the intestine affected). As subscores the following parameters are used:

mononuclear infiltrate (0: absent; 1: mild, 2: moderate 3: severe), crypt hyperplasia, epithelial

injury/erosion, polymorphonuclear infiltrate and transmural inflammation.

<u>Flow Cytometric Analysis of ROS Levels</u>

250,000 iCtrl or i*Xbp1* Mode-K cells were seeded in six-well plates and treated the subsequent

day as indicated in the figure legends. After an incubation time of 24h, cells were stained with

Zombie NIR (BioLegend, 1:1000 diluted in PBS) for 20 min at 37°C (to discriminate live and

dead cells in the later analysis), detached with trypsin, centrifuged at 350 g for 5 min and

washed with warm DMEM. Then, cells were stained in 100 μl DMEM supplemented with

10 μM DCF-DA (PromoCell) for 30 min at 37°C. After the incubation step, cells were

centrifuged at 350 g for 5 min, washed twice with ice-cold PBS and resuspended in ice-cold

FACS buffer (PBS with 10 % FBS and 0.5 mM EDTA). Samples were measured using the BD

FACSCanto™ system and BD FACSDiva software. Data analysis was performed using

FlowJo v10.6.2 software. All measurements were performed as independent, biological

experiments (n ≥ 3) indicated in the respective figure legend.

<u>Flow Cytometric Analysis of LPO Levels</u>

Image-iT™ Lipid Peroxidation Kit (ThermoFisher) was used according to manufacturer`s

instructions. In brief, 250,000 iCtrl or i*Xbp1* Mode-K cells were cultivated in 2 ml medium and

treated after 24h as stated in the figure legends. Following 24h incubation, cells were harvest

via trypsinization, centrifuged at 350 g for 5 min and washed with warm DMEM. Then, cells

were stained in 100 µl DMEM supplemented with 10 µM Lipid peroxidation sensor

(BODIPY® 581/591 C11) for 30 min at 37°C. After two washing steps with ice-cold PBS at

350 g for 5 min, cells were resuspended in ice-cold 400 µl FACS buffer and measured using

NovoCyte Quanteon system (Aligent). Data analysis was performed using FlowJo v10.6.2

software. All measurements were performed as independent, biological experiments (n ≥ 3).

<u>Determination of GSH/GSSG ratio</u>

GSH/GSSG ratios were obtained via a GSH/GSSG-GLO^{TM} assay kit (Promega, V6611)

following the manufacturer`s instructions. For the experiments, $4x10^4$ iCtrl or iXbp1 cells were

plated out in opaque, white 96 well plates (Greiner, CELLSTAR, 655083), incubated for 24h

in DMEM (10% FBS) and processed according to the manufacture`s protocol. GSH/GSSG

ratios were determine by the following equation: total GSH-(GSSGx2)/GSSG. All

measurements were performed as technical replicates (N = 3) and repeated as independent,

biological experiments (n = 3) indicated in the respective figure legend.

<u>Stable isotope tracing and metabolite extraction</u>

Stable isotope tracing experiments were performed as previously described in Kiweler *et al.*,

2020 (57). Briefly, [U-^{13}C]-glucose or [U-^{13}C]-glutamine (Cambridge Isotope Laboratories,

CLM-1396 and CLM-1822) tracing was performed in DMEM supplemented with 10 % FBS

and the respective tracers (2 mM glutamine or 17 mM glucose). $2x10^5$ cells were seeded in 12-

well plates in triplicates for each experimental condition. Furthermore, an identical set of

triplicate wells were plate-out to calculate the packed cell volume (PCV) by determining cell

count and cell volume at the start and end of each tracing experiment (42). After 24h, growth

medium was replaced by ^{13}C tracer medium and cells were cultured for an additional 24 h at

37°C. In parallel, starting PCV was determined by counting three wells per condition. To assess

PCV at the end of the experiment, one set of triplicate wells was counted at the experimental

endpoint, whereas another triplicates set was used for subsequent metabolite extraction. In

addition, medium was collected, centrifuged at 350 g for 10 min at 4°C and supernatants were

stored at -80°C until medium quantification via GC-MS or YSI.

For GC-MS analysis, cells from [U-^{13}C]-glucose and [U-^{13}C]-glutamine tracer experiments

were washed once with 0.9 % ice-cold NaCl. Thereafter, metabolites were extracted by adding

400 μl ice-cold MeOH/mqH$_2$O (ratio, 1:1) to each well. In addition, the extraction solvent

contains the internal standards pentanedioc-d6 acid and [U-^{13}C]-ribitol at a final concentration

of 1 μg/ml as well as Tridecanoid-d25 acid at a final concentration of 5 μg/ml. After the plates

were incubated for 5 min at 4°C on a rocking shaker, supernatant was transferred to 1.5-ml

Eppendorf tubes, mixed with 200 μl CHCl$_3$, and centrifuged for 5 min at 13,000 g and at 4°C.

Finally, the upper polar phase was collected, fully dried via SpeedVac and stored at -20°C for

subsequent GC-MS analysis.

For LC-MS analysis, metabolites extraction was performed as previously described by Meiser

et al., 2016 (42) and extracted metabolites were stored at -80°C until LC-MS analysis.

<u>GC-MS: Determination of MIDs of intracellular metabolites following stable isotope tracing</u>

Polar metabolites were derivatized for 90 min at 55 °C with 20 μl of methoxyamine (c = 20

mg/ml) in pyridine under continuous shaking and subsequently for 60 min at 55 °C with 20 μl

of MTBSTFA w/ 1% TBDMCS. GC-MS analysis was performed using an Agilent 7890B GC

coupled to an Agilent 5977A Mass Selective Detector (Agilent Technologies). A sample

volume of 1 μl was injected into a Split/Splitless inlet, operating in splitless mode at 270 °C.

Gas chromatograph was equipped with a 30 m (I.D. 250 μm, film 0.25 μm) ZB-35MS capillary

column with 5 m guard column (Phenomenex). Helium was used as carrier gas with a constant

flow rate of 1.2 ml/min. GC oven temperature was held at 100 °C for 2 min and increased to

300 °C at 10 °C/min and held for 4 min. Total run time was 26 min. Transfer line temperature

was set to 280 °C. Mass selective detector (MSD) was operating under electron ionization at

70 eV. MS source was held at 230 °C and the quadrupole at 150 °C. For precise quantification

of the MID, measurements were performed in selected ion monitoring mode. Target ions (m/z)

and Dwell times are shown in Table 2.

The MetaboliteDetector software package (v. 3.220180913) was used for mass spectrometric

data post processing, quantification, MID calculations, correction of natural isotope abundance,

and determinations of fractional carbon contributions (58).

Table 2: Target ions and Dwell times of metabolites quantified in this study.

Derivative (m/z of selected fragment)	Quant-Ions (m/z)	Dwell Time (ms/ for each)	Sum Formula (fragment)	MID size
Pyruvic Acid 1MeOX 1TBDMS (174)	174.0 – 180.1	15	C6H12O3NSi	4
Lactic acid 2TBDMS (261)	261.1 – 267.1	15	C11H25O3Si2	4
Alanine 2TBDMS (260)	260.1 – 266.1	15	C11H26NO2Si2	4
Glycine 2TBDMS (246)	246.1 – 252.1	15	C10H24NO2Si2	3
Valine 2TBDMS (288)	288.2 – 295.2	15	C13H30NO2Si2	6
Leucine 2TBDMS (302)	302.2 – 310.2	15	C14H32NO2Si2	7
Isoleucine 2TBDMS (302)	302.2 – 310.2	15	C14H32NO2Si2	7
Proline 2TBDMS (286)	286.2 – 294.2	15	C13H28NO2Si2	6
Succinic Acid 2TBDMS (289)	289.1 – 296.1	15	C12H25O4Si2	5
Fumaric Acid 2TBDMS (287)	287.1 – 294.1	15	C12H23O4Si2	5
(Internal Standard) Pentanedioic-D6 Acid	235.2, 309.2, 351.3	30		
Serine 3TBDMS (390)	390.2 – 396.2	15	C17H40NO3Si3	4
Methionine 2TBDMS (320)	320.2 – 328.2	10	C13H30NO2SSi2	6
a-Ketoglutaric Acid 1MeOX 2TBDMS (346)	346.2 – 354.2	10	C14H28NO5Si2	6
Malic Acid 2TBDMS (419)	419.2 – 426.2	15	C18H39O5Si3	5
Aspartic Acid 3TBDMS (418)	418.2 – 425.2	15	C18H40NO4Si3	5
Glutamic Acid 3TBDMS (432)	432.3 – 440.3	10	C19H42NO4Si3	6
Lysine 3TBDMS (431)	431.3 – 440.3	15	C20H47N2O2Si3	7
Glutamine 3TBDMS (431)	431.3 – 439.3	10	C19H43N2O3Si3	6
Citric Acid 4TBDMS (591)	591.3 – 600.3	10	C26H55O7Si4	7

3-Phosphoglyceric acid 4TBDMS (585)	585.4 – 592.4	15	C27H63O7PSi4	4
Histidine 3TBDMS (440)	440.2 – 449.2	15	C20H42N3O2Si3	7
Tryptophan 2TBDMS (375)	375.2 – 389.2	10	C19H31N2O2Si2	12

Analysis of Medium Exchange Rates

Medium quantification and exchange rate calculation was performed as described in Meiser et

al., 2016 (42). For metabolite extraction, 20 µl of the collected medium sample was mixed with

180 µl extraction solvent (80 % MeOH/20 % mqH$_2$O) containing the internal standards [U-

13C]-ribitol (50 µg/ml) and pentanedioc-d6 acid (20 µg/ml), vortexed for 10 s and

subsequently incubated for 15 min at 4°C and 1350 rpm in a thermomixer. After centrifugation

for 5 min at max speed, 50 µl of the supernatant was transferred in GC-vial and dried on

SpeedVac overnight.

Metabolite derivatization was performed by using a multi-purpose sample preparation robot

(Gerstel). Dried medium extracts were dissolved in 30 µl pyridine, containing 20 mg/mL

methoxyamine hydrochloride (Sigma-Aldrich), for 120 min at 45 °C under shaking. After

adding 30 µl N-methyl-N-trimethylsilyl-trifluoroacetamide (Macherey-Nagel), samples were

incubated for 30 min at 45 °C under continuous shaking.

GC-MS analysis was performed by using an Agilent 7890A GC coupled to an Agilent 5975C

inert XL Mass Selective Detector (Agilent Technologies). A sample volume of 1 µl was

injected into a Split/Splitless inlet, operating in split mode (10:1) at 270 °C. The gas

chromatograph was equipped with a 30 m (I.D. 0.25 mm, film 0.25 µm) DB-5ms capillary

column (Agilent J&W GC Column) with 5 m guard column in front of the analytical column.

Helium was used as carrier gas with a constant flow rate of 1.2 ml/min. The GC oven

temperature was held at 90 °C for 1 min and increased to 220 °C at 10 °C/min. Then, the

temperature was increased to 280 °C at 20 °C/min followed by 5 min post run time at 325 °C.

The total run time was 22 min. The transfer line temperature was set to 280 °C. The MSD was

operating under electron ionization at 70 eV. The MS source was held at 230 °C and the

quadrupole at 150 °C. Mass spectra were acquired in selected ion monitoring (SIM) mode for

precise quantification of medium components. In Table 3, masses used for quantification and

qualification of the derivatized target analytes (dwell times between 25 and 70 ms) are

illustrated.

Table 3: List of quantification and qualification ions used for metabolite detection in medium.

Analyte Name	Quantification Ions (*m/z*)	Qualification Ion I (m/z)	Qualification Ion II (m/z)
Pyruvic acid 1MEOX 1TMS	174-179	158.1	189.1
Lactic acid 2TMS	219.1-224.1	190.1	117.1
Alanine 2TMS	218.1-223.1	190.1	116.1
Valine 2TMS	144.1	218.1	246.2
Leucine 2TMS	158.1	218.1	232.2
Isoleucine 2TMS	158.1	218.1	232.2
Glycine 3TMS	276.1-280.1	248.1	174
Serine 3TMS	306.1-311.1	218.1	204.1
Threonine 3TMS	218.1	291.2	320.2
IS Pentanedioic acid-D6 2TMS	267.1	163.1	239.1
Methionine 2TMS	176.1	250.1	293.1
Glutamic acid 3TMS	363.2-371.2	246.1	348.2
Phenylalanine 2TMS	192.1	218.1	266.1
Asparagine 3TMS	231.1-236.1	132.1	348.1
IS [UL-13C5]-Ribitol 5TMS	220.1	310.2	323.2
Glutamine 3TMS	347.2-354.2	245.1	156.1
Glucose 1MEOX 5TMS	319.2, 323.2	217.1	220.1
Lysine 4TMS	174.1	317.2	434.3
Tyrosine 3TMS	218.1	179.1	280.2
Tryptophan 2TMS	218.1-223.1	130	348.2
Cystine 4TMS	218.1-223.1	297.1	411.2

All GC-MS chromatograms were processed using MetaboliteDetector (v. 3.2.20190704).

Compounds were annotated by retention time and mass spectrum using an in-house mass

spectral (SIM) library (overall similarity: >0.60). The following deconvolution settings were

applied: Peak threshold: 2; Minimum peak height: 2; Bins per scan: 10; Deconvolution width:

8 scans; no baseline adjustment; Minimum 1 peaks per spectrum; no minimum required base

peak intensity.

Peak areas of all isotopologues of defined quantification ions were summed up and divided by

the peak area of the internal standard for normalization. In addition, a calibration curve was

prepared to calculate absolute concentrations. Absolute uptake and release rates were

calculated as described in Hiller et al. 2009 (58).

<u>LC-MS measurements</u>

Untargeted LC-MS analysis was carried out as previously (42). For calculating intracellular

metabolite level, measured peak areas were initially corrected for natural isotope abundance,

normalized to internal standards and subsequently divided by PCV to obtain relative metabolite

levels.

<u>YSI measurements</u>

Collected medium samples were initially filtrated (Phenex-RC 4mm Syringe Filters 0.2 μm)

prior measurement to remove particles. Absolute quantitative values for Glutamine and

Glutamic acid were acquired using the YSI 2950D Biochemistry Analyzer (Kreienbaum

KWM). For a precise and reliable quantification, external calibration curves of each compound

were prepared and measured in duplicates. To get absolute consumption and release rate

(CORE, fmol/PCV/h), measured concentrations were divided by PCV (determined at start and

end of the experiment, see stable isotope tracing for details) and the duration of the experiment.

<u>IFN and Serine related genes in IBD bulk and scRNA-Seq datasets</u>

The publicly available GSE109142 (59) (RNA sequencing TPM counts of pediatric ulcerative

colitis patients) and GSE73661 (60) (microarray of UC patients treated with either infliximab

or vedolizumab) were downloaded from the Gene Expression Omnibus (GEO) repository and

processed using *R* version 4.1.0. TPM counts were log2 transformed after adding half of the

smallest non-zero value.

Digital gene expression matrices for SCP259 (45) (colon mucosa of 18 UC patients and 12

healthy individuals) were downloaded from the Single Cell Portal and processed in *R* using the

Seurat package. Following UMAP dimensional reduction, cells were clustered using the

FindClusters() function in Seurat at a resolution of 0.5. Main cell types were labelled using the

SingleR package and the Encode+Blueprint reference dataset. Epithelial cells were further

subclustered at a resolution of 0.5 and the 24 identified clusters labelled using known epithelial

markers published by Elmentaite et al., 2021 (61). Cells extracted from the lamina propria were

considered for subsequent analyses.

Gene expression scores were calculated as the mean of scaled and centered expression values

(z-scores) for the bulk datasets or on the mean normalized counts for the scRNA-Seq dataset.

ISG composing signature genes were *IFITM1*, *MX1*, *OAS3*, *IFIT1*, *IFI44L* and *IFI16* as

previously described (23). The serine score was calculated on the serine pathway related genes

PHGDH, *MTHFD2*, *SHMT2*, *MTHFD1L* and *SLC7A11*.

Quantification and statistical analysis

The GraphPad Prism 5 software package (GraphPad Software Inc., La Jolla, USA) was used

for statistical analysis. If not otherwise specified, the Student's unpaired t-test was performed.

Data is visualized as mean ± standard error of the mean (SEM), if not otherwise stated. A

significance threshold of a p-value ≤ 0.05 (*) was used, a p-value of ≤ 0.01, respectively ≤

0.001, were considered as strongly significant (**) and highly significant (***), if not

otherwise specified.

We define one *n* as one independent biological experiment (in some cases further consisting of

several wells, e.g. triplicate wells for all stable isotope tracing experiments). The technical

mean of one biological experiment was considered as one *n*. The mean values of several

independent, biological experiments (as indicated in figure legends) were plotted and used for

statistical analysis as indicated.

Funding & Acknowledgement

This work was supported by the DFG Research Training Group 1743 (P.R.), the BMBF

iTREAT project (P.R.), DFG Cluster of excellence "Precision medicine in chronic

inflammation" RTF III and TI-1, the DFG CRC 1182 C2 (P.R.) the EKFS research grant

#2019_A09 (K.A.), the Wilhelm Sander-Stiftung #2019.046.1 (K.A.), the BMBF (eMED

Juniorverbund "Try-IBD" 01ZX1915A), the DFG RU5042 (P.R., K.A.), the FNR-ATTRACT

program A18/BM/11809970 (J.M.), the "European Union under the Horizon 2020 Research

and Innovation Program" (S.J.W) and the MSCA-Innovative Training Networks Programme

MSCA-ITN (EDGE, 675278) (S.J.W.). Stable isotope tracers were provided as part of a

Research Award by Cambridge Isotope Laboratories Inc. (B.B. and J.M.).

We thank M. Rohm, M. Hansen, S. Kock, D. Oelsner, S. Baumgarten, M. Reffelmann, J.

Ohrndorf for perfect technical assistance. We also thank the LCSB Metabolomics Platform,

especially Xiangyi Dong and Floriane Vanhalle, for providing technical and analytical support;

the National Cytometry Platform (Quantitative Biology Unit, LIH) and especially Thomas

Cerutti for support with flow cytometric analyses.

<u>**References**</u>

1.   Kaplan GG. The global burden of IBD: from 2015 to 2025. Nat Rev Gastroenterol

Hepatol. 2015 Dec;12(12):720–7.

2.   Schenk W, Klugmann T, Borkenhagen A, Klecker C, Dietel P, Kirschner R, et al. The

detection of the cytomegalovirus DNA in the colonic mucosa of patients with ulcerative colitis

is associated with increased long-term risk of proctocolectomy: results from an outpatient IBD

clinic. Int J Colorectal Dis. 2019 Mar;34(3):393–400.

3.   Qin Y, Wang G, Kong D, Li G, Wang H, Qin H, et al. Risk Factors of Cytomegalovirus

Reactivation in Ulcerative Colitis Patients: A Meta-Analysis. Diagn Basel Switz. 2021 Oct

21;11(11):1952.

4.   Grootjans J, Kaser A, Kaufman RJ, Blumberg RS. The unfolded protein response in

immunity and inflammation. Nat Rev Immunol. 2016 Aug;16(8):469–84.

5.   Adolph TE, Tomczak MF, Niederreiter L, Ko HJ, Böck J, Martinez-Naves E, et al.

Paneth cells as a site of origin for intestinal inflammation. Nature. 2013 Nov;503(7475):272–

6.

6.   Kaser A, Lee AH, Franke A, Glickman JN, Zeissig S, Tilg H, et al. XBP1 Links ER

Stress to Intestinal Inflammation and Confers Genetic Risk for Human Inflammatory Bowel

Disease. Cell. 2008 Sep 5;134(5):743–56.

7.   Heazlewood CK, Cook MC, Eri R, Price GR, Tauro SB, Taupin D, et al. Aberrant

Mucin Assembly in Mice Causes Endoplasmic Reticulum Stress and Spontaneous

Inflammation Resembling Ulcerative Colitis. PLOS Med. 2008 Mar 4;5(3):e54.

8.   Tréton X, Pédruzzi E, Cazals–Hatem D, Grodet A, Panis Y, Groyer A, et al. Altered

Endoplasmic Reticulum Stress Affects Translation in Inactive Colon Tissue From Patients

With Ulcerative Colitis. Gastroenterology. 2011 Sep 1;141(3):1024–35.

9.   Zhao F, Edwards R, Dizon D, Afrasiabi K, Mastroianni JR, Geyfman M, et al.

Disruption of Paneth and goblet cell homeostasis and increased endoplasmic reticulum stress

in Agr2−/− mice. Dev Biol. 2010 Feb 15;338(2):270–9.

10.   Stengel S, Messner B, Falk-Paulsen M, Sommer N, Rosenstiel P. Regulated proteolysis

as an element of ER stress and autophagy: Implications for intestinal inflammation. Biochim

Biophys Acta BBA - Mol Cell Res. 2017 Nov 1;1864(11):2183–90.

11.   Vanhove W, Nys K, Arijs I, Cleynen I, Noben M, De Schepper S, et al. Biopsy-derived

Intestinal Epithelial Cell Cultures for Pathway-based Stratification of Patients With

Inflammatory Bowel Disease. J Crohns Colitis. 2018 Jan 24;12(2):178–87.

12.   Powell N, Pantazi E, Pavlidis P, Tsakmaki A, Li K, Yang F, et al. Interleukin-22

orchestrates a pathological endoplasmic reticulum stress response transcriptional programme

in colonic epithelial cells. Gut. 2020 Mar;69(3):578–90.

13.   Welz L, Kakavand N, Hang X, Laue G, Ito G, Silva MG, et al. Epithelial X-Box Binding

Protein 1 Coordinates Tumor Protein p53-Driven DNA Damage Responses and Suppression

of Intestinal Carcinogenesis. Gastroenterology. 2022 Jan;162(1):223-237.e11.

14.   Tschurtschenthaler M, Adolph TE, Ashcroft JW, Niederreiter L, Bharti R, Saveljeva S,

et al. Defective ATG16L1-mediated removal of IRE1α drives Crohn's disease–like ileitis. J

Exp Med. 2017 Feb 1;214(2):401–22.

15.   Motwani M, Pesiridis S, Fitzgerald KA. DNA sensing by the cGAS–STING pathway

in health and disease. Nat Rev Genet. 2019 Jul 29;1–18.

16.   Hopfner KP, Hornung V. Molecular mechanisms and cellular functions of cGAS–

STING signalling. Nat Rev Mol Cell Biol. 2020 May 18;1–21.

17.   Wottawa F, Bordoni D, Baran N, Rosenstiel P, Aden K. The role of cGAS/STING in

intestinal immunity. Eur J Immunol. 2021;51(4):785–97.

18.   Lio CWJ, McDonald B, Takahashi M, Dhanwani R, Sharma N, Huang J, et al. cGAS-

STING Signaling Regulates Initial Innate Control of Cytomegalovirus Infection. J Virol. 2016

Sep 1;90(17):7789–97.

19.   Piersma SJ, Poursine-Laurent J, Yang L, Barber GN, Parikh BA, Yokoyama WM. Virus

infection is controlled by hematopoietic and stromal cell sensing of murine cytomegalovirus

through STING. eLife. 2020 Jul 29;9.

20.   Stempel M, Chan B, Juranić Lisnić V, Krmpotić A, Hartung J, Paludan SR, et al. The

herpesviral antagonist m152 reveals differential activation of STING-dependent IRF and NF-

κB signaling and STING's dual role during MCMV infection. EMBO J. 2019 Mar

1;38(5):e100983.

21.   Guimarães ES, Gomes MTR, Campos PC, Mansur DS, Dos Santos AA, Harms J, et al.

Brucella abortus Cyclic Dinucleotides Trigger STING-Dependent Unfolded Protein Response

That Favors Bacterial Replication. J Immunol Baltim Md 1950. 2019 May 1;202(9):2671–81.

22.   Moretti J, Roy S, Bozec D, Martinez J, Chapman JR, Ueberheide B, et al. STING

Senses Microbial Viability to Orchestrate Stress-Mediated Autophagy of the Endoplasmic

Reticulum. Cell. 2017 Nov 2;171(4):809-823.e13.

23.   Aden K, Tran F, Ito G, Sheibani-Tezerji R, Lipinski S, Kuiper JW, et al. ATG16L1

orchestrates interleukin-22 signaling in the intestinal epithelium via cGAS–STING. J Exp Med.

2018 Nov 5;215(11):2868–86.

24.   Isler JA, Skalet AH, Alwine JC. Human Cytomegalovirus Infection Activates and

Regulates the Unfolded Protein Response. J Virol. 2005 Jun;79(11):6890–9.

25.   Stengel ST, Fazio A, Lipinski S, Jahn MT, Aden K, Ito G, et al. Activating

Transcription Factor 6 Mediates Inflammatory Signals in Intestinal Epithelial Cells Upon

Endoplasmic Reticulum Stress. Gastroenterology. 2020 Jul;S0016508520349246.

26.   Coleman OI, Lobner EM, Bierwirth S, Sorbie A, Waldschmitt N, Rath E, et al.

Activated ATF6 Induces Intestinal Dysbiosis and Innate Immune Response to Promote

Colorectal Tumorigenesis. Gastroenterology. 2018 Nov 1;155(5):1539-1552.e12.

27.   Dambacher J, Beigel F, Seiderer J, Haller D, Goke B, Auernhammer CJ, et al.

Interleukin 31 mediates MAP kinase and STAT1/3 activation in intestinal epithelial cells and

its expression is upregulated in inflammatory bowel disease. Gut. 2007 Apr 5;56(9):1257–65.

28.   Onyeagocha C, Hossain MS, Kumar A, Jones RM, Roback J, Gewirtz AT. Latent

cytomegalovirus infection exacerbates experimental colitis. Am J Pathol. 2009

Nov;175(5):2034–42.

29. Hansen K, Prabakaran T, Laustsen A, Jørgensen SE, Rahbæk SH, Jensen SB, et al.

Listeria monocytogenes induces IFNβ expression through an IFI16-, cGAS- and STING-

dependent pathway. EMBO J. 2014 Aug 1;33(15):1654–66.

30. Zhang J, Pavlova NN, Thompson CB. Cancer cell metabolism: the essential role of the

nonessential amino acid, glutamine. EMBO J. 2017 May 15;36(10):1302–15.

31. Meister A. New aspects of glutathione biochemistry and transport--selective alteration

of glutathione metabolism. Nutr Rev. 1984 Dec;42(12):397–410.

32. Chen Y, Shertzer HG, Schneider SN, Nebert DW, Dalton TP. Glutamate cysteine ligase

catalysis: dependence on ATP and modifier subunit for regulation of tissue glutathione levels.

J Biol Chem. 2005 Oct 7;280(40):33766–74.

33. Bannai S, Sato H, Ishii T, Taketani S. Enhancement of glutathione levels in mouse

peritoneal macrophages by sodium arsenite, cadmium chloride and glucose/glucose oxidase.

Biochim Biophys Acta. 1991 Apr 17;1092(2):175–9.

34. Liu T, Sun L, Zhang Y, Wang Y, Zheng J. Imbalanced GSH/ROS and sequential cell

death. J Biochem Mol Toxicol [Internet]. 2021 Nov 2 [cited 2021 Nov 4]; Available from:

https://onlinelibrary.wiley.com/doi/10.1002/jbt.22942

35. Owen JB, Butterfield DA. Measurement of Oxidized/Reduced Glutathione Ratio. In:

Bross P, Gregersen N, editors. Protein Misfolding and Cellular Stress in Disease and Aging

[Internet]. Totowa, NJ: Humana Press; 2010 [cited 2021 Nov 4]. p. 269–77. (Methods in

Molecular Biology; vol. 648). Available from: http://link.springer.com/10.1007/978-1-60761-

756-3_18

36. Liu Y, Adachi M, Zhao S, Hareyama M, Koong AC, Luo D, et al. Preventing oxidative

stress: a new role for XBP1. Cell Death Differ. 2009 Jun;16(6):847–57.

37. Mayr L, Grabherr F, Schwärzler J, Reitmeier I, Sommer F, Gehmacher T, et al. Dietary

lipids fuel GPX4-restricted enteritis resembling Crohn's disease. Nat Commun. 2020 Apr

14;11(1):1775.

38.   Forcina GC, Dixon SJ. GPX4 at the Crossroads of Lipid Homeostasis and Ferroptosis.

PROTEOMICS. 2019 Sep;19(18):1800311.

39.   Benzarti M, Delbrouck C, Neises L, Kiweler N, Meiser J. Metabolic Potential of Cancer

Cells in Context of the Metastatic Cascade. Cells. 2020 Sep 5;9(9):2035.

40.   Yang M, Vousden KH. Serine and one-carbon metabolism in cancer. Nat Rev Cancer.

2016 Oct;16(10):650–62.

41.   Maddocks ODK, Athineos D, Cheung EC, Lee P, Zhang T, van den Broek NJF, et al.

Erratum: Corrigendum: Modulating the therapeutic response of tumours to dietary serine and

glycine starvation. Nature. 2017 Aug;548(7665):122–122.

42.   Meiser J, Tumanov S, Maddocks O, Labuschagne CF, Athineos D, Van Den Broek N,

et al. Serine one-carbon catabolism with formate overflow. Sci Adv. 2016 Oct;2(10):e1601273.

43.   Weinstabl H, Treu M, Rinnenthal J, Zahn SK, Ettmayer P, Bader G, et al. Intracellular

Trapping of the Selective Phosphoglycerate Dehydrogenase (PHGDH) Inhibitor **BI-4924**

Disrupts Serine Biosynthesis. J Med Chem. 2019 Sep 12;62(17):7976–97.

44.   Kiweler N, Delbrouck C, Neises L, Pozdeev VI, Soriano-Baguet L, Xian F, et al.

Mitochondrial One-Carbon Flux has a Growth-Independent Role in Promoting Breast Cancer

Metastasis [Internet]. Cancer Biology; 2021 May [cited 2022 Feb 14]. Available from:

http://biorxiv.org/lookup/doi/10.1101/2021.05.27.445928

45.   Smillie CS, Biton M, Ordovas-Montanes J, Sullivan KM, Burgin G, Graham DB, et al.

Intra- and Inter-cellular Rewiring of the Human Colon during Ulcerative Colitis. Cell. 2019

Jul;178(3):714-730.e22.

46.   Le-Trilling VTK, Ebel JF, Baier F, Wohlgemuth K, Pfeifer KR, Mookhoek A, et al.

Acute Cytomegalovirus Infection Modulates the Intestinal Microbiota and Targets Intestinal

Epithelial Cells [Internet]. Microbiology; 2022 Mar [cited 2022 May 17]. Available from:

http://biorxiv.org/lookup/doi/10.1101/2022.03.17.484735

47.   Rodríguez-Sánchez I, Munger J. Meal for Two: Human Cytomegalovirus-Induced

Activation of Cellular Metabolism. Viruses. 2019 Mar 19;11(3):273.

48.   Shen L, Hu P, Zhang Y, Ji Z, Shan X, Ni L, et al. Serine metabolism antagonizes

antiviral innate immunity by preventing ATP6V0d2-mediated YAP lysosomal degradation.

Cell Metab. 2021 May;33(5):971-987.e6.

49.   Madison BB, Dunbar L, Qiao XT, Braunstein K, Braunstein E, Gumucio DL. Cis

elements of the villin gene control expression in restricted domains of the vertical (crypt) and

horizontal (duodenum, cecum) axes of the intestine. J Biol Chem. 2002 Sep 6;277(36):33275–

83.

50.   Jin L, Hill KK, Filak H, Mogan J, Knowles H, Zhang B, et al. MPYS Is Required for

IFN Response Factor 3 Activation and Type I IFN Production in the Response of Cultured

Phagocytes to Bacterial Second Messengers Cyclic-di-AMP and Cyclic-di-GMP. J Immunol.

2011 Sep 1;187(5):2595–601.

51.   Vidal K, Grosjean I, evillard JP, Gespach C, Kaiserlian D. Immortalization of mouse

intestinal epithelial cells by the SV40-large T gene. Phenotypic and immune characterization

of the MODE-K cell line. J Immunol Methods. 1993 Nov 5;166(1):63–73.

52.   Lee AH, Iwakoshi NN, Anderson KC, Glimcher LH. Proteasome inhibitors disrupt the

unfolded protein response in myeloma cells. Proc Natl Acad Sci. 2003 Aug 19;100(17):9946–

51.

53.   Sato T, Vries RG, Snippert HJ, Wetering M van de, Barker N, Stange DE, et al. Single

Lgr5 stem cells build crypt-villus structures in vitro without a mesenchymal niche. Nature.

2009 May;459(7244):262–5.

54.   Yin X, Farin HF, van Es JH, Clevers H, Langer R, Karp JM. Niche-independent high-

purity cultures of Lgr5+ intestinal stem cells and their progeny. Nat Methods. 2014

Jan;11(1):106–12.

55. Sorbara MT, Ellison LK, Ramjeet M, Travassos LH, Jones NL, Girardin SE, et al. The Protein ATG16L1 Suppresses Inflammatory Cytokines Induced by the Intracellular Sensors Nod1 and Nod2 in an Autophagy-Independent Manner. Immunity. 2013 Nov 14;39(5):858–73.

56. Chan B, Magalhães VG, Lemmermann NAW, Lisnić VJ, Stempel M, Bussey KA, et al. The murine cytomegalovirus M35 protein antagonizes type I IFN induction downstream of pattern recognition receptors by targeting NF-κB mediated transcription. PLOS Pathog. 2017 May 25;13(5):e1006382.

57. Kiweler N, Delbrouck C, Pozdeev VI, Neises L, Soriano-Baguet L, Eiden K, et al. Mitochondria preserve an autarkic one-carbon cycle to confer growth-independent cancer cell migration and metastasis [Internet]. Cancer Biology; 2021 May [cited 2022 Apr 2]. Available from: http://biorxiv.org/lookup/doi/10.1101/2021.05.27.445928

58. Hiller K, Hangebrauk J, Jäger C, Spura J, Schreiber K, Schomburg D. MetaboliteDetector: comprehensive analysis tool for targeted and nontargeted GC/MS based metabolome analysis. Anal Chem. 2009 May 1;81(9):3429–39.

59. Haberman Y, Karns R, Dexheimer PJ, Schirmer M, Somekh J, Jurickova I, et al. Ulcerative colitis mucosal transcriptomes reveal mitochondriopathy and personalized mechanisms underlying disease severity and treatment response. Nat Commun. 2019 Jan 3;10(1):38.

60. Arijs I, De Hertogh G, Lemmens B, Van Lommel L, de Bruyn M, Vanhove W, et al. Effect of vedolizumab (anti-α4β7-integrin) therapy on histological healing and mucosal gene expression in patients with UC. Gut. 2018 Jan;67(1):43–52.

61. Elmentaite R, Kumasaka N, Roberts K, Fleming A, Dann E, King HW, et al. Cells of the human intestinal tract mapped across space and time. Nature. 2021 Sep 9;597(7875):250–

1056   5.

1057

Fig. 1

Fig. S1

Fig. 2

Fig. S2

Fig. 3

Fig. S3

Fig. S4