Midbrain-Hippocampus Structural Integrity Selectively Predicts Motivated Memory Encoding

Blake L. Elliott[1]

Kimberlee D'Ardenne[2]

Vishnu P. Murty[1]

Gene A. Brewer[2]

Samuel M. McClure[2]

1. Department of Psychology, Temple University, Philadelphia, PA, USA

2. Department of Psychology, Arizona State University, Tempe, AZ, USA

Correspondence: Blake L. Elliott; Blake.Elliott@temple.edu

Abstract

Motivation is a powerful driver of learning and memory. Functional MRI studies show that interactions between the dopaminergic midbrain (SN/VTA), hippocampus, and nucleus accumbens (NAc) are critical for motivated memory encoding. However, it is not known if these effects are transient and purely functional, or if individual differences in the structure of this circuit underlie motivated memory encoding. To quantify individual differences in structure, diffusion-weighted MRI and probabilistic tractography were used to quantify SN/VTA-striatum and SN/VTA-hippocampus pathways associated with motivated memory encoding in humans. Participants completed a motivated source memory paradigm. During encoding, words were randomly assigned to one of three conditions: reward ($1.00), control ($0.00), or punishment (-$1.00). During retrieval, participants were asked to retrieve item and source information of the previously studied words and were rewarded or penalized according to their performance. Source memory for words assigned to both reward and punishment conditions was greater than control words, while there were no differences in item memory based on value. Anatomically, probabilistic tractography results revealed a heterogeneous, topological arrangement of the SN/VTA. Related to memory, tract integrity measures of SN/VTA-hippocampus pathways were positively correlated with individual differences in reward and punishment modulated memory performance, while tract integrity measures of SN/VTA-striatum pathways showed no association. This novel finding suggests that pathways emerging from the human SV/VTA are anatomically separable and functionally heterogeneous. Individual differences in structural integrity of the dopaminergic hippocampus-VTA loop are selectively associated with motivated memory encoding.

Keywords: value-directed remembering, episodic memory, source memory, dopamine, DTI, MRI

Significance Statement

Functional MRI studies show that interactions between the dopaminergic midbrain (SN/VTA), hippocampus, and nucleus accumbens (NAc) are critical for motivated memory encoding. This has led to competing theories that posit either SN/VTA-NAc reward prediction errors or SN/VTA-hippocampus signals underlie motivated memory encoding. Additionally, it is not known if these effects are transient and purely functional, or if individual differences in the structure of these circuits underlie motivated memory encoding. Using Diffusion-weighted MRI and probabilistic tractography, we show that tract integrity measures of SN/VTA-hippocampus pathways are positively correlated with motivated memory performance, while tract integrity measures of SN/VTA-striatum pathways show no association. This finding suggests that anatomical individual differences of the dopaminergic hippocampus-VTA loop are selectively associated with motivated memory encoding.

Midbrain-Hippocampus Structural Integrity Selectively Predicts Motivated Memory

Encoding

The motivation to obtain rewards and avoid punishments is a powerful driver of learning and memory. Scientists have investigated the effect of reward motivation on human episodic memory for nearly a century (Heyer Jr & O'Kelly, 1949; Kahneman & Peavler, 1969; Loftus & Wickens, 1970; Weiner & Walker, 1966). In cognitive neuroscience, most studies have focused on functional properties of the dopamine system rather than more stable neuroanatomical differences. We used diffusion weighted imaging to investigate structural connectivity of neural circuits involved in reward- and punishment-motivated memory encoding.

The neural mechanisms underlying reward-motivated memory encoding remain unresolved. Modulatory effects of the dopaminergic reward system on memory-related brain regions of the medial temporal lobe are hypothesized to be critical (Knowlton & Castel, 2022; Shohamy & Adcock, 2010). Neuroimaging studies in humans have found the involvement of the midbrain substantia nigra/ventral tegmental area (SN/VTA) and the nucleus accumbens (NAc) in the prediction of both rewards and punishments (Carter et al., 2009; McClure et al., 2003; Samanez-Larkin et al., 2007). Results from studies implementing functional magnetic resonance imaging (fMRI) during reward-motivated memory encoding have shown that increased activation in the medial temporal lobe and the reward system (i.e. SN/VTA and NAc) are predictive of better memory prioritization for higher-valued stimuli (Adcock et al., 2006; Shigemune et al., 2014; Wittmann et al., 2005). However, it is not clear if this pattern of activation represents canonical SN/VTA-NAc reward prediction errors observed during reinforcement learning (Calderon et al., 2021; Jang et al., 2019; Schultz, 1998), SN/VTA-hippocampus reward/novelty signals (Lisman et al., 2011; Lisman & Grace, 2005), or some

combination of the two (Ergo et al., 2020). To test between these possibilities, the present study tested whether the structural integrity of one or both pathways is associated with behavioral measures of motived memory encoding.

Anatomically, animal research has revealed that the midbrain SN/VTA is not a uniform structure. The SN/VTA has a distinct topological organization, with heterogeneous parallel pathways from dopamine nuclei that support separate cognitive processes. The medial regions (VTA, medial SN) innervate the ventral (limbic) striatum, while more lateral regions innervate the dorsal (executive and motor) striatum (Hedreen & Delong, 1991; Lynd-Balta & Haber, 1994; Selemon & Goldman-Rakic, 1990; Szabó, 1979). The SN/VTA-striatum connections form an upward-spiraling circuit that integrates signals from affective, executive control, and motor regions to produce goal-directed behavior (Haber, 2016). The SN/VTA also forms a circuit with the hippocampus that is responsible for encoding information into long-term memory (i.e. the hippocampus-VTA loop; Lisman & Grace, 2005). The upward arc of the hippocampus-VTA loop consists of direct dopaminergic projections to the hippocampus (Gasbarri et al., 1991; Gasbarri et al., 1997; Lewis et al., 2001; Swanson, 1982). These neurons release dopamine into the hippocampus, which promotes synaptic plasticity by enhancing long-term potentiation (Lisman et al., 2011; Lisman & Grace, 2005).

Although the relationship between the anatomical circuitry of the SN/VTA and its functions has been documented in rodents and nonhuman primates, research in human populations has been limited to functional activation and functional connectivity. We used diffusion weighted MRI and probabilistic tractography to assess structural connectivity of the SN/VTA to reward- and punishment-motivated memory encoding. We hypothesize that the SN/VTA will be anatomically and functionally heterogeneous. Anatomically, we aim to resolve

the topological heterogeneity of the SN/VTA non-invasively in a neurotypical human population. Medial regions (VTA) were hypothesized to have higher connectivity with ventral (limbic) regions of the striatum and with the hippocampus. More lateral regions of the SN/VTA were hypothesized to have higher connectivity with dorsal (executive and motor) regions of the striatum. Given the fMRI findings implicating the SN/VTA, NAc, and hippocampus in reward-motivated memory encoding, we hypothesized that structural integrity of one or both tracts would be associated with reward- and punishment-motivated memory encoding.

Method

Participants

Fifty participants were recruited via an online newsletter and flyers posted around the Arizona State University campus (mean age 20.36 years, SD = 1.94, min = 18.21 years, max = 28.17 years). Participants were told that they could be compensated up to $100 for an MRI study investigating memory and decision-making. Participants were all right-handed and were screened for neurological or psychiatric disorders. Two subjects were excluded from the DTI analysis. One subject was excluded because they were missing the reverse phase-encoding scan needed for FSL topup distortion correction. The other subject was excluded because they did not have a usable structural scan. Two additional subjects were excluded from the final analyses because they scored at or below chance level on their overall memory performance (i.e. hit rates).

Experimental Paradigm

The experimental task closely replicated Shigemune et al. (2014). The task consisted of 6 alternating encoding and retrieval phases (3 of each, with a retrieval period always following an encoding period). The stimuli consisted of 704 words drawn randomly from the Penn

Electrophysiology of Encoding and Retrieval Study word pool (an edited version of the Kucera-Francis word pool, available at http://memory.psych.upenn.edu/; Lohnas & Kahana, 2013). fMRI data were acquired during both the encoding and retrieval periods. The encoding phases consisted of 30 words each randomly assigned to one of three conditions: reward ($1.00), control ($0.00), or punishment (-$1.00), 10 of each value. The word and value pairs were shown concurrently for 2s on either the left or right side of the screen with an interstimulus interval randomly jittered between 3s and 5s. Participants were instructed to encode both the word (item) and location (source) of the word. The paradigm was presented using fMRI compatible video goggles.

The retrieval phases consisted of 45 words (30 from the most recent list and 15 new words, randomly intermixed) presented one at a time in the center of the screen without values. The words were presented for 2s with an interstimulus interval randomly jittered between 3s and 5s. Participants were asked to judge whether the word was previously presented on the left side of the screen during the study phase, the right side of the screen, or whether the word was a "new" word that had not been studied before. If the word was previously in the reward condition, the participants were told they would gain 50¢ for correctly recognizing the word but getting the location incorrect, and an additional 50¢ if both the word and location were correct. If the word was previously in the punishment condition, the participants were told they would avoid losing 50¢ for correctly recognizing the word but getting the location incorrect and avoid losing an additional 50¢ if both the word and location were correct. If the word was previously in the control condition the participant neither gained nor lost any money based on their memory performance. The participants were told they would lose 50¢ for incorrectly classifying "new"

words as "old" (i.e. false alarms). Participants were paid the total amount of money earned in the

task at the end of the session. Responses were made using a five-button optic response box.

Behavioral Data Analysis

Items with correct source memory (item with source hits: IWS) and items that were

correctly classified as old but with incorrect source memory (item only hits: IO) were aggregated

into proportions of total correct responses in that category by dividing by total number of studied

items. The experimental design is a 3×2 repeated measures ANOVA with the following factors:

condition (reward, control, punishment) and memory judgment (IWS, IO). Following Shigemune

et al. (2014), we expected a main effect of condition and memory judgment. Importantly, we also

expected a significant interaction between the two factors. Reward and punishment motivation

are hypothesized to selectively enhance associative memory and not item memory. Therefore, we

also expected a polynomial trend analysis to show a quadratic effect for the interaction between

the two factors (i.e. reward > control < punishment in the IWS condition only).

MRI Acquisition, Preprocessing, and Analysis

Structural MRI

MRI scans were acquired on a 3.0 T GE Discovery MR750 using a Nova Medical 32-

channel head coil (General Electric, Milwaukee, WI). For each participant, high-resolution, T1-

weighted structural images were collected using a co-planar, single shot, interleaved, 3D

magnetization-prepared rapid gradient-echo (BRAVO) in the sagittal plane (TR = 1900 ms, TE =

3.06 ms, FOV 23.0 cm, slice thickness 0.9 mm, slices per slab=192).

Diffusion-weighted MRI

Diffusion-weighted images were collected using 48 directions, with a total gradient

diffusion sensitivity of b = 1000 s/mm^2 with 6 repeats of the B0 (no diffusion weighting) image,

resulting in a 4D volume (dimension=256×256×80×54). An additional B0 image was collected

in the reverse phase encoding direction for use with FSL topup EPI distortion correction.

The diffusion-weighted data analysis closely follows the procedures from our previous

study in which we quantified structural integrity of SN/VTA-striatum tracts in a clinical sample,

establishing the reliability of this protocol (Elliott et al., 2021). The diffusion-weighted data were

processed using FSL (www.fmrib.ox.ac.uk/fsl). After correcting for movement and eddy current

artifacts, the diffusion parameters were calculated for each voxel using FSL bedpostX. Measures

of tract strength were calculated using probabilistic tractography, which is based on a

probabilistic Bayesian framework (Behrens et al., 2007; Behrens et al., 2003). Fiber tracking was

conducted in parallel for each voxel within a predefined SN/VTA seed mask. We used 5,000

samples per voxel, a curvature threshold of 0.2, and a step length of 0.5 mm. Target areas in the

striatum were defined using a connectivity-based segmentation atlas with subdivisions for

sensorimotor, executive, and limbic regions; this atlas is freely available with the FSL software

(http://fsl.fmrib.ox.ac.uk/fsl/fslwiki/Atlases/striatumconn). These striatal subdivisions also likely

reflect separable pathways from midbrain dopaminergic regions. The homogeneity of dopamine

release measured with positron emission tomography in response to the administration of

amphetamine is significantly higher within these connectivity-based subdivisions of the striatum

compared to within anatomical subdivisions (i.e., putamen, caudate, and nucleus accumbens

(Tziortzi et al., 2014). Hippocampus masks were defined using the Harvard-Oxford subcortical

atlas integrated within FSL (Figure 1). The MNI space target masks were normalized to each

participant's native space using the inverse of the spatial normalization parameters. To tailor the

ROIs to each individual's anatomy, we used individually segmented gray matter (GM) and

fractional anisotropy (FA) images to mask the ROIs. Following Tziortzi et al. (2014), the lower

threshold for the GM mask was set at 0.25, and the FA mask upper threshold was set at 0.40. The

seed ROI for the SN/VTA was defined using a probabilistic atlas of human SN/VTA (Murty et

al., 2014). We used a 50% probability threshold, and the mask was normalized to each

participant's native space using the inverse of the normalization parameters.

Because the dorsal and ventral tiers of the SN and VTA have not yet been reliably

resolved in humans with MRI, we use the term "SN/VTA-striatum tracts" to describe meso- and

nigro-striatal tracts. We note that the use of this term is not meant to imply directionality because

diffusion-weighted imaging data do not include information about whether tracts are efferent or

afferent. We chose to use the larger SN/VTA combined mask due to anatomical variability and

the relatively small size of the SN and VTA (Halliday & Törk, 1986; Keuken et al., 2014).

However, the results of the tractography analysis should follow our a priori predictions of the

topology of the SN/VTA. Higher probability of connection with the hippocampus and limbic

striatum should be localized in the medial SN and VTA. Higher probability of connection with

the executive and motor striatum should be localized primarily in the SN.

All tractography analyses were conducted in the participants' native anatomical space.

The FSL FDT toolbox was used to perform probabilistic tractography with a partial volume

model (Behrens et al., 2003), allowing for up to two fiber directions in each voxel (Behrens et

al., 2007). We generated 5,000 sample tracts from each voxel in the SN/VTA seed mask. Visual

inspection was used to ensure that the tractography maps were successful and acceptable for

further analysis. Tractography was performed separately for the left and right striatum and

hippocampus, and possible tracts were restricted to the hemisphere of origin using an exclusion

mask of the contralateral hemisphere. Following standard procedures, the seed-based

classification maps were first thresholded so that only voxels with at least 10 tracts terminating

in one of the target regions were kept (Cohen et al., 2009; Forstmann et al., 2012). Next, the

voxel values were converted into proportions of the number of tracts reaching the target mask

from one voxel, divided by the number of tracts generated from that voxel (maximum 5,000).

We used the mean of these value maps as the measure of SN/VTA-striatum and SN/VTA-

hippocampus tract strength.

SN/VTA Topology

To determine the organization of SN/VTA-striatum and SN/VTA- hippocampus across

participants, individual probability maps were normalized into Montreal Neurological Institute

space. Each map was thresholded so that only voxels that had at least 50 samples reaching any

target region were included (Tziortzi et al., 2014). The resulting maps were averaged to create 8

total for the entire sample (one for each target region for each hemisphere – limbic, executive,

and motor striatum and hippocampus, Figure 4). Visual inspection and similarity metrics (Dice

and dilated Dice coefficients) were used to quantify the topological arrangement of the

probability maps in relation to the SN and VTA, respectively. Because the larger combined

SN/VTA probability maps completely encapsulated both the SN and VTA, the maps were

restricted to include only voxels that were greater than the mean value of the entire map. Thus,

only voxels with the highest probability of connection were retained for comparison. Similarity

metrics (Dice and dilated Dice coefficients) were calculated for the thresholded maps and the SN

and VTA separately for each hemisphere (Table 1).

The Dice coefficient represents the degree of spatial overlap between two ROIs and was

calculated as follows: DICE $= \frac{2 \ x \ ROI_1 \cap ROI_2}{ROI_1 + ROI_2}$. The dilated Dice coefficient is a suitable measure

for small structures with complex shapes such as the VTA. This measure is similar to the Dice

coefficient except that it dilates each mask by one voxel. This means that a single-voxel offset is

not penalized as it is for the classical Dice coefficient, an important factor for small, complex

structures. Both coefficients range from 0 to 1, with 0 representing no overlap and 1 representing

complete overlap. Both measures were derived using the "segmentation_statistics" function from

the Nighres python toolbox (Huntenburg et al., 2018). We expected that limbic and hippocampus

maps would have higher Dice and dilated Dice values with the VTA and lower values with the

SN. Conversely, the executive and motor maps were expected to have higher Dice and dilated

Dice values with the SN and lower values with the VTA.

Tract Integrity and Motivated-Memory

The integrity of each fiber tract was calculated by taking the mean of each probability

map corresponding to each target region for each subject (four per hemisphere, eight in total).

Hierarchical regressions were used to test the association of SN/VTA-hippocampus tracts and

SN/VTA-limbic tracts with behavioral measures of reward and punishment prioritization (i.e. the

difference between correct reward and control IWS hits, and the difference between correct

punishment and control IWS hits). The diffusion measures for both the hippocampus and limbic

tracts were averaged across the left and right hemispheres in order to circumvent the issue of

multicollinearity in our regression models. Intracranial volume was entered in the first step of the

model. The second step of the model included SN/VTA-hippocampus tracts and SN/VTA-limbic

tracts, with intracranial volume treated as a covariate.

We further tested the association of SN/VTA-hippocampus tracts and SN/VTA-limbic

tracts on reward and punishment prioritization separately for each hemisphere. Tract integrity

measures were correlated across subjects with behavioral measures of reward and punishment

prioritization. Partial correlations were conducted, with intracranial volume included as a

covariate. Based on previous fMRI evidence implicating the hippocampus-VTA loop in reward-

motivated memory encoding, we expected positive correlations between SN/VTA-hippocampus tract integrity and individual differences in reward- and punishment-motivated encoding. The SN/VTA and the NAc have also been implicated in reward-motivated memory, possibly due to reward prediction error signaling (Ergo et al., 2020). We therefore also conducted correlations between SN/VTA-limbic tract integrity and individual differences in reward- and punishment-motivated encoding. Correlations between SN/VTA and other striatal regions were exploratory and are provided in Extended Data Figure 6-1.

Results

Behavioral Results

Memory performance as a function of value and response type (correct item with source, IWS, and item only, IO, performance) is summarized in Figure 2. The results replicate Shigemune et al. (2014). Specifically, a 3 (condition: reward, control, punishment) × 2 (memory judgment: IWS, IO) repeated measures ANOVA with Greenhouse-Geisser corrections revealed a marginal effect of condition ($F_{(1.427,47)} = 3.44$, $p = 0.053$, $\eta_p^2 = .07$), and a main effect of memory judgment ($F_{(1,47)} = 152.8$, $p < 0.001$, $\eta_p^2 = .77$). We also found a significant interaction between the two factors ($F_{(1.92,47)} = 4.24$, $p < 0.05$, $\eta_p^2 = .08$). A quadratic trend analysis revealed a significant interaction of value and response type ($F_{(1,47)} = 6.78$, $p < 0.05$, $\eta_p^2 = .13$). These results demonstrate that both reward and punishment motivated encoding conditions increased memory for items that were accurately remembered with the correct source location (Figure 2).

We quantified individual differences in memory prioritization for reward and punishment as the difference between reward and control IWS hits and the difference between punishment and control IWS hits (Figure 3). We chose to use only IWS hits because this is where the effect of value was isolated in previous experiments with similar tasks (Elliott et al., 2020; Shigemune

et al., 2014). This method has also been shown to form a stable latent factor with reward-modulated "remember" response in a VDR remember-know paradigm (Elliott et al., 2020). A one sample t-test showed that both the reward (M = .06, SD = .17) and punishment (M = .07, SD = .17) effects were statistically greater than 0: $t(47) = 2.36$, $p < 0.05$, Cohen's d = .341; $t(47) = 2.795$, $p < 0.05$, Cohen's d = .403. These effects were also strongly correlated (Figure 2; $r(47) = .723$, $p < .001$). Due to concerns that these effects may be driven by a participant with large reward and punishment effects (greater than 3 SD from the mean in both categories, Mahalanobis distance = 1.9; Extended Data Figure 3-1), we performed bootstrapped one sample t-tests with 1,000 samples for reward (95% CI .014 to .109, $p < .05$) and punishment (95% CI .024 to .117, $p < .05$). We also performed one sample t-tests with this subject removed. Both the reward and punishment effects were statistically greater than 0: $t(46) = 2.16$, $p < 0.05$, Cohen's d = .315; $t(46) = 2.75$, $p < 0.01$, Cohen's d = .401, and the measures were still highly correlated (Figure 3; $r(46) = .549$, $p < .001$). We chose to remove this outlier from the following connectivity correlation analyses. For completeness and transparency, scatterplots and statistics with this participant included can be found in the Extended Data Figure 5-1 and 6-2.

Structural Organization of the SN/VTA

Before relating structural integrity to motivated memory, we first wanted to confirm that the anatomical organization of the SN/VTA corroborates with what has been shown in rodents and non-human primates. When tracts with endpoints in the midbrain SN/VTA, functionally-parcellated striatal regions, and hippocampus were classified, four subdivisions emerged within the midbrain SN/VTA (Figure 4). The organization of these subdivisions was consistent with known anatomy in animals and non-human primates (Haber & Knutson, 2010). The subdivisions revealed an inverse dorsal-ventral and medial-lateral connectivity with the striatum. Projections

from the SN/VTA to the limbic striatum were primarily located in the VTA and medial SN.

Projections from the SN/VTA to the executive striatum were located more laterally, primarily in

the SN. Projections from the SN/VTA the motor striatum were located most laterally in the SN.

Additionally, connectivity between the SN/VTA and hippocampus was primarily localized

within voxels located in the VTA.

To further assess the topology of the SN/VTA projections, Dice and dilated Dice

coefficients were calculated between each target region and the SN and VTA separately. Because

the larger combined SN/VTA probability maps would completely encapsulate both the SN and

VTA, these maps were restricted to include only voxels that were greater than the mean value of

the entire map, leaving only the voxels with the highest probability of connection with each

target region. The results are shown in Table 1. Both Dice and dilated Dice similarity metrics

revealed a linear pattern of overlap with the SN and the target regions in both hemispheres.

Motor tracts had the largest Dice coefficients with the SN, followed by executive, limbic, and

hippocampus. The opposite pattern was shown with the VTA. Limbic and hippocampus tracts

had the largest Dice coefficients, followed by executive and lastly, motor tracts. These findings

comport with our a priori hypothesis and what is known from animal histology. We were able to

demonstrate non-invasively in humans a topological arrangement of the midbrain SN/VTA.

Tract Strength and Motivated Memory Performance

We next conducted analyses to determine whether individual differences in reward-

motivated memory prioritization were related to SN/VTA-hippocampus tract integrity using

hierarchical multiple regression (Table 2). The first step of the hierarchical regression model

included intracranial volume, so that the amount of variance explained by the structural integrity

measures can be isolated in the next step by examining the change in R^2. Based on this first step,

10.7% of the variance in reward prioritization was accounted for by intracranial volume. Results

from the second step revealed that SN/VTA-hippocampus tracts alone were associated with

reward prioritization. Based on this additive model, 15.1% of the variance in reward

prioritization could be accounted for by SN/VTA- hippocampus tract integrity, controlling for

intracranial volume. Cohen's f^2 indicated that this was a medium effect ($f^2 = 0.18$).

A similar hierarchical regression was conducted for punishment prioritization (Table 3).

The first step of the hierarchical regression model included intracranial volume, so that the

amount of variance explained by the structural integrity measures can be isolated in the next step

by examining the change in R^2. Based on this first step, 0.7% of the variance in reward

prioritization was accounted for by intracranial volume. Results from the second step revealed

that SN/VTA-hippocampus tracts alone were associated with punishment prioritization. Based

on this additive model, 17.6% of the variance in reward prioritization could be accounted for by

SN/VTA-hippocampus tract integrity, controlling for intracranial volume. Cohen's f^2 indicated

that this was a medium effect ($f^2 = 0.21$).

We further tested the association of SN/VTA- hippocampus tracts and SN/VTA-limbic

tracts on reward and punishment prioritization separately for each hemisphere. We computed

partial correlations for the reward effect and SN/VTA-hippocampus tract integrity and the

punishment effect and SN/VTA-hippocampus tract integrity for each hemisphere (controlling for

intracranial volume). The results revealed that both the reward ($r(42) = .315, p < .05$) and

punishment ($r(42) = .325, p < .05$) memory effects were positively correlated with the left

SN/VTA-hippocampus tract. Additionally, both the reward ($r(42) = .325, p < .05$) and

punishment ($r(42) = .340, p < .05$) were positively correlated with the right SN/VTA-

hippocampus tract as well (Figure 5). We also conducted analyses to determine whether

individual differences in memory prioritization were related to SN/VTA-limbic tract integrity. We computed partial correlations for the reward effect and SN/VTA-limbic tract integrity and the punishment effect and SN/VTA-limnic tract integrity for each hemisphere (controlling for intracranial volume). The results revealed that neither the reward ($r(42) = -.159$, *N.S.*) nor punishment ($r(42) = -.122$, *N.S.*) memory effects were correlated with the left SN/VTA-limbic tracts. Additionally, both the reward ($r(42) = .221$, *N.S.*) and punishment ($r(42) = .112$, *N.S.*) were not correlated with the right SN/VTA-hippocampus tracts as well (Figure 6).

It is difficult to interpret null results when comparing two experimental effects. Therefore, we conducted fisher r-to-z transformations to directly compare correlations between SN/VTA-hippocampus and SN/VTA-NAc pathways for both reward and punishment (Fisher, 1921). This analysis revealed a significant difference in the correlations for both reward ($Z = 2.23$, $p < 0.05$, two-tailed) and punishment ($Z = 2.11$, $p < 0.05$, two-tailed) in the left hemisphere. No significant difference was found for reward ($Z = 0.52$, *N.S.*, two-tailed) or punishment ($Z = 1.11$, *N.S.*, two-tailed) in the right hemisphere.

We also conducted analyses to test whether individual differences in memory prioritization were related to the other SN/VTA-striatum tracts (SN/VTA-executive, SN/VTA-motor). The measures were exploratory, and we expected no significant correlations to emerge. Statistics and figures can be found in Extended Data Figure 6-1. We computed partial correlations between the reward effect and SN/VTA-striatum tract integrity and the punishment effect and SN/VTA- striatum tract integrity for each hemisphere (controlling for intracranial volume). The results revealed that neither reward nor punishment effects were correlated to SN/VTA-striatum tract integrity (Extended Data Figure 6-1).

Discussion

The current study used diffusion weighted MRI and probabilistic tractography to map the topology of the midbrain SN/VTA and to relate structural connectivity measures to individual differences in reward- and punishment-motivated memory encoding. Behaviorally, both reward and punishment effects were localized to associative memory (correct source recognition) and not found for item memory. These findings support previous findings that reward- and punishment-motivated encoding selectively enhances deeper, recollective memories (Elliott & Brewer, 2019; Gruber & Otten, 2010; Hennessee et al., 2017). Memories that rely on recollection afford access to rich associative and contextual information from the study episode, whereas memories that rely upon familiarity only afford access to information of the study items. Different neural regions have been shown to dissociate recollection and familiarity. Recollection relies more upon the hippocampus, whereas familiarity relies more upon the perirhinal cortex (Diana et al., 2007). These results fit with the "binding of items and contexts" model. This model posits that the perirhinal cortex processes item information whereas the parahippocampal cortex processes context information. Finally, the hippocampus binds item and context information to support episodic memories (Ranganath, 2010). Our findings that reward- and punishment-motivated memory encoding selectively enhances source memory may be indicative of enhanced hippocampal processing, thereby permitting more enriched encoding episodes.

The neural mechanisms underlying motivated memory remain opaque. It is hypothesized that interactions between the dopaminergic reward system (SN/VTA and NAc) and the hippocampus are critical. Previous neuroimaging studies in humans have found the involvement of the SN/VTA and the NAc in the prediction of both rewards and punishments (Carter et al., 2009, McClure et al., 2003; Samanez-Larkin et al., 2007), and in learning from these predictions via reinforcement learning (Schultz et al., 1997). Previous fMRI studies in humans during

motivated memory encoding tasks have shown increased activation in the medial temporal lobe, SN/VTA and NAc are predictive of better memory prioritization for reward-predicting stimuli (Adcock et al., 2006; Wittmann et al., 2005) and punishment-predicting stimuli (Shigemune et al., 2014). Individual differences in functional connectivity between the SN/VTA and hippocampus have also been shown to predict individual differences in motivated memory performance (Wolosin et al., 2012).

Interpretations of the SN/VTA, NAc, and hippocampal activation observed during motivated memory encoding vary. Some researchers suggest that canonical SN/VTA-ventral striatum reward prediction errors underlie the reward-driven gain in episodic memory performance (Calderon et al., 2021; De Loof et al., 2018; Jang et al., 2019; Rouhani et al., 2018). Other researchers have theorized that the hippocampus-VTA loop is critical for motivated memory encoding (Adcock et al., 2006). This latter hypothesis suggests that dopamine released into the hippocampus via midbrain projections increases hippocampal plasticity, thereby enhancing encoding and consolidation of information into long-term memory (Lisman et al., 2011). However, these studies have all focused on functional activation within these circuits. Our findings suggest that neuroanatomical differences across people determine the strength of this reward-related effect on memory. We found that individual differences in tract integrity of the SN/VTA-hippocampus were positively correlated with both reward and punishment effects on human episodic memory. Additionally, Fisher's r-to-z transformation revealed that these correlations were specific to the SN/VTA-hippocampus tracts and not the SN/VTA-NAc tracts (in the left hemisphere). The results provide evidence that structural connectivity of dopaminergic midbrain-hippocampus projections underlie reward-motivated memory encoding, rather, or in addition to, canonical SN/VTA-NAc reward prediction error signaling.

Although other researchers have theorized SN/VTA-limbic striatal reward prediction

error signaling can drive memory performance (Calderon et al., 2021), we believe this may be a

separate neural system that can influence episodic memory, depending on the task demands.

Studies that find traditional SN/VTA-limbic striatum reward prediction errors support episodic

memory encoding typically implement a reinforcement learning paradigm with either

instrumental or Pavlovian learning via feedback. Memoranda in these tasks are almost always

incidentally encoded (Ergo et al., 2020). Studies have also found that hippocampal-driven

episodic memory can compete for processing resources and impair striatal feedback-based

learning (Foerde et al., 2013; Wimmer et al., 2014). The motivated source memory task

implemented here requires participants to use effortful memory encoding strategies, which could

bias activity in favor of the hippocampal-VTA episodic memory. Future studies are needed to

further dissociate the exact contributions of the SN/VTA reward system to effortful episodic

encoding versus conditioned learning.

Further evidence that motivated-memory encoding may have different neural generators

from standard SN/VTA-striatum reward prediction errors comes from a recent

electroencephalography study (Elliott et al., 2020). This study parametrically manipulated

reward value during the encoding period of a recognition memory task. The study found that a

P3 event-related potential during encoding scaled linearly with the value of the studied item.

Furthermore, individual differences in this P3 component were positively correlated with

individual differences in reward-motivated memory performance. However, when we analyzed

the feedback related negativity (FRN) component during encoding (an event-related potential

component theorized to reflect canonical reward prediction error signals transmitted from the

midbrain to the ACC; Sambrook & Goslin, 2014; San Martín, 2012; Walsh & Anderson, 2012)

we failed to find an effect of value. The findings suggest that different neural generators may code different types of reward value depending on the task (motivated memory versus reinforcement learning).

Studies have shown that connectivity between frontal cortex and striatum is defined by functionally disparate circuits that subserve sensorimotor, executive, and limbic functions. These parallel circuits are topographically organized in a dorsal-ventral configuration, with each structural circuit maintaining the functional characteristics of the target cortical region. Midbrain dopamine neurons display similar functional heterogeneous circuits with their projections to the striatum. The midbrain-striatum projections have a spiraling medial/lateral and inverse ventral/dorsal topographical arrangement from the VTA and SN. The dorsomedial regions of the SN/VTA innervate the ventral striatum and the ventrolateral regions innervate the dorsal striatum (Fallon & Moore, 1978; Lynd-Balta & Haber, 1994; McRitchie et al., 1996). This spiraling architecture is hypothesized to allow affective limbic signals to influence executive and motor actions (Haber & Knutson, 2010; Haber, 2016). We used tract integrity measures to test if connections to the hippocampus and striatum from the SN/VTA regions display the same anatomy known from primate and rodent histology. The SN/VTA-striatum connections revealed an inverse dorsal-ventral and medial-lateral configuration. The dorsomedial SN/VTA showed greatest connectivity to the limbic striatum, with more ventrolateral regions of the SN/VTA showing highest connectivity to the dorsal striatum. Importantly, connections to the hippocampus also fell primarily within the VTA and medial SN region. These findings demonstrate that it is feasible to non-invasively measure heterogeneous midbrain circuits in a neurotypical human population.

Anatomically, the hippocampus receives direct dopaminergic projections from the VTA (Bethus et al., 2010; Gasbarri et al., 1997; Luo et al., 2011; Zubair et al., 2021). To our knowledge, we are the first group to use probabilistic tractography to quantify SN/VTA-hippocampus integrity in a group of participants. Although tractography does not have the ability to resolve afferent and efferent connections, the hippocampus has no known direct efferent connections to the SN/VTA. Additionally, our results found that the hippocampus showed greatest connectivity with the dorsomedial SN/VTA, corroborating rodent and primate findings (Lisman & Grace, 2005, Zubair et al., 2021). We believe this is further evidence that the tracts quantified here do, in fact, represent dopaminergic VTA-hippocampus projections. The tractography results extend what is known from the animal literature to a neurotypical human population.

We found no significant correlations between reward and punishment effects and SN/VTA-striatum connectivity. Although previous fMRI studies implementing motivated-memory tasks have found NAc activation in addition to SN/VTA and hippocampus activation, this may not reflect the SN/VTA-limbic pathways quantified here. The hippocampus-VTA loop consists of two primary pathways: (1) the upward arc (VTA-hippocampus) that we quantified in this study and (2) the downward arc (hippocampus-NAc-ventral pallidum-VTA). If the hippocampus-VTA loop is requisite for motivated (reward and punishment) effects on episodic memory, it is possible that the NAc activation observed in previous studies is representative of activation along the downward arc (hippocampus-NAc connections), and not direct SN/VTA-NAc connectivity associated with reward prediction errors, as often interpreted. Therefore, the NAc activation observed in other motivated memory paradigms could be epi-phenomenal, resulting from hippocampal signaling.

Memory is critical for goal-directed behavior, but human memory has a finite capacity. The excess of sensory information humans encounter daily requires that we selectively prioritize and encode important or rewarding information at the expense of less important information, allowing our memory systems to be adaptive. It is hypothesized that interactions between the dopaminergic reward system and hippocampus underlie memory selectivity. We found that both reward- and punishment-motivated memory encoding preferentially enhance associative memory and not item memory. Additionally, structural integrity of the midbrain-hippocampus connections was predictive of individual differences in motivated memory prioritization for both rewards and punishments. These results provide compelling evidence to support theories of SN/VTA heterogeneity and the involvement of the dopaminergic reward system in motivated memory encoding.

Acknowledgements

This work was supported by an NSF GRFP (B.L.E.), NSF grant 1634179 (S.M.M.), and NIH grant R01 MH091068 (S.M.M.).

Conflicts of Interest

The authors declare no conflicts of interest.

Figure 1: Seed and Target Regions for the probabilistic tractography analysis. The target regions of the striatum were functionally segmented based on projections from motor, executive, and limbic cortices (Tziortzi et al., 2014). The seed region of the midbrain was generated from a combined SN/VTA probabilistic atlas (Murty et al., 2014).

Figure 2: Memory performance as a function of encoding condition. IWS = item with source memory, IO = item only memory. Error bars represent 95% CI. * *p* < .05

Figure 3: Reward and punishment effects (difference scores). A) Violin plot B) Scatterplot (* *p* < 0.001).

Figure 4: Group average projections from the SN/VTA to the four target regions (motor, executive, limbic, hippocampus).

Figure 5: Scatterplots of reward and punishment effects and hippocampus tract integrity. A) Left hippocampus tract integrity and reward effect B) Right hippocampus tract integrity and reward effect C) Left hippocampus tract integrity and punishment effect D) Right hippocampus tract integrity and punishment effect (* *p* < 0.05).

Figure 6: Scatterplots of reward and punishment effects and limbic tract integrity. A) Left limbic tract integrity and reward effect B) Right limbic tract integrity and reward effect C) Left limbic tract integrity and punishment effect D) Right limbic tract integrity and punishment effect.

Table 1. Dice and dilated Dice statistics for the SN and VTA with each target region

	Dice		Dilated Dice	
Target Region	SN	VTA	SN	VTA
Left Motor	0.31	0.31	0.83	0.60
Left Executive	0.32	0.37	0.79	0.67
Left Limbic	0.23	0.52	0.67	0.79
Left Hippocampus	0.14	0.52	0.58	0.79
Right Motor	0.34	0.28	0.82	0.61
Right Executive	0.32	0.42	0.76	0.71
Right Limbic	0.30	0.49	0.71	0.80
Right Hippocampus	0.12	0.46	0.58	0.77

Table 2. Hierarchical linear regression models predicting reward prioritization

	B	SE	β	t	p
Step 1: Covariate					
Intracranial Volume	$1.05e^{-7}$	<0.001	0.107	0.703	0.486
Step 2: Additive Effect					
SN/VTA-Limbic	-0.071	0.504	-0.021	-0.140	0.889
SN/VTA-Hippocampus*	2.408	0.888	0.406	2.713	0.010

Table 3. Hierarchical linear regression models predicting punishment prioritization

	B	SE	β	t	p
Step 1: Covariate					
Intracranial Volume	$6.84e^{-9}$	<0.001	0.007	0.049	0.961
Step 2: Additive Effect					
SN/VTA-Limbic	-0.212	0.465	-0.067	-0.457	0.650
SN/VTA-Hippocampus*	2.422	0.818	0.439	2.960	0.005

References

Adcock, R. A., Thangavel, A., Whitfield-Gabrieli, S., Knutson, B., & Gabrieli, J. D. (2006). Reward-motivated learning: mesolimbic activation precedes memory formation. *Neuron*, *50*(3), 507-517.

Behrens, T. E., Berg, H. J., Jbabdi, S., Rushworth, M. F., & Woolrich, M. W. (2007). Probabilistic diffusion tractography with multiple fibre orientations: What can we gain? *neuroimage*, *34*(1), 144-155.

Behrens, T. E., Woolrich, M. W., Jenkinson, M., Johansen-Berg, H., Nunes, R. G., Clare, S., . . . Smith, S. M. (2003). Characterization and propagation of uncertainty in diffusion-weighted MR imaging. *Magnetic Resonance in Medicine: An Official Journal of the International Society for Magnetic Resonance in Medicine*, *50*(5), 1077-1088.

Bethus, I., Tse, D., & Morris, R. G. (2010). Dopamine and memory: modulation of the persistence of memory for novel hippocampal NMDA receptor-dependent paired associates. *Journal of Neuroscience*, *30*(5), 1610-1618.

Calderon, C. B., De Loof, E., Ergo, K., Snoeck, A., Boehler, C. N., & Verguts, T. (2021). Signed reward prediction errors in the ventral striatum drive episodic memory. *Journal of Neuroscience*, *41*(8), 1716-1726.

Carter, R. M., MacInnes, J. J., Huettel, S. A., & Adcock, R. A. (2009). Activation in the VTA and nucleus accumbens increases in anticipation of both gains and losses. *Frontiers in behavioral neuroscience*, *3*, 21.

Castel, A. D., Benjamin, A. S., Craik, F. I., & Watkins, M. J. (2002). The effects of aging on selectivity and control in short-term recall. *Memory & Cognition*, *30*(7), 1078-1085.

Cohen, M. X., Schoene-Bake, J.-C., Elger, C. E., & Weber, B. (2009). Connectivity-based segregation of the human striatum predicts personality characteristics. *Nature neuroscience*, *12*(1), 32-34.

Davachi, L., Mitchell, J. P., & Wagner, A. D. (2003). Multiple routes to memory: distinct medial temporal lobe processes build item and source memories. *Proceedings of the National Academy of Sciences*, *100*(4), 2157-2162.

De Loof, E., Ergo, K., Naert, L., Janssens, C., Talsma, D., Van Opstal, F., & Verguts, T. (2018). Signed reward prediction errors drive declarative learning. *PLoS One*, *13*(1), e0189212.

Delgado, M. R., Jou, R. L., & Phelps, E. A. (2011). Neural systems underlying aversive conditioning in humans with primary and secondary reinforcers. *Frontiers in neuroscience*, *5*, 71.

Diana, R. A., Yonelinas, A. P., & Ranganath, C. (2007). Imaging recollection and familiarity in the medial temporal lobe: a three-component model. *Trends in cognitive sciences*, *11*(9), 379-386.

Elliott, B. L., Blais, C., McClure, S. M., & Brewer, G. A. (2020). Neural correlates underlying the effect of reward value on recognition memory. *NeuroImage*, *206*, 116296.

Elliott, B. L., & Brewer, G. A. (2019). Divided attention selectively impairs value-directed encoding. *Collabra: Psychology*, *5*(1).

Elliott, B. L., D'Ardenne, K., Mukherjee, P., Schweitzer, J. B., & McClure, S. M. (2021). Limbic and Executive Meso-and Nigro-striatal Tracts Predict Impulsivity Differences in ADHD. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*.

Elliott, B. L., McClure, S. M., & Brewer, G. A. (2020). Individual differences in value-directed remembering. *Cognition*, *201*, 104275.

Ergo, K., De Loof, E., & Verguts, T. (2020). Reward prediction error and declarative memory. *Trends in Cognitive Sciences*, *24*(5), 388-397.

Fallon, J. H., & Moore, R. Y. (1978). Catecholamine innervation of the basal forebrain IV. Topography of the dopamine projection to the basal forebrain and neostriatum. *Journal of Comparative Neurology*, *180*(3), 545-579.

Floresco, S. B., West, A. R., Ash, B., Moore, H., & Grace, A. A. (2003). Afferent modulation of dopamine neuron firing differentially regulates tonic and phasic dopamine transmission. *Nature neuroscience*, *6*(9), 968-973.

Fisher, R. A. (1921). 014: On the" Probable Error" of a Coefficient of Correlation Deduced from a Small Sample.

Foerde, K., Braun, E. K., & Shohamy, D. (2013). A trade-off between feedback-based learning and episodic memory for feedback events: evidence from Parkinson's disease. *Neurodegenerative Diseases*, *11*(2), 93-101.

Forstmann, B. U., Keuken, M. C., Jahfari, S., Bazin, P.-L., Neumann, J., Schäfer, A., . . . Turner, R. (2012). Cortico-subthalamic white matter tract strength predicts interindividual efficacy in stopping a motor response. *Neuroimage*, *60*(1), 370-375.

Gasbarri, A., Campana, E., Pacitti, C., Hajdu, F., & Tömböl, T. (1991). Organization of the projections from the ventral tegmental area of Tsai to the hippocampal formation in the rat. *Journal fur Hirnforschung*, *32*(4), 429-437.

Gasbarri, A., Sulli, A., & Packard, M. G. (1997). The dopaminergic mesencephalic projections to the hippocampal formation in the rat. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, *21*(1), 1-22.

Gruber, M. J., & Otten, L. J. (2010). Voluntary control over prestimulus activity related to encoding. *Journal of Neuroscience*, *30*(29), 9793-9800.

Haber, S. N. (2016). Corticostriatal circuitry. *Dialogues in clinical neuroscience*, *18*(1), 7.

Halliday, G. M., & Törk, I. (1986). Comparative anatomy of the ventromedial mesencephalic tegmentum in the rat, cat, monkey and human. *Journal of Comparative Neurology*, *252*(4), 423-445.

Hedreen, J. C., & Delong, M. R. (1991). Organization of striatopallidal, striatonigral, and nigrostriatal projections in the macaque. *Journal of Comparative Neurology*, *304*(4), 569-595.

Hennessee, J. P., Castel, A. D., & Knowlton, B. J. (2017). Recognizing what matters: Value improves recognition by selectively enhancing recollection. *Journal of memory and language*, *94*, 195-205.

Heyer Jr, A., & O'kelly, L. (1949). Studies in motivation and retention: II. Retention of nonsense syllables learned under different degrees of motivation. *The Journal of psychology*, *27*(1), 143-152.

Huntenburg, J. M., Steele, C. J., & Bazin, P.-L. (2018). Nighres: processing tools for high-resolution neuroimaging. *GigaScience*, *7*(7), giy082.

Jang, A. I., Nassar, M. R., Dillon, D. G., & Frank, M. J. (2019). Positive reward prediction errors during decision-making strengthen memory encoding. *Nature human behaviour*, *3*(7), 719-732.

Johnson, J. D., & Rugg, M. D. (2007). Recollection and the reinstatement of encoding-related cortical activity. *Cerebral cortex*, *17*(11), 2507-2515.

Kahneman, D., & Peavler, W. S. (1969). Incentive effects and pupillary changes in association learning. *Journal of Experimental Psychology*, *79*(2p1), 312.

Keuken, M. C., Bazin, P.-L., Crown, L., Hootsmans, J., Laufer, A., Müller-Axt, C., . . . Turner, R. (2014). Quantifying inter-individual anatomical variability in the subcortex using 7 T structural MRI. *NeuroImage*, *94*, 40-46.

Kim, H., Shimojo, S., & O'Doherty, J. P. (2006). Is avoiding an aversive outcome rewarding? Neural substrates of avoidance learning in the human brain. *PLoS Biol*, *4*(8), e233.

Knowlton, B. J., & Castel, A. D. (2022). Memory and reward-based learning: A value-directed remembering perspective. *Annual review of psychology*, *73*, 25-52.

Knutson, B., Westdorp, A., Kaiser, E., & Hommer, D. (2000). FMRI visualization of brain activity during a monetary incentive delay task. *Neuroimage*, *12*(1), 20-27.

Lewis, D. A., Melchitzky, D. S., Sesack, S. R., Whitehead, R. E., Auh, S., & Sampson, A. (2001). Dopamine transporter immunoreactivity in monkey cerebral cortex: regional, laminar, and ultrastructural localization. *Journal of Comparative Neurology*, *432*(1), 119-136.

Lisman, J., Grace, A. A., & Duzel, E. (2011). A neoHebbian framework for episodic memory; role of dopamine-dependent late LTP. *Trends in neurosciences*, *34*(10), 536-547.

Lisman, J. E., & Grace, A. A. (2005). The hippocampal-VTA loop: controlling the entry of information into long-term memory. *Neuron*, *46*(5), 703-713.

Lodge, D. J., & Grace, A. A. (2011). Hippocampal dysregulation of dopamine system function and the pathophysiology of schizophrenia. *Trends in pharmacological sciences*, *32*(9), 507-513.

Loftus, G. R., & Wickens, T. D. (1970). Effect of incentive on storage and retrieval processes. *Journal of Experimental Psychology*, *85*(1), 141.

Lohnas, L. J., & Kahana, M. J. (2013). Parametric effects of word frequency in memory for mixed frequency lists. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, *39*(6), 1943.

Luo, A. H., Tahsili-Fahadan, P., Wise, R. A., Lupica, C. R., & Aston-Jones, G. (2011). Linking context with reward: a functional circuit from hippocampal CA3 to ventral tegmental area. *Science*, *333*(6040), 353-357.

Lynd-Balta, E., & Haber, S. (1994). The organization of midbrain projections to the striatum in the primate: sensorimotor-related striatum versus ventral striatum. *Neuroscience*, *59*(3), 625-640.

McClure, S. M., Berns, G. S., & Montague, P. R. (2003). Temporal prediction errors in a passive learning task activate human striatum. Neuron, 38(2), 339-346.

McRitchie, D., Hardman, C., & Halliday, G. (1996). Cytoarchitectural distribution of calcium binding proteins in midbrain dopaminergic regions of rats and humans. *Journal of comparative neurology*, *364*(1), 121-150.

Murty, V. P., & Adcock, R. A. (2014). Enriched encoding: reward motivation organizes cortical networks for hippocampal detection of unexpected events. *Cerebral Cortex*, *24*(8), 2160-2168.

Murty, V. P., Shermohammed, M., Smith, D. V., Carter, R. M., Huettel, S. A., & Adcock, R. A. (2014). Resting state networks distinguish human ventral tegmental area from substantia nigra. *Neuroimage*, *100*, 580-589.

Otten, L. J. (2007). Fragments of a larger whole: retrieval cues constrain observed neural correlates of memory encoding. *Cerebral Cortex*, *17*(9), 2030-2038.

Ranganath, C. (2010). Binding items and contexts: The cognitive neuroscience of episodic memory. *Current Directions in Psychological Science*, *19*(3), 131-137.

Ranganath, C., Yonelinas, A. P., Cohen, M. X., Dy, C. J., Tom, S. M., & D'Esposito, M. (2004). Dissociable correlates of recollection and familiarity within the medial temporal lobes. *Neuropsychologia*, *42*(1), 2-13.

Rouhani, N., Norman, K. A., & Niv, Y. (2018). Dissociable effects of surprising rewards on learning and memory. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, *44*(9), 1430.

Samanez-Larkin, G. R., Gibbs, S. E., Khanna, K., Nielsen, L., Carstensen, L. L., & Knutson, B. (2007). Anticipation of monetary gain but not loss in healthy older adults. *Nature neuroscience*, *10*(6), 787-791.

Sambrook, T. D., & Goslin, J. (2014). Mediofrontal event-related potentials in response to positive, negative and unsigned prediction errors. *Neuropsychologia*, *61*, 1-10.

San Martín, R. (2012). Event-related potential studies of outcome processing and feedback-guided learning. *Frontiers in human neuroscience*, *6*, 304.

Schultz, W. (1998). Predictive reward signal of dopamine neurons. *Journal of neurophysiology*, *80*(1), 1-27.

Schultz, W., Dayan, P., & Montague, P. R. (1997). A neural substrate of prediction and reward. *Science*, *275*(5306), 1593-1599.

Schwartz, S. T., Siegel, A. L., & Castel, A. D. (2020). Strategic encoding and enhanced memory for positive value-location associations. *Memory & cognition*, *48*, 1015-1031.

Selemon, L., & Goldman-Rakic, P. (1990). Topographic intermingling of striatonigral and striatopallidal neurons in the rhesus monkey. *Journal of Comparative Neurology*, *297*(3), 359-376.

Shigemune, Y., Tsukiura, T., Kambara, T., & Kawashima, R. (2014). Remembering with gains and losses: effects of monetary reward and punishment on successful encoding activation of source memories. *Cerebral cortex*, *24*(5), 1319-1331.

Shohamy, D., & Adcock, R. A. (2010). Dopamine and adaptive memory. *Trends in cognitive sciences*, *14*(10), 464-472.

Stefanidi, A., Ellis, D. M., & Brewer, G. A. (2018). Free recall dynamics in value-directed remembering. *Journal of memory and language*, *100*, 18-31.

Swanson, L. (1982). The projections of the ventral tegmental area and adjacent regions: a combined fluorescent retrograde tracer and immunofluorescence study in the rat. *Brain research bulletin*, *9*(1-6), 321-353.

Szabó, J. (1979). Strionigral and nigrostriatal connections. *Stereotactic and Functional Neurosurgery*, *42*(1-2), 9-12.

Tziortzi, A. C., Haber, S. N., Searle, G. E., Tsoumpas, C., Long, C. J., Shotbolt, P., . . . Rabiner, E. A. (2014). Connectivity-based functional analysis of dopamine release in the striatum using diffusion-weighted MRI and positron emission tomography. *Cerebral cortex*, *24*(5), 1165-1177.

Uncapher, M. R., & Rugg, M. D. (2005). Encoding and the durability of episodic memory: a functional magnetic resonance imaging study. *Journal of Neuroscience*, *25*(31), 7260-7267.

Walsh, M. M., & Anderson, J. R. (2012). Learning from experience: event-related potential correlates of reward processing, neural adaptation, and behavioral choice. *Neuroscience & Biobehavioral Reviews*, *36*(8), 1870-1884.

Watkins, M., & Bloom, L. (1999). Selectivity in memory: An exploration of willful control over the remembering process. *Unpublished manuscript*.

Weiner, B., & Walker, E. L. (1966). Motivational factors in short-term retention. *Journal of Experimental Psychology*, *71*(2), 190.

Wimmer, G. E., Braun, E. K., Daw, N. D., & Shohamy, D. (2014). Episodic memory encoding interferes with reward learning and decreases striatal prediction errors. *Journal of Neuroscience*, *34*(45), 14901-14912.

Wittmann, B. C., Schott, B. H., Guderian, S., Frey, J. U., Heinze, H.-J., & Düzel, E. (2005). Reward-related FMRI activation of dopaminergic midbrain is associated with enhanced hippocampus-dependent long-term memory formation. *Neuron*, *45*(3), 459-467.

Wolosin, S. M., Zeithamova, D., & Preston, A. R. (2012). Reward modulation of hippocampal subfield activation during successful associative encoding and retrieval. *Journal of Cognitive Neuroscience*, *24*(7), 1532-1547.

Wrase, J., Kahnt, T., Schlagenhauf, F., Beck, A., Cohen, M. X., Knutson, B., & Heinz, A. (2007). Different neural systems adjust motor behavior in response to reward and punishment. *NeuroImage*, *36*(4), 1253-1262.

Yonelinas, A. P. (2002). The nature of recollection and familiarity: A review of 30 years of research. *Journal of memory and language*, *46*(3), 441-517.

Zubair, M., Murris, S. R., Isa, K., Onoe, H., Koshimizu, Y., Kobayashi, K., . . . Isa, T. (2021). Divergent whole brain projections from the ventral midbrain in macaques. *Cerebral Cortex*, *31*(6), 2913-2931.