Lipidomic profiling reveals concerted temporal patterns of functionally related

lipids in *Aedes aegypti* females following blood feeding

Meng-Jia Lau[1*]; Shuai Nie[2*]; Qiong Yang[1]; Lawrence G. Harshman[1]; Cungui Mao[3]; Nicholas

A. Williamson[2]; Ary A. Hoffmann[1]

[1] Pest and Environmental Adaptation Research Group, Bio21 Institute and the School of

BioSciences, The University of Melbourne, Parkville, Victoria, Australia

[2] Mass Spectrometry and Proteomics Facility, Bio21 Institute, The University of Melbourne,

Parkville, Victoria, Australia

[3] Department of Medicine and Stony Brook Cancer Center, Renaissance School of Medicine,

Stony Brook University, Stony Brook, New York, United States

[*] corresponding author:

Email: lau.mengjia@gmail.com; shuai.nie@unimelb.edu.au

**Abstract**

We conducted a lipidomic analysis of the whole body of female *Aedes aegypti* mosquitoes at

different time points over the course of feeding and reproduction. There were temporal

biphasic increases of more than 80% of lipids identified at the time of feeding and from 16 h

to 30 h post blood meal. During these two increases, the abundance of many lipids dropped

while body weight remained stable, probably reflecting blood lipid digestion and the

synthesis of vitellogenin in this period. A concerted temporal pattern was particularly strong

at the second peak for membrane and signalling lipids such as PE, PI, CL, HexCer and LPA.

Lyso-glycerophospholipids showed three distinct change patterns which are functionally

related: LPE and LPC, which are membrane lipids, showed little change; LPA, a signalling

lipid, showed a dramatic increase from 16 to 30 h PBM; LPI, a bioactive lipid, and both LPG

and LPS which are bacterial membrane lipids, showed one significant increase from the time

of feeding to 16 hours post blood meal. The result of our study on the anautogenous insect *Ae.*

*aegypti* point to specific lipids likely to be important in the reproductive process with a role

in the formation and growth of ovarian follicles.

**Keywords**

Lipidomics; *Aedes aegypti*; reproduction; blood digestion

**Introduction**

The yellow fever mosquito, *Aedes aegypti*, is one of the major vectors that transmits

arbovirus diseases in tropical and subtropical areas, such as dengue and Zika [1-3]. *Ae. aegypti*

females are anautogenous: they must ingest human/vertebrate blood to obtain unique

nutrients to produce their eggs, and diseases are transmitted through the blood ingestion

process [4,5].

Reproduction in mosquito females is generally regulated through the provision of nutrition

and the modulation of hormones [6]. Essential nutrients such as amino acids and cholesterol

mainly come from human blood [7,8], and supports gene products involved in yolk

development [9]. The source and the quality of blood that females are fed can directly impact

their fertility [10-13], though their nutritional components such as amino acids and cholesterol

are considered similar [14,15]. Apart from these changes of amino acids, cholesterol and yolk

protein, the vittellogenic cycle is also expected to involve lipid metabolism; lipids not only

act as major energy resources in supporting egg development [16] and contribute 35% of

ovarian dry weight [17,18], but also exhibit a variety of functions in biological system [19,20].

The development of ovaries can be separated into the pre-vitellogenic (before blood meal,

BBM) and vitellogenic (post blood meal, PBM) stages depending on the timing of the blood

meal. In the pre-vitellogenic stages, lipids are reserved in the fat body [16], which is essential

to support *Ae. aegypti* reproduction [21], particularly when a sugar source is compromised.

Starving female mosquitoes before or after their blood feeding will not reduce their fecundity

but will significantly reduce the amount of lipids in mothers after laying eggs [22], suggesting

a trade-off between reproduction and female survival.

During the vitellogenic stages, consumption of blood meal triggers insulin-like peptide 3

(ILP3) from the brain to stimulate oogenesis [23], which represents the insulin signalling

pathway [24] The secretion of 20-hydroxyecdysone then increases the expression of

vitellogenin genes in the fat body to support yolk protein accumulation and ovarian follicle

development [25], which is stimulated by gonadotropin [26], and lipids are transferred from

the fat body to follicles [16]. In the meantime, *Ae. aegypti* females also digest and absorb

essential amino acids from human blood, where the concentration of amino acids increasing

in haemolymph acts as a signal to activate egg development [27] through the TOR pathway [7].

In this study, we used liquid chromatography coupled with the mass spectrometry (LCMS)-

based lipidomic method and a comprehensive lipid standard mixture covering near all major

lipid groups to identify and analyse lipid components of whole female individuals before the

blood meal and post blood meal. Our study aims to understand lipid metabolism in *Ae.*

*aegypti* during the reproductive process, and how lipids with different functions act to

support the development of eggs.

**Material and methods**

**Sample preparation**

*Aedes aegypti* mosquitoes in this experiment were collected from *Wolbachia*-uninfected

populations from Cairns, QLD, Australia and maintained in laboratory conditions. Samples

were prepared by hatching *Ae. aegypti* eggs stored in an incubator (PG50 Plant Growth

Chambers, Labec Laboratory Equipment, Marrickville, NSW, Australia) at 22-30 ± 1 °C,

with a 12 h : 12 h photoperiod. Larvae were reared in reverse osmosis (RO) water with yeast

and fed with Hikari Sinking Wafers fish food tablets until adult. All female mosquitoes were

blood fed by the same volunteer's forearm. The first blood meal was provided to adult

females at 4 ± 1 days post-emergence. Females were isolated individually and observed for

seven days to discard those that laid small numbers of eggs (< 50). At the seventh day, a

second blood meal was provided. Non- engorged individuals were removed from the

experiment. Samples were then stored as different groups in their second vitellogenic cycle

before blood meal (BBM) and at different time post blood meal (PBM): 0 h PBM (30 mins

after blood meal), 4 h PBM, 16 h PBM, 30 h PBM, 72 h PBM, and 120 h PBM. Sugar and

oviposition cups were provided during the entire period of the experiment.

**Female mosquito measurement**

Apart from analysis of the lipidome, we measured the weight of different female individuals

(four to seven) at each time point from the additional collected samples using a Sartorius

Analytical balance BP 210 D (Sartorius, Gottigen, Germany, Readability: 0.01 mg). After

weighing, eight mosquitoes were randomly selected from all time points and their wing

lengths were measured to indicate female body size [28]. To further understand the

development of female ovaries, we hatched batches of mosquito eggs and took photos of

ovaries using an NIS Elements BR imaging microscope (Nikon Instruments, Tokyo, Japan)

every day PBM.

**Lipid extraction**

For each group, we extracted the whole body of six mosquito females, keeping individuals

separate. We added 10 µL of Mouse SPLASH® LIPIDOMIX® Mass Spec Standard (cat.

330707, Avanti) after 1 in 4 dilution to each sample prior the extraction and then

homogenised samples in 200 µL ice-cold 60% methanol (MeOH) containing 0.01% (w/v)

butylated hydroxytoluene (BHT) by using TissueLyserII (Qiagen, Hilden, Germany) at 25

Hertz for one minute and incubating for 20 minutes in an ultrasonic bath (Soniclean Pty Ltd.,

Thebarton, Australia). Following the method previously reported by Lydic, *et al.* [29], 120 µL

of MilliQ water, 420 µl of MeOH with 0.01% (w/v) BHT, and 260 µL of chloroform ($CHCl_3$)

were added to each tube and vortexed thoroughly before incubating in a Bioshake iQ

(Quantifoil Instrument GmbH, Germany) setting at 1,400 rpm for 30 minutes and centrifuged

at 14,000 rpm for 15 minutes. Supernatant containing lipids was transferred to a new 2 mL

tube. The remaining sediments were re-extracted with 100 µL of MilliQ water and 400 µL of

$CHCl_3$ : MeOH (1:2, v:v) containing 0.01% (w/v) BHT as described above. Supernatants

from the first and second extraction were pooled in the same 2 mL tube and then dried by

evaporation under vacuum using a GeneVac miVac sample concentrator (SP Scientific,

Warminster, PA, USA). Before HPLC, dried lipid extracts were resuspended in 50 µL of

$CHCl_3$ : MeOH (1:9, v:v, containing 0.01% BHT).

**Mass spectrometry analyses**

Samples were analysed by ultrahigh performance liquid chromatography (UHPLC) coupled

to tandem mass spectrometry (MS/MS) employing a Vanquish UHPLC linked to an Orbitrap

Fusion Lumos mass spectrometer (Thermo Fisher Scientific), with separate runs in positive

and negative ion polarity. Solvent A was 6/4 (v/v) acetonitrile/water with 10 mM ammonium

acetate and 5 µM medronic acid and solvent B was 9/1 (v/v) isopropanol/acetonitrile with 10

mM ammonium acetate. Each sample was injected into an RRHD Eclipse Plus C18 column

(2.1 × 100 mm, 1.8 µm, Agilent Technologies) at 50°C at a flow rate of 350 µl/min for 3 min

using 3% solvent B. During separation, the percentage of solvent B was increased from 3% to

70% in 5 min, from 70% to 99% in 16 min, from 99% to 99% in 3 min, from 99% to 3% in

0.1 min and maintained at 3% for 3.9 min.

All experiments were performed using a Heated Electrospray Ionization source. The spray

voltages were 3.5 kV in positive ionisation mode and 3.0 kV in negative ionisation mode. In

both polarities, the flow rates of sheath, auxiliary and sweep gases were 20 and 6 and 1

'arbitrary' unit(s), respectively. The ion transfer tube and vaporizer temperatures were

maintained at 350 °C and 400 °C, respectively, and the ion funnel RF level was set at 50%. In

both positive and negative ionisation modes from 3 to 24 min, top speed data-dependent scan

with a cycle time of 1 s was used. Within each cycle, a full-scan MS spectrum was acquired

firstly in the Orbitrap at a mass resolving power of 120,000 (at $m/z$ 200) across a $m/z$ range of

300–2000 using quadrupole isolation, an automatic gain control (AGC) target of 4e5 and a

maximum injection time of 50 milliseconds, followed by higher-energy collisional

dissociation (HCD)-MS/MS at a mass resolving power of 15,000 (at $m/z$ 200), a normalised

collision energy (NCE) of 27% at positive mode and 30% at negative mode, an $m/z$ isolation

window of 1, a maximum injection time of 22 milliseconds and an AGC target of 5e4. For

the improved structural characterisation of glycerophosphocholine (PC) in positive mode, a

data-dependent product ion ($m/z$ 184.0733)-triggered collision-induced dissociation (CID)-

MS/MS scan was performed in the cycle using a q-value of 0.25 and a NCE of 30%, with

other settings being the same as that for HCD-MS/MS. For the improved structural

characterisation of triacylglycerol (TG) lipid in positive mode, the fatty acid + $NH_3$ neutral

loss product ions observed by HCD-MS/MS were used to trigger the acquisition of the top-3

data-dependent CID-MS3 scans in the cycle using a q-value of 0.25 and a NCE of 30%, with

other settings being the same as that for HCD-MS/MS.

**Lipid identification and quantification**

LC-MS/MS data was searched through MS Dial 4.80 [30,31]. The mass accuracy settings are

0.005 Da and 0.025 Da for MS1 and MS2. The minimum peak height is 50000 and mass slice

width is 0.05 Da. The identification score cut off is 80%. Post identification was done with a

text file containing name and *m/z* of each standard in Mouse SPLASH® LIPIDOMIX® Mass

Spec Standard. In positive mode, $[M+H]^+$ and $[M+NH_4]^+$ were selected as ion forms. In

negative mode, $[M-H]^-$ were selected as ion forms. All lipid classes available were selected

for the search. PC, LPC, DG, TG, CE, SM were identified and quantified at positive mode

while PE, LPE, PS, LPS, PG, LPG, PI, LPI, PA, LPA, Cer, CL were identified and quantified

at negative mode. The retention time tolerance for alignment is 0.1 min. Lipids with

maximum intensity less than 5-fold of average intensity in blank were removed. All other

settings were default. All lipid LC-MS features were manually inspected and re-integrated

when needed. We also removed four other types of lipids: (1) lipids with only sum

composition except PC and SM; (2) lipids identified due to peak tailing; (3) retention time

outliers within each lipid class; and (4) LPA and PA artifacts generated by in-source

fragmentation of LPS and PS. The shorthand notation used for lipid classification and

structural representation follows the nomenclature proposed previously [32].

Quantification of lipid species was achieved by comparison of the LC peak areas of identified

lipids against peak areas and quantities of the corresponding internal lipid standards in the

same or similar lipid class (Table 1), and the concentrations of all lipids were reported at

nmol/individual. Given that the commercially available stable isotope-labelled lipid standards

are limited, some of the identified lipids were normalised against a standard from a different

class or sub-class, and no attempts were made to quantitatively correct for different ESI

responses of individual lipids due to concentration, acyl chain length, degree of unsaturation,

or matrix effects caused by differences in chromatographic retention times compared with the

relevant standards.

**Data analysis**

All data were analysed through R v. 3.6.0. Female weight was analysed by ANOVA across

time followed by posthoc Tukey HSB tests. The concentrations of different lipid groups

(nmol/individual) were used as variables for principal component analysis (PCA). For

specific comparisons between treatments for particular lipids, concentration was natural log-

transformed before undertaking paired t-tests, and p values were corrected for lipid number

considered in a comparison by the Benjamin-Hochberg method.

**Results**

**Female body weight and wing length measurement**

*Aedes aegypti* female mosquitoes were stored for lipidomic study in their second vitellogenic

cycle before their blood meal (BBM) and at different time post blood meal (PBM): (0 h (30

mins after blood meal), 4 h, 16 h, 30 h, 72 h, and 120 h). We first weighed female mosquitoes

from samples collected at each time point (Figure 1), with the difference between BBM and 0

h PBM representing the amount of blood a female engorged. Sampling time had a significant

effect on female weight (ANOVA, $F_{7,41} = 13.06$, $P < 0.001$). The weights of engorged female

mosquitoes were approximately double their weights BBM. The females then started to

digest blood, and their weight dropped significantly from 0 h PBM to 4 h PBM, thereafter

stabilising until 30 h PBM, and then dropping again further to the BBM level at 72 h PBM;

Figure 1. **Weights** ~~of female Aegypti females across different time points~~ **over the course of feeding and reproduction**. Weights were measured at eight time points before blood meal (BBM) and post blood meal (PBM). The same letter at the top of the boxplots represents groups that are not significantly different following Tukey HSD post-hoc tests.

this change in weight reflects the process of blood digestion, which takes around 32 hours

after females are engorged [33,34]. The appearance of female ovaries was captured every day

under a microscope (Supplementary Material 1). We also calculated the coefficient of

variation (CV, CV = (Standard Deviation/Mean) × 100) for female wing length for the eight

samples we measured (mean ± se = 2.07 ± 0.06 mm, CV = 8.71) to indicate the spread of

mosquito body sizes in this study.

**Lipids identification and their changes during female reproductive process**

We measured lipidomic changes averaged to individual female BBM and at different time points PBM, covering the entire vitellogenic cycle. In total, we identified 456 lipids (Table 1) and divided them into six major groups: glycerolipids, diacyl-glycerophospholipids, lyso-glycerophospholipids, Cardiolipin (CL), Carnitine (CAR) and sphingolipids. The most abundant lipid group in female mosquitoes was the neutral glycerolipids, with 55 molecules making up approximately 80% of the total lipid amount. Glycerophospholipids accounted for 55% of the molecules identified in this class, with 193 diacyl-glycerophospholipids and 48 lyso-glycerophospholipids. We also identified 66 CL, 15 CAR and 69 sphingolipids.

Table 1. List of lipid standards, adduct type and number of lipids that were identified in each group.

| Name (Abbreviation) | Standard | Adduct type | Number of lipids identified |
|---|---|---|---|
| Diacylglycerol (DG) | 15:0-18:1(d7) DAG | $[M+NH_4]^+$ | 14 |
| Triglycerides (TG) | 15:0-18:1(d7)-15:0 TAG | $[M+NH_4]^+$ | 41 |
| Phosphatidylethanolamines (PE) | 15:0-18:1(d7) PE | $[M-H]^-$ | 43 |
| Ether-linked PE (EtherPE) | 15:0-18:1(d7) PE | $[M-H]^-$ | 18 |
| Oxidized PE (OxPE) | 15:0-18:1(d7) PE | $[M-H]^-$ | 8 |
| Phosphatidylinositol (PI) | 15:0-18:1(d7) PI | $[M-H]^-$ | 33 |
| Ether-linked PI (EtherPI) | 15:0-18:1(d7) PI | $[M-H]^-$ | 3 |
| Phosphatidylserine (PS) | 15:0-18:1(d7) PS | $[M-H]^-$ | 10 |
| Phosphatidic acids (PA) | 15:0_18:1(d7) PA | $[M-H]^-$ | 5 |
| Phosphatidylglycerol (PG) | 15:0_18:1(d7) PG | $[M-H]^-$ | 24 |
| Phosphatidylcholines (PC) | 15:0_18:1(d7) PC | $[M+H]^+$ | 52 |
| Ether-linked PC (EtherPC) | 15:0_18:1(d7) PC | $[M+H]^+$ | 7 |
| Lyso-PE (LPE) | 18:1(d7) Lyso PE | $[M-H]^-$ | 13 |
| Ether-linked lyso-PE (EtherLPE) | 18:1(d7) Lyso PE | $[M-H]^-$ | 2 |
| Lyso-PI (LPI) | 15:0-18:1(d7) PI | $[M-H]^-$ | 7 |
| Lyso-PS (LPS) | 15:0-18:1(d7) PS | $[M-H]^-$ | 6 |
| Lyso-PA (LPA) | 15:0_18:1(d7) PA | $[M-H]^-$ | 4 |
| Lyso-PG (LPG) | 15:0_18:1(d7) PG | $[M-H]^-$ | 7 |
| Lyso-PC (LPC) | 18:1(d7) Lyso PC | $[M+H]^+$ | 9 |
| Cardiolipin (CL) | 15:0_18:1(d7) PG | $[M-H]^-$ | 66 |
| Carnitine (CAR) | 15:0_18:1(d7) PC | $[M+H]^+$ | 15 |
| Sphingomyelin (SM) | d18:1-18:1(d9) SM | $[M+H]^+$ | 26 |
| Ceramides (Cer) | d18:1-18:1(d9) SM | $[M-H]^-$ | 23 |
| Hexosylceramides (HexCer) | d18:1-18:1(d9) SM | $[M-H]^-$ | 4 |
| Sulfatides (SHexCer) | d18:1-18:1(d9) SM | $[M-H]^-$ | 16 |

213 In order to obtain an overview of lipidomic changes during the reproductive process, we first

214 undertook a principal component analysis (PCA) to describe overall patterns (Figure 2). This

215 indicated that lipid components of female mosquitoes went through two main changes: the

216 first occurred immediately after a blood meal (0 h PBM), and the second change occurred

217 between 16 and 30 h PBM. Further analysis showed that the amounts of most lipids increased

218 at 0 h and 16 h PBM respectively (Figure 3), except for lyso-glycerophospholipids, in which

219 case the total amount remained relatively stable during the reproduction process. These

results are also reflected in PCA analyses on specific classes of lipids (Figure 4); for most

classes the first PC explained more than half of the variation, with the exception of the lyso-

glycerophospholipids where it only explained 31.8% of variation and there was no obvious

change directly after blood feeding. However there was a shift at a later time point for this

group, indicating that most changes in lyso-glycerophospholipid reflect mosquito metabolism

rather than direct blood ingestion.

Figure 2. **Principal Component Analysis suggested lipidome differences of** *Aedes aegypti*
**females at different time points over the course of feeding and reproduction.** Colour and
shape symbols correspond to the eight time points before blood meal (BBM) and post blood
meal (PBM). The first two principal axes explained 76 % of the variance.

Figure 3. **Changes of lipid amounts in *Aedes aegypti* females over the course of feeding and reproduction.** The total amount (nmol/individual) of glycerolipids, diacyl-glycerophospholipids, lyso-glycerophospholipids, cardiolipins, carnitines and sphingolipids were measured at eight time points before blood meal (BBM) and post blood meal (PBM).

**Identification of change patterns for specific lipids**

The amount of blood a female mosquito takes in when fully engorged can be similar to the

weight of the individual, or even nearly twice the individual's weight [35,36]. By comparing

the time points BBM and 0 h PBM for individual lipid, we were able to determine which

lipids were taken in with human blood. We found 86% (392 molecules) of all the lipids

identified significantly increased at the time of blood meal (Supplementary Material 2). An

exception occurred for glycerophospholipids and carnitines. We found 29 lyso-

glycerophospholipids and 19 diacyl-glycerophospholipids that did not increase significantly

after the blood meal, which might indicate that these lipids originated from mosquitoes

instead of the human blood. These lipids include most of the LPA, LPG identified, and all

LPI identified. The carnitines showed a mild increase generally. The carnitines

tetradecasphingenine (C14) and hexadecasphingenine (C16) CAR did not change

significantly, while most of the other carnitines (7 out of 11) increased only slightly (adjusted

$0.05 > p > 0.01$); only one molecule (CAR 20:1) in this class increased highly significantly

(adjusted $p < 0.001$).

From 0 h to 16 h PBM, the amount of lipids dropped while the weight of the mosquitoes did

not decrease correspondingly. From 16 h to 30 h PBM, 87.7% of lipids (400 molecules)

identified increased significantly (Supplementary Material 3), of which 392 showed at least a

two-fold increase, and 139 showed an increase of at least ten-fold. There were 34 lyso-

glycerophospholipids which did not increase significantly and these lipids were distributed in

all sub-classes of lyso-glycerophospholipids except for LPA. All LPA identified increased

more than ten-fold, while none of the individual lipids identified from LPC and LPE

significantly increased.

Unlike lyso-glycerophospholipids, almost all diacyl-glycerophospholipids had significantly

increased, including all ether PI, oxidized PE, PE, PA, PG, PI and PS, In particular, 30 out of

43 PE and 18 out of 33 PI identified increased more than ten-fold. Besides, 42 out of 66 CL

increased more than ten-fold and the four HexCer identified increased about ten-fold. In

summary, lipids that increased by more than ten-fold include most of PE, PI and CL that act

as membrane components, and all of HexCer and LPA that act as signalling molecules and

participate in a variety of physiological functions. These lipids are expected to be particularly

important in mosquito egg production.

In general, most individual lipids have similar change patterns, regardless of the length of

fatty acyl chain and number of double bonds. Outliers (that did not increase significantly

from BBM to 0 h PBM or from 16 to 30 h PBM) were found mostly in lyso-

glycerophospholipids. Looking at the overall pattern, 21 lipid molecules identified did not

increase from BBM to 0 h PBM or from 16 h to 30 h PBM, and 19 of these are lyso-

glycerophospholipids.

Figure 4. **Principal Component Analysis of different groups of lipids in *Aedes aegypti* females at different time points over the course of feeding and reproduction**. Colour and shape symbols correspond to the eight time points before blood meal (BBM) and post blood meal (PBM).

We then further analysed the change patterns of lyso-glycerophospholipids, which represent a

group composed of various sub-groups of lipids with different patterns of change (Figure 5),

and diverse metabolic roles such as metabolic intermediates, extracellular signalling

molecules and immune system activators. Specifically, the two most abundant lyso-

glycerophospholipids LPE and LPC, which play important roles in membrane composition,

were relatively stable in terms of total amount during reproduction. LPA, an important group

of lipids in regulating various signalling pathways, showed a dramatic increase from 16 to 30

h PBM, which is similar to the change patterns for other groups (other than lyso-

glycerophospholipids) described previously. LPI, LPG and LPS, on the other hand, only had

one significant increase at a time period from 0 to 16 h PBM, when most lipids in other

groups decreased. LPI, a bioactive lipid, can initiate a variety of functions such as cell growth,

differentiation, and immunity [37,38]. While LPS and LPG, membrane lipids in gram-positive

bacteria and gram-negative bacteria respectively, can act as microbial toxins to trigger

mosquito immune response by activating the Toll and Imd pathway [39,40]. In this study, the

changes of these two lipid groups most likely indicate participation of gut bacteria in

mosquito blood digestion.

Figure 5. **Changes of lipid amounts in lyso-glycerophospholipids in** *Aedes aegypti* **females over the course of feeding and reproduction.** The total amount (nmol/individual) of LPE; LPI; LPS; LPA; LPG and LPC were measured at eight time points before blood meal (BBM) and post blood meal (PBM).

By comparing the differences between the two periods (from BBM to 0 h PBM and from 16

h to 30 h PBM) when most increases occurred, we found that most lipids we identified were

present in both human blood and mosquitoes, and also being affected by mosquito ovarian

follicle development. Nevertheless, some lipids seemed to be present at a stable level in

mosquitoes and were not significantly increased after blood feeding, such as

tetradecasphingenine (C14) and hexadecasphingenine (C16) CAR, and most of the LPGs.

Some lipids seemed to exist at a much higher abundance in human blood but were not used in

mosquitoes to support egg development, which we define here as those that increased more

than ten-fold after blood feeding but did not increase significantly (adjust p > 0.05) at the

later period. These include some Oxidized PE and sphingomyelin.

**Discussion**

In the current study, we focused on lipidomic changes of whole female mosquitoes during the

reproductive process. Whole-body investigations provide an overall picture of lipid changes

that might not be evident when focusing only on organs or cells. We found biphasic increases

in more than 80% of lipids identified at the time of feeding and from 16 to 30 h PBM. The

amount of lipids dropped between these periods of increase, as also observed in the fat body

[41]. The weight of the mosquitoes did not decrease at the first time point when lipids

decreased, suggesting that lipids were transformed into different metabolites. Many

metabolites can be produced within this period such as glycogen, and fatty acids [20,42], but it

is more likely that the decrease of lipids from 0 to 16 h PBM reflects the synthesis of

vitellogenin in the fat body, which is triggered by blood feeding. Previous studies found that

the secretion of vitellogenin in the fat body reaches its highest peak at around 18 h PBM, and

decreases to the previtellogenic level at around 30 h PBM [43,44].

Pitch et al. [41] observed a relatively lower level of TG in the *Ae. aegypti* female fat body at

around 30 h PBM, while our study of whole body lipidome showed highest level of TG at 30

h PBM.  The differences may reflect the transportation of lipids from the fat body or midgut

into oocytes during the insect reproductive process through lipophorin, a major lipid-carrying

protein [45]. *Ae. aegypti* and some other Diptera predominantly transport TG via lipophorin,

not DG as is the case of most other insects [46-48]. Moreover, a general observation for

insects is that the midgut can convert DG to TG in the presence of high concentrations of

potentially toxic fatty acids; this conversion could maintain low intracellular levels of DG

and fatty acids to prevent lipotoxicity [20,49,50]. Additionally, there are other possibilities for

lipid movement not based on DG transport including evolutionarily conserved mechanisms

for movement of fatty acids [51].

Lyso-glycerophospholipids were highlighted as a lipid group of interest by the present study

due to their unusual temporal pattern in abundance.  Lyso-glycerophospholipids have diverse

functions and they are also found in cell membranes [52]; as precursor of phospholipid

biosynthesis, they may act as signal molecules involved in a broad range of biological

processes [53,54]. However, they might also reflect a highly regulated reservoir of lipids with

accessible utility and concentration-dependent lipid toxicity liability [55], which can result

from lipid overload generated by dietary excess [56] as might be the case when mosquitoes

blood feed. LPE and LPC incorporate into membranes and change membrane permeability

and stability. The change in permeability and stability can result in membrane toxicity unless

concentrations of lyso-glycerophospholipids are kept below a cytotoxic limit [55], which may

explain the stability of these two lipid groups. Also, we found substantial increases in cell

membrane lipids between 16 and 30 h PBM as in the cases of PE and PC, reflecting the

growth of embryos during the reproductive process. Moreover, lyso-glycerophospholipids

could act as a collective reserve which provides acyl chains for beta-oxidation when needed

[51], or they could be a source of partially assembled molecules for further biosynthesis,

which may explain the unusual increase of LPI between 0 and 16 h PBM followed by

increase of nearly all major lipid groups from 16 to 30 h PBM. LPS and LPG also increase

between 0 and 16 h PBM, which might indicate the growth of gut bacteria during the blood

digestion process [57]. LPA is the only group from lyso-glycerophospholipids showing a

bimodal pattern with increases, and these lipids are essential signalling molecules [58].

We also found that many lipids from groups PI and CL increased dramatically from 16 to 30

h PBM. These lipids are closely connected with metabolic activity and signalling. For

example, PIs are important in regulating phototransduction, cell growth and the

developmental process of fruit flies [59-61]. And CLs are integral to mitochondrial membranes,

with important roles in ATP-synthase and phospholipid remodelling [62,63]. Moreover, SM

play significant roles in mammal cells being involved in diverse signalling pathways [64,65],

whereas for dipteran insects such as *Drosophila*, there is a lack of SM and instead they

synthesize the SM analogue phosphoethanolamine ceramide [66]. Though we did not look

into phosphoethanolamine ceramide in the present study, we also note that most SMs were

either relatively constant or changed only slightly between 16 h and 30 h PBM.

Finally, it is worth noting that approximately a quarter of the lipids, including almost all

SHexCer we identified, contained odd chains. Pinch et al. [41] investigated the fat body of

mosquitoes feeding on bovine blood and considered that odd-chain lipids may come from

ruminal microflora or microbes in larval rearing water or branched chain amino acids from

haemoglobin. Our study showed that the last hypothesis is most likely correct, as we

observed these odd-chain lipids from human blood and these also increased in mosquitoes

between 16 h and 30 h PBM. These odd chain fatty acid-containing lipids did not show any

distinct change patterns comparable to even chain fatty acid-containing lipids from the same

lipid group.

**Supplementary Materials**

Supplementary Material 1. Ovarian appearance of female mosquitoes dissected at (A) BBM,

(B) 1 d PBM, (C) 2 d PBM, (D) 3 d PBM, (E) 4 d PBM and (F) 5 d PBM. There is some

variation in the stages of development of the ovaries among individuals as captured by

variation in weight (Fig. 1).

Supplementary Material 2. List of individual lipids and their changes before and after female

*Aedes aegypti* mosquitoes blood fed (between BBM and 0 h PBM).

Supplementary Material 3. List of individual lipids and their changes between 16 hours and

30 hours after female *Aedes aegypti* mosquitoes blood fed (between 16 h and 30 h PBM).

**Funding**

This research was funded by the National Health and Medical Research Council (1132412,

1118640, www.nhmrc.gov.au). The funders had no role in the study design, data collection

and analysis, decision to publish, or preparation of the manuscript.

**Ethics statement**

Female mosquitoes were blood fed on a volunteer in this experiment, with ethics approval

from the University of Melbourne Human Ethics committee (approval 0723847).

**Acknowledgment**

We thank Cameron Patrick from Statistical Consulting Centre and Melbourne Statistical

Consulting Platform, the University of Melbourne, for statistical advice. We also thank

Perran A. Ross and Moshe Jasper for their assistance and advice.

**Reference**

1.    Gould, E.A.; Higgs, S. Impact of climate change and other factors on emerging
arbovirus diseases. *Trans R Soc Trop Med Hyg* **2009**, *103*, 109-121.
2.    Bhatt, S.; Gething, P.W.; Brady, O.J.; Messina, J.P.; Farlow, A.W.; Moyes, C.L.;
Drake, J.M.; Brownstein, J.S.; Hoen, A.G.; Sankoh, O. The global distribution and
burden of dengue. *Nature* **2013**, *496*, 504-507.
3.    Brown, J.E.; Evans, B.R.; Zheng, W.; Obas, V.; Barrera-Martinez, L.; Egizi, A.; Zhao,
H.; Caccone, A.; Powell, J.R. Human impacts have shaped historical and recent
evolution in *Aedes aegypti*, the dengue and yellow fever mosquito. *Evolution* **2014**, *68*,
514-525, doi:10.1111/evo.12281.
4.    Gulia-Nuss, M.; Elliot, A.; Brown, M.R.; Strand, M.R. Multiple factors contribute to
anautogenous reproduction by the mosquito *Aedes aegypti*. *J Insect Physiol* **2015**, *82*,
8-16, doi:10.1016/j.jinsphys.2015.08.001.
5.    Scott, T.W.; Naksathit, A.; Day, J.F.; Kittayapong, P.; Edman, J.D. A fitness
advantage for *Aedes aegypti* and the viruses it transmits when females feed only on

human blood. *Am J Trop Med Hyg* **1997**, *57*, 235-239,
doi:10.4269/ajtmh.1997.57.235.
6.  Attardo, G.M.; Hansen, I.A.; Raikhel, A.S. Nutritional regulation of vitellogenesis in
mosquitoes: implications for anautogeny. *Insect Biochem Mol Biol* **2005**, *35*, 661-675.
7.  Hansen, I.A.; Attardo, G.M.; Park, J.H.; Peng, Q.; Raikhel, A.S. Target of rapamycin-
mediated amino acid signaling in mosquito anautogeny. *Proc Natl Acad Sci U S A*
**2004**, *101*, 10626-10631, doi:10.1073/pnas.0403460101.
8.  Geoghegan, V.; Stainton, K.; Rainey, S.M.; Ant, T.H.; Dowle, A.A.; Larson, T.;
Hester, S.; Charles, P.D.; Thomas, B.; Sinkins, S.P. Perturbed cholesterol and
vesicular trafficking associated with dengue blocking in *Wolbachia*-infected *Aedes*
*aegypti* cells. *Nat Commun* **2017**, *8*, 526, doi:10.1038/s41467-017-00610-8.
9.  Price, D.P.; Nagarajan, V.; Churbanov, A.; Houde, P.; Milligan, B.; Drake, L.L.;
Gustafson, J.E.; Hansen, I.A. The fat body transcriptomes of the yellow fever
mosquito *Aedes aegypti*, pre- and post- blood meal. *PLoS One* **2011**, *6*, e22573,
doi:10.1371/journal.pone.0022573.
10.  Phasomkusolsil, S.; Tawong, J.; Monkanna, N.; Pantuwatana, K.; Damdangdee, N.;
Khongtak, W.; Kertmanee, Y.; Evans, B.P.; Schuster, A.L. Maintenance of mosquito
vectors: effects of blood source on feeding, survival, fecundity, and egg hatching rates.
*J Vector Ecol* **2013**, *38*, 38-45, doi:10.1111/j.1948-7134.2013.12006.x.
11.  Ross, P.A.; Lau, M.-J.; Hoffmann, A.A. Does membrane feeding compromise the
quality of *Aedes aegypti* mosquitoes? *PLoS One* **2019**, *14*, e0224268.
12.  Dutra, H.L.C.; Rodrigues, S.L.; Mansur, S.B.; de Oliveira, S.P.; Caragata, E.P.;
Moreira, L.A. Development and physiological effects of an artificial diet for
*Wolbachia*-infected *Aedes aegypti*. *Sci Rep* **2017**, *7*, 15687, doi:10.1038/s41598-017-
16045-6.
13.  Talyuli, O.A.; Bottino-Rojas, V.; Taracena, M.L.; Soares, A.L.; Oliveira, J.H.;
Oliveira, P.L. The use of a chemically defined artificial diet as a tool to study *Aedes*
*aegypti* physiology. *J Insect Physiol* **2015**, *83*, 1-7,
doi:10.1016/j.jinsphys.2015.11.007.
14.  Caragata, E.P.; Rancès, E.; Hedges, L.M.; Gofton, A.W.; Johnson, K.N.; O'Neill, S.L.;
McGraw, E.A. Dietary cholesterol modulates pathogen blocking by *Wolbachia*. *PLoS*
*Pathog* **2013**, *9*, e1003459, doi:10.1371/journal.ppat.1003459.
15.  Caragata, E.P.; Rancès, E.; O'Neill, S.L.; McGraw, E.A. Competition for amino acids
between *Wolbachia* and the mosquito host, *Aedes aegypti*. *Microb Ecol* **2014**, *67*,
205-218, doi:10.1007/s00248-013-0339-4.
16.  Ziegler, R.; Ibrahim, M.M. Formation of lipid reserves in fat body and eggs of the
yellow fever mosquito, *Aedes aegypti*. *J Insect Physiol* **2001**, *47*, 623-627,
doi:10.1016/s0022-1910(00)00158-x.
17.  Briegel, H.; Hefti, M.; DiMarco, E. Lipid metabolism during sequential gonotrophic
cycles in large and small female *Aedes aegypti*. *J Insect Physiol* **2002**, *48*, 547-554,
doi:10.1016/s0022-1910(02)00072-0.
18.  Troy, S.; Anderson, W.A.; Spielman, A. Lipid content of maturing ovaries of *Aedes*
*aegypti* mosquitoes. *Comp Biochem Physiol B* **1975**, *50*, 457-461, doi:10.1016/0305-
0491(75)90258-8.
19.  Vrablik, T.L.; Watts, J.L. Polyunsaturated fatty acid derived signaling in reproduction
and development: insights from Caenorhabditis elegans and Drosophila melanogaster.
*Mol Reprod Dev* **2013**, *80*, 244-259, doi:10.1002/mrd.22167.
20.  Toprak, U.; Hegedus, D.; Doğan, C.; Güney, G. A journey into the world of insect
lipid metabolism. *Arch Insect Biochem Physiol* **2020**, *104*, e21682,
doi:10.1002/arch.21682.

21. Zhou, G.; Pennington, J.E.; Wells, M.A. Utilization of pre-existing energy stores of
female *Aedes aegypti* mosquitoes during the first gonotrophic cycle. *Insect Biochem*
*Mol Biol* **2004**, *34*, 919-925, doi:10.1016/j.ibmb.2004.05.009.
22. Zhou, G.; Flowers, M.; Friedrich, K.; Horton, J.; Pennington, J.; Wells, M.A.
Metabolic fate of [14C]-labeled meal protein amino acids in *Aedes aegypti*
mosquitoes. *J Insect Physiol* **2004**, *50*, 337-349, doi:10.1016/j.jinsphys.2004.02.003.
23. Brown, M.R.; Clark, K.D.; Gulia, M.; Zhao, Z.; Garczynski, S.F.; Crim, J.W.;
Suderman, R.J.; Strand, M.R. An insulin-like peptide regulates egg maturation and
metabolism in the mosquito *Aedes aegypti*. *Proc Natl Acad Sci U S A* **2008**, *105*,
5716-5721, doi:10.1073/pnas.0800478105.
24. Hansen, I.A.; Attardo, G.M.; Rodriguez, S.D.; Drake, L.L. Four-way regulation of
mosquito yolk protein precursor genes by juvenile hormone-, ecdysone-, nutrient-,
and insulin-like peptide signaling pathways. *Front Physiol* **2014**, *5*, 103,
doi:10.3389/fphys.2014.00103.
25. Belles, X.; Piulachs, M.D. Ecdysone signalling and ovarian development in insects:
from stem cells to ovarian follicle formation. *Biochim Biophys Acta* **2015**, *1849*, 181-
186, doi:10.1016/j.bbagrm.2014.05.025.
26. Klowden, M.J. Endocrine aspects of mosquito reproduction. *Arch Insect Biochem*
*Physiol* **1997**, *35*, 491-512.
27. Uchida, K.; Ohmori, D.; Yamakura, F.; Suzuki, K. Changes in free amino acid
concentration in the hemolymph of the female *Culex pipiens* pallens (Diptera:
Culicidae), after a blood meal. *J Med Entomol* **1990**, *27*, 302-308.
28. Briegel, H. Metabolic relationship between female body size, reserves, and fecundity
of *Aedes aegypti*. *J Insect Physiol* **1990**, *36*, 165-172.
29. Lydic, T.A.; Townsend, S.; Adda, C.G.; Collins, C.; Mathivanan, S.; Reid, G.E. Rapid
and comprehensive 'shotgun' lipidome profiling of colorectal cancer cell derived
exosomes. *Methods* **2015**, *87*, 83-95, doi:10.1016/j.ymeth.2015.04.014.
30. Tsugawa, H.; Cajka, T.; Kind, T.; Ma, Y.; Higgins, B.; Ikeda, K.; Kanazawa, M.;
VanderGheynst, J.; Fiehn, O.; Arita, M. MS-DIAL: data-independent MS/MS
deconvolution for comprehensive metabolome analysis. *Nat Methods* **2015**, *12*, 523-
526, doi:10.1038/nmeth.3393.
31. Tsugawa, H.; Ikeda, K.; Takahashi, M.; Satoh, A.; Mori, Y.; Uchino, H.; Okahashi, N.;
Yamada, Y.; Tada, I.; Bonini, P.; et al. A lipidome atlas in MS-DIAL 4. *Nat*
*Biotechnol* **2020**, *38*, 1159-1163, doi:10.1038/s41587-020-0531-2.
32. Liebisch, G.; Fahy, E.; Aoki, J.; Dennis, E.A.; Durand, T.; Ejsing, C.S.; Fedorova, M.;
Feussner, I.; Griffiths, W.J.; Köfeler, H.; et al. Update on LIPID MAPS classification,
nomenclature, and shorthand notation for MS-derived lipid structures. *J Lipid Res*
**2020**, *61*, 1539-1555, doi:10.1194/jlr.S120001025.
33. Oshaghi, M.A.; Chavshin, A.R.; Vatandoost, H.; Yaaghoobi, F.; Mohtarami, F.;
Noorjah, N. Effects of post-ingestion and physical conditions on PCR amplification of
host blood meal DNA in mosquitoes. *Exp Parasitol* **2006**, *112*, 232-236,
doi:10.1016/j.exppara.2005.11.008.
34. Mukabana, W.R.; Takken, W.; Seda, P.; Killeen, G.F.; Hawley, W.A.; Knols, B.G.
Extent of digestion affects the success of amplifying human DNA from blood meals
of *Anopheles gambiae* (Diptera: Culicidae). *Bull Entomol Res* **2002**, *92*, 233-239,
doi:10.1079/ber2002164.
35. Harrington, L.C.; Edman, J.D.; Scott, T.W. Why do female *Aedes aegypti* (Diptera:
Culicidae) feed preferentially and frequently on human blood? *J Med Entomol* **2001**,
*38*, 411-422, doi:10.1603/0022-2585-38.3.411.

36. Roy, D. On the role of blood in ovulation in *Aedes aegypti*, Linn. *Bull Entomol Res* **1936**, *27*, 423-429.

37. Arifin, S.A.; Falasca, M. Lysophosphatidylinositol Signalling and Metabolic Diseases. *Metabolites* **2016**, *6*, doi:10.3390/metabo6010006.

38. Piñeiro, R.; Falasca, M. Lysophosphatidylinositol signalling: new wine from an old bottle. *Biochim Biophys Acta* **2012**, *1821*, 694-705, doi:10.1016/j.bbalip.2012.01.009.

39. Kim, T.; Kim, Y.J. Overview of innate immunity in *Drosophila*. *J Biochem Mol Biol* **2005**, *38*, 121-127, doi:10.5483/bmbrep.2005.38.2.121.

40. Sohlenkamp, C.; Galindo-Lagunas, K.A.; Guan, Z.; Vinuesa, P.; Robinson, S.; Thomas-Oates, J.; Raetz, C.R.; Geiger, O. The lipid lysyl-phosphatidylglycerol is present in membranes of Rhizobium tropici CIAT899 and confers increased resistance to polymyxin B under acidic growth conditions. *Mol Plant Microbe Interact* **2007**, *20*, 1421-1430, doi:10.1094/mpmi-20-11-1421.

41. Pinch, M.; Mitra, S.; Rodriguez, S.D.; Li, Y.; Kandel, Y.; Dungan, B.; Holguin, F.O.; Attardo, G.M.; Hansen, I.A. Fat and happy: profiling mosquito fat body lipid storage and composition post-blood meal. *Front Insect Sci* **2021**, *1*, doi:10.3389/finsc.2021.693168.

42. Hou, Y.; Wang, X.L.; Saha, T.T.; Roy, S.; Zhao, B.; Raikhel, A.S.; Zou, Z. Temporal Coordination of Carbohydrate Metabolism during Mosquito Reproduction. *PLoS Genet* **2015**, *11*, e1005309, doi:10.1371/journal.pgen.1005309.

43. Hagedorn, H.H.; Fallon, A.M.; Laufer, H. Vitellogenin synthesis by the fat body of the mosquito *Aedes aegypti*: evidence of transcriptional control. *Dev Biol* **1973**, *31*, 285-294, doi:10.1016/0012-1606(73)90265-0.

44. Sun, J.; Hiraoka, T.; Dittmer, N.T.; Cho, K.H.; Raikhel, A.S. Lipophorin as a yolk protein precursor in the mosquito, *Aedes aegypti*. *Insect Biochem Mol Biol* **2000**, *30*, 1161-1171, doi:10.1016/s0965-1748(00)00093-x.

45. Kawooya, J.K.; Law, J.H. Role of lipophorin in lipid transport to the insect egg. *J Biol Chem* **1988**, *263*, 8748-8753.

46. Ford, P.S.; Van Heusden, M.C. Triglyceride-rich lipophorin in *Aedes aegypti* (*Diptera: Culicidae*). *J Med Entomol* **1994**, *31*, 435-441, doi:10.1093/jmedent/31.3.435.

47. Pennington, J.E.; Nussenzveig, R.H.; Van Heusden, M.C. Lipid transfer from insect fat body to lipophorin: comparison between a mosquito triacylglycerol-rich lipophorin and a sphinx moth diacylglycerol-rich lipophorin. *J Lipid Res* **1996**, *37*, 1144-1152.

48. Arrese, E.L.; Canavoso, L.E.; Jouni, Z.E.; Pennington, J.E.; Tsuchida, K.; Wells, M.A. Lipid storage and mobilization in insects: current status and future directions. *Insect Biochem Mol Biol* **2001**, *31*, 7-17, doi:10.1016/s0965-1748(00)00102-8.

49. Canavoso, L.E.; Wells, M.A. Metabolic pathways for diacylglycerol biosynthesis and release in the midgut of larval *Manduca sexta*. *Insect Biochem Mol Biol* **2000**, *30*, 1173-1180, doi:10.1016/s0965-1748(00)00094-1.

50. Canavoso, L.E.; Jouni, Z.E.; Karnas, K.J.; Pennington, J.E.; Wells, M.A. Fat metabolism in insects. *Annu Rev Nutr* **2001**, *21*, 23-46, doi:10.1146/annurev.nutr.21.1.23.

51. Black, P.N.; DiRusso, C.C. Transmembrane movement of exogenous long-chain fatty acids: proteins, enzymes, and vectorial esterification. *Microbiol Mol Biol Rev* **2003**, *67*, 454-472, table of contents, doi:10.1128/mmbr.67.3.454-472.2003.

52. Rietschel, E.T.; Kirikae, T.; Schade, F.U.; Mamat, U.; Schmidt, G.; Loppnow, H.; Ulmer, A.J.; Zähringer, U.; Seydel, U.; Di Padova, F.; et al. Bacterial endotoxin: molecular relationships of structure to activity and function. *Faseb j* **1994**, *8*, 217-225, doi:10.1096/fasebj.8.2.8119492.

53. Ishii, I.; Fukushima, N.; Ye, X.; Chun, J. Lysophospholipid receptors: signaling and
biology. *Annu Rev Biochem* **2004**, *73*, 321-354,
doi:10.1146/annurev.biochem.73.011303.073731.
54. Tan, S.T.; Ramesh, T.; Toh, X.R.; Nguyen, L.N. Emerging roles of lysophospholipids
in health and disease. *Prog Lipid Res* **2020**, *80*, 101068,
doi:10.1016/j.plipres.2020.101068.
55. Arouri, A.; Mouritsen, O.G. Membrane-perturbing effect of fatty acids and lysolipids.
*Prog Lipid Res* **2013**, *52*, 130-140, doi:10.1016/j.plipres.2012.09.002.
56. Leventhal, K.R.; Malmberg, E.; Symons, J.L.; Fan, Y.Y.; Chapkin, R.S.; Ernst, R.;
Leventhal, I. Lipidomic and biophysical homeostasis of mammalian membranes
counteracts dietary lipid perturbations to maintain cellular fitness. *Nat Commun* **2020**,
*11*, 1339, doi:10.1038/s41467-020-15203-1.
57. Minard, G.; Mavingui, P.; Moro, C.V. Diversity and function of bacterial microbiota
in the mosquito holobiont. *Parasit Vectors* **2013**, *6*, 146, doi:10.1186/1756-3305-6-
146.
58. Luquain, C.; Sciorra, V.A.; Morris, A.J. Lysophosphatidic acid signaling: how a small
lipid does big things. *Trends Biochem Sci* **2003**, *28*, 377-383, doi:10.1016/s0968-
0004(03)00139-7.
59. Gupta, A.; Toscano, S.; Trivedi, D.; Jones, D.R.; Mathre, S.; Clarke, J.H.; Divecha, N.;
Raghu, P. Phosphatidylinositol 5-phosphate 4-kinase (PIP4K) regulates TOR
signaling and cell growth during *Drosophila* development. *Proc Natl Acad Sci U S A*
**2013**, *110*, 5963-5968, doi:10.1073/pnas.1219333110.
60. Milligan, S.C.; Alb, J.G., Jr.; Elagina, R.B.; Bankaitis, V.A.; Hyde, D.R. The
phosphatidylinositol transfer protein domain of *Drosophila* retinal degeneration B
protein is essential for photoreceptor cell survival and recovery from light stimulation.
*J Cell Biol* **1997**, *139*, 351-363, doi:10.1083/jcb.139.2.351.
61. Garcia-Murillas, I.; Pettitt, T.; Macdonald, E.; Okkenhaug, H.; Georgiev, P.; Trivedi,
D.; Hassan, B.; Wakelam, M.; Raghu, P. lazaro encodes a lipid phosphate
phosphohydrolase that regulates phosphatidylinositol turnover during *Drosophila*
phototransduction. *Neuron* **2006**, *49*, 533-546, doi:10.1016/j.neuron.2006.02.001.
62. Schlame, M.; Ren, M. The role of cardiolipin in the structural organization of
mitochondrial membranes. *Biochim Biophys Acta* **2009**, *1788*, 2080-2083,
doi:10.1016/j.bbamem.2009.04.019.
63. Acehan, D.; Malhotra, A.; Xu, Y.; Ren, M.; Stokes, D.L.; Schlame, M. Cardiolipin
affects the supramolecular organization of ATP synthase in mitochondria. *Biophys J*
**2011**, *100*, 2184-2192, doi:10.1016/j.bpj.2011.03.031.
64. Hannun, Y.A.; Obeid, L.M. Principles of bioactive lipid signalling: lessons from
sphingolipids. *Nat Rev Mol Cell Biol* **2008**, *9*, 139-150, doi:10.1038/nrm2329.
65. Hannun, Y.A.; Bell, R.M. Functions of sphingolipids and sphingolipid breakdown
products in cellular regulation. *Science* **1989**, *243*, 500-507,
doi:10.1126/science.2643164.
66. Acharya, U.; Acharya, J.K. Enzymes of sphingolipid metabolism in *Drosophila*
*melanogaster*. *Cell Mol Life Sci* **2005**, *62*, 128-142, doi:10.1007/s00018-004-4254-1.
594