Gene Novelties in Amphioxus Illuminate the Early Evolution of Cephalochordates

Authors: Qing Xiong[1,2,3]†, Kevin Yi Yang[1,2]†, Xi Zeng[4]†, Mingqiang Wang[1,5]†, Patrick Kwok-Shing Ng[1], Jun-Wei Zhou[1], Judy Kin-Wing Ng[1], Cherie Tsz-Yiu Law[1], Qiao Du[4], Kejin Xu[4], Bingyu Mao[6], Stephen Kwok-Wing Tsui[1,2,7] *

Affiliations:
School of Biomedical Sciences, The Chinese University of Hong Kong, Hong Kong.
Hong Kong Bioinformatics Centre, The Chinese University of Hong Kong, Hong Kong.
Department of Health Technology and Informatics, Faculty of Health and Social Science, The Hong Kong Polytechnic University, Hong Kong.
Agricultural Bioinformatics Key Laboratory of Hubei Province, 3D Genomics Research Centre, College of Informatics, Huazhong Agricultural University, Wuhan, China.
Stanford Cardiovascular Institute, Stanford University School of Medicine, Stanford, CA, 94305, USA.
State Key Laboratory of Genetic Resources and Evolution, Kunming Institute of Zoology, Chinese Academy of Sciences, 650223, Kunming, Yunnan, China.
Centre for Microbial Genomics and Proteomics, The Chinese University of Hong Kong, Hong Kong.

* Correspondence to Professor Stephen Kwok-Wing Tsui, School of Biomedical Sciences, The Chinese University of Hong Kong, Shatin, N.T., Hong Kong. E-mail: kwtsui@cuhk.edu.hk; Tel: 852-3943 6381; Fax: 852-2603 5123.

† These authors contributed equally to this work.

Abstract

Amphioxus is considered the best-known living proxy to the chordate ancestor and an irreplaceable model organism for evolutionary studies of chordates and deuterostomes. In this study, a high-quality genome of the Beihai amphioxus, *Branchiostoma belcheri* beihai, was *de novo* assembled and annotated. Within four amphioxus genomes, twenty-eight groups of gene novelties were identified, revealing new genes that lack homologs in non-deuterostome metazoa, but share unexpectedly high similarities with those from non-metazoan species. These gene innovation events have played roles in amphioxus adaptations, including innate immunity responses, glycolysis, and regulation of calcium balance. The gene novelties related to innate immunity, such as a group of lipoxygenases and a DEAD-box helicase, boosted amphioxus immune responses. The novel genes for alcohol dehydrogenase and ferredoxin could aid in the glycolysis of amphioxus. A proximally arrayed cluster of EF-hand calcium-binding protein genes were identified to resemble those of bacteria. The copy number of this gene cluster was negatively correlated to the sea salinity of the collection region, suggesting that it may enhance their survival at different calcium concentrations. This comprehensive study collectively reveals insights into adaptive evolution of cephalochordates and provides valuable resources for research on early evolution of deuterostomes.

Keywords

amphioxus, cephalochordate, deuterostome, gene novelty, genome evolution

Background

Amphioxus, a group of non-vertebrate chordates within Cephalochordata, diverged from the

common ancestor of other two chordate groups, urochordates (tunicates) and vertebrates,

around 550 million years ago (Bertrand and Escriva, 2011; Bourlat et al., 2006; Delsuc et al.,

2006). Compared with tunicates, which were considered as a rapidly-evolving group (Bourlat

et al., 2006), amphioxus shares a basic chordate body plan, undergoes slower molecular

evolution (Benito-Gutiérrez et al., 2021; Putnam et al., 2008). Therefore, amphioxus is

considered an irreplaceable model organism for evolutionary studies. With the advent of

high-throughput sequencing, genome-based investigation has revolutionized the study of

amphioxus evolution and greatly expanded our knowledge of the origin and evolution of

vertebrates (Acemel et al., 2016; Brasó-Vives et al., 2022a; Holland et al., 2008; Huang et al.,

2014; Huang et al., 2023; Marlétaz et al., 2018; Putnam et al., 2008; Simakov et al., 2020).

Whole-genome comparison of the three chordate groups provided evidence supporting the 2R

hypothesis, which posits that two rounds of whole-genome duplication occurred in the early

vertebrate lineage (Holland et al., 2008; Putnam et al., 2008). The application of functional

genomics to amphioxus has provided a wealth of information on the origins of the complex

gene regulatory landscapes that are characteristic of vertebrates, shedding new light on the

evolutionary history of this important group of animals (Marlétaz et al., 2018).

In this study, we successfully constructed a high-quality genome of the Beihai amphioxus,

Branchiostoma (*B.*) *belcheri* beihai, which was collected from the nearshore area of South

China. Although amphioxus is widely regarded as the most basally divergent group of

chordates and lacks the specializations and genome innovations of vertebrates (Escriva, 2018;

Holland, 2017; Kassahn et al., 2009; Louis et al., 2012), we performed a comparative

genomic analysis of amphioxus and identified a number of gene novelties, where the genes

involved markedly differ from those found in other non-deuterostome metazoa, but

surprisingly exhibit similarities to those found in non-metazoan species such as bacteria,

fungi, and plants. Our analysis suggests that these novel genes may play crucial roles in a

variety of amphioxus adaptations, including innate immunity, glycolysis, and calcium

balance. Overall, our comprehensive genomic analysis of the gene novelties in amphioxus

provided insights into the genetic mechanisms that underlie the adaptations and

diversification of chordates and deuterostomes.

Methods

Sample collection

The adults of the Beihai amphioxus, *B. belcheri* beihai, were obtained from the sea near Dianbai District, Maoming City, Guangdong Province, China, cultured at 24–28 °C with air-pumped circulating artificial seawater in Beihai Marine Station of Nanjing University in Beihai City, Guangxi Province, China, and fed with seawater and sea alga (Chen et al., 2009; Du et al., 2023; Yang et al., 2016). This *B. belcheri* beihai amphioxus population has been sustained for several years to attain a low degree of polymorphism.

Genomic DNA extraction and sequencing

To obtain the genome information of *B. belcheri* Beihai, genomic DNA was extracted from a pool of amphioxus samples and subjected to both PacBio long-read and Illumina short-read sequencing. Genomic DNAs were extracted using the Qiagen Blood & Cell Culture DNA Maxi Kit (Qiagen, Germany). Firstly, adult bodies of the Beihai amphioxus were washed twice with phosphate-buffered saline (PBS, pH 7.4), immediately frozen in liquid nitrogen, and then homogenized into powder form using a mortar and pestle with liquid nitrogen to maintain a low temperature. The homogenized samples were then incubated at 50 °C for proteolysis following the manufacturer's protocol. Genomic DNA was bound to the column, washed, and eluted by washing buffer and elution buffer respectively, and precipitated in 70% ethanol. Finally, air-dry pellets of genomic DNA were dissolved overnight in UltraPure™ DNase/RNase-free distilled water (Thermo Fisher Scientific, USA) overnight at room

temperature. The integrity of genomic DNA was determined by electrophoresis in 0.5%

agarose gel and analyzed by Agilent 2100 Bioanalyzer (Agilent Technologies, USA), and the

quantity was detected by Nanodrop 2000 spectrophotometer (Thermo Fisher Scientific, USA)

and Qubit Fluorometer (Thermo Fisher Scientific, USA). The genomic DNA of *B. belcheri*

beihai was sequenced by Illumina HiSeq 2000 in 500-bp and 3,000-bp libraries to generate

NGS reads and PacBio RS system for third-generation long reads.

De novo genome assembly and annotation

The initial genome assembly of *B. belcheri* beihai was constructed with PacBio long reads

using Canu v1.8 (Koren et al., 2017). Then, scaffolding was performed by SSPACE Basic

v2.0 (Boetzer et al., 2011) with paired-end Illumina short reads (Bentley et al., 2008) and by

SSPACE LongRead v1.1 (Boetzer and Pirovano, 2014) with PacBio long reads (Eid et al.,

2009). The final sequence polishing was finished by Pilon v1.22 (Walker et al., 2014) with all

Illumina short reads to generate the final genome assembly. The genome assembly was

assessed by QUAST v5.0.2 (Gurevich et al., 2013) in continuity and the completeness of

genome assembly was assessed by BUSCO v3.1.0 (Simão et al., 2015) with database

metazoa_odb9.

With genome assembled, genome annotation was performed. Firstly, repeat masking was

performed by *de novo* prediction with RepeatModeler v2.0.1 (Flynn et al., 2020) and masking

with RepeatMasker v4.0.8 (RepBase edition 20181026) (Tarailo-Graovac and Chen, 2009).

In the *de novo* prediction with RepeatModeler v2.0.1 (Flynn et al., 2020), RECON v1.05

(Bao and Eddy, 2002) and RepeatScout v1.0.6 (Price et al., 2005) were used for the

prediction of repeat family in the genome. Then, genome annotation was performed by Maker

pipeline v2.31 (Cantarel et al., 2008). In the Maker pipeline, transcriptome assemblies and

homologous proteins were used as evidence for alignment by Exonerate v2.4.0 (Slater and

Birney, 2005), whilst gene prediction was completed by SNAP (lib v2017-03-01) (Korf,

2004), GeneMark v4.38 (Borodovsky and Lomsadze, 2011) and Augustus v3.3.1 (Hoff and

Stanke, 2019). The quality of genome annotation was assessed by BUSCO v3.1.0 (Simão et

al., 2015) with database metazoa_odb9.

Data collection and access

Besides of the *de novo* assembled and annotated genome of *B. belcheri* beihai, genome

assemblies and annotations of other three amphioxus were downloaded from NCBI database

(NCBI Assembly accession GCF_001625305.1 for *B. belcheri* xiamen, GCF_000003815.2

for *B. floridae* and GCA_900088365.1 for *B. lanceolatum*). The genome sequencing data,

assembly, and annotation of *B. belcheri* beihai have been uploaded to NCBI database under

the BioProject accession: PRJNA804338. The transcriptome data of *B. belcheri* beihai were

from the BioProject accession: PRJNA310680.

Despite the recent reports of several high-quality *B. lanceolatum* genomes (Brasó-Vives et

al., 2022b), the absence of transcriptome data hindered our ability to use the genome in

manual curation. We did not include other chromosome-level amphioxus genomes produced

from hybrid breeding due to concerns about the phasing step and subsequent transcriptome

analysis (Huang et al., 2023).

Phylogenetic analysis of deuterostomes

The genome sequences of deuterostome species were phylogenetically analyzed with those of other species. The overlapped single and complete BUSCO protein sequences were extracted by BUSCO v3.1.0 (Simão et al., 2015) with database metazoa_odb9, then aligned by MAFFT (Katoh et al., 2002) and edited in Gblocks (Castresana, 2000) with the option '-t=p' to generate sequence alignment of conserved amino-acid residues. Finally, the sequence alignment was used to construct the phylogenetic tree in maximum likelihood algorithm and 100 bootstrap replicates by RAxML v8.2.12 (Stamatakis, 2014) with the options '-m PROTCATWAG -f a -# 100'. The phylogenetic tree was edited by the online tool Interactive Tree of Life (iTOL) (Letunic and Bork, 2021).

To construct a ultrametric time tree, the alignment of codon sequences of conserved amino-acid residues were extracted and used for estimating divergence time by the MCMCTree program from the PAML package v4.9j (Yang, 2007). The ultrametric time tree was final edited by the online tool Interactive Tree of Life (iTOL) (Letunic and Bork, 2021).

Phylogenomic orthology analysis

Phylogenomic orthology and gene gain/loss analysis (Hahn et al., 2005) was performed among protein sequences in 21 annotated genomes by OrthoFinder v2.5.4 (Emms and Kelly, 2015,

2019) and CAFÉ v4.2 (De Bie et al., 2006) based on the ultrametric time tree. All the protein

sequences were assigned into orthogroups according to the protein similarities on sequence

level. These orthogroups were also considered as gene families. Specific orthogroups were

summarized and further analyzed. Functional clusters of orthologous groups (COGs) of genes

in specific orthogroups were performed by eggnog-mapper v2.1.5 (Cantalapiedra et al., 2021;

Huerta-Cepas et al., 2019).

Identification of gene novelties

Our study primarily focuses on exploring the gene novelties of amphioxus within the

framework of deuterostomes. Specifically, we concentrated on genes that exhibit notable

differences from those present in other non-deuterostome metazoans, while also sharing

similarities with genes from non-metazoan species, including bacteria, fungi, plants, and

other eukaryotes. The whole workflow can be seen in Fig. S3.

To identify gene novelties in amphioxus, we compared the best hits of all the annotated

proteins of *B. belcheri* beihai to those from different taxonomic groups in UniRef50 database

(UniProt, last updated on Dec. 15[th], 2020) (Suzek et al., 2015) as follows. Although

amphioxus specific orthogroups have been extracted, we input all the annotated proteins of *B.*

belcheri beihai, to avoid missing important gene novelties.

1) In UniRef50 database, collecting the proteins of different taxonomic groups according to

the taxon identifiers, namely non-deuterostome metazoa, bacteria, fungi, plants, and other

eukaryotes (excluding fungi, plants, and metazoa), and building BLAST databases.

2)   Searching the best hits of 44,745 annotated proteins of *B. belcheri* beihai in the five BLAST databases built in Step 1, using BLASTP v2.9.0 (McGinnis and Madden, 2004) with the options '-evalue 1e-6 -max_hsps 1 -max_target_seqs 1 -outfmt 6'.

3)   The best hits in different taxonomic groups were compared and identified two groups of candidates of gene novelties. Firstly, when one protein had hit in the databases of bacteria, fungi, plants, or other eukaryotes, but not that of non-deuterostome metazoan, this gene was identified as group I candidates of gene novelties. Secondly, when the gene novelty index of a protein was calculated as dividing the bit score of the best hit in the databases of bacteria, fungi, plants, and other eukaryotes, by that of non-deuterostome metazoan; if the index ≥ 1.5, this gene was identified as group II candidates. This method has been adapted from the one employed to identify horizontal gene transfer events (Xiong et al., 2022).

4)   Collecting protein sequences of two groups of candidates, performing BLASTP online search in the NCBI non-redundant protein (NR) database (last updated on Nov. 1[st], 2021), and then manually checking the top 100 hits. Except those from Amphioxiformes (including all amphioxus), if the best hits were from unusual taxonomic groups like bacteria, fungi, plants, and other eukaryotes (excluding fungi, plants, and metazoa), this gene was confirmed as an amphioxus specific gene; if some of the best hits were from other deuterostome groups except jawed vertebrates, this gene was considered as a novel gene conserved with other

deuterostomes. Additionally, only the DBH and ADH genes had homologs from the soft coral

Dendronephthya gigantea and the sponge *Amphimedon queenslandica*, respectively (Table

1).

Therefore, the dominantly abundant and well annotated proteins of jawed vertebrates were

used to filter out the candidate genes possibly introduced at stem chordates and

deuterostomes, to avoid repeated study in the previous report on in stem chordates (Holland

et al., 2008) and deuterostomes (Simakov et al., 2015). The absence of the candidate genes in

jawed vertebrates was double confirmed by searching using TBLASTN in the genomic DNA

sequences and BLASTP in the annotated proteins at E-value cutoff of 1E-6. Those candidates

supported by poor hits (e.g., low bit scores or repeated domains) were discarded either.

5) Once a candidate gene was confirmed, homologous genes of four amphioxus were

searched by TBLASTN in the genomic DNA sequences and BLASTP in the annotated

proteins at E-value cutoff of 1E-6. After again checking those homologous genes at Step 4

and filtering out false-positive genes; manual checking and curating based on transcriptome

data, especially in splice sites (locations of exon–intron boundaries) using the program

Integrative Genomics Viewer (IGV) v2.9.4 (Thorvaldsdóttir et al., 2013); finally, all

homologous genes among four amphioxus (Table 1) were collected and further analyzed.

Most candidates were filtered out and the final 28 groups of gene novelties were summarized

in Table 1. Since those involved genes were conserved among up to deuterostome groups

(except the VDCC gene among three groups, Table S3) and often had highly similar genes

from unusual taxonomic groups (Table 1), we excluded the possibility of assembly

contamination and false-positiveness as implausible. For better reproducibility, the protein

sequences coded by these representative genes can be found in Table S4. Some gene novelties

were further validated by phylogenetic analysis.

Transcriptome and gene expression level

To explore the expression level of target genes, we collected the transcriptome data of

different tissues of *B. lanceolatum* as follows. The SRA accessions of the transcriptome data

in the NCBI database are epidermis: SRR6246024; cirri: SRR6246021; gill 1: SRR6246026;

gill 2: SRR6246027; gut 1: SRR6246028; gut 2: SRR6246029; hepatic diverticulum 1:

SRR6246030; hepatic diverticulum 2: SRR6246031; male gonad 1: SRR6246032; male

gonad 2: SRR6246033; female gonad: SRR6246025; muscle 1: SRR6246034; muscle 2:

SRR6246035; neural tube 1: SRR6246036; neural tube 2: SRR6246037. The transcript

sequences of all the target genes and Glyceraldehyde 3-phosphate dehydrogenase (*GAPDH*)

or elongation factor 1 alpha (*EF1A*) of *B. lanceolatum* were built as index and calculated in

transcripts per million (TPM) values by the program Salmon v0.12.0 with the options '--type

quasi -k 31'. The values of log2 (1+ TPM) appeared to be disorganized, likely due to varying

transcriptome data sizes among tissues. This suggests the need for data normalization. To

generate normalized TPM values, the TPM values of all target genes of *B. lanceolatum* were

divided by that of the reference genes, *GAPDH* or *EF1A (Zhang et al., 2016)*, and then the

log2 (1+ normalized TPM) values were calculated and presented in the heatmaps generated

by the program GraphPad Prism v9.0.0.

Comparative analysis of genes

All protein sequences of target genes were searched TBLASTN in the genomic DNA

sequences and BLASTP in the annotated proteins of the four amphioxus genomes at E-value

cutoff of 1E-6, to acquire all the homologs. After manual check and validation based on

transcriptome data, all protein sequences of target genes were collected in four amphioxus.

The sequence alignment of genes was performed by the online tool Clustal Omega (Sievers

and Higgins, 2014). The gene synteny and splice sites (locations of exon–intron boundaries)

of target genes were identified and checked in the program Integrative Genomics Viewer

(IGV) v2.9.4 (Thorvaldsdóttir et al., 2013). The conserved sites were defined with at least

one of two flanking amino acids identical among protein sequences. The domain structure of

target genes was manually drawn based on the online analysis of BLASTP with the NCBI NR

database, InterProScan (Jones et al., 2014) and Pfam search (Mistry et al., 2021). Gene

synteny alignment was manually drawn in PowerPoint 2021, in which the gene distance

below 5 Kb defined as tandemly arrayed and below 20 Kb as proximally arrayed.

In basic phylogenetic analysis of genes, protein sequences were aligned by CLUSTAL W

(Thompson et al., 1994) and MUSCLE (Edgar, 2004), and phylogenetic trees were

constructed by MEGA v10.2.2 (Kumar et al., 2018) with maximum likelihood (ML)

algorithm in the JTT (Jones-Taylor-Thornton) model, 80% site coverage and 100 bootstrap

replicates, and then edited by online tool Interactive Tree of Life (iTOL) (Letunic and Bork,

2006).

To validate the novelty of these genes, another phylogenetic method was adopted as follows.

The protein sequences of the target genes were subjected to phylogenetic analysis with the

best-hit protein sequences of other taxonomical groups, including bacteria, fungi, other

eukaryotes (excluding metazoa, plants, and fungi) and other metazoa (excluding

Amphioxiformes), in UniRef50 database and some sequences in the NCBI GenBank

database. In addition, protein sequences of the multi-copy genes of amphioxus were

phylogenetically analyzed to explore their duplication and divergence. All protein sequences

were aligned by MAFFT (Katoh et al., 2002) and edited in Gblocks (Castresana, 2000) with

the option '-b4=5 -b5=h' to generate sequence alignment of conserved amino-acid residues.

Then, the sequence alignment was used to construct the phylogenetic trees in maximum

likelihood algorithm by IQ-TREE v2.2.0 (Minh et al., 2020) was used with options "-seqtype

AA -T AUTO --merge -rclusterf 10 -m MFP -alrt 1000 -bb 1000 -safe" to generate

phylogenetic trees dually tested by Shimodaira-Hasegawa approximate likelihood-ratio test

(SH-aLRT) and ultrafast bootstrap values. For poorly supported branches with SH-aLRT or

ultrafast bootstrap values below 50 that may cause misinterpretation to the gene origin, the

branches were discarded and manually converted into polytomies. Phylogenetic noise like

long branch attraction effect (Townsend et al., 2012) was carefully excluded using the classic

SAW method (Siddall and Whiting, 1999). The phylogenetic tree was edited by the online

tool Interactive Tree of Life (iTOL) (Letunic and Bork, 2021).

To perform selection pressure analysis for one-to-one orthologs, we utilized the wgd pipeline

(Zwaenepoel and Van de Peer, 2018) and the CodeML program from the PAML package

v4.9j (Yang, 2007) to determine the nonsynonymous-to-synonymous substitution ratio

(*Ka*/*Ks*). The annotated proteins of the genomes of *B. belcheri xiamen* and *B. lanceolatum*

were selected for the analysis, to enable more one-to-one orthologs of the genes (Table 1).

Data availability

The dataset of *B. belcheri* beihai supporting the results of this article is available in the NCBI

BioProject repository, with the accession PRJNA804338.

Results and Discussion

Genome assembly and annotation

A high-quality genome of the Beihai amphioxus, *B. belcheri* beihai (Du et al., 2023; Yang et al., 2016), was *de novo* assembled, annotated, and compared with three other genome assemblies of the Xiamen amphioxus (*B. belcheri* xiamen), Florida amphioxus (*B. floridae*) and European amphioxus (*B. lanceolatum*) (Supplementary Note 1.1, Table S1, Fig. S1). For the genome assembly of *B. belcheri* beihai, the scaffold N50 length (4,185,906 bp) was second only to that of *B. floridae* (25,441,410 bp), the gap content was the lowest among the four amphioxus and the BUSCO completeness was one of the highest (97.2%) (Supplementary Note 1.2, Table S1). In the genome annotation of *B. belcheri* beihai, 44,745 protein-coding genes were annotated with 95.2% BUSCO completeness. The genome of *B. lanceolatum* was annotated with 38,190 protein-coding genes with 91.9% completeness (Supplementary Note 1.3, Table S1).

Phylogeny and comparative genomics

Along with other genome assemblies (Supplementary Note 2.1, Table S2), a phylogenetic analysis was performed, with the sponge *Amphimedon queenslandica* used as the outgroup. The deuterostome phylogeny of this organism was consistent with current opinions (Kapli et al., 2021; Satoh et al., 2014). The phylogeny supported the monophylies of both chordates and deuterostomes, and amphioxus was suggested as the outermost or early branching chordate group (Fig. 1).

To gain insights into the genome evolution of amphioxus, we performed a comprehensive comparative genomics analysis of annotated animal genomes (Table S2). Based on the ultrametric time tree and phylogenomic orthology analysis (Supplementary Note 2.3), we observed significantly more assigned orthogroups in the four amphioxus genomes compared to other animal genomes, including over 2,000 amphioxus-specific orthogroups (Fig. S2). This observation piqued our interest in exploring the gene novelties that underlie the genome evolution of amphioxus. While it would be also interesting to explore the divergence of these amphioxus genomes, these fall outside the scope of this study, which focuses on gene novelties.

Gene novelties

Within four annotated amphioxus genomes (Table S1), we identified a number of gene novelties conserved among them (Table 1), in which the involved genes resemble those from bacteria, fungi, plants or other eukaryotes, and markedly differ from those of other non-deuterostome metazoa (Fig. S3). In addition to cautious filtering and rigorous validation to avoid false positivity in assembled sequences, the high conservation of those genes among four independently collected, sequenced, and assembled genomes of amphioxus (Table 1) precluded the occurrence of contamination. Previous studies have identified a number of gene novelties in stem chordates (Holland et al., 2008) and deuterostomes (Simakov et al., 2015). To avoid repeating these efforts and to focus on novel gene features in amphioxus, we excluded candidate novelties that are also conserved in jawed vertebrates, including the

sialidase gene, which may have arisen in the deuterostome stem (Fig. S4, Supplementary

Note 3.1). In the scope of this study, we also excluded novel genes that may originate from

extreme sequence variation, protein domain recombination or shuffling events.

A gene novelty could generate many novel genes via gene duplications; for example, in *B.*

belcheri beihai, 28 groups of gene novelties generated 108 new genes (Table 1). According to

their biological functions, these involved genes were classified into six categories, namely,

innate immunity, glycolysis, calcium balance, amino acid metabolism, other metabolic genes,

and others without explicit functional domains (Table 1). The expression level of these genes

was analyzed in different tissues of *B. lanceolatum* because of its high-quality transcriptome

data, and all the genes were expressed in at least one tissue, although some gene expression

levels are quite low (Fig. S5). Regardless of the inconsistent gene expression in tissue

duplicates of muscle and male gonads, the most marked expression feature was that the genes

coding for calcium-binding proteins were highly expressed in the guts and hepatic diverticula

(Fig. S5, Supplementary Note 3.2). Subsequently, we explored these novel genes in three

important functional groups, namely innate immunity, glycolysis, and calcium balance. More

gene novelties were discussed in Supplementary Note 3.

Innate immunity

Innate immunity is the first-line nonspecific immune system against infections, and this

immune system is observed in jawed and jawless vertebrates as well as invertebrates (Bartl et

al., 2003). Successful immunity evolution confers survival traits to a species. In the absence

of an adaptative immune system in amphioxus (Bartl et al., 2003), innate immunity is the

most important weapon of amphioxus in response to various pathogens.

Lipoxygenases (LOXs) are important in arachidonic acid metabolism, participate in

inflammatory lipid signaling and further regulate inflammatory and allergic responses (Yuan

and Xu, 2016; Yuan et al., 2014). A group of duplicated LOX genes of amphioxus were

identified similar to those of cyanobacteria or proteobacteria (Table 1). Notably, these LOX

genes have unexpected homologs from only two living groups of jawless vertebrates

(Cyclostomata) (Janvier, 2006), the sea lamprey *Petromyzon marinus* and the hagfish

Eptatretus burger (Fig. S6); however, such homologs are not observed in other metazoa. In

the phylogenetic tree, these LOX genes of amphioxus are close to those of sea lamprey,

bacteria and other eukaryotes (Fig. 2A). Three splice sites (locations of exon–intron

boundaries) were conserved among the LOX genes of four amphioxus and the sea lamprey

(Fig. S6). In the absence of genome annotation, two fragments of the LOX gene were

recovered from the hagfish genome, and three other conserved splice sites were identified

between those of the two jawless vertebrates (Fig. S6).

The phylogenetic analysis revealed that these LOX genes were duplicated and diversified

among the four amphioxus and *LOX_Bf_9* was an outlier gene (Fig. 2B). All the LOXs

contain a C2-like domain (Fig. 2B). Except for the five LOXs of *B. floridae* and *B.*

lanceolatum, the LOXs possessed one to five domains of the PYR (pyrabactin

resistance)/PYL (PYR1-like)/RCAR (regulatory components of ABA receptor) family in the

SRPBCC superfamily (SRPBCC domains, Fig. 2B) and these SRPBCC domains could be

further divided into three groups (Fig. S7). In Fig. 2C, the gene synteny alignment revealed

that many LOXs (12 of 27 genes) were proximally located (distance <20 Kb) and possibly

generated by tandem gene duplications. *LOX_Bl_5* had an apparently wide expression in the

gills, gut, and hepatic diverticula (Fig. S5).

These unusual LOX genes containing a C2-like domain were reported in amphioxus (Yuan

and Xu, 2016; Yuan et al., 2014) and first in jawless vertebrates, although the SRPBCC

domains were not identified. Since SRPBCC superfamily proteins contain a deep

hydrophobic ligand-binding pocket and can bind diverse ligands (Chen et al., 2015), we

consider that these SRPBCC domains may also function as receptor domains in these LOX

genes in response to various signals. A multifunctional LOX containing a SRPBCC domain

was reported in the red alga *Pyropia haitanensis* and presented substrate flexibility,

oxygenation position flexibility, and functional versatility (Chen et al., 2015). These LOX

genes may enrich the inflammatory lipid signaling pathway of amphioxus.

DEAD-box helicases (DBHs) are known as sensors for nucleic acids and mediators of

antiviral innate immunity (Taschuk and Cherry, 2020). A DBH gene was identified as a

single-copy gene that resembles those from bacteria, and it was conserved among four

amphioxus (Table 1). In the phylogenetic tree (Fig. 3A), this DBH gene of amphioxus is close

to those of the charophyte alga *Klebsormidium* (*K.*) *nitens*, soft coral *Dendronephthya*

gigantea, hemichordate *Saccoglossus kowalevskii*, and a group of bacteria. Intriguingly, the

DBH genes of amphioxus shared three conserved splice sites with those of charophyte alga, soft coral, and hemichordate (Fig. 3B and S7), which suggested that the DBH genes of four amphioxus, soft coral and hemichordate shared a common source with that of the charophyte alga. A homologous DBH gene was also identified in the genome of the other hemichordate, *Ptychodera flava*, in the absence of annotation. This DBH gene was widely expressed at low levels in the tissues of *B. lanceolatum* (Fig. S5). The single-copy DBH gene may serve as a novel sensor in the antiviral innate immunity of amphioxus. Another DBH gene was reported as a horizontally transferred gene in coral species (Bhattacharya et al., 2016).

In addition, another nine conserved splice sites in orange lines (Fig. 3B and S7) indicated close phylogenetic relationships among the DBHs of the four amphioxus, soft coral and hemichordate; another four conserved splice sites in blue lines (Fig. 3B and S7) suggested that the DBHs of the soft coral and hemichordate originated from the same ancestral gene, while two unusually conserved splice sites marked with black lines (S1 and S2, Fig. 3B and S7) could be explained by intron loss during gene evolution, which indicated S1 loss in ancestral amphioxus but S2 loss in both the ancestral amphioxus and hemichordate. Additional gene novelties related to innate immunity, such as Ricin B-like lectins, can be found in Supplementary Note 3.3.

Glycolysis

Two novel genes, Alcohol Dehydrogenases (ADHs) and Ferredoxin (FDX), have been identified as being involved in the glycolysis pathway of amphioxus (Table 1).

ADHs facilitate the oxidation of alcohols into aldehydes and ketones during the glycolysis

process (Di et al., 2021). In the phylogenetic tree (Fig. 3C), an ADH gene containing zinc-

binding sites and belonging to the conserved medium-chain dehydrogenase/reductase (MDR)

family was much closer to those of the sponge *Amphimedon queenslandica*, the copepod

Eurytemora affinis, and a bacterial group than to those of other metazoa. The splice sites of

the ADH gene of amphioxus are highly conserved with those of the sponge but not the

copepod (Fig. 3D and S10), which suggested that the ADH genes of amphioxus and

Amphimedon queenslandica have the same eukaryotic origin. Among the 100 best hits of this

ADH gene in the NCBI NR database, we noticed over ten hits from anaerobic bacteria and

the ADH gene of *B. belcheri* beihai shared 64% identity with that of *Anaerolineae bacterium*

(GenBank accession: MCD6286762.1).

FDX, which contains an iron-sulfur cluster, is recognized as a mediator of electron transfer in

numerous metabolic pathways, including glycolysis (Meyer, 1988). The FDX gene of

amphioxus was identified to be highly conserved with the genes in certain plants, bacteria, or

other eukaryotes (identities over 68%) but not with that of other metazoa (excluding

Amphioxiformes, Table 1). This FDX gene was tandemly duplicated into two copies on

opposite strands in *B. floridae*. All these FDX genes contain a Ubl_FUBI (Ubiquitin-like

protein FUBI) domain, which may protect the FDX gene from ubiquitination and subsequent

degradation (Rossman et al., 2003).

Calcium balance

Calcium balance and osmoregulation of fishes are important for their physiological functions (Guerreiro, 2007). A voltage-dependent calcium channel (VDCC), EF-hand calcium-binding protein (EF-CaBP) group and calcium-binding protein (CML26) were identified as gene novelties in amphioxus (Table 1).

VDCC is a complex of several different subunits (Dolphin, 2006); however, whether this VDCC gene functions independently as a calcium channel deserves further investigation. The VDCC genes of amphioxus were more similar to those of other eukaryotes than to those of other metazoans (Table 1). Additionally, the VDCC genes of amphioxus have homologs in the hemichordate *Saccoglossus kowalevskii* and three starfish species, all of which share three conserved splice sites (Fig. S12). This VDCC gene was proposed to be related to electrophysiological function and was widely expressed among tissues, but no apparent tissue-specific expression (Fig. S5).

EF-CaBPs may function in maintaining a high calcium concentration in the endoplasmic reticulum (Camacho et al., 2003) and could be involved in bone formation (Allen and Hall, 1978). Of the 73 newly identified EF-CaBP genes in four amphioxus, the bit scores of the best hits to bacteria were significantly higher than those to other metazoa (Fig. 4A and S12).

This group of EF-CaBP genes was proximally arrayed in four amphioxus and presented high copy number variation. *B. belcheri* xiamen has 25 EF-CaBP genes, *B. belcheri* beihai has 19

genes, *B. floridae* has 17 genes, and *B. lanceolatum* has only 12 genes. Marine salinity is

linearly correlated with the calcium ion concentration (He et al., 2020). Therefore, we further

explored the relationship of the gene number variation of these proximally arrayed EF-CaBP

genes and the sea surface salinity of the collection locations (Fig. 4B). The Mediterranean

Sea has a much higher than average seawater salinity (approximately 35 parts per thousand)

(Millero et al., 2008; Millot and Taupier-Letage, 2005), and the Atlantic Ocean is globally

saltier than the Pacific Ocean (Jones and Cessi, 2017). Although high-quality salinity data for

the nearshore regions where the four amphioxus were collected is lacking, a negative

correlation between seawater salinity and the number of EF-CaBP genes can be observed. *B.*

lanceolatum, collected from the Mediterranean Sea, has the fewest genes, while the two *B.*

belcheri from the Pacific Ocean have a higher number of genes (Fig. 4B).

In the phylogenetic tree (Fig. 4C), except for the outlier gene *Bf_10*, the EF-CaBP genes were

classified into two large groups, Clade A and Clade B. Clade A was further classified into

three clusters, Cluster 1, Cluster 2, and Cluster 3. In Cluster 1, no genes of *B. floridae* were

identified while eight genes of two *B. belcheri* were assigned to four pairs, and they included

four highly conserved genes, *Bbb_2*, *Bbb_3*, *Bbx_21* and *Bbx_22*. However, three and two

tandemly arrayed genes in Cluster 2 were identified in *B. floridae* and *B. lanceolatum*,

respectively, and variable duplications were observed in Cluster 3. The phylogenetic analysis

indicated that at least three genes in Clade A appeared in the ancestral amphioxus species. For

Clade B, five relatively unique genes of *B. floridae*, *Bf_1-5* in Cluster 4, were proximally

arrayed, and many paired genes were identified in two *B. belcheri*, such as three pairs in

Cluster 5. All 73 EF-CaBPs were proximally arrayed in four amphioxus (Fig. 4D) and all

spaces between genes were below 20 Kb. Notably, four longer genes, *Bbb_6*, *Bbx_18*, *Bf_11*

and *Bl_6*, contained more EF-hand motifs (Cluster 3, Fig. 4C).

Except for the large group of proximally arrayed EF-CaBPs above, the CML26 gene was a

calcium-binding gene with a shorter length (Table 1). This CML26 gene was tandemly

duplicated in *B. floridae* and *B. lanceolatum* (Table 1). Initially, we considered the CML26

gene (Table 1) to belong to another event, but the conserved splice sites (Fig. S14) suggested

that the CML26 gene and the large group of EF-CaBPs possibly originated from a common

ancestor.

Along with the two CML26 genes, these EF-CaBP genes were apparently highly expressed in

the gut and hepatic diverticulum, which are both considered digestive organs for calcium

absorption (Fig. S5). Thus, the expansion of these EF-CaBP genes was proposed to occur in

response to marine salinity and calcium concentration changes. The *EF-CaBP_Bl_11* and the

two CML26 genes had the highest expression levels (Fig. S5).

Conclusions

As an important group of invertebrate chordates, amphioxus play a pivotal role in the

evolutionary study of cephalochordates, chordates, and deuterostomes. In this study, the

genome of the Beihai amphioxus, *B. belcheri* beihai, was *de novo* assembled and annotated

(Table S1). Combined with the three amphioxus genomes (Table S1), we performed a

phylogenetic analysis of the deuterostome genomes (Fig. 1), which supported the

monophylies of deuterostomes and chordates.

We identified a number of gene novelties conserved among the four amphioxus genomes

(Table 1, Supplementary Note 3.1). Among these gene novelties, except for the TIR domains

of amphioxus, which was mentioned in a previous report (Huang et al., 2008), all other genes

were newly identified and reported. Certain involved genes had homologs from other

deuterostome groups except for the jawed vertebrate group (Table 1 and S3). That is, the

LOX gene had homologs from two jawless vertebrates; 8 genes had homologs from either

tunicates or echinoderms or hemichordates; and only the VDCC gene had homologs from

both echinoderms and hemichordates. The conservation of these genes can be well explained

by frequent gene gain and loss in the early and rapid evolution of deuterostomes (Table S3)

(Kapli et al., 2021). For example, the LOX gene was suggested to be a novel gene that was

introduced at the deuterostome stem but decayed in tunicates and jawed vertebrates. Those

gene novelties could be well dated according to their conservation among different

deuterostome groups, such as the three genes conserved with urochordates (SCP24, MCO and

UF-TM3) were proposed as results of events occurred at the chordate stem, while their

absence in vertebrates is due to gene decay. While it is true that certain genes, such as DBH

and ADH, have homologs in ancient metazoa like coral or sponge, we cannot definitively

conclude that these genes originated in the common ancestor of all metazoa. This is because

the homology was observed only in specific coral or sponge species, suggesting independent

acquisition events rather than inheritance from a common ancestor. For example, while the

alignment of splice sites within the DBH gene (Fig. 3B) may suggest a close relationship

among these genes, it cannot support that this gene was introduced in a common ancestor

before the emergence of the ancient soft coral. Although these genes with restricted

distribution are not exclusive to the amphioxus lineage, their surprisingly high similarity to

non-metazoan genes is interesting and deserves further investigation. Additionally, the

absence of high-quality genomes from the other amphioxus genera, including *Epigonichthys*

and *Asymmetron*, hinders our comprehensive investigation of more cephalochordate species.

The gene novelties without any homologs in deuterostomes were suggested to be acquired by

amphioxus lineage-specific horizontal gene transfer (HGT) from non-metazoan species. A

number of adaptation innovations were enabled by these genes, such as innate immunity-

related events that contributed more complexity to the immune system in amphioxus,

especially for amphioxus has no adaptative immune system. Although the function of these

novel genes has not been validated in any experimental model, either the gene duplications

(Table 1) or the negative selection indicated by their low Ka/Ks values (Fig. S25) suggest that

they may indeed be functional essential genes.

Proposed mechanisms

Although it is difficult to rule out gene loss in other groups like protostomes and/or extreme divergence, those gene novelties, especially the genes with phylogenetic supports, can be explained as results of HGT from cross-kingdom origins such as marine bacteria, considering many genes shared high similarity with those from unusual taxonomic groups. Notably, reports have documented similar HGT events of the RAG transposon and the F-type lectins in amphioxus (Bishnoi et al., 2015; Carmona and Schatz, 2017).

The HGT of amphioxus could be explained by the 'you are what you eat hypothesis' (Doolittle, 1998). This hypothesis suggests that the HGT genes in amphioxus originate from its food sources, which is particularly plausible considering that amphioxus can perform phagocytic intercellular digestion in the diverticulum lumen (He et al., 2018). All modern deuterostomes were considered to have evolved from a filter-feeding ancestor (Simakov et al., 2015). Along with other deuterostomes, such as tunicates, echinoderms, and hemichordates, amphioxus is known as a filter feeder (Peterson and Eernisse, 2016; Riisgård and Svane, 1999) and may obtain genes from marine microbes, including bacteria and algae, which are potential foods. For example, the charophyte alga *K. nitens* frequently appeared in the analysis of the HGT genes, including the DBH, SCP24 and UF-TM1 genes (Table 1). The DBH gene was unusually identified in the charophyte alga *K. nitens* (Fig. 3A). The highly conserved splice sites among the DBH genes of the charophyte algae, soft coral, hemichordate, and amphioxus (Fig. 3B and S7) suggested that their DBH genes originated from a common ancient eukaryote, such as some marine algae. Although the charophyte alga

K. nitens is considered a freshwater/terrestrial and nonmarine species (Hori et al., 2014), we

proposed that the ancestor of *K. nitens* was possibly distributed in marine environments and

served as food for these marine animals, especially because of the sister marine species *K.*

marinum (no available genome sequences). Similarly, the SCP24 gene (Fig. S15) and UF-

TM1 gene (Table 1) possibly originated from the marine ancestor of *K. nitens*.

New gene generation

The origin of new genes is a fundamental process of genome evolution (Long et al., 2003). In

this study, the comparative analysis of the gene novelties revealed insights into the new gene

generation. Regardless of whether common gene duplication and diversification occurred,

many genes of amphioxus underwent possible gene fusion or fission to generate new genes.

Three novel models of new gene generation were summarized, including gene fusion (Fig.

S26A), gene fission (Fig. S26B) and partial duplication or deletion models (Fig. S26C).

In the gene fusion model (Fig. S26A), we revealed that two distinct TIR domains fused with a

number of endogenous genes (or domains) with immune functions (Fig. S10) and generated a

series of new immune genes. Both gene fission and partial duplication or deletion were

observed in the LOX family of amphioxus (Fig. 2B). Since we found that the LOX genes of

amphioxus were mostly similar to bacterial LOX genes containing up to three SRPBCC

domains and three groups of SRPBCC domains could be clustered with three SRPBCC

domains of the cyanobacterium *Coleofasciculus sp. FACHB-125* (GenBank accession:

WP_190414990.1) (Fig. S7), we considered that the original LOX gene of amphioxus

contains three SRPBCC domains. Then, they were split into LOX genes with only C2

domains and genes with only SRPBCC domain via gene fission (Fig. S26B) and generated a

series of LOX genes with variable numbers of SRPBCC domains (Fig. 2B) via partial gene

duplication or deletion (Fig. S26C). The investigation of the origins of the functional domains

of these novel genes through phylogenetic analysis would enhance the credibility of the

findings. However, the substantial challenges posed by the relatively low sequence

similarities resulting from extensive evolutionary divergence, coupled with the lack of

comparable outgroup genes from other taxonomic groups, complicate this analysis

significantly.

Limitations of this study

During the identification of gene novelties, we ignored those candidates with homologous

genes in jawed vertebrates. Although this avoided repeating efforts on the gene novelties of

chordates (Holland et al., 2008) and deuterostomes (Simakov et al., 2015), we may miss

some important genes. Meanwhile, the filter standard of gene novelties may be too high to

achieve highly comprehensive results, and many genes may be missed during filtration.

The lack of animal model experiments impeded our validation of the metabolic functions of

some interesting genes. The exploration of genomic evolution following these gene novelties

presents an intriguing direction, except the gene duplications, such as LOXs and EF-CaBPs.

To fully explore this direction, delineating the functional pathways and gene networks in

amphioxus would be necessary, but represents a substantial challenge that falls beyond the

scope of this study. In the case of the ADH gene, three intrinsic ADH like genes in *B. lanceolatum*, which share homology with those in other metazoan species, were identified with high similarity to the novel ADH gene, exhibiting low E-values below 1E-30 in BLASTP. In muscle tissue, the novel ADH gene could be expressed at levels over the combined expression of the three intrinsic genes, indicating its functional importance and warranting further investigation into its role within the related gene network.

Despite the methodological challenges and difficulties in pinpointing the precise origins of gene novelties, the phylogenetic trees support our conclusion that these notable genes are more closely related to those of non-metazoan species and represent gene novelties in amphioxus. However, the analysis included only a limited number of genes, as many were excluded due to relatively low similarities resulting from long-term evolution or the absence of comparable outgroup genes from other taxonomic groups.

Table 1. Twenty-eight groups of gene novelties identified in amphioxus

Functional category	Abbrev.	Representative gene ID[①]	Amino acid length	Best hit, metazoan[②]	Best hit[③]			Copy number[④]			
					E-value	Identity	Taxonomy	*Bbb*	*Bbx*	*Bf*	*Bl*
Innate immunity	LOX*	Bbb_152860.1[⑤]	696	3E-62	7E-168	39.6%	Bacteria, cyanobacteria/proteobacteria	6	5	9	7
	DBH*	Bbb_034850.1	1253	2E-141	0.0	39.1%	Bacteria, cyanobacteria/proteobacteria	1	1	1	1
	RBL	Bbb_161350.1	282	8E-17	4E-23	29.7%	Bacteria, Terrabacteria group	2	1	3	1
	TIR-a	Bbb_152643.1[⑥]	126	1E-4	4E-21	42.5%	Bacteria, *Solibacter usitatus*	4	5	2	2
	TIR-b	Bbb_258280.1[⑥]	95	N. A.	1E-10	43.3%	Bacteria, cyanobacteria	11	14	18	9
Glycolysis	ADH*	Bbb_340330.1	355	3E-78	1E-129	51.3%	Bacteria, Gammaproteobacteria	1	1	1	1
	FDX	Bbb_233460.1	171	1E-12	6E-46	74.5%	Plants, *Micractinium conductrix*	1	1	2	1
Calcium balance	VDCC*	Bbb_084320.1	524	3E-36	9E-83	37.2%	Other eukaryotes, *Salpingoeca rosetta*	1	1	1	1
	EF-CaBP	Bbb_077420.1	248	2E-6	5E-11	27.4%	Bacteria, Gammaproteobacteria	19	25	17	12
	CML26	Bbb_165620.1	153	4E-6	3E-11	36.2%	Bacteria, Gammaproteobacteria	1	1	2	2
Amino acid metabolism	SCP24*	Bbb_236620.1	474	1E-83	8E-136	46.6%	Plants, *Pycnococcus provasolii*	1	1	1	1
	GSA*	Bbb_006430.1	438	2E-32	2E-113	53.6%	Bacteria, Deltaproteobacteria	2	1	1	1
	GNAT	Bbb_266880.1	325	3E-26	7E-60	39.3%	Bacteria, Alphaproteobacteria	3	5	5	5
	DDC*	Bbb_001110.1	586	4E-49	2E-141	43.4%	Bacteria, *Bacteroidetes bacterium*	9	8	16	7
	DOD	Bbb_303310.1	173	3E-8	3E-36	42.5%	Other eukaryotes, *Tetrahymena*	2	1	1	1
	FR	Bbb_207740.1	241	9E-24	3E-72	54.0%	Other eukaryotes, Chlorarachniophyte	2	3	5	4
	DHFR	Bbb_115460.1	186	2E-29	1E-43	44.0%	Bacteria, *Ignavibacteriae bacterium*	1	1	1	1
Other metabolic genes	GH27[⑦]	Bbb_359430.1	419	8E-32	6e-82	40.1%	Bacteria, Terrabacteria group	1	1	1	1
	MCO*	Bbb_246040.1	640	4E-10	7E-67	30.7%	Plants, *Tetraselmis astigmatica*	5	3	8	4
	anmK	Bbb_029670.1	292	7E-29	2E-51	52.2%	Bacteria, cyanobacteria	1	1	1	1
Others	TPR	Bbb_379800.1	884	1E-86	0.0	37.6%	Other eukaryotes, Amoeba	1	1	1	1
	ApoLp	Bbb_173200.1	1205	N. A.	1E-158	36.8%	Bacteria, Gamma/Deltaproteobacteria	3	2	3	3
	HintN	Bbb_173190.1	789	2E-5	1E-118	36.3%	Bacteria, Gamma/Deltaproteobacteria	3	1	2	3
	SDR_a4	Bbb_177610.1	263	N. A.	5E-35	34.6%	Bacteria, *Methylomarinum vadi*	3	3	3	2
	CytP460	Bbb_256630.1	435	N. A.	2E-94	41.3%	Fungi, *Punctularia strigosozonata*	5	3	8	2

UF-TM1	Bbb_158830.1	341	N. A.	6E-39	35.0%	Plants, *Klebsormidium nitens*	6	6	7	12
UF-TM2*	Bbb_202120.1	1108	8E-7	4E-34	23.9%	Fungi, *Aspergillus thermomutatus*	1	1	1	1
UF-TM3*	Bbb_320160.1	268	N. A.	1E-8	29.8%	Bacteria, *Streptomyces*	12	10	6	7

① To describe the gene features, a gene of *B. belcheri* beihai was selected as the representative gene. The protein sequences are listed in Table S4.

② The best hits of metazoan were of matches to genes of metazoa (excluding Amphioxiformes) in UniRef50 database. N.A., Not applicable, because no best hit has E-value smaller than the cutoff 0.001. In some cases, the gene has homologs in other metazoan species and marked in star (*); then these metazoan species were excluded.

③ The best hits in UniRef50 database were listed and displayed in detail. In many cases, the best matches were from a group of microbes, but not exactly a species, so the group name like cyanobacteria was listed.

④ Since many genes underwent gene duplications, copy number of four amphioxus were compared. Bbb, *B. belcheri* beihai; Bbx, *B. belcheri* xiamen; Bf, *B. floridae*; Bl, *B. lanceolatum*.

⑤ For the LOX gene Bbb_152860.1 contains many domains, only the partial 696-aa sequence containing its LOX domain was analyzed.

⑥ For the TIR-a and TIR-b, only their TIR domains were analyzed, which covered 35-160 of Bbb_152640.3 and 107-201 of Bbb_258280.1, respectively.

⑦ For the GH27 gene Bbb_359430.1, the best hit in NR database (6E-82 from Terrabacteria) has a much lower E-value than the best hit in UniRef50 database (6E-74 from plants), therefore we chose to present the best hit in NR database.

Abbreviation:

Innate immunity

LOX *   lipoxygenase; homologs in the sea lamprey *Petromyzon marinus* and the hagfish *Eptatretus burger*;

DBH *   DEAD/DEAH-box helicase; homologs in the soft coral *Dendronephthya gigantea* and hemichordates;

RBL   ricin B-like lectin;

TIR-a   Toll/interleukin-1 receptor domain a;

TIR-b   Toll/interleukin-1 receptor domain b;

Glycolysis

ADH *   alcohol dehydrogenase; homolog in the sponge *Amphimedon queenslandica*;

FDX   ferredoxin;

Calcium balance
VDCC *   voltage-dependent calcium channel (subunit alpha-2/delta-4-like); homologs in hemichordates and echinoderms;
EF-CaBP   EF-hand calcium-binding protein;
CML26   calmodulin-like protein 26;

Amino acid metabolism
SCP24 *   serine carboxypeptidase 24; homologs in tunicates;
GSA *   glutamate-1-semialdehyde aminotransferase; homologs in echinoderms;
GNAT   GNAT family N-acetyltransferase;
DDC *   DOPA decarboxylase; homologs in hemichordates;
DOD   DOPA 4,5-dioxygenase;
FR   folate receptor;
DHFR   dihydrofolate reductase;

Other metabolic genes
GH27   glycoside hydrolase family 27;
MCO *   multicopper oxidase; homologs in tunicates;
anmK   anhydro-N-acetylmuramic acid kinase;

Others
TPR   tetratricopeptide repeat protein;
ApoLp   apolipophorin III-like domain-containing protein;
HintN   Hedgehog/Intein domain-containing protein;
SDR_a4   atypical SDR subgroup 4, containing a glycine-rich NAD(P)-binding motif;
CytP460   cytochrome P460 family like protein;
UF-TM1   transmembrane protein 1 of unknown function;
UF-TM2 *   transmembrane protein 2 of unknown function; homologs in hemichordates;
UF-TM3 *   transmembrane protein 3 of unknown function; homologs in tunicates.

Appendix A. Supplementary Materials

Supplementary Notes 1–3

Tables S1–4

Figures S1–26

Acknowledgements

This work was supported by the General Research Fund from Research Grants Council (Reference numbers: 464710, 475113, 14119219, 14119420, 14175617) and Health and Medical Research Fund from Food and Health Bureau (Reference numbers: 06171016, 07181266) of Hong Kong. We would like to thank Prof. Jun-Yuan Chen from the State Key Laboratory of Palaeobiology and Stratigraphy, China, for providing the Beihai amphioxus samples and for reviewing our manuscript. Additionally, we extend our gratitude to Dr. Laura Jane Falkenberg from the Simon FS Li Marine Science Laboratory, The Chinese University of Hong Kong, Hong Kong, for her valuable comments on the adaptation of amphioxus to ocean environments.

Competing interests

The authors declare no competing interests.

References

Acemel, R.D., Tena, J.J., Irastorza-Azcarate, I., Marlétaz, F., Gómez-Marín, C., de la Calle-Mustienes, E., Bertrand, S., Diaz, S.G., Aldea, D., Aury, J.M., Mangenot, S., Holland, P.W., Devos, D.P., Maeso, I., Escrivá, H., Gómez-Skarmeta, J.L., 2016. A single three-dimensional chromatin compartment in amphioxus indicates a stepwise evolution of vertebrate Hox bimodal regulation. Nat Genet 48, 336-341.

Allen, L.H., Hall, T.E., 1978. Calcium Metabolism, Intestinal Calcium-binding Protein, and Bone Growth of Rats Fed High Protein Diets. The Journal of Nutrition 108, 967-972.

Bao, Z., Eddy, S.R., 2002. Automated de novo identification of repeat sequence families in sequenced genomes. Genome Res 12, 1269-1276.

Bartl, S., Baish, M., Weissman, I.L., Diaz, M., 2003. Did the Molecules of Adaptive Immunity Evolve from the Innate Immune System? Integrative and Comparative Biology 43, 338-346.

Benito-Gutiérrez, È., Gattoni, G., Stemmer, M., Rohr, S.D., Schuhmacher, L.N., Tang, J., Marconi, A., Jékely, G., Arendt, D., 2021. The dorsoanterior brain of adult amphioxus shares similarities in expression profile and neuronal composition with the vertebrate telencephalon. BMC Biol 19, 110.

Bentley, D.R., Balasubramanian, S., Swerdlow, H.P., Smith, G.P., Milton, J., Brown, C.G., Hall, K.P., Evers, D.J., Barnes, C.L., Bignell, H.R., Boutell, J.M., Bryant, J., Carter, R.J., Keira Cheetham, R., Cox, A.J., Ellis, D.J., Flatbush, M.R., Gormley, N.A., Humphray, S.J., Irving, L.J., Karbelashvili, M.S., Kirk, S.M., Li, H., Liu, X., Maisinger, K.S., Murray, L.J., Obradovic, B., Ost, T., Parkinson, M.L., Pratt, M.R., Rasolonjatovo, I.M., Reed, M.T., Rigatti, R., Rodighiero, C., Ross, M.T., Sabot, A., Sankar, S.V., Scally, A., Schroth, G.P., Smith, M.E., Smith, V.P., Spiridou, A., Torrance, P.E., Tzonev, S.S., Vermaas, E.H., Walter, K., Wu, X., Zhang, L., Alam, M.D., Anastasi, C., Aniebo, I.C., Bailey, D.M., Bancarz, I.R., Banerjee, S., Barbour, S.G., Baybayan, P.A., Benoit, V.A., Benson, K.F., Bevis, C., Black, P.J., Boodhun, A., Brennan, J.S., Bridgham, J.A., Brown, R.C., Brown, A.A., Buermann, D.H., Bundu, A.A., Burrows, J.C., Carter, N.P., Castillo, N., Chiara, E.C.M., Chang, S., Neil Cooley, R., Crake, N.R., Dada, O.O., Diakoumakos, K.D., Dominguez-Fernandez, B., Earnshaw, D.J., Egbujor, U.C., Elmore, D.W., Etchin, S.S., Ewan, M.R., Fedurco, M., Fraser, L.J., Fuentes Fajardo, K.V., Scott Furey, W., George, D., Gietzen, K.J., Goddard, C.P., Golda, G.S., Granieri, P.A., Green, D.E., Gustafson, D.L., Hansen, N.F., Harnish, K., Haudenschild, C.D., Heyer, N.I., Hims, M.M., Ho, J.T., Horgan, A.M., Hoschler, K., Hurwitz, S., Ivanov, D.V., Johnson, M.Q., James, T., Huw Jones, T.A., Kang, G.D., Kerelska, T.H., Kersey, A.D., Khrebtukova, I., Kindwall, A.P., Kingsbury, Z., Kokko-Gonzales, P.I., Kumar, A., Laurent, M.A., Lawley, C.T., Lee, S.E., Lee, X., Liao, A.K., Loch, J.A., Lok, M., Luo, S., Mammen, R.M., Martin, J.W., McCauley, P.G., McNitt, P., Mehta, P., Moon, K.W., Mullens, J.W., Newington, T., Ning, Z., Ling Ng, B., Novo, S.M., O'Neill, M.J., Osborne, M.A., Osnowski, A., Ostadan, O., Paraschos, L.L., Pickering, L., Pike, A.C., Pike, A.C., Chris Pinkard, D., Pliskin, D.P., Podhasky, J., Quijano, V.J., Raczy, C., Rae, V.H., Rawlings, S.R., Chiva Rodriguez, A., Roe, P.M., Rogers, J., Rogert Bacigalupo, M.C., Romanov, N., Romieu, A., Roth, R.K., Rourke, N.J., Ruediger, S.T., Rusman, E., Sanches-Kuiper, R.M., Schenker, M.R., Seoane, J.M., Shaw, R.J., Shiver, M.K., Short, S.W., Sizto, N.L., Sluis, J.P., Smith, M.A., Ernest Sohna Sohna, J., Spence, E.J., Stevens, K., Sutton, N., Szajkowski, L., Tregidgo, C.L., Turcatti, G., Vandevondele, S., Verhovsky, Y., Virk, S.M., Wakelin, S., Walcott, G.C., Wang, J., Worsley, G.J., Yan, J., Yau, L., Zuerlein, M., Rogers, J., Mullikin, J.C., Hurles, M.E., McCooke, N.J., West, J.S., Oaks, F.L., Lundberg, P.L., Klenerman, D., Durbin, R., Smith, A.J., 2008. Accurate whole human genome sequencing using reversible terminator chemistry. Nature 456, 53-59.

Bertrand, S., Escriva, H., 2011. Evolutionary crossroads in developmental biology: amphioxus. Development 138, 4819-4830.

Bhattacharya, D., Agrawal, S., Aranda, M., Baumgarten, S., Belcaid, M., Drake, J.L., Erwin, D., Foret, S., Gates, R.D., Gruber, D.F., Kamel, B., Lesser, M.P., Levy, O., Liew, Y.J., MacManes, M., Mass, T., Medina, M., Mehr, S., Meyer, E., Price, D.C., Putnam, H.M., Qiu, H., Shinzato, C., Shoguchi, E., Stokes, A.J., Tambutté, S., Tchernov, D., Voolstra, C.R.,

Wagner, N., Walker, C.W., Weber, A.P.M., Weis, V., Zelzion, E., Zoccola, D., Falkowski, P.G., 2016. Comparative

genomics explains the evolutionary success of reef-forming corals. eLife 5, e13288.

Bishnoi, R., Khatri, I., Subramanian, S., Ramya, T.N.C., 2015. Prevalence of the F-type lectin domain. Glycobiology 25,

888-901.

Boetzer, M., Henkel, C.V., Jansen, H.J., Butler, D., Pirovano, W., 2011. Scaffolding pre-assembled contigs using

SSPACE. Bioinformatics 27, 578-579.

Boetzer, M., Pirovano, W., 2014. SSPACE-LongRead: scaffolding bacterial draft genomes using long read sequence

information. BMC Bioinformatics 15, 211.

Borodovsky, M., Lomsadze, A., 2011. Eukaryotic gene prediction using GeneMark.hmm-E and GeneMark-ES. Current

protocols in bioinformatics Chapter 4, Unit 4.6.1-10.

Bourlat, S.J., Juliusdottir, T., Lowe, C.J., Freeman, R., Aronowicz, J., Kirschner, M., Lander, E.S., Thorndyke, M.,

Nakano, H., Kohn, A.B., Heyland, A., Moroz, L.L., Copley, R.R., Telford, M.J., 2006. Deuterostome phylogeny reveals

monophyletic chordates and the new phylum Xenoturbellida. Nature 444, 85-88.

Brasó-Vives, M., Marlétaz, F., Echchiki, A., Mantica, F., Acemel, R.D., Gómez-Skarmeta, J.L., Hartasánchez, D.A., Le

Targa, L., Pontarotti, P., Tena, J.J., Maeso, I., Escriva, H., Irimia, M., Robinson-Rechavi, M., 2022a. Parallel evolution of

amphioxus and vertebrate small-scale gene duplications. Genome Biology 23, 243.

Brasó-Vives, M., Marlétaz, F., Echchiki, A., Mantica, F., Acemel, R.D., Gómez-Skarmeta, J.L., Hartasánchez, D.A., Le

Targa, L., Pontarotti, P., Tena, J.J., Maeso, I., Escriva, H., Irimia, M., Robinson-Rechavi, M., 2022b. Parallel evolution of

amphioxus and vertebrate small-scale gene duplications. Genome Biol 23, 243.

Camacho, P., John, L., Li, Y., Paredes, R.M., Roderick, H.L., 2003. Calnexin and calreticulin, ER associated modulators

of calcium transport in the ER. Calreticulin. Springer, pp. 126-132.

Cantalapiedra, C.P., Hernández-Plaza, A., Letunic, I., Bork, P., Huerta-Cepas, J., 2021. eggNOG-mapper v2: Functional

Annotation, Orthology Assignments, and Domain Prediction at the Metagenomic Scale. Molecular Biology and

Evolution 38, 5825-5829.

Cantarel, B.L., Korf, I., Robb, S.M., Parra, G., Ross, E., Moore, B., Holt, C., Sánchez Alvarado, A., Yandell, M., 2008.

MAKER: an easy-to-use annotation pipeline designed for emerging model organism genomes. Genome Res 18, 188-196.

Carmona, L.M., Schatz, D.G., 2017. New insights into the evolutionary origins of the recombination-activating gene

proteins and V(D)J recombination. The FEBS Journal 284, 1590-1605.

Castresana, J., 2000. Selection of Conserved Blocks from Multiple Alignments for Their Use in Phylogenetic Analysis.

Molecular Biology and Evolution 17, 540-552.

Chen, H.-m., Zhu, Z.-j., Chen, J.-J., Yang, R., Luo, Q.-j., Xu, J.-l., Shan, H., Yan, X.-J., 2015. A multifunctional

lipoxygenase from Pyropia haitanensis— The cloned and functioned complex eukaryotic algae oxylipin pathway

enzyme. Algal Research 12, 316-327.

Chen, X., Li, Q., Wang, J., Guo, X., Jiang, X., Ren, Z., Weng, C., Sun, G., Wang, X., Liu, Y., Ma, L., Chen, J.-Y., Wang,

J., Zen, K., Zhang, J., Zhang, C.-Y., 2009. Identification and characterization of novel amphioxus microRNAs by Solexa

sequencing. Genome Biology 10, R78.

De Bie, T., Cristianini, N., Demuth, J.P., Hahn, M.W., 2006. CAFE: a computational tool for the study of gene family

evolution. Bioinformatics 22, 1269-1271.

Delsuc, F., Brinkmann, H., Chourrout, D., Philippe, H., 2006. Tunicates and not cephalochordates are the closest living

relatives of vertebrates. Nature 439, 965-968.

Di, L., Balesano, A., Jordan, S., Shi, S.M., 2021. The role of alcohol dehydrogenase in drug metabolism: beyond ethanol

oxidation. The AAPS Journal 23, 1-21.

Dolphin, A.C., 2006. A short history of voltage-gated calcium channels. Br J Pharmacol 147 Suppl 1, S56-S62.

Doolittle, W.F., 1998. You are what you eat: a gene transfer ratchet could account for bacterial genes in eukaryotic

nuclear genomes. Trends in Genetics 14, 307-311.

Du, Q., Peng, F., Xiong, Q., Xu, K., Yang, K.Y., Wang, M., Wu, Z., Li, S., Cheng, X., Rao, X., Wang, Y., Tsui, S.K.-W., Zeng, X., 2023. Genomic Analysis of Amphioxus Reveals a Wide Range of Fragments Homologous to Viral Sequences. Viruses 15, 909.

Edgar, R.C., 2004. MUSCLE: a multiple sequence alignment method with reduced time and space complexity. BMC Bioinformatics 5, 113.

Eid, J., Fehr, A., Gray, J., Luong, K., Lyle, J., Otto, G., Peluso, P., Rank, D., Baybayan, P., Bettman, B., Bibillo, A., Bjornson, K., Chaudhuri, B., Christians, F., Cicero, R., Clark, S., Dalal, R., Dewinter, A., Dixon, J., Foquet, M., Gaertner, A., Hardenbol, P., Heiner, C., Hester, K., Holden, D., Kearns, G., Kong, X., Kuse, R., Lacroix, Y., Lin, S., Lundquist, P., Ma, C., Marks, P., Maxham, M., Murphy, D., Park, I., Pham, T., Phillips, M., Roy, J., Sebra, R., Shen, G., Sorenson, J., Tomaney, A., Travers, K., Trulson, M., Vieceli, J., Wegener, J., Wu, D., Yang, A., Zaccarin, D., Zhao, P., Zhong, F., Korlach, J., Turner, S., 2009. Real-time DNA sequencing from single polymerase molecules. Science 323, 133-138.

Emms, D.M., Kelly, S., 2015. OrthoFinder: solving fundamental biases in whole genome comparisons dramatically improves orthogroup inference accuracy. Genome Biol 16, 157.

Emms, D.M., Kelly, S., 2019. OrthoFinder: phylogenetic orthology inference for comparative genomics. Genome Biol 20, 238.

Escriva, H., 2018. My Favorite Animal, Amphioxus: Unparalleled for Studying Early Vertebrate Evolution. Bioessays 40, e1800130.

Flynn, J.M., Hubley, R., Goubert, C., Rosen, J., Clark, A.G., Feschotte, C., Smit, A.F., 2020. RepeatModeler2 for automated genomic discovery of transposable element families. Proceedings of the National Academy of Sciences 117, 9451-9457.

Guerreiro, P.F., Juan, 2007. Control of calcium balance in fish. Fish Osmoregulation. Science Publisher, USA.

Gurevich, A., Saveliev, V., Vyahhi, N., Tesler, G., 2013. QUAST: quality assessment tool for genome assemblies. Bioinformatics 29, 1072-1075.

Hahn, M.W., De Bie, T., Stajich, J.E., Nguyen, C., Cristianini, N., 2005. Estimating the tempo and mode of gene family evolution from comparative genomic data. Genome Res 15, 1153-1160.

He, C., Han, T., Liao, X., Zhou, Y., Wang, X., Guan, R., Tian, T., Li, Y., Bi, C., Lu, N., He, Z., Hu, B., Zhou, Q., Hu, Y., Lu, Z., Chen, J.Y., 2018. Phagocytic intracellular digestion in amphioxus (Branchiostoma). Proc Biol Sci 285.

He, H., Li, Y., Wang, S., Ma, Q., Pan, Y., 2020. A High Precision Method for Calcium Determination in Seawater Using Ion Chromatography. Frontiers in Marine Science 7.

Hoff, K.J., Stanke, M., 2019. Predicting Genes in Single Genomes with AUGUSTUS. Current protocols in bioinformatics 65, e57.

Holland, L.Z., Albalat, R., Azumi, K., Benito-Gutiérrez, E., Blow, M.J., Bronner-Fraser, M., Brunet, F., Butts, T., Candiani, S., Dishaw, L.J., Ferrier, D.E., Garcia-Fernàndez, J., Gibson-Brown, J.J., Gissi, C., Godzik, A., Hallböök, F., Hirose, D., Hosomichi, K., Ikuta, T., Inoko, H., Kasahara, M., Kasamatsu, J., Kawashima, T., Kimura, A., Kobayashi, M., Kozmik, Z., Kubokawa, K., Laudet, V., Litman, G.W., McHardy, A.C., Meulemans, D., Nonaka, M., Olinski, R.P., Pancer, Z., Pennacchio, L.A., Pestarino, M., Rast, J.P., Rigoutsos, I., Robinson-Rechavi, M., Roch, G., Saiga, H., Sasakura, Y., Satake, M., Satou, Y., Schubert, M., Sherwood, N., Shiina, T., Takatori, N., Tello, J., Vopalensky, P., Wada, S., Xu, A., Ye, Y., Yoshida, K., Yoshizaki, F., Yu, J.K., Zhang, Q., Zmasek, C.M., de Jong, P.J., Osoegawa, K., Putnam, N.H., Rokhsar, D.S., Satoh, N., Holland, P.W., 2008. The amphioxus genome illuminates vertebrate origins and cephalochordate biology. Genome Res 18, 1100-1111.

Holland, P., 2017. The dawn of amphioxus molecular biology - a personal perspective. Int J Dev Biol 61, 585-590.

Hori, K., Maruyama, F., Fujisawa, T., Togashi, T., Yamamoto, N., Seo, M., Sato, S., Yamada, T., Mori, H., Tajima, N., Moriyama, T., Ikeuchi, M., Watanabe, M., Wada, H., Kobayashi, K., Saito, M., Masuda, T., Sasaki-Sekimoto, Y.,

Mashiguchi, K., Awai, K., Shimojima, M., Masuda, S., Iwai, M., Nobusawa, T., Narise, T., Kondo, S., Saito, H., Sato, R., Murakawa, M., Ihara, Y., Oshima-Yamada, Y., Ohtaka, K., Satoh, M., Sonobe, K., Ishii, M., Ohtani, R., Kanamori-Sato, M., Honoki, R., Miyazaki, D., Mochizuki, H., Umetsu, J., Higashi, K., Shibata, D., Kamiya, Y., Sato, N., Nakamura, Y., Tabata, S., Ida, S., Kurokawa, K., Ohta, H., 2014. Klebsormidium flaccidum genome reveals primary factors for plant terrestrial adaptation. Nature Communications 5, 3978.

Huang, S., Chen, Z., Yan, X., Yu, T., Huang, G., Yan, Q., Pontarotti, P.A., Zhao, H., Li, J., Yang, P., Wang, R., Li, R., Tao, X., Deng, T., Wang, Y., Li, G., Zhang, Q., Zhou, S., You, L., Yuan, S., Fu, Y., Wu, F., Dong, M., Chen, S., Xu, A., 2014. Decelerated genome evolution in modern vertebrates revealed by analysis of multiple lancelet genomes. Nat Commun 5, 5896.

Huang, S., Yuan, S., Guo, L., Yu, Y., Li, J., Wu, T., Liu, T., Yang, M., Wu, K., Liu, H., Ge, J., Yu, Y., Huang, H., Dong, M., Yu, C., Chen, S., Xu, A., 2008. Genomic analysis of the immune gene repertoire of amphioxus reveals extraordinary innate complexity and diversity. Genome Res 18, 1112-1126.

Huang, Z., Xu, L., Cai, C., Zhou, Y., Liu, J., Xu, Z., Zhu, Z., Kang, W., Cen, W., Pei, S., Chen, D., Shi, C., Wu, X., Huang, Y., Xu, C., Yan, Y., Yang, Y., Xue, T., He, W., Hu, X., Zhang, Y., Chen, Y., Bi, C., He, C., Xue, L., Xiao, S., Yue, Z., Jiang, Y., Yu, J.-K., Jarvis, Erich D., Li, G., Lin, G., Zhang, Q., Zhou, Q., 2023. Three amphioxus reference genomes reveal gene and chromosome evolution of chordates. Proceedings of the National Academy of Sciences 120, e2201504120.

Huerta-Cepas, J., Szklarczyk, D., Heller, D., Hernández-Plaza, A., Forslund, S.K., Cook, H., Mende, D.R., Letunic, I., Rattei, T., Jensen, L.J., von Mering, C., Bork, P., 2019. eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated orthology resource based on 5090 organisms and 2502 viruses. Nucleic acids research 47, D309-d314.

Janvier, P., 2006. Modern look for ancient lamprey. Nature 443, 921-923.

Jones, C.S., Cessi, P., 2017. Size Matters: Another Reason Why the Atlantic Is Saltier than the Pacific. Journal of Physical Oceanography 47, 2843-2859.

Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H., Maslen, J., Mitchell, A., Nuka, G., Pesseat, S., Quinn, A.F., Sangrador-Vegas, A., Scheremetjew, M., Yong, S.-Y., Lopez, R., Hunter, S., 2014. InterProScan 5: genome-scale protein function classification. Bioinformatics 30, 1236-1240.

Kapli, P., Natsidis, P., Leite, D.J., Fursman, M., Jeffrie, N., Rahman, I.A., Philippe, H., Copley, R.R., Telford, M.J., 2021. Lack of support for Deuterostomia prompts reinterpretation of the first Bilateria. Science Advances 7, eabe2741.

Kassahn, K.S., Dang, V.T., Wilkins, S.J., Perkins, A.C., Ragan, M.A., 2009. Evolution of gene function and regulatory control after whole-genome duplication: comparative analyses in vertebrates. Genome Res 19, 1404-1418.

Katoh, K., Misawa, K., Kuma, K.i., Miyata, T., 2002. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. Nucleic Acids Research 30, 3059-3066.

Koren, S., Walenz, B.P., Berlin, K., Miller, J.R., Bergman, N.H., Phillippy, A.M., 2017. Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. Genome Res 27, 722-736.

Korf, I., 2004. Gene finding in novel genomes. BMC Bioinformatics 5, 59.

Kumar, S., Stecher, G., Li, M., Knyaz, C., Tamura, K., 2018. MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. Mol Biol Evol 35, 1547-1549.

Letunic, I., Bork, P., 2006. Interactive Tree Of Life (iTOL): an online tool for phylogenetic tree display and annotation. Bioinformatics 23, 127-128.

Letunic, I., Bork, P., 2021. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display and annotation. Nucleic Acids Research 49, W293-W296.

Long, M., Betrán, E., Thornton, K., Wang, W., 2003. The origin of new genes: glimpses from the young and old. Nature Reviews Genetics 4, 865-875.

Louis, A., Roest Crollius, H., Robinson-Rechavi, M., 2012. How much does the amphioxus genome represent the

ancestor of chordates? Brief Funct Genomics 11, 89-95.

Marlétaz, F., Firbas, P.N., Maeso, I., Tena, J.J., Bogdanovic, O., Perry, M., Wyatt, C.D.R., de la Calle-Mustienes, E., Bertrand, S., Burguera, D., Acemel, R.D., van Heeringen, S.J., Naranjo, S., Herrera-Ubeda, C., Skvortsova, K., Jimenez-Gancedo, S., Aldea, D., Marquez, Y., Buono, L., Kozmikova, I., Permanyer, J., Louis, A., Albuixech-Crespo, B., Le Petillon, Y., Leon, A., Subirana, L., Balwierz, P.J., Duckett, P.E., Farahani, E., Aury, J.-M., Mangenot, S., Wincker, P., Albalat, R., Benito-Gutiérrez, È., Cañestro, C., Castro, F., D'Aniello, S., Ferrier, D.E.K., Huang, S., Laudet, V., Marais, G.A.B., Pontarotti, P., Schubert, M., Seitz, H., Somorjai, I., Takahashi, T., Mirabeau, O., Xu, A., Yu, J.-K., Carninci, P., Martinez-Morales, J.R., Crollius, H.R., Kozmik, Z., Weirauch, M.T., Garcia-Fernàndez, J., Lister, R., Lenhard, B., Holland, P.W.H., Escriva, H., Gómez-Skarmeta, J.L., Irimia, M., 2018. Amphioxus functional genomics and the origins of vertebrate gene regulation. Nature 564, 64-70.

McGinnis, S., Madden, T.L., 2004. BLAST: at the core of a powerful and diverse set of sequence analysis tools. Nucleic Acids Research 32, W20-W25.

Meyer, J., 1988. The evolution of ferredoxins. Trends in Ecology & Evolution 3, 222-226.

Millero, F.J., Feistel, R., Wright, D.G., McDougall, T.J., 2008. The composition of Standard Seawater and the definition of the Reference-Composition Salinity Scale. Deep Sea Research Part I: Oceanographic Research Papers 55, 50-72.

Millot, C., Taupier-Letage, I., 2005. Circulation in the Mediterranean Sea. In: Saliot, A. (Ed.), The Mediterranean Sea. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 29-66.

Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., von Haeseler, A., Lanfear, R., 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. Molecular Biology and Evolution 37, 1530-1534.

Mistry, J., Chuguransky, S., Williams, L., Qureshi, M., Salazar, Gustavo A., Sonnhammer, E.L.L., Tosatto, S.C.E., Paladin, L., Raj, S., Richardson, L.J., Finn, R.D., Bateman, A., 2021. Pfam: The protein families database in 2021. Nucleic Acids Research 49, D412-D419.

Peterson, K.J., Eernisse, D.J., 2016. The phylogeny, evolutionary developmental biology, and paleobiology of the Deuterostomia: 25 years of new techniques, new discoveries, and new ideas. Organisms Diversity & Evolution 16, 401-418.

Price, A.L., Jones, N.C., Pevzner, P.A., 2005. De novo identification of repeat families in large genomes. Bioinformatics 21, i351-i358.

Putnam, N.H., Butts, T., Ferrier, D.E., Furlong, R.F., Hellsten, U., Kawashima, T., Robinson-Rechavi, M., Shoguchi, E., Terry, A., Yu, J.K., Benito-Gutiérrez, E.L., Dubchak, I., Garcia-Fernàndez, J., Gibson-Brown, J.J., Grigoriev, I.V., Horton, A.C., de Jong, P.J., Jurka, J., Kapitonov, V.V., Kohara, Y., Kuroki, Y., Lindquist, E., Lucas, S., Osoegawa, K., Pennacchio, L.A., Salamov, A.A., Satou, Y., Sauka-Spengler, T., Schmutz, J., Shin, I.T., Toyoda, A., Bronner-Fraser, M., Fujiyama, A., Holland, L.Z., Holland, P.W., Satoh, N., Rokhsar, D.S., 2008. The amphioxus genome and the evolution of the chordate karyotype. Nature 453, 1064-1071.

Riisgård, H.U., Svane, I., 1999. Filter Feeding in Lancelets (Amphioxus), Branchiostoma lanceolatum. Invertebrate Biology 118, 423-432.

Rossman, T.G., Visalli, M.A., Komissarova, E.V., 2003. fau and its ubiquitin-like domain (FUBI) transforms human osteogenic sarcoma (HOS) cells to anchorage-independence. Oncogene 22, 1817-1821.

Satoh, N., Rokhsar, D., Nishikawa, T., 2014. Chordate evolution and the three-phylum system. Proc Biol Sci 281, 20141729.

Siddall, M.E., Whiting, M.F., 1999. Long-branch abstractions. Cladistics 15, 9-24.

Sievers, F., Higgins, D.G., 2014. Clustal Omega, accurate alignment of very large numbers of sequences. Methods Mol Biol 1079, 105-116.

Simakov, O., Kawashima, T., Marlétaz, F., Jenkins, J., Koyanagi, R., Mitros, T., Hisata, K., Bredeson, J., Shoguchi, E.,

Gyoja, F., Yue, J.-X., Chen, Y.-C., Freeman, R.M., Sasaki, A., Hikosaka-Katayama, T., Sato, A., Fujie, M., Baughman, K.W., Levine, J., Gonzalez, P., Cameron, C., Fritzenwanker, J.H., Pani, A.M., Goto, H., Kanda, M., Arakaki, N., Yamasaki, S., Qu, J., Cree, A., Ding, Y., Dinh, H.H., Dugan, S., Holder, M., Jhangiani, S.N., Kovar, C.L., Lee, S.L., Lewis, L.R., Morton, D., Nazareth, L.V., Okwuonu, G., Santibanez, J., Chen, R., Richards, S., Muzny, D.M., Gillis, A., Peshkin, L., Wu, M., Humphreys, T., Su, Y.-H., Putnam, N.H., Schmutz, J., Fujiyama, A., Yu, J.-K., Tagawa, K., Worley, K.C., Gibbs, R.A., Kirschner, M.W., Lowe, C.J., Satoh, N., Rokhsar, D.S., Gerhart, J., 2015. Hemichordate genomes and deuterostome origins. Nature 527, 459-465.

Simakov, O., Marlétaz, F., Yue, J.X., O'Connell, B., Jenkins, J., Brandt, A., Calef, R., Tung, C.H., Huang, T.K., Schmutz, J., Satoh, N., Yu, J.K., Putnam, N.H., Green, R.E., Rokhsar, D.S., 2020. Deeply conserved synteny resolves early events in vertebrate evolution. Nat Ecol Evol 4, 820-830.

Simão, F.A., Waterhouse, R.M., Ioannidis, P., Kriventseva, E.V., Zdobnov, E.M., 2015. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. Bioinformatics 31, 3210-3212.

Slater, G.S.C., Birney, E., 2005. Automated generation of heuristics for biological sequence comparison. BMC Bioinformatics 6, 31.

Stamatakis, A., 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. Bioinformatics 30, 1312-1313.

Suzek, B.E., Wang, Y., Huang, H., McGarvey, P.B., Wu, C.H., UniProt, C., 2015. UniRef clusters: a comprehensive and scalable alternative for improving sequence similarity searches. Bioinformatics 31, 926-932.

Tarailo-Graovac, M., Chen, N., 2009. Using RepeatMasker to identify repetitive elements in genomic sequences. Current protocols in bioinformatics Chapter 4, Unit 4.10.

Taschuk, F., Cherry, S., 2020. DEAD-Box Helicases: Sensors, Regulators, and Effectors for Antiviral Defense. Viruses 12.

Thompson, J.D., Higgins, D.G., Gibson, T.J., 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. Nucleic acids research 22, 4673-4680.

Thorvaldsdóttir, H., Robinson, J.T., Mesirov, J.P., 2013. Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. Brief Bioinform 14, 178-192.

Townsend, J.P., Su, Z., Tekle, Y.I., 2012. Phylogenetic Signal and Noise: Predicting the Power of a Data Set to Resolve Phylogeny. Systematic Biology 61, 835-835.

Walker, B.J., Abeel, T., Shea, T., Priest, M., Abouelliel, A., Sakthikumar, S., Cuomo, C.A., Zeng, Q., Wortman, J., Young, S.K., Earl, A.M., 2014. Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. PloS one 9, e112963.

Xiong, Q., Wan, A.T.-Y., Liu, X., Fung, C.S.-H., Xiao, X., Malainual, N., Hou, J., Wang, L., Wang, M., Yang, K.Y., Cui, Y., Leung, E.L.-H., Nong, W., Shin, S.-K., Au, S.W.-N., Jeong, K.Y., Chew, F.-T., Hui, J.H.-L., Leung, T.-F., Tungtrongchitr, A., Zhong, N., Liu, Z., Tsui, S.K.-W., 2022. Comparative Genomics Reveals Insights into the Divergent Evolution of Astigmatic Mites and Household Pest Adaptations. Molecular Biology and Evolution 39.

Yang, K.Y., Chen, Y., Zhang, Z., Ng, P.K.-S., Zhou, W.J., Zhang, Y., Liu, M., Chen, J., Mao, B., Tsui, S.K.-W., 2016. Transcriptome analysis of different developmental stages of amphioxus reveals dynamic changes of distinct classes of genes during development. Scientific Reports 6, 23195.

Yang, Z., 2007. PAML 4: phylogenetic analysis by maximum likelihood. Molecular biology and evolution 24, 1586-1591.

Yuan, D., Xu, A., 2016. Chapter 10 - Evolution of Inflammation-Related Lipid Metabolism Pathway and Inflammatory Lipid Signaling in Amphioxus. In: Xu, A. (Ed.), Amphioxus Immunity. Academic Press, pp. 189-205.

Yuan, D., Zou, Q., Yu, T., Song, C., Huang, S., Chen, S., Ren, Z., Xu, A., 2014. Ancestral genetic complexity of

arachidonic acid metabolism in Metazoa. Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids 1841, 1272-1284.

Zhang, Q.-L., Zhu, Q.-H., Liao, X., Wang, X.-Q., Chen, T., Xu, H.-T., Wang, J., Yuan, M.-L., Chen, J.-Y., 2016. Selection of reliable reference genes for normalization of quantitative RT-PCR from different developmental stages and tissues in amphioxus. Scientific Reports 6, 37549.

Zwaenepoel, A., Van de Peer, Y., 2018. wgd—simple command line tools for the analysis of ancient whole-genome duplications. Bioinformatics 35, 2153-2155.

Fig. 1. Phylogenetic analysis of major deuterostome groups

The completeness of 29 genome assemblies (Table S2) was assessed by Benchmarking Universal Single-Copy Orthologs (BUSCO) using database metazoa_odb9. 112 single-copy BUSCOs overlapped by 29 genome assemblies were extracted in protein sequences and aligned by MAFFT for the phylogenetic analysis by RAxML with 100 bootstrap replicates (Supplementary Note 2.2).

Fig. 2. A group of lipoxygenase (LOX) genes in amphioxus

(**A**) The amphioxus LOX genes and their closest genes in the UniRef50 database from other taxonomic categories, including bacteria, fungi, other eukaryotes (excluding metazoa, plants, and fungi) and other metazoa (excluding Amphioxiformes) were collected for phylogenetic analysis. In this maximum likelihood phylogenomic tree, SH-aLRT and ultrafast bootstrap support values are given in that order on the branches. (**B**) Phylogenetic analysis of all LOX genes of four amphioxus and their domain architecture. The LOX gene of the sea lamprey *Petromyzon marinus* (GenBank accession: XP_032817193.1) was used as outgroup. C2, C2-like domain (universal calcium-binding domain); SRPBCC, PYR/PYL/RCAR family domain in SRPBCC superfamily. (**C**) Gene synteny alignment of the LOX genes of four amphioxus. The distance precision was set as 5 Kb, e.g., 30 Kb represents for 30-35 Kb. If not noted, a slash (/) between two genes represents 5 Kb, e.g., one slash represents 5-10 Kb. Except the outlier (*LOX_Bf_9*), LOXs located on clades supported by high bootstrap value (>95) in the phylogenetic tree were painted in the same color. The suffixes stand for four amphioxus, *Bbb* for *B. belcheri* beihai, *Bbx* for *B. beicheri* xiamen, *Bf* for *B. floridae* and *Bl* for *B. lanceolatum*. The grey turnover arrow means the gene alignment is after reverse complement.

Fig. 3. Comparative analysis of two gene novelties in amphioxus

(A) The DEAD/DEAH-box helicase (DBH) genes of amphioxus and their closest genes in the UniRef50 database from other taxonomic categories, including bacteria, fungi, other eukaryotes (excluding metazoa, plants, and fungi) and other metazoa (excluding Amphioxiformes) were collected for phylogenetic analysis. **(B)** Alignment of splice sites of the DBH genes of amphioxus, the charophytic algae *K. nitens* (GenBank accession: GAQ83283.1), the soft coral *Dendronephthya gigantea* (GenBank accession: XP_028406807.1) and the hemichordate *Saccoglossus kowalevskii* (GenBank accession: XP_002737002.1). Conserved splice sites were marked by vertical lines (Fig. S8), and three conserved splice sites of all seven sequences were highlighted in yellow triangles. Two unusual splice sites were labeled as S1 and S2. **(C)** The alcohol dehydrogenase (ADH) genes of amphioxus and their closest genes in the UniRef50 database from other taxonomic categories were collected for phylogenetic analysis. **(D)** Alignment of splice sites of the ADH genes of amphioxus, the sponge *Amphimedon queenslandica* (GenBank accession: XP_011406967.2) and the copepod *Eurytemora affinis* (UniProt accession: UPI000C762174) (Fig. S11). In these maximum likelihood phylogenomic trees, SH-aLRT and ultrafast bootstrap support values are given in that order on the branches.

Fig. 4. A group of proximally arrayed EF-hand calcium-binding proteins (EF-CaBPs)
(A) Of a group of EF-CaBP genes of amphioxus, bit scores of the best hits in the UniRef50 database to other metazoa, other deuterostomes and other chordates (excluding Amphioxiformes respectively) were compared with those to bacteria in violin plot with quartile lines. **** p <0.0001, when comparing the best hits to bacteria with other three groups, respectively. **(B)** Correlation of the gene number and the sea surface salinity of collection locations of four amphioxus. A darker color indicates higher sea salinity. **(C)** Phylogenetic analysis of these EF-CaBPs of four amphioxus. Two large clades, Clade A and B, were highlighted and five specific clusters were marked. **(D)** Gene synteny alignment of the proximally arrayed EF-CaBP genes. A slash (/) represents 5 Kb, e.g., one slash represents 5-10 Kb, and three slashes represent 15-20 Kb, while none represents below 5 Kb. The grey turnover arrow means after reverse complement.