Article Category:

Original article

Title:

Extremely low level of genetic diversity in *Gentiana yakushimensis*, an endangered species in

Yakushima Island, Japan

Naohiro I. Ishii[1,2*], Shun K. Hirota[1], Yoshihiro Tsunamoto[1,3], Ayumi Matsuo[1], Harue Abe[4],

Yoshihisa Suyama[1]

[1]Field Science Center, Graduate School of Agricultural Science, Tohoku University, 232-3

Yomogida, Naruko-onsen, Osaki, Miyagi, 989-6711, Japan

[2]Graduate School of Environment and Information Sciences, Yokohama National University,

79-7 Tokiwadai, Hodogaya, Yokohama, Kanagawa, 240-8501, Japan

[3]Research Institute of Energy, Environment, and Geology, Hokkaido Research Organization,

Kita19-Nishi12, Kita-ku, Sapporo, Hokkaido, 060-0819, Japan

[4]Sado Station, Sado Island Center for Ecological Sustainability, Niigata University, 94-2 Koda,

Sado, Niigata 952-2206, Japan

[*]**Correspondence to:**

Naohiro I. Ishii

Graduate School of Environment and Information Sciences, Yokohama National University,

79-7 Tokiwadai, Hodogaya, Yokohama, Kanagawa 240-8501, Japan

E-mail: na0hiro1shi1@gmail.com

Abstract

Rare or endangered species largely contributing to global biodiversity are essential components of ecosystems. Because genetic diversity depends not only on apparent population size but also demographic processes, the conservation priority of endangered species based on population genetics needs to be assessed by comparison with common congeners and demographic events. In this study, we performed population genetic analysis for *Gentiana yakushimensis* Makino, a rare and endangered plant species with an isolated distribution on Yakushima Island, Japan. We performed comparison of genetic diversity with the congener (*Gentiana triflora* Pall. var. *japonica* (Kusn.) H.Hara), demographic inference, and bottleneck test based on genome-wide single nucleotide polymorphisms. The result was that *G. yakushimensis* had an extremely low genetic diversity level and a high inbreeding coefficient level compared to those of the congener. The population of *G. yakushimensis* was estimated to experience an expansion during the Last Glacial Maximum and a recent bottleneck by the demographic inference and the bottleneck test. Therefore, the low level of genetic diversity of this species might have resulted from the impact of the recent bottleneck and the long-term maintenance of a small population size. Due to the long lifespan of the genus *Gentiana*, the current level of genetic diversity probably does not reflect a large part of recent demographic events, which requires long-term monitoring of changes in genetic diversity to identify the time lag between the reduction in apparent population size and genetic diversity as a future outlook.

Keywords

Endangered species, endemic species, genome-wide SNPs, Gentiana, MIG-seq

Introduction

Rare or endangered species, largely contributing to global biodiversity, are essential

components of various ecosystems (Enquist et al., 2019). Most previous studies have demonstrated that rare species with limited geographic distributions tend to have lower levels of genetic diversity than common species because of their small effective population size (Broadhurst et al., 2017; Chung et al., 2018; Gibson, Rice, & Stucke, 2008; Hamrick & Godt, 1996; Nybom & Bartish, 2000). The low level of genetic diversity in a species or a population can trigger its vulnerability because it could generally result in the reduction in adaptive potential to environmental changes and resilience or resistance to environmental extremes (Reusch, Ehlers, Hammerli, & Worm, 2005; Willi, Van Buskirk, & Hoffmann, 2006). Inbreeding depression associated with low genetic diversity can also reduce reproductive fitness (Losdat, Chang, & Reid, 2014; O'Grady et al., 2006; Reed & Frankham, 2003). Thus, it is reasonable to globally evaluate the conservation priority of rare or endangered species based on apparent population size and genetic diversity (Garner, Hoban, & Luikart, 2020). Accumulation of genetic information for thousands of endangered species can lay the foundation for biodiversity conservation and conservation prioritization.

The genetic diversity of rare species is not always low, and it can be species-specific because it depends on apparent population size and demographic processes or life history traits (Ellegren & Galtier, 2016; Hamrick & Godt, 1996; Nybom & Bartish, 2000; Thiel-Egenter et al., 2009). For example, previous studies have demonstrated that species with a long lifespan tend to have higher levels of genetic diversity than those with short lifespans (Kort et al., 2021; Reisch & Rosbakh, 2020). Past and current demographic events, such as population bottlenecks or expansion due to environmental and anthropogenic fluctuations, can affect the current genetic diversity while comparing closely related species with similar life history traits (Ellegren & Galtier, 2016). For example, the species widely distributed in the past can become rare or endangered due to disturbances, resulting in the low level of genetic diversity (e.g. Li et al., 2014). Conversely, in isolated habitats such as islands, a small population size can be

maintained for long term due to limited immigrants, which was estimated to be a factor of low

genetic diversity (Gonzalez, Cerón-Souza, Mateo, & Zardoya, 2014). Therefore, estimating

demographic history allows us to examine which current level of genetic diversity is determined

by historical or recent events.

Alpine plants, including many rare or endangered species, are the targets for which

conservation prioritization based on genetic aspects is required (Noroozi et al., 2018; Perrigo,

Hoorn, & Antonelli, 2020), as these plants are vulnerable to climate change and anthropogenic

disturbances (Guisan & Theurillat, 2000; Thuiller, Lavorel, Araújo, Sykes, & Prentice, 2005).

Previous studies have demonstrated that the current levels of genetic diversity of alpine plants

are strongly affected by several demographic events, such as expansion or reduction of

distribution range after the Last Glacial Maximum (LGM; 21,000 years ago) (Loïc Pellissier et

al., 2016), geographical isolation among populations by the distribution patterns restricted to

high-altitude areas (Reisch & Rosbakh, 2020), and direct human activities such as trampling by

tourists and illegal digging (Banks et al., 2013; Feng, Liu, Chiang, & Gong, 2017). Therefore,

inferring historical demography allows us to examine the influence of historical demographic

events on current genetic diversity levels and accurately assess the conservation status of rare

alpine species.

Gentiana yakushimensis Makino (Figure 1) is an endangered plant species in Japan

(Japanese Ministry of Environment, 2020) and endemic to Yakushima Island (Iwatsuki,

Yamazaki, Boufford, & Ohba, 1993). On Yakushima Island, designated a UNESCO World

Heritage Site, this species grows only in limited alpine zones covered by granite rocks with low

vegetation cover in the central mountains (Figure 2). Illegal digging for ornamental horticulture

has drastically reduced its population (Japanese Ministry of Environment, 2020). *G.*

yakushimensis, or its ancestor, was estimated to colonize Japan 4–1 million years ago with the

Qinghai–Tibet Plateau as the origin (Appendix S3 in Favre et al., 2016). The taxonomic position

of *G. yakushimensis* is discussed. Recently, phylogenetic analysis based on chloroplast DNA

regions suggested that *G. yakushimensis* is close to the section *Pneumonanthe* (Mishiba et al.,

2009). Moreover, phylogenetic analysis covering all *Gentiana* sections supported the inclusion

of *G. yakushimensis* in *the Pneumonanthe*/*Cruciata* lineage (Favre et al., 2016). The previous

phylogeographic studies of alpine species have demonstrated range shifts or persistence during

and after the LGM not only in Japan (e.g Ikeda, Higashi, Yakubov, Barkalov, & Setoguchi,

2014; Ikeda, Yakubov, Barkalov, Sato, & Fujii, 2020), but also globally (e.g Allen, Marr,

Mccormick, & Hebda, 2015; Ikeda et al., 2017; Schönswetter, Stehlik, Holderegger, & Tribsch,

2005). Thus, by estimating the population demography of *G. yakushimensis*, which distributed

in the subalpine region, we can examine whether the current level of genetic diversity resulted

from recent and/or historical demographic events.

Here, we assessed the genetic diversity and demographic history of *G. yakushimensis*

based on genome-wide single nucleotide polymorphisms (SNPs) to identify their conservation

status. We performed 1) the comparison of genetic diversity with a congener (*Gentiana triflora*

Pall. var. *japonica* (Kusn.) H.Hara), 2) the evaluation of genetic differentiation across

Yakushima Island, and 3) the demographic inference of *G. yakushimensis* to identify whether

the determinant of the current level of genetic diversity is recent or historical, aiming to provide

the genetic information of the endangered species, supporting decision-making in conservation

implications, and prioritizing.

Materials and Methods

Studied species

G. yakushimensis (Gentianaceae) is a perennial herb endemic to Yakushima Island,

Japan (Chen & Wang, 1999; Ho & Liu, 2001; Iwatsuki et al., 1993). Although Ho (1988)

recorded the occurrence of *G. yakushimensis* in Taiwan, China based on a single specimen (PE

No. 81854; https://www.cvh.ac.cn/spms/detail.php?id=ec90580f), Chen and Wang (1999) denied the distribution of this species in Taiwan because the specimen lacks collection labels and no other specimens from Taiwan exist. *G. yakushimensis* has a minimal and patchy distribution; it frequently appears on granite rocks around the tops of the central mountains, at altitudes ranging from 1,700 to 1,900 m (Figure 2). *G. yakushimensis* was pollinated by bees, which were likely to be Andrenidae and Halictidae (H. Abe, personal observation), and its seeds with no wings were assumed to be dispersed by gravity and wind. The population size of *G. yakushimensis* has decreased due to illegal digging in recent years (Japanese Ministry of the Environment, 2014). In addition, feeding damage by sika deer (*Cervus nippon yakushimae*) might also be a factor in population size reduction. In Yakushima Island, grazing by sika deer has affected vegetation from lowlands to subalpine zones (Takatsuki, 1990; Tsujino & Yumoto, 2004). Takatsuki (1990) demonstrated that sika deer relied on forbs for about 5 % of their total diet in the subalpine area of Yakushima Island, indicating that *G. yakushimensis* could be affected by grazing. Thus, *G. yakushimensis* has been listed in the Japanese Red List as Endangered (EN) (Japanese Ministry of Environment, 2020) and conserved under the "Act on Conservation of Endangered Species of Wild Fauna and Flora" since 2016.

Gentiana triflora var. *japonica* was used to compare the genetic diversity of *G. yakushimensis*. *G. triflora* var. *japonica* is a relatively common species widely distributed in the mountainous area from Hokkaido to the Kinki region of Japan (Figure 2). This species is one of the most closely related to *G. yakushimensis* based on the chloroplast DNA region (Favre et al., 2016; Mishiba et al., 2009).

We collected leaf samples from all individuals of *G. yakushimensis* (560 individuals), which had grown in accessible habitats, for DNA analysis at Yakushima Island in August from 2009 to 2010. Small individuals or individuals with few leaves were not collected to conserve *G. yakushimensis*. When multiple individuals grew at the same coordinates, one individual was

151   selected for DNA analysis. Leaf samples were collected with permission from the prefecture

152   and ministries. The leaf samples of *G. triflora* were collected from nine individuals as congeners

153   from an area of approximately 100 m^2 in Osaki, Miyagi Prefecture, in the northern area of Japan

154   (Figure 2).

156   ***DNA extraction, amplification, and sequencing***

157   Total genomic DNA was extracted from frozen leaf samples using the modified

158   cetyltrimethylammonium bromide (CTAB) method (Doyle, 1991). For DNA analysis, we used

159   353 and 9 samples of *G. yakushimensis* and *G. triflora*, respectively (Table 1). The samples of

160   *G. yakushimensis* for the analysis were at least 3 m away from each other because whether the

161   different shoots in granite crevices were from the same individual could not be clearly

162   distinguished.

163   To evaluate the genetic diversity of *G. yakushimensis* and *G. triflora*, we performed *de*

novo single-nucleotide polymorphism (SNP) detection using multiplexed inter-simple sequence

repeat (ISSR) genotyping by sequencing (MIG-seq) (Suyama et al., 2021; Suyama & Matsuki,

2015). We prepared the MIG-seq library and sequenced it following the protocol described by

Suyama et al. (2021). Briefly, ISSRs were amplified using primer set-1 (Suyama & Matsuki,

2015) in the first PCR, and then the index and adaptor sequences were added to both ends of

the 1st PCR products in the second PCR. We sequenced 80 base pair-end reads using the

Illumina MiSeq platform and MiSeq Reagent Kit v3 (150 cycles) (Illumina, San Diego, CA,

USA). We skipped detection of the first 17 bases of Reads 1 and 2 using the "DarkCycle" option.

Read assembling and SNP marker filtering

174   One base at each end of the processed reads was trimmed, and the reads were filtered

based on the quality of reads and the adaptor sequences using Trimmomatic v.0.3.2 (Bolger,

Lohse, & Usadel, 2014). Therefore, we performed the subsequent analysis using filtered reads

with a length of 78 bp. Stacks v2.52 (Catchen, Amores, Hohenlohe, Cresko, & Postlethwait,

2011) which was used to process the filtered reads with the default parameter settings: the

minimum number of identical reads required to create a stack (m = 3), the nucleotide

mismatches between loci within a single individual (M = 2), and mismatches between loci when

building the catalog (n = 1). The SNP genotype for each individual was exported with the

following parameter setting using the 'populations' command in Stacks v2.52: the maximum

observed heterozygosity, --max-obs-het = 0.6; minimum minor allele frequency, --min-maf = 2

/ the number of individuals. For the dataset of both species (common loci) and individual

species (common and species-specific loci; hereafter, referred to as "all loci"), the SNP

genotype was exported by setting r (the parameter of the minimum sharing rate of SNP among

individuals per population) at 0.5, 0.6, and 0.7, in addition to the above parameter settings to

compare the genetic diversity of *G. yakushimensis* and *G. triflora*. We set up each species as a

population to meet the r threshold for each species (p = 2 for the dataset of common loci and p

= 1 for all loci in the 'populations' command). The comparison with the different values of the

parameter r was performed because the number of SNPs of the common loci was low (less than

100) when the parameter r was 0.7. The estimation of genetic indices with low parameter r

values leads to increase in missing data but ensures accuracy by increasing the number of SNPs

(Hodel et al., 2017).

Comparison of genetic diversity between G. yakushimensis and G. triflora

Genetic diversity was compared between *G. yakushimensis* and *G. triflora* for both the

common and all loci. We calculated nucleotide diversity (π) based on fixed and variant sites

using the 'populations' command of Stacks v2.52. The other indices, observed and expected

heterozygosity (H_O and H_E), and inbreeding coefficient (F_{IS}), were determined using GenAlEx

v6.51 (Peakall & Smouse, 2006). Wilcoxon rank sum tests were performed on these indices to

test the significance of the difference between the species.

Genetic differentiation of G. yakushimensis

Subsequent analyses were performed using the dataset of all *G. yakushimensis* loci.

Kosman & Leonard's (2005) pairwise genetic distance between *G. yakushimensis* individuals

was calculated using the R package PopGenReport (Adamack & Gruber, 2014). Based on

genetic and geographic distance matrices among individuals, we assessed spatial

autocorrelation (isolation by distance, IBD) across the distribution area using the

'mantel.correlog' function in the R package vegan (Oksanen et al., 2020). To test for the

presence of genetic structure, we performed principal coordinate analysis (PCoA) based on the

genetic distance matrix using the 'pcoa' function in the R package ape (Paradis & Schliep,

2019).

Inference of population demographic history

We inferred the population demographic history of *G. yakushimensis*. In the

'populations' command in Stacks v2.52, we exported SNP genotypes for the dataset of the all

loci of *G. yakushimensis* with the change of the following parameter: minimum minor allele

frequency, --min-maf = 0. Population demography was inferred by Stairway plot v2.1 (Liu &

Fu, 2015), a model-flexible method of inferring historical changes in population size based on

folded site frequency spectrums (SFS). SFSs were calculated using easySFS

(https://github.com/isaacovercast/easySFS), and samples were projected downward to

maximize the number of loci without missing data and number of individuals retained (Kreiner

et al., 2019). The parameter settings were as follows: 67% of sites for training; 15 years per

generation; 4.089×10^{-8} of mutation rate. We assumed that the lifespan (generation time) of *G.*

yakushimensis is 15 years, according to which the lifespan was at least 15 years in a congener,

G. pneumonanthe (Oostermeijer, Veer, & Nijs, 1994). The mutation rate was calculated by the

relationship between genome size and mutation rate (Lynch, 2010), following Garot, Joët,

Combes, Severac, & Lashermes (2019). The mean genome size of the genus *Gentiana* (3640.7

Mb) in the plant DNA C-values database (Pellicer & Leitch, 2020) used for this analysis.

Finally, the signatures of recent genetic bottleneck were assessed using the

heterozygote excess test implemented in the software BOTTLENECK (Cornuet & Luikart,

1996; Piry, Luikart, & Cornuet, 1999) by INEST v2.2 (Chybicki & Burczyk, 2009). This

method compares H_E in an empirical population to the heterozygosity that is expected in a

population at mutation-drift equilibrium (H_{Eq}). A transitory excess in H_E (measured as $\Delta H = H_E$

- H_{Eq}) is assumed in recently bottlenecked populations. The significance of the heterozygote

excess was evaluated using Wilcoxon signed-rank tests under an infinite alleles model (IAM)

(Maruyama & Fuerst, 1985). Although IAM is not entirely appropriate for biallelic SNPs, this

test was exploratorily conducted following recent studies (Camak, Osborne, & Turner, 2021;

Kogura et al., 2011; Setzke, Wong, & Russello, 2022; Stojanova, Šurinová, Zeisek,

Münzbergová, & Pánková, 2020).

Results

The mean number of filtered reads per sample of *G. yakushimensis* and *G. triflora* was

92,960 and 58,900, respectively. The number of SNPs in the common loci between the two

species was 72, 51, and 34, with r = 0.5, 0.6, and 0.7, respectively (Table 1). In the common

and species-specific (all) loci of *G. yakushimensis*, the number of SNPs was 784, 562, and 402

with r = 0.5, 0.6, and 0.7, respectively (Table 1). On the other hand, the number of SNPs in all

loci of *G. triflora* was 401, 297, and 202, with r = 0.5, 0.6, and 0.7, respectively (Table 1).

Compared to *G. triflora*, the genetic diversity indices (π, H_O, and H_E) of *G.*

yakushimensis were significantly lower in all parameter settings (Table 1). When comparing *G.*

yakushimensis with *G. triflora* for common loci with setting r = 0.7 in the 'populations'

command, the former was 0.05 times for π and H_E, and 0.03 times for H_O as much as the latter

(Table 1). The inbreeding coefficients (F_{IS}) of *G. yakushimensis* were significantly greater than

zero for all parameter settings (Table 1). Also, it was significantly greater than those of *G.*

triflora except for the common loci with setting r = 0.5 in the 'populations' command.

We observed a significant positive correlation between genetic and geographic

distances among the individuals of *G. yakushimensis* (Mantel correlation coefficient, *r* = 0.25–

0.27, *P* < 0.001) (Figure 3a). However, an increase in genetic differentiation among the

individuals was not detected when the pairwise geographical distance was over approximately

500 m. As a result of the Mantel correlogram, calculating Mantel correlation coefficients for

each distance class, consistently positive correlations were shown up to approximately 140–

160 m (Figure 3b). PCoA did not demonstrate population genetic structure across the

distribution area, although the first three axes of PCoA explained only 4–6 % of the whole

genetic variation of *G. yakushimensis* (Figure 4, and figure S1).

As a result of the demographic inference by the Stairway plot, the expansion of

population size at 10,000–40,000 years ago and the constant population size until recent years

were estimated (Figure 5). The median of the population size of 5,000–6,000 individuals was

inferred to be maintained for at least 10,000 years. In addition, the 95% confidence intervals of

population size were expanded downward from 1,700 to 10,000 at about 50 years ago (34

generations ago). These patterns were consistent among with the different setting of the

parameter (r = 0.5, 0.6, and 0.7 in the 'populations' command in Stacks). As the results of the

test of genetic bottleneck, the median of ΔH was significantly higher than zero (H_E was higher

than H_{Eq}) at each setting of the parameter (r = 0.5, 0.6, and 0.7), which exhibited that the

population of *G. yakushimensis* showed significant heterozygote excess (Figure 6).

Discussion

In this study, we revealed 1) the extremely low level of genetic diversity of *G. yakushimensis*, 2) the gentle genetic differentiation across the Yakushima population despite the patchy distribution, and 3) the small population size due to the recent bottleneck and no dramatic increase in population size over the long term.

We demonstrated that *G. yakushimensis* had extremely low levels of genetic diversity and a high level of inbreeding coefficient compared to those of its congener (Table 1). Surprisingly, the genetic diversity across only nine individuals from a single population of *G. triflora* exceeded that across 353 individuals of *G. yakushimensis*. This result was consistent with that of previous studies that showed that relatively rare species have lower genetic diversity than that of common species (Broadhurst et al., 2017; Chung et al., 2018; Gibson et al., 2008; Hamrick & Godt, 1996; Nybom & Bartish, 2000). The limited gene flow across a narrow range (details in the next paragraph) might resulted in inbreeding among closely related individuals. Because inbreeding can lead to the reduction in reproductive fitness (Losdat et al., 2014; O'Grady et al., 2006; Reed & Frankham, 2003), the negative impacts on population dynamics of *G. yakushimensis* related to inbreeding depression is a concern in the near future. The decrease in patch size and connectivity between patches might result in further excellence in inbreeding.

There was less genetic differentiation among the fragmented patches of *G. yakushimensis* across Yakushima Island. IBD was significant only across the narrow range (up to 140–160 m), and the constant pattern of pairwise genetic distance was detected over this range, suggesting that gene flow occurs across the 140–160 m range. The granitic rocks, the habitats of *G. yakushimensis*, are densely distributed in the center of Yakushima Island. Thus, there is gene flow between adjacent habitats, and the patchy habitat network of *G.*

yakushimensis might be maintained. These results indicate that *G. yakushimensis* on Yakushima

Island comprises almost a genetically single population. *Gentiana yakushimensis* was mainly

pollinated by bees (H. Abe, personal observation). The foraging range of bees was at least 100

m for species with the smallest body size (about 1 mm of intertegular span) (Greenleaf,

Williams, Winfree, & Kremen, 2007). Generally, bees prioritize the foraging of neighboring

flower patches after foraging within a patch (Woodgate, Makinson, Lim, Reynolds, & Chittka,

2017). Thus, the pollination behavior by bees might gently connect patchy habitats and maintain

a low level of genetic differentiation.

The population of *G. yakushimensis* was likely to experience the range expansion

during or after the LGM and the recent reduction of effective population size according to the

demographic inferences and the bottleneck tests (Figure 5 and figure 6). This result was

consistent with the previous studies which demonstrated the range expansion of alpine plants

during or after the LGM (Ikeda et al., 2020; Ikeda, Yakubov, Barkalov, & Setoguchi, 2018;

Schönswetter et al., 2005). Although these previous studies showed the range expansion into

the areas that were glaciated during the LGM, Yakushima Island was not glaciated during the

LGM (Tsukada, 1985). Based on the results of the present study, it was not possible to

distinguish whether the increase in population size was due to a decrease in the vegetation and

forest limit associated with the cold climate during the LGM and/or an expansion into the area

with the highest altitude related to the warming after the LGM. The low level of genetic

diversity in *G. yakushimensis* was assumed to be due to the significant impact of the recent

bottleneck. There are three possible reasons for this result: illegal digging (Japanese Ministry

of Environment, 2020), grazing by sika deer (Takatsuki, 1990, 2009; Tsujino & Yumoto, 2004),

and a habitat area reduction by the uplift of forest and vegetation lines by warming (Japanese

Ministry of the Environment, 2017). The high intensity of grazing by mammalian herbivores

negatively affected the maintenance of population in the congener of *G. yakushimensis*

(Moschetti, Besnard, Couturier, & Fonderflick, 2020). The high density of the sika deer in

Yakushima Island (Tsujino & Yumoto, 2004) might lead to the decrease in population size. In

Yakushima Island, because the forest cover and density of *C. japonica* increased in the high-

altitude area from 1963 to 2014 (Japanese Ministry of the Environment, 2017), it is inferred

that the increase in the vegetation limit of shrubs and dwarf bamboos reduces the habitat area

of *G. yakushimensis*. Because the seedling establishments in the *Gentiana* species can be

limited by the high level of light competition (Ekrtová & Košnar, 2012; Miller, Geddes, &

Mardon, 1998), the uplift of vegetation cover is one of the factors strongly influencing the

population size via competitions. In addition, because the maximum population size of *G.*

yakushimensis estimated to be approximately 9,000 (Figure 5), one of the possible reasons for

low genetic diversity is the long-term maintenance of a small population size. While the recent

bottleneck was supported by the demographic inferences, the reduction in population size based

on the estimated median was low level (Figure 5), which might be attributed to the long lifespan

of the genus *Gentiana* (more than 15 years) (Oostermeijer, Veer, et al., 1994). The longevity

generally causes a time lag between reducing the apparent population size and the decline in

genetic diversity (Fuller & Doyle, 2018; Reisch et al., 2017). This time lag might lead to a

further reduction in the genetic diversity of *G. yakushimensis* in the near future.

G. yakushimensis has an extraordinarily low level of genetic diversity, resulting in

vulnerability to environmental changes and human disturbances (Reusch et al., 2005; Willi et

al., 2006). Moreover, the effective range of gene flow in this species is extremely narrow. In

species with limited dispersal ability, a high level of patch connectivity can enhance gene flow,

decreasing the extinction probability (Dornier & Cheptou, 2012). In the congener of *G.*

yakushimensis, the low level of genetic diversity resulted in the reduction in offspring fitness

(Oostermeijer, van Eijck, & den Nijs, 1994). Thus, the loss of small patches may threaten the

maintenance of fitness in the entire population. Therefore, we emphasize the conservation of

large and small patches of *G. yakushimensis*. In the present study, we provided information on

the genetic diversity and population demography of *G. yakushimensis*. The accumulation of the

genetic information of a lot of endangered species can be useful for the conservation

prioritization (Garner et al., 2020), such as the conservation project of plant species with

extremely small populations in China (Ma et al., 2013; Ren et al., 2012). For an outlook, we

require the assessment of population stability by long-term monitoring for the changes in

genetic diversity and population size to identify the factors influencing the reduction in genetic

diversity and the length of the time lag between the decrease in apparent population size and

effective population size, since a drastic reduction in effective population size can occur in the

near future.

Acknowledgments

The authors are grateful to Yuji Isagi, the project leader of the Environment Research

and Technology Development Fund. The authors would like to thank Kenshi Tetsuka, Arata

Yoichi, Toshihiro Saito, Tatsuko Tetsuka, and Kengo Fuse for sampling plant materials. We also

thank our laboratory colleagues for helping with the analysis and an earlier draft of this

manuscript, especially Kenji Seiwa, Yu Fukasawa, Kimiyo Matsukura, Takuya Konuma, and

Rimpei Nagaoka. This research was supported by the Environment Research and Technology

Development Fund (F-091, 4-1605 and 4-1902) of the Environmental Restoration and

Conservation Agency of Japan. Permission to conduct research in the study area was provided

by Kagoshima Prefecture, the Japanese Ministry of Environment, the Japanese Ministry of

Education, Culture, Sports, Science and Technology, and the Japanese Forestry Agency.

Conflict of Interest

The authors have no conflicts of interest directly relevant to the content of this article.

Author's Contributions

N.I.I. performed the molecular experiments and analyses and led the writing of the manuscript; Y.T., A.M., and H.A. contributed to the molecular experiments and analyses; S.K.H. and Y.S. contributed to the analyses and writing; and N.I.I., H.A., and Y.S. designed the research.

References

Adamack, A. T., & Gruber, B. (2014). PopGenReport: Simplifying basic population genetic analyses in R. *Methods in Ecology and Evolution*, *5*, 384–387. doi: 10.1111/2041-210X.12158

Allen, G. A., Marr, K. L., Mccormick, L. J., & Hebda, R. J. (2015). Geographical origins, migration patterns and refugia of *Sibbaldia procumbens*, an arctic-alpine plant with a fragmented range. *Journal of Biogeography*, *42*, 1665–1676. doi: 10.1111/jbi.12543

Banks, S. C., Cary, G. J., Smith, A. L., Davies, I. D., Driscoll, D. A., Gill, A. M., … Peakall, R. (2013). How does ecological disturbance influence genetic diversity? *Trends in Ecology and Evolution*, *28*, 670–679. doi: 10.1016/j.tree.2013.08.005

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics*, *30*, 2114–2120. doi: 10.1093/bioinformatics/btu170

Broadhurst, L., Breed, M., Lowe, A., Bragg, J., Catullo, R., Coates, D., … Byrne, M. (2017). Genetic diversity and structure of the Australian flora. *Diversity and Distributions*, *23*, 41–52. doi: 10.1111/ddi.12505

Camak, D. T., Osborne, M. J., & Turner, T. F. (2021). Population genomics and conservation of Gila Trout (*Oncorhynchus gilae*). *Conservation Genetics*, *22*, 729–743. doi: 10.1007/s10592-021-01355-0

Catchen, J. M., Amores, A., Hohenlohe, P., Cresko, W., & Postlethwait, J. H. (2011). Stacks: Building and genotyping loci *de novo* from short-read sequences. *G3: Genes, Genomes, Genetics*, *1*, 171–182. doi: 10.1534/g3.111.000240

Chen, C.-H., & Wang, J.-C. (1999). Revision of the genus *Gentiana* L. (Gentianaceae) in Taiwan. *Botanical Bulletin of Academia Sinica*, *40*, 9–38.

Chung, M. Y., López-Pujol, J., Son, S., Suh, G. U., Yukawa, T., & Chung, M. G. (2018). Patterns of genetic diversity in rare and common orchids focusing on the Korean peninsula: Implications for conservation. *Botanical Review*, *84*, 1–25. doi:

10.1007/s12229-017-9190-5

Chybicki, I. J., & Burczyk, J. (2009). Simultaneous estimation of null alleles and inbreeding

coefficients. *Journal of Heredity*, *100*, 106–113. doi: 10.1093/jhered/esn088

Cornuet, J. M., & Luikart, G. (1996). Description and power analysis of two tests for

detecting recent population bottlenecks from allele frequency data. *Genetics*, *144*, 2001–

2014. doi: 10.1093/genetics/144.4.2001

Dornier, A., & Cheptou, P. O. (2012). Determinants of extinction in fragmented plant

populations: *Crepis sancta* (Asteraceae) in urban environments. *Oecologia*, *169*, 703–

712. doi: 10.1007/s00442-011-2229-0

Doyle, J. (1991). DNA protocols for plants. *Molecular Techniques in Taxonomy*, *57*, 283–

293. doi: 10.1007/978-3-642-83962-7_18

Ekrtová, E., & Košnar, J. (2012). Habitat-related variation in seedling recruitment of

Gentiana pannonica. *Acta Oecologica*, *45*, 88–97. doi: 10.1016/j.actao.2012.10.003

Ellegren, H., & Galtier, N. (2016). Determinants of genetic diversity. *Nature Reviews*

Genetics, *17*, 422–433. doi: 10.1038/nrg.2016.58

Enquist, B. J., Feng, X., Boyle, B., Maitner, B., Newman, E. A., Jørgensen, P. M., … McGill,

B. J. (2019). The commonness of rarity: Global and future distribution of rarity across

land plants. *Science Advances*, *5*, 1–14. doi: 10.1126/sciadv.aaz0414

Favre, A., Michalak, I., Chen, C. H., Wang, J. C., Pringle, J. S., Matuszak, S., … Muellner-

Riehl, A. N. (2016). Out-of-Tibet: The spatio-temporal evolution of *Gentiana*

(Gentianaceae). *Journal of Biogeography*, *43*, 1967–1978. doi: 10.1111/jbi.12840

Feng, X., Liu, J., Chiang, Y. C., & Gong, X. (2017). Investigating the genetic diversity,

population differentiation and population dynamics of *Cycas segmentifida* (Cycadaceae)

endemic to southwest china by multiple molecular markers. *Frontiers in Plant Science*,

8, 1–14. doi: 10.3389/fpls.2017.00839

Fuller, M. R., & Doyle, M. W. (2018). Gene flow simulations demonstrate resistance of long-

lived species to genetic erosion from habitat fragmentation. *Conservation Genetics*, *19*,

1439–1448. doi: 10.1007/s10592-018-1112-5

Garner, B. A., Hoban, S., & Luikart, G. (2020). IUCN Red List and the value of integrating

genetics. *Conservation Genetics*, *21*, 795–801. doi: 10.1007/s10592-020-01301-6

Garot, E., Joët, T., Combes, M. C., Severac, D., & Lashermes, P. (2019). Plant population

dynamics on oceanic islands during the Late Quaternary climate changes: genetic

evidence from a tree species (*Coffea mauritiana*) in Reunion Island. *New Phytologist*,

224, 974–986. doi: 10.1111/nph.16052

Gibson, J. P., Rice, S. A., & Stucke, C. M. (2008). Comparison of population genetic

diversity between a rare, narrowly distributed species and a common, widespread species

of *Alnus* (Betulaceae). *American Journal of Botany*, *95*, 588–596. doi:

10.3732/ajb.2007316

Gonzalez, E. G., Cerón-Souza, I., Mateo, J. A., & Zardoya, R. (2014). Island survivors:

Population genetic structure and demography of the critically endangered giant lizard of

La Gomera, *Gallotia bravoana. BMC Genetics*, *15*. doi: 10.1186/s12863-014-0121-8

Greenleaf, S. S., Williams, N. M., Winfree, R., & Kremen, C. (2007). Bee foraging ranges

and their relationship to body size. *Oecologia*, *153*, 589–596. doi: 10.1007/s00442-007-

0752-9

Guisan, A., & Theurillat, J.-P. (2000). Assessing alpine plant vulnerability to climate change:

a modeling perspective. *Integrated Assessment*, *1*, 307–320. doi:

10.1023/A:1018912114948

Hamrick, J. L., & Godt, M. J. W. (1996). Effects of life history traits on genetic diversity in

plant species. *Philosophical Transactions of the Royal Society in London Series B*, *351*,

1291–1298. doi: 10.1098/rstb.1996.0112

Ho, T. N. (1988). *Flora reipublicae popularis sinicae*. Beijing: Science Press.

Ho, T. N., & Liu, S. W. (2001). *A worldwide monograph of Gentiana*. Beijing: Science Press.

Hodel, R. G. J., Chen, S., Payton, A. C., McDaniel, S. F., Soltis, P., & Soltis, D. E. (2017).

Adding loci improves phylogeographic resolution in red mangroves despite increased

missing data: Comparing microsatellites and RAD-Seq and investigating loci filtering.

Scientific Reports, *7*, 1–14. doi: 10.1038/s41598-017-16810-7

Ikeda, H., Eidesen, P. B., Yakubov, V., Barkalov, V., Brochmann, C., & Setoguchi, H.

(2017). Late Pleistocene origin of the entire circumarctic range of the arctic-alpine plant

Kalmia procumbens. Molecular Ecology, *26*, 5773–5783. doi: 10.1111/mec.14325

Ikeda, H., Higashi, H., Yakubov, V., Barkalov, V., & Setoguchi, H. (2014).

Phylogeographical study of the alpine plant *Cassiope lycopodioides* (Ericaceae) suggests

a range connection between the Japanese archipelago and Beringia during the

Pleistocene. *Biological Journal of the Linnean Society*, *113*, 497–509. doi:

10.1111/bij.12342

Ikeda, H., Yakubov, V., Barkalov, V., Sato, K., & Fujii, N. (2020). East Asian origin of the

widespread alpine snow-bed herb, *Primula cuneifolia* (Primulaceae), in the northern

Pacific region. *Journal of Biogeography*, *47*, 2181–2193. doi: 10.1111/jbi.13918

Ikeda, H., Yakubov, V., Barkalov, V., & Setoguchi, H. (2018). Post-glacial East Asian origin

of the alpine shrub *Phyllodoce aleutica* (Ericaceae) in Beringia. *Journal of*

Biogeography, *45*, 1261–1274. doi: 10.1111/jbi.13230

Iwatsuki, K., Yamazaki, T., Boufford, D. E., & Ohba, H. (1993). *Flora of Japan vol. IIIa*.

Tokyo: Kodansha Ltd.

Japanese Ministry of the Environment. (2014). *Japanese ministry of the environment red data*

book. 272. Retrieved from https://ikilog.biodic.go.jp/rdbdata/files/envpdf/植物I_270.pdf

Japanese Ministry of the Environment. (2017). *Consideration of adaptation strategies to*

climate change in the World Natural Heritage area of Yakushima Island. Retrieved from

https://www.env.go.jp/park/yakushima/ywhcc/wh/kagaku/17/180217-2-4-2.pdf

Japanese Ministry of the Environment. (2020). *Japanese ministry of the environment red list*.

Retrieved from http://www.env.go.jp/press/files/jp/113667.pdf

Kogura, Y., Seeb, J. E., Azuma, N., Kudo, H., Abe, S., & Kaeriyama, M. (2011). The genetic

population structure of lacustrine sockeye salmon, *Oncorhynchus nerka*, in Japan as the

endangered species. *Environmental Biology of Fishes*, *92*, 539–550. doi:

10.1007/s10641-011-9876-1

Kort, H. De, Prunier, J. G., Ducatez, S., Honnay, O., Baguette, M., Stevens, V. M., &

Blanchet, S. (2021). Life history, climate and biogeography interactively affect

worldwide genetic diversity of plant and animal populations. *Nature Communications*,

12, 516. doi: 10.1038/s41467-021-20958-2

Kreiner, J. M., Giacomini, D. A., Bemm, F., Waithaka, B., Regalado, J., Lanz, C., … Wright,

S. I. (2019). Multiple modes of convergent adaptation in the spread of glyphosate-

resistant *Amaranthus tuberculatus*. *Proceedings of the National Academy of Sciences of*

the United States of America, *116*, 21076–21084. doi: 10.1073/pnas.1900870116

Li, S., Li, B., Cheng, C., Xiong, Z., Liu, Q., Lai, J., … Yan, J. (2014). Genomic signatures of

near-extinction and rebirth of the crested ibis and other endangered bird species. *Genome*

Biology, *15*, 557. doi: 10.1186/s13059-014-0557-1

Liu, X., & Fu, Y. X. (2015). Exploring population size changes using SNP frequency spectra.

Nature Genetics, *47*, 555–559. doi: 10.1038/ng.3254

Loïc Pellissier, Eidesen, P. B., Ehrich, D., Descombes, P., Schönswetter, P., Tribsch, A., …

Alsos, I. G. (2016). Past climate-driven range shifts and population genetic diversity in

arctic plants. *Journal of Biogeography*, *43*, 461–470. doi: 10.1111/jbi.12657

Losdat, S., Chang, S. M., & Reid, J. M. (2014). Inbreeding depression in male gametic

performance. *Journal of Evolutionary Biology*, *27*, 992–1011. doi: 10.1111/jeb.12403

Lynch, M. (2010). Evolution of the mutation rate. *Trends in Genetics*, *26*, 345–352. doi:

10.1016/j.tig.2010.05.003

Ma, Y., Chen, G., Edward Grumbine, R., Dao, Z., Sun, W., & Guo, H. (2013). Conserving

plant species with extremely small populations (PSESP) in China. *Biodiversity and*

Conservation, *22*, 803–809. doi: 10.1007/s10531-013-0434-3

Maruyama, T., & Fuerst, P. A. (1985). Population bottlenecks and nonequilibrium models in

population genetics. II. Number of alleles in a small population that was formed by a

recent bottleneck. *Genetics*, *111*, 675–689. doi: 10.1093/genetics/111.3.675

Miller, G. R., Geddes, C., & Mardon, D. K. (1998). Response of the alpine gentian *Gentiana*

nivalis L. to protection from grazing by sheep. *Biological Conservation*, *87*, 311–318.

doi: 10.1016/S0006-3207(98)00076-7

Mishiba, K. I., Yamane, K., Nakatsuka, T., Nakano, Y., Yamamura, S., Abe, J., … Nishihara, M. (2009). Genetic relationships in the genus *Gentiana* based on chloroplast DNA sequence data and nuclear DNA content. *Breeding Science*, *59*, 119–127. doi: 10.1270/jsbbs.59.119

Moschetti, M., Besnard, A., Couturier, T., & Fonderflick, J. (2020). Grazing intensity negatively affects the maintenance of *Gentiana pneumonanthe* and the survival of Phengaris alcon egg-laying. *Journal of Insect Conservation*, *24*, 343–351. doi: 10.1007/s10841-020-00220-8

Noroozi, J., Talebi, A., Doostmohammadi, M., Rumpf, S. B., Linder, H. P., & Schneeweiss, G. M. (2018). Hotspots within a global biodiversity hotspot-areas of endemism are associated with high mountain ranges. *Scientific Reports*, *8*, 1–10. doi: 10.1038/s41598-018-28504-9

Nybom, H., & Bartish, I. V. (2000). Effects of life history traits and sampling strategies on genetic diversity estimates obtained with RAPD markers in plants. *Perspectives in Plant Ecology, Evolution and Systematics*, *3*, 93–114. doi: 10.1078/1433-8319-00006

O'Grady, J. J., Brook, B. W., Reed, D. H., Ballou, J. D., Tonkyn, D. W., & Frankham, R. (2006). Realistic levels of inbreeding depression strongly affect extinction risk in wild populations. *Biological Conservation*, *133*, 42–51. doi: 10.1016/j.biocon.2006.05.016

Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., Mcglinn, D., … Maintainer, H. W. (2020). Package "vegan." Retrieved from https://cran.ism.ac.jp/web/packages/vegan/vegan.pdf

Oostermeijer, J. G. B., van Eijck, M. W., & den Nijs, J. C. M. (1994). Offspring fitness in relation to population size and genetic variation in the rare perennial plant species *Gentiana pneumonanthe* (Gentianaceae). *Oecologia*, *97*, 289–296. doi: 10.1007/BF00317317

Oostermeijer, J. G. B., Veer, R. V., & Nijs, J. C. M. Den. (1994). Population structure of the rare, long-lived perennial *Gentiana pneumonanthe* in relation to vegetation and management in the Netherlands. *The Journal of Applied Ecology*, *31*, 428. doi: 10.2307/2404440

Paradis, E., & Schliep, K. (2019). Ape 5.0: An environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics*, *35*, 526–528. doi: 10.1093/bioinformatics/bty633

Peakall, R., & Smouse, P. E. (2006). GENALEX 6: Genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes*, *6*, 288–295. doi: 10.1111/j.1471-8286.2005.01155.x

Pellicer, J., & Leitch, I. J. (2020). The Plant DNA C-values database (release 7.1): an updated

online repository of plant genome size data for comparative studies. *New Phytologist*,

226, 301–305. doi: 10.1111/nph.16261

Perrigo, A., Hoorn, C., & Antonelli, A. (2020). Why mountains matter for biodiversity.

Journal of Biogeography, *47*, 315–325. doi: 10.1111/jbi.13731

Piry, S., Luikart, G., & Cornuet, J. M. (1999). BOTTLENECK: A computer program for

detecting recent reductions in the effective population size using allele frequency data.

Journal of Heredity, *90*, 502–503. doi: 10.1093/jhered/90.4.502

Reed, D. H., & Frankham, R. (2003). Correlation between fitness and genetic diversity.

Conservation Biology, *17*, 230–237. doi: 10.1046/j.1523-1739.2003.01236.x

Reisch, C., & Rosbakh, S. (2020). Patterns of genetic variation in European plant species

depend on altitude. *Diversity and Distributions*, *27*, 157–163. doi: 10.1111/ddi.13177

Reisch, C., Schmidkonz, S., Meier, K., Schöpplein, Q., Meyer, C., Hums, C., … Schmid, C.

(2017). Genetic diversity of calcareous grassland plant species depends on historical

landscape configuration. *BMC Ecology*, *17*, 1–13. doi: 10.1186/s12898-017-0129-9

Ren, H., Zhang, Q., Lu, H., Liu, H., Guo, Q., Wang, J., … Bao, H. (2012). Wild plant species

with extremely small populations require conservation and reintroduction in China.

Ambio, *41*, 913–917. doi: 10.1007/s13280-012-0284-3

Reusch, T. B. H., Ehlers, A., Hammerli, A., & Worm, B. (2005). Ecosystem recovery after

climatic extremes enhanced by genotypic diversity. *Proceedings of the National

Academy of Sciences of the United States of America*, *102*, 2826–2831. doi:

10.1073/pnas.0500008102

Schönswetter, P., Stehlik, I., Holderegger, R., & Tribsch, A. (2005). Molecular evidence for

glacial refugia of mountain plants in the European Alps. *Molecular Ecology*, *14*, 3547–

3555. doi: 10.1111/j.1365-294X.2005.02683.x

Setzke, C., Wong, C., & Russello, M. A. (2022). Genome-wide assessment of kokanee

salmon stock diversity, population history and hatchery representation at the northern

range margin. *Conservation Genetics*, *23*, 243–253. doi: 10.1007/s10592-021-01418-2

Stojanova, B., Šurinová, M., Zeisek, V., Münzbergová, Z., & Pánková, H. (2020). Low

genetic differentiation despite high fragmentation in the endemic serpentinophyte

Minuartia smejkalii (M. verna agg., Caryophyllaceae) revealed by RADSeq SNP

markers. *Conservation Genetics*, *21*, 187–198. doi: 10.1007/s10592-019-01239-4

Suyama, Y., Hirota, S. K., Matsuo, A., Tsunamoto, Y., Mitsuyuki, C., Shimura, A., & Okano,

K. (2021). Complementary combination of multiplex high-throughput DNA sequencing

for molecular phylogeny. *Ecological Research*, 1–11. doi: 10.1111/1440-1703.12270

Suyama, Y., & Matsuki, Y. (2015). MIG-seq: An effective PCR-based method for genome-

wide single-nucleotide polymorphism genotyping using the next-generation sequencing

platform. *Scientific Reports*, *5*, 1–12. doi: 10.1038/srep16963

Takatsuki, S. (1990). Summer dietary compositions of sika deer on Yakushima Island,
southern Japan. *Ecological Research*, *5*, 253–260. doi: 10.1007/BF02346995

Takatsuki, S. (2009). Effects of sika deer on vegetation in Japan: A review. *Biological
Conservation*, *142*, 1922–1929. doi: 10.1016/j.biocon.2009.02.011

Thiel-Egenter, C., Gugerli, F., Alvarez, N., Brodbeck, S., Cieślak, E., Colli, L., …
Holderegger, R. (2009). Effects of species traits on the genetic diversity of high-
mountain plants: A multi-species study across the Alps and the Carpathians. *Global
Ecology and Biogeography*, *18*, 78–87. doi: 10.1111/j.1466-8238.2008.00421.x

Thuiller, W., Lavorel, S., Araújo, M. B., Sykes, M. T., & Prentice, I. C. (2005). Climate
change threats to plant diversity in Europe. *Proceedings of the National Academy of
Sciences of the United States of America*, *102*, 8245–8250. doi:
10.1073/pnas.0409902102

Tsujino, R., & Yumoto, T. (2004). Effects of sika deer on tree seedlings in a warm temperate
forest on Yakushima Island, Japan. *Ecological Research*, *19*, 291–300. doi:
10.1111/j.1440-1703.2004.00638.x

Tsukada, M. (1985). Map of vegetation during the last glacial maximum in Japan. *Quaternary
Research*, *23*, 369–381. doi: 10.1016/0033-5894(85)90041-9

Willi, Y., Van Buskirk, J., & Hoffmann, A. A. (2006). Limits to the adaptive potential of
small populations. *Annual Review of Ecology, Evolution, and Systematics*, *37*, 433–458.
doi: 10.1146/annurev.ecolsys.37.091305.110145

Woodgate, J. L., Makinson, J. C., Lim, K. S., Reynolds, A. M., & Chittka, L. (2017).
Continuous radar tracking illustrates the development of multi-destination routes of
bumblebees. *Scientific Reports*, *7*, 1–15. doi: 10.1038/s41598-017-17553-1

Common/ all loci	Species	No. of samples	r	No. of loci	No. of polymorphic loci	No. of Sites	No. of polymorphic sites	π Mean	SE	P	H_O Mean	SE	P	H_E Mean	SE	P	F_{IS} Mean	SE	P_0	P
Common	*G. yakushimensis*	353	0.5	89	47	7,237	72	0.00020	0.00008	***	0.013	0.005	***	0.020	0.008	***	0.222	0.045	***	
Common	*G. triflora*	9		89	47	7,237	72	0.00135	0.00030		0.121	0.030		0.125	0.023		0.048	0.076		
Common	*G. yakushimensis*	353	0.6	69	35	5,623	51	0.00009	0.00002	**	0.006	0.002	**	0.009	0.002	***	0.263	0.054	***	*
Common	*G. triflora*	9		69	35	5,623	51	0.00103	0.00028		0.100	0.030		0.105	0.025		0.058	0.075		
Common	*G. yakushimensis*	353	0.7	51	25	4,150	34	0.00005	0.00002	**	0.004	0.001	*	0.006	0.002	**	0.274	0.069	***	*
Common	*G. triflora*	9		51	25	4,150	34	0.00095	0.00031		0.113	0.041		0.108	0.031		-0.004	0.098		
All	*G. yakushimensis*	353	0.5	2,124	487	172,094	784	0.00055	0.00003	***	0.084	0.004	***	0.121	0.005	***	0.341	0.014	***	***
All	*G. triflora*	9		1,475	302	116,730	401	0.00162	0.00008		0.329	0.009		0.436	0.002		0.230	0.022	***	
All	*G. yakushimensis*	353	0.6	1,758	387	142,510	562	0.00041	0.00003	***	0.070	0.005	***	0.103	0.005	***	0.358	0.017	***	***
All	*G. triflora*	9		1,181	222	93,622	297	0.00147	0.00009		0.311	0.009		0.430	0.003		0.263	0.023	***	
All	*G. yakushimensis*	353	0.7	1,429	292	115,877	402	0.00033	0.00003	***	0.058	0.005	***	0.095	0.006	***	0.387	0.019	***	***
All	*G. triflora*	9		945	155	74,977	202	0.00125	0.00009		0.315	0.011		0.435	0.004		0.264	0.027	***	

623
624
625

626   **FIGURE 1**. Photograph of (a) *Gentiana yakushimensis* and (b-c) its habitat. The white arrows
627   in the panels (b) and (c) point to *Gentiana yakushimensis*.
628

629
630

FIGURE 2. Map of Yakushima Island and the distribution area of *Gentiana yakushimensis*.

The gray part in the right map denotes the area above 1,700 m above sea level. In the left map,

the gray part denotes the distribution area of *G. triflora*. The contour interval is 50 m.

FIGURE 3. Relationship between pairwise genetic and geographic distance. (a) Relationship
between individual-based pairwise genetic and geographic distance of *Gentiana yakushimensis*
when the parameter r is 0.5, 0.6, and 0.7. (b) Mantel correlograms for the association between
individual-based genetic and geographic distances. Filled symbols in the figure denote
statistical significance.

643
644

FIGURE 4. Principal coordinate analysis (PCoA) score plots based on individual-based genetic

distance of *Gentiana yakushimensis* when the parameter r is 0.5. (a), axis 1-2; (b), axis 1-3.

Colors denote the location (longitude) of each individual.

FIGURE 5. Demographic inference using stairway plot. The solid lines denote the median
estimate of the effective population size when the parameter r is 0.5, 0.6, and 0.7. The
translucence bands indicate 95% confidence intervals.

655
656

657 **FIGURE 6**. Heterozygosity-excess ($\Delta H = H_E - H_{Eq}$) based on INEST. The values of ΔH in all
658 SNPs with the setting of parameter r = 0.5, 0.6, and 0.7 are represented by dots. The
659 significances (P values) in heterozygote excess ($\Delta H > 0$) based on Wilcoxon signed-rank test
660 are denoted.

662
663

664 **FIGURE S1**. Principal coordinate analysis (PCoA) score plots based on individual-based
665 genetic distance of *Gentiana yakushimensis* when the parameter r is 0.6 (a, b) and 0.7 (c, d). (a,
666 c), axis 1-2; (b, d), axis 1-3. Colors denote the location (longitude) of each individual.
667

668
669