A deeply conserved protease, acylamino acid-releasing enzyme (AARE), acts in ageing in Physcomitrella and Arabidopsis

Sebastian N.W. Hoernstein[1], Buğra Özdemir[1,5], Nico van Gessel[1], Alessandra A. Miniera[1], Bruno Rogalla von Bieberstein[1,6], Lars Nilges[1], Joana Schweikert Farinha[1,7], Ramona Komoll[1,8], Stella Glauz[1], Tim Weckerle[1,9], Friedrich Scherzinger[1,10], Marta Rodriguez-Franco[2], Stefanie J. Müller-Schüssele[3], Ralf Reski[1,4,*]

[1] Plant Biotechnology, Faculty of Biology, University of Freiburg, Schaenzlestrasse 1, 79104 Freiburg, Germany

[2] Cell Biology, Faculty of Biology, University of Freiburg, Schaenzlestrasse 1, 79104 Freiburg, Germany

[3] Molecular Botany, Department of Biology, Technical University of Kaiserslautern, Erwin-Schrödinger-Strasse 70, 67663 Kaiserslautern, Germany

[4] Signalling Research Centres BIOSS and CIBSS, Schaenzlestrasse 18, 79104 Freiburg, Germany

[5] Euro-BioImaging Bio-Hub, EMBL Heidelberg, Meyerhofstraße 1, 69117 Heidelberg, Germany

[6] Department of Anesthesiology, University Hospital Würzburg, Oberduerrbacher Strasse 6, 97072 Würzburg, Germany

[7] Institute for Molecular Biosciences, Goethe University Frankfurt, Max-von-Laue-Str. 9, 60438 Frankfurt, Germany

[8] Heraeus Medical GmbH, Philipp-Reis-Straße 8-13. 61273 Wehrheim, Germany

[9] Eurofins GeneScan GmbH, Engesserstraße 4, 79108 Freiburg, Germany

[10] Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Puschstrasse 4, 04103 Leipzig, Germany

* Corresponding author: ralf.reski@biologie.uni-freiburg.de

ORCID-IDs S.N.W.H.: 0000-0002-2095-689X, B.Ö.: 0000-0001-9823-0581, N.v.G.: 0000-0002-0606-246X, A.A.M.: 0000-0003-1638-5438, B.R.v.B.: 0000-0002-9650-4064, L.N.: 0000-0003-

29 0716-7236, J.S.F.: 0000-0002-5602-3901, R.K.: 0000-0002-3142-0863, S.G.: 0000-0002-2305-
30 4525, T.W.: 0000-0002-9584-5449, F.S.: 0000-0001-5825-6621, M.R.F.: 0000-0003-1183-2075,
31 S.J.M.S.: 0000-0003-4061-1175, R.R.: 0000-0002-5496-6711

Abstract

Reactive oxygen species (ROS) are constant by-products of aerobic life. In excess, ROS lead to
cytotoxic protein aggregates, which are a hallmark of ageing in animals and linked to age-related
pathologies in humans. Acylamino acid-releasing enzymes (AARE) are bifunctional serine
proteases, acting on oxidized proteins. AARE are found in all domains of life, albeit under different
names, such as acylpeptide hydrolase (APEH/ACPH), acylaminoacyl peptidase (AAP), or
oxidized protein hydrolase (OPH). In humans, AARE malfunction is associated with age-related
pathologies, while their function in plants is less clear. Here, we provide a detailed analysis of
AARE genes in the plant lineage and an in-depth analysis of AARE localization and function in
the moss Physcomitrella and the angiosperm Arabidopsis. *AARE* loss-of-function mutants have
not been described for any organism so far. We generated and analysed such mutants and
describe a connection between AARE function, aggregation of oxidized proteins and plant ageing,
including accelerated developmental progression and reduced life span. Our findings complement
similar findings in animals and humans, and suggest a unified concept of ageing may exist in
different life forms.

Introduction

Reactive oxygen species (ROS) are by-products of O_2 metabolism and represent a challenge to all aerobic life. ROS play a dual role as potentially lethal oxidants and as signalling molecules[1]. Therefore, aerobic organisms possess sophisticated redox systems to scavenge and detoxify excess ROS. The major sources of ROS in plant cells are the electron transport chains in mitochondria and plastids, but also in peroxisomes and at the plasma membrane[2]. Environmental stresses such as heat, drought or intense light are factors for increasing ROS production to detrimental levels. Plants possess a repertoire of detoxifying enzymes such as catalases, superoxide dismutases, ascorbate peroxidases, glutathione peroxidase-like proteins and peroxiredoxins. Electrons for reduction are largely provided *via* non-enzymatic components such as ascorbic acid, glutathione and NADPH[3,4,5,6,7]. In addition, a range of heat-shock proteins assist in disaggregation or refolding of damaged proteins[8,9,10].

Despite conversion into non-toxic derivatives, the continuous exposure to ROS results in oxidation of DNA, lipids, and proteins[5]. On the protein level, ROS lead to irreversible cysteine oxidation, advanced glycation end-products, adducts with ROS-generated reactive aldehydes, and carbonylation of amino-acid side-chains[11,12]. If not cleared *via* proteolysis, an excess of oxidized proteins accumulates to cytotoxic protein aggregates. Plant antioxidant systems and the role of ROS as signalling molecules in abiotic stress responses are well studied[13]. Yet, factors involved in a plant cell's last line of defence, such as the proteolytic systems for the clearance of irreversibly oxidized proteins, are still underexplored.

A class of serine proteases is evolutionary deeply conserved as their activity is found in bacteria, archaea, animals and plants, and can degrade irreversibly oxidized proteins[14,15,16,17]. These proteases have different names in different organisms, e.g. acylamino acid-releasing enzyme (AARE), acylpeptide hydrolase (APEH/ACPH), acylaminoacyl peptidase (AAP), or oxidized protein hydrolase (OPH)[18] but are collectively addressed as AARE here. AARE acts in multimeric complexes[14,17] as a bifunctional protease as it cleaves N^α-acetylated amino acids from oligopeptides *via* an exopeptidase mode, but also cleaves oxidized proteins *via* an endopeptidase mode[16,19,20,21,22]. AARE isoforms from various organisms show different specificities towards N^α-acetylated amino-acids, with bacterial and archaeal enzymes preferring AcLeu and AcPhe substrates[15,23,24,25] and plant and animal isoforms preferring AcAla, AcMet or AcGly substrates[16,17,26,27]. Besides substrate specificities, their subcellular localization appears to be conserved among eukaryotes, as human (HsACPH) and Arabidopsis (AtAARE) AAREs are reported as cytosolic enzymes[20,21].

In humans, AARE malfunction is linked to different types of cancer[26,28,29] and sarcoma cell

viability[30]. Moreover, AARE and proteasomal activity correlate and cooperatively prevent cytotoxic

aggregate formation[31,32,33]. Several selective inhibitors have been identified[34,35] and blocking of

AARE function is considered as anti-cancer treatment[28]. Despite an increasing number of studies

on AARE functionality in humans, AAREs in plants are far less characterized.

AARE from *Arabidopsis thaliana* (AtAARE) and from cucumber have endo- and exopeptidase

functions[17], and silencing of *AtAARE* increased the levels of oxidized proteins[21]. AtAARE activity

was also detected in plastid stroma fractions, although a fusion with a fluorescent reporter did not

co-localize with chloroplasts[21]. Suppression of *AtAARE via* RNAi resulted in an enhanced

accumulation of oxidized proteins in roots and enhanced electrolyte leakage in leaves, but a

further impact on plant physiology was not described[21].

Moreover, the complete loss of function of this protease has not yet been reported for any

organism, neither bacterium, archaeon, animal, or plant. Thus, although the deep evolutionary

conservation of AARE suggests its pivotal role in all major life forms, experimental evidence is far

from optimal.

In a proteomics study on protein arginylation we identified an AARE homolog from the moss

Physcomitrella[36]. Our further analysis revealed altogether three Physcomitrella AARE homologs

(PpAARE1-3). Here, we analysed the subcellular localization of these PpAARE isoforms and of

their homolog from the angiosperm Arabidopsis (AtAARE). We show that an alternative splicing

event is targeting PpAARE1 to chloroplasts, mitochondria and the cytosol. We provide evidence

that an alternative translation initiation is sufficient to localize AtAARE to the same three

subcellular compartments. Bioinformatic analyses of several genomes suggest that the

localization of AARE in chloroplasts and mitochondria is conserved across the plant lineage.

Employing combinatorial gene knockouts and protein co-immunoprecipitation we found distinct

interactions between these three isoforms and their concerted action on progressive ageing in

Physcomitrella. Likewise, an Arabidopsis *AARE* loss-of-function mutant exhibits enhanced levels

of oxidized proteins and accelerated bolting, as a hallmark of plant ageing.

Results

AARE gene family expansion and splice variants

Previously, PpAARE1 (Pp1s619_3V6.1) was identified as the most prominent target for N-terminal arginylation in Physcomitrella[36,37]. N-terminal arginylation mediates poly-ubiquitination *via* the N-degron pathway, thus triggering subsequent proteasomal degradation[38]. Simultaneously, two homologs (PpAARE2: Pp1s108_134V6.1 and PpAARE3: Pp1s97_68V6.1) were identified, although those were not proven arginylation targets[36]. Meanwhile, a new Physcomitrella genome version with chromosome assembly and updated gene versions was released[39]. Consequently, the gene accessions used here are *PpAARE1* (Pp3c2_30720V3.1), *PpAARE2* (Pp3c12_21080V3.1), and *PpAARE3* (Pp3c7_25140V3.1). According to *OrthoMCL*[40,41], all three proteins are homologs of the *Arabidopsis thaliana* acylamino acid-releasing enzyme (AtAARE: AT4G14570). According to publicly available data (https://peatmoss.online.uni-marburg.de)[42] *PpAARE1-3* are expressed in all major Physcomitrella tissues and developmental stages, although at varying levels (Fig. S1a). Except for leaves (phylloids) and spores, *PpAARE1* is the most strongly expressed gene of this family (between 4 and 20 times). In contrast, *PpAARE2* is expressed considerably stronger than *PpAARE1* and *PpAARE3* in spores (Fig. S1a). Likewise, *AtAARE* is expressed in all major Arabidopsis tissues (Fig. S1b, data utilized from Mergner et al. (2020)[43] and downloaded from http://athena.proteomics.wzw.tum.de/). These data indicate a strong, positive correlation between transcript level and protein abundance across all tissues (Fig. S1b). Stress conditions decrease *AARE* expression in Arabidopsis shoots and in Physcomitrella protonemata (Fig. S1c-e).

To investigate whether other plants also possess multiple AARE homologs and to infer their phylogenetic relation, we conducted *BLASTP*[44] searches against protein models from selected species using the protein sequence of AtAARE as a query. We selected the alga *Chlamydomonas reinhardtii*[45], the liverwort *Marchantia polymorpha*[46], the peat moss *Sphagnum fallax* (*Sphagnum fallax* v1.1, DOE-JGI, http://phytozome.jgi.doe.gov/)*, the lycophyte *Selaginella moellendorffii*[47], the monocot *Oryza sativa*[48] and the dicot *Populus trichocarpa*[49], all available at the Phytozome12 database (https://phytozome.jgi.doe.gov). Additionally, we performed a NCBI *BLASTP* search against the charophyte *Klebsormidium nitens* proteome[50], and identified a single homolog (GAQ80280.1) in this species. We also included proteins of *Funaria hygrometrica*[51], a close relative to Physcomitrella from the Funariacea family[52], in our search. Finally, the AtAARE sequence was again searched against the Physcomitrella[39] and Arabidopsis[53] proteomes.

Homology of the resulting *BLAST* hits was confirmed if the reciprocal best *BLAST* hit against *A.*

thaliana was again AtAARE.

In *P. trichocarpa*, we identified a single homolog for which three distinct splice variants are

annotated (Potri.008G160400.1, Potri.008G160400.2, Potri.008G160400.3). These encode

different protein isoforms, but two variants seem to encode non-catalytic proteins. AARE enzymes

are prolyl-oligopeptidases with a conserved catalytic triad (Ser/Asp/His) in the C-terminal

peptidase domain[54,55]. In splice variant 2 (Fig. S2, Potri.008G160400.2) alternative splicing results

in the deletion of the catalytic Asp whereas the whole catalytic triad is lacking in splice variant 3

(Potri.008G160400.3). Hence, we consider these splice variants as non-active and disregard

them from further discussion.

In rice, we identified two homologs (LOC_Os10g28020.3, LOC_Os10g28030.1), with an

additional splice variant (LOC_Os10g28020.1) at one locus which encodes an N-terminal

extension.

In *C. reinhardtii, M. polymorpha, S. fallax* and *S. moellendorffii*, we identified a single ortholog

each. In *M. polymorpha*, three distinct splice variants are annotated (Mapoly0111s0036.1,

Mapoly0111s0036.2, Mapoly0111s0036.3). The latter two are UTR (untranslated region) splice

variants, thus resulting in the same protein sequence, whereas Mapoly0111s0036.1 encodes an

N-terminal extension of 97 aa compared to the other two variants. In *F. hygrometrica* we identified

three distinct isoforms.

Finally, our *BLASTP* searches using the latest Physcomitrella protein models[39] confirmed three

homologs of AtAARE (Pp3c2_30720V3.1, Pp3c12_21080V3.1, Pp3c7_25140V3.1). Additionally,

this search revealed another hit, Pp3c1_2590V3.1. This gene is composed of a single exon and

encodes a 131 aa protein which harbours the AARE N-terminal domain (PF19283). However, it

lacks a catalytic peptidase domain and is hardly expressed across different culture conditions and

tissues[56]. We also did not find any proteomics evidence across several Physcomitrella

analyses[36,57,58,59] for this protein. Therefore, we excluded this gene from further analysis.

We then used phylogenetic reconstruction to investigate the origin of gene duplications within the

gene family. As an outgroup, we included the well-characterized rat[16] and human[26] AARE and

two isoforms of the Antarctic icefish *Chionodraco hamatus*[27]. Physcomitrella and *F. hygrometrica*

share three distinct pairs of orthologs hinting at an expansion in the common ancestor of the two

species. Our phylogenetic analysis did not resolve AARE subfamilies across kingdoms (Fig. 1a)

and we conclude that the gene family expansions observed in rice and in the Funariaceae are
lineage-specific events.

In addition, this analysis reveals a closer relationship between *PpAARE1* and *PpAARE3*, which
presumably originate from a more recent gene duplication event, compared to *PpAARE2*. This is
supported by the fact that the open reading frames (ORFs) of *PpAARE1* and *PpAARE3* are
represented by a single exon whereas the ORF of *PpAARE2* is split across 17 exons, similar to
AtAARE (Fig. 1b). This is in congruence with a more recent emergence of intron-poor genes in
intron-rich families linked to stress response and developmental processes[60] and in line with
intron-less orphan Physcomitrella genes as earliest responders to abiotic stress[61].

For the three *PpAARE* genes, several splice variants are annotated, but only two splice variants
of *PpAARE1* give rise to distinct protein isoforms (Fig. 1b; PpAARE1_1, PpAARE1_2). Both splice
variants are present in Physcomitrella protonema (Fig. 1b, c), according to RT-PCR with splice
variant-specific primers (Supplementary Table S1). Likewise, for *AtAARE* two different ORF
definitions exist. With Araport11[62], a new version of the gene model was introduced exhibiting a
longer ORF at the 5' end (Fig. 1b). We detected the full-length transcript *via* reverse transcription
polymerase chain reaction (RT-PCR, Fig. 1c).

For *PpAARE1*, alternative splicing in the 5' end results in an N-terminal truncated variant whereas
the longer non-spliced variant encodes an N-terminal plastid transit peptide (cTP) according to
TargetP2.0[63]. A cleavage of the transit peptide at the predicted cleavage site (Ala^{72}-M^{73},
Supplementary Table S2) of PpAARE1 would release exactly the protein encoded by the short
splice variant. In contrast, PpAARE2 and PpAARE3 do not harbour any predicted N-terminal
targeting signals. Moreover, PpAARE3 is also lacking the WD40 domain that is present in
PpAARE1 and PpAARE2 (Fig. 1d).

The extension of the originally annotated ORF of *AtAARE* also encodes a plastid transit peptide
(Fig. 1d). To our knowledge, the longer variant of AtAARE has not yet been investigated, whereas
the short variant of AtAARE localizes to the nucleus and the cytosol[21]. In agreement with the latter
findings, we could predict a nuclear localization sequence (NLS, KKKK) with *LOCALIZER*[64]. Thus,
targeting of AtAARE to the cytosol and the nucleus, but also to plastids could be enabled by
alternative translation initiation. Likewise, PtAARE harbours a plastid transit peptide and a
potential alternative translation initiation site downstream of the predicted cTP cleavage site.

Accordingly, we checked for NLS in PpAARE isoforms and found one (KRRP, Supplementary
Table S2) in PpAARE1 and PpAARE3, whereas PpAARE2 has none, further supporting our

hypothesis that PpAARE1 and PpAARE3 originate from a relatively recent gene duplication event.
Accordingly, alternative splicing also generates two distinct transcripts for AARE1 in rice
(OsAARE1, Supplementary Table S2), where one variant encodes a potential plastid transit
peptide.

For all other plant species, no plastid targeting sequence was predicted, while the *C. reinhardtii*
AARE harbours a mitochondrial targeting sequence (Supplementary Table S2).

PpAARE1 and AtAARE in mitochondria, chloroplasts and cytoplasm

Organellar targeting of AARE has not yet been reported, although AARE activity was observed in
plastid-enriched fractions of cucumber[17]. However, chloroplasts, peroxisomes and mitochondria
are major hubs of ROS generation[65], and thus are likely organelles with elevated levels of oxidized
proteins. Thus, we investigated whether PpAARE1 and AtAARE would localize to chloroplasts *in*
vivo.

We generated fusion constructs of the PpAARE isoforms and of AtAARE with eGFP for transient
expression in Physcomitrella protoplasts. Due to the presence of a predicted plastid targeting
peptide for PpAARE1, eGFP was fused in frame to the 3' end of all coding sequences (CDS).
Since also peroxisomes are ROS-producing organelles, we used *PlantPredPTS1*[66,67] to check for
the presence of C-terminal positioned peroxisomal targeting signals. None of the selected AARE
isoforms were predicted to localize to peroxisomes (Supplementary Table S2). Although AtAARE
has a C-terminal CKL tripeptide, which is experimentally verified to mediate peroxisomal
targeting[66,68], the properties of its other C-terminal amino acids most likely prevent peroxisomal
targeting. A more recent prediction approach for PTS1-mediated targeting for Arabidopsis
proteins[69] further supports this conclusion. Peroxisomal targeting can also be mediated *via* N-
terminal nona-peptides[70,71], but these motifs are also not present within the first 100 aa in any of
the selected AARE sequences. In agreement with these predictions eGFP was fused to the 3'
end of the CDSs.

For PpAARE1 three different fusion constructs were assembled. Among these, a fusion of the
CDS of the short splice variant (Pp3c2_30720V3.1) and eGFP was cloned, as well as a fusion of
the CDS of the longer splice variant (Pp3c2_30720V3.2) and eGFP. Additionally, we cloned a
fusion of eGFP and the sequence part in which both variants differ (M^1-A^{72}, Pp3c2_30720V3.2).
This part harbours the plastid transit peptide predicted by *TargetP2.0*. All fusion constructs were
expressed under the control of the Physcomitrella Actin5 promoter[57,72] in a pMAV4 plasmid
backbone[73].

The PpAARE1 isoform derived from the short splicing variant (PpAARE1_1, Fig. 1b) clearly

localized to the cytoplasm (Fig. 2). The same localization was observed for PpAARE2 and

PpAARE3 (Fig. 2). Despite a predicted NLS, we did not observe a nuclear localization, either for

PpAARE1 or for PpAARE3.

The isoform encoded by the longer splice variant of *PpAARE1* (PpAARE1_2, Fig. 1b) localized to

chloroplasts and surprisingly also to mitochondria (Fig. 2). In contrast to the diffuse cytosolic

distribution of PpAARE1, specific foci were observed in chloroplasts. To investigate whether the

N-terminal sequence differing between the two PpAARE1 variants (M^1-A^{72}, Pp3c2_30720V3.2) is

sufficient to confer dual targeting, we fused this N-terminal sequence 5' to eGFP and observed

again dual localization (Fig. 2, PpAARE1_Nt). Full-length PpAARE1 was necessary to localize

eGFP to foci within chloroplasts, as the PpAARE1_Nt:eGFP fusion led to a uniform distribution.

However, full-length PpAARE1 was also homogeneously distributed throughout the cytoplasm.

This indicates the presence of interactors that recruit PpAARE1 to specific sites or complexes

within the chloroplasts. Further, we conclude that the N-terminal extension of PpAARE1_2

encodes an ambiguous targeting signal for import into chloroplasts and mitochondria as it is

capable of directing the fusion protein simultaneously to both organelles.

Simultaneous localization of proteins to chloroplasts and mitochondria can be mediated *via*

ambiguous targeting signals which are recognized by both translocation machineries. We

evaluated whether *ATP2*[75], a tool for the prediction of ambiguous targeting, would recognize

PpAARE1 but this was not predicted to be dually targeted. In contrast, AtAARE was predicted to

be dually targeted *via* an ambiguous signal. Thus, we cloned the analogous three fusion

constructs for AtAARE and investigated the subcellular localization of their encoded proteins

accordingly.

The AtAARE isoform translated from the shorter ORF (AtAARE, Fig. 1b) localized to the

cytoplasm (AtAARE_SV, Fig. 2), as observed for PpAARE1_1. This result is partially in agreement

with Nakai et al. (2012)[21] since we could not observe nuclear localization. Using the fusion

construct of the longer AtAARE variant, we observed clear dual targeting of the protein to

chloroplasts and mitochondria (AtAARE_LV, Fig. 2), as observed for PpAARE1_2. Here, the

eGFP signal was distributed homogeneously in the chloroplasts, in contrast to the foci of

PpAARE1:eGFP.

Next, we cloned only the N-terminal sequence differing between both variants (M^1-A^{55}, longer

ORF definition, Fig. 1b) and fused it to eGFP. In order to investigate whether the exact N-terminal

difference between the two AtAARE variants would be sufficient for targeting, the M^{56} (same as

M^1 in the shorter variant), which is the P1 aa at the predicted cleavage site, was deleted. Using

this construct, the eGFP signal localized to chloroplasts and mitochondria (AtAARE_Nt, Fig. 2).

The signal within chloroplasts was homogeneously distributed, similar to the longer AtAARE

variant. Thus, we conclude that the N-terminal extension of both long PpAARE and AtAARE

variants is sufficient for dual targeting of proteins *in vivo.* Intriguingly, the longer variant of AtAARE

localized exclusively to chloroplasts and mitochondria although alternative translation initiation

should be possible. This is interesting as alternative translation initiation is also possible in the

longer splice variant of *PpAARE1* (PpAARE1_2). In the latter also, the fusion protein localizes

exclusively to chloroplasts and mitochondria, which excludes the possibility of an alternative

translation initiation, at least in protoplasts. There are numerous transcripts in mammals where

translation of an upstream positioned ORF suppresses the translation of the downstream main

ORF[76]. A similar scenario is conceivable in Physcomitrella. However, it remains unclear how and

if translation from the internal start codons is controlled. It is also possible that factors controlling

alternative translation initiation of AtAARE are absent in Physcomitrella, at least in a

spatiotemporal manner, or they might only be triggered in specific physiological situations.

According to our data, the translation of the two variants of PpAARE1 is mainly controlled by

alternative splicing and not by alternative translation initiation.

In summary, PpAARE1 and AtAARE localize to three subcellular compartments *via* an ambiguous

targeting signal. In contrast, PpAARE2 and PpAARE3 localize solely to the cytoplasm.

Double knockout of *PpAARE1/2* reduces lifespan

Null mutants of *AARE* have not been described in any organism, and the biological role of this

protease apart from its catalytic function remained unclear. Hence, we created *AARE* knockouts

in Physcomitrella by deleting the full CDS of each gene *via* gene targeting (Fig. 3a) according

to[77]. To enable subsequent knockout (KO) of other AARE genes, different selection markers were

used. Since three distinct AARE genes exist in Physcomitrella which result in cytosolic proteases,

we generated all possible combinations of double KOs and triple KOs to avoid potential

compensation of the loss of function.

Plants surviving the selection procedure were screened *via* PCR for the absence of the respective

CDS and correct integration of the KO construct in the target locus (Fig. S4a). Additionally, the

number of genomic integrations of the KO construct was measured *via* quantitative PCR (qPCR)

as described[78] (Fig. S4b). At least three independent lines were identified for all single, double

and triple KOs (Fig. S4c-i) and a line with only a single integration in the genome was detected

for each KO (j-p, line numbers with stars). Further, haploidy of all lines was confirmed *via* flow

cytometry (Fig. S5a-c) as described[79]. These precautions were made as the transformation

procedure may generate plants with multiple integrations[80], possibly leading to off-target effects.

Further, the transformation procedure may lead to diploid plants with altered gene expression[81].

We used haploid lines with a single integration of the KO construct (Fig. S4) for subsequent

experiments.

Typically, AARE exopeptidase activity is assayed *via* N^α-acetylated amino acids like AcAla or

AcMet coupled to a reporter such as para-nitro-anilide (pNA) or 7-amido-4-methylcoumarin

(AMC). From these, AcAla-pNA was tested for several eukaryotic AAREs[22,26], including

AtAARE[17]. Here, we analysed the impact of AARE loss of function on the activity towards AcAla-

pNA and AcLeu-pNA. The latter is a substrate of bacterial and archaeal AARE isoforms but also

eukaryotic isoforms exhibit cleavage activity on this substrate[16,22,23,24,25]. On the single KO level,

the exopeptidase activity on both substrates was significantly reduced in the ΔPpAARE1 mutant

whereas the single KO of the other isoforms did not affect the activity (Fig. 3b). This strong impact

of PpAARE1 on the exopeptidase activity was consistent across all transgenic mutant lines (Fig.

S6).

One important step in Physcomitrella development is the transition from protonema to

gametophores, a developmental progression regulated among others by plant age and the

nutrient status[82]. The single KOs were phenotypically inconspicuous on the gametophore level

(Fig. 3c). In contrast, gametophores of Δ*PpAARE1/2* and Δ*PpAARE1/3/2* were severely stunted

and colonies were denser compared to wild type (WT) or the other mutants (Fig. 3c). This growth

effect is restricted to gametophores since protonema growth on solid medium did not differ

between WT and the KOs (Fig. 3d, e). Intriguingly, Δ*PpAARE1/2* and Δ*PpAARE1/3/2* mutants

showed accelerated developmental transition, as they developed gametophores from protonema

earlier (Fig. 3d). Since other KO lines did not show this effect, we attributed this to the double KO

of *PpAARE1/2* and performed a quantitative comparison with WT. Here, gametophores were

already observed after 6 days in the double KO of *PpAARE1/2*, while in WT a similar number of

gametophores per colony was observed only after 13 days (Fig. 3f). Consequently, the double

KO of *PpAARE1/2* causes accelerated developmental progression but gametophores remained

ultimately smaller than in WT (Fig. 3c). These effects are not linked to AARE exopeptidase activity

since the exopeptidase activity in gametophores was significantly reduced in all lines with a KO of PpAARE1 (Fig. 3g), which mimics the activity profile in protonema (Fig. S6).

To analyse AARE endopeptidase activity, we assessed the total levels of oxidized proteins in gametophores. In this assay, protein carbonyl groups derived from oxidation are derivatized with 2,4-dinitrophenylhydrazine (DNPH) to 2,4-dinitrophenylhydrazone (DNP). This irreversible modification is then recognized on Western blots by a primary anti-DNP antibody. Since DNPH can also react with oxidation states of cysteine side chains[83], this assay detects not only protein carbonylation but general protein oxidation.

With this assay we found that PpAARE2 had the strongest impact on the level of oxidized proteins in gametophores (Fig. 3h) and thus is not linked to exopeptidase activity. This was consistently observed in three independent analyses (Fig. 3h, Fig. S7). Apparently, PpAARE3 does not have any impact on exopeptidase and endopeptidase activity in gametophores under standard conditions (Fig. 3g, h).

Taken together, PpAARE1 predominantly acts as exopeptidase, while PpAARE2 predominantly acts as endopeptidase, and only the simultaneous loss of both activities in the double knockout mutants has the severest phenotypical consequences.

We found another remarkable difference between WT and mutants with a double KO of *PpAARE1/2* in older plants. After 5 months of cultivation, Δ*PpAARE1/2* and Δ*PpAARE1/3/2* were only viable at the tip of the gametophores (Fig. 4a), whilst most of the colony was already dead. In contrast, gametophores of WT and the other KOs were fully viable. After 8 months, Δ*PpAARE1/2* and Δ*PpAARE1/3/2* were already dead, in contrast to WT and the respective parental lines, which only showed some dead gametophores (Fig. 4b, c).

In summary, mutants with a double KO of *PpAARE1/2* exhibit accelerated developmental transition from protonema to gametophore (Fig. 3f), while size and life span of gametophores is strikingly reduced (Fig. 3c, Fig. 4a-c). In contrast, these effects are not visible in Δ*PpAARE1/3*. Therefore, these ageing phenotypes are linked to the concurrent loss of major AARE endopeptidase and exopeptidase activity.

Distinct *in vivo* interactions of PpAARE isoforms

In different organisms, AARE forms different homomeric complexes such as dimers[14], tetramers[17], or hexamers[24]. Thus, we analysed whether the PpAARE isoforms can interact with each other. Previously, all three isoforms were identified, although the protein modification used

for pulldown (N-terminal arginylation) was only identified on PpAARE1[36]. This gave rise to two
hypotheses: First, PpAARE2 and PpAARE3 are also targets for N-terminal arginylation, but
modified peptides were not identified for these isoforms. Second, the isoforms interact in
complexes which were pulled down due to the N-terminal arginylation of PpAARE1. We generated
Citrine fusion lines for each isoform *via* in-frame tagging at the native locus (knock-in, Fig. 5a).
The original stop codons of the respective PpAARE CDS were deleted. The Citrine-tag was
intended for two different analyses: First, it should enable *in vivo* monitoring of PpAARE isoforms
expressed from the native promoter, and second, it is a valid tag for co-immunoprecipitation (Co-
IP) *via* commercially available trap-beads.

In plants with a detectable fusion transcript (Fig. S8a-c) the presence of the target protein was
checked *via* IP and subsequent MS analysis. For *PpAARE3:Citrine* lines, we detected transcripts
in one line and obtained only insufficient coverage and intensity at the MS across several Co-IPs.
Thus, these Physcomitrella lines were excluded from further analysis. The co-precipitation of
other PpAARE isoforms with the respective bait isoforms was observed in test IPs (Fig. S8d-e)
confirming previous MS-Data[36]. All plants harbouring the Citrine fusion were phenotypically
inconspicuous (Fig. S8f) and haploid (Fig. S8g). Although the fusion proteins were detected in
two independent lines for each of the two isoforms (PpAARE1, PpAARE2), we could not observe
any Citrine signal within Physcomitrella protonemata or gametophores, probably due to the low
abundance of the PpAARE isoforms. Nevertheless, MS intensities and sequence coverage
enabled quantitative Co-IPs. The MS data have been deposited in PRIDE with the accession
codes PXD033854 and PXD038742[84,85].

When targeting PpAARE1:Citrine, both other isoforms appeared as significant interacting
partners (Fig. 5b, $p < 0.01$, FDR = 0.01). In a reciprocal Co-IP targeting PpAARE2:Citrine only,
PpAARE1 appeared as significant interacting partner. PpAARE3 was not detected in this
pulldown. Although lacking a reciprocal IP targeting PpAARE3:Citrine, the data show that
PpAARE1 interacts with PpAARE2 and PpAARE3, whereas PpAARE2 only interacts with
PpAARE1. Consequently, there are distinct interactions of PpAARE1 with PpAARE2 and
PpAARE3 *in vivo*, possibly resulting in cytosolic heteromeric AARE complexes in Physcomitrella.

AARE affects bolting time in Arabidopsis

In Physcomitrella three *AARE* genes exist, and the concerted action and interaction of the
enzymes affect plant ageing. To evaluate if this is an evolutionary conserved function, we
analysed the situation in Arabidopsis. Here, it was known that silencing of the single *AARE* gene

leads to an accumulation of oxidized proteins, whereas overexpression did not affect their levels[21].

To gain deeper insights, we screened for available Arabidopsis T-DNA mutants.

We identified two T-DNA insertion lines (SALK_071125 and GK-456A10) in the T-DNA Express

database at the SIGnAL website (http://signal.salk.edu). SALK_071125 (s68) has a T-DNA

insertion in an intron at the 5´UTR, whereas the T-DNA insertion in GK-456A10 (GK) maps to an

intron in the region encoding the catalytic domain (Fig. 6a). We obtained these mutants, identified

homozygous populations of GK-456A410 containing the T-DNA insertion by their resistance to

sulfadiazine, and confirmed their genotype by PCR. In the case of the Salk line (s68), homozygous

plants had lost their resistance to kanamycin, but we confirmed their genotype by PCR.

Homozygous plants of both genotypes were back-crossed with WT Col-0, and brought to

homozygosis for subsequent experiments. Primers for screening and validation of both T-DNA

lines are listed in Supplementary Table S3. Additionally, we analysed *AARE* gene expression *via*

RT-PCR. The transcript was still detectable in both lines (Fig. 6a), although very reduced in s68,

while the protein was not detectable *via* Western blot (Fig. 6b). Surprisingly, in WT AARE was

detected at around 100 kDa, although the estimated molecular weight is approximately 84 kDa

(90 kDa for the longer ORF variant without cleavage of the plastid targeting peptide) which

indicates the presence of posttranslational modifications. Phenotypically, neither seedlings nor

adult mutant plants showed obvious deviations from WT (Col, Fig. S9)

Next, we assayed AARE exopeptidase function in Arabidopsis WT and the two mutants. The

exopeptidase activity on AcAla-pNA was significantly reduced in both T-DNA mutants. In contrast,

the activity on AcLeu-pNA did not change significantly, although a slight reduction was observed

in the GK line (Fig. 6c). Thus, in agreement with Western blot and activity data, the remaining

transcripts detected *via* RT-PCR (Fig. 6a) are not translated to functional enzymes, suggesting

complete loss of AARE function in the mutants. Based on this characterization, we concentrated

on T-DNA mutant s68.

Subsequently, we assayed the AARE endopeptidase activity in Arabidopsis. The most striking

feature reported were increased levels of oxidized proteins in AARE-silenced Arabidopsis plants[21]

which is in line with our findings in Physcomitrella AARE mutants. To corroborate this, we

investigated levels of oxidized proteins in the Arabidopsis T-DNA mutant (s68) in comparison to

WT (Col) cultivated under short day conditions. In both genotypes, the distribution of oxidized

proteins was different to Physcomitrella gametophores (Fig. 6d). Using the same extraction

protocol, oxidized proteins in Arabidopsis were mainly of lower molecular weight, whereas in

Physcomitrella they were mainly at higher molecular weight (Fig. 3h). Despite these size differences, the level of oxidized proteins was higher in the Arabidopsis mutant than in WT (Fig. 6d), mimicking the situation in Physcomitrella. Together, we found AtAARE exopeptidase and endopeptidase activity to be reduced in our experiments.

Oxidized proteins accumulate during vegetative growth of Arabidopsis and are depleted at the transition to bolting[86]. It was not clear from this study whether the level of oxidized proteins is a signal for bolting, or if the reset of protein oxidation at the bolting transition is a side-effect of enhanced protein turnover during this process. To address this, we assessed the bolting behaviour of Arabidopsis WT and mutant plants and found that bolting in mutant plants differed significantly from WT. In WT, bolting started at 24±1 rosette leaves whereas it started at 21±2 rosette leaves in the mutants (Fig. 6e). Consequently, this accelerated developmental transition in the mutants correlates with enhanced levels of oxidized proteins.

Discussion

A universal definition of ageing is difficult due to strongly differing characteristics between and within the domains of life. In humans, the term ageing is inherently linked to degenerative diseases whereas plant ageing is considered as the progression of developmental transitions from seeds to senescence. Nevertheless, a common feature of ageing is the execution of a genetic program that controls growth, development and maturation. In turn, the progression of this genetic program depends on the metabolic state and environmental factors. Among the molecules responsible for ageing are ROS. Their involvement in ageing and diseases was first postulated by Harman[87] and was extended by considering mitochondria as central sources[88]. In humans, mitochondrial malfunction in line with increased ROS levels is a central determinant of ageing and associated pathologies[89]. In plants, mitochondrial ROS increase during ageing of seeds[90] and are major determinants of germination capacity[91]. In photosynthetic tissues chloroplasts are the major source of ROS and their levels increase during ageing[92,93,94]. Plants cannot escape from stress situations that increase ROS production to detrimental levels, and despite several layers of ROS defence, oxidized proteins constitute the major share of modified molecules under stress. Consequently, protein oxidation and subsequent aggregate deposition are hallmarks of ageing[95,96]. The degradation of artificially oxidized proteins by AARE has been demonstrated repeatedly, and silencing of *AtAARE* in turn increased the levels of oxidized proteins[19,20,21,22].

However, the contribution of AARE to the progress of ageing remained elusive, although several

studies associate AARE-function to age-related diseases[97,98,99]. Here, we identified three *AARE*

genes in Physcomitrella. This gene family expansion is an outstanding feature of the moss family

Funariaceae, as most organisms we analysed contain only a single *AARE* gene, and only a few

have two. We analysed these three isoforms in Physcomitrella and compared selected features

to the single isoform in Arabidopsis. Our data reveal specific functions in age-related

developmental transitions and life span determination.

Surprisingly, we found triple localization of AtAARE and PpAARE1 to chloroplasts, mitochondria

and the cytosol *in vivo*, suggesting a functional role for AARE in these cellular compartments. In

Physcomitrella, the triple localization is mediated *via* alternative splice variants and in Arabidopsis

likely *via* alternative translation initiation. Although there have been indications of AtAARE being

associated to chloroplasts[17,21], there was no clear evidence for organellar targeting of this

protease until now. It is remarkable that the triple localization of AARE is evolutionary conserved

between Arabidopsis and Physcomitrella, although likely executed *via* different molecular

mechanisms. This suggests an essential and evolutionary conserved function of AARE activity in

the cytoplasm, chloroplasts and mitochondria. As mosses and seed plants diverged more than

500 million years ago[100], this is a deep evolutionary conservation.

Previously, AARE exopeptidase activity was observed in cucumber chloroplasts[17], and AARE

peptides were found in proteomes of Arabidopsis chloroplasts[101] and mitochondria[102]. In contrast,

AARE was not identified in plastid or mitochondrial proteomes of Physcomitrella[57,59]. We found

AARE exopeptidase activity in chloroplasts and mitochondria of Physcomitrella. Chloroplasts are

a major source of ROS in photosynthetic tissues exposed to light, whereas mitochondria are the

major source of ROS in non-photosynthetic tissues or in the dark[93,94,103,104]. Until now, it remained

unresolved how plants deplete oxidized proteins from these organelles.

In yeast and mammals, the ATP-dependent LON and AAA proteases are involved in clearance

of misfolded and oxidized mitochondrial proteins. Intriguingly, mutants of plant homologs of LON

proteases did not show clear effects on the levels of oxidized proteins, but AAA-type FTSH

proteases may play a role[105]. Nevertheless, stressors such as heat, drought or intense light

compromise photosynthesis[106,107] and mitochondrial respiration[108], leading to a depletion of ATP

and to mitochondrial dysfunction[109]. In turn, energy supply for ATP-dependent defence systems

such as heat-shock proteins and AAA-type proteases is severely compromised, leaving the

question unanswered how oxidized proteins in chloroplasts and mitochondria can be cleared.

Because AARE is an ATP-independent protease, our data suggest that organellar-targeted AARE

may act as an ATP-independent defence to prevent or attenuate protein aggregation in the major
ROS-producing organelles.

Based on localization prediction, 70% of our selected plant species possess one AARE isoform
that localizes to either plastids or to mitochondria. Whether dual targeting *via* ambiguous targeting
also occurs in these species remains to be experimentally validated. Further, we do not exclude
that the remaining species also have organellar AARE isoforms, because our predictions may be
compromised by incomplete gene models. Such incomplete AARE gene models without a transit
peptide were present in earlier genome annotations of Arabidopsis and Physcomitrella[41,53], whilst
complete gene models with transit peptides were only introduced with later versions[39,62].

It is not yet clear how translation of both variants in Arabidopsis is regulated, but a recent study
highlights the importance of alternative translation initiation in shaping different subcellular
localizations or functions of proteoforms[110]. This mechanism is also present in Physcomitrella,
where dual targeting of FtsZ isoforms to the cytoplasm and chloroplasts is enabled *via* alternative
translation initiation[111]. Thus, localization of PpAARE1 to the cytoplasm is also possible from the
longer splice variant. Alternative translation initiation of *AtAARE* is further evidenced by
proteomics data (www.peptideatlas.org/builds/arabidopsis/)[112]. Here, we found evidence for N-
terminal acetylation of M^{56} which is the initiator methionine of the short variant (PXD012708[113]).

A reporter fusion of the shorter *AtAARE* ORF was observed in the cytoplasm and in the nucleus[21].
We did not detect a nuclear localization of any AARE isoform. Using *LOCALIZER*[64] we identified
an NLS in AtAARE, in PpAARE1 and PpAARE3, but not in PpAARE2. In contrast, the human
AARE homolog HsACPH does not have a predictable NLS, but nuclear import is mediated *via*
interaction with XRCC1 under stress, where it acts in DNA-damage repair[114]. Similarly, a nuclear
localization of AARE might also occur under stress *in planta*.

In Physcomitrella, AARE1 is the dominant exopeptidase, whereas AARE2 acts predominantly as
endopeptidase, and the operation mode of AARE3 remains unresolved. Crystal structures from
bacterial and archaeal AAREs revealed two possible entrances for substrates to the catalytic
centres[14,115] but the mode of substrate entry is not fully understood. Although the quaternary
arrangements of subunits differ between species[115,116], the secondary structure arrangement is
conserved across kingdoms and specific subunit interactions (multimerization) are likely a
mechanism to mediate substrate specificity and modulate activity. This could be further used as
a switch between endopeptidase and exopeptidase activity and additionally enable access of the

catalytic centre *via* structural re-arrangements. Accordingly, the interaction between the distinct PpAARE isoforms may modify substrate specificity and activity.

Our data indicate a partial compensation between PpAARE1 and PpAARE2. Both single KO mutants are phenotypically inconspicuous under normal conditions, although a significant reduction in exopeptidase activity was observed in ∆*PpAARE1*. PpAARE2 had the strongest effect on the accumulation of oxidized proteins in gametophores on the single isoform KO level, whereas the double KO of *PpAARE1* and *PpAARE2* showed the most striking effect in all lines as accelerated transition from protonema to gametophores. This phenotype is similar to mutants with disturbed auxin transport[117]. However, colony growth in these mutants was reduced which is different from our double KO mutant (*PpAARE1/2*). Intriguingly, this double KO results in stunted gametophores and a reduced life span. This phenotype partially resembles a loss-of-function mutant of a central component of autophagy (PpATG3) in Physcomitrella[118]. In ∆*PpATG3*, gametophores exhibit a reduced life span and colonies are smaller than in WT. In contrast, gametophore size was not affected. Further, photosynthetic capacity in ∆*PpATG3* was also reduced, an effect which is apparently not caused by AARE depletion[21] and *PpAARE* genes were not differentially expressed in ∆*PpATG3*[118]. We conclude that the reduced life span observed in ∆*PpAARE1/2* and ∆*PpAARE1/3/2* is not due to an impaired autophagy system causing nitrogen starvation. This is in line with data which opposes autophagy at the onset of senescence in Arabidopsis[119].

In mammals, elevated levels of oxidized proteins are associated with age-related pathologies, such as Alzheimer´s disease, diabetes and different types of carcinomas[120]. If proteolytic clearance fails, further accumulation of oxidized proteins causes protein aggregation, which is a hallmark of ageing in animals[96,121,122]. A connection between protein oxidation and ageing was less well studied in plants. Plastid ROS levels increase during ageing[92], which is in line with strong oxidation of plastid proteins in ageing leaves[86]. Likewise, protein oxidation marks the developmental transition between vegetative growth and flowering in Arabidopsis[86]. Physcomitrella mutants ∆*PpAARE1/2* and ∆*PpAARE1/3/2* showed accelerated developmental transition from protonema to gametophores, reduced life span and increased levels of oxidized proteins as signs of accelerated ageing. This is supported by the fact that gametophore tips, which is younger tissue, are viable longer than the older stems in both mutants. In the Arabidopsis *AARE* T-DNA mutants we found increased levels of oxidized proteins under normal cultivation conditions and an accelerated developmental transition, in this case premature bolting. These findings suggest an evolutionary conserved connection between protein oxidation and ageing.

We provide here a detailed analysis of *AARE* genes in the plant lineage and an in-depth analysis

of AARE localization and function in the moss Physcomitrella and the annual angiosperm

Arabidopsis. *AARE* loss-of-function mutants have not been described for any organism so far.

We generated and analysed such mutants and describe a connection between AARE function,

aggregation of oxidized proteins and plant ageing, including accelerated developmental

progression and reduced life span. Our findings complement similar findings in humans and

animals where AARE malfunction is associated with protein aggregation and age-related

pathologies.

To solidify the role of AARE in ageing in different life forms, particularly in plants with contrasting

maximum life spans and in animals of different complexity, loss-of-function mutants should be

established and analysed in selected model species. Likewise, a deeper understanding of AARE-

function in human diseases is desirable. Together, such analyses may contribute to a unified

concept of ageing in different life forms.

Methods

Cultivation of Physcomitrella

Physcomitrella WT (new species name: *Physcomitrium patens* (Hedw.) Mitt.[123]); ecotype
"Gransden 2004" and AARE KO lines were cultivated in Knop medium[124] supplemented with
microelements. Knop medium (pH 5.8) containing 250 mg/L KH_2PO_4, 250 mg/L KCl, 250 mg/L
$MgSO_4$ x 7 H_2O, 1,000 mg/L $Ca(NO_3)_2$ x 4 H_2O and 12.5 mg/L $FeSO_4$ x 7 H_2O supplemented with
10 mL per litre of a microelement (ME) stock solution[125,126] (309 mg/L H_3BO_3, 845 mg/L
$MnSO_4$ x 1 H_2O, 431 mg/L $ZnSO_4$ x 7 H_2O, 41.5 mg/L KI, 12.1 mg/L Na_2MoO_4 x 2 H_2O, 1.25 mg/L
$CoSO_4$ x 5 H_2O, 1.46 mg/L $Co(NO_3)_2$ x 6 H_2O). For cultivation on solid medium, 12 g/L Agar was
added to the KnopME medium. Moss suspension cultures were disrupted weekly with an ULTRA-
TURRAX (IKA) at 18,000 rpm for 90 s. If not indicated otherwise, moss was grown under standard
light conditions (55 μmol photons/m^{2}s) at 22°C in a 16h/8h light/dark cycle.

Hydroponic Physcomitrella gametophore cultures were assembled as described[36,127]. Here, a thin
layer of protonema from suspension was distributed on gauze mesh (PP, 250 m mesh, 215 m
thread, Zitt Thoma GmbH, Freiburg, Germany) capped on a glass ring. The glass rings with
protonema-covered mesh gauze were placed in Magenta®Vessels (Sigma-Aldrich, St. Louis,
USA) and KnopME medium was added until the protonema-covered gauze mesh was moist. The
medium was exchanged every 4 weeks. Gametophores were harvested after 12 weeks.

Gametophore colonies on Agar plates (KnopME) were generated by transplanting single
gametophores to new plates. Plates were sealed with Parafilm®.

Generation of AARE knockout lines

Knockout constructs for each *PpAARE* gene were generated by amplifying genomic regions as
homologous flanks. The PCR products were fused to a neomycin (*PpAARE1*), hygromycin
(*PpAARE2*) and zeocin (*PpAARE3*) resistance cassettes, respectively, employing triple template
PCR as described[128] with primer sequences listed in Supplementary Table S1 using the *Q5*
polymerase (New England Biolabs, Ipswich, USA). The knockout constructs were released from
their vector backbones with XhoI (*PpAARE1*), BglII (*PpAARE2*) and DraI (*PpAARE3*),
respectively. Enzymes were purchased from Thermo Fisher Scientific. Digested plasmids were
precipitated and sterilized prior to transfection using standard ethanol precipitation method[129].
Transfection of Physcomitrella WT or KO (for consecutive knockouts) protoplasts was conducted
via PEG-mediated procedure[77,130]. The WT strain as well as the identified *PpAARE* KO lines are

accessible via the International Moss Stock Center (IMSC, www.moss-stock-center.org). IMSC
accession numbers for the mutant and WT are available in Supplementary Table S4.

Screening of plants surviving antibiotics selection was done by PCR. KO mutant plants surviving
the antibiotics selection were checked for the presence of a genomic region which should be
removed upon homologous recombination-based integration of the knockout construct (Fig. S4a).
In case of the absence of this WT signal, plants were further checked for correct 5' and 3'
integration of the respective knockout construct using primers listed in Supplementary Table S1.

Protonema growth and gametophore induction

Suspension cultures were started at the same day and disrupted weekly as described. Dry weight
was measured in triplicates and suspension density was adjusted to 440 dry weight (DW) per litre
(mg DW/L) as described[131]. Droplets of 15 µL were distributed on solid medium (Knop ME, 0.02%
glucose). Sealed plates were cultivated as described above. Three droplets each per line were
distributed on one plate and all lines were grown on the same plate. 3 plates (12 colonies) were
used per assay. Colony areas were measured with *ImageJ*. White pixels counted from binarized
colony images were used as quantitative values. Gametophores were counted upon visibility of
the first leafy structure on buds.

Generation of PpAARE-Citrine knock-in lines

Knock-in constructs to fuse the coding sequence of Citrine to the C-terminus of PpAARE isoforms
via an 18 bp linker[128] at the endogenous genomic locus were cloned via Gibson assembly[132]. All
necessary parts were amplified using primers listed in Supplementary Table S1. Additionally, XhoI
(*PpAARE1*), SalI (*PpAARE2*) and BamHI (*PpAARE3*) restriction sites were added to the 5' and 3'
ends of the respective knock-in construct. All parts were assembled into pJet1.2 vector backbone
(Thermo Scientific) using the Gibson Assembly®Cloning Kit from New England Biolabs (Ipswich,
Massachusetts, USA). Transfection of Physcomitrella WT protoplasts were conducted as
described[77,130] by co-transfecting a plasmid containing a neomycin phosphotransferase
resistance (nptII) cassette as transient selection marker (pBSNNNEV, Mueller et al., 2014)[57]. The
linearized plasmid and the co-transfection vector were purified and sterilized *via* ethanol
precipitation[129] prior to transfection.

The presence of Citrine was checked with primers listed in Supplementary Table S1 and resulting
positive lines were further checked for correct 5' and 3' integration by PCR using the Plant Phire
Kit with primers listed in Supplementary Table S1. All identified fusion lines are available *via* the

International Moss Stock Center (IMSC, www.moss-stock-center.org). IMSC accessions are

listed in Supplementary Table S4.

qPCR analysis

The copy number of the integrated KO constructs was determined using a qPCR-based method[78].

Genomic DNA was extracted from 100 mg frozen protonema using the GeneJEt Plant Genomic

DNA Purification Kit (Thermo Scientific, Waltham, USA). DNA concentrations were adjusted to 3

ng/µl for each sample and qPCR was performed with primers for the 5' and 3' flanks as well as

with primers for the corresponding selection cassette. Additionally, primers for the single copy

gene *CLF* (Pp3c22_2940V3) were used as internal control. Reference lines were WT as well as

in-house lines with known single integrations of the used selection cassettes. Primers are listed

in Supplementary Table S5. PCR reactions were done using 2x SensiFAST Mix (Bioline, London,

UK) and analysed in a Lightcycler 480 II (Roche, Basel, Schweiz).

cDNA preparation

RNA was extracted using the innuPREP Plant RNA Kit (Analytik Jena, Jena, Germany). The

extracted RNA was treated with DNAse I (Thermo Scientific) and subsequently reverse

transcribed into cDNA using Superscript III Reverse Transcriptase (Life Technologies, Carlsbad,

USA).

Fusion constructs for subcellular localization

All constructs were generated using Gibson assembly[132] and integrated into a

PpAct5:Linker:eGFP-MAV4 vector backbone[73]. Integration of the different coding sequences was

done in frame in front of the Linker:eGFP. All parts for the Gibson assembly (inserts and

corresponding vector backbones) were amplified either with Phusion™ polymerase (Thermo

Fisher Scientific) or with HiFi polymerase (PCR Biosystems Ltd) according to the manufacturer´s

instructions. The primers were designed to have a 25 bp overlap to the corresponding fragment

to be fused with. All primers and combinations are listed in Supplementary Table S6. In the case

of the N-terminal difference of AtAARE (M^1-A^{55} of AT4G14570.1, gene model provided by TAIR

(https://www.arabidopsis.org/)) the Actin5 promoter was replaced by the CaMV35S promoter[133]

previously validated in Physcomitrella[134].

Cloned plasmids were purified using the PureYield™ Plasmid Midiprep kit (Promega, Wisconsin,

USA) according to the manufacturer´s instructions. The plasmids were purified and sterilized *via*

ethanol precipitation[129].

Confocal microscopy

Confocal imaging was performed on transiently transfected live protoplasts using Leica TCS SP8

(Leica Microsystems, Wetzlar, Germany). Immediately before microscopy, MitoTracker™ Orange

CMTMRos (ThermoFisher Scientific) was added to protoplast suspensions to a final concentration

of 100 nM. For all imaging experiments, an HC PL APO CS2 63x/1.40 OIL objective was used

with a zoom factor of 4. The pinhole was set to 35.4 µm. For excitation, a WLL laser (70%) was

used. In a sequential acquisition setup, eGFP and chlorophyll were excited with the same laser

beam (488 nm, 2%) and their signals were detected simultaneously, whereas MitoTracker™ was

excited with a different laser beam (550 nm, 2%) and its signal was detected separately. The

detection ranges were specified as 502 nm - 546 nm for eGFP, 662 nm - 732 nm for chlorophyll

and 597 nm - 636 nm for MitoTracker™. The images were acquired as z-stacks with the number

of optical sections varying with the protoplast size. The voxel sizes of the resulting z-stacks were

0.0903, 0.0903, 0.239 µm in the x-y-z order. For visual representation and analysis, single slices

with the best signal-to-noise ratio were selected and extracted from each z-stack using FIJI

software.

Cultivation of Arabidopsis

Seeds were surface-sterilized for 4 min. in 80% ethanol and subsequently for 1 min. in 100%

ethanol. Seeds were placed on plates containing ½ MS supplemented with 1% (D+) sucrose and

0.8% Agar. Alternatively, seeds were directly placed on sterilized soil. Seeds were incubated at

8°C for 2-4 days in the dark for stratification before placing them in growth chambers. Plants were

further cultivated under short day conditions at 22°C and 70 µmol photons/m^{2}s in an 8h/16h

light/dark cycle. For bolting assays, plants were placed in a phyto-chamber at 22°C and 70 µmol

photons/m^{2}s in a 16h/8h light/dark cycle (long day condition). Rosette leaves and days since

sowing were counted upon appearance of the shoot in the middle of the rosette.

Screening of Arabidopsis AARE mutants

Arabidopsis thaliana lines with T-DNA insertions in the At4G14570 locus were identified from the

public T-DNA Express database at the SIGnAL website (Salk Institute Genomic Analysis

Laboratory). Lines GK-456A10, SALK_080653C, SALK_071125C, and SALK_205137C, were

obtained from the Nottingham Arabidopsis Stock Centre. Homozygous mutant alleles were

verified   by   PCR   using   the   following   primers: forward   LB   GK   5´-

ATATTGACCATCATACTCATTGC-3´ and reverse GK-456 5´-CTTCAAAGAAACACCAATCAG-

3´for the GK-456A10 line, and forward LB-pROK 5´-GCGTGGACCGCTTGCTGCAACT-3´ and

reverse Salk_53 5´-TCTTTAGCCGAATCAGTTCCAGA-3´for the SALK_080653C,

SALK_071125C, and SALK_205137C lines. Identified homozygous mutant plants were back-

crossed with Arabidopsis Col-0. The F2 generation was screened for homozygous mutant plants

using the above listed primer-sets to identify the mutant allele or substituting the forward primers

with forward WT-GK-456 5´-AAGATGCTTTGCAGTCTCTA-3´ and forward WT-Salk 5´-

ACTGCCTTATGATCCATTGTCTC-3´, to identify the GK and SALK lines WT alleles, respectively.

RT-PCR was additionally performed to check for the presence of AtAARE transcripts using Taq

polymerase on cDNA prepared as described above with primers At1-At4. All primer combinations

are listed in Supplementary Table S4.

AARE exopeptidase activity

The enzyme activity assay according to [17] was modified. Here, tissue (80-100 mg) was

homogenized in liquid nitrogen and dissolved in 1 mL extraction buffer (50 mM PBS, 1 mM EDTA,

2 mM DTT). After centrifugation at 20,000 x g for 20 min at 4°C, 300 µl supernatant was mixed

with 700 µL reaction buffer (50 mM HEPES-KOH, pH 7.5 containing 1 mM Ac-Ala-pNA (Bachem,

Bubendorf, Switzerland) or 50 mM HEPES-KOH, pH 7.5, 10% DMSO containing 1mM AcLeu-

pNA (Bachem) and incubated at 37°C for 30-120 min. The reaction was stopped by the addition

of 500 µL 30% acetic acid. Absorbance was measured at 410 nm in a photospectrometer. This

approach was used to generate the data of Fig. 3b and Fig. S6a-d. Later the method was modified

(Fig. 3g, Fig. S6e-g). Here, tissue was homogenized in liquid nitrogen and dissolved in 100 µL

extraction buffer (50 mM PBS, 1 mM EDTA, 2 mM DTT) per 10 mg FW. After centrifugation at

20,000 x g for 20 min at 4°C, 5 µL supernatant was mixed with 195 µL reaction buffer (50 mM

HEPES-KOH, pH 7.5 containing 1 mM Ac-Ala-pNA (Bachem, Bubendorf, Switzerland)) in a 96

well micro titer plate and incubated at 37°C for 30-120 min. In the case of Ac-Leu-pNA as

substrate, 50 mM HEPES-KOH, pH 7.5 with 10% DMSO containing 1mM AcLeu-pNA (Bachem)

was used as reaction buffer. Every biological sample was measured in three technical replicates.

Absorbance was measured at 410 nm. Activity was calculated using a molar absorbance

coefficient[21] of 8.8 mM*cm^{-1} and represents the release of pNA [µmol] per minute normalized to

the total protein concentration of the sample. The protein concentration was determined using the

A_{280} method of a NanoDrop^{TM} (Thermo Fisher Scientific) or with a Bradford assay[135].

Western Blots

Western Blots were performed as described[74] using the *ECL Advance detection kit* (GE

Healthcare). The primary antibody against *A. thaliana* AARE was kindly provided by Dr. Yasuo

Yamauchi[17]. The primary antibody was diluted 1:10,000 in TBST-Buffer with 2% Blocking (GE

Healthcare) and incubated on the membrane for 2 h. As secondary antibody anti-Guinea pig

antibody, coupled to a horseradish peroxidase (Agrisera, AS 10 1496) diluted 1:10,000 in 2%

TBST with 2% Blocking (GE Healthcare), was applied for 1 h.

Detection of oxidized proteins

Plant tissues were homogenized in liquid nitrogen and proteins were extracted in 50 mM PBS, 50

mM DTT, 1 mM EDTA. 3-6 µg total protein was derivatized with DNPH and subsequently detected

with an anti-DNP antibody according to the manufacturer's instruction of the *OxyBlot Protein*

*Oxidation Detectio*n Kit (S7150, Sigma-Aldrich). Equal amounts of the derivatized protein samples

were employed as loading control on a separate SDS-gel and stained with Coomassie or silver

staining.

Flow cytometry

Flow cytometry analysis was performed as described[79]. Here, protonemata were chopped with a

razor blade in a small petri dish (6 cm diameter) in 2 mL of DAPI-buffer containing 0.01 mg/L 4′,6-

Diamidin-2-phenylindol (DAPI), 1.07 g/L $MgCl_2$ x 6 H_2O, 5 g/L NaCl, 21.11g/L Tris, 0.1% Triton,

pH 7. The solution was filtered using 30 µm pore size filters and the fluorescence intensity was

measured using a Cyflow®Space flow cytometry system (Partec, Munich, Germany).

Computational predictions

Predictions for the presence of cleavable targeting peptides were performed with *TargetP2.0*[63].

Additional predictions of subcellular localizations were performed with *LOCALIZER*[64]. The

presence of peroxisomal targeting signals was predicted with *PredPlantPTS1*[66,67]. Prediction of

protein domains was performed using *InterProScan*[135] and protein domain annotations according

to PFAM[137] were used.

Co-Immunoprecipitation

Co-immunoprecipitation was performed using GFP-Trap Magnetic Particles M-270 (Chromotek,

Planegg-Martinsried, Germany) as recommended by the manufacturer with modifications. 300

mg protonema was homogenized in a 2 mL reaction tube using a tungsten and a glass bead. For

each line three biological replicates were realized. The extraction buffer was chosen according to

the manufacturer's recommendations for plant samples and contained 25 mM HEPES-KOH,

pH7.5, 2 mM EDTA, 100 mM NaCl, 200 nM DTT, 0.5% Triton X-100, 1% plant protease inhibitor

cocktail (PPI, P9599, Sigma Aldrich). Ground plant material was dissolved in a final volume of 2

mL ice-cold extraction buffer and incubated for 10 min in a water quench ultrasonic device.

Samples were centrifuged at 4°C at 20,000xg for 30 min. For each sample 25 µL magnetic particle

slurry was washed with 500 µL extraction buffer. The sample supernatant was transferred to the

cleaned beads and incubated, rotating for 1 h at 6°C. Subsequently, beads were washed with 1

mL extraction buffer without Triton and PPI and again with 500 µL. Beads were then dissolved in

500 µL wash buffer (10 mM Tris-HCl, pH7.5, 150 mM NaCl, 0.5 mM EDTA), transferred to a new

reaction tube and washed again in 500 µL wash buffer. A RapiGest solution (0.2% in 50 mM Tris-

HCl, pH7.5; RapiGest SF Surfactant (Waters, Milford, MA, USA) was mixed 3:1 with 5 mM DTT

in 50 mM Tris-HCl, pH7.5. 30 µL of the resulting mixture was applied to each sample. Samples

were incubated at 95°C for 5 min under continuous shaking. Samples were cooled down to RT

and 5 µL of a trypsin (V5117, P) solution (0.025 µg/µL in 50 mM Tris-HCl, pH 7.5) were added to

each sample. Digestion on the beads was performed for 30 min at 32°C under continuous

shaking. Supernatants were transferred to new reaction tubes and the remaining beads were

washed twice with 50 µL 5 mM Iodoacetamide solution (in 50 mM Tris-HCl, pH7.5). The wash

supernatants were combined with the trypsin-containing supernatant and incubated over night at

30°C under continuous shaking. Acid-catalysed cleavage of the RapiGest surfactant was

performed as recommended by the manufacturer. Samples were purified using C18-STAGE-Tips

as described[138] and eluted from the Tip in 30% ACN in 0.1% FA.

Mass spectrometry measurement and data analysis

MS analysis was performed on an Orbitrap Q-Exactive Plus instrument (Thermo Fisher Scientific)

coupled to an UltiMate 3000 RSLCnano system (Dionex LC Packings/Thermo Fisher Scientific)

as described[139]. Database search and label-free quantitation was performed using *MaxQuant*

software V 1.6.0.16[36140]. For each Co-IP a specific database was employed containing all V3.3

proteins of Physcomitrella[39] as well as the sequence of the respective fusion protein. Additionally,

the contaminant list provided by the software was included. Decoys were generated on the fly by

reverting the given protein sequences. Variable modifications were formation of pyro Glu (N term

Q, −17.026549 Da), oxidation (M, +15.994915 Da), acetylation (N-term, +42.010565 Da) and

deamidation (N, +0.9 84016 Da). Carbamidomethylation (C, +57. 021464 Da) was specified as

fixed modification. Enzymatic specificity was set to tryptic with semi-specific free N-terminus. An

FDR of 0.01 was set for protein identification. LFQ values[141] were used as quantitative values.

Interaction analysis was performed in Perseus V 1.6.12.0[142]. Missing values were imputed from

a normal distribution with a down-shift of 1.8 and distribution width of 0.3. Interaction partners

were accepted at an FDR of 0.01 and a p-value less than 0.01.

Raw files of the test-IP measurements (Fig. S8e) were processed using Mascot Distiller V2.7.10

and searched against all Physcomitrella protein models V3.3[39] using Mascot Server V2.7.0

(Matrix Science).Processed mgf files from immunoprecipitation experiments targeting N-terminal

arginylation (PXD003232[36,143]) and the test-IPs were searched again against all Physcomitrella

protein models V3.3[39] using Mascot Server V2.7.0 (Matrix Science). The precursor mass

tolerance was 5 ppm and the fragment mass tolerance was 0.02 Da. Variable modifications were

formation of pyro Glu (N term Q, −17.026549 Da), oxidation (M, +15.994915 Da), acetylation (N-

term, +42.010565 Da) and deamidation (N, +0.9 84016 Da). Carbamidomethylation (C, +57.

021464 Da) was specified as fixed modification. Enzymatic specificity was set to tryptic with semi-

specific free N-terminus. Search results were loaded in Scaffold™ 5 (V5.0.1, Proteome Software)

and proteins were accepted at an FDR = 1 and peptides at an FDR = 0.5. A table of identified

proteins is accessible in Supplementary Table S7.

Multiple sequence alignment and phylogenetic reconstruction

Homologous protein sequences were aligned with UPP[144] (version 4.4.0) using default

parameters and subsequently translated into a codon-aware CDS alignment with PAL2NAL[145]

(version 1.4). Based on this multiple coding sequence alignment we reconstructed a maximum

likelihood tree with RAxML[146] (version 8.2.12) using the GTRCAT model with 1000 rapid bootstrap

samples. The tree was rooted at the split between animal and plant sequences and plotted in R[147]

using the packages ape[148] and ggtree[149].

Statistics and reproducibility

Statistical differences in datasets were analysed with one-way Anova with subsequent post-hoc

test if different lines at same conditions were compared. Two-way Anova with subsequent post-

hoc test was performed to analyse differences between lines at different conditions. Sample sizes

of biological replicates are specified in the figure legends. Anova and post-hoc analysis was

performed in R[147]. Statistical significance was accepted at $p < 0.05$ (*), $p < 0.01$ (**) and $p < 0.001$

(***).

Data availability

The authors confirm that all relevant data supporting the findings of this study are available within

the paper and its supplementary files. The mass spectrometry proteomics data have been

deposited at the ProteomeXchange Consortium via the PRIDE partner repository[84,85] with the

dataset identifier PXD033854 and 10.6019/PXD038742. Plant lines used in this study are

available upon reasonable request from the corresponding author (R.R.) or *via* the International
Moss Stock Center (IMSC, www.moss-stock-center.org). IMSC accessions are listed in
Supplementary Table S4. Full Blot images for Fig. 6b and Fig. S3b are available in Fig. S10.
Supplementary Table S8 contains all numeric source data used to generate the graphs and charts
in this study. Plasmids generated in this study are available from the International Moss Stock
Center IMSC (https://www.moss-stock-center.org) with the accession numbers P1519 (*PpAARE1*
KO), P1841 (*PpAARE2* KO (pJet)), P1520 (*PpAARE3* KO (pJet)), P1655 (*PpAARE1:Citrine* KI
(pJet)), P1813 (*PpAARE2:Citrine* KI (pJet)), P1814 (*PpAARE3:Citrine* KI), P1833
(*PpAARE1_1:eGFP* (pMAV4)), P1834 (*PpAARE1_2:eGFP* (pMAV4)), P1853
(*PpAARE1_Nt:eGFP* (pMAV4)), P1855 (*PpAARE3:eGFP* (pMAV4)), P1856 (*PpAARE3:eGFP*
(pMAV4)), P1854 (*AtAARE_SV:eGFP* (pMAV4)), P1881 (*AtAARE_LV:eGFP* (pMAV4)) and
P1862 (*AtAARE_Nt:eGFP* (pMAV4)).

Acknowledgements

We thank Christine Glockner, Agnes Novakovic and Eija Schulze for expert technical assistance
and Anne Katrin Prowse for language editing. Support with the Arabidopsis assays from Dr.
Philipp Schwenk is gratefully acknowledged. We thank Prof. Dr. Bettina Warscheid for the
possibility to use the QExactive Plus instrument and Prof. Dr. Yasuo Yamauchi for the primary
antibody against *A. thaliana* AARE. We gratefully acknowledge funding by the Deutsche
Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence
Strategy EXC-2189 (CIBSS to R.R.) and by the Wissenschaftliche Gesellschaft Freiburg. Open
Access funding enabled and organized by Projekt DEAL.

Author contributions

845 S.N.W.H. designed research, performed experiments, analysed data, and wrote the manuscript.
846 B.Ö. and N.v.G. analysed data and helped writing the manuscript. A.A.M. analysed data. B.R.v.B.,
847 L.N., J.S.F., R.K., S.G., T.W., and F.S. performed experiments. M.R.F. and S.J.M.S. designed
848 research and helped writing the manuscript. R.R. designed and supervised research, acquired
849 funding, and wrote the manuscript. All authors approved the final version of the manuscript.

Competing interests

851 All authors declare to have no competing interests.

References

1. Mittler, R., Zandalinas, S. I., Fichman, Y. & Van Breusegem, F. Reactive oxygen species signalling in plant stress responses. *Nature Reviews Molecular Cell Biology* **23**, 663-679 (2022).

2. Tola, A. J., Jaballi, A. & Missihoun, T. D. Protein carbonylation: Emerging roles in plant redox biology and future prospects. *Plants* **10**, 1451 (2021).

3. Gratão, P. L., Polle, A., Lea, P. J. & Azevedo, R. A. Making the life of heavy metal-stressed plants a little easier. *Functional Plant Biology* **32**, 481-494 (2005).

4. Smirnoff, N. Ascorbic acid: metabolism and functions of a multi-facetted molecule. *Current Opinion in Plant Biology* **3**, 229-235 (2000).

5. Sharma, P., Jha, A. B., Dubey, R. S. & Pessarakli, M. Reactive oxygen species, oxidative damage, and antioxidative defense mechanism in plants under stressful conditions. *Journal of Botany* **2012**: 217037 (2012).

6. Müller-Schüssele, S. J., et al. Chloroplasts require glutathione reductase to balance reactive oxygen species and maintain efficient photosynthesis. *Plant Journal* **103**, 1140-1154 (2020).

7. Soares, C., Carvalho, M. E., Azevedo, R. A. & Fidalgo, F. Plants facing oxidative challenges—A little help from the antioxidant networks. *Environmental and Experimental Botany* **161**, 4-25 (2019)

8. Parsell, D. A., Kowal, A. S., Singer, M. A. &, Lindquist, S. Protein disaggregation mediated by heat-shock protein Hspl04. *Nature* **372**, 475-478 (1994).

9. Queitsch, C., Hong, S. W., Vierling, E. & Lindquist, S. Heat shock protein 101 plays a crucial role in thermotolerance in Arabidopsis. *Plant Cell* **12**, 479-492 (2000).

10. Mogk, A., Bukau, B. &, Kampinga, H. H. Cellular handling of protein aggregates by disaggregation machines. *Molecular Cell* **69**, 214-226 (2018).

11. Møller, I. M., Jensen, P. E. & Hansson, A. Oxidative modifications to cellular components in plants. *Annual Review of Plant Biology* **58**, 459-481 (2007).

12. Ciacka, K., Tymiński, M., Gniazdowska, A. & Krasuska, U. Carbonylation of proteins—an element of plant ageing. *Planta* **252**, 12 (2020).

13. Farooq, M. A., et al. Acquiring control: The evolution of ROS-Induced oxidative stress and redox signaling pathways in plant stress responses. *Plant Physiology and Biochemistry* **141**, 353-369 (2019).

14. Bartlam, M., et al. Crystal structure of an acylpeptide hydrolase/esterase from *Aeropyrum pernix K1*. *Structure* **12**, 1481-1488 (2004).

15. Brunialti, E. A., Gatti-Lafranconi, P. & Lotti, M. Promiscuity, stability and cold adaptation
of a newly isolated acylaminoacyl peptidase. *Biochimie* **93**, 1543-1554 (2011).

16. Tsunasawa, S., Narita, K. & Ogata, K. Purification and properties of acylamino acid-
releasing enzyme from rat liver. *Journal of Biochemistry* **77**, 89-102 (1975).

17. Yamauchi, Y., Ejiri, Y., Toyoda, Y. & Tanaka, K. Identification and biochemical
characterization of plant acylamino acid–releasing enzyme. *Journal of Biochemistry* **134**,
251-257 (2003).

18. Rawlings, N. D., et al. The MEROPS database of proteolytic enzymes, their substrates
and inhibitors in 2017 and a comparison with peptidases in the PANTHER database.
Nucleic Acids Research **46**, D624-D632 (2018).

19. Fujino, T., et al. Identification of the cleavage sites of oxidized protein that are susceptible
to oxidized protein hydrolase (OPH) in the primary and tertiary structures of the protein.
Journal of Biochemistry **127**, 1087-1093 (2000).

20. Shimizu, K., et al. Overexpression of oxidized protein hydrolase protects COS-7 cells from
oxidative stress-induced inhibition of cell growth and survival. *Biochemical and Biophysical*
Research Communications **304**, 766-771 (2003).

21. Nakai, A., Yamauchi, Y., Sumi, S. & Tanaka, K. Role of acylamino acid-releasing
enzyme/oxidized protein hydrolase in sustaining homeostasis of the cytoplasmic
antioxidative system. *Planta* **236**, 427–436 (2012).

22. Gogliettino, M., et al. A novel class of bifunctional acylpeptide hydrolases–potential role in
the antioxidant defense systems of the Antarctic fish *Trematomus bernacchii*. *FEBS*
Journal **281**, 401-415 (2014).

23. Kiss, A. L., et al. The acylaminoacyl peptidase from *Aeropyrum pernix* K1 thought to be
an exopeptidase displays endopeptidase activity. *Journal of Molecular Biology* **368**, 509-
520 (2007).

24. Szeltner, Z., et al. Characterization of a novel acylaminoacyl peptidase with hexameric
structure and endopeptidase activity. *Biochimica et Biophysica Acta (BBA)-Proteins and*
Proteomics **1794**, 1204-1210 (2009).

25. Gogliettino, M., et al. Identification and characterisation of a novel acylpeptide hydrolase
from *Sulfolobus solfataricus*: structural and functional insights. *PLOS ONE* **7**, e37921
(2012).

26. Scaloni, A., et al. Deficiency of acylpeptide hydrolase in small-cell lung carcinoma cell
lines. *Journal of Laboratory and Clinical Medicine* **120**, 546-552 (1992).

27. Riccio, A., et al. A new APEH cluster with antioxidant functions in the antarctic hemoglobinless icefish *Chionodraco hamatus*. *PLOS ONE* **10**, e0125594 (2015).

28. Gogliettino, M., et al. Selective inhibition of acylpeptide hydrolase in SAOS-2 osteosarcoma cells: is this enzyme a viable anticancer target? *Molecular Biology Reports* **48**, 1505-1519 (2021).

29. Tangri, A., et al. Deubiquitinase UCHL1 maintains protein homeostasis through the PSMA7–APEH–Proteasome axis in high-grade serous ovarian carcinoma. *Molecular Cancer Research* **19**, 1168-1181 (2021).

30. Palumbo, R., et al. APEH inhibition affects osteosarcoma cell viability via downregulation of the proteasome. *International Journal of Molecular Sciences* **17**, 1614 (2016).

31. Shimizu, K., Kiuchi, Y., Ando, K., Hayakawa, M. & Kikugawa, K. Coordination of oxidized protein hydrolase and the proteasome in the clearance of cytotoxic denatured proteins. *Biochemical and Biophysical Research Communications* **324**, 140–146 (2004).

32. Palmieri, G., et al. Acylpeptide hydrolase inhibition as targeted strategy to induce proteasomal down-regulation. *PLOS ONE* **6**, e25888 (2011).

33. Bergamo, P., et al. RedOx status, proteasome and APEH: Insights into anticancer mechanisms of t10, c12-conjugated linoleic acid isomer on A375 melanoma cells. *PLOS ONE* **8**, e80900 (2013).

34. Yamin, R., Zhao, C., O'Connor, P. B., McKee, A. C. & Abraham, C. R. Acyl peptide hydrolase degrades monomeric and oligomeric amyloid-beta peptide. *Molecular Neurodegeneration* **4**, 33 (2009).

35. Bachovchin, D. A., et al. A high-throughput, multiplexed assay for superfamily-wide profiling of enzyme activity. *Nature Chemical Biology* **10**, 656-663 (2014).

36. Hoernstein, S. N. W., et al. Identification of targets and interaction partners of arginyl-tRNA protein transferase in the moss *Physcomitrella patens*. *Molecular & Cellular Proteomics* **15**, 1808-1822 (2016).

37. Schuessele, C., et al. Spatio-temporal patterning of arginyl-tRNA protein transferase (ATE) contributes to gametophytic development in a moss. *New Phytologist* **209**, 1014–1027 (2016).

38. Varshavsky, A. N-degron and C-degron pathways of protein degradation. *Proceedings of the National Academy of Sciences U.S.A.* **116**, 358-366 (2019).

39. Lang, D., et al. The *Physcomitrella patens* chromosome-scale assembly reveals moss genome structure and evolution. *Plant Journal* **93**, 515-533 (2018).

40. Chen, F., Mackey, A. J., Stoeckert Jr, C. J. & Roos DS OrthoMCL-DB: querying a
comprehensive multi-species collection of ortholog groups. *Nucleic Acids Research* **34**,
D363-D368 (2006).

41. Zimmer, A. D., et al. Reannotation and extended community resources for the genome of
the non-seed plant *Physcomitrella patens* provide insights into the evolution of plant gene
structures and functions. *BMC Genomics* **14**, 498 (2013).

42. Fernandez-Pozo, N., et al. PEATmoss (Physcomitrella Expression Atlas Tool): a unified
gene expression atlas for the model plant *Physcomitrella patens. Plant Journal* **102**, 165-
177 (2020).

43. Mergner, J., et al. Mass-spectrometry-based draft of the *Arabidopsis* proteome. *Nature*
579, 409-414 (2020).

44. Altschul, S. F., et al. Gapped BLAST and PSI-BLAST: a new generation of protein
database search programs. *Nucleic Acids Research* **25**, 3389-3402 (1997).

45. Merchant, S. S., et al. The *Chlamydomonas* genome reveals the evolution of key animal
and plant functions. *Science* **318**, 245-250 (2007).

46. Bowman, J. L., et al. Insights into land plant evolution garnered from the *Marchantia*
polymorpha genome. *Cell* **171**, 287-304 (2017).

47. Banks, J. A., et al. The *Selaginella* genome identifies genetic changes associated with the
evolution of vascular plants. *Science* **332**, 960-963 (2011).

48. Ouyang, S., et al. The TIGR rice genome annotation resource: improvements and new
features. *Nucleic Acids Research* **35**, D883-D887 (2007).

49. Tuskan, G. A., et al. The genome of black cottonwood, *Populus trichocarpa* (Torr. & Gray).
Science **313**, 1596-1604 (2006).

50. Hori, K., et al. *Klebsormidium flaccidum* genome reveals primary factors for plant terrestrial
adaptation. *Nature Communications* **5**, 3978 (2014).

51. Kirbis, A., et al. Transcriptional landscapes of divergent sporophyte development in two
mosses, *Physcomitrium* (Physcomitrella) *patens* and *Funaria hygrometrica. Frontiers in*
Plant Science **11**, 747 (2020).

52. Ostendorf, A. K., et al. Polyploidization within the Funariaceae – a key principle behind
speciation, sporophyte reduction and the high variance of spore diameters? *Bryophyte*
Diversity and Evolution **43**, 164-179 (2021).

53. Lamesch, P., et al. The Arabidopsis Information Resource (TAIR): improved gene
annotation and new tools. *Nucleic Acids Research* **40**, D1202-D1210 (2012).

54. Rawlings, N. D., Polgar, L. & Barrett, A. J. A new family of serine-type peptidases related
to prolyl oligopeptidase. *Biochemical Journal* **279**, 907-908 (1991).

55. Polgar, L. The prolyl oligopeptidase family. *Cellular and Molecular Life Sciences* **59**, 349-
362 (2002).

56. Perroud, P. F., et al. The *Physcomitrella patens* gene atlas project: large-scale RNA-seq
based expression data. *Plant Journal* **95**, 168-182 (2018).

57. Mueller, S. J., et al. Quantitative analysis of the mitochondrial and plastid proteomes of
the moss *Physcomitrella patens* reveals protein macrocompartmentation and
microcompartmentation. *Plant Physiology* **164**, 2081-2095 (2014).

58. Fesenko, I. A., et al. Specific pools of endogenous peptides are present in gametophore,
protonema, and protoplast cells of the moss *Physcomitrella patens*. *BMC Plant Biology*
15, 87 (2015).

59. Fesenko, I., et al. The *Physcomitrella patens* chloroplast proteome changes in response
to protoplastation. *Frontiers in Plant Science* **7**, 1661 (2016).

60. Liu, H., et al. The emergence and evolution of intron-poor and intronless genes in intron-
rich plant gene families. *Plant Journal* **105**, 1072-1082 (2021).

61. Beike, A. K. et al. Insights from the cold transcriptome of *Physcomitrella patens*: global
specialization pattern of conserved transcriptional regulators and identification of orphan
genes involved in cold acclimation. *New Phytologist* **205**, 869-881 (2015).

62. Cheng, C. Y., et al. Araport11: a complete reannotation of the *Arabidopsis thaliana*
reference genome. *Plant Journal* **89**, 789-804 (2017).

63. Armenteros, J. J. A., et al. Detecting sequence signals in targeting peptides using deep
learning. *Life Science Alliance* **2**, e201900429 (2019).

64. Sperschneider, J., et al. LOCALIZER: subcellular localization prediction of both plant and
effector proteins in the plant cell. *Scientific Reports* **7**, 44598 (2017).

65. Waszczak, C., Carmody, M. & Kangasjärvi, J. Reactive oxygen species in plant signaling.
Annual Review of Plant Biology **69**, 209-236 (2018).

66. Lingner, T., et al. Identification of novel plant peroxisomal targeting signals by a
combination of machine learning methods and *in vivo* subcellular targeting analyses. *Plant*
Cell **23**, 1556-1572 (2011).

67. Reumann, S., Buchwald, D. & Lingner, T. PredPlantPTS1: a web server for the prediction
of plant peroxisomal proteins. *Frontiers in Plant Science* **3**, 194 (2012).

68. Gould, S. J., Keller, G. A., Hosken, N., Wilkinson, J. & Subramani, S. A conserved
tripeptide sorts proteins to peroxisomes. *Journal of Cell Biology* **108**, 1657-1664 (1989).

69. Wang, J., Wang, Y., Gao, C., Jiang, L. & Guo, D. PPero, a computational model for plant
PTS1 type peroxisomal protein prediction. *PLOS ONE* **12**, e0168912 (2017).

70. Petriv, I., Tang, L., Titorenko, V. I. & Rachubinski, R. A. A new definition for the consensus
sequence of the peroxisome targeting signal type 2. *Journal of Molecular Biology* **341**,
119-134 (2004).

71. Reumann, S. Specification of the peroxisome targeting signals type 1 and type 2 of plant
peroxisomes by bioinformatics analyses. *Plant Physiology* **135**, 783-800 (2004).

72. Weise, A., et al. Use of *Physcomitrella patens actin* 5′ regions for high transgene
expression: importance of 5′ introns. *Applied Microbiology and Biotechnology* **70**, 337-345
(2006).

73. Özdemir, B., et al. Cytological analysis and structural quantification of FtsZ1-2 and FtsZ2-
1 network characteristics in *Physcomitrella patens*. *Scientific Reports* **8**, 11165 (2018).

74. Lang, E. G. E., et al. Simultaneous isolation of pure and intact chloroplasts and
mitochondria from moss as the basis for sub-cellular proteomics. *Plant Cell Reports* **30**,
205–215 (2011).

75. Fuss, J., Liegmann, O., Krause, K. & Rensing, S. A. Green targeting predictor and
ambiguous targeting predictor 2: the pitfalls of plant protein targeting prediction and of
transient protein expression in heterologous systems. *New Phytologist* **200**, 1022-1033
(2013).

76. Orr, M. W., Mao, Y., Storz, G. & Qian, S. B. Alternative ORFs and small ORFs: shedding
light on the dark proteome. *Nucleic Acids Research* **48**, 1029-1042 (2020).

77. Hohe, A., et al. An improved and highly standardised transformation procedure allows
efficient production of single and multiple targeted gene-knockouts in a moss,
Physcomitrella patens. *Current Genetics* **44**, 339-347 (2004).

78. Noy-Malka, et al. A single CMT methyltransferase homolog is involved in CHG DNA
methylation and development of *Physcomitrella patens*. *Plant Molecular Biology* **84**, 719-
735 (2014).

79. Heck, M. A., et al. Axenic *in vitro* cultivation of 19 peat moss (*Sphagnum* L.) species as a
resource for basic biology, biotechnology, and paludiculture. *New Phytologist* **229**, 861-
876 (2021).

80. Kamisugi, Y., et al. The mechanism of gene targeting in *Physcomitrella patens*:
homologous recombination, concatenation and multiple integration. *Nucleic Acids*
Research **34**, 6205-6214 (2006).

81. Rempfer, C., et al. Autopolyploidization affects transcript patterns and gene targeting
frequencies in Physcomitrella. *Plant Cell Reports* **41**, 153-173 (2022).

82. Lorenz, S., Tintelnot, S., Reski, R. & Decker, E. L. Cyclin D-knockout uncouples
developmental progression from sugar availability. *Plant Molecular Biology* **53**, 227-236
(2003).

83. Dalle-Donne, I., et al. Protein carbonylation: 2, 4-dinitrophenylhydrazine reacts with both
aldehydes/ketones and sulfenic acids. *Free Radical Biology and Medicine* **46**, 1411-1419
(2009).

84. Deutsch, E. W., et al. The ProteomeXchange consortium in 2020: enabling 'big data'
approaches in proteomics. *Nucleic Acids Research* **48**, D1145-D1152 (2020).

85. Perez-Riverol, Y., et al. The PRIDE database resources in 2022: a hub for mass
spectrometry-based proteomics evidences. *Nucleic Acids Research* **50**, D543-D552
(2022).

86. Johansson, E., Olsson, O. & Nyström, T. Progression and specificity of protein oxidation
in the life cycle of *Arabidopsis thaliana*. *Journal of Biological Chemistry* **279**, 22204-22208
(2004).

87. Harman, D. Aging: A theory based on free radical and radiation chemistry. *Journal of
Gerontology* **11**, 298–300 (1956).

88. Harman, D. The biologic clock: the mitochondria? *Journal of the American Geriatrics
Society* **20**, 145-147 (1972).

89. Giorgi, C., et al. Mitochondria and reactive oxygen species in aging and age-related
diseases. *International Review of Cell and Molecular Biology* **340**, 209-344 (2018).

90. Mao, C., et al. Nitric oxide regulates seedling growth and mitochondrial responses in aged
oat seeds. *International Journal of Molecular Sciences* **19**, 1052 (2018).

91. Ratajczak, E., Małecka, A., Ciereszko, I. & Staszak, A. M. Mitochondria are important
determinants of the aging of seeds. *International Journal of Molecular Sciences* **20**, 1568
(2019).

92. Munné-Bosch, S. & Alegre, L. Plant aging increases oxidative stress in chloroplasts.
Planta **214**, 608-615 (2002).

93. Noctor, G., Veljovic-Jovanovic, S., Driscoll, S., Novitskaya, L. & Foyer, C. H. Drought and
oxidative load in the leaves of C3 plants: a predominant role for photorespiration? *Annals
of Botany* **89**, 841-850 (2002).

94. Foyer, C. H. & Noctor, G. Redox sensing and signalling associated with reactive oxygen in chloroplasts, peroxisomes and mitochondria. *Physiologia Plantarum* **119**, 355-364 (2003).

95. Sohal, R. S. & Weindruch, R. Oxidative stress, caloric restriction, and aging. *Science,* **273**, 59-63 (1996).

96. López-Otín, C., Blasco, M. A., Partridge, L., Serrano, M. & Kroemer, G. The hallmarks of aging. *Cell* **153**, 1194-1217 (2013).

97. Senthilkumar, R., Reddy, P. N. & Sharma, K. K. Studies on trypsin-modified bovine and human lens acylpeptide hydrolase. *Experimental Eye Research* **72**, 301-310 (2001).

98. Palmieri, G., et al. Low erythrocyte levels of proteasome and acyl-peptide hydrolase (APEH) activities in Alzheimer's disease: a sign of defective proteostasis? *Journal of Alzheimer's Disease* **60**, 1097-1106 (2017).

99. Marshall, I., et al. Analyzing the activity and expression of acyl peptide enzyme hydrolase (APEH) in the blood serum of patients with type II diabetes. *Bios* **90**, 70-78 (2019).

100. Resemann, H. C., et al. Convergence of sphingolipid desaturation across over 500 million years of plant evolution. *Nature Plants* **7**, 219-232 (2021).

101. Zybailov, B., et al. Sorting signals, N-terminal modifications, and abundance of the chloroplast proteome. *PLOS ONE* **3**, e1994 (2008).

102. Rugen, N., et al. Protein interaction patterns in *Arabidopsis thaliana* leaf mitochondria change in dependence to light. *Biochimica et Biophysica Acta (BBA)-Bioenergetics* **1862**, 148443 (2021).

103. Maxwell, D. P., Wang, Y. & McIntosh, L. The alternative oxidase lowers mitochondrial reactive oxygen production in plant cells. *Proceedings of the National Academy of Sciences U.S.A.* **96**, 8271-8276 (1999).

104. Møller, I. M. Plant mitochondria and oxidative stress: electron transport, NADPH turnover, and metabolism of reactive oxygen species. *Annual Review of Plant Biology* **52**, 561-591 (2001).

105. Smakowska, E., Czarna, M. & Janska, H. Mitochondrial ATP-dependent proteases in protection against accumulation of carbonylated proteins. *Mitochondrion* **19**, 245-251 (2014).

106. Al-Khatib, K. & Paulsen, G. M. Enhancement of thermal injury to photosynthesis in wheat plants and thylakoids by high light intensity. *Plant Physiology* **90**, 1041-1048 (1989).

107.   Wang, Q. L., Chen, J. H., He, N. Y. & Guo, F. Q. Metabolic reprogramming in chloroplasts under heat stress in plants. *International Journal of Molecular Sciences* **19**, 849 (2018).

108.   Sweetlove, L. J., et al. The impact of oxidative stress on Arabidopsis mitochondria. *Plant Journal* **32**, 891-904 (2002).

109.   Tiwari, B. S., Belenghi, B. & Levine, A. Oxidative stress increased respiration and generation of reactive oxygen species, resulting in ATP depletion, opening of mitochondrial permeability transition, and programmed cell death. *Plant Physiology* **128**, 1271-1281 (2002).

110.   Willems, P., Ndah, E., Jonckheere, V., van Breusegem, F. & van Damme, P. To new beginnings: Riboproteogenomics discovery of N-terminal proteoforms in *Arabidopsis thaliana*. *Frontiers in Plant Science* **12**, 778804 (2021).

111.   Kiessling, J., et al. Dual targeting of plastid division protein FtsZ to chloroplasts and the cytoplasm. *EMBO Reports* **5**, 889-894 (2004).

112.   van Wijk, K. J., et al. The Arabidopsis PeptideAtlas: Harnessing worldwide proteomics data to create a comprehensive community proteomics resource. *Plant Cell* **33**, 3421-3453 (2021).

113.   Zhang, H., et al. Arabidopsis proteome and the mass spectral assay library. *Scientific Data* **6**, 278 (2019).

114.   Zeng, Z., et al. Acylpeptide hydrolase is a component of the cellular response to DNA damage. *DNA Repair* **58**, 52-61 (2017).

115.   Brocca, S., et al. A bacterial acyl aminoacyl peptidase couples flexibility and stability as a result of cold adaptation. *FEBS Journal* **283**, 4310-4324 (2016).

116.   Menyhárd, D. K., et al. A self-compartmentalizing hexamer serine protease from *Pyrococcus horikoshii*: substrate selection achieved through multimerization. *Journal of Biological Chemistry* **288**, 17884-17894 (2013).

117.   Viaene, T., et al. Directional auxin transport mechanisms in early diverging land plants. *Current Biology* **24**, 2786-2791 (2014).

118.   Chen, Z., et al. Comprehensive analysis of the *Ppatg3* mutant reveals that autophagy plays important roles in gametophore senescence in *Physcomitrella patens*. *BMC Plant Biology* **20**, 440 (2020).

119.   Jia, M., et al. Noncanonical ATG8-ABS3 interaction controls senescence in plants. *Nature Plants* **5**, 212-224 (2019).

120. Dalle-Donne, I., Giustarini, D., Colombo, R., Rossi, R. & Milzani, A. Protein carbonylation in human diseases. *Trends in Molecular Medicine* **9**, 169-176 (2003).

121. Hipp, M. S., Kasturi, P. & Hartl, F. U. The proteostasis network and its decline in ageing. *Nature Reviews Molecular Cell Biology* **20**, 421-435 (2019).

122. Chen, Y. R., et al. Tissue-specific landscape of protein aggregation and quality control in an aging vertebrate. Preprint at: https://doi.org/10.1101/2022.02.26.482120 (2022).

123. Medina, R., et al. Phylogenomic delineation of *Physcomitrium* (Bryophyta: Funariaceae) based on targeted sequencing of nuclear exons and their flanking regions rejects the retention of *Physcomitrella*, *Physcomitridium* and *Aphanorrhegma*. *Journal of Systematics and Evolution* **57**, 404-417 (2019).

124. Reski, R. & Abel, W. O. Induction of budding on chloronemata and caulonemata of the moss, *Physcomitrella patens*, using isopentenyladenine. *Planta* **165**, 354-358 (1985).

125. Egener, T., et al. High frequency of phenotypic deviations in *Physcomitrella patens* plants transformed with a gene-disruption library. *BMC Plant Biology* **2**, 6 (2002).

126. Schween, G., Hohe, A., Koprivova, A. & Reski, R. Effects of nutrients, cell density and culture techniques on protoplast regeneration and early protonema development in a moss, *Physcomitrella patens*. *Journal of Plant Physiology* **160**, 209–212 (2003).

127. Erxleben, A., Gessler, A., Vervliet-Scheebaum, M. & Reski, R. Metabolite profiling of the moss *Physcomitrella patens* reveals evolutionary conservation of osmoprotective substances. *Plant Cell Reports* **31**, 427-436 (2012).

128. Tian, G. W., et al. High-throughput fluorescent tagging of full-length *Arabidopsis* gene products in planta. *Plant Physiology* **135**, 25-38 (2004).

129. Sambrook, J. & Russell, D. W. Standard ethanol precipitation of DNA in microcentrifuge tubes. *Cold Spring Harbor Protocols* **2006**, pdb-prot4456 (2006).

130. Hohe, A. & Reski, R. Optimisation of a bioreactor culture of the moss *Physcomitrella patens* for mass production of protoplasts. *Plant Science* **163**, 69-74 (2002).

131. Decker, E. L., et al. Strigolactone biosynthesis is evolutionarily conserved, regulated by phosphate starvation and contributes to resistance against phytopathogenic fungi in a moss, *Physcomitrella patens*. *New Phytologist* **216**, 455-468 (2017).

132. Gibson, D. G., et al. Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nature Methods* **6**, 343–345 (2009).

1184   133.   Amack, S. C. & Antunes, M. S. CaMV35S promoter–a plant biology and
1185   biotechnology workhorse in the era of synthetic biology. *Current Plant Biology* **24**, 100179
1186   (2020).

1187   134.   Horstmann, V., Huether, C. M., Jost, W., Reski, R. & Decker, E. L. Quantitative
1188   promoter analysis in *Physcomitrella patens*: a set of plant vectors activating gene
1189   expression within three orders of magnitude. *BMC Biotechnology* **4**, 13 (2004).

1190   135.   Bradford, M. M. A rapid and sensitive method for the quantitation of microgram
1191   quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*
1192   **72**, 248-254 (1976).

1193   136.   Blum, M., et al. The InterPro protein families and domains database: 20 years on.
1194   *Nucleic Acids Research* **49**, D344-D354 (2021).

1195   137.   Mistry, J., et al. Pfam: The protein families database in 2021. *Nucleic Acids*
1196   *Research* **49**, D412-D419 (2021).

1197   138.   Hoernstein, S. N. W., et al. Host cell proteome of *Physcomitrella patens* harbors
1198   proteases and protease inhibitors under bioproduction conditions. *Journal of Proteome*
1199   *Research* **17**, 3749-3760 (2018).

1200   139.   Top, O., et al. Expression of a human cDNA in moss results in spliced mRNAs and
1201   fragmentary protein isoforms. *Communications Biology* **4**, 964 (2021).

1202   140.   Cox, J. & Mann, M. MaxQuant enables high peptide identification rates,
1203   individualized ppb-range mass accuracies and proteome-wide protein quantification.
1204   *Nature Biotechnology* **26**, 1367-1372 (2008).

1205   141.   Cox, J., et al. Accurate proteome-wide label-free quantification by delayed
1206   normalization and maximal peptide ratio extraction, termed MaxLFQ. *Molecular & Cellular*
1207   *Proteomics* **13**, 2513-2526 (2014).

1208   142.   Tyanova, S., et al. The Perseus computational platform for comprehensive
1209   analysis of (prote)omics data. *Nature Methods* **13**, 731-740 (2016).

1210   143.   Hoernstein, S. N. W., et al. Identification of targets and interaction partners of
1211   arginyl-tRNA protein transferase in the moss *Physcomitrella patens*.
1212   http://dx.doi.org/10.6019/PXD003232 (2016).

1213   144.   Nguyen, N. D., Mirarab, S., Kumar, K. & Warnor, T. Ultra-large alignments using
1214   phylogeny-aware profiles. *Genome Biology* **16**, 124 (2015).

1215   145.   Suyama, M., Torrents, D. & Bork, P. PAL2NAL: robust conversion of protein
1216   sequence alignments into the corresponding codon alignments. *Nucleic Acids Research*
1217   **34**, W609–W612 (2006).

146.   Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis
of large phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

147.   R Core Team. R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/
(2022).

148.   Paradis, E. & Schliep, K. ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics* **35**, 526-528 (2018).

149.   Yu, G., Smith, D., Zhu, H., Guan, Y. & Lam, T. T. ggtree: an R package for
visualization and annotation of phylogenetic trees with their covariates and other
associated data. *Methods in Ecology and Evolution* **8**, 28-36 (2017).

1228

Supplementary Information

Figure S1 Expression levels of PpAARE genes and AtAARE in different tissues and under stress
conditions.

Figure S2 Sequence alignment of AtAARE and PtAARE isoforms.

Figure S3 AARE activity in Physcomitrella plastids and mitochondria.

Figure S4 Identification of PpAARE single, double and triple KO mutants by PCR on genomic
DNA.

Figure S5 FCM analysis of all confirmed PpAARE KO lines to analyse the ploidy.

Figure S6 AARE exopeptidase activity protonema of KO mutants.

Figure S7 Plate overview and additional OxyBlots from Physcomitrella gametophore colonies
with loading controls.

Figure S8 Screening for haploid Physcomitrella lines with a citrine fusion transcript and overview
on identification of the three PpAARE isoforms in different IP experiments.

Figure S9 Phenotypes of seedlings and adult AtAARE T-DNA mutant lines (GK, s68) and WT
(Col).

Supplementary Table S1 List of primers used to assemble KO and KI constructs and primers
used to screen for transgenic plants in Physcomitrella.

Supplementary Table S2 List of gene accession numbers for all sequences used for

phylogenetic analysis and corresponding subcellular localization predictions.

Supplementary Table S3 List of primers used to screen possible T-DNA mutant plants in

Arabidopsis.

Supplementary Table S4 List of accession numbers for the International Moss Stock Center

(IMSC) of all transgenic Physcomitrella lines.

Supplementary Table S5 List of primers used for qPCR on genomic DNA to determine copy

numbers of KO constructs in transgenic Physcomitrella lines.

Supplementary Table S6 List of primers used to assemble the constructs for transient

localization analysis in Physcomitrella protoplasts.

Supplementary Table S7 Spectrum report of the revised database search of the anti-Arg(Nt) IP

published in[36].

Supplementary Table S8 This table contains all numeric source data used to generate the

graphs in Fig. 3b, Fig. 3e-g, Fig. 5b, Fig. 6c, Fig. 6e, Fig. S1, Fig. S3c, Fig. S4j-p, Fig. S5, Fig.

S6, Fig. S8d, Fig. S8e and Fig. S8g.

1261

Fig. 1 Phylogenetic tree for AARE isoforms from selected organisms, gene structures and proteins domains of *P. patens* and *A. thaliana* AARE isoforms. (a) Maximum likelihood tree based on an alignment of multiple *AARE* coding sequences. Duplication events in several species appear to be lineage-specific. Node values reflect percentage confidence values based on 1000 bootstrap replicates. Species abbreviations: At: *Arabidopsis thaliana*; Pt: *Populus trichocarpa*; Os: *Oryza sativa*; Pp: *Physcomitrium patens*; Fh: *Funaria hygrometrica*; Sf: *Sphagnum fallax*; Mp: *Marchantia polymorpha*; Sm: *Selaginella moellendorffii*; Kn: *Klebsormidium nitens*; Cr: *Chlamydomonas reinhardtii*; Rn: *Rattus norvegicus*; Hs: *Homo sapiens*; Ch: *Chionodraco hamatus*. (b) Gene structure of *PpAARE1-3* and *AtAARE*. For *PpAARE1* two splice variants exist. For *AtAARE* two different 5' UTR definitions are present (upper: Araport11[62]; lower: TAIR10[53]). (c) Left: Both splice variants of *PpAARE1* were amplified from complementary DNA (cDNA; +: with cDNA template; -: without reverse transcriptase). Expected amplicon size: *PpAARE1_1*: 2512 bp; *PpAARE1_2*: 2596 bp. Primers were designed to be specific for the splice variants (Table S1). Right: The longer open reading frame of *AtAARE* was amplified from cDNA. Expected amplicon size: 2457 bp (Table S3). (d) Protein structures showing PFAM-domains for PpAARE1-3 and AtAARE. All isoforms contain an AARE N-terminal domain (PF19283) and a catalytic Peptidase S9 domain (PF00326). PpAARE1, PpAARE2 and AtAARE additionally contain a WD40 domain (PF07676). The long splice variant of PpAARE1 and the longer open reading frame of AtAARE encode a predicted N-terminal plastid transit peptide (cTP). AA: amino acid. Cleavable N-terminal sequences were predicted by *TargetP2.0*[63].

Fig. 2 Confocal microscopy images showing the localization of PpAARE isoforms and AtAARE with C-terminal fused eGFP in Physcomitrella protoplasts. Fusion proteins with domain structures are depicted. PpAARE1_1, PpAARE2 and PpAARE3 localize to the cytoplasm. PpAARE1_2 localizes to specific foci in plastids (white arrows) and to mitochondria. The N-terminal extension of PpAARE1_2 encoding a predicted plastid transit peptide (PpAARE1_Nt) directs eGFP to plastids and mitochondria. The short variant of AtAARE (SV) localizes to the cytoplasm. The long variant (LV) localizes to plastids and mitochondria. The N-terminal extension of the long variant of AtAARE localizes to plastids and mitochondria. Left image: chlorophyll autofluorescence (red) and eGFP (green). Middle image: chlorophyll autofluorescence (red) and MitoTracker™ (magenta). Right image: chlorophyll autofluorescence (red), eGFP (green), Mitotracker™ (magenta) and co-localization of eGFP and MitoTracker™ (white). Bars = 7 µm.

Based on this data, PpAARE1 is targeted to three organelles: chloroplasts, mitochondria and the cytosol. To independently scrutinize these findings, we measured AARE enzyme activity in the

organelles. Using organelle fractionation as previously described[74], we detected AARE activity in chloroplasts, the cytosol, and mitochondria, although to a lesser extent in the latter (Fig. S3). Thus, *in vivo* localization of fusion proteins and enzyme measurements after cell fractionation independently confirm the predicted triple localization of AARE in Physcomitrella.

Fig. 3 Phenotype analyses, AARE activity and level of oxidized proteins in Physcomitrella WT and KO mutants. (a) Each isoform was knocked out using a different selection marker. The selection markers confer resistance against G418 (npt), hygromycin (hpt) and zeocin (zeo), respectively. Homologous regions (HR) were chosen in order to remove the full CDS of the corresponding gene upon integration of the selection marker. (b) Mean AARE activity on AcAla-pNA in Physcomitrella protonema with standard deviation (n= three biological replicates). Significance levels are based on a one-way Anova and subsequent post-hoc test (***: p<0.001) and indicate differences compared to WT. (c) Phenotypes of gametophores of WT and the different KOs cultivated on KnopME for 4 months. All colonies were grown on the same plate. Upper panel: Bar = 0.5 mm; lower panel: Bar = 2mm. (d) Protonema colonies grown on KnopME with 0.02% glucose taken from suspension culture. Bar = 1 mm. Images were taken 9 days after spotting. (e) Box plot showing the distribution of colony areas (n = 12 colonies per boxplot; colony images taken 7 days after spotting). No significant difference of colony areas between WT and the KOs was observed (tested via one-way Anova). (f) Box plot showing the number of gametophores per colony and day of WT and ΔPpAARE1/2. Counting was performed at indicated days after spotting protonema suspension culture on solid medium (KnopME with 0.02% glucose). Indicated significance levels are based on a two-way Anova with subsequent post-hoc test (***: p<0.001). The boxplots (e, f) depict the mean (horizontal bold line) of the data, the interquartile range (box) and the 1.5x interquartile range (whiskers). (g) Mean AARE exopeptidase activity on AcAla-pNA in gametophores. Depicted is the mean of three independent colonies per line with standard deviations. All colonies were cultivated on the same plate (Fig. S7a). Significance levels are based on one-way Anova and indicate differences compared to WT (***: p<0.001). (h) Levels of oxidized proteins in gametophores of WT and the different KOs (Coomassie-stained loading control is shown in Fig. S7b). The analysis was repeated three times (additional Blots available in Fig. S7c, d). Detection of oxidized proteins was performed with the OxyBlot™ *Protein Oxidation Detection Kit* (Merck).

Fig. 4 Physcomitrella gametophore colonies of varying age. (a) Colonies of all representative knockout mutants after 5 months on solid medium (KnopME). Bar = 5 mm. Gametophores of ΔPpAARE1/2 and ΔPpAARE1/3/2 are only viable at the tip, whereas plant material at the base

(red arrow) is dead. (b) Colonies after 8 months on solid medium. ΔPpAARE1 is the parental line

for ΔPpAARE1/2. Gametophores of WT and ΔPpAARE1 are still viable whereas ΔPpAARE1/2

gametophores are mostly dead. Bar = 2 mm. (c) Colonies after 8 months on solid medium.

ΔPpAARE1 is the parental line for ΔPpAARE1/3 and ΔPpAARE1/3 is the parental line for

ΔPpAARE1/3/2. Colonies of WT, ΔPpAARE1 and ΔPpAARE1/3 still have viable gametophores,

whereas ΔPpAARE1/3/2 gametophores are dead. Bar = 2 mm.

Fig. 5 Reciprocal co-immunoprecipitation (Co-IP) with Citrine-tagged PpAARE isoforms.

(a) Tagging of PpAARE isoforms was realized by in-frame fusion with a linker-*Citrine* CDS at the

respective native locus *via* homologous recombination. Original stop codons were deleted. (b)

Volcano plots showing the result of the Co-IPs against each of the PpAARE:Citrine fusion

proteins. Left panel: Pulldown of PpAARE1:Citrine. Right panel: Pulldown of PpAARE2:Citrine.

Co-IP was performed with GFP-Trap Magnetic Particles M-270 (Chromotek) and protonema from

suspension culture. Log_2 ratios of normalized label-free quantitation values (LFQ) are plotted

against $–log_{10}$ of adjusted p-values. Proteins significantly enriched in the Citrine tagged pulldown

are shown in blue circles (p < 0.01, FDR = 0.01). Significantly enriched PpAARE isoforms are

depicted as red crosses.

Fig. 6 Molecular and phenotypical characterization of *A. thaliana* T-DNA mutant lines. (a)

Structure of the AtAARE gene and positions of T-DNA insertions (red arrows indicate positions of

T-DNA insertion, s68 (SALK_071125) and GK (GK-456A10)) and primers (At1, At2) used for RT-

PCR (below). Transcription analysis of *AtAARE* was performed by RT-PCR in WT (Col), s68, and

GK plants. Negative controls without reverse transcriptase (-), a water control (H_2O) and genomic

DNA control (gDNA) are indicated. Expected amplicon sizes: *AtAARE*: cDNA 739 bp, gDNA: 1204

bp; *PP2AA3* (reference gene): cDNA: 517 bp, gDNA: 911 bp. Primers are listed in Table S3. (b)

Western blot analysis on extracts of the two T-DNA mutants and WT probed with a polyclonal

@AARE antibody. In both T-DNA lines AtAARE is not detectable. Full blot images are available

in Fig. S10. (c) Mean exopeptidase activity in *A. thaliana* extracts on AcAla-pNA and AcLeu-pNA

with standard deviation (n = 3 biological replicates). Significance levels are based on one-way

Anova and subsequent post-hoc test (***: p<0.001) and indicate differences compared to WT

(Col). (d) Detection of oxidized proteins in WT (Col) and *AtAARE* mutant (s68). From three

independent plants one exterior rosette leaf was taken after cultivation for 8 weeks in short-day

conditions (8h light/16h dark) Bars correspond to 2 cm. Protein loading control by Coomassie

staining is depicted below. (e) Boxplot of the bolting analysis of WT (Col, n = 14) and *AtAARE*

mutant plants (s68, n=13). The boxplot (e, f) depicts the mean (horizontal bold line) of the data,

the interquartile range (box) and the 1.5x interquartile range (whiskers). Outliers are depicted as

white dots. Significance level is based on a one-way Anova and subsequent post-hoc test (***:

p<0.001). Exemplary plants (45-day-old plants grown in long-day conditions, 16h light/8h dark)

are depicted below the boxplot. Bars correspond to 1 cm.

1365

PpAARE1_1:eGFP

AtAARE_SV:eGFP

PpAARE1_2:eGFP

AtAARE_LV:eGFP

PpAARE1_Nt:eGFP

AtAARE_Nt:eGFP

PpAARE2:eGFP

PpAARE3:eGFP

AARE N-terminal domain   WD40 domain   S9 Peptidase   Plastid transit peptide   GFP   50 AA