1 **Ancestral state reconstruction suggests repeated losses of**

2 **recruitment communication during ant evolution (Hymenoptera:**

3 **Formicidae)**

Simone M Glaser[1] and Christoph Grüter[1,2]

[1] Institute of Organismic and Molecular Evolutionary Biology, Johannes-Gutenberg

University of Mainz, Germany

[2] Present address: School of Biological Sciences, University of Bristol, 24 Tyndall

Avenue, BS8 1TQ Bristol, UK

Correspondence: simone-glaser1@gmx.de

Acknowledgement

We thank the Department of Behavioural Ecology and Social Evolution, Johannes

Gutenberg-University Mainz for feedback and advice. C.G. and S.M.G. were funded

by the German Research Foundation (DFG: GR 4986/1-1).

17 **Author contributions**

18 S.M.G and C.G. developed the study question. S.M.G. performed the literature

19 research, analyzed the data and wrote the primarily manuscript. S.M.G and C.G.

20 contributed critically to the drafts and gave final approval for publication.

Abstract

Eusocial insects have evolved different strategies to share information about their environment and workers can recruit nestmates to food sources or new nest sites. Ants are the most species-rich social insect group and are known to use pheromones, visual and tactile signals to communicate and inform nestmates about resources. However, how these different strategies evolved and whether there was a predominant evolutionary sequence that led to present day recruitment strategies is not well understood. In our study we explored two competing hypotheses about the ancestral recruitment communication: (1) ant ancestors did not recruit nestmates and species evolved more complex recruitment strategies over time vs. (2) early ants used mass-recruitment, which was lost repeatedly in some lineages. We combined an extensive search of the scientific literature and ancestral state reconstruction to estimate the ancestral recruitment strategy, focusing on the categories (i) no recruitment, (ii) tandem running, (iii) group-recruitment and (iv) chemical mass-recruitment. Stochastic character mapping suggests that mass-recruitment was ancestral in ants (59-61%), whereas "no recruitment" was unlikely to be the ancestral condition (21%). Similarly, marginal ancestral state reconstruction suggests that mass-recruitment (44-81%) or group-recruitment (48-50%) represented the original state. Our results are consistent with the finding that early ants lived in colonies containing up to several thousand individuals, which are typically associated with mass-recruiting in ants. However, our ability to robustly identify patterns in the evolution of communication in ants remains hampered by a lack of natural history information for most ant species.

Keywords: ancestral state, communication, Formicidae, recruitment, tandem running

Introduction

Communication about resources is widespread in social insects and different species have evolved a variety of strategies to communicate with other individuals (Czaczkes et al., 2015; Franklin, 2014; Grüter & Leadbeater, 2014; Hölldobler & Wilson, 1990; Leadbeater & Chittka, 2007; von Frisch, 1967; Wilson, 1971; Grüter, 2020). Senders can use visual, tactile (body contact in honeybees, Grooters, 1987; antennation, Hölldobler & Wilson, 1990; von Frisch, 1967) or chemical (e.g. pheromones, Czaczkes et al., 2015; Hölldobler & Wilson, 1990) signals to share information about food locations, nesting locations, dangers or the needs of individuals and the colony.

Ants (Family: Formicidae) are an extraordinarily diverse, widespread and ecologically important group of social insects, containing over 13,000 extant species (Reeves & Moreau, 2019). Ants probably first appeared in the early Cretaceous, more than 100 million years ago, most likely from a lineage of wasps (Hölldobler & Wilson 2009; Moreau et al., 2006; Ward, 2014). Ants have evolved a variety of different communication strategies, some of which are behaviourally highly complex (Hölldobler, 1999). There are also species that employ several strategies during recruitment (see below).

Several classification systems for recruitment behaviours have been proposed (Beckers et al., 1989; Jaffe, 1984; Lanan, 2014). The most common strategies are the use of pheromone trails (Czaczkes et al., 2015; Hölldobler & Wilson, 1990, 2009), group-recruitment (Hölldobler, 1971; Liefke et al., 2001), tandem recruitment (Franklin, 2014; Glaser & Grüter, 2018; Grüter et al., 2018) or no recruitment at all (solitary/individual) (Beckers et al., 1989; Jaffe, 1984). In tandem recruitment, a scout with knowledge of a valuable resource recruits a single nestmate to that resource. The follower maintains contact with the leader by antennating the gaster and hind legs of

the leader. In group-recruitment, a scout leads a group of up to thirty nestmates to a

resource (Beckers et al. 1989; Liefke et al., 2001). A short-lived, volatile pheromone

emitted by the leader helps recruited ants stay close to the leader (Möglich et al., 1974).

In mass-recruitment, successful scouts lay pheromone trails between the nest and the

resource, which is followed and strengthened by nestmates. Some species use a

particular strategy in one context but not in another. For example, *Neoponera* or

Diacamma species perform tandem runs when they relocate to a new nest site, but

when they are foraging for prey they do so solitarily (Fresneau, 1985; Kaur et al., 2017).

Furthermore, recruitment communication can depend on the type of food source that

is collected. When collecting small prey, ants often do not recruit nestmates as they

can carry home the food item by themselves, but use recruitment communication when

finding larger resources, like honey dew-secreting aphid colonies or large prey items

(Czaczkes et al., 2011; Czaczkes & Ratnieks, 2012; Detrain & Deneubourg, 2008;

Lach, 2005). These examples highlight that the use of recruitment communication

often depends on the ecological context.

Species with small colonies (<1000 individuals) often do not use recruitment or they

perform tandem runs, whereas medium size colonies (up to a few thousand individuals)

tend to recruit nestmates by group-recruitment or by pheromone trails and large

colonies use mainly pheromone trails (Beckers et al., 1989). Recruitment *via*

pheromone trails requires colonies to have a minimum number of foragers to deposit

pheromones in order to maintain the trail (Beekman et al., 2001). The link between

colony size and recruitment strategy is not rigid, however, and species with similar

colony sizes can differ in the strategies they employ (Beckers et al., 1989).

This diversity in recruitment strategies raises the question how these strategies

evolved and whether certain forms of recruitment tend to precede other methods. One

hypothesis is that recruitment communication increased in complexity over

evolutionary time. According to this scenario, early ants would have foraged solitarily,

like many present-day ponerine ants (Beckers et al., 1989; Maschwitz & Schönegge,

1983; Villet, 1990). Subsequently, small-scale communication mechanisms evolved,

like tandem-running and group-recruitment. From these forms of communication,

mass-recruitment with longer-lasting chemical trails may have evolved (Beckers et al.,

1989; Hölldobler & Wilson, 1990; Traniello, 1989). Hingston (1929) and Wilson (1959),

for instance, suggested that tandem running is an ancestral form of communication

that was used by early ants. Short-lived pheromones released by the tandem leader

play a potentially important role for the cohesion between the leader and follower in a

tandem run (Basari et al., 2014). It would then have been a small evolutionary step to

produce longer lasting trail pheromones that allowed mass-recruitment in species with

larger colony sizes. The hypothesis of an increase in scale – from small-scale to large-

scale recruitment – has recently been supported by Reeves & Moreau (2019) who

suggest solitary foraging as the ancestral recruitment strategy.

A phylogenetic analysis by Burchill & Moreau (2016), on the other hand, suggested

that early ant species had medium colony sizes, with colonies containing up to several

thousand individuals, which is typically associated with mass-recruitment in extant ants

(Beckers et al., 1989). This suggests that mass-recruitment may have been a more

likely strategy used by early ants. Following this argument, recruitment would have

been lost over time in some lineages as ant species with small colony sizes evolved

(Burchill & Moreau 2016). The antiquity of tandem running has also been questioned

by the fact that this behaviour is found in species that are considered more derived,

such as in *Temnothorax* and *Leptothorax* (Planque et al. 2010). This suggests that

tandem running may be a derived behaviour, at least in some groups.

The main aim of our study was to estimate the ancestral state of recruitment

communication of the Formicidae. While Reeves & Moreau (2019) focused on

recruitment communication during foraging, we also considered whether species use

recruitment during emigrations. We included these cases because numerous ant

species do not communicate during foraging but use recruitment communication in

other ecological situations (Fresneau, 1985; Grüter et al., 2018; Kaur et al., 2017). The

value of communication in foraging depends on the foraging ecology of a species, such

as the kind of food that is exploited and food source distribution (Anna Dornhaus et al.,

2006; l'Anson Price et al., 2019; Sherman & Visscher, 2002). Thus, the strategies used

during foraging reflect foraging ecology and provide an incomplete picture of the

recruitment strategies used by a species. To better understand the evolution of

recruitment communication mechanisms, it is instructive to consider whether a species

uses recruitment communication, irrespective of the type of resource that is exploited.

In addition, we explored whether tandem running was indeed an early recruitment

strategy that preceded group- and mass-recruitment (Hingston, 1929; Wilson, 1959).

Material & Methods

Literature research for recruitment strategies

Data were collected on the recruitment strategies used by extant ant species *via* an

extensive search of the published scientific literature (from October 2019 to March

2020). For many ant species, information about recruitment was collected from reviews

or articles about recruitment (Beckers et al., 1989; Jaffe, 1984; Silvestre et al., 1999).

Furthermore, we searched in Google Scholar using the search terms (ant species or

genus in combination with "recruit", "forag", "prey", "individual", "solitary", "tandem",

"group", "trail", "pheromone"). We included species-level information when the

recruitment strategy was described based on observations or collected in controlled experiments. Data were coded as discrete character traits. Each species was allocated to one of four different recruitment strategies, similar to Jaffé (1984), Beckers et al. (1989) and Lanan (2014): no recruitment, tandem running, group-recruitment and mass-recruitment (Table 1). Reeves & Moreau (2019) found that the different classification systems led to very similar outcomes in their ancestral state reconstruction.

Table 1. Recruitment classifications and definitions, largely based on Jaffé (1984).

Recruitment strategy	Definition
Solitary/individual	No recruitment, no information transfer between nestmates
Tandem running	A single ant (scout) attracts a single nestmate using antennal contact and then physically leads a nestmate to the goal. Physical contact is often maintained between scout and nestmate, chemical signals may be used
Group-recruitment	A scout recruits "up to thirty nestmates" and leads them to the goal. Chemical signals are often used for short-distance attraction but physical contact between scout and the group is also used
Mass-recruitment	Groups are guided via chemical trails alone. Large numbers of ants can be recruited by a small number of recruiters. Chemical trails are laid on the substrate.

The literature search highlighted that there is a relative scarcity of detailed information about foraging strategies in the ant literature (see also Reeves & Moreau 2019). There are numerous examples of studies that mention recruitment of nestmates, but without providing details of the strategy and the context when this was observed. These studies were not included in our analysis. If possible, we also collected and compared the recruitment data to Reeves and Moreau (2019), who collected data on ant foraging strategies. In some cases, we were unable to recover the information from the cited primary literature (e.g. *Buniapone amblyops*, *Mayaponera constricta* or *Megaponera*

analis). *Tetramorium caespitum* uses both group and mass-recruitment (Collignon &

Detrain, 2010). We considered this to be a mass-recruiting species for our analysis.

Additionally, we used a subset that included only ant species that perform tandem runs

and analysed if recruitment to nest sites or food sources was more likely to be the

ancestral state.

Phylogenetic comparative methods

We modified the phylogenetic tree of Branstetter et al. (2017), which contains ~1000

ant species and is a phylogram based on molecular data. In our literature research we

found information for 161 species, 82 genera and 11 sub-families (Table S1) that were

also present in the phylogenetic tree. Overall, the species included in our study

represent 25% of genera and 65% of sub-families. Species with no recruitment data

available were removed from the dataset with the *drop.tip* function in the R package

"ape" (Paradis & Schliep, 2019).

We performed marginal ancestral state reconstructions (ASR) for the dataset, using

the functions *fitMk* and *ace* from the R package "phytools" (Revell, 2012) and "ape"

(Paradis & Schliep, 2019) to estimate the transition rates and the ancestral states for

our tested character using a maximum likelihood (ML) approach. The *fitMk* function

assumes that the probability to change from one state to another depends only on the

current state and not on the state that has come before. Furthermore, every character

state is equally likely to change to one of the other states. The *ace* function utilizes

marginal reconstruction and returns the marginal ancestral state likelihood of all nodes

within a phylogeny.

Additionally, we performed a stochastic character mapping (SCM) by using

make.simmap from the R package "phytools". For the stochastic reconstructions of

character states we used an MCMC approach, to explore the posterior probabilities of

all nodes and provided the number of changes between the character states (1000

simulations performed).

Three commonly used transition rate models were analysed for the ancestral state

reconstructions: "equal rates" (ER), "symmetrical rates" (SYM) and "all rates different"

(ARD) with names referring to transition rates between each state. We used the Akaike

information criterion (AIC) values corrected for small sample sizes (AICc values) for

the three transition rates. We calculated the AIC-weights which standardize the AIC

scores of the fitted models and measured the relative weight of evidence for the three

models used in our data (Harmon, 2019). We visualized the results by mapping the

ancestral state on the phylogeny with the function *plotTree*.

Results

Evolution of recruitment strategy

The ancestral state reconstruction results were mapped to our phylogeny (Fig. 1). The

log-likelihood values, AIC values, AICc values and the number of free parameters per

model are presented in Table 2. We compared the AIC and AICc values, which

revealed that simple ER models were inferior and, thus, were rejected. A "symmetric

model" and "all-rates-different model" best explained our transition between

recruitment states. Thus, both transition models were used to analyse the ancestral

state of recruitment strategies in ants.

A

PONEROIDS

FORMICOIDS

strategy

- 🟡 group
- 🔵 mass
- 🩵 solitary
- 🔴 tandem

B

PONEROIDS

FORMICOIDS

strategy

- group
- mass
- solitary
- tandem

213 **Figure 1**. Ant phylogeny including recruitment strategy and (**A**) marginal ancestral
214 state reconstruction or (**B**) stochastic character mapping. Nodes provide estimates
215 based on Markov chain models. The phylogeny is based on Branstetter et al. (2017).
216

Table 2. Results of the transition rate models. Log likelihoods, Akaike information criterion values, number of free parameters and Akaike-weights are shown.

Model	LogL	AIC	AICc	free parameters	AICcW
ER	-170.0	336.9	342.0	1	0.03
SYM	-161.4	331.6	335.4	6	0.45
ARD	-155.7	331.4	337.6	12	0.52

Marginal ancestral state reconstruction analyses and stochastic character mapping

both suggest that mass-recruitment is the most probable strategy at the root of the

phylogeny (71.2% and 60.2%, respectively) (Table 3). Also, internal nodes (lineage

splitting events) were dominated by high probabilities for the mass-recruitment

category. Mass-recruitment was the most likely ancestral state in both the Poneroids

and the Formicoids (Figure 1). The stochastic character mapping revealed that there

were an estimated 81.3 changes between recruitment strategies (Table 4). The most

common transitions were from mass-recruitment to solitary/individual behaviour

(33.7%) or to group-recruitment (18.5%) (Figure 2). Furthermore, there were

transitions from no recruitment to tandem running (14.8%) or mass-recruitment

(12.7%). Tandem running evolved several times independently in the subfamilies

Ponerinae, Pseudomyrmecinae, Formicinae and Myrmecinae. Furthermore, it seems

that recruitment was lost at least once in all subfamilies, except in the Dorylinae (army

ants). Similarly, we found group-recruitment in nearly all included sub-families, except

in the Paraponerinae. The Myrmeciinae were the only group without species that

perform mass-recruitment.

Table 3. Ancestral character estimation using marginal ancestral state reconstruction and stochastic character mapping. Values represent likelihoods of recruitment strategies at the root.

	Character states	Scaled root likelihood (fitMk)	Scaled root likelihood (ace)	Stochastic charac mapping (make.simmap)
SYM	No recruitment	0.15	0.16	0.21
	Tandem running	0.01	0.01	0.10
	Group-recruitment	0.02	0.02	0.07
	Mass-recruitment	0.81	0.81	0.61
ARD	No recruitment	0.05	0.06	0.21
	Tandem running	0.01	0.01	0.09
	Group-recruitment	0.50	0.48	0.12
	Mass-recruitment	0.44	0.45	0.59

Table 4. Changes from stochastic character mapping. GR = group-recruitment, MS = mass-recruitment, NR = no recruitment, TR = tandem running

	SYM		ARD	
Total changes	75.783		104.472	
Type	Number	Percentage	Number	Percentage
GR → MS	3.44	4.5%	32.013	30.6%
GR → NR	0.595	0.7%	0	0%
GR → TR	1.224	1.6%	0	0%
MS → GR	14.902	19.7%	23.572	22.6%
MS → NR	26.57	35.1%	27.514	26.3%
MS → TR	0.489	0.6%	0.952	0.9%
NR → GR	1.412	1.9%	7.409	7.1%
NR → MS	10.061	13.3%	0	0%
NR → TR	9.8	12.9%	9.572	9.2%

TR → GR	2.626	3.4%	3.44	3.3%
TR → MS	0.117	0.2%	0	0%
TR → NR	4.547	6.0%	0	0%

Figure 2. Transitions between states from stochastic character mapping for the methods SYM and ARD. The thickness of the arrow reflects the relative commonness of a transition. GR = group-recruitment, MS = mass-recruitment, NR = no recruitment, TR = tandem running

Evolution of tandem running

Twenty-one species included in our study perform tandem runs. For some species it is not known if they recruit to food sources and during nest relocations. For other species it is known that they perform tandem runs to new nest sites, but forage solitarily for food sources (*Neoponera, Diacamma* or *Paltothyreus*) (Table S1).

After mapping the recruitment strategy onto the phylogenetic tree (Figure 3), we found that all species that can perform tandem runs do so during colony emigrations to new nest sites. Several species (43%) that use recruitment via tandem running do not do so during foraging.

Figure 3. Ancestral state reconstruction for ant species that perform tandem running to
either new nest sites, food sources or both. Nodes provide estimates based on Markov
chain models.

Discussion

Our ancestral state analyses indicate that mass-recruitment or group-recruitment were

likely used for recruitment by the last common ancestor of present-day ants. During

the course of their evolutionary history, all included subfamilies show switches from

mass-recruitment to other recruitment strategies. Strikingly, most transitions occurred

from mass recruitment to group or no recruitment. The repeated loss of communication

seems puzzling, but probably coincided with the emergence of lineages with small

colony sizes (e.g. in Amblyoponinae, Myrmeciinae, Ponerinae, Pseudomyrmecinae)

(Burchill & Moreau 2016), where communication is less beneficial (Beckers et al.

1989). Transitions between mass- and group-recruitment were also more frequent than

between tandem running and no recruitment. Tandem running evolved independently

in 4 of 11 subfamilies. Furthermore, there were several transitions from no recruitment

to tandem running or mass-recruitment. These findings highlight that recruitment

strategies are an evolutionarily flexible and reversible trait.

Many lineages appear to have lost the ability to recruit nestmates. The putative

loss of chemical mass-recruitment seems puzzling at first, but could be explained by

the constraints of living in a small colony (Beckers et al., 1989; Dornhaus et al., 2012)

and/or a switch to a diet or foraging strategy that does not require pheromone trails.

Smaller colonies tend to exploit resources solitarily or they use tandem running. One

reason could be that smaller colonies do not have sufficiently large colonies to maintain

pheromone trails (Beckers et al., 1989; Beekman et al., 2001; Planqué et al., 2010). A

recent study has suggested that the ancestral Formicidae had medium colony sizes

containing up to several thousand individuals (Burchill & Moreau, 2016). These

findings in combination with the findings that medium sized colonies often use group

or mass-recruitment (Beckers et al. 1989) are consistent with our results that the best-

supported ancestral recruitment strategy was mass- or group-recruitment.

Another reason for the loss of mass-recruitment could be that recruitment by

pheromone trails can have disadvantages in changing foraging landscapes.

Pheromone trails can persist for relatively long periods (up to several hours), which

can make it difficult for the colony to re-allocate workers to a newly available higher-

quality resource due to the strong positive feedback created by a pheromone trail

(Beckers et al., 1989; Grüter et al. 2012; Czaczkes et al., 2015; l'Anson Price et al.,

2016). This makes recruitment less flexible and colonies are more likely to miss out on

new food sources when the environment changes.

Reeves & Moreau (2019) found evidence that solitary foraging, rather than mass

recruitment, represented the original state in terms of recruitment strategies. Our and

their results are not necessarily contradictory. In our study, we considered whether a

species uses recruitment communication during colony emigrations and/or during

foraging. It is well known that foraging strategies in ants strongly depend on the

foraging ecology (Davidson, 1977; Dejean et al., 2012; Dornhaus et al., 2006;

Hölldobler & Wilson, 1990; Lanan, 2014). For example, in many ant species, foragers

follow a solitary foraging strategy when collecting insect prey, but they use recruitment

communication when the colony emigrates to a new nest-site (Lanan, 2014). In other

words, these species possess the ability to recruit, but foragers do not perform

recruitment because this would not be an adaptive strategy given their foraging

ecology.

The observation that numerous species recruit to new nest sites, but do not use

recruitment communication during foraging (e.g. *Neoponera* or *Diacamma* species,

Fresneau, 1985; Grüter et al., 2018) raises the question if recruitment communication

evolved first to help colonies during emigrations rather than to communicate the

location of food sources, as has also been suggested in the case of the honeybee

waggle dance (Beekman et al. 2008; I'Anson Price & Grüter, 2015). This seems

plausible given that during nest relocations of cavity nesting species, nest locations

have to be communicated very precisely. If the old nest is damaged or destroyed, a

fast and precise relocation is critical (Dornhaus et al., 2004; Franks et al., 2003). During

foraging, on the other hand, communication might often be less important. Especially

when food sources are abundant and evenly distributed, communication might not

provide benefits or even be disadvantageous due to time costs (Dechaume-

Moncharmont et al., 2005; Dornhaus et al., 2006; Goy et al., 2021; I'Anson Price et al.,

2019). The hypothesis that recruitment evolved first in colony emigrations and was co-

opted by some species in a foraging context is also supported by our results suggesting

group- or mass-recruitment as the ancestral state and those of Reeves & Moreau

(2019) who suggested solitary foraging as the ancestral condition.

It has been suggested that tandem running is a "primitive" recruitment strategy

(*i.e.* ancestral) (Hingston 1929; Hölldobler et al., 1974; Schultheiss et al., 2015; Wilson,

1959). Our results do not support this view. We found that tandem running evolved

repeatedly and independently in the subfamilies Ponerinae, Pseudomyrmecinae,

Formicinae and Myrmecinae. Transitions to tandem running occurred most often from

no recruitment and, more rarely, from mass or group-recruitment. One benefit of

tandem running is that it allows small colonies to defend resources against competitors

when competition for nest sites or food sources is intense(Glaser et al., 2021).

In summary, our results suggest that mass or group-recruitment were the most

likely recruitment strategies used by the last common ancestor of present-day ants.

There were repeated, independent transitions to different strategies, such as tandem

running or no recruitment, but also transitions back to group or mass-recruitment. It

should be noted that our analysis is restricted to a small proportion of ant species and

we currently lack information about recruitment behaviours for the majority of species

(see also Reeves & Moreau 2019). We echo the call of Reeves & Moreau (2019) to

pay more attention to ant behaviour and ecology as this will allow us to better

understand the links between different traits of ant behaviour, ecology and natural

history.

References

Basari, N., & Laird-Hopkins, B. (2014). Trail laying during tandem-running recruitment

in the ant *Temnothorax albipennis*. *Naturwissenschaften*, *101*(7), 549–556.

https://doi.org/10.1007/s00114-014-1191-1

Beckers, R., Goss, S., Deneubourg, J. L., & Pasteels, J. M. (1989). Colony size,

communication and ant foraging strategy. *Psyche: A Journal of Entomology*, *96*(3–

4), 239–256. https://doi.org/10.1155/1989/94279

Beekman, M., Sumpter, D. J. T., & Ratnieks, F. L. W. (2001). Phase transition between

disordered and ordered foraging in Pharaoh ' s ants. *PNAS*, *98*(17), 9703–9706.

Burchill, A. T., & Moreau, C. S. (2016). Colony size evolution in ants:

macroevolutionary   trends.   *Insectes   Sociaux*,   *63*(2),   291–298.

https://doi.org/10.1007/s00040-016-0465-3

Collignon, B., & Detrain, C. (2010). Distributed leadership and adaptive decision-

making in the ant *Tetramorium caespitum*. *Proceedings of the Royal Society B:*

Biological   Sciences,   277(1685),   1267–1273.

https://doi.org/10.1098/rspb.2009.1976

Czaczkes, T. J., Grüter, C., & Ratnieks, F. L. W. (2015). Trail pheromones: An

integrative view of their role in social insect colony organization. *Annual Review of*

Entomology,   *60*(1),   581–599.   https://doi.org/10.1146/annurev-ento-010814-

020627

Czaczkes, T. J., Nouvellet, P., & Ratnieks, F. L. W. (2011). Cooperative food transport

in the Neotropical ant, Pheidole oxyops. *Insectes Sociaux*, *58*(2), 153–161.

https://doi.org/10.1007/s00040-010-0130-1

Czaczkes, T. J., & Ratnieks, F. L. W. (2012). Pheromone trails in the Brazilian ant

Pheidole oxyops: extreme properties and dual recruitment action. *Behavioral*

Ecology and Sociobiology, *66*(8). https://doi.org/10.1007/s00265-012-1367-7

Davidson, D. W. (1977). Foraging ecology and community organization in desert seed-

eating ants. *Ecology*, *58*(4), 725–737.

Dechaume-Moncharmont, F. X., Dornhaus, A., Houston, A. I., McNamara, J. M.,

Collins, E. J., & Franks, N. R. (2005). The hidden cost of information in collective

foraging. *Proceedings of the Royal Society B: Biological Sciences*, *272*(1573),

1689–1695. https://doi.org/10.1098/rspb.2005.3137

Dejean, A., Delabie, J. H. C., Corbara, B., Azémar, F., Groc, S., Orivel, J., & Leponce,

M. (2012). The ecology and feeding habits of the arboreal trap-jawed ant Daceton

armigerum. *PLoS ONE*, *7*(6). https://doi.org/10.1371/journal.pone.0037683

Detrain, C., & Deneubourg, J.-L. (2008). Collective decision-making and foraging

patterns in ants and honeybees. *Advances in Insect Physiology*, *35*, 123–173.

Dornhaus, A., Franks, N. R., Hawkins, R. M., & Shere, H. N. S. (2004). Ants move to

improve: colonies of *Leptothorax albipennis* emigrate whenever they find a

superior nest site. *Animal Behaviour*, *67*(5), 959–963.

https://doi.org/10.1016/j.anbehav.2003.09.004

Dornhaus, Anna, Klügl, F., Oechslein, C., Puppe, F., & Chittka, L. (2006). Benefits of

recruitment in honey bees: effects of ecology and colony size in an individual-

based model. *Behavioral Ecology*, *17*(3), 336–344.

https://doi.org/10.1093/beheco/arj036

Dornhaus, Anna, Powell, S., & Bengston, S. (2012). Group size and its effects on

collective organization. *Annual Review of Entomology*, *57*, 123–141.

https://doi.org/10.1146/annurev-ento-120710-100604

Franklin, E. L. (2014). The journey of tandem running: the twists, turns and what we

have learned. *Insectes Sociaux*, *61*(1), 1–8. https://doi.org/10.1007/s00040-013-

0325-3

Franks, N. R., Dornhaus, A., Fitzsimmons, J. P., & Stevens, M. (2003). Speed versus

accuracy in collective decision making. *Proceedings of the Royal Society B:*
Biological Sciences, *270*(1532), 2457–2463.
https://doi.org/10.1098/rspb.2003.2527

Fresneau, D. (1985). Individual foraging and path fidelity in a ponerine ant. *Insectes Sociaux*, *32*(2), 109–116. https://doi.org/10.1007/BF02224226

Glaser, S. M., & Grüter, C. (2018). Ants (*Temnothorax nylanderi*) adjust tandem running when food source distance exposes them to greater risks. *Behavioral Ecology and Sociobiology*, *72*(3), 1–8. https://doi.org/10.1007/s00265-018-2453-2

Glaser, S. M., Feitosa, R.M., Koch, A., Goss, N., Nascimento, F.S. & Grüter, C. (2021). Tandem communication improves ant foraging success in a highly competitive tropical environment. *Insectes Sociaux*, *68*, 161–172. https://doi.org/10.1007/s00040-021-00810-y

Goy, N., Glaser, S. M., & Grüter, C. (2021). The adaptive value of tandem communication in ants: insights from an agent-based model. *Journal of Theoretical Biology*, *526*, 110762.

Grooters, H. J. (1987). Influences of queen piping and worker behaviour on the timing of emergence of honey bee queens. *Insectes Sociaux*, *34*(3), 181–193. https://doi.org/10.1007/BF02224083

Grüter, C., & Leadbeater, E. (2014). Insights from insects about adaptive social information use. *Trends Ecol. Evol.*, *29*(3), 177–184. https://doi.org/10.1016/j.tree.2014.01.004

Grüter, C., Wüst, M., Cipriano, A. P., & Nascimento, F. S. (2018). Tandem recruitment and foraging in the ponerine ant *Pachycondyla harpax* (Fabricius). *Neotropical*

Entomology, *47*(6), 742–749. https://doi.org/10.5194/hess-16-3075-2012

Grüter, C, Czaczkes, T. J., & Ratnieks, F. L. W. (2011). Decision making in ant foragers
(Lasius niger) facing conflicting private and social information. *Behavioral Ecology*
and Sociobiology, *65*(2), 141–148. https://doi.org/10.1007/s00265-010-1020-2

Grüter, C., Schürch, R., Czaczkes, T. J., Taylor, K., Durance, T., Jones, S.M. &
Ratnieks, F. L. W. (2012). Negative feedback enables fast and flexible collective
decision-making   in   ants.   *PLoS   One,   7(9)*,   e44501.
https://doi.org/10.1371/journal.pone.0044501

Grüter, Christoph. (2020). *Stingless Bees: Their Behaviour, Ecology and Evolution*.
Springer International Publishing.

Harmon, L. J. (2019). *Phylogenetic Comparative Methods*.

Hingston, R. (1929). *Instinct and intelligence*. Macmillan.

Hölldobler, B. (1971). Recruitment behavior in Camponotus socius (Hym. Formicidae).
Zeitschrift   Für   Vergleichende   Physiologie,   *75*(2),   123–142.
https://doi.org/10.1007/BF00335259

Hölldobler, B. (1999). Multimodal signals in ant communication. *Journal of*
Comparative Physiology A, *184*(2), 129–141.

Hölldobler, B., Möglich, M., & Maschwitz, U. (1974). Communication by tandem
running in the ant Camponotus sericeus. *Journal of Comparative Physiology*,
90(2), 105–127. https://doi.org/10.1007/BF00694481

Hölldobler, B., & Wilson, E. O. (1990). *The Ants*. Harvard University Press, Springer-
Verlag Berlin Heidelberg.

Hölldobler, B., & Wilson, E. O. (2009). *The Superorganism*. W. W. Norton & Company.

l'Anson Price, R., Dulex, N., Vial, N., Vincent, C., & Grüter, C. (2019). Honeybees

forage more successfully without the "dance language" in challenging

environments. *Science Advances*, *5*(2). https://doi.org/10.1126/sciadv.aat0450

l'Anson Price, R., Grüter, C., Hughes, W. O. H., & Evison, S. E. F. (2016). Symmetry

breaking in mass-recruiting ants: extent of foraging biases depends on resource

quality. *Behavioral Ecology and Sociobiology*, *70*(11), 1813–1820.

https://doi.org/10.1007/s00265-016-2187-y

Jaffe, K. (1984). Negentropy and the evolution of chemical recruitment in ants

(Hymenoptera: Formicidae). *Journal of Theoretical Biology*, *106*(4), 587–604.

https://doi.org/10.1016/0022-5193(84)90009-2

Kaur, R., Joseph, J., Anoop, K., & Sumana, A. (2017). Characterization of recruitment

through tandem running in an Indian queenless ant *Diacamma indicum*. *Royal*

Society Open Science, *4*(1), 160476. https://doi.org/10.1098/rsos.160476

Lach, L. (2005). Interference and exploitation competition of three nectar-thieving

invasive ant species. *Insectes Sociaux*, *52*(3), 257–262.

https://doi.org/10.1007/s00040-005-0807-z

Lanan, M. C. (2014). Spatiotemporal resource distribution and foraging strategies of

ants (Hymenoptera: Formicidae). *Myrmecological News*, *20*, 53–70.

https://doi.org/10.1016/j.biotechadv.2011.08.021.

Leadbeater, E., & Chittka, L. (2007). Social learning in insects - from miniature brains

to consensus building. *Current Biology*, *17*(16), 703–713.

https://doi.org/10.1016/j.cub.2007.06.012

Liefke, C., Hölldobler, B., & Maschwitz, U. (2001). Recruitment behavior in the ant

genus *Polyrhachis* (Hymenoptera, Formicidae). *Journal of Insect Behavior*, *14*(5),

637–657.

Maschwitz, U., & Schönegge, P. (1983). Forage communication, nest moving

recruitment, and prey specialization in the oriental ponerine *Leptogenys chinensis*.

Oecologia, *57*, 175–182.

Möglich, M., Maschwitz, U., & Hölldobler, B. (1974). Tandem calling: a new kind of

signal in ant communication. *Science*, *186*(4168), 1046–1047.

Moreau, C. S., Bell, C. D., Vila, R., Archibald, S. B., & Pierce, N. E. (2006). Phylogeny

of the ants: Diversification in the age of angiosperms. *Science*, *312*(5770), 101–

104. https://doi.org/10.1126/science.1124891

Paradis, E., & Schliep, K. (2019). ape 5.0: an environment for modern phylogenetics

and evolutionary analyses in R. *Bioinformatics*, *35*(3), 526–528.

https://doi.org/10.1093/bioinformatics/bty633

Planqué, R., van den Berg, J. B., & Franks, N. R. (2010). Recruitment strategies and

colony size in ants. *PLoS ONE*, *5*(8), e11664.

https://doi.org/10.1371/journal.pone.0011664

Reeves, D. D., & Moreau, C. S. (2019). The evolution of foraging behavior in ants

(Hymenoptera: Formicidae). *Arthropod Systematics and Phylogeny*, *77*(2), 351–

363. https://doi.org/10.26049/ASP77-2-2019-10

Revell, L. J. (2012). phytools: an R package for phylogenetic comparative biology (and

other things). *Methods in Ecology and Evolution*, *3*(2), 217–223.

https://doi.org/10.1111/j.2041-210X.2011.00169.x

Schultheiss, P., Raderschall, C. A., & Narendra, A. (2015). Follower ants in a tandem

pair are not always naïve. *Scientific Reports*, *5*, 10747.

https://doi.org/10.1038/srep10747

Sherman, G., & Visscher, P. K. (2002). Honeybee colonies achieve fitness through

dancing. *Nature*, *419*(6910), 920–922. https://doi.org/10.1038/nature01127

Silvestre, R., Brandão, C. R. F., & Paiva, R. V. S. (1999). Ant guilds in the Brazilian

"Cerrados." *Ant   Guilds   in   the   Brazilian   "Cerrados."*

https://doi.org/10.13140/RG.2.2.22720.74246

Traniello, J. F. A. (1989). Foraging strategies of ants. *Annual Review of Entomology*,

34(1), 191–210. https://doi.org/10.1146/annurev.ento.34.1.191

Villet, M. H. (1990). Social organization of *Platythyrea lamellosa* (Roger)

(Hymenoptera: Formicidae): II. Division of labour. *South African Journal of*

Zoology, *25*(4), 254–259. https://doi.org/10.1080/02541858.1990.11448222

von Frisch, K. (1967). *The dance language and orientation of bees*. Harvard University

Press.

Ward, P. S. (2014). The phylogeny and evolution of ants. *Annual Review of Ecology,*

Evolution, and Systematics, *45*, 23–43. https://doi.org/10.1146/annurev-ecolsys-

120213-091824

Wilson, E. O. (1959). Communication by tandem running in the ant genus

Cardiocondyla. *Psyche*, *66*(3), 29–34.

Wilson, E. O. (1971). *The insect societies*. Belknap Press of Harvard University Press.