Disentangling the historical routes to community assembly in the global epicentre

of biodiversity

Running title: Sympatry and elevational divergence

Abstract

Aim

The coexistence and turnover of species along elevation gradients makes tropical

mountains hotspots of biodiversity. However, understanding the historical processes

through which species arising in geographic isolation (i.e. allopatry) assemble along the

same mountain slope (i.e. sympatry) remains a major challenge. Multiple models have

been proposed including, (1) the sorting of already elevationally divergent species, (2)

the displacement of elevation upon secondary contact, potentially followed by

convergence, or (3) elevational conservatism, in which ancestral elevational ranges are

retained. However, the relative contribution of these processes to generating patterns of

elevational overlap and turnover is unknown.

Location

Tropical mountains of Central- and South-America.

Time period

The last 12 myr.

Major taxa studied

Birds.

Methods

We collate a dataset of 165 avian sister pairs containing estimates of phylogenetic age,

geographical and regional elevational range overlap. We develop a framework based on

continuous-time Markov models to infer the relative frequency of different historical

pathways in explaining present day overlap and turnover of sympatric species along

elevational gradients.

Results

We show that the turnover of closely related bird species across elevation can

predominantly be explained by displacement of elevation ranges upon contact (81%)

rather than elevational divergence in allopatry (19%). In contrast, overlap along

elevation gradients is primarily (88%) explained by conservatism of elevational ranges

rather than displacement followed by elevational expansion (12%).

Main conclusions

Bird communities across tropical elevation gradients are assembled through a mix of

processes, including the sorting, displacement and conservatism of species elevation

ranges. The dominant role of conservatism in explaining co-occurrence of species on

mountain slopes rejects more complex scenarios requiring displacement followed by

subsequent expansion. The ability of closely related species to coexist without

elevational divergence provides a direct and thus faster pathway to sympatry and may

help explain the exceptional species richness of tropical mountains.

Keywords: biotic interactions, character displacement, community assembly,

elevational gradient, niche conservatism, species sorting, sympatry

1 Introduction

Explaining the combination and diversity of species that co-occur within ecological communities remains a major challenge. In part, this is because the patterns of spatial overlap observed at the present depend not only on current ecological interactions between species, but also on historical processes operating over much longer timescales and that are beyond the reach of direct observation or experimental manipulation (Weber *et al.*, 2017). These historical processes include speciation, niche evolution, dispersal, and range expansions (Mittelbach & Schemske, 2015), with differences in the dynamics of these processes, and the ecological factors controlling them, underlying different theoretical models for how communities are assembled.

One of the main models for how communities assemble is based on the idea that new species arising in spatial isolation (i.e. allopatry) diverge in their ecological niche (including habitat), and only those that happen to diverge sufficiently to minimise competition are able to co-occur when they come back into secondary contact (Pfennig & Pfennig, 2010; Stuart & Losos, 2013). According to this model, community assembly largely involves the 'sorting' of pre-existing variation arising due to geographically variable selection pressures. An alternative to the ecological sorting model, proposes that niche differences between co-occurring species arise upon secondary contact, with competition between species driving divergent selection and a displacement of their ecological niches (Brown & Wilson, 1956). As with ecological sorting, this 'ecological displacement' model (termed 'character displacement' when considering heritable traits) assumes that niche similarity limits co-occurrence, but differs in its predictions of when and why niche differences evolve. A final possibility, is that species may re-assemble into communities whilst retaining their ancestral niche. This 'niche conservatism' model (Cadena *et al.*, 2011) is expected to predominate if ecological niche overlap is not limiting, either because of other constraints on co-occurrence (e.g. dispersal limitation) or because species have diverged along alternative niche dimensions (Pigot *et al.*, 2018).

A classic system and spatial parallel for studying these historical processes concerns the distribution of species along tropical mountain slopes. Tropical mountains are renowned for their exceptional diversity. For instance, in the tropical Andes over 800 species of birds can occur on a single mountain slope (Walker *et al.*, 2006). Since most speciation events involve the geographic isolation of populations on different mountains

(i.e. allopatry)(Price, 2008; Cadena & Céspedes, 2020; Linck *et al.*, 2020), the main

problem is to understand how these species subsequently assemble on the same

mountain slope (i.e. sympatry). In this system, sympatry may involve species co-

occurring at the same elevation, what is often referred to as syntopy. Alternatively,

species may occupy different elevation ranges. Indeed, many species have very narrow

elevation distributions (e.g. a few hundred metres in vertical distance) and replace one

another across elevational gradients (Diamond, 1973; Terborgh & Weske, 1975).

Previous studies have variously provided evidence that conservatism (Cadena *et al.*,

2011), sorting (Cadena, 2007), and displacement (Freeman, 2015; Freeman *et al.*, 2022)

of elevational ranges may individually be involved in shaping these patterns of co-

occurrence and turnover along mountain slopes, but progress in disentangling their

relative contributions has been limited.

One problem is that different assembly models can lead to the same present day

pattern. In particular, according to the elevational conservatism (EC) model, species

living on the same mountain slope have overlapping elevation ranges because they

retain the adaptations to specific environmental conditions inherited from their

ancestor (Cadena *et al.*, 2011) (Figure 1d). However, while some studies have argued

that elevational conservatism is a pervasive process (Linck *et al.*, 2021), the same

pattern could also arise under the elevational displacement (ED) model. Under ED,

competition on secondary contact first forces species to occupy different elevations, but

as species diverge along alternative niches dimensions (e.g., resource or microhabitat

use) they may subsequently expand their elevation ranges and overlap along the

gradient (Figure 1c). This ED model has long been regarded as the dominant process

explaining the build-up of species within elevation zones on tropical mountains

(Diamond, 1973; Freeman, 2015).

A similar challenge exists when trying to discriminate elevational displacement from

elevational sorting (ES). Both of these models predict that species living on the same

mountain slope should have non-overlapping elevation ranges, at least prior to any

subsequent expansion in elevation range. Thus, discriminating between these pathways

depends critically on inferences about whether elevational divergence precedes (i.e. ES)

or coincides with (i.e. ED) the attainment of secondary contact (Figure 1b, c). Studies

attempting to address this issue have variously used phylogenetic information to

compare the relative age of sister pairs that are elevationally divergent and those that

are sympatric, or have compared elevation divergence between sister species in zones of

sympatry to places where they remain allopatric (Freeman, 2015; Cadena & Céspedes,

2020; Figure S1). However, while such tests can provide evidence that elevational

displacement does or does not occur, critically they do not reveal how frequently this

process occurs and thus its relative contribution to the build-up of sympatric diversity

and elevational turnover (Anderson & Weir, 2021). Progress in understanding

community assembly, thus requires new approaches that can reliably infer the relative

frequencies of different historical processes underlying current patterns of species

distributions.

Here, we investigate the relative frequency of elevational displacement, sorting,

and conservatism and how they contribute to the turnover and co-occurrence of

passerine birds across elevation gradients in the Neotropics. We first compiled a new

dataset consisting of the divergence time, geographical and regional elevational

distributions of n = 165 sister species. We then apply a novel modelling approach that

builds on previous studies examining the transition to sympatry using continuous-time

Markov models (Pigot & Tobias, 2013; Pigot *et al.*, 2018), but extend this framework by

considering the multiple pathways through which species can attain sympatry via the

evolution of elevational ranges. Focussing on avian sister species, where we can

confidently assume that speciation involves the geographic isolation of populations

(Phillimore *et al.*, 2008; Price, 2008), allows us to use information on their current age

and state (e.g. allopatric or sympatric, and overlapping in elevation or not) to model the

historical pathways through which patterns of geographic and elevational range overlap

arise. Using this approach, we address three main questions. First, what is the relative

importance of elevational sorting, displacement and conservatism in the build-up of

sympatry among Neotropical montane sister species? Second, what is the relative

importance of elevational sorting and displacement in generating turnover between

sister species across elevation gradients? Third, how does co-occurrence between

species at the same elevation on the same mountain slope arise? We address these

questions by modelling the historical pathways through which current distributions

have arisen and conduct sensitivity tests and simulations to identify potential biases

arising from limitations in the data or methods. Our results show that it is possible to

reliably recover the historical pathways through which present day patterns of

sympatry and co-occurrence have arisen, and we provide estimates of the relative

frequency of these different pathways in explaining the assembly of Neotropical

montane bird communities.

2 Material and methods

2.1 Elevational and geographic data

We compiled a dataset consisting of avian sister species occurring in the mountain

ranges of the Neotropics. Sister species and their respective divergence times were

extracted from two recently published phylogenies covering respectively >95% and

>98% of the oscines and suboscines from North and South America (Barker *et al.*, 2015;

Harvey *et al.*, 2020). We only retained sister pairs that met the following conditions: (1)

At least one of the species occupies montane habitat, defined as areas >500m above sea-

level (Freeman, 2015). (2) Both species occupy predominantly humid environments

such that species pairs could realistically live on the same mountain slope and

macrohabitat (e.g. we removed species pairs if one sister occupied predominantly arid

habitats and not humid forest) as assessed with information from field guides. (3) Both

species live in the Neotropics, which included the tropical mountain ranges of Central

and South America. Species that had ranges stretching into North America were

included in the dataset if they were primarily Neotropical. The final dataset consisted of

n = 165 sister pairs.

We scored whether each sister pair was allopatric or sympatric and whether they

had overlapping or non-overlapping elevational distributions using geographical and

elevational range data gathered from a number of sources, including regional field

guides and surveys (Hilty & Brown, 1986; Ridgely & Greenfield, 2001; Hilty, 2003;

Walker *et al.*, 2006; Vallely & Dyer, 2018). We supplemented this with *Birds of the World*

(BW) (Billerman *et al.*, 2020) that typically reflects information found in regional field

guides but incorporates more recent taxonomic changes. We note that, although not

primary scientific literature, field guides offer a wealth of expert knowledge on the

natural history of birds, and are frequently used in studies of bird elevation distributions

(e.g. Quintero & Jetz, 2018).

For each species, we used these sources to record elevational ranges on a region-

by-region basis, usually corresponding to countries, but in greater detail where available

(e.g. Choco vs. Amazonian slope in Colombia). This is important, not only to detect

differences in elevation range overlap between zones of allopatry versus sympatry, but

also because species can vary in their elevation limits across their geographic range

more generally due to differences in climate, mountain height and biotic interactions

(Janzen, 1967; Londoño *et al.*, 2017; Pujolar *et al.*, 2022). In some cases, elevation range

limits in a particular region were not available, but field guides would report that

elevational ranges of the sister species were non-overlapping. We used this expert

information when assigning species as overlapping or not in that region. Observations of

single specimens at single elevations were not incorporated in our dataset.

2.2 Assigning geographic and elevational range overlap

Although sister species can overlap in their geographic and elevational range to varying

degrees, our modelling approach requires treating overlap as a discrete state. We

defined species as having overlapping elevation ranges if overlap (range in vertical

height, metres) was ≥20% of the species with the narrower elevation range. This

threshold was chosen to avoid classing species which only meet marginally along

narrow contact zones as overlapping. Sympatry was defined as when sister species were

present on the same mountain slope, regardless of whether the geographic extent of

overlap was widespread (i.e. thousands of kms) or minimal (i.e. a few kms). We did not

use metrics of absolute or proportional geographic range overlap to define sympatry

because this is unsuitable for montane systems where one species can have a very small

geographic range. In practice, defining sympatry was unambiguous as allopatric pairs

typically occurred on different mountain summits or regions separated by obvious

geographical barriers (e.g. valleys).

We assigned species pairs to one of five possible discrete states, defined by the

combination of sympatry/allopatry and elevation overlap/non-overlap (Figure 1a). For

sister species occurring in allopatry, they may have overlapping (state 1) or non-

overlapping (state 3) elevational ranges. Sister species occurring in sympatry may have

non-overlapping elevational ranges (state 4), overlapping elevational ranges (state 5), or

elevational ranges that overlap in allopatry but not sympatry (state 2). Pairs were not

classified in a separate state if they showed elevational divergence in allopatry but not

sympatry (opposite of state 2). Under this scenario, sympatry has been established

irrespective of elevational differentiation, making the distinction of limited relevance to

our analysis (and classified in state 5). Regardless, under the protocol discussed below

merely one pair would be classified in such a state.

We found that for most species, variation among regions in their elevation range

limits was relatively minor (Figure S2). However, we incorporated the regional variation

that does exist, by using the mean pairwise overlap between sister species elevation

ranges among the regions where they occur, doing this for the allopatric and sympatric

parts of their geographic ranges separately. Thus, allopatric sister species are assigned

to state 1 if their average regional elevation overlap is ≥20%. Otherwise, allopatric pairs

are assigned to state 3. If pairs are sympatric, they are assigned to state 2, 4 or 5

according to the following rules. Sister species are assigned to state 2 if they show on

average ≥20% overlap in elevation ranges between allopatric regions, and <20%

overlap between sympatric regions. If sister species show on average <20% overlap

between their respective allopatric ranges compared to the other species' entire range,

as well as an average overlap in sympatry of <20%, they are assigned to state 4. All

remaining pairs, which show on average ≥20% elevation overlap in both sympatry and

allopatry, are assigned to state 5. We refer to this method of using the mean pairwise

regional overlap as the 'average overlap' protocol to distinguish this from alternative

approaches used in our sensitivity analyses.

2.3 Sensitivity analyses

To ensure our results were robust to the way elevation range overlap was defined we

performed a series of sensitivity tests. First, we repeated our analysis using 1 and 40%

thresholds to define sister pairs as overlapping (Figure S3, S4, S5, S6). Second, we

accounted for regional variation in elevation ranges in different ways. For zones of

allopatry and sympatry we used: (i) the maximum upper and minimum lower

elevational range limit of a species across regions to define its elevation range (ii) the

minimum overlap observed among regional elevation ranges and (iii) the maximum

overlap observed among regional elevation ranges. These different protocols which we

refer to as the 'global', 'minimum' and 'maximum' approach, will variously tend to

increase ('global' and 'maximum') or decrease ('minimum') the number of sister pairs

assigned as having overlapping elevation ranges (Figure S3, S4, S5, S6). Third, we

explored different ways of detecting the displacement of elevation ranges in sympatry

(Figure S7).

In our main analysis we assigned sister species to state 2 if elevation ranges

overlap in allopatry but not (i.e. <20%) in sympatry, a pattern expected if competition

causes a displacement of species elevation ranges on secondary contact. However, this is

a rather strict definition because displacement could be substantial but still not

eliminate overlap. For example, sister species elevation ranges could overlap by 90% in

allopatry and 50% in sympatry, a decrease in overlap indicative of competitive

displacement but one that would not be detected by our assignment. To address this we

perform an additional analysis in which species are assigned to state 2 if they show

≥20% reduction in overlap in sympatry compared to allopatry (Figure S7).

2.4 Statistical analysis using Markov models

To infer the relative importance of different community assembly pathways in

explaining patterns of geographic and elevation overlap and turnover, we constructed a

continuous time multi-state Markov model (Figure 1b). In this model, the initial state for

sister species is allopatry with overlapping elevational ranges (state 1), which reflects

the situation expected at the time of their initial divergence (Coyne & Orr, 2004). Sister

pairs then stochastically transition between states, with different transitions

corresponding to the different assembly pathways (Figure 1). Under the EC pathway,

sister species transition directly from state 1 to state 5 (i.e. sympatry and overlapping

elevational ranges; Figure 1d) while under the ED and ES pathways species must pass

through intermediate states. Under the ED pathway, sister species transition from state

1 to state 2 (i.e. retaining overlapping elevation ranges in allopatry but non-overlapping

elevation ranges in sympatry), before then converging or expanding in elevation in

sympatry to reach state 5 (Figure 1c). Under the ES pathway, sister species transition to

state 3 (i.e. non-overlapping elevation ranges whilst remaining in allopatry), then state 4

(i.e. attaining sympatry and retaining non-overlapping elevation ranges) and finally to

state 5 (Figure 1b). Thus, our model allows for the elevation overlap of sister pairs in

sympatry (state 5), which is the final absorbing state, to be attained through any of the

three community assembly pathways.

Within this framework, we considered two additional scenarios. First, in our

model we also allow sister pairs in state 2 to transition to state 4, representing a process

where species that have attained sympatry via elevation range displacement

subsequently diverge in their elevation range in the allopatric part of their distribution

(Figure 1c). Such divergence in allopatry, was hypothesised by Diamond (1973) to be

the result of the spread of adaptations to the compressed elevational range arising in

sympatry, resulting in displaced elevation ranges even where the sister species is not

present. Second, in an additional analysis we relax the assumption that all sister pairs

arise in state 1, by estimating an additional parameter, γ, representing the probability

that species arise in state 3 (Supplementary Information Figure S8, S9). This accounts

for the possibility that newly formed sister species may immediately occur at different

elevations (e.g. due to differences in climate and thus heights of habitat zones between

regions).

Based on the estimated time since divergence and states of sister pairs at present

we used maximum likelihood (ML) (Jackson, 2011) to estimate the transition rates

between the states. The full model contains 7 rate parameters corresponding to the 7

possible state transitions. We also considered simpler models by constraining

combinations of parameters to have identical rates. The simplest model has only a single

rate parameter, corresponding to identical rates for all transitions. We used AIC to

compare model fit across all (n = 877) possible model simplifications and report both

the best model and the model-averaged parameter values of all highly supported models

(ΔAIC $\leq$ 2 of best model).

2.5 Relative frequency of community assembly pathways

Having inferred the transition rates between states, we then used these to estimate the

relative contribution of elevational displacement (ED), elevational sorting (ES), and

elevational conservatism (EC) to pairs leaving state 1 and arriving in states 2, 4 and 5.

Specifically, we used the inferred rates to perform 1000 posterior predictive simulations

using the Gillespie algorithm for constant rates (Gillespie, 1977). Under the Gillespie

algorithm, transitions between states correspond to events. The simulation starts at

time t = 0, indicating the time at divergence for all sister species. The waiting time (δ) to

the next event is determined by a random draw from an exponential distribution with

the mean equal to the sum of all transition rates across all sister pairs. For example, if

the estimated transition rate from state 1 to 2 (r12) is 0.05 and there are currently 5

pairs in state 1, then this transition adds 0.25 to the total rate. Transition rates are

assumed to be constant through time, but older species will have more time to

experience a transition. The event that occurs at time $t + \delta$ is selected with a probability

equal to the relative contribution of each rate to the total rate. This transition applies to

a single species pair and this pair is chosen at random with equal weighting across all

pairs that are currently in the relevant state. As we simulate forward in time, species can

no longer transition if their age is less than t and thus the simulation continues for a

period equal to the age of the oldest sister pair. During the simulation we record the

percentage of sister pairs passing through each of the three different community

assembly pathways (i.e. ED, ES, and EC), and report the median and 95% confidence

intervals (CI) in these %'s across simulation runs.

2.6 Assessing model fit

Although the model is optimised using maximum likelihood, model fit may be poor if the

underlying assumptions of the model are not met. If that is the case, estimated rates will

poorly reflect the empirical observations. We assessed how well the predicted transition

rates can predict (*i*) the change in frequency of states through time, and (ii) the

distribution of sister species divergence times for each state using the output of the

posterior predictive simulations. To examine if the model can adequately predict state

changes through time we binned species pairs in three equal sized age bins of 55-56

sister pairs each. We used three bins to ensure we would approximately capture

nonlinear changes in the prevalence of states through time beyond just increases and

decreases. The prevalence of each of the five states per bin is then compared between

the empirical data and 95% CI's constructed from the final prevalence of the simulated

states. If the empirical prevalence falls within the 95% CI this would indicate good model

fit (Figure 3a), but large CI's likewise indicate high uncertainty. Finally, we also

compared the empirical age distribution of each state to the average distribution of ages

across the posterior simulations (Figure 3b).

2.7 Simulation tests of accuracy and precision

Using simulations, we further evaluated the model by assessing if we can both

accurately and precisely recover the true transition rates. High accuracy indicates that

the model is not biased towards over- or under-estimating transition rates. High

precision indicates that the estimated rates are close to the true rates. We explored

multiple scenarios (Supplementary Information Table S1-S4), designed to characterise

the three different community assembly pathways (S1-S3), as well as a scenario

assuming identical transition rates among states (S4) and one corresponding to the

transition rates inferred from the model-averaged fit to the empirical data (S5). For each

scenario, we performed 100 replicate simulations using the observed number and ages

of sister pairs. For each simulation, we then performed an identical model fitting

procedure as for our empirical data, resulting in transition rate estimates according to

the best and model-averaged approach for each simulation. We constructed 95% CIs

from the estimates across the simulations for the best and model-averaged approaches

to evaluate the accuracy and precision of the rate estimation. Additionally, we

determined the coverage of the models (*sensu* Morris *et al.*, 2019) by inferring how often

the (a) 95%CI of the best model and (b) unconditional CI (Anderson, 2008) of the

model-averaged estimates for an individual simulation captured the true rates as

predetermined for every scenario ('rate coverage'). We also investigated how potential

error in rate estimates impacts the estimation of the relative frequency of community

assembly pathways taken by sister pairs ('pathway coverage', Table S3, S4).

3 Results

3.1 Empirical distribution of sister pair states

Across the n = 165 sister species pairs, the majority are currently in allopatry with

overlapping elevation ranges (state 1: 58%). The next most common state is sympatry

with overlapping elevation ranges (state 5: 18%), with fewer pairs having non-

overlapping elevational ranges and occurring in sympatry (state 4: 7%), or allopatry

(state 3: 8%), or having non-overlapping elevational ranges in sympatry but overlapping

elevational ranges in allopatry (state 2: 10%). The median age of pairs that are in

allopatry with overlapping elevation ranges (state 1: 2.09Myr) is younger than all other

states (state 2: 3.36, state 3: 2.74, state 4: 3.98, and state 5: 2.73Myr), consistent with

our assumption that it is the initial state at the time of species divergence.

3.2 Transition rates between states

We found that our best model contains two parameters and constrains r12, r13, r15,

r24, r25, and r45 to 0.07 (95% CI: 0.06-0.09), and r34 to 0.28 (95% CI: 0.13-0.63)

transitions per pair per million years (Myr^{-1}). While there are 66 models that are highly

supported (≤2 ΔAIC), the model-averaged rate estimates are very similar to the best

model. Because of this, we focus on the model-averaged results below. The model

averaged results show that the transition rate from the initial state of allopatry with

overlapping elevation ranges (state 1) to non-overlapping elevation ranges, either while

in allopatry (r13 = 0.06 Myr^{-1}) or upon the attainment of sympatry (r12 = 0.08 Myr^{-1}) is

relatively slow, and similar to attaining sympatry while conserving elevational ranges

(r15 = 0.08 Myr^{-1}). Once elevational differentiation has occurred in either allopatry

(state 3) or upon secondary contact (state 2), the transition to elevational differentiation

in both allopatry and sympatry (state 4) is relatively fast (r34 = 0.18 Myr^{-1} and r24 =

0.17 Myr^{-1}). In contrast, the transition from a state where species have non-overlapping

ranges in sympatry (state 2 and 4) to one where they co-occur at the same elevation in sympatry (state 5) is relatively slow ($r25 = 0.06$ Myr^{-1} and $r45 = 0.08$ Myr^{-1}).

3.3 The relative contribution of community assembly pathways

The posterior-predictive simulations show that following speciation a similar proportion of pairs embark on the ES (28.36%, 95% CI: 18.52-38.89%), ED (35.06%, 95% CI: 24.66-47.30%) and EC pathways (36.00%, 95% CI: 25.71-45.95%) (Figure 2b). Of the pairs with currently non-overlapping elevation ranges in sympatry (states 2 and 4), substantially more are inferred to be generated through ED (81.48%, 95% CI: 65.51-95.83%) than ES (18.52%, 95% CI: 4.17-34.49%) (Figure 2c). Of the species pairs that attain overlapping elevation ranges in sympatry (state 5) almost all of these (88.46%, 95% CI: 74.98-97.68%) attain this state via the EC pathway rather than the ES (0.00%, CI: 0.00-9.09%)or ED (9.09%, 95% CI: 0.00-21.74%) pathways (Figure 2d). This can be explained because although a similar proportion of pairs embark on each pathway (i.e. $r12 \approx r13 \approx r15$), species pairs taking the ES (1→3→4→5) and ED (1→2→4→5 or 1→2→5) pathway must pass through a number of intermediate states to attain overlapping elevation ranges in sympatry and this takes much longer than the direct EC pathway (1→5). We note that these key results remained qualitatively the same when using (i) alternative thresholds for defining elevation range overlap, (ii) different approaches for incorporating regional variation in elevation ranges, (iii) different criteria for detecting elevational displacement on secondary contact and (iv) relaxing assumptions about the initial state of sister pairs at speciation (Figure S3, S4, S5, S6, S7, S8).

3.4 Assessing model fit and the accuracy and reliability of estimated transition rates

Our simulations show that transition rates from state 1 (r12, r13, and r15) (Figure 1), can be reliably and accurately estimated irrespective of the simulation scenario (Table S1, S2). Accordingly, our simulations also show that we can reliably recover the contribution of ED and ES pathways in generating sympatric sister pairs with non-overlapping elevation ranges (states 2 and 4) (Table S3-S4). Later transitions (r24, r25, r34, r45) are estimated with less accuracy, probably because there are relatively few old sister pairs and thus less information to reliably estimate these rates. As a result, when it comes to explaining the species that are currently sympatric and have overlapping elevation ranges (state 5) there is a bias towards overestimating the contribution of ED and underestimating EC (Table S3, S4) (Figure 2d). We note that this does not influence

our conclusions, because despite a bias against detecting EC, we still infer this as by far
the most common pathway to state 5. Overall, our simulations show that we can predict
the incidence of the five states well through time (Figure 3). The simulations capture the
main patterns, namely: an increase with age of sympatric pairs with overlapping
elevation ranges (state 5), a decrease in allopatric pairs with overlapping elevation
ranges (state 1), and slight increases for pairs that are either in allopatry or sympatry
but have diverged in elevation (state 2, 3 and 4).
412
4 Discussion
Multiple historical processes have been proposed to explain the patterns of range
overlap and turnover across tropical elevational gradients. However, because previous
studies have focused on static biogeographic patterns and treated elevational sorting,
displacement and conservatism as alternative explanations, their relative contribution
has remained unknown. Through our analysis of the dynamics of sympatry and
elevation overlap among Neotropical montane birds, we present three key findings.
First, we show that the relative contributions of different historical assembly models can
be reliably inferred given current phylogenetic and geographic data among sister
species. Second, our results show that elevational displacement upon secondary contact
(>81%) contributes substantially more to the turnover of closely related bird species
across elevation gradients than sorting (<19%) of species that diverged to occupy
different elevations in allopatry. Finally, we show that the overlap of sister species along
elevation gradients is almost entirely (>88%) explained by species attaining sympatry
while retaining their ancestral elevational range, rejecting more complex scenarios
requiring displacement upon secondary contact followed by subsequent shifts to occupy
the same elevational range.
The high biodiversity of tropical mountains is associated with rapid turnover in
community composition, as ecologically similar species replace one another up the
mountain. Previous studies of montane birds have argued that elevational displacement
rather than sorting would be the dominant process explaining such elevational
replacements, but these studies have been unable to quantify the relative contribution of
these pathways (Diamond, 1973; Freeman, 2015; Freeman *et al.*, 2022). Our results
show that both sorting and displacement contribute to the turnover in sister species
across elevations, and that current patterns of sympatry cannot be explained by either

process acting alone. However, we also show that elevational displacement upon contact
is far more common that sorting, accounting for ~81% of sympatric sister pairs with
non-overlapping elevation ranges. These results are consistent with the idea that
negative interactions between species (e.g. competition or apparent competition) cause
displacement to occupy different elevations, and that this is the dominant process
driving the layering of species across mountain slopes.

While our analysis shows that both elevational sorting and displacement are
involved in generating turnover of sister species across elevation gradients, we find that
sympatry is frequently reached without differentiation in elevation. We estimate that
~45% of sister species living on the same mountain slope and ~88% of species
additionally living at the same elevation attain sympatry while conserving their
ancestral elevation range. Such a high frequency of elevational range conservatism may
not seem surprising given that previous studies focussed on the drivers of speciation
have shown that most vertebrate sister species have overlapping elevation ranges
(Cadena *et al.*, 2011). However, a high frequency of elevation overlap among sister
species is by itself inconclusive regarding how communities assemble because the same
pattern can arise under the elevational displacement pathway if species subsequently
expand their elevation ranges as they diverge across alternative niche axis (e.g. resource
use) and competition is relaxed. Indeed, based on his studies of New Guinea birds,
elevation displacement followed by expansion was the pathway proposed by Diamond
(1973) for how diverse communities at a given elevation had assembled. Our
phylogenetic approach to modelling the dynamics of elevation range overlap, enables us
to exclude this possibility. Indeed, we estimate that displacement and subsequent
overlap in species elevational ranges contributes little (9.09%) to current patterns of co-
occurrence among sister species along elevation slopes.

The finding that a high proportion of allopatric sister species directly transition
to occupy overlapping elevations in sympatry need not suggest that competition or
other negative species interactions are unimportant in limiting coexistence for these
species. Indeed, evidence that interspecific competition limits elevational ranges is
widespread in birds (Terborgh & Weske, 1975; Freeman *et al.*, 2022). Instead, species
attaining sympatry without diverging in their elevation range may have diverged across
alternative niche dimensions (either in allopatry or upon secondary contact) such as
resource or microhabitat use. Such an explanation would be consistent with previous

evidence that sympatry among Neotropical bird species is limited by divergence in key

trophic traits, such as beak size (Pigot & Tobias, 2013; Pigot *et al.*, 2018). Furthermore,

while our analysis is limited to sister species and does not consider how patterns of

elevation overlap and turnover arise among more distantly related lineages, it is among

recently diverged species that competition is expected to be strongest (Pigot & Tobias,

2013). Therefore, while extending our models to deeper phylogenetic scales is an

important direction for future research we do not expect this to overturn our finding

that conservatism in elevation ranges is a more common route to co-occurrence than

elevational range displacement followed by expansion.

In addition to the effects of phylogenetic scale, it is important to consider the

possibility that the relative mix of different assembly processes may not be static over

geological time. Given the relatively young age of many Neotropical montane radiations,

and that there is little evidence for a slowdown in the rates of diversification (Weir,

2006; Harvey *et al.*, 2020), local niche space at any point along the elevational gradient

may currently be far from saturated. As niches become increasingly densely packed over

time, it is possible that elevational sorting and displacement may become increasingly

important pathways to sympatry as has been suggested for New Guinean (Diamond,

1973) and Himalayan birds (Price *et al.*, 2014). It would therefore be interesting to apply

our modelling approach to other organisms and tropical mountain systems to test

whether the relative mix of different assembly pathways is consistent or varies across

different contexts and stages of evolutionary radiations.

Fitting our model requires reliable information on both geographic and elevation

overlap and for these spatial properties to each be encoded as simple binary states. In

reality, elevation ranges can vary in complex ways, for example, showing overlap in

some places but not in others, and with species shifting their elevation ranges across

regions to match changes in climate and habitats. Here, we focussed on Neotropical

birds occupying humid forests for which distributions are relatively well known and

where we could incorporate regional variation in species elevation limits into our

metrics of overlap. While we found that estimates of different community assembly

pathways were robust to the way elevation range overlap was scored, accounting for the

full complexity of overlap patterns may best be addressed using a spatially explicit

model (Reijenga *et al.*, 2021; Hagen, 2022). Such a model could incorporate the actual

environmental conditions (and their variation across space) that limit geographic ranges

as well as biotic interactions beyond those occurring between only sister species

(Rangel *et al.*, 2018; Hagen *et al.*, 2021). Due to their complexity, such spatially explicit

models are difficult to fit to empirical data, but the parameters that we estimate here

could be used as priors to inform the rates of key processes (e.g. transition times to

sympatry or elevation divergence).

In our model, sympatric species that co-occur at the same elevation (state 5)

represent the final stage in the pathway of community assembly. The fact that not all

sister pairs exist in this state is explained by new speciation events re-setting the cycle,

so that insufficient time has passed since speciation for most sister pairs to transition to

this state. Our model uses the observed distribution of sister pair ages which are

determined by rates of speciation and extinction (Weir & Schluter, 2007), but it does not

consider how the geographic or elevational overlap of species could in turn feedback to

influence these diversification dynamics. Because speciation and extinction events

prune or replace the set of sister pairs in a phylogeny, if these rates depend on the

geographic or elevation overlap of a species, this could bias the estimated transition

rates. For example, we speculate that sympatric species may have a higher

instantaneous probability of speciating because the expansion of their geographic range

makes them susceptible to renewed rounds of geographic isolation (Weir & Price, 2011).

In this case, our model may underestimate transition rates to sympatry. While we think

it is unlikely that such biases would also apply to elevation overlap, developing models

that incorporate the reciprocal evolution between geographic/elevation overlap and

diversification dynamics is an important avenue for future research (Goldberg *et al.*,

2011; van Els *et al.*, 2021; Landis *et al.*, 2022).

The enormous diversity of species in tropical mountain ranges appear as stark

outliers in statistical models seeking to explain global variation in species richness on

the basis of climate and topography (Rahbek *et al.*, 2019b). This is often attributed to the

rapid turnover of species across elevations and mountains (Rahbek *et al.*, 2019a). Our

results provide a different perspective, showing that the rate of range expansions

leading to sympatry are substantially accelerated by the capacity for species to occur on

the same slope without having to first diverge in their elevation range. Specifically, the

estimated mean lag time to sympatry following speciation is 6.01Myr, substantially

shorter than the 9.46Myr that would be expected if the direct transition from allopatry

to co-occurrence in sympatry at the same elevation (i.e. 1◻5) were precluded (see

Supplementary Information). Thus, in addition to the turnover of species across

elevations or mountains, a key additional ingredient explaining the high diversity of

tropical mountain ranges is the capacity for species to coexist locally without having to

diverge into different elevational zones.

Our model represents a simplification of the complex processes governing the

assembly of montane biotas and is limited to explaining the patterns of sympatry and

elevation overlap among sister species. However, to our knowledge this is the first study

to quantify the relative frequency of elevational sorting, displacement, and conservatism

in shaping species distributions across montane gradients. A key next step will be to test

how well our results generalise to different mountain regions or taxa that vary in their

ecology. Our model could also be applied to disentangle the dynamics of assembly across

other ecological gradients, such as the vertical layering of foraging niches among

rainforest birds (MacArthur, 1958), perch height among Anolis lizards (Lister, 1976), or

depth zonation in the Cichlid fish of East African rift lakes (Rodríguez & Lewis, 1997).

Funding information

This research was supported by a Royal Society University Research Fellowship

awarded to XX4 and studentship to XX1.

Data Accessibility Statement

Data used in this study was compiled from previously published sources. Code and the

data set are made publicly available on acceptance.

Author contributions

Conceptualisation, XX1, XX3, XX4; data curation XX1, XX2; formal analysis, XX1; funding

acquisition, XX4; investigation, XX1; visualisation, XX1; writing – original draft, XX1;

writing – review and editing, XX1, XX2, XX3, XX4. All authors have read and agreed to

submit the current version of the manuscript.

References

Anderson, D.R. (2008) *Model based inference in the life sciences: A primer on evidence*,

Springer, New York, NY, USA.

Anderson, S.A.S. & Weir, J.T. (2021) Character displacement drives trait divergence in a

continental fauna. *Proceedings of the National Academy of Sciences of the United*

States of America, **118**.

Barker, K., Burns, K.J., Klicka, J., Lanyon, S.M. & Lovette, I.J. (2015) New insights into New

World biogeography: An integrated view from the phylogeny of blackbirds,

cardinals, sparrows, tanagers, warblers, and allies. *Auk*, **132**, 333–348.

Billerman, S.M., Keeney, B.K., Rodewald, P.G. & Schulenberg, T.S. eds. (2020) *Birds of the*

World, Cornell Laboratory of Ornithology, Ithaca, NY, USA.

Brown, W.L. & Wilson, E.O. (1956) Character displacement. *Systematic Zoology*, **5**, 49–64.

Cadena, C.D. (2007) Testing the role of interspecific competition in the evolutionary

origin of elevational zonation: An example with buarremon brush-finches (aves,

emberizidae) in the neotropical mountains. *Evolution*, **61**, 1120–1136.

Cadena, C.D. & Céspedes, L.N. (2020) *Origin of Elevational Replacements in a Clade of*

Nearly Flightless Birds: Most Diversity in Tropical Mountains Accumulates via

Secondary Contact Following Allopatric Speciation. Neotropical diversification:

Patterns and processes, pp. 635–659. Springer.

Cadena, C.D., Kozak, K.H., Gómez, J.P., Parra, J.L., Mccain, C.M., Bowie, R.C.K., Carnaval,

A.C., Moritz, C., Rahbek, C., Roberts, T.E., Sanders, N.J., Schneider, C.J., Vanderwal, J.,

Zamudio, K.R. & Graham, C.H. (2011) Latitude, elevational climatic zonation and

speciation in New World vertebrates. *Proceedings of the Royal Society B: Biological*

Sciences, **279**, 194–201.

Coyne, J.A. & Orr, H.A. (2004) *Speciation*, Sinauer Associates, Inc, Sunderland, MA.

Diamond, J.M. (1973) Distributional ecology of New Guinea Birds. *Science*, **179**, 759–769.

van Els, P., Herrera-Alsina, L., Pigot, A.L. & Etienne, R.S. (2021) Evolutionary dynamics of

the elevational diversity gradient in passerine birds. *Nature Ecology and Evolution*,

5, 1259–1265.

Freeman, B.G. (2015) Competitive Interactions upon Secondary Contact Drive

Elevational Divergence in Tropical Birds. *The American Naturalist*, **186**, 470–479.

Freeman, B.G., Strimas-mackey, M. & Miller, E.T. (2022) Interspecific competition limits

bird species' ranges in tropical mountains. **377**, 416–420.

Gillespie, D.T. (1977) Exact stochastic simulation of coupled chemical reactions. *Journal*

of Physical Chemistry, **81**, 2340–2361.

Goldberg, E.E., Lancaster, L.T. & Ree, R.H. (2011) Phylogenetic inference of reciprocal

effects between geographic range evolution and diversification. *Systematic Biology*,

60, 451–465.

Hagen, O. (2022) Coupling eco-evolutionary mechanisms with deep-time environmental

dynamics to understand biodiversity patterns. *Ecography*, 1–16.

Hagen, O., Flück, B., Fopp, F., Cabral, J.S., Hartig, F., Pontarp, M., Rangel, T.F. & Pellissier, L.

(2021) *gen3sis: A general engine for eco-evolutionary simulations of the processes*

that shape Earth's biodiversity,.

Harvey, M.G., Bravo, G.A., Claramunt, S., Cuervo, A.M., Derryberry, G.E., Battilana, J.,

Seeholzer, G.F., Shearer McKay, J., O'Meara, B.C., Faircloth, B.C., Edwards, S. V., Pérez-

Emán, J., Moyle, R.G., Sheldon, F.H., Aleixo, A., Smith, B.T., Chesser, R.T., Silveira, L.F.,

Cracraft, J., Brumfield, R.T. & Derryberry, E.P. (2020) The evolution of a tropical

biodiversity hotspot. *Science*, **370**, 1343–1348.

Hilty, S.L. (2003) *Birds of Venezuela*, Princeton University Press, Princeton, NJ, USA.

Hilty, S.L. & Brown, W.L. (1986) *A guide to the birds of Colombia*, Princeton University

Press.

Jackson, C.H. (2011) Multi-state models for panel data: The msm package for R. *Journal*

of Statistical Software, **38**, 1–28.

Janzen, D.H. (1967) Why Mountain Passes are Higher in the Tropics. *The American*

Naturalist, **101**, 233–249.

Landis, M.J., Quintero, I., Muñoz, M.M., Zapata, F. & Donoghue, M.J. (2022) Phylogenetic

inference of where species spread or split across barriers. *Proceedings of the*

National Academy of Sciences of the United States of America, **119**.

Linck, E., Freeman, B.G. & Dumbacher, J.P. (2020) Speciation and gene flow across an

elevational gradient in New Guinea kingfishers. *Journal of Evolutionary Biology*, **33**,

1643–1652.

Linck, E.B., Freeman, B.G., Cadena, C.D. & Ghalambor, C.K. (2021) Evolutionary

conservatism will limit responses to climate change in the tropics. *Biology Letters*,

17.

Lister, B.C. (1976) The Nature of Niche Expansion in West Indian Anolis Lizards I:

Ecological Consequences of Reduced Competition. *Evolution*, **30**, 659–676.

Londoño, G.A., Chappell, M.A., Jankowski, J.E. & Robinson, S.K. (2017) Do

thermoregulatory costs limit altitude distributions of Andean forest birds?

Functional Ecology, **31**, 204–215.

MacArthur, R.H. (1958) Population Ecology of Some Warblers of Northeastern

Coniferous Forests. *Ecology*, **39**, 599–619.

Mittelbach, G.G. & Schemske, D.W. (2015) Ecological and evolutionary perspectives on

community assembly. *Trends in Ecology and Evolution*, **30**, 241–247.

Morris, T.P., White, I.R. & Crowther, M.J. (2019) Using simulation studies to evaluate

statistical methods. *Statistics in Medicine*, **38**, 2074–2102.

Pfennig, D.W. & Pfennig, K.S. (2010) Character displacement and the origins of diversity.

American Naturalist, **176**.

Phillimore, Orme, Thomas, Blackburn, Bennett, Gaston & Owens (2008) Sympatric

Speciation in Birds Is Rare: Insights from Range Data and Simulations. *The American*

Naturalist, **171**, 646.

Pigot, A.L., Jetz, W., Sheard, C. & Tobias, J.A. (2018) The macroecological dynamics of

species coexistence in birds. *Nature Ecology and Evolution*, **2**, 1112–1119.

Pigot, A.L. & Tobias, J.A. (2013) Species interactions constrain geographic range

expansion over evolutionary time. *Ecology Letters*, **16**, 330–338.

Price, T. (2008) *Speciation in birds*, Roberts and Co., Greenwood village, Colorado.

Price, T.D., Hooper, D.M., Buchanan, C.D., Johansson, U.S., Tietze, D.T., Alström, P., Olsson,

U., Ghosh-Harihar, M., Ishtiaq, F., Gupta, S.K., Martens, J., Harr, B., Singh, P. & Mohan,

D. (2014) Niche filling slows the diversification of Himalayan songbirds. *Nature*,

509, 222–225.

Pujolar, J.M., Blom, M.P.K., Reeve, A.H., Kennedy, J.D., Marki, P.Z., Korneliussen, T.S.,

Freeman, B.G., Sam, K., Linck, E., Haryoko, T., Iova, B., Koane, B., Maiah, G., Paul, L.,

Irestedt, M. & Jønsson, K.A. (2022) The formation of avian montane diversity across

barriers and along elevational gradients. *Nature Communications*, **13**, 1–13.

Quintero, I. & Jetz, W. (2018) Global elevational diversity and diversification of birds.

Nature, **555**, 246–250.

Rahbek, C., Borregaard, M.K., Antonelli, A., Colwell, R.K., Holt, B.G., Nogues-Bravo, D.,

Rasmussen, C.M.Ø., Richardson, K., Rosing, M.T., Whittaker, R.J. & Fjeldså, J. (2019a)

Building mountain biodiversity: Geological and evolutionary processes. *Science*,

365, 1114–1119.

Rahbek, C., Borregaard, M.K., Colwell, R.K., Dalsgaard, B., Holt, B.G., Morueta-Holme, N.,

Nogues-Bravo, D., Whittaker, R.J. & Fjeldså, J. (2019b) Humboldt's enigma: What

causes global patterns of mountain biodiversity? *Science*, **365**, 1108–1113.

Rangel, T.F., Edwards, N.R., Holden, P.B., Diniz-Filho, J.A.F., Gosling, W.D., Coelho, M.T.P.,

Cassemiro, F.A.S., Rahbek, C. & Colwell, R.K. (2018) Modeling the ecology and

evolution of biodiversity: Biogeographical cradles, museums, and graves. *Science*,

361, 1–14.

Reijenga, B.R., Murrell, D.J. & Pigot, A.L. (2021) Priority effects and the

macroevolutionary dynamics of biodiversity. *Ecology Letters*, **24**, 1455–1466.

Ridgely, R.S. & Greenfield, P.J. (2001) *The Birds of Ecuador*, Christopher Helm.

Rodríguez, M.A. & Lewis, W.M. (1997) Structure of fish assemblages along

environmental gradients in floodplain lakes of the Orinoco River. *Ecological*

Monographs, **67**, 109–128.

Stuart, Y.E. & Losos, J.B. (2013) Ecological character displacement: Glass half full or half

empty? *Trends in Ecology and Evolution*, **28**, 402–408.

Terborgh, J. & Weske, J.S. (1975) The role of competition in the distribution of Andean

birds. *Ecology*, **56**, 1237-562–576.

Vallely, A.C. & Dyer, D. (2018) *Birds of Central America*, Princeton University Press.

Walker, B., Stotz, D.F., Pequeno, T. & Fitzpatrick, J.W. (2006) Birds of the Manu Biosphere

Reserve. *Fieldiana Zoology*, **110**, 23–49.

Weber, M.G., Wagner, C.E., Best, R.J., Harmon, L.J. & Matthews, B. (2017) Evolution in a

Community Context: On Integrating Ecological Interactions and Macroevolution.

Trends in Ecology and Evolution, **32**, 291–304.

Weir, J.T. (2006) Divergent Timing and Patterns of Species Accumulation in Lowland and

Highland Neotropical Birds. *Evolution*, **60**, 842.

Weir, J.T. & Price, T.D. (2011) Limits to Speciation Inferred from Times to Secondary

Sympatry and Ages of Hybridizing Species along a Latitudinal Gradient. *The*

American Naturalist, **177**, 462–469.

Weir, J.T. & Schluter, D. (2007) The latitudinal gradient in recent speciation and

extinction rates of birds and mammals. *Science*, **315**, 1574–1576.

Figure 1 Hypothetical pathways towards sympatry in montane sister species. a) All

possible transitions (arrows) in our model between the different states (numbers) are

shown. States are defined by the geographical and elevational patterns of overlap

between sister species ranges (coloured ovals) on mountains. Species pairs are assumed

to start in a state of allopatry with overlapping elevational ranges (i.e. state 1). From this

initial state, pairs may follow different possible community assembly pathways

depending on which states they transition through: Elevational sorting (ES), Elevational

displacement (ED) and Elevational conservatism (EC). b-d) The relative dominance of

each pathway is expected to leave distinct signatures in how the prevalence of the five

states across sister pairs changes with time since divergence. Here, for illustrating these

distinct signals, curves of the expected prevalence are based on simulations in which

90% of sister pairs are assumed to pass through the (b) ES, (c) ED and (d) EC pathways.

Note that while in (a) state 2 is represented as the shift in the elevation range of one

sister species, both species could shift or contract relative to their range in allopatry.

Figure 2 The inferred contribution of each community assembly pathway in explaining

patterns of sympatry, turnover and overlap among sister species. a) estimated

prevalence of the five states with time since divergence (Myr), as represented in Figure

1 but based on rates inferred by our model from the empirical data. Note the prevalence

of state 2 (black) and 3 (yellow) are very similar. b) Percentage of sister pairs that have

left state 1 and have transitioned to state 3, 2, or 5 representing the elevational sorting

(ES), displacement (ED), or conservatism (EC) pathways respectively. c) the percentage

of pairs that currently occupy non-overlapping elevational ranges in sympatry (states 2

and 4) that have arisen through the ES or ED pathway. d) The percentage of pairs that

are sympatric with overlapping elevational distributions (state 5) that have arisen

through the ES, ED, or EC pathway. Results are obtained from posterior-predictive

simulations of the model-averaged parameter estimates and using a threshold of 20%

overlap and the 'average overlap' protocol to define elevational overlap.

Figure 3 Tests of model adequacy. Empirical and predicted prevalence of states through
time and age distributions of states. a) Empirical (circles) and predicted (triangles)
prevalence (triangles) represent the prevalence of the states within three age bins of
approximately equal size (n = 56, 55, and 55). Brackets represent the 95% CI of the
prevalence of each state across 1000 posterior-predictive simulations. b) Empirical and
predicted age distributions per state. Brackets indicate the mean and 95% CI for the
empirical age distributions and the average mean and 95% CI over 1000 simulations for
the predictions. Results are obtained from the model-averaged parameter estimates and
using a threshold of 20% overlap and the 'average overlap' protocol to define
elevational overlap.