TITLE:

Structural analysis of cholesterol binding and sterol selectivity by ABCG5/G8.

AUTHOR NAMES:

Danny Farhat[1,‡], Fatemeh Rezaei[1,‡], Milica Ristovski[1,2,‡!], Yidai Yang[1], Albert Stancescu[1,3,#],

Lucia Dzimkova[1,4], Sabrina Samnani[1,4], Jean-François Couture[1] and Jyh-Yeuan Lee[1,*]

AUTHOR ADDRESSES:

[1] Department of Biochemistry, Microbiology and Immunology, Faculty of Medicine, University

of Ottawa, Ottawa, Ontario, Canada

[2] Translational and Molecular Medicine Program, Faculty of Medicine, University of Ottawa,

Ontario, Ottawa, Canada

[3] Biochemistry Program, Faculty of Health Sciences, McMaster University, Hamilton, Ontario,

Canada

[4] Biochemistry Program, Faculty of Science, University of Ottawa, Ottawa, Ontario, Canada

AUTHOR INFORMATION

* Correspondence should be addressed to Jyh-Yeuan (Eric) Lee (Jyh-Yeuan.Lee@uOttawa.ca).

‡ Contributed equally.

Current address: MBiotech Program, University of Toronto-Mississauga

! Current address: Undergraduate Medical Education, Faculty of Medicine, University of Ottawa

ABSTRACT:

The ATP-binding cassette (ABC) sterol transporters are responsible for maintaining cholesterol homeostasis in mammals by participating in reverse cholesterol transport (RCT) or transintestinal cholesterol efflux (TICE). The heterodimeric ABCG5/G8 carries out selective sterol excretion, preventing the abnormal accumulation of plant sterols in human bodies, while homodimeric ABCG1 contributes to the biogenesis and metabolism of high-density lipoproteins. A sterol-binding site on ABCG5/G8 was proposed at the interface of the transmembrane domain and the core of lipid bilayers. In this study, we have determined the crystal structure of ABCG5/G8 in a cholesterol-bound state. The structure combined with amino acid sequence analysis shows that in the proximity of the sterol-binding site, a highly conserved phenylalanine array supports functional implications for ABCG cholesterol/sterol transporters. Lastly, *in silico* docking analysis of cholesterol and stigmasterol (a plant sterol) suggests sterol-binding selectivity on ABCG5/G8, but not ABCG1. Together, our results provide a structural basis for cholesterol binding on ABCG5/G8 and the sterol selectivity by ABCG transporters.

KEYWORDS:

ABCG5, ABCG8, ABCG1, ABCG sterol transporters, cholesterol, sterol selectivity, X-ray crystallography, molecular docking

ABREVIATIONS:

ABC, Adenosine-triphosphate-binding cassette; NBD, Nucleotide binding domain; TMD, Transmembrane domain; TMH, Transmembrane helix; RCT, Reverse cholesterol transport; PH, Phenylalanine Highway

1. INTRODUCTION

Lipid homeostasis in mammalian cells is a critical process regulated in part by the ATP binding cassette (ABC) lipid transporters, particularly the ABCG and ABCA cholesterol transporters, such as ABCG1, ABCG4, ABCG5/G8 and ABCA1. As a critical component of cellular membranes, cholesterol comprises >50% of total cellular lipid content and is further utilized as the precursor molecule for steroid hormones that modulate gene regulation as well as bile acids that are required for nutrient absorption. To maintain cholesterol balance, excess cholesterol must be eliminated from cells and tissues via the reverse cholesterol transport (RCT) pathway by circulating high-density lipoproteins [1–3]. Within the human ABCG protein subfamily, all members function as lipid floppases, with the notable exception of ABCG2, that opts for a more relaxed approach to substrate efflux by working as the sole multidrug transporter (MDR) of the subfamily [4,5]. In addition, the PDR subfamily of full-length transporters (ABCG2 homologues in *Saccharomyces cerevisiae*) are related to ABCG and sharing the structural fold and inverted topology. Moreover, ABCG transporters are becoming increasingly important with new findings uncovering their physiological roles in severe chronic diseases, notably such as cancer, atherosclerosis, diabetes and Alzheimer's disease [6–11]. While these proteins have gained more notoriety, the efflux mechanisms have remained a mystery, with various proposed hypotheses that are yet to be agreed upon.

In mammals, ABCG lipid transporters consist of the heterodimeric ABCG5/G8 and the homodimeric ABCG1 or ABCG4 complexes. ABCG5/G8 is localized on the canalicular membranes of the bile ducts in the liver and the brush border of enterocytes in the small intestines, where it mediates the final step of the RCT [9,12–14]. ABCG1 and ABCG4 translocate cholesterol within the plasma membrane and in endosomes [15,16] and play a key role in the regulating

cholesterol balance in the lung, the brain and in macrophage-rich tissues [17–20]. ABCG5/G8

undergoes obligatory heterodimerization and is unique in its capability of preferential efflux for

dietary plant sterols over cholesterol [8,9,21,22]. The mechanisms that govern such substrate

selectivity within this protein subfamily, however, remains elusive. Based on the crystal structure

of ABCG5/G8, we can now theorize that such substrate specificity among ABCG transporters may

be the result of the difference in hydrophobic valve composition between the multidrug transporter:

ABCG2 and the lipid transporters: ABCG1, ABCG4, ABCG5/G8 [23,24]. In addition, a sterol-

binding site was postulated at the membrane-transporter interface based on the crystal structure of

ABCG5/G8 [25], whereas studies in recent ABCG1 structures [26,27] have suggested sterol

binding at the cytoplasmic end of the transmembrane domain (TMD).

The general TMD topology is shared among ABCG subfamily proteins. Each half

transporter consists of six transmembrane helices (TMH) and the re-entry helices that are part of

the extracellular domains. A connecting helix bridges the TMD to the nucleotide-binding domain

(NBD), forming an amphipathic interface on the cytoplasmic leaflet of the lipid bilayers.

Meanwhile a triple-helical bundle brings the TMDs and the NBDs together to create a compact

and short-stacked transporter conformation [23–25]. At the cellular level, ABC lipid transporters

have been shown to alter the composition of their local membrane environment. Such revisions

may come from the subduing of lipid rafts by redistributing the high cholesterol content towards

the surrounding membrane regions [16,28]. Using *in vivo* and *in vitro* functional assays, we have

previously characterized a loss-of-function (LOF) mutation ABCG5$_{A540F}$, a conserved residue near

a putative sterol-binding site in ABCG5/G8 [25,29]. These findings suggested the likelihood of an

access site that allows the binding of cholesterol binding prior to its re-localization within the

membrane-embedded portion of the sterol transporters.

One of the significant challenges in studying the cholesterol-transporter interaction is to accurately identify the sterol molecules as either the transport substrates or the intrinsic component of cell membranes. The latter can also play regulatory roles in membrane protein functions, such as ABCG2 [30] or G-protein coupled receptors [31]. Besides experimental approaches to obtain atomic models by crystallography or electron microscopy, *in silico* molecular docking can allow researchers to predict the ligand-binding sites on a protein by using available structural models and ligand libraries. The availability of either ABCG5/G8 or ABCG1 structural can now make it possible to provide a molecular framework and to allow further structural analyses by using *in silico* tools, such as molecular docking or molecular dynamics simulation.

In this study, we first carried out amino acid sequence analysis of ABCG transporters, highlighting key conserved aromatic residues arranged in a spatially conserved pattern on TMH2. We solved the crystal structure of ABCG5/G8 in complex with cholesterol. The structure shows that an orthogonal cholesterol molecule fitting horizontally in front of A540, a conserved ABCG5 residue at this orthogonal sterol-binding site. *In vitro* cholesterol-binding assay showed an inhibition of sterol binding activity by the mutant $ABCG5_{A540F}$/G8. To determine the differential sterol binding activities on ABCG sterol transporters, blind molecular docking analysis was conducted on wild-type (WT) ABCG5/G8 or ABCG1 in the presence of cholesterol or stigmasterol, as well as selective mutants $ABCG5_{Y432F}$, $ABCG8_{N564P}$ and $ABCG5_{A540F}$. We also analyzed the cholesterol-binding patterns on the ATP bound states by using both experimental and homology models, showing the sterols only accumulated on the surface of the extracellular domain, possibly an intermediate state during the ATP-coupled sterol-transport cycle. Together with previous *in vivo* and *in vitro* observations, our results provide a structural basis for sterol selectivity of ABCG lipid transporters and a visual component to the sterol-transport mechanism.

2. RESULTS

2.1 Conserved phenylalanine cluster on THM2 of ABCG transporters:

Recent structural determination of ABCG transporters by cryo-electron microscopy [26,27,32,33] confirms the general molecular architecture as initially revealed by the crystal structure of an apo ABCG5/G8 (Lee et al., 2016). The TMH2 and TMH5 of both subunits are adjacent to one another, forming a dimer interface within the transmembrane domains. This interface has been hypothesized to be a sterol translocation pathway based on sterol localization as observed in the EM structures. However, to what extent this pathway contributes to sterol translocation remains unclear. In this study, we used multiple sequence alignment to analyze the amino acid sequence conservation in the TMH2 of human ABCG1/2/4/5/8, PDR5 of the yeast *Saccharomyces cerevisiae* [34], and White protein of *Drosophila melanogaster* [35,36](**Fig. 1**). Among these seven ABCG transporters, White is a close homologue to ABCG1, while the drug resistant PDR5 is an ABC drug transporter and an ABCG2 homologue. Using ABCG1 as sequence reference and analyzing 300 mammalian orthologs, each ABCG member includes a highly conserved phenylalanine (or tyrosine in ABCG5) at position F1 (**Fig 1A**). The adjacent position F2 shows similar conservation of hydrophobic residues, albeit with a more relaxed sequence homology than F1 (**Fig 1A**). Position F3 reveals a conserved phenylalanine in ABCG1, ABCG4 and the N-terminal half of PDR5 (PDR5-1), although the C-terminal half of PDR5 (PDR5-2) has a conserved phenylalanine just one residue off from F3. Position F4 shows the last highly conserved phenylalanine found on TMH2 (**Fig 1A**). These conserved phenylalanine residues are visualized through a Logo plot (**Fig 1B**), indicating at least two phenylalanine residues being flanked between positions F1 and F4 on the hydrophobic residue-rich TMH2.

When inspecting the spatial relationship of these phenylalanine residues, we observed an

open space immediately below the hydrophobic valve and above a cavity that is open to the

cytoplasm in the inward-facing conformation of ABCG transporters (**Fig. 1C, left**). Not only are

the F1-4 residues shared among the eight homologous monomers, but two or three aromatic side

chains also point inwards the TMD dimer interface, creating a highway-shaped open space (**Fig**

1C, right). We have coined this open space as the Phenylalanine Highway (PH) to highlight this

unique structural motif in the transmembrane domain. ABCG1 and ABCG4 share the most closely

spaced phenylalanine residues in their iteration of the PH, while the F2 in White is one residue off

from that in ABCG1. ABCG2, ABCG8 and PDR5 (either the N- or C-terminal half) have similar

spacing of their phenylalanine residues, whereas the equivalent residues on ABCG5 are replaced

by aromatic tyrosine residues. In addition, the phenylalanine side chains may be adjusted based on

the ligand placements (**Suppl Fig. 1**). Moreover, position F1 is spatially adjacent to the aromatic

residues of the previously described hydrophobic valve (or ABCG2's di-leucine valve/gate),

forming a closed interface in the inward-facing protein conformation (**Fig 1D & 1E**) [23]. It may

be possible to speculate that with such close spatial proximity, interaction between the

hydrophobic valve and the phenylalanine residues could occur during the catalytic cycle via certain

conformational changes. Further investigation is needed to address this question.

2.2 Crystal structure of ABCG5/G8 in an orthogonal cholesterol-binding state:

152   Several *bona fide* sterol molecules were observed on recent cryo-EM structures of ABCG1

or ABCG5/G8, which led to various models of their sterol-transport (or binding) mechanisms, but

more structural data and further mechanistic analysis are still required to establish a convergent

model of cholesterol transport. We have previously crystallized human ABCG5/G8 in an apo state

by bicelle crystallization. To gain the structural basis of the sterol-transport (or binding)

mechanism, in this study, we determined a cytoplasmic-facing crystal structure of cholesterol-

bound ABCG5/G8 at 4Å by X-ray crystallography. Three X-ray diffraction datasets were

integrated and scaled with acceptable statistics. The initial model was obtained by molecular

replacement using a previously resolved cryo-EM structure (PDB ID: 7JR7) [33], and the motifs

involved in ATP binding were adjusted based on the previous crystal structure (PDB ID: 5DO7)

[25]. Two ABCG5/G8 heterodimers are observed in the asymmetric unit. The protein model in

this report includes a corrected amino acid registry in the nucleotide-binding domain as previously

highlighted by Zhang *et al.* Additionally, it shows a consistent overall architecture and key

structural features as described previously, overall root mean square deviation (RMSD) of ~0.9 Å

when superimposed [12,25,33](**Suppl. Fig. 1A**). After initial refinement, we observed two clear

electron densities at the interface of symmetrically located transmembrane domains of these two

transporters, which resembled the polycyclic characteristic of cholesterol molecules. Therefore,

we added two cholesterol molecules to the initial model for further refinement and obtained a

significant improvement of the Rfree from 0.338 to 0.302 and reasonable geometrical validation

at current resolution. The final model shows an orthogonal cholesterol molecule in close contact

with $ABCG5_{A540}$ (namely the ABCG5-dominant side), with the 3-β-hydroxyl group forming a

weak hydrogen bonding with the side chain of $ABCG8_{T430}$ (4.3-4.5 Å). No sterol was observed on

the equivalent side of ABCG8 (namely the ABCG8-dominant side) (**Fig. 2A**). Selected regions of

the electron densities are shown in Suppl Fig. 2. The statistics of data processing and model

refinement are summarized in **Table 1**.

177
With an *in vivo* analysis of biliary cholesterol secretion and an *in vitro* ATPase assay, we

have previously characterized a loss-of-function (LOF) mutant $ABCG5_{A540F}$/G8 (A540F),

predicting a putative sterol-binding at this transporter-membrane interface [25,29]. In this study,

we developed an *in vitro* cholesterol-binding assay using tritiated cholesterol (see Materials and

Methods) and characterized cholesterol binding on purified ABCG5/G8 WT and A540F mutant

proteins (**Fig. 2B**). The reaction reached a saturation of cholesterol at ~3 μM with the B_{MAX} as

50.64±0.77 CPM, where the mutant showed about two-fold inhibition of the maximal cholesterol-

binding rate. The binding constant K_D and Hill coefficient h were estimated as 1.46 ± 0.28 μM (h

= 2.91) and 1.55 ± 0.01 μM (h = 2.98) for the WT and mutant proteins, respectively. Given the

ATPase activity of A540F mutation was suppressed by ~90% with similar K_M for sterols [29],

both assays suggest a functionally defective conformation by this LOF mutant. Together with the

new structural data, our analysis supports the role of the orthogonal cholesterol-binding site in

proximity to ABCG5$_{A540}$ (**Fig. 2**).

2.3 Asymmetric binding of cholesterol or stigmasterol on ABCG5/G8, but not ABCG1:

The ABC transporter superfamily translocate a broad spectrum of substrates across cellular

membranes, with individual members conveying their unique substrate specificities. To determine

how ABCG transporters bind sterol ligands, we used previously published apo models ABCG5/G8

(PDB ID: 5DO7 or 7JR7) and ABCG1 (PDB ID: 7R8D) [25,33] and carried out molecular docking

on the TMD with cholesterol or stigmasterol, a plant sterol previously shown to have the highest

accumulation rate in an Abcg5/g8-deficient mouse model [37]. Using UCSF DOCK6 suite [38],

we first optimized the docking protocol by carrying out a redocking of the ligands (topotecan,

mitoxantrone and imatinib) to reproduce their complex with ABCG2. Additional redocking was

done on a distinct ABCG2 model to further evaluate docking optimization. (**Suppl Fig. 3**) [39,40].

Figure 3 shows the docking results of all acceptable (or viable) poses for the two sterol

ligands, cholesterol and stigmasterol, on the ABCG sterol transporters according to the standard

flexible docking protocol as implemented in DOCK6 suites [38]. On ABCG5/G8, poses of sterols

are distributed asymmetrically at the opposing membrane-transporter interfaces, having more

docking conformations on the ABCG5-dominant side (**Fig. 3A/B**). The predicted orthogonal

cholesterol (*i.e.*, one of the top-3 poses) has a nearly identical position to the cholesterol molecule

in the newly determined cholesterol-bound structure (**Fig. 2 & 3D/E**). On the ABCG8-dominant

side, no orthogonal sterol was obtained, and either ligand are clustered on the inner leaflet of the

membrane's lipid-bilayered and with the topmost sterols in close distance from the lower end of

the PH (*i.e.*, ABCG5$_{Y432}$ or ABCG1$_{F455}$). Such asymmetric sterol binding on ABCG5/G8 suggests

another level of functional asymmetry by ABCG5/G8-mediated sterol transport. On the ABCG1

homodimer, acceptable poses are only observed on the inner leaflet of the lipid-bilayered

membranes, resembling those on the ABCG8-dominant side, and bound sterols can be equally

distributed on either side of the transporter-membrane interface (**Fig. 3A/B, bottom**). Such

symmetric sterol binding is distinct from ABCG5/G8, suggesting different mechanisms of using

the PH between ABCG1 and ABCG5/G8.

To analyze sterol docking with ATP-bound structures, we performed the analysis on an

ATP-bound ABCG1 and a homology model of ATP-bound ABCG5/G8 (see Materials and

Methods) [41]. Surprisingly, given the same docking grid as used for apo structures, sterol docking

on the ATP-bound ABCG1 or ABCG5/G8 only showed clusters on top of the extracellular

domains of the extracellular-facing structures and far away from the PH (**Fig. 4A/B**). As current

apo or ATP-bound structures of ABCG sterol transporters are rather limited to early and end states

of catalysis [25–27,33], the sterol poses as observed here may represent the post-efflux

conformations where the previously observed ligand binding sites have become inaccessible due

to the compression of protein's interfacial cavity in these ATP-bound structures. For this reason,

binding sites within the protein were unable to be detected and subsequently docked. We

additionally utilized MOLE2.5 [42] to determine the volume of cavities detected within the TMD.

In ABCG1 cavities substantially large were found in between the transmembrane helices, however

these were away from the central canal of the protein (**Suppl Fig. 5B**, left). In ABCG5/G8, while

one big cavity appeared within the central canal of the protein, it was found to be a conglomeration

of multiple spheroidal shaped cavities bridged together by very thin regions. Each cavity on its

own was presented to be far too small of a volume at around ~$200Å^3$ comparing to the volume of

cholesterol (**Suppl Fig. 5B**, Right), being ~$627Å^3$ [43]. Our data thus emphasizes the need for

occluded turnover state models, as seen in ABCG2 models [44].

2.4 Sterol selectivity by ABCG1 and ABCG5/G8:

Among mammalian ABCG sterol transporters, ABCG5/G8 is known to promote

absorption of cholesterol but not dietary sterols in the gastrointestinal tract [9]. To study such sterol

selectivity, we analyzed the distributions of the sterol-binding poses from the docking experiments

by using ABCG1 and ABCG5/G8 as model systems. On ABCG1, cholesterol cluster was gathered

uninterruptedly around F455, *i.e.*, F4 of PH, and docked similarly on both sides of the transporter-

membrane interface. The sterol cloud is angled towards the putative sterol binding cavity at the

NBD-TMD interface, in the membranes lower leaflet. (**Fig. 3A, bottom**). Stigmasterol equally

shows the same angling into the binding site, but with reduced possible poses when compared to

cholesterol on ABCG1 (**Fig. 3B, bottom**). The largest number of positions for cholesterol or

stigmasterol are located immediately below the PH and above the NBD-TMD interface, suggesting

a common sterol-binding cavity in ABCG sterol transporters. However, the orientation the

hydroxyl group on the sterols is not necessarily exposed to the aqueous milieu (**Fig. 3A/B,**

bottom).

On the ABCG5-dominant side of ABCG5/G8, stigmasterol binds in two clusters

partitioned between each monomer, leaving a gap in the middle, which is opposite to the

continuous cloud as revealed by cholesterol (**Fig. 3A/B, top**). In general, there are more possible

binding poses in the vicinity of ABCG5$_{A540}$ and ABCG8's PH, where interestingly all sterol

molecules appear to orient in a similar direction, *i.e.*, the hydroxyl group pointing to the NBD-

TMD interface. On the other hand, on the ABCG8-dominant side and near F4 of ABCG5's PH,

the sterol molecules are oriented in various but seemingly unspecific directions (or conformations)

(**Fig. 3A/B**). When compared to cholesterol which binds in two separate pockets (**Fig. 3A, top**),

stigmasterol has three binding pockets across both sides of the transporter-membrane interfaces

(**Fig. 3B, top**). Interestingly, the highest clusters of stigmasterol and cholesterol are located on the

opposite sides of the transporter, suggesting preferred association of sterol species on either side

of the transporter-membrane interface.

ABCG1$_{F455}$ and ABCG5$_{Y432}$ are at the equivalent position F4 of the PH. When comparing

the sterol binding patterns around these key residues, more cholesterol poses can be found around

ABCG1$_{F455}$ in an increasingly linear shape, while more stigmasterol poses were observed around

ABCG5$_{Y432}$ in a "C" shape (**Fig. 3A/B**). In the nucleotide-free structures as used for analysis here,

sterol binding in general is positioned more towards the orthogonal sterol-binding site in

ABCG5/G8 than in ABCG1. Sterol poses in ABCG1 cluster closer to the central cavity, around

PH's F4, while sterols bind ABCG5/G8 closer to the transporter-membrane interface (**Fig 3C**).

The sterols of the top binding poses all have hydroxyl groups pointing outwards and away from

PH's F4 position. Interestingly, the cholesterol molecules are similarly angled in both ABCG1 and

ABCG5/G8, *i.e.,* in parallel to the lipid bilayers.

2.5 Alteration of ABCG5/G8 cholesterol-binding activity by the LOF mutant A540F:

By exploiting the LOF mutant A540F, *in vivo* and *in vitro* functional studies support the

notion that it is exposed to the orthogonal cholesterol-binding site [25,29] (**Fig. 2**). Using

molecular docking and homology modelling, it is possible to develop hypotheses for the structural

basis of how this mutation impacts ABCG5/G8 function. In this study, we analyzed the effects of

inducing the A540F mutation on the binding positions of cholesterol as compared to in WT

ABCG5/G8 (**Fig. 5**). In the WT protein, the cholesterol cloud stretches from proline 415 (P415,

end of the connecting helix of ABCG8 and a landmark position) to A540 (**Fig. 5A**), with one of

the top poses being the orthogonal sterol adjacent to A540. The remaining top-three are vertically

oriented, exposing the hydroxyl group to the orthogonal sterol-binding site (**Fig. 5B, right & Fig.**

5C). Using a homology model containing the A540F mutation, the second and third best binding

positions were missing, leaving a gap between PH's F4 and the orthogonal sterol-binding site (**Fig.**

5B, left & 5C). As mutation was induced, the orientation of the orthogonal cholesterol is flipped

180° at the sterol-binding site (**Fig. 5D/E**), and the remaining non-top three poses have their

hydroxyl groups pointing to the cytoplasmic side of the lipid bilayer (**Fig. 5B**). Overall, lower

number of different cholesterol poses were docked on the mutant as compared to the WT protein.

The microenvironment surrounding the orthogonal cholesterol on A540F mutant are highlighted

in **Figure 5E**, showing that the phenylalanine substitution prevents the interactions between the

cholesterol and polar residues (*e.g.*, ABCG8$_{T430}$) that may attract the hydroxyl group in the WT

protein.

2.6 Lower transmembrane sterol-binding cavity in ABCG5/G8:

The recent cryo-EM structural determination of ABCG1 suggested a cholesterol-binding

cavity in the cytoplasmic leaflet of the bilayer [26,27]. The protein's surface within the binding

cavity in ABCG1 shows a narrow gap that is wider than corresponding cavity found in ABCG5/G8

(**Fig. 6A**); additionally, this cavity is located immediately below F4 of the PH (**Fig. 6B**). In the

WT proteins, $ABCG5_{Y432}$ and $ABCG8_{N564}$ are separated by 4.0 Å, whereas the equivalent positions

in $ABCG1_{F455}$ and $ABCG1_{P558}$ are farther separated by 6.9Å. We created an ABCG5/G8 homology

model that consists of mutations $ABCG5_{Y432F}$ and $ABCG8_{N564P}$ to mimic an ABCG1-like cavity.

The interaction between the mutant residues in ABCG5/G8 are at a similar distance and positions

to the corresponding residues in ABCG1 (**Fig. 6B/C**). This distance is significantly widened from

4.0 Å to 6.9 Å in the mutant. Due to the increased distance between the residues, the double mutant

lacks the ability to form such N-Y blockade and results in a deeper cavity than that of WT. We

then performed sterol docking analysis on this double-mutant model and analyzed the possible

sterol conformations in this cytoplasmic sterol-binding cavity. In the WT, stigmasterol appears to

bind more readily than cholesterol, where the cluster of docked stigmasterol poses is much bigger

than the cholesterol cloud (**Fig. 6D/E**). Both sterol clouds semi-encircle Y432 in a "C" shape.

When Y432 is mutated to phenylalanine, the number of docked sterols decreased significantly as

compared to the WT, simply localizing overhead of position 432 and suggesting lower sterol-

binding ability in the mutant. Interestingly, the size of viable poses becomes more alike for either

cholesterol or stigmasterol. Despite using $ABCG5_{Y432F}/G8_{N564P}$ mutations to mimic the width of

the cavity found in WT ABCG1, the binding pattern in the mutant does not resemble that of

ABCG1 (**Fig. 2A and Fig. 2B, bottom**), suggesting the possibility of either sterol transporter

adopting multiple conformations to bind sterols throughout the transport cycle. Further functional

assays are necessary to define the role of these residues and how they can obstruct sterol

translocation trough the cavity.

3. DISCUSSION

Atomic models of ABCG sterol transporters have provided the molecular framework to study how individual transporter proteins translocate different substrates. Yet, the location and methods by which these transporters bind sterols remains elusive and controversial. This is particularly true in the case of ABCG5/G8, which is believed to identify and bind plant sterols differently from cholesterol. Understanding how sterols binds on ABCG5/G8 and its homologs will allow us to study the impact of such sterol selectivity on the regulation of sterol transport. In this study, we have determined a cholesterol-bound crystal structure of ABCG5/G8, showing an orthogonal cholesterol binding on the ABCG5-dominant side at the transporter-membrane interface (**Fig. 2**). A mutant ABCG5$_{A540F}$ was previously demonstrated to downregulate biliary cholesterol efflux *in vivo* [25] and to inhibit sterol-coupled ATPase activity *in vitro* [29]. The crystal structure herein provides a structural evidence of this peripheral sterol binding site where a sterol molecule makes direct contact with the conserved ABCG5$_{A540}$ residue (**Fig. 2 & 5**). The sterol fits into an indented planar cleft within the upper region of the vestibule formed by the re-entry and the transmembrane helices. The captured cholesterol extends across the interfacial space, with its hydroxyl group forming a hydrogen bond with ABCG8$_{T430}$, while the hydrocarbon tail remains straight and in contact with hydrophobic residues on ABCG5, such as L516, L537, I562, F565 and F567 (**Fig. 5D**). We also showed an inhibition of the cholesterol-binding activity by the A540F mutant (**Fig. 2C**). The K_D value is similar for both WT and mutant proteins, echoing the similar K_M for sterol-coupled ATPase activity in previous studies, given the mutant suppressed ATP hydrolysis by ~90% [29]. Taken together with a 180° flipping of cholesterol by this mutant (**Fig. 5E**), current structural and functional data suggest a functionally defective conformation by

this LOF mutant, inhibiting both sterol and ATP associations to the transporter. The Hill

coefficient was estimated as ~3, although no evidence so far shows whether the transporter bind

one or multiple sterol molecules. Further structural and functional studies will be necessary to

determine how these sterols bind ABCG5/G8.

In a separate experiment, cholesterol was docked to the TMD of previously published

ABCG5/G8 structures (PDB ID: 5DO7 or 7JR7) [25,33], where one of the top-3 predicted poses

binds to the transporter in similar conformation as revealed by the crystal structure (**Fig. 3D/E**).

The core steroid rings and the hydroxyl group matched in similar tilt, although the hydrocarbon

tail varied slightly with minimal rotation. The latter may be explained by its relatively high

mobility, such that the deviation was expected. Using a homology model containing the A540F

mutation, the large aromatic group in position 540 introduced steric hindrance, consequently

disabling cholesterol docking onto the observed position as shown in the crystal structure or

predicted docking on WT protein (**Fig. 2 & 3**). In addition, cholesterol docking with the A540F

mutant resulted in a loss of top-3 poses as seen on the WT protein. Altogether, past and present

data support the existence of an orthogonal cholesterol binding site located around residue

$ABCG5_{A540}$, and the orthogonal sterol poses may be part of intermediate states of sterol-

translocation trajectory along the TMD.

Via a protein sequence analysis of primarily mammalian ABCG transporters, we have

identified a structural motif, "phenylalanine highway" along the TMH2 that consists of conserved

aromatic residues (usually phenylalanine) (**Fig. 1**). This motif is flanked by two highly conserved

aromatic residues (*i.e.*, positions F1 and F4), where position F4 appears to involve in sterol

capturing from the cytoplasmic end of the membranes (**Suppl Fig. 1)** and the PH may provide a

pathway for docked sterols to enter the dimer's interface (**Fig. 3**). Although not fully defined,

certain PH residues are believed to form a molecular clamp previously shown to mediate $\pi-\pi$

interactions with sterol molecules [39,41,45,46]. As substrates for ABCG tend to be polycyclic

with more or less an aromatic group that can interact with aromatic amino acids, the PH motif

possibly serves as a conserved molecular clamp of ABCG sterol transporters (**Fig. 1C**). The clamp-

like role of the lower PH terminus was recently suggested by structural studies of ABCG1 and

ABCG2. The ATPase activity of ABCG1 was inhibited by $ABCG1_{F455A}$ mutation [27], while

ABCG2-driven drug efflux was downregulated by $ABCG2_{F439A}$ mutation [46]. Interestingly,

mutation $ABCG2_{F439Y}$ or $ABCG2_{F439W}$ did not hamper the efflux function, emphasizing the

importance of aromatic residues in contributing to a substrate-binding route from the inner leaflet

of the membranes [46]. In ABCG5/G8, while the clamp-like function of $ABCG5_{Y432}$ has yet to be

showcased, our docking analysis and a previous *in vivo* data suggest its biochemical nature as a

key feature for the transporter function, presumably sterol binding. In this study, a double-mutant

model ($ABCG5_{Y432F}$/$ABCG8_{N564P}$) revealed a loss of the ability to form a hydrogen bonding at the

TMD dimer interface of the protein model (**Fig. 6B**) and a deeper cavity was formed to imitate

ABCG1, yet the overall binding patterns were adversely affected (**Fig. 6D/E**). While the

physiological impact of this double mutation has yet to be determined, such pair nonetheless

created a blockade in the cavity of ABCG5/G8's model. Since docking utilizes static protein

models, widening of the cavity through mutation helped to mimic the effects of possible dynamic

protein movement causing the subsequent fracture of the physical hindrance. This alteration in

binding conformations and obstruction of the molecular clamp may help explain why *in vivo*,

$ABCG5_{Y432A}$/$ABCG8_{WT}$ caused a significant decrease in biliary cholesterol efflux [25]. The

mutations we had modelled retained the aromatic group; the *in vivo* data thus suggested a loss of

the molecular clamp at the lower PH terminus. It is important to note that $ABCG5_{Y432}$ is also part

of the TMD polar relay network [23,29]. Unlike the ABCG8 equivalent (ABCG8$_{F461}$), ABCG5$_{Y432}$ may play a dual role in the sterol binding/translocation and the allosteric regulation that couples ATPase and sterol-transport activities [29].

The upper PH terminus (*i.e.*, ABCG5$_{Y424}$ and ABCG8$_{F453}$) lies immediately under the hydrophobic valve (**Fig. 1D/E**), suggesting a cooperativity between PH's F1 position and the hydrophobic valve in the sterol-transport function. ABCG5$_{Y424}$, ABCG8$_{F453}$ and ABCG1$_{F447}$ all extend their side chains towards the valve's aromatic residues, forming aromatic-aromatic interactions as suggested previously [27]. In ABCG1 and ABCG4, the di-phenylalanine valve may grant the possibility of aromatic interactions between the neighboring six phenylalanine residues. In contrast, the leucine and methionine from ABCG5/G8's hydrophobic valve provide an asymmetric gating of sterol transport, as opposed to the symmetric valve in ABCG1 or ABCG4. In ABCG2, the lack of any phenylalanine within its di-leucine valve denies $\pi-\pi$ interactions, although ABCG2$_{F431}$ (PH's F1) was shown to engage in drug-protein interactions with ABCG2 inhibitors [40]. Position F2 in the PH is moderately conserved **(Fig. 1A)**, although residues at this position has been proposed to involve sterol interactions, as suggested by decreased cholesterol efflux in the presence of ABCG1$_{F448A}$ mutant [32] or drug interaction in ABCG2 [47]. Interestingly, such a PH-like motif may not be limited to ABCG sterol transporters. We have observed similar phenylalanine spacing on the structures of other cholesterol-binding or -transport proteins, including ABCA1, ABCA7, SIDT1, SIDT2, PTCH1, and NPC1 (**Suppl Fig. 4**) [34,48–51]. The PH structural motif may be more common than expected, and further studies will be needed to define its functional roles.

The docking experiments in this study have allowed us to analyze different binding profiles of cholesterol or stigmasterol on ABCG1 and ABCG5/G8 (**Fig. 3**). Previous *in vivo* studies showed

that stigmasterol efflux by ABCG5/G8 was the fastest among dietary plant sterols [37]; hence

stigmasterol was used to analyze phytosterol/plant sterol binding to ABCG transporters. Both

proteins share cholesterol as a substrate yet deviate in phytosterol transporting abilities.

ABCG5/G8 is a known phytosterol transporter, and each transporter-membrane interface has a

distinct selectivity for one of the two docked sterols (**Fig. 3A/B**). In comparison, ABCG1 showed

similar binding conformations between stigmasterol and cholesterol, with a decreased amount of

stigmasterol binding poses on ABCG1 than ABCG5/G8. Together, our results indicate that sterol

selectivity not only occurs on the opposite transporter-membrane interfaces but also differentiate

ABCG5/G8 from other ABCG sterol transporters. In addition, our simulation showed the

hydrocarbon tail of sterol molecules leading to the PH along with the TMD dimer interface, rather

than the hydroxyl group as seen in previous cryo-EM structures of ABCG1 [27]. However, at least

one docking pose on the ABCG8-dominant side horizontally points its hydroxyl group to the dimer

interface, similar to that of a recent cryo-EM structure (**Suppl Fig. 5A**). We speculate that the

cholesterol orientation as seen in the cryo-EM structures may represent an intermediate

cholesterol-binding state among the docked poses (**Fig. 5**).

Studies on cholesterol redistribution in lipid rafts have suggested a TMD sterol access site

in ABCG1 (**Fig. 4**) [16,28,52]. Theoretically, ABCG1 intakes sterols from enriched lipid

microdomains, relocating them towards the surrounding sterol-deprived membranes. The docking

pattern follows the amphipathic nature of the lipid bilayers with cholesterol's polar head localizing

around the cytoplasmic matrix (**Fig. 3**), where. cholesterol's hydroxyl group is believed to generate

hydrogen bonds with the surrounding polar lipid regions, orienting the sterols head towards the

hydrophilic region [53]. Our docking analysis suggests ABCG1's apparent preference for the inner

leaflet sterols [16], where the transporters then translocate cholesterol to the upper leaflet, similar

to the recently proposed mechanism [26,32]. In ABCG5/G8, both crystallographic and docking

results indicated that sterol binding is not limited to the TMD cavity open towards to the

cytoplasmic side, but an orthogonal sterol-binding state is a possible intermediate state to drive

sterol efflux. As the TMD dimer interface in ABCG5/G8 is much narrower than that in ABCG1,

it is possible that ABCG5/G8 similarly uptakes sterol substrates as ABCG1. Still, sterols would

need to undergo a series of yet-to-be characterized movements from their longitudinal orientations

to an orthogonal binding state as shown in the cholesterol-bound structure (**Fig. 2 &3**). Among

ABCG5/G8 substrates, plant sterols may show higher binding preferences as suggested by

stigmasterol poses (**Fig. 3**).

In conclusion, our crystallographic and computational analysis have highlighted possible

regions of sterol binding and interactions on ABCG5/G8. This study shows a conserved

phenylalanine highway motif on TMH2 that may play an essential role in substrate recruitment

and binding and a crystal structure revealing an orthogonal cholesterol bound on the ABCG5-

dominant transporter-membrane interface, consistent with the leading cholesterol poses predicted

by molecular docking. The docking analysis of cholesterol and stigmasterol also suggested the

asymmetric sterol-binding potential on ABCG5- and ABCG8-dominant transporter-membrane

interfaces, and difference in sterol-docking patterns may contribute to different substrate

specificity between ABCG1 and ABCG5/G8. Further structural and functional studies will reveal

more insight into sterol selectivity and translocation trajectory by ABCG sterol transporters.

4. MATERIALS AND METHODS

4.1 Protein purification and crystallization

For purification of human WT ABCG5/G8 from *Pichia pastoris* yeast, we have followed the procedures based on our previously reported protocol [25]. With minor modifications, the procedures are briefly described in the following. For each purification, six litres of yeast cell culture were used for ~200 mL of the microsomal membrane preparation at the total protein content of ~10 mg/mL. The membranes were solubilized by equal volume of the solubilization buffer (50 mM Tris-HCl, pH 8.0, 100 mM NaCl, and 10% glycerol, 2% (w/v) β-dodecyl maltoside (β-DDM, Anatrace), 0.5% (w/v) cholate (Sigma-Aldrich), 0.25% (w/v) cholesteryl hemisuccinate Tris (CHS-Tris, Anatrace), 5 mM imidazole, 5 mM β-mercaptoethanol (β-ME), 2 μg/ml leupeptin, 2 μg/ml pepstatin A, 2 mM PMSF). Insoluble membranes were removed by ultracentrifugation, and the solubilized supernatant was treated with a final concentration and 0.1 mM Tris (2-carboxylethylcarboxyethyl) phosphine (TCEP). Following that, tandem affinity chromatography was performed. The soluble membrane proteins were separated using a 20-mL nickel-nitrilotriacetic acid (Ni-NTA) column. Then, the peak fractions of ABCG5/G8 from the Ni-NTA eluates were mixed with the equal amount of CBP column wash buffer (50 mM HEPES, pH 7.5, 100 mM NaCl, 0.1% (w/v) β-DDM, 0.05% (w/v) cholate, 0.01% (w/v) CHS (Steraloids), 0.1 mM TCEP ,1 mM $CaCl_2$, 1 mM $MgCl_2$) and then loaded onto a 5-mL CBP column. The CBP column was washed sequentially to exchange detergents to DMNG, and ABCG5/G8 proteins were finally collected using an elution buffer (50 mM HEPES, pH 7.5, 300 mM NaCl, 2 mM EGTA, 0.1% (w/v) DMNG, 0.05% (w/v) cholate, 0.01% (w/v) CHS, 1 mM TCEP). The N-linked glycans and the CBP tag were cleaved by endoglycosidase H (Endo H, ~0.2 mg per 10–15 mg purified protein, New England Biolabs) and HRV-3C protease (~2 mg per 10–15 mg purified proteins) for

overnight at 4 °C. The CBP tag-free proteins were further purified by gel filtration chromatography

using an ÄKTA Purifier and a Superdex 200 30/100 GL column in the buffer containing 10 mM

HEPES, pH 7.5, 100 mM NaCl, 0.1% (w/v) DMNG, 0.05% (w/v) cholate, 0.01% (w/v) CHS. The

peak fractions were pooled together and treated by reductive methylation. The methylated proteins

were relipidated by overnight incubation with 0.5 mg/ml DOPC: DOPE (3:1, w/w) and further

purified by a 2-mL Ni-NTA column, followed by iodoacetamide treatment for cysteine alkylation.

The relipidated proteins were treated with 1 mM AMPPNP (sodium salt, Roche) for overnight at

4 °C and desalted with a PD-10 column. The desalted and lipidated proteins were incubated

overnight with cholesterol (prepared in isopropanol) to a final concentration of ~20 μM, and

protein precipitants were removed by ultracentrifugation for 10 min at 4 °C. Using 100 kDa cutoff

filters, the supernatants were concentrated to a final protein concentration of 30-50 mg/ml and

crystallized within a week.

Protein crystal growth was carried out by bicelle crystallization as described previously.

Briefly, the cholesterol-treated ABCG5/G8 proteins were reconstituted into 10% DMPC/CHAPSO

bicelles (lipids and detergents in a 3:1 (w/w) ratio, with the lipids containing 5 mol% cholesterol

and 95 mol% DMPC). After that, the proteins and bicelle stock were gently mixed in a 1:4 (v/v)

ratio, resulting in a final protein concentration of ~10 mg/ml. At 20 °C, the crystallization was

done in a hanging-drop vapor diffusion technique by mixing protein/bicelle preparation with

equal-volume crystallization reservoir solution containing 1.8–2.0 M ammonium sulfate, 100 mM

MES pH 6.5, 2–5% PEG400, and 1 mM TCEP. Protein crystals were harvested within 1-2 weeks

of incubation by first submerging the crystals with 0.2 M sodium malonate and then flash-freezing

the crystals in liquid nitrogen with Mitegen cryoloops.

4.2 Data collection of X-ray diffraction, structural determination and refinement

X-ray diffraction data were collected using beamlines 19-ID on a Quantum 315r detector

at the Advanced Photon Source (APS). Scaling and integration of three datasets were carried out

by HKL2000 software package, and the final data was scaled between 30Å and 4Å resolutions to

ensure I/σ >1 [54]. Using Phaser program, we obtain the phase information and a structural

solution by molecular replacement using a cryo-EM structure (PDB ID: 7JR7) as the initial

template [33,55]. Before model refinement by Phenix, the Phaser solution was corrected at the

Walker A, the Walker B, and the Signature motifs by the more defined registry in the crystal

structure (PDB ID: 5DO7) [25,56]. The initial refinement resulted in a refined model with Rfree

and Rwork of 0.338 and 0.309, respectively. Close inspection of the Fo-Fc map, two orthogonal

electron densities that clearly include the nature of polycyclic rings were observed between the

crystallographic dimer interface and on the ABCG5-dominant side where each electron density is

within the van der Waals' distance from ABCG5$_{A540}$. No such electron densities were observed on

ABCG8-dominant side. We thus modeled a cholesterol molecule per electron density, and we

refined the structure in the presence of one cholesterol molecule on each ABCG5/G8 heterodimer

by testing a series of sterol orientations using the program COOT [57]. Following our final

refinement, Rfree and Rwork were determined to be 0.302 and 0.244, respectively. The model

quality was validated by MolProbity as implemented in Phenix [56,58]. As a note, given the

proteins were co-crystallized with AMPPNP, we were not able to observe obvious electron

densities for the nucleotides from the current datasets at the current resolution.

4.3 Protein sequence analysis

All FASTA files were acquired using PSI-BLAST (position-specific iterated basic local

alignment search tool) as implemented in the NCBI protein sequence database. All queries were

acquired such that the organism belonged to the Mammalia family, and all partial and incomplete

queries were filtered out. FASTA sequences were then surveyed one by one to ensure that the

search criteria were executed properly, and no incomplete sequences were passed. FASTA

sequences were acquired for ABCG1, ABCG2, ABCG4, ABCG5, and ABCG8 (P45844,

Q9UNQ0, Q9H172, Q9H222, Q9H221) with the following PSI-BLAST search query:

"ABCXY"[Gene] AND "Mammalia"[Organism] NOT "partial"[All Fields], Whereby X denotes

the ABC subfamily, and Y denotes the member protein. Including 250-300 species, all alignments

and analyses were run locally using the Bio Python repository [59]. A multiple sequence alignment

(MSA) was conducted using all the FASTA sequences using a BLOSUM62 substituation matrix

[60], ClustalW [61] or PROMALS3D [62]. To score conservation, alignment files were made of

the FASTA sequence library using the Multiple Sequence Comparison by Log-Expectation

(MUSCLE) tool [63]. To generate a consensus sequence, a "dumb consensus" was conducted with

a default threshold of 70% and a required multiple of 1. Consensus sequences were acquired for

mammalian ABCG1, ABCG2, ABCG4, ABCG5, and ABCG8. Consensus sequences for each

protein subfamily and member were compared to one another to determine highly conserved amino

acids and amino acid sequences. For example, the amino acids present in the consensus sequence

of ABCG1 was individually aligned to ABCG2, ABCG4, ABCG5, and ABCG8. Conservation

scores above the set threshold (80%) were extracted for analytical purposes. An 80% similarity

threshold, as opposed to 70%, was used using AMAS method of multiple sequence alignment

because this analysis compared two consensus sequences together, 80% was chosen arbitrarily to

ensure the amino acids chosen were highly similar without extracting ambiguous amino acid [64].

For further analysis, the results from the conservation scores were compared to each other (*i.e.,* the

conservation scores of ABCG5 and ABCG2 was compared to conservation scores of ABCG5 and

ABCG8). This was done across all comparisons to generate a comprehensive list of the amino

acids that are conserved across all subfamily and member combinations. The same procedure was

used to analyze ABCA1, ABCA7, SIDT1, SIDT2, PTCH1, and NPC1, of which experimental

structures have been reported [48–51].

4.4 Homology Modelling

UCSF Modeller was used for homology modelling [65]. The sequence alignment was

created between the template and the mutant sequences using PROMALS3D [62]. The template is

the currently accepted PDB ID: 5DO7 in conjunction with X-ray data to create a more accurate

model with a better energy DOPE (discrete optimized protein energy) score. The mutant sequences

included ABCG5$_{Y432F}$, ABCG8$_{N564P}$ and ABCG5$_{A540F}$. After the alignment, UCSF Modeller was

used to generate 400 models and to rank them based on their DOPE scores. The best model was

taken and visualized using PyMOL or UCSF Chimera. For the ATP-bound model, the cryo-EM

structure of ATP-bound ABCG2 E211Q (PDB ID: 6HBU) was used as the template [41]. 100

models were generated with MODELLER, and the model with the best DOPE score was used for

the sterol docking experiments in this study. All models were evaluated using MolProbity web

server [58] to ensure no severe geometric violation in the Ramachandran plot [66].

4.5 Protein and ligand model preparation for docking

Protein models ABCG1, ABCG2, ABCG5/G8, PTCH1, NPC1, ABCA1 and PDR5

gathered from RCSB PDB database, PDB ID: 7OZ1/7R8D, 6VXI/6VXF/6VXJ,

5DO7/7R8E/7JR7, 6OEU, 6UOX, 5XJY, 7P03, respectively. ABCG4, White protein, ABCA7,

SIDT1/2 structures were downloaded from AlphaFold2 predictions [25–27,33,34,39,48–51].

Regions of interest in these proteins had very high confidence scoring (pLDDT>90). Visualization

of the structures done through PyMOL and Chimera. Double mutant $ABCG5_{Y432F}$ and

$ABCG8_{N564P}$ and $ABCG5_{A540F}$ were induced and modelled in MODELLER based on PDB ID:

5DO7. The proteins docked (ABCG1, ABCG5/G8, ATP-bound, ABCG5/G8 mutants) were cut at

the elbows, excluding the nucleotide-binding domain due to the large size of the full protein (>999

amino acids). Cholesterol was gathered from PDB ID: 7R8D, while stigmasterol was obtained in

PubChem [27]. ABCG2 ligands used for docking (imatinib and SN38) were previously modelled

within the structure [39,40]. Dunbrack 2010 rotamer library [67] and ANTECHAMBER [68] were

used in dockprep charge and hydrogen bond additions.

4.6 Molecular Docking

UCSF Dock6.9 [38] was utilized through SBGrid [69] to perform molecular docking.

UCSF Chimera [70] was used as the graphic software for the simulated models. Protein binding

site was determined by "sphgen" estimation specified at 10 Å distance from the ligand, located

roughly in the centre of the protein in ABCG transporters. The "gridbox" and "grid" were

generated and analyzed prior to the dock. To avoid potential mixing-up, we first performed

docking with each sterol ligand on either side of transporter-membrane interface, whose results

were consistent with those when placing the ligands at the centre of the TMD interface. Grid was

built at a distance of 8 Å from the selected "sphgen" spheres, and ligand's energy was minimized

through minor conformational changes. A flexible docking procedure was then followed by

calculation of 1000 or 5000 orientations, yielding no difference in results. The best scoring poses

were compiled and individually viewed through UCSF Chimera's viewdock tool, these were then

ranked by van der Waals energy (vdw_energy) and root mean square deviation (RMSD).

Interacting amino acids were identified and plotted by both Chimera and PyMOL [70].

To validate our protocol for sterol docking on ABCG1 or ABCG5/G8, we performed a

control experiment using ABCG2 and three of its known drugs (i.e., imatinib, and SN38) [39].

Using a previously published ABCG2 structure (PDB ID: 6FFC, 6VXH and 6VXJ), ABCG2 was

docked with the drug ligand (one at a time) and weighed again against the originally modelled

pose [40]. The simulated results showed consistent orientation of the drugs between predicted and

experimental data, and the best pose for each drug fits the originally modelled ligand with the best

RMSD and the lowest van vdw_energy value. The "best pose" in this study was therefore defined

as the lowest van der Waal's (vdw) energy, which was used to rank all of the docking results.

4.7 Cholesterol binding assay

A direct cholesterol-binding assay using detergent-purified ABCG5/G8 was developed

based on the previously described [71]. Briefly, 0.24µM of His-tagged WT ABCG5/G8, the

mutant or the control protein (*i.e.*, BSA) was incubated with 0-4 µM of cholesterol that was

prepared in the presence of [^{3}H]Cholesterol with a specificity of 110cpm/fmol (Perkein Elmer) in

50 µl of binding buffer containing 50mM Tris-HCl pH7.5, 100mM NaCl, and 0.05%(w/v) β-DDM

(Anatrace), 0.05% (w/v) sodium cholate (Sigma-Aldrich), 1mM TCEP, 30µl of pre-equilibrated

Ni-NTA bead (QIAGEN) on ice for overnight. The supernatant was removed by centrifugation at

600 rpm for 1min in a bench-top centrifuge. The bead was extensively washed with 600 µl of ice-

cold binding buffer plus 0.01% (w/v) CHS (Steraloids) for 4 times. To elute protein, 600 µl of

binding buffer plus 0.01% (w/v) CHS and 200mM imidazole was added and incubated with the

washed bead on ice for 30min. The bound [^{3}H]Cholesterol was quantified by measuring the

radioactivity of 500µl of elution fraction in a liquid scintillation counter (Perkin Elmer Tri-Carb

4910TR).

5. Data Availability

Diffraction data have been deposited in PDB under the accession code 8CUB.

AUTHOR CONTRIBUTIONS:

DF carried out docking analysis of ABCG transporters and other cholesterol-binding proteins; JYL

performed crystallization and collected X-ray diffraction data; FR processed the X-ray data and

determined the crystal structure. DF, MR and AS performed sequence analysis; MR and LD

generated homology models. YY, JFC and JYL designed the cholesterol-binding assay; YY

optimized the assay and analyzed the data. JYL oversaw the project, and DF, MR, FR, SS and JYL

wrote the manuscript. All authors have read and agreed to the submitted version of the manuscript.

FUNDING:

This research was supported by a startup grant from the University of Ottawa, a Natural Sciences

and Engineering Research Council Discovery Grant (RGPIN 2018-04070), a National New

Investigator Award from the Heart and Stroke Foundation of Canada, and a Canadian Institutes of

Health Research Project Grant (PJT-180640) to JYL.

ACKNOWLEDGEMENTS:

We thank the technical help from the staff of both the Common Equipment & Technical Services

and the Protein Biophysics Core Facilities, as well as the critical feedback from Drs. Thomas

Stockner, Gregory Graf, and Ms. Annahat Kochhar. We also thank Aiman Zein, Angelica Venes

and Toka Hussein for technical assistance. This work is partially based on the theses that were

submitted to fulfill in part the requirement for the degrees of Honors Bachelor of Science (DF &

MR). FR is partially supported by an Academia Sinica-University of Ottawa Travel Support for

Research Projects. The Advanced Photon Source is a US Department of Energy Office of Science

User Facility operated for the Department of Energy Office of Science by Argonne National

Laboratory (DE-AC02-06CH11357).

REFERENCES

[1]   W. Nan, L. Debin, C. Wengen, M. Fumihiko, T.A. R., ATP-binding cassette transporters G1 and G4 mediate cellular cholesterol efflux to high-density lipoproteins, Proc. Natl. Acad. Sci. 101 (2004) 9774–9779. https://doi.org/10.1073/pnas.0403506101.

[2]   A.M. Vaughan, J.F. Oram, ABCG1 Redistributes Cell Cholesterol to Domains Removable by High Density Lipoprotein but Not by Lipid-depleted Apolipoproteins *, J. Biol. Chem. 280 (2005) 30150–30157. https://doi.org/10.1074/jbc.M505368200.

[3]   L. Yu, S. Gupta, F. Xu, A.D.B. Liverman, A. Moschetta, D.J. Mangelsdorf, J.J. Repa, H.H. Hobbs, J.C. Cohen, Expression of ABCG5 and ABCG8 Is Required for Regulation of Biliary Cholesterol Secretion *, J. Biol. Chem. 280 (2005) 8742–8747. https://doi.org/10.1074/jbc.M411080200.

[4]   D.L. Austin, Y. Weidong, A.L. V., K. Tammy, G. Yongming, R.A. K., R.D. D., A multidrug resistance transporter from human MCF-7 breast cancer cells, Proc. Natl. Acad. Sci. 95 (1998) 15665–15670. https://doi.org/10.1073/pnas.95.26.15665.

[5]   J.D. Allen, A. van Loevezijn, J.M. Lakhai, M. van der Valk, O. van Tellingen, G. Reid, J.H.M. Schellens, G.-J. Koomen, A.H. Schinkel, Potent and Specific Inhibition of the Breast Cancer Resistance Protein Multidrug Transporter in Vitro and in Mouse Intestine by a Novel Analogue of Fumitremorgin C1, Mol. Cancer Ther. 1 (2002) 417–425.

[6]   Y. Namba, C. Sogawa, Y. Okusha, H. Kawai, M. Itagaki, K. Ono, J. Murakami, E. Aoyama, K. Ohyama, J. Asaumi, M. Takigawa, K. Okamoto, S.K. Calderwood, K. Kozaki, T. Eguchi, Depletion of Lipid Efflux Pump ABCG1 Triggers the Intracellular Accumulation of Extracellular Vesicles and Reduces Aggregation and Tumorigenesis of Metastatic Cancer Cells   , Front. Oncol. . 8 (2018).

664 https://www.frontiersin.org/article/10.3389/fonc.2018.00376.

[7] C. Tian, D. Huang, Y. Yu, J. Zhang, Q. Fang, C. Xie, ABCG1 as a potential oncogene in
lung cancer, Exp Ther Med. 13 (2017) 3189–3194. https://doi.org/10.3892/etm.2017.4393.

[8] M.-H. Lee, K. Lu, S. Hazard, H. Yu, S. Shulenin, H. Hidaka, H. Kojima, R. Allikmets, N.
Sakuma, R. Pegoraro, A.K. Srivastava, G. Salen, M. Dean, S.B. Patel, Identification of a
gene, ABCG5, important in the regulation of dietary cholesterol absorption, Nat. Genet.
27 (2001) 79–83. https://doi.org/10.1038/83799.

[9] K.E. Berge, T. Hui, G.G. A., Y. Liqing, G.N. V., S. Joshua, K. Peter, S. Bei, B. Robert,
H.H. H., Accumulation of Dietary Cholesterol in Sitosterolemia Caused by Mutations in
Adjacent ABC Transporters, Science (80-.). 290 (2000) 1771–1775.
https://doi.org/10.1126/science.290.5497.1771.

[10] I. Meurs, B. Lammers, Y. Zhao, R. Out, R.B. Hildebrand, M. Hoekstra, T.J.C. Van
Berkel, M. Van Eck, The effect of ABCG1 deficiency on atherosclerotic lesion
development in LDL receptor knockout mice depends on the stage of atherogenesis,
Atherosclerosis. 221 (2012) 41–47.
https://doi.org/https://doi.org/10.1016/j.atherosclerosis.2011.11.024.

[11] T.M. Do, M.-S. Noel-Hudson, S. Ribes, C. Besengez, M. Smirnova, S. Cisternino, M.
Buyse, F. Calon, G. Chimini, H. Chacun, J.-M. Scherrmann, R. Farinotti, F. Bourasset,
ABCG2- and ABCG4-Mediated Efflux of Amyloid-β Peptide 1-40 at the Mouse Blood-
Brain Barrier, J. Alzheimer's Dis. 30 (2012) 155–166. https://doi.org/10.3233/JAD-2012-
112189.

[12] A.A. Zein, R. Kaur, T.O.K. Hussein, G.A. Graf, J.-Y. Lee, ABCG5/G8: a structural view
to pathophysiology of the hepatobiliary cholesterol secretion, Biochem. Soc. Trans. 47

(2019) 1259–1268. https://doi.org/10.1042/BST20190130.

[13] Y. Liqing, H.R. E., L.-H. Jia, von B. Klaus, L. Dieter, C.J. C., H.H. H., Disruption of

Abcg5 and Abcg8 in mice reveals their crucial role in biliary cholesterol secretion, Proc.

Natl. Acad. Sci. 99 (2002) 16237–16242. https://doi.org/10.1073/pnas.252582399.

[14] S.B. Patel, A. Honda, G. Salen, Sitosterolemia: exclusion of genes involved in reduced

cholesterol biosynthesis, J. Lipid Res. 39 (1998) 1055–1061.

https://doi.org/10.1016/S0022-2275(20)33874-8.

[15] E. Pandzic, I.C. Gelissen, R. Whan, P.J. Barter, D. Sviridov, K. Gaus, K.-A. Rye, B.J.

Cochran, The ATP binding cassette transporter, ABCG1, localizes to cortical actin

filaments, Sci. Rep. 7 (2017) 42025. https://doi.org/10.1038/srep42025.

[16] O. Sano, S. Ito, R. Kato, Y. Shimizu, A. Kobayashi, Y. Kimura, N. Kioka, K. Hanada, K.

Ueda, M. Matsuo, ABCA1, ABCG1, and ABCG4 are distributed to distinct membrane

meso-domains and disturb detergent-resistant domains on the plasma membrane, PLoS

One. 9 (2014). https://doi.org/10.1371/journal.pone.0109886.

[17] S. Savary, F. Denizot, M.-F. Luciani, M.-G. Mattei, G. Chimini, Molecular cloning of a

mammalian ABC transporter homologous to Drosophila white gene, Mamm. Genome. 7

(1996) 673–676. https://doi.org/10.1007/s003359900203.

[18] D.D. Bojanic, P.T. Tarr, G.D. Gale, D.J. Smith, D. Bok, B. Chen, S. Nusinowitz, A.

Lövgren-Sandblom, I. Björkhem, P.A. Edwards, Differential expression and function of

ABCG1 and ABCG4 during development and aging, J. Lipid Res. 51 (2010) 169–181.

https://doi.org/10.1194/M900250-JLR200.

[19] T.Q. de Aguiar Vallim, E. Lee, D.J. Merriott, C.N. Goulbourne, J. Cheng, A. Cheng, A.

Gonen, R.M. Allen, E.N.D. Palladino, D.A. Ford, T. Wang, Á. Baldán, E.J. Tarling,

ABCG1 regulates pulmonary surfactant metabolism in mice and men[S], J. Lipid Res. 58 (2017) 941–954. https://doi.org/https://doi.org/10.1194/jlr.M075101.

[20] D. Sag, C. Cekic, R. Wu, J. Linden, C.C. Hedrick, The cholesterol transporter ABCG1 links cholesterol homeostasis and tumour immunity, Nat. Commun. 6 (2015) 6354. https://doi.org/10.1038/ncomms7354.

[21] G.A. Graf, L. Yu, W.-P. Li, R. Gerard, P.L. Tuma, J.C. Cohen, H.H. Hobbs, ABCG5 and ABCG8 Are Obligate Heterodimers for Protein Trafficking and Biliary Cholesterol Excretion*, J. Biol. Chem. 278 (2003) 48275–48282. https://doi.org/https://doi.org/10.1074/jbc.M310223200.

[22] C. Yang, L. Yu, W. Li, F. Xu, J.C. Cohen, H.H. Hobbs, Disruption of cholesterol homeostasis by plant sterols, J. Clin. Invest. 114 (2004) 813–822. https://doi.org/10.1172/JCI22186.

[23] N. Khunweeraphong, J. Mitchell-White, D. Szöllősi, T. Hussein, K. Kuchler, I.D. Kerr, T. Stockner, J.Y. Lee, Picky ABCG5/G8 and promiscuous ABCG2 - a tale of fatty diets and drug toxicity, FEBS Lett. 594 (2020) 4035–4058. https://doi.org/10.1002/1873-3468.13938.

[24] N. Khunweeraphong, D. Szöllősi, T. Stockner, K. Kuchler, The ABCG2 multidrug transporter is a pump gated by a valve and an extracellular lid, Nat. Commun. 10 (2019) 5433. https://doi.org/10.1038/s41467-019-13302-2.

[25] J.Y. Lee, L.N. Kinch, D.M. Borek, J. Wang, J. Wang, I.L. Urbatsch, X.S. Xie, N. V. Grishin, J.C. Cohen, Z. Otwinowski, H.H. Hobbs, D.M. Rosenbaum, Crystal structure of the human sterol transporter ABCG5/ABCG8, Nature. 533 (2016) 561–564. https://doi.org/10.1038/nature17666.

[26] L. Skarda, J. Kowal, K.P. Locher, Structure of the Human Cholesterol Transporter

ABCG1, J. Mol. Biol. 433 (2021) 167218.

https://doi.org/https://doi.org/10.1016/j.jmb.2021.167218.

[27] Y. Sun, J. Wang, T. Long, X. Qi, L. Donnelly, N. Elghobashi-Meinhardt, L. Esparza, J.C.

Cohen, X.-S. Xie, H.H. Hobbs, X. Li, Molecular basis of cholesterol efflux via ABCG

subfamily transporters, Proc. Natl. Acad. Sci. U. S. A. 118 (2021) e2110483118.

https://doi.org/10.1073/pnas.2110483118.

[28] M. Ristovski, D. Farhat, S.E.M. Bancud, J.-Y. Lee, Lipid Transporters Beam Signals from

Cell Membranes, Membranes (Basel). 11 (2021) 562.

https://doi.org/10.3390/membranes11080562.

[29] B.M. Xavier, A.A. Zein, A. Venes, J. Wang, J.Y. Lee, Transmembrane polar relay drives

the allosteric regulation for ABCG5/G8 sterol transporter, Int. J. Mol. Sci. 21 (2020) 1–20.

https://doi.org/10.3390/ijms21228747.

[30] Á. Telbisz, C. Özvegy-Laczka, T. Hegedűs, A. Váradi, B. Sarkadi, Effects of the lipid

environment, cholesterol and bile acids on the function of the purified and reconstituted

human ABCG2 protein, Biochem. J. 450 (2013) 387–395.

https://doi.org/10.1042/BJ20121485.

[31] G. Gimpl, Interaction of G protein coupled receptors and cholesterol, Chem. Phys. Lipids.

199 (2016) 61–73. https://doi.org/10.1016/j.chemphyslip.2016.04.006.

[32] D. Xu, Y. Li, F. Yang, C.-R. Sun, J. Pan, L. Wang, Z.-P. Chen, S.-C. Fang, X. Yao, W.-T.

Hou, C.-Z. Zhou, Y. Chen, Structure and transport mechanism of the human cholesterol

transporter ABCG1, Cell Rep. 38 (2022) 110298.

https://doi.org/https://doi.org/10.1016/j.celrep.2022.110298.

[33]   H. Zhang, C.-S. Huang, X. Yu, J. Lee, A. Vaish, Q. Chen, M. Zhou, Z. Wang, X. Min,

757   Cryo-EM structure of ABCG5/G8 in complex with modulating antibodies, Commun. Biol.

758   4 (2021) 526. https://doi.org/10.1038/s42003-021-02039-8.

759   [34]   A. Harris, M. Wagner, D. Du, S. Raschka, L.-M. Nentwig, H. Gohlke, S.H.J. Smits, B.F.

760   Luisi, L. Schmitt, Structure and efflux mechanism of the yeast pleiotropic drug resistance

761   transporter Pdr5, Nat. Commun. 12 (2021) 5254. https://doi.org/10.1038/s41467-021-

762   25574-8.

763   [35]   M.J. Ferreiro, C. Pérez, M. Marchesano, S. Ruiz, A. Caputi, P. Aguilera, R. Barrio, R.

764   Cantera, Drosophila melanogaster White Mutant w(1118) Undergo Retinal Degeneration,

765   Front. Neurosci. 11 (2018) 732. https://doi.org/10.3389/fnins.2017.00732.

766   [36]   Y. Mahé, Y. Lemoine, K. Kuchler, The ATP Binding Cassette Transporters Pdr5 and

767   Snq2 of Saccharomyces cerevisiae Can Mediate Transport of Steroids

768   in Vivo*, J. Biol. Chem. 271 (1996) 25167–25172.

769   https://doi.org/10.1074/jbc.271.41.25167.

770   [37]   L. Yu, K. von Bergmann, D. Lutjohann, H.H. Hobbs, J.C. Cohen, Selective sterol

771   accumulation in ABCG5/ABCG8-deficient mice, J. Lipid Res. 45 (2004) 301–307.

772   https://doi.org/10.1194/jlr.M300377-JLR200.

773   [38]   W.J. Allen, T.E. Balius, S. Mukherjee, S.R. Brozell, D.T. Moustakas, P.T. Lang, D.A.

774   Case, I.D. Kuntz, R.C. Rizzo, DOCK 6: Impact of new features and current docking

775   performance, J. Comput. Chem. 36 (2015) 1132–1156. https://doi.org/10.1002/jcc.23905.

776   [39]   B.J. Orlando, M. Liao, ABCG2 transports anticancer drugs via a closed-to-open switch,

777   Nat. Commun. 11 (2020) 2264. https://doi.org/10.1038/s41467-020-16155-2.

778   [40]   S.M. Jackson, I. Manolaridis, J. Kowal, M. Zechner, N.M.I. Taylor, M. Bause, S. Bauer,

779  R. Bartholomaeus, G. Bernhardt, B. Koenig, A. Buschauer, H. Stahlberg, K.-H. Altmann,

780  K.P. Locher, Structural basis of small-molecule inhibition of human multidrug transporter

781  ABCG2, Nat. Struct. Mol. Biol. 25 (2018) 333–340. https://doi.org/10.1038/s41594-018-

782  0049-1.

783 [41] I. Manolaridis, S.M. Jackson, N.M.I. Taylor, J. Kowal, H. Stahlberg, K.P. Locher, Cryo-

784  EM structures of a human ABCG2 mutant trapped in ATP-bound and substrate-bound

785  states, Nature. 563 (2018) 426–430. https://doi.org/10.1038/s41586-018-0680-3.

786 [42] D. Sehnal, R. Svobodová Vařeková, K. Berka, L. Pravda, V. Navrátilová, P. Banáš, C.-M.

787  Ionescu, M. Otyepka, J. Koča, MOLE 2.0: advanced approach for analysis of

788  biomacromolecular channels., J. Cheminform. 5 (2013) 39. https://doi.org/10.1186/1758-

789  2946-5-39.

790 [43] A.I. Greenwood, S. Tristram-Nagle, J.F. Nagle, Partial molecular volumes of lipids and

791  cholesterol., Chem. Phys. Lipids. 143 (2006) 1–10.

792  https://doi.org/10.1016/j.chemphyslip.2006.04.002.

793 [44] Q. Yu, D. Ni, J. Kowal, I. Manolaridis, S.M. Jackson, H. Stahlberg, K.P. Locher,

794  Structures of ABCG2 under turnover conditions reveal a key step in the drug transport

795  mechanism, Nat. Commun. 12 (2021) 4376. https://doi.org/10.1038/s41467-021-24651-2.

796 [45] N.M.I. Taylor, I. Manolaridis, S.M. Jackson, J. Kowal, H. Stahlberg, K.P. Locher,

797  Structure of the human multidrug transporter ABCG2, Nature. 546 (2017) 504–509.

798  https://doi.org/10.1038/nature22345.

799 [46] T. Gose, T. Shafi, Y. Fukuda, S. Das, Y. Wang, A. Allcock, A. Gavan McHarg, J. Lynch,

800  T. Chen, I. Tamai, A. Shelat, R.C. Ford, J.D. Schuetz, ABCG2 requires a single aromatic

801  amino acid to "clamp" substrates and inhibitors into the binding pocket, FASEB J. 34

(2020) 4890–4903. https://doi.org/10.1096/fj.201902338RR.

[47]   T. Nagy, Á. Tóth, Á. Telbisz, B. Sarkadi, H. Tordai, A. Tordai, T. Hegedűs, The transport

pathway in the ABCG2 protein and its regulation revealed by molecular dynamics

simulations, Cell. Mol. Life Sci. 78 (2021) 2329–2339. https://doi.org/10.1007/s00018-

020-03651-3.

[48]   H. Qian, X. Zhao, P. Cao, J. Lei, N. Yan, X. Gong, Structure of the Human Lipid Exporter

ABCA1, Cell. 169 (2017) 1228-1239.e10. https://doi.org/10.1016/j.cell.2017.05.020.

[49]   X. Qi, P. Schmiege, E. Coutavas, J. Wang, X. Li, Structures of human Patched and its

complex with native palmitoylated sonic hedgehog, Nature. 560 (2018) 128–132.

https://doi.org/10.1038/s41586-018-0308-7.

[50]   T. Long, X. Qi, A. Hassan, Q. Liang, J.K. De Brabander, X. Li, Structural basis for

itraconazole-mediated NPC1 inhibition, Nat. Commun. 11 (2020) 152.

https://doi.org/10.1038/s41467-019-13917-5.

[51]   J. Jumper, R. Evans, A. Pritzel, T. Green, M. Figurnov, O. Ronneberger, K.

Tunyasuvunakool, R. Bates, A. Žídek, A. Potapenko, A. Bridgland, C. Meyer, S.A.A.

Kohl, A.J. Ballard, A. Cowie, B. Romera-Paredes, S. Nikolov, R. Jain, J. Adler, T. Back,

S. Petersen, D. Reiman, E. Clancy, M. Zielinski, M. Steinegger, M. Pacholska, T.

Berghammer, S. Bodenstein, D. Silver, O. Vinyals, A.W. Senior, K. Kavukcuoglu, P.

Kohli, D. Hassabis, Highly accurate protein structure prediction with AlphaFold, Nature.

596 (2021) 583–589. https://doi.org/10.1038/s41586-021-03819-2.

[52]   O. Sano, A. Kobayashi, K. Nagao, K. Kumagai, N. Kioka, K. Hanada, K. Ueda, M.

Matsuo, Sphingomyelin-dependence of cholesterol efflux mediated by ABCG1, J. Lipid

Res. 48 (2007) 2377–2384. https://doi.org/10.1194/jlr.M700139-JLR200.

[53]   I. Ermilova, A.P. Lyubartsev, Cholesterol in phospholipid bilayers: positions and

orientations inside membranes with different unsaturation degrees, Soft Matter. 15 (2019)

78–93. https://doi.org/10.1039/C8SM01937A.

[54]   Z. Otwinowski, W.B.T.-M. in E. Minor, [20] Processing of X-ray diffraction data

collected in oscillation mode, in: Macromol. Crystallogr. Part A, Academic Press, 1997:

pp. 307–326. https://doi.org/https://doi.org/10.1016/S0076-6879(97)76066-X.

[55]   M.J. Grayling, PhaseR: An R package for phase plane analysis of autonomous ODE

systems, R J. 6 (2014) 43–51. https://doi.org/10.32614/rj-2014-023.

[56]   P.D. Adams, P. V Afonine, G. Bunkóczi, V.B. Chen, I.W. Davis, N. Echols, J.J. Headd,

L.-W. Hung, G.J. Kapral, R.W. Grosse-Kunstleve, A.J. McCoy, N.W. Moriarty, R.

Oeffner, R.J. Read, D.C. Richardson, J.S. Richardson, T.C. Terwilliger, P.H. Zwart,

PHENIX: a comprehensive Python-based system for macromolecular structure solution,

Acta Crystallogr. D. Biol. Crystallogr. 66 (2010) 213–221.

https://doi.org/10.1107/S0907444909052925.

[57]   P. Emsley, K. Cowtan, {\it Coot}: model-building tools for molecular graphics, Acta

Crystallogr. Sect. D. 60 (2004) 2126–2132. https://doi.org/10.1107/S0907444904019158.

[58]   V.B. Chen, W.B. Arendall 3rd, J.J. Headd, D.A. Keedy, R.M. Immormino, G.J. Kapral,

L.W. Murray, J.S. Richardson, D.C. Richardson, MolProbity: all-atom structure validation

for macromolecular crystallography, Acta Crystallogr. D. Biol. Crystallogr. 66 (2010) 12–

21. https://doi.org/10.1107/S0907444909042073.

[59]   P.J.A. Cock, T. Antao, J.T. Chang, B.A. Chapman, C.J. Cox, A. Dalke, I. Friedberg, T.

Hamelryck, F. Kauff, B. Wilczynski, M.J.L. de Hoon, Biopython: freely available Python

tools for computational molecular biology and bioinformatics, Bioinformatics. 25 (2009)

1422–1423. https://doi.org/10.1093/bioinformatics/btp163.

[60]   S. Henikoff, J.G. Henikoff, Amino acid substitution matrices from protein blocks, Proc.

Natl. Acad. Sci. U. S. A. 89 (1992) 10915–10919.

https://doi.org/10.1073/pnas.89.22.10915.

[61]   J.D. Thompson, D.G. Higgins, T.J. Gibson, CLUSTAL W: improving the sensitivity of

progressive multiple sequence alignment through sequence weighting, position-specific

gap penalties and weight matrix choice, Nucleic Acids Res. 22 (1994) 4673–4680.

https://doi.org/10.1093/nar/22.22.4673.

[62]   J. Pei, B.-H. Kim, N. V Grishin, PROMALS3D: a tool for multiple protein sequence and

structure alignments, Nucleic Acids Res. 36 (2008) 2295–2300.

https://doi.org/10.1093/nar/gkn072.

[63]   R.C. Edgar, MUSCLE: multiple sequence alignment with high accuracy and high

throughput, Nucleic Acids Res. 32 (2004) 1792–1797. https://doi.org/10.1093/nar/gkh340.

[64]   C.D. Livingstone, G.J. Barton, Protein sequence alignments: a strategy for the hierarchical

analysis of residue conservation, Bioinformatics. 9 (1993) 745–756.

https://doi.org/10.1093/bioinformatics/9.6.745.

[65]   B. Webb, A. Sali, Protein Structure Modeling with MODELLER BT - Protein Structure

Prediction, in: D. Kihara (Ed.), Springer New York, New York, NY, 2014: pp. 1–15.

https://doi.org/10.1007/978-1-4939-0366-5_1.

[66]   C.J. Williams, J.J. Headd, N.W. Moriarty, M.G. Prisant, L.L. Videau, L.N. Deis, V.

Verma, D.A. Keedy, B.J. Hintze, V.B. Chen, S. Jain, S.M. Lewis, W.B. Arendall 3rd, J.

Snoeyink, P.D. Adams, S.C. Lovell, J.S. Richardson, D.C. Richardson, MolProbity: More

and better reference data for improved all-atom structure validation, Protein Sci. 27 (2018)

293–315. https://doi.org/10.1002/pro.3330.

[67] M. V Shapovalov, R.L. Dunbrack Jr, A smoothed backbone-dependent rotamer library

for proteins derived from adaptive kernel density estimates and regressions, Structure. 19

(2011) 844–858. https://doi.org/10.1016/j.str.2011.03.019.

[68] J. Wang, W. Wang, P.A. Kollman, D.A. Case, Automatic atom type and bond type

perception in molecular mechanical calculations, J. Mol. Graph. Model. 25 (2006) 247–

260. https://doi.org/https://doi.org/10.1016/j.jmgm.2005.12.005.

[69] A. Morin, B. Eisenbraun, J. Key, P.C. Sanschagrin, M.A. Timony, M. Ottaviano, P. Sliz,

Collaboration gets the most out of software, Elife. 2 (2013) e01456.

https://doi.org/10.7554/eLife.01456.

[70] E.F. Pettersen, T.D. Goddard, C.C. Huang, G.S. Couch, D.M. Greenblatt, E.C. Meng, T.E.

Ferrin, UCSF Chimera - A visualization system for exploratory research and analysis, J.

Comput. Chem. 25 (2004) 1605–1612. https://doi.org/10.1002/jcc.20084.

[71] A. Radhakrishnan, L.-P. Sun, H.J. Kwon, M.S. Brown, J.L. Goldstein, Direct Binding of

Cholesterol to the Purified Membrane Region of SCAP: Mechanism for a Sterol-Sensing

Domain, Mol. Cell. 15 (2004) 259–268. https://doi.org/10.1016/j.molcel.2004.06.019.

 Table 1: Crystallographic data collection and refinement.

Beamline	APS 19-ID
Number of crystals	3
Wavelength	0.9794
Resolution range	30.58 - 4.051 (4.195 - 4.051)*
Space group	I 2 2 2
Unit cell (*a, b, c* (Å))	173.268 230.11 249.819 90 90 90
Total reflections	427824 (30232)
Unique reflections	38173 (915)
Multiplicity	11.2 (14.4)
Completeness (%)	83.96 (22.75)
Mean I/sigma(I)	5.90 (1.41)
R-merge	3.392 (5.094)
CC1/2	0.377 (0.137)
CC*	0.74 (0.491)
Reflections used in refinement	34374 (915)
Reflections used for R-free	2016 (54)
R-work	0.2459 (0.2832)
R-free	0.3029 (0.3039)
CC(work)	0.864 (0.703)
CC(free)	0.817 (0.715)
Number of non-hydrogen atoms	18498

macromolecules	18442
ligands	56
RMS(bonds)	0.005
RMS(angles)	0.93
Ramachandran favored (%)	93.54
Ramachandran allowed (%)	5.90
Ramachandran outliers (%)	0.56
Rotamer outliers (%)	0.00
Clashscore	11.97
Average B-factor	52.45
macromolecules	52.45
ligands	49.48
Number of TLS groups	1

889
890 * Statistics for the highest-resolution shell are shown in parentheses.

Figure 1. Conservation of aromatic amino acids on TMH2 of human ABCG family protein and homologues, including their localization in relation to key regions in the ABCG5/G8 heterodimer. (A) Sequence analysis of transmembrane helix five (TMH2) was done through PROMALS3D. Residues were screened for high conservation through BLAST sequence analysis. Red highlights perfect (or near perfect) aromatic residue conservation. Orange highlights conservation with ± 1 residue displacement. Green shows moderate conservation. Black shows non-sequence, but rather structural conservation. **(B)** Logo plot of TMH2 residues are colored according to their chemical nature. Red: acidic amino acids; Black: hydrophobic amino acids; Purple: neutral residues; Green: polar residues. **(C)** Structural conservation of highlighted aromatic amino acids from (A), using ABCG1 amino acid numeration as a reference. ABCG1 (PDB ID: 7OZ1, green), ABCG2 (PDB ID: 6VXI, pink, ABCG5/G8 (PDB ID: 5DO7; G5 in blue, G8 in

yellow), PDR5-1 and 2 (PDB ID: 7PO3, light grey). *Drosophila* White (purple) and ABCG4
(orange) are the only two Alphafold models, both have helices with predicted local distance
difference test score of >90. F447 (F1) and F455 (F4) residues in relation to ABCG1, have the
highest conservation. The residues directions are maintained, with F447 stacking under the
hydrophobic valve and F455 pointing towards the lumenal space between the TMD dimer. ABCG5
is the only monomer which has tyrosine's instead of phenylalanine's. ABCG1 and ABCG4 have
large hydrophobic amino acids pointing lumenally causing a mass hydrophobic region spanning
most of the transmembrane domain. **(D, E)** ABCG5/G8 heterodimer is split and shown in a top-
down view, highlighting the location of conserved F453 and Y424 (red) in relation to hydrophobic
valves (black). **(D)** In ABCG8, F453 (F1) stacks underneath F576, which (along M577) comprise
the hydrophobic valve shown in black. **(E)** In ABCG5, Y424 (F1) is stacked underneath both
hydrophobic valve residues; F548 and L549, shown in black.

915
Figure 2. Crystal structure of cholesterol-bound ABCG5/G8 and cholesterol-binding assay.
(A) Cholesterol-bound ABCG5/G8 structure is plotted in cartoon presentation. The first zoomed-
in picture shows the electron density related to the cholesterol molecule which was observed at
the interface of transmembrane domains of the crystallographic dimers (top right). The
cholesterol molecule was modeled to the primary structure as shown in the second zoomed-in
picture (bottom right). Alanine 540 (A540) is marked in a black sphere as a residue involved in
the cholesterol-binding site. **(B)** The cholesterol-binding activities of the WT and the A540F
mutant were measured by using [^{3}H]-cholesterol, showing the binding parameters for the WT
and the mutant with Bmax = 50.64±0.77 and 27.65±0.01 μCi/ml, K_D = 1.46 ±0.28 and 1.55±0.01
μM, and h = 2.91 and 2.98, respectively. A P-value = 0.043 was obtained by paired t-test
between WT and mutant. All calculations were performed using GraphPad Prism programs.

Docking cholesterol | Crystal structure cholesterol

927
Figure 3. Cholesterol and stigmasterol docking in ABCG5/G8 and ABCG1. (A-B) Cholesterol
(blue, left) and stigmasterol (violet, right) are bound to either side of the interface between the two
dimers. TMH2 and TMH5 are labeled as II and V. **(A)** Cholesterol or stigmasterol in ABCG5/G8
docks differently on either side (shown only ABCG5 here). *Left of Y432*, cholesterol docks in three
conformations: lightly angled upwards above Y432, horizontal on the X-axis below Y432 and
vertical. Stigmasterol (right) poses to the left of Y432 has a "C" shaped pattern that surrounds
Y432. *Right of Y432*, cholesterol binds in a continuous pattern, starting at A540 and ending at the
lower edge of the transmembrane domain (TMD). A single cholesterol conformation binds
horizontally near A540. Stigmasterol binds in two separated clusters with the top cluster showing
horizontal poses ahead of A540 and the bottom cluster containing horizontally oriented sterols.
(B) On ABCG1 (PDB ID: 7OZ1), either cholesterol or stigmasterol includes similar docking poses
on both sides. The sterols are angled towards the putative sterol binding cavity on the cytoplasmic
end of the bilayers. Given similar binding pattern, more cholesterol poses are obtained than
stigmasterol. **(C)** Top sterol poses on both transporters are illustrated as a superposition. ABCG1
cholesterol and stigmasterol are colored blue and violet respectively. ABCG5/G8 cholesterol and
stigmasterol are colored dark blue and dark red respectively. In general, sterols in ABCG5/G8

dock higher, near the orthogonal sterol-binding site, whereas sterols in ABCG1 binds under F455
(below F4 of the phenylalanine highway). **(D, E)** The cholesterol docking pose (orange) is aligned
with either cholesterol in the asymmetric unit of the crystal structure (light blue) with an RMSD
of 0.4 Å (D) and 0.9 Å (E).

948
Figure 4. (A, B) Sterol docking on ATP-bound ABCG1 (A) and ABCG5/G8 (B) models. ABCG1
(PDB ID: 7R8E) in gray; ABCG5/G8 in pale-green and tan. Cholesterol (blue) and stigmasterol
(violet) docked shown in a front view (lower) and a top-down view (upper). In brackets is the
extent of the phenylalanine highway (PH) within the transmembrane domain (TMD) of the protein.

Figure 5. Wild-type and mutant A540F ABCG5/G8 docking predictions. (A-C) ABCG5 and
ABCG8 are colored in pale-green and tan respectively, and the cholesterol docking on WT results
are shown in (A). In comparison, WT docking is shown in blue while mutant is shown in teal (B).
Landmark residues ABCG5$_{A540}$ and ABCG8$_{P415}$ are highlighted in tan and orange respectively for
the orientation (B), showing that WT has a continuous cholesterol binding pattern from A540 to
P415 while the mutant has a break in the docking prediction, separating the two clusters. Two of
the top 3 poses in the wild type docking appear to be missing in the A540F mutant (C). **(D, E)** The
micro-environment surrounding cholesterol highlights sterol-protein interaction. Residues are
shown as sticks. From left to right, ABCG8: L426, M427, T430; ABCG5: F559, F540, L536, I562,
L537 Q568, T566, F567, L516, L637. The side chain of F540 is shown in spheres to highlight the
mutation.

966
Figure 6. Lower transmembrane cavity in ABCG5/G8 and ABCG1 with cholesterol and
stigmasterol docking on wild-type and double mutant (Y432F, N564P) ABCG5/G8. (A)
ABCG1's (PDB ID: 7OZ1, gray) hollow transmembrane region was previously denoted as a
putative cholesterol binding cavity contrasted with ABCG5 (pale-green)/G8(tan). All atoms are
included (but not displayed) with the surface fit over the protein. **(B)** WT ABCG5/G8 (PDB ID:
7RJ7, pale-green/tan) with residues Y432 and N564 shown as sticks. Distance between the two
amino acids is ~4.0 Å, and for the equivalent residues in ABCG1 (PDB ID: 7OZ1, gray), F455
and P558, they are ~6.9 Å apart. A homology model of the double-mutant ABCG5/G8, shown in
white, has Y432 and N564 substituted for phenylalanine and proline, respectively, as observed in
ABCG1. **(C, D)** Cholesterol (blue) and stigmasterol (violet) docking was simulated on WT and
double mutant (white) ABCG5/G8, with only ABCG5 shown here. Yellow: residue 432 on
ABCG5. Wild type has more potential binding poses than the double mutant that only binds a
handful sterols above residue 432.

Supplemental Figure 1. Topological representation of the ABCG5/G8 and Superposition of the phenylalanine highway on ABCG1 and ABCG2. (A) *Left,* Topological representation of the ABCG5/G8 crystal structure (PDB ID: 8CUB) is depicted in empty cylinders, showing the transmembrane structural segments of ABCG5/G8 including connecting helixes (CnH), TM helixes, coupling helixes (CpH), and re-entry helixes. *Right,* The cartoon illustration of the ABCG5/G8 heterodimer with ABCG5 in pale-green and ABCG8 in tan colors. The N-glycosylation sites are shown in solid circles: $ABCG5_{N592}$ or $ABCG8_{N619}$ in red and black, $ABCG5_{N584}$ in black. **(B)** Side view of ABCG1 (PDB ID: 7OZ1), showing the difference in height of CHS (green) and cholesterol (purple). **(C)** Top-down view ABCG1 bound to CHS (PDB ID: 7OZ1, green) aligned with ABCG1 bound to cholesterol (7R8D, purple) emphasizing the difference in F455 rotation when interacting with different length ligands. Rotation occurs when longer ligands (such as CHS) reach deep enough into the transmembrane cavity, causing the alpha carbon and phenyl side chain of F455 to rotate. F447 is also seen stacked under hydrophobic valve. **(D)** Side view of ABCG2, showing the difference in localization of Imatinib (orange), Topotecan (pink), Mitoxantrone (green). **(E)**Top-down view of ABCG2 bound with Imatinib (PDB ID:

6VXH, orange), Topotecan (PDB ID :6VXJ, pink) and Mitoxantrone (PDB ID: 6VXI, green). Rotation of F439 is seen when substrates are deep enough into the cavity to interact with the residue. Contrarily, when they are not, such rotation events do not happen, as observed with Imatinib.

(A) ABCG5 TMH

TMH1   TMH2   TMH3   TMH4   TMH5   Re-entry helices   TMH6

(B) ABCG8 TMH

TMH1   TMH2   TMH3   TMH4   TMH5   Re-entry helices   TMH6

(C)   ABCG5   ABCG8   **(D)**

CnH   CpH   CnH   CpH   Cholesterol Molecules

Supplemental Figure 2. The fitting of the structural model in the electron densities (2Fo-Fc omit map). The representative segments of the human ABCG5/G8 proteins, including transmembrane (A/B), connecting (C), coupling helixes (C), and cholesterol molecules (D) are shown separately for G5 and G8 chains in the electron density with a mesh contoured at 1 σ and the thickness of 1.5 as implemented in PyMOL.

Supplemental Figure 3. Resolved ligand structures compared to dock prediction in ABCG2 (A-E) and ABCG5/G8 (F). (**A-C**) **Redocking of** resolved ligands on ABCG2 (blue): Topotecan/SN38 (PDB ID :6VXJ), Mitoxantrone (PDB ID: 6VXI), and Imatinib (PDB ID: 6VXH). Green highlights the minimized rigid conformation as calculated by Dock6.9. Orange highlights the best predicted pose. Blue ligands were observed in the experimentally determined structures. RMSD of all superpositions is (0.57Å – 1.5Å). (**D/E**) Ligands were aligned and docked to ABCG2 (PDB ID: 6FFC). Pink highlights the best RMSD predicted pose. The RMSD of all superpositions is (2.1Å – 2.4Å). (**F**) Redocking of cholesterol ligand on ABCG5/G8 (RMSD 2.5Å, present study). See details in Materials and Methods.

Supplemental Figure 4. Aromatic amino acid pattern in non-ABCG lipid transporters.
Conserved aromatic residues along the putative (or observed) sterol-binding/interacting TMH
(cartoon presentation) show a similar pattern of phenylalanine highway as observed in ABCG
transporters. SIDT2 (**A**, yellow), SIDT1 (**B**, dark blue), ABCA1 (**C**, light blue), ABCA7 (**D**, pink),
PTCH1 (**E**, orange) and NPC1 (**F**, green). The aromatic amino acids: phenylalanine, tyrosine and
tryptophan are highlighted in red sticks. Only the most central and luminal helices are modelled in
detail.

Supplemental Figure 5. Cavities and binding regions in ABCG structures. (A) *Left.* ABCG5/G8 (PDB ID: 5DO7) aligned with cryo-EM structure (PDB ID: 7R8D). *Right.* Comparison of cholesterol docking pose (light blue) with the cholesterol (green) in the cryo-EM structure [28]. **(B)** Volumes of transmembrane domain (TMD) cavities found in the of ATP-bound ABCG1 (yellow, PDB ID: 7R8E) and ABCG5/G8 (blue) calculated by MOLE2.5. *Left*, two cavities in ABCG1 found near the interface between the TMD and extracellular domain (ECD). These cavities ($1629Å^3$ each) curve over the central canal of the protein. *Right*, one large cavity found spanning throughout the TMD. This cavity represents a conglomeration of sphere-shaped cavities connected by thin bridging regions. Relevant cavities in the central canal of the protein are represented as spherical and estimated to be near the size of $\sim200Å^3$. *Bottom*, the volume of a cholesterol molecule is $627Å^3$.