1　# Direct and indirect phenotypic effects on sociability indicate potential to evolve
2

David N. Fisher

School of Biological Sciences, University of Aberdeen, King's College, Aberdeen, AB243FX, UK

david.fisher@abdn.ac.uk

Abstract

The decision to leave or join a group is important as group size influences many aspects of
organisms' lives and their fitness. This tendency to socialise with others, sociability, should be
influenced by genes carried by focal individuals (direct genetic effects) and by genes in partner
individuals (indirect genetic effects), indicating the trait's evolution could be slower or faster than
expected. However, estimating these genetic parameters is difficult. Here, in a laboratory population
of the cockroach *Blaptica dubia*, I estimate phenotypic parameters for sociability: repeatability (R)
and repeatable influence (RI), that indicate whether direct and indirect genetic effects respectively
are likely. I also estimate the interaction coefficient (Ψ), which quantifies how strongly a partner's
trait influences the phenotype of the focal individual and is key in models for the evolution of
interacting phenotypes. Focal individuals were somewhat repeatable for sociability across a three-
week period (R = 0.083), and partners also had marginally consistent effects on focal sociability (RI =
0.055). The interaction coefficient was non-zero, although in opposite sign for the sexes; males
preferred to associate with larger individuals (Ψ_{male} = -0.129) while females preferred to associate
with smaller individuals (Ψ_{female} = 0.068). Individual sociability was consistent between dyadic trials
and in social networks of groups. These results provide phenotypic evidence that direct and indirect
genetic effects have limited influence on sociability, with perhaps most evolutionary potential
stemming from heritable effects of the body mass of partners. Sex-specific interaction coefficients
may produce sexual conflict and the evolution of sexual dimorphism in social behaviour.

Key words: cockroach, group size, heritability, indirect genetic effects, interaction coefficient,
personality, repeatability, sociability, social network

Introduction
30

Many animals form groups and aggregations to find food, avoid predators, and to be buffered from
environmental stressors (Krause & Ruxton, 2002). Individual sociability is therefore an important
trait that can influence access to resources, mating opportunities, predators, and disease (Gartland
et al., 2022). This importance means it is often linked to fitness. Further, in aggregate individual
sociability determines group size, which in its own right can have influences on individuals' fitness
components (Silk, 2007). These links with fitness imply sociability is frequently under selection, and
therefore would be expected to evolve if heritable. Predicting how individual sociability, and
therefore also group size, will evolve requires us to estimate the genetic variance underpinning the
trait i.e. its heritability (Scott *et al.*, 2018). Typically, when estimating the heritability of a trait we
consider its direct additive genetic variance i.e., how much variance among individuals in their own
genes relates to variance in their phenotypes (hereafter direct genetic effects, "DGEs"). However,
alongside its own social tendencies an individual's sociability will likely depend on the traits of others

in the groups it may join. For example, a normally sociable individual may be less willing to join a
group with particularly aggressive individuals. As the traits of others will be at least partly influenced
by genes, the heritable variation in sociability is likely to stem not only from DGEs, but also indirect
genetic effects (IGEs), where genes in an interacting individual influence the focal individual's trait
(Griffing, 1967; Moore *et al.*, 1997a). The presence of IGEs (and their covariance with DGEs) can
accelerate evolutionary change, retard it, remove it completely, or even reverse it (Moore *et al.*,
1997a; Wolf *et al.*, 1998), potentially leading to non-linear responses to selection (Trubenová *et al.*,
2015), responses to selection in the opposite direction to that of direct selection (Bijma & Wade,
2008; Fisher & Pruitt, 2019) and even maladaptation (Fisher & McAdam, 2019; McGlothlin & Fisher,
2021). Indirect genetic effects are widely appreciated in animal and plant breeding for their ability to
prevent the evolution of higher yields (Muir, 2005; Ellen *et al.*, 2014; Costa e Silva *et al.*, 2017), and
are becoming increasingly well appreciated in evolutionary ecology (Baud *et al.*, 2022). If we want to
understand how evolution shapes variation in sociability, the diversity of group sizes in nature, and
how these traits might evolve in the future, we need to estimate how important both DGEs and IGEs
are for individual sociability.

Despite the clear need to measure DGEs and IGEs on sociability, estimates of DGEs are quite rare
(Lea *et al.*, 2010; Brent *et al.*, 2013; Staes *et al.*, 2016; Knoll *et al.*, 2018; Scott *et al.*, 2018), and
estimates of IGEs are completely absent (although Lea *et al.* did estimate DGEs for the tendency to
receive interactions in a social network of marmots, which should be very similar to IGEs for initiating
interactions). This can be partially attributed to two factors: 1) Experimental design to quantify
individual sociability and how it is influenced by both direct and indirect effects can be difficult
(Gartland *et al.*, 2022) and 2) Estimating DGEs and IGEs in any context requires large amounts of
both phenotypic data and information on genetic relatedness (Moore *et al.*, 1997a; Bijma, 2014;
Kruuk & Wilson, 2018). While 1) can be solved with appropriate experimental design, solving 2) can
be logistically challenging. One partial (and temporary) solution is to estimate parameters that
represent DGEs and IGEs at the phenotypic level, which does not require data on genetic relatedness
and may also require less data overall as phenotypic variances are typically larger than genetic
variances. Although not ideal, these parameters can still give insight into the evolutionary potential
of the trait of interest as the relative magnitude of phenotypic and genetic variances (and
covariances) are normally aligned (Hadfield *et al.*, 2007; Dochtermann, 2011; Dochtermann *et al.*,
2015).

For DGEs, the phenotypic parameter that (in most cases) sets the upper limit for heritability is
repeatability (R, but see: Dohm, 2002). Repeatability is defined as the portion of phenotypic variance
attributable to among individual differences (V_I; Nakagawa & Schielzeth, 2010). This parameter can
be decomposed into additive genetic variance and permanent environmental variance ($V_I = V_A + V_{PE}$); where for behavioural traits on average 52% of V_I stems from V_A (Dochtermann *et al.*, 2015).
V_{PE}); where for behavioural traits on average 52% of V_I stems from V_A (Dochtermann *et al.*, 2015).
We can therefore think of R as a phenotypic proxy for DGEs (as well as providing useful information
about the relative balance between among- and within-individual variation in the population).
Regarding IGEs, an analogous phenotypic equivalent in dyadic interactions would be the variance
attributed to the identity of the interaction partner (V_S). We could then calculate "repeatable
influence" (RI) as the portion of phenotypic variance in the focal individual's trait attributable to the
among partner differences. For interactions with multiple partners, V_S is multiplied by the average
number of interaction partners (Bijma, 2011), and so can lead to the variance attributable to social
interactions being greater than the phenotypic variance (Bijma *et al.*, 2007; see: Ellen *et al.*, 2014 for
empirical examples of the total hertiable variation of a trait being greater than the phenotypic
variance, due to IGEs), which makes it less directly relatable to R. Estimating both R and RI should

give us a good indication of the *relative* contribution of DGEs and IGEs to a trait, while also giving
some indication of the likely *absolute* magnitude of these parameters.

Alternatively to estimating DGEs and IGEs, a parameter widely used to infer their importance of
social interactions in evolution is the interaction coefficient (Ψ; Moore *et al.*, 1997a; Bailey &
Desjonquères, 2022). This term is the coefficient from a regression of the focal individual's
phenotype on an interacting individual's trait. It therefore does not require data on genetic
relatedness. The Ψ term is key in the "trait-based" approach for understanding the role of social
interactions in evolution, as opposed to the "variance-based" approach, which relies on DGEs, IGEs,
and their covariance (McGlothlin & Brodie, 2009). Ψ can alter the direction and steepness of
evolutionary trajectories, lead to feedback between interacting traits, and result in non-linear
change (Bailey & Desjonquères, 2022). Additionally, Ψ can be converted into a direct-indirect
covariance if the genetic variances of the traits of interest are known (McGlothlin & Brodie, 2009).
We can therefore think of Ψ as both an important evolutionary parameter in its own right and as a
phenotypic indicator of the likely magnitude of key genetic covariances. Together, R, RI, and Ψ give
us useful gauges of the likely importance of DGEs and IGEs for a trait's evolution, and so estimating
them for sociability will give us a reasonable indicator for how this trait, and therefore group size,
may evolve in the absence of genetic information.

Here, I estimated R, RI, and Ψ for sociability in the gregarious cockroach *Blaptica dubia* (Blattodea:
Blaberidae). This is a communally living species who form aggregations in refuges to access
resources, avoid predators such as ants, and to buffer environmental perturbations (Grandcolas,
1998) – hence their sociability is an important trait for their survival and fitness. Further, cockroach
species are often sources of zoonoses (Zarchi & Vatani, 2009; Patel *et al.*, 2022), and so a better
understanding of the evolution of their aggregation formation could inform how the disease risk to
humans they pose changes in the future under selection pressures such as changing climates and
different management strategies. I measured sociability in dyadic trials and validated this assay in a
group context using social networks of up to 24 individuals. I assayed individuals in the dyadic trials
repeatedly to allow me to estimate consistency in sociability (R) and to isolate the consistent effect
of a partner individual on the focal (RI). I also tested how a trait of the interaction partner influences
the focal individual's sociability to quantify Ψ. I used body mass as the trait in interacting individuals
as it is typically heritable; Clark and Moore (1995) estimated the full-sibling heritability (likely to be
an overestimate) of body mass in the Madagascar hissing cockroach (*Gramphadorhina portentosa*),
which like *B. dubia* is in the Blaberidae family, as 0.93, while Moore et al. (2004) estimated the
heritability of pronotum width in the speckled cockroach (*Nauphoeta cinerea*, also a Blaberid) as
0.62. Therefore, a clear estimate of Ψ for body mass would indicate social interactions are likely to
be important for the evolution of sociability. I predicted that 1) there would be a correlation
between the measures of sociability in the dyadic and group context, 2) that sociability will be
repeatable, 3) that sociability will be repeatably influenced by the identity of the partner individual,
and 4) that individuals will prefer to interact with larger partners (as smaller values in my sociability
assay indicate more sociable, this means $\Psi < 0$) as they represent better protection from predators
and the elements.

Methods

Experimental animals

Blaptica dubia is a quite large (up to 45 mm in length) sexually dimorphic blaberid cockroach (Wu,
2013). They typically live in aggregations at high temperature and humidity in central and south
America (Alamer & Hoffmann, 2014), consuming vegetative matter, and are ovoviviparous. They are
described as "gregarious" (Grandcolas, 1998) or "communal" as individuals of the same generation
cohabit (without shared parental care; Bewick *et al.*, 2017). I purchased an initial colony of *B. dubia*
online in March 2021. I maintained them at the University of XXX [removed to preserve anonymity]
at 28°C, 50% humidity, with a 50:50 light:dark cycle. I provided them with cardboard egg trays for
shelter, carrot for hydration, and Sainsbury's Complete Nutrition Adult Small Dog Dry Dog Food
(approx. nutritional composition = 1527 kJ energy, 24g protein, 12g fat per 100g) for nutrition.
Mortality was very low at all life stages (0.31% of the adults in the colony died per half week)
indicating the colony was healthy. I moved newly born nymphs every few days to a container of
dimensions 610 x 402 x 315 mm of similar aged individuals (density ranged from a few hundred of
the earliest instars to 10-80 of later instars) and maintained them in mixed groups until adulthood
(seven instars which takes approx. 250 days at this temperature; Wu, 2013). Upon reaching
adulthood I moved them to either single sex groups (again in containers of 610 x 402 x 315 mm) or in
small groups of two males and four to eight females in a container of dimensions 340 x 200 x 125
mm for breeding to maintain the stock population. For this experiment I selected 48 unmated males
and 48 unmated females from the single-sex adult groups. Individuals were all more than five days
old, and so presumed to be sexually mature (Hintze-Podufal & Vetter, 1996). I transferred each
individual to a clear plastic box (79 x 47 x 22 mm) labelled with its unique ID to allow individual
recognition. I gave individuals a small piece of carrot for hydration which was replaced weekly.

Data collection

I tested individuals in two blocks of 48, treating all individuals in each block once as a focal individual
and once as a partner for a member of the same sex over two days. This means that in the first two
days 24 males and 24 females were each assayed for sociability once and each acted as a partner
individual once. On days three and four I repeated this with a second block of 24 males and 24
females. In this way individuals only ever acted as focal or partner individuals with members of the
same sex in the same block (either first or second) and were each assayed for sociability and acted
as a partner once per week. I repeated this for three weeks, so each individual was assayed up to
three times as a focal individual as acted as a partner up to three times. Some individuals received
fewer than three trials if they died (n = 5 males and 0 females), in which case I replaced them with a
member of the same sex from the stock population (who did *not* inherit the same ID and was
therefore another unique individual). Individuals might also record fewer than three measures for
sociability if the mesh was breached by either the partner or the focal before the trial began (11
females and eight males recorded one measure, 48 females and 42 males recorded two measures,
30 females and 54 males recorded three measures).

I assayed sociability in medium sized plastic boxes (200 x 100 x 70 mm) where I glued a fine
polypropylene mesh (mesh size 0.6 x 0.6 mm, Micromesh, Haxnicks) across the interior 50 mm from
one end. This creates an arena with a small compartment (50 x 100 x 70 mm) and a large
compartment (150 x 100 x 70 mm) separated by the mesh (Fig. 1A). Separating by mesh was
necessary to prevent a partner individual imposing close proximity on the focal individual by
constantly following or attempting to dominate it (Clark *et al.*, 1995), and therefore my assay
captures the focal individual's willingness to socialise, rather than the partner's (Gartland *et al.*,
2022). For the first block I randomly placed 12 females and 12 males alone, each into their own
plastic box, in the large compartment. These were the focal individuals. I then randomly placed an

179 individual of the same sex into the small compartment; these were the partner individuals. I used
180 individuals of the same sex to ensure I was measuring sociability rather than willingness to mate. I
181 then placed these 24 arenas into four large plastic boxes (six in each) which I placed underneath a
182 video camera (ABUS IP video surveillance 8MPx mini tube camera), so that each video camera
183 recorded six arenas simultaneously. I maintained the room the video recordings occurred in at 20-
184 22°C using portable heaters, while I used a thermometer to record the temperature at the start and
185 end of each trial. I was not able to control or monitor humidity during the trials. Once all arenas
186 were in position and cameras focused, I started the recording and left the room. The lights
187 automatically switched off after 40 minutes, and so the trial began 40 minutes after I left the room,
188 in darkness, which is when *B. dubia* is active (Bouchebti *et al.*, 2022). I returned two hours after
189 leaving to end the trial, meaning the trials lasted 80 minutes. In darkness the cameras automatically
190 switch to infra-red filming using infra-red LEDs.

191 For each trial, every ten minutes I recorded the proximity of the focal individual (in the larger
192 compartment) to the mesh that separated it from the partner individual (Fig. 1A), giving a maximum
193 of eight measures per trial. The distance of an individual to a conspecific in this manner is often used
194 to measure sociability (reviewed in: Gartland *et al.*, 2022) and by using the distance to the mesh
195 rather than the partner I have a measure solely under the influence of the focal individual; the
196 partner cannot directly influence the distance by moving closer or further away. If the focal was sat
197 directly on the mesh (perpendicular to the floor) I recorded a distance of zero, otherwise I used the
198 hexagons on the bottom of the box to record how far the focal individual's head was from the mesh
199 (Fig. 1A). Smaller values mean a focal individual closer to the partner individual which indicates
200 higher sociability. Individuals were in some cases able to bypass the mesh (this occurred 95 times
201 before the lights went out and 33 times after they did out of 288 trials, the 33 breaches after lights
202 out are still included in the analyses with only the measurements before the breach used, see Data
203 analysis). I used the video recordings to determine when this happened and stopped recording data
204 from the video as soon as either individual bypassed the mesh. If either individual bypassed the
205 mesh in the 40 minutes the lights were on before the trial started then I recorded no data from that
206 trial. To avoid mixing individuals up at the end of the trial I dotted either the partner or the focal
207 with white paint (Edding Extra-fine paint markers). At the end of the trial, I returned all individuals to
208 their unique boxes. I then weighed all partner individuals to the nearest 0.01 g (Fisherbrand
209 Analytical Balances, readability 0.0001 g). In the first two days I also weighed the focal individual, but
210 since the correlation between an individual's mass as a focal and its mass as a partner either the
211 following or preceding day was 0.994 (Pearson correlation, t_{38} = 56.894, p < 0.001) I stopped doing
212 this to save time. Instead, I entered the mass of the focal individual as its mass as a partner
213 individual recorded the same week (always within one day). As described above, each individual in
214 the two blocks was assayed once as a focal and acted once as a partner for another individual of the
215 same sex in that block per week, and this was repeated for three weeks.

Figure 1. Pictures of experimental set-up (both XX [removed to preserve anonymity]). A. Assay
for sociability. The position of the focal individual (on the right in the larger compartment, no
white dot) in relation to the mesh is recorded every ten minutes to assess willingness to
socialise. This individual would record a score of two. B. Social network trials. Marked individuals
(here showing green-blue, red-white, white-blue, and green-white, starting at the top left and
moving clockwise) can chose among four equal shelters (the cardboard egg trays taped to the
sides of the box). Co-occurring at a shelter with the same individual regularly indicates a social
association. A close up of red-white is shown in the upper right corner.

After the third trial I aggregated individuals into four groups of 21-24; all the individuals of the same
sex from the same block were together, with groups having fewer than 24 individuals if any died (I
did not replace individuals that died with stock individuals as I was only interested in the social
network position of those with a known sociability from the dyadic trials). I gave each individual a
unique combination of two colours (red, green, blue, white, gold) on their wing cases using paint
pens (Edding Extra-fine paint markers; Fig. 1B) which allowed me to track them individually
(combinations were repeated between groups i.e., red-blue featured in each of the four groups). I
then placed each group into new plastic boxes (340 x 200 x 125 mm) along with four shelters made
from cardboard egg tray (approx. 100 x 120 mm), each placed vertically at each corner on a long side
(Fig. 1B). Shelters were taped to the walls of the box, creating clear space between both the shelter
opposite it (on the opposite long side) and next to it (on the same long side). I placed 2 g dog food
and 10 g carrot in the centre of each box. Each shelter was large enough to accommodate many but
not all of the individuals, and the number of shelters was considerably less than the number of
individuals. Therefore, the formation of aggregations in shelters was enforced, but individuals could
move between shelters and therefore could exert some influence on who they co-habited with.
Regularly after placing the individuals into these groups (after 3, 10, 14, 18, and 21 days) I recorded
which individuals were using the same shelter. Individuals who could not be identified were
recorded as such but they were not used to build the networks. Collecting data in this way gives a
group-by-individual matrix analogous to those generated by observing flocks of birds or herds of
ungulates in the wild, and further is similar to methods than have been used to generate social
networks in forked fungus beetles *(Bolitotherus cornutus*; Formica *et al.*, 2012, 2016, 2020) and
maritime earwigs *(Anisolabis maritima*; Vipperman, 2021). While a single incidence of sharing a
shelter could be due to chance, by aggregating these observations I can infer consistent social
associations. When recording these data, I also updated any paint markings that were starting to

wear, maintaining individually-recognisable marks for the duration of the experiment, and replaced
carrot and dog food as necessary.

Data analysis

All analyses were conducted in R (version 4.1.3; R Development Core Team, 2016). To analyse
sociability, I summed each individuals' distances from the mesh across the 1-8 records per trial and
entered that as a response variable in a generalised linear mixed effects model using *glmmTMB*
(Brooks *et al.*, 2017). To account for the different number of measures contributing to this sum (if
individuals "breached" the barrier during the trial) I included an offset of the log of the number of
records the individual recorded from the trial and used a Poisson error distribution and a log link
function. While the response variable entered into the model is still the sum of the distances (and
hence an integer), this approach effectively models the mean distance the individual is from the
mesh (sum / n. trials) but is preferable from directly using this variable as it can be used with a
Poisson error distribution, which requires integers and so is incompatible with the mean distance
(the residuals are also greatly improved, see Fig. S1). I included fixed effects of the temperature in
the room (scaled to a mean of zero and a standard deviation of one), the sex of the individual (and
therefore also its partner), the body mass of the focal individual and the body mass of the partner,
both scaled to a mean of zero and a standard deviation of one, and the interaction between both
focal and partner (scaled) body mass and sex. The effect of the partner mass is key for testing
prediction 4 as its coefficient is our (unstandardised) estimate of Ψ, while the interaction with sex
tests whether this differs between the sexes. I included random effects of individual ID, partner ID,
and date, to estimate the variance among focal individuals, partner individuals, and dates
respectively. To estimate R for sociability (testing prediction 2) I extracted the model intercept, the
among-focal individual variance, and the sum of all variance components, and entered them into the
'QGicc' function in the package *QGglmm* (de Villemereuil *et al.*, 2016), using the 'model =
"Poisson.log"' setting. This calculates R for sociability on the original scale as opposed to the latent
scale (Nakagawa & Schielzeth, 2010; de Villemereuil *et al.*, 2016); the former is necessary to
compare to estimates of R from traits analysed assuming a Gaussian distribution. I repeated this
with the among-partner individual variance instead of the among-focal individual variance to obtain
the estimate of RI (testing prediction 3). Alongside the magnitudes of R, I demonstrated the
importance of accounting for differences among individuals in sociability by comparing the AIC of
the model described above to a model identical except that the random effect of focal individual
was removed. I did the same for RI i.e., comparing models with and without the partner ID term (the
models were otherwise identical to the one described above). To determine the clarity of fixed
effects I used the 'Anova' function in the package *car* (Fox & Weisberg, 2019) with a type three sum
of squares to generate p values (see: Dushoff *et al.*, 2019 for a discussion on the use of "clarity" over
"significance").

After finding a clear interaction between body mass and sex (see Results), I wished to obtain sex-
specific estimates of Ψ that were standardised to facilitate comparisons across studies (note this was
a decision made after viewing the initial results and so should be interpreted more cautiously). To do
this I refitted the above model to the sexes separately (minus the fixed effect of sex and its
interactions with focal and partner body mass), this time with the mean distance the focal individual
was from the mesh, scaled to a mean of zero and a standard deviation of one, as the response
variable, with no offset and assuming a Gaussian error distribution with a normal link function. Using
this transformed variable generates estimates of Ψ that are comparable across traits and studies
(Bailey & Desjonquères, 2022).

To generate a social network per group I used each groups' group-by-individual matrix, which
contains records of which individual was in which shelter at each time point. From this I created four
networks where individuals were linked with their relative association strengths, which is undirected
i.e., individual A's interaction with individual B is equal to individual B's interaction with individual A.
I calculated relative associations strengths as the simple ratio index (Cairns & Schwager, 1987), using
the package *asnipe* (Farine, 2013). This index is the count of all times individuals shared a shelter,
divided by the number of occasions both individuals were recorded (this could be less than five if an
individual died during this phase of the study), and so indicates the relative strength of the
association between any two individuals. A score of one indicates two individuals who were always
seen sharing a shelter, and zero two individuals who never shared the same shelter. I summed each
individual's association scores to gives that individual's 'strength', a measure of network centrality
that captures an individual's overall engagement in social interactions and in this case is analogous
to the average group size an individual was found in.

To test whether sociability as determined by the dyadic assays correlated with sociability in the
social network (prediction 1), I followed the suggestions of Hadfield et al. (2010) in the guide from
Houslay and Wilson (2017) to estimate the among individual correlation between the two traits (see
also: Dingemanse *et al.*, 2012; Dingemanse & Dochtermann, 2013). This approach excludes the
residual variance from the correlation, specifically addressing our question of interest (are
individuals that are more sociable in the dyadic assay more sociable in the social network). To do this
I fitted a bivariate mixed-effects model in MCMCglmm (Hadfield, 2010). The response variables were
each of an individual's sum of locations from the dyadic trials, and its strength as quantified in the
social network. As there is only one value of the latter it is repeated each time an individual records
a sociability score i.e., 1-3 times. I included the fixed effect of the log of the number of observations
and modified the prior to constrain the relationship between this fixed effect and the sum of
locations or network strength to 1 or 0 respectively (by setting the coefficients to 1 and 0
respectively and setting both variances as 1×10^{-9}). This approach is equivalent to fitting number of
observations as an offset for the sum of locations, and as having no relationship at all with network
strength. The random effect was a 2 x 2 covariance matrix estimating the among individual variance
in each trait and the among-individual covariance between them (our parameter of interest). I
allowed the residual variance for sociability in the dyadic assay to be non-zero (as there are multiple
measures on individuals) while I fixed it at 0.0001 for network strength (as there is only a single
measure it does not vary within individuals), and I fixed the residual covariance at zero. I set a
Poisson error distribution for sociability in the dyadic assay and a Gaussian error distribution for
network strength. I used 550,000 iterations, with the first 50,000 discarded and 1 in 100 of each
subsequent iteration retained. I confirmed the model had converged by running three chains and
calculating the Gelman and Rubin convergence diagnostic (Gelman & Rubin, 1992; point estimates
were all 1.01 or lower), as well as assessing the trace plots. I calculated the among individual
correlation between the two measures of sociability as the among-individual covariance between
the two traits divided by the square root of the product of their two among-individual variances and
extracted the mode and 95% credible intervals from the resulting posterior distribution.

Results

I collected 193 observations of sociability across 92 individuals in dyadic trials, and quantified the
social network position of each of these individuals (Fig. 2). Individuals that were more sociable in

the pairwise trials had higher strength in the social network trials (Among-individual correlation
between sociability and network strength, posterior distribution mode = -0.758, 95% credible
intervals = -0.948 to -0.265), confirming prediction 1 and validating the use of the dyadic trials to
assay sociability. Full model results are shown in Table S1.

Individuals are weakly repeatable in how sociable they are, with an R of 0.083 (prediction 2), while
the ΔAIC of model with vs without the random effect of focal individual ID was 767. Individuals
exerted a small amount of repeatable influence on the sociability of their partners, with an RI of
0.055 (ΔAIC = 774; prediction 3). The body mass of the partner individual influenced how sociable
the focal individual was (prediction 4), with males being more sociable with larger individuals and
females being more sociable with smaller individuals (Fig. 3; main effect β = 1.350, se = 0.463, χ2 =
8.493, p = 0.004, interaction β = -2.833, se = 0.686, χ2 = 17.038, p < 0.001). Full model results are
presented in Table 1. Given this clear interaction, I decided post-hoc to fit separate models to each
sex to generate sex-specific and standardised estimates of Ψ, giving Ψ_{female} = 0.068 (se = 0.103) and
Ψ_{male} = -0.129 (se = 0.107; recall that lower scores in the pairwise trial indicate higher sociability as
they represent shorter distances from the partner individual). There are therefore phenotypic
indicators that there are IGEs on individual sociability, but the direction of the effect is opposite in
sign for the sexes and both are quite near zero.

Table 1. Full model output for analysis of sociability in dyadic trials. Females were set as the default
sex and so the effect of sex is the contrast between males and females.

Variable	Estimate	Standard Error	Chi-squared	P value
Intercept	-0.476	0.584	0.665	0.415
Sex (Male contrast)	-1.787	0.986	3.283	0.070
Body mass of focal	-0.940	0.450	4.359	0.037
Body mass of partner	1.350	0.463	8.493	0.004
Temperature	0.168	0.178	0.897	0.343
Sex : Body mass of focal	-0.345	0.586	0.347	0.556
Sex : Body mass of partner	-2.833	0.686	17.038	0.000

	Random effect	Variance
	Individual	2.503
	Partner	2.130
	Date	0.281

Figure 2. Social networks of *Blaptica dubia* individuals for each group (females on top row, males on bottom row, node size indicates sociability in pairwise trial [larger = more sociable], thickness of lines indicates strength of association). I have removed edge weights below 0.12 to reduce visual clutter.

Discussion

I found that the measure of sociability in the dyadic trials was correlated among-individuals with a measure of centrality from a social network, validating the use of the dyadic trials for assaying sociability. Sociability in the dyadic trials was repeatable, and partners had a repeatable influence on the sociability of others, but both values were small. This implies that the direct and indirect genetic variance in the trait is likely to be low, and hence selection on this trait will be converted into limited evolutionary change. I also found the body mass of the partner influenced the sociability of the focal individual, but in sex-specific ways; males were more sociable with large individuals, while females were sociable with small individuals. As body mass is typically heritable this suggests that there is some evolutionary potential in sociability through the effect of body size; if body mass evolves, so will sociability, but in opposite directions for males and females.

Figure 3. The body mass of the partner individual influences the sociability of the focal individual, with males (blue) preferring to be near heavier individuals, while females (orange) preferring to be nearer lighter individuals. Points are individual scores while lines indicate the mean effect estimated by the plotting function. Grey areas indicate the 95% confidence intervals around the mean.

Finding sociability had non-zero repeatability is to be somewhat expected as behaviours typically are repeatable (Bell *et al.*, 2009; Holtmann *et al.*, 2017). Further, a low repeatability might be expected given social behaviours often depend on the phenotype of interaction partners which can be expected to vary substantially in short periods of time (Holtmann *et al.*, 2017). Further, I found relatively little variance attributed to the identity of the partner individual. To explore this further, I re-ran the model with the mass of the partner (and its interaction with sex) removed. This reduced model gave a slightly higher estimate of V_S (2.19 vs. 2.13). Therefore, while body mass exerts some consistent effect on other individuals, other traits also contribute. In any case, most of the variation in sociability was not explained by the variance components or fixed effects, indicating sociability is highly labile.

Given that, on average, 52% of the among-individual variance of behaviours stems from direct additive genetic variance (Dochtermann *et al.*, 2015), we can expect sociability to have a low heritability even when the indirect genetic variance is included. A low heritability means selection would be translated into a small amount of evolutionary change, and so sociability and therefore mean group size may show limited response to direct selection. In contrast, Dochtermann *et al.* (2019) used a meta-analysis across taxa to infer that social behaviours had heritabilities typical of behavioural traits; similar to the overall mean of 0.235. This estimate however only considered

direct heritability and not indirect genetic effects, the total heritability may well be even higher (or

lower, depending on whether the direct-indirect genetic covariance is positive or negative; Bijma,

2011). Estimating the direct and indirect heritability of sociability and obtaining ecologically relevant

estimates of selection (or relationships between sociability and fitness components such as survival

or longevity; Blumstein *et al.*, 2018; Brodin *et al.*, 2019; Montero *et al.*, 2020) are logical next steps

to better understand the microevolution of this trait (see also the artificial selection experiment of

Scott *et al.*, 2022 who successful increased sociability in *Drosophila melanogaster* over 25

generations).

Alongside the repeatability of sociability, I found that how sociable an individual was depended on

its own mass and the mass of the mass of the partner individual. While heavier individuals of both

sexes were more sociable, females preferred to be near smaller females, while males preferred to be

near larger males i.e., Ψ was sex specific. While I predicted an effect of partner mass, I had not

predicted a sex-specific effect. Females are larger than males and require protein for egg production

which males do not (Maklakov *et al.*, 2008; Jensen *et al.*, 2015), and so females may be in more

intense competition for resources with each other than males are. This competition could lead to

them preferring to associate with smaller individuals who are presumably less competitive. In

contrast, males may prefer larger individuals as they offer better protection from predators, more

protection from desiccation, and possibly if larger males attract females, the so called "hotshot"

effect (Beehler & Foster, 1988). Alternatively, both sexes may be seeking mating partners (I used

unmated individuals), which are always smaller or larger than themselves for females and males

respectively (see how the masses of the sexes do not overlap in Fig. 3). This would lead to females

preferring to be near smaller females who are perhaps harder to distinguish from males, and vice

versa for males (suggested by Han *et al.*, 2016, although they found no effect of partner body size on

same-sex behaviour in water striders *Gerris lacustris*). While we would expect chemical

communication to be important for mate choice cockroaches (Schal *et al.*, 1984; as well as for other

social interactions: Moore, 1997; Moore *et al.*, 1997b), it is possible individuals use both chemical

cues and morphological traits when searching for a partner. Testing these ideas, and the fitness

consequences for both males and females for associating with large and small individuals ("social

selection"; Wolf *et al.*, 1999; e.g.: Santostefano *et al.*, 2019; Fisher *et al.*, 2021), represent key next

steps. Furthermore, it is worth highlighting that body mass represents only one trait that influences

sociability; there may well be other morphological, behavioural, or chemical traits of partners that

affect focal individual behaviour. While these are accounted for in the estimate of V_S, identifying the

causal traits is a useful step for understanding the mechanisms underpinning social interactions.

Indirect genetic effects can fundamentally alter the direction and magnitude of evolutionary change,

and so finding opposing estimates of Ψ for the sexes implies that sociability in the sexes could follow

quite different evolutionary trajectories. Whether they will do so or not depends on the genetic

variance underpinning mass (which is likely to be non-zero, see Introduction) and sociability and the

inter-sex genetic correlation for sociability (McGlothlin & Brodie, 2009). Further, given preferences

differ, selection on partner choice may well differ between the sexes (Parker, 1979; "sexually

discordant selection"; Westneat & Sih, 2009). Such selection can be expected when sexual

dimorphism cannot evolve, possibly due to inter-sex correlations typically being large and positive

(Poissant *et al.*, 2010). In the case of *B. dubia*, any quantitative predictions at this stage would be

premature given the number of assumptions I would be required to make, but it is interesting that

same-sex social interactions potentially facilitate sexual conflict thanks to estimates of Ψ which are

opposite for the sexes. Results here show that if population mean body mass increases, male

sociability will increase, while female sociability will decrease. In the short term at least (before

female preferences change) reduced sociability could reduce female mating rate, influencing the

costs and benefits of polyandry such as male harassment, sexual infection transmission, direct and
indirect benefits from mating with multiple males. In general, however, the estimates of Ψ (which
typically range from -1 to 1; Bailey & Desjonquères, 2022) in this study are quite near zero, indicating
only modest deviations from a situation where individuals do not impact each other's traits through
social interactions (this is quite common for estimates of Ψ between different traits; Bailey &
Desjonquères, 2022).

My results add to increasing evidence that Ψ varies among- (Kent *et al.*, 2008; Bleakley & Brodie III,
2009; Bailey & Zuk, 2012; Edenbrow *et al.*, 2017; Marie-Orleach *et al.*, 2017; Culumber *et al.*, 2018;
Kraft *et al.*, 2018) and within-populations (Edenbrow *et al.*, 2017; Signor *et al.*, 2017; Han *et al.*,
2018). Ψ itself can evolve (Chenoweth *et al.*, 2010; Bailey & Zuk, 2012; Rebar *et al.*, 2020), and the
evolution of Ψ can both increase or decrease the speed of evolutionary change (Kazancioğlu *et al.*,
2012). It is therefore clearly important to study this parameter in an evolutionary context to better
understand the evolution of interacting phenotypes (Bailey & Desjonquères, 2022). This is especially
true when the majority of the evolution potential of a trait may stem not from genetic variance in
the trait (which is what is typically assumed), but from associations with other heritable traits (such
as body mass in this study). Additionally, it is especially important to estimate Ψ in systems where is
expected to be large. Bailey & Desjonquères (2022) highlight that large and positive values of Ψ
occur when individuals interact through the same trait, suggesting a positive feedback loop in trait
expression. This would mean small initial evolved changes in a trait would lead to increased
expression of the same trait in the partner, and so the trait mean would increase rapidly (Moore *et*
al., 1997a). High and positive values of Ψ are observed in shoaling fish (*Poecilia reticulata*; Bleakley &
Brodie III, 2009; Marie-Orleach *et al.*, 2017; *Gambusia holbrooki*; Kraft *et al.*, 2018), fruit flies
(*Drosophila melanogaster* and *D. simulans*; Bailey & Hoskins, 2014; Signor *et al.*, 2017) and burying
beetles (*Nicrophorus vespilloides*; Rebar *et al.*, 2020). More estimates of Ψ need to be accumulated
before more general patterns in when Ψ strongly influences evolutionary potential can be identified
(Bailey & Desjonquères, 2022).

The sociability of an individual estimated through repeated pairwise trials over three weeks was
related to the individual's centrality, in terms of its overall number and strength of associations, in a
social network formed over 21 days. This result indicates that my assay for sociability accurately
captures a facet of individual social behaviour, and this social behaviour is trait of an individual that
is consistent across time and across contexts, and hence could be associated with lifetime
reproductive success (Kluen & Brommer, 2013; although this association cannot be very strong as
the trait's repeatability is low). What maintains consistent among-individual differences in sociability
and social network position are open questions (Wilson *et al.*, 2012; Wilson & Krause, 2014; Gartland
et al., 2022). If the behaviour is indeed heritable, then different levels of sociability should give
similar fitness on average, as otherwise selection would remove the variation in sociability from the
population. Different levels of sociability may therefore represent different strategies that bring
both benefits (e.g., a higher sociability decreases water loss) and costs (e.g., a higher sociability
decreases access to resources). Furthermore, switching between strategies must impose costs in
some way so that individuals cannot be completely plastic (Dall *et al.*, 2004; Snell-Rood, 2013). A
second, not mutually exclusive, mechanism that could maintain variation in social strategies is if
selection on the strategy is negative frequency dependent (Bergmüller & Taborsky, 2010). Better
understanding of patterns of selection for this and similar traits is therefore key for predicting
evolutionary responses in natural populations, and, ultimately, population dynamics and viability.

In summary, sociability in *B. dubia* shows a small degree of repeatability, some consistent influence
from the identity of a partner, and is correlated among-individuals between trials in pairs and trials

in groups. Males preferred to associate with larger individuals while females preferred to associate
with smaller individuals. The latter result suggests that the evolution of sociability, and therefore the
evolution of group size, may fundamentally depend on evolutionary change in body mass, and could
drive sexual dimorphism in social behaviour. These sex-specific estimates of Ψ will be important for
informing our models predicting microevolutionary change and for understanding sexual conflict.
Future work will need to assess the fitness consequences of social behaviour and identifying the
factors that predict patterns of social interactions in various more ecologically relevant settings.

Acknowledgements
504 I would like to thank Keith Lockhart for his invaluable work maintaining the stock population. Maria
505 Moiron and Francesca Santostefano provided numerous useful comments and suggestions on an
506 earlier draft. Two anonymous reviewers made constructive comments. Funding was provided by the
507 University of Aberdeen "Internal Funding to Pump-Prime Interdisciplinary Research and Impact
508 Activities" fund. I have no conflicts of interest.

Data accessibility
511 R code to conduct the analyses in this manuscript can be accessed at
512 https://github.com/DFofFreedom/Direct-and-indirect-phenotypic-effects-on-sociability-. The
513 associated data will be added upon acceptance.

References

Alamer, A.H. & Hoffmann, K.H. 2014. Endocrine aspects of ovary growth and gestation in the Argentinian cockroach, Blaptica dubia (Dictyoptera, Blaberidae). *http://dx.doi.org/10.1080/07924259.2013.797935* **58**: 34–41. Taylor & Francis.

Bailey, N.W. & Desjonquères, C. 2022. The Indirect Genetic Effect Interaction Coefficient ψ: Theoretically Essential and Empirically Neglected. *J. Hered.* **113**: 79–90. Oxford Academic.

Bailey, N.W. & Hoskins, J.L. 2014. Detecting cryptic indirect genetic effects. *Evolution* **68**: 1871–82.

Bailey, N.W. & Zuk, M. 2012. Socially flexible female choice differs among populations of the Pacific field cricket: geographical variation in the interaction coefficient psi (Ψ). *Proc. Biol. Sci.* **279**: 3589–96.

Baud, A., McPeek, S., Chen, N. & Hughes, K.A. 2022. Indirect Genetic Effects: A Cross-disciplinary Perspective on Empirical Studies. *J. Hered.* **113**: 1–15. Oxford Academic.

Beehler, B.M. & Foster, M.S. 1988. Hotshots, Hotspots, and Female Preference in the Organization of Lek Mating Systems. *Am. Nat.* **131**: 203–219.

Bell, A.M., Hankison, S.J. & Laskowski, K.L. 2009. The repeatability of behaviour: a meta-analysis. *Anim. Behav.* **77**: 771–783. Elsevier Ltd.

Bergmüller, R. & Taborsky, M. 2010. Animal personality due to social niche specialisation. *Trends Ecol. Evol.* **25**: 504–511.

Bewick, A.J., Vogel, K.J., Moore, A.J. & Schmitz, R.J. 2017. Evolution of DNA Methylation across Insects. *Mol. Biol. Evol.* **34**: 654. Oxford University Press.

Bijma, P. 2011. A general definition of the heritable variation that determines the potential of a population to evolve. *Genetics* **189**: 1347–1359.

Bijma, P. 2014. The quantitative genetics of indirect genetic effects: a selective review of modelling issues. *Heredity (Edinb).* **112**: 61–69. The Genetics Society.

Bijma, P., Muir, W.M. & Van Arendonk, J.A.M. 2007. Multilevel selection 1: Quantitative genetics of inheritance and response to selection. *Genetics* **175**: 277–88.

Bijma, P. & Wade, M.J. 2008. The joint effects of kin, multilevel selection and indirect genetic effects on response to genetic selection. *J. Evol. Biol.* **21**: 1175–88.

Bleakley, B.H. & Brodie III, E.D. 2009. Indirect genetic effects influence antipredator behavior in guppies: Estimates of the coefficient of interaction psi and the inheritance of reciprocity. *Evolution (N. Y).* **63**: 1796–1806. Blackwell Publishing Inc.

Blumstein, D.T., Williams, D.M., Lim, A.N., Kroeger, S. & Martin, J.G.A. 2018. Strong social relationships are associated with decreased longevity in a facultatively social mammal. *Proc. R. Soc. B Biol. Sci.* **285**. Royal Society Publishing.

Bouchebti, S., Cortés-Fossati, F., Estepa, Á.V., Lozano, M.P., Calovi, D.S. & Arganda, S. 2022. Sex-Specific Effect of the Dietary Protein to Carbohydrate Ratio on Personality in the Dubia Cockroach. *Insects 2022, Vol. 13, Page 133* **13**: 133. Multidisciplinary Digital Publishing Institute.

Brent, L.J.N., Heilbronner, S.R., Horvath, J.E., Gonzalez-Martinez, J., Ruiz-Lambides, A., Robinson, A.G., *et al.* 2013. Genetic origins of social networks in rhesus macaques. *Sci. Rep.* **3**: 1042.

Brodin, T., Fogarty, S., Sih, A. & Cote, J. 2019. Personality-dependent survival of the invasive
mosquitofish: being social can be deadly. *Aquat. Invasions* **14**: 465–477.

Brooks, M.E., Kristensen, K., van Benthem, K.J., Magnusson, A., Berg, C.W., Nielsen, A., *et al.* 2017.
glmmTMB Balances speed and flexibility among packages for modeling. *R J.* **9**: 378–400.

Cairns, S.J. & Schwager, S.J. 1987. A comparison of association indices. *Anim. Behav.* **35**: 1454–1469.

Chenoweth, S.F., Rundle, H.D. & Blows, M.W. 2010. Experimental evidence for the evolution of
indirect genetic effects: changes in the interaction effect coefficient, psi (Psi), due to sexual
selection. *Evolution (N. Y).* **64**: 1849–1856.

Clark, D.C., Beshear, D.D. & Moore, A.J. 1995. Role of Familiarity in Structuring Male–Male Social
Interactions in the Cockroach Gromphadorhina portentosa (Dictyoptera: Blaberidae). *Ann.*
Entomol. Soc. Am. **88**: 554–561. Oxford Academic.

Clark, D.C. & Moore, A.J. 1995. Genetic aspects of communication during male-male competition in
the madagascar hissing cockroach: Honest signalling of size. *Heredity (Edinb).* **75**: 198–205.

Costa e Silva, J., Potts, B.M., Gilmour, A.R. & Kerr, R.J. 2017. Genetic-based interactions among tree
neighbors: identification of the most influential neighbors, and estimation of correlations
among direct and indirect genetic effects for leaf disease and growth in Eucalyptus globulus.
Heredity (Edinb)., doi: 10.1038/hdy.2017.25.

Culumber, Z.W., Kraft, B., Lemakos, V., Hoffner, E., Travis, J. & Hughes, K.A. 2018. GxG epistasis in
growth and condition and the maintenance of genetic polymorphism in Gambusia holbrooki.
Evolution (N. Y). **72**: 1146–1154. John Wiley & Sons, Ltd.

Dall, S.R.X., Houston, A.I. & McNamara, J.M. 2004. The behavioural ecology of personality: consistent
individual differences from an adaptive perspective. *Ecol. Lett.* **7**: 734–739.

de Villemereuil, P., Schielzeth, H., Nakagawa, S. & Morrissey, M. 2016. General methods for
evolutionary quantitative genetic inference from generalized mixed models. *Genetics* **204**:
1281–1294.

Dingemanse, N.J. & Dochtermann, N. a. 2013. Quantifying individual variation in behaviour: mixed-
effect modelling approaches. *J. Anim. Ecol.* **82**: 39–54.

Dingemanse, N.J., Dochtermann, N.A. & Nakagawa, S. 2012. Defining behavioural syndromes and the
role of 'syndrome deviation' in understanding their evolution. *Behav. Ecol. Sociobiol. 2012 6611*
66: 1543–1548. Springer.

Dochtermann, N.A. 2011. Testing Cheverud's conjecture for behavioral correlations and behavioral
syndromes. *Evolution* **65**: 1814–20.

Dochtermann, N.A., Schwab, T., Berdal, M.A., Dalos, J. & Royauté, R. 2019. The Heritability of
Behavior: A Meta-analysis. *J. Hered.* **110**: 403–410.

Dochtermann, N.A., Schwab, T. & Sih, A. 2015. The contribution of additive genetic variation to
personality variation: heritability of personality. *Proc. Biol. Sci.* **282**: 20142201. Royal Society of
London.

Dohm, M.R. 2002. Repeatability estimates do not always set an upper limit to heritability. *Funct.*
Ecol. **16**: 273–280.

Dushoff, J., Kain, M.P. & Bolker, B.M. 2019. I can see clearly now: Reinterpreting statistical
significance. *Methods Ecol. Evol.* **10**: 756–759. British Ecological Society.

Edenbrow, M., Bleakley, B.H., Darden, S.K., Tyler, C.R., Ramnarine, I.W. & Croft, D.P. 2017. The
evolution of cooperation: Interacting phenotypes among social partners. *Am. Nat.* **189**: 630–
643. University of Chicago Press.

Ellen, E.D., Rodenburg, T.B., Albers, G.A.A., Bolhuis, J.E., Camerlink, I., Duijvesteijn, N., *et al.* 2014.
The prospects of selection for social genetic effects to improve welfare and productivity in
livestock. *Front. Genet.* **5**: 377. Frontiers Research Foundation.

Farine, D.R. 2013. Animal social network inference and permutations for ecologists in R using asnipe.
Methods Ecol. Evol. **4**: 1187–1194.

Fisher, D.N., Legrice, R.J. & Painting, C.J. 2021. Social selection is density dependent but makes little
contribution to total selection in New Zealand giraffe weevils. *Proc. R. Soc. B* **288**. The Royal
Society.

Fisher, D.N. & McAdam, A.G. 2019. Indirect genetic effects clarify how traits can evolve even when
fitness does not. *Evol. Lett.*, doi: 10.1002/evl3.98. John Wiley & Sons, Ltd.

Fisher, D.N. & Pruitt, J.N. 2019. Opposite responses to selection and where to find them. *J. Evol. Biol.*
32: 505–518. John Wiley & Sons, Ltd (10.1111).

Formica, V. a, Wood, C.W., Larsen, W.B., Butterfield, R.E., Augat, M.E., Hougen, H.Y., *et al.* 2012.
Fitness consequences of social network position in a wild population of forked fungus beetles
(Bolitotherus cornutus). *J. Evol. Biol.* **25**: 130–137.

Formica, V., Donald, H., Marti, H., Irgebay, Z. & Brodie, E. 2020. Social network position experiences
more variable selection than weaponry in wild subpopulations of forked fungus beetles. *J.
Anim. Ecol.* **90**: 168–182. Blackwell Publishing Ltd.

Formica, V., Wood, C., Cook, P. & Brodie, E. 2016. Consistency of animal social networks after
disturbance. *Behav. Ecol.* **28**: 85–93. Oxford University Press.

Fox, J. & Weisberg, S. 2019. *An R Companion to Applied Regression*, Third. Sage, Thousand Oaks (CA).

Gartland, L.A., Firth, J.A., Laskowski, K.L., Jeanson, R. & Ioannou, C.C. 2022. Sociability as a
personality trait in animals: methods, causes and consequences. *Biol. Rev.* **97**: 802–816. John
Wiley & Sons, Ltd.

Gelman, A. & Rubin, D.B. 1992. Inference from Iterative Simulation Using Multiple Sequences. *Stat.
Sci.* **7**: 457–472. Institute of Mathematical Statistics.

Grandcolas, P. 1998. The Evolutionary Interplay of Social Behavior, Resource Use and Anti-Predator
Behavior in Zetoborinae+ Blaberinae+Gyninae+Diplopterinae Cockroaches: A Phylogenetic
Analysis. *Cladistics* **14**: 117–127. John Wiley & Sons, Ltd.

Griffing, B. 1967. Selection in reference to biological groups. I. Individual and group selection applied
to populations of unordered groups. *Aust. J. Biol. Sci.* **20**: 127–39.

Hadfield, J.D. 2010. MCMC methods for multi-response generalized linear mixed models⁠: The
MCMCglmm R package. *J. Stat. Softw.* **33**: 1–22.

Hadfield, J.D., Nutall, A., Osorio, D. & Owens, I.P.F. 2007. Testing the phenotypic gambit: Phenotypic,
genetic and environmental correlations of colour. *J. Evol. Biol.* **20**: 549–557. John Wiley & Sons,
Ltd (10.1111).

Hadfield, J.D., Wilson, A.J., Garant, D., Sheldon, B.C. & Kruuk, L.E.B. 2010. The misuse of BLUP in
ecology and evolution. *Am. Nat.* **175**: 116–25. The University of Chicago Press.

Han, C.S., Santostefano, F. & Dingemanse, N.J. 2016. Do social partners affect same-sex sexual
behaviour in male water striders? *Anim. Behav.* **116**: 53–59. Academic Press.

Han, C.S., Tuni, C., Ulcik, J. & Dingemanse, N.J. 2018. Increased developmental density decreases the
magnitude of indirect genetic effects expressed during agonistic interactions in an insect.
Evolution (N. Y). **72**: 2435–2448.

Hintze-Podufal, C. & Vetter, R. 1996. Hormonal Control of Courtship Behavior and Reproductive
Cycle in the Cockroach Species Blaptica dubia (Blattoidea: Blaberoidea: Blaberidae). *Entomol.*
Gen. **20**: 169–175. Schweizerbart'sche Verlagsbuchhandlung.

Holtmann, B., Lagisz, M. & Nakagawa, S. 2017. Metabolic rates, and not hormone levels, are a likely
mediator of between-individual differences in behaviour: a meta-analysis. *Funct. Ecol.* **31**: 685–
696.

Houslay, T.M. & Wilson, A.J. 2017. Avoiding the misuse of BLUP in behavioural ecology. *Behav. Ecol.*
12: S9.

Jensen, K., McClure, C., Priest, N.K. & Hunt, J. 2015. Sex-specific effects of protein and carbohydrate
intake on reproduction but not lifespan in Drosophila melanogaster. *Aging Cell* **14**: 605. Wiley-
Blackwell.

Kazancioğlu, E., Klug, H. & Alonzo, S.H. 2012. The evolution of social interactions changes predictions
about interacting phenotypes. *Evolution (N. Y).* **66**: 2056–2064.

Kent, C., Azanchi, R., Smith, B., Formosa, A. & Levine, J.D. 2008. Social Context Influences Chemical
Communication in D. melanogaster Males. *Curr. Biol.* **18**: 1384–1389. Cell Press.

Kluen, E. & Brommer, J.E. 2013. Context-specific repeatability of personality traits in a wild bird: a
reaction-norm perspective. *Behav. Ecol.* **24**: 650–658.

Knoll, A.T., Jiang, K. & Levitt, P. 2018. Quantitative trait locus mapping and analysis of heritable
variation in affiliative social behavior and co-occurring traits. *Genes, Brain Behav.* **17**: e12431.
John Wiley & Sons, Ltd.

Kraft, B., Lemakos, V.A., Travis, J. & Hughes, K.A. 2018. Pervasive indirect genetic effects on
behavioral development in polymorphic eastern mosquitofish. *Behav. Ecol.* **29**: 289–300.
Oxford Academic.

Krause, J. & Ruxton, G.D. 2002. *Living in groups*. Oxford University Press.

Kruuk, L.E.B. & Wilson, A.J. 2018. The challenge of estimating indirect genetic effects on behavior: A
comment on Bailey et al. *Behav. Ecol.* **29**: 13–14. Oxford University Press.

Lea, A.J., Blumstein, D.T., Wey, T.W. & Martin, J.G.A. 2010. Heritable victimization and the benefits
of agonistic relationships. *Proc. Natl. Acad. Sci. U. S. A.* **107**: 21587–92. National Academy of
Sciences.

Maklakov, A.A., Simpson, S.J., Zajitschek, F., Hall, M.D., Dessmann, J., Clissold, F., *et al.* 2008. Sex-
Specific Fitness Effects of Nutrient Intake on Reproduction and Lifespan. *Curr. Biol.* **18**: 1062–
1066. Cell Press.

Marie-Orleach, L., Vogt-Burri, N., Mouginot, P., Schlatter, A., Vizoso, D.B., Bailey, N.W., *et al.* 2017.
Indirect genetic effects and sexual conflicts: Partner genotype influences multiple
morphological and behavioral reproductive traits in a flatworm. *Evolution (N. Y).* **71**: 1232–
1245. Wiley/Blackwell (10.1111).

McGlothlin, J.W. & Brodie, E.D. 2009. How to measure indirect genetic effects: The congruence of

680   trait-based and variance-partitioning approaches. *Evolution (N. Y).* **63**: 1785–1795.

681   McGlothlin, J.W. & Fisher, D.N. 2021. Social Selection and the Evolution of Maladaptation. *J. Hered.,*
682   doi: 10.1093/JHERED/ESAB061. Oxford University Press (OUP).

683   Montero, A.P., Williams, D.M., Martin, J.G.A. & Blumstein, D.T. 2020. More social female yellow-
684   bellied marmots, Marmota flaviventer, have enhanced summer survival. *Anim. Behav.*

685   Moore, A.J. 1997. The evolution of social signals: Morphological, functional, and genetic integration
686   of the sex pheromone in Nauphoeta cinerea. *Evolution (N. Y).* **51**: 1920–1928. John Wiley &
687   Sons, Ltd.

688   Moore, A.J., Brodie, E.D.I. & Wolf, J.B. 1997a. Interacting phenotypes and the evolutionary process: I.
689   Direct and indirect genetic effects of social interactions. *Evolution (N. Y).* **51**: 1352–1362.

690   Moore, P.J., Harris, W.E., Montrose, V.T., Levin, D. & Moore, A.J. 2004. Constraints on Evolution and
691   Postcopulatory Sexual Selection: Trade-offs among Ejaculate Characteristics. *Evolution (N. Y).*
692   **58**: 1773–1780.

693   Moore, P.J., Reagan-Wallin, N.L., Haynes, K.F. & Moore, A.J. 1997b. Odour conveys status on
694   cockroaches. *Nat. 1997 3896646* **389**: 25–25. Nature Publishing Group.

695   Muir, W.M. 2005. Incorporation of competitive effects in forest tree or animal breeding programs.
696   *Genetics* **170**: 1247–1259.

697   Nakagawa, S. & Schielzeth, H. 2010. Repeatability for Gaussian and non-Gaussian data: a practical
698   guide for biologists. *Biol. Rev.* **85**: 935–956.

699   Parker, G.A. 1979. Sexual selection and sexual conflict. In: *Sexual Selection and Reproductive*
700   *Competition in Insects* (M. S. Blum & N. A. Blum, eds), pp. 123–166. Academic Press, New York,
701   USA.

702   Patel, A.;, Jenkins, M.;, Rhoden, K.;, Barnes, A.N.A., Giangaspero, A., Barlaam, A., *et al.* 2022. A
703   Systematic Review of Zoonotic Enteric Parasites Carried by Flies, Cockroaches, and Dung
704   Beetles. *Pathog. 2022, Vol. 11, Page 90* **11**: 90. Multidisciplinary Digital Publishing Institute.

705   Poissant, J., Wilson, A.J. & Coltman, D.W. 2010. Sex-specific genetic variance and the evolution of
706   sexual dimorphism: A systematic review of cross-sex genetic correlations. *Evolution (N. Y).* **64**:
707   97–107. John Wiley & Sons, Ltd.

708   R Development Core Team. 2016. R: A language and environment for statistical computing. R
709   Foundation for Statistical Computing, Vienna, Austria.

710   Rebar, D., Bailey, N.W., Jarrett, B.J.M. & Kilner, R.M. 2020. An evolutionary switch from sibling rivalry
711   to sibling cooperation, caused by a sustained loss of parental care. *Proc. Natl. Acad. Sci. U. S. A.*
712   **117**: 2544–2550. National Academy of Sciences.

713   Santostefano, F., Garant, D., Bergeron, P., Montiglio, P. & Réale, D. 2019. Social selection acts on
714   behavior and body mass but does not contribute to the total selection differential in eastern
715   chipmunks. *Evolution (N. Y).* evo.13875. John Wiley & Sons, Ltd (10.1111).

716   Schal, C., Gautier, J.Y. & Bell, W.J. 1984. Behavioral ecology of cockroaches. *Biol. Rev.* **59**: 209–254.
717   John Wiley & Sons, Ltd.

718   Scott, A.M., Dworkin, I. & Dukas, R. 2022. Evolution of sociability by artificial selection. *Evolution (N.*
719   *Y).* **76**: 541–553. John Wiley & Sons, Ltd.

720   Scott, A.M., Dworkin, I. & Dukas, R. 2018. Sociability in Fruit Flies: Genetic Variation, Heritability and

721   Plasticity. *Behav. Genet.* **48**: 247–258.

722 Signor, S.A., Abbasi, M., Marjoram, P. & Nuzhdin, S. V. 2017. Social effects for locomotion vary
723   between environments in Drosophila melanogaster females. *Evolution (N. Y).* **71**: 1765–1775.

724 Silk, J.B. 2007. The adaptive value of sociality in mammalian groups. *Philos. Trans. R. Soc. Lond. B.*
725   *Biol. Sci.* **362**: 539–59. The Royal Society.

726 Snell-Rood, E.C. 2013. An overview of the evolutionary causes and consequences of behavioural
727   plasticity. *Anim. Behav.* **85**: 1004–1011.

728 Staes, N., Weiss, A., Helsen, P., Korody, M., Eens, M. & Stevens, J.M.G. 2016. Bonobo personality
729   traits are heritable and associated with vasopressin receptor gene 1a variation. *Sci. Reports*
730   *2016 61* **6**: 1–8. Nature Publishing Group.

731 Trubenová, B., Novak, S. & Hager, R. 2015. Indirect genetic effects and the dynamics of social
732   interactions. *PLoS One* **10**: e0126907.

733 Vipperman, A.C. 2021. Uninvited guests: Response of social networks to changing group composition
734   in the maritime earwig (Anisolabis maritima). Villanova University.

735 Westneat, D.F. & Sih, A. 2009. Sexual conflict as a partitioning of selection. *Biol. Lett.* **5**: 675–677. The
736   Royal Society.

737 Wilson, A.D.M. & Krause, J. 2014. Personality and social network analysis in animals. In: *Animal*
738   *Social Networks* (J. Krause, R. James, D. W. Franks, & D. P. Croft, eds). Oxford University Press,
739   Oxford.

740 Wilson, A.D.M., Krause, S., Dingemanse, N.J. & Krause, J. 2012. Network position: a key component
741   in the characterization of social personality types. *Behav. Ecol. Sociobiol.* **67**: 163–173.

742 Wolf, J.B., Brodie III, E.D., Cheverud, J.M., Moore, A.J. & Wade, M.J. 1998. Evolutionary
743   consequences of indirect genetic effects. *Trends Ecol. Evol.* **13**: 64–69.

744 Wolf, J.B., Brodie III, E.D. & Moore, A.J. 1999. Interacting Phenotypes and the Evolutionary Process.
745   II. Selection Resulting from Social Interactions. *Am. Nat.* **153**: 254–266.

746 Wu, H. 2013. Biology of Blaptica dubia (Blattodea: Blaberidae). Auburn University.

747 Zarchi, A.A.K. & Vatani, H. 2009. A Survey on Species and Prevalence Rate of Bacterial Agents
748   Isolated from Cockroaches in Three Hospitals. *https://home.liebertpub.com/vbz* **9**: 197–200.
749   Mary Ann Liebert, Inc. 2 Madison Avenue Larchmont, NY 10538 USA .