1 **Title:** Strength of selection potentiates distinct adaptive responses in an evolution experiment
2 with outcrossing yeast
3

4 **Authors:** Mark A. Phillips, Rupinderjit K. Briar, Marcus Scaffo, Shenghao Zhou, Molly K.
5 Burke

6
Abstract

Combining experimental evolution with whole-genome sequencing is now a well-
established method for studying the genetics of adaptation and complex traits. In this type of
work that features sexually-reproducing populations, studies consistently find that adaptation is
highly polygenic and fueled by standing genetic variation. Less consistency is observed with
respect to general evolutionary dynamics however; for example, investigators remain ambivalent
about whether selection produces repeatable versus idiosyncratic responses, or whether small
shifts in allele frequencies at many loci drive adaptation, versus selective sweeps at fewer loci.
Resolving these open questions is a crucial next step as we move toward extrapolating findings
from laboratory evolution experiments to populations inhabiting natural environments. We
propose that subtle differences in experimental parameters between studies can influence
evolutionary dynamics in meaningful ways, and here we empirically assess how one of those
parameters – selection intensity – shapes these dynamics. We subject populations of outcrossing
Saccharomyces cerevisiae to zero, moderate, and high ethanol stress for ~200 generations and
ask: 1) does stronger selection intensity lead to greater changes in allele frequencies, and a higher
likelihood of selective sweeps at sites driving adaptation; and (2) do targets of selection vary
with selection intensity? We find some evidence for positive correlations between selection
intensity and allele frequency change, but no evidence for more sweep-like patterns at high
intensity. While we do find genomic regions that suggest some shared genetic architecture across
treatments, we also identify distinct adaptive responses in each selection treatment. Combined
with evidence of phenotypic trade-offs between treatments, our findings support the hypothesis
that selection intensity might influence evolutionary outcomes due to pleiotropic and epistatic
interactions. As such, we conclude that details of a selection regime should be a major point of
consideration when attempting to generalize inferences across experimental evolution studies.
Finally, our results demonstrate the importance of clearly defined controls when associating
genomic changes with adaptation to specific selective pressures. Despite working with a
presumably lab-adapted model system, without this element we would not have been able to
distinguish between genomic responses to ethanol stress and laboratory conditions.

Introduction

Evolve and resequence ("E&R") experiments are now commonly used to study the genetics of adaptation and complex traits (Long et al. 2015; Schlötterer et al. 2015). In these studies, experimental populations are subjected to carefully designed selective regimes under controlled conditions, and sampled for DNA or RNA sequencing over many generations. This experimental framework allows for powerful statistical associations between observed, presumably adaptive, phenotypic responses with shifts in genetic variation or changes in gene expression. In addition to linking genotypes to phenotypes, the ability to control population size, level of starting genetic variation, and environmental conditions make these experiments a powerful tool for studying broad adaptive dynamics. It is now well understood that adaptation in E&R studies featuring outcrossing systems is fueled by standing genetic variation (Long et al. 2015; Barghi et al. 2020). However, the specific evolutionary dynamics observed across these studies vary considerably. We believe achieving a better understanding of what causes these differences is a key step towards extending general findings from E&R to real populations (Phillips and Burke 2021).

Major points of discrepancy in the literature today include topics like evolutionary repeatability, the role of contingency in the genetics of adaptation, and what some term "sweeps versus shifts" (Barghi et al. 2020). Using evolutionary repeatability as an example, while many studies report high levels of evolutionary repeatability, as evidenced by parallel responses to selection across replicate populations (e.g. Linder et al. 2021), others find more idiosyncratic and replicate-specific responses (e.g. Barghi et al. 2019. We suggest that this and other observed discrepancies are likely due to differences in experimental design among studies, as well as the genetic variation segregating in ancestral populations. In the case of evolutionary repeatability, differences in starting allele frequency distributions and effective population size stand out as potentially relevant factors (i.e., these factors directly influence the probability that selection will favor a specific beneficial allele in a given replicate). More broadly, we believe these discrepancies point to a need for deeper understanding of how experimental parameters influence adaptive processes before attempting to extend inferences from E&R studies to populations outside the lab. Here we focus on how a specific experimental choice – selection intensity – impacts evolutionary outcomes in large outcrossing populations with abundant standing genetic variation.

How a given selective pressure manifests (e.g. intensity, duration, constant versus dynamic, etc.) has clear ecological relevance that likely shapes evolutionary outcomes in both natural populations and laboratory experiments. While this topic has been explored across a number of theoretical studies (e.g. Kessner and Novembre 2015; Stetter et al. 2018; Christodoulaki et al. 2019; Vlachos and Kofler 2019; Hayward and Sella 2021), few E&R studies have sought to address it empirically. Here we aim to address this gap by directly testing the effects of selection intensity on the genetic architecture of adaptation using populations of outcrossing *S. cerevisiae*. Specifically, we compare the genomic response to moderate and high ethanol stress in replicate populations all derived from a single ancestral population. We are

primarily interested in how selection intensity shapes signatures of selection and whether
treatments featuring different selection intensities show distinct genomic responses to selection.
Christodoulaki et al. (2019) explored the idea that selection intensity influences
signatures of selection using simulations designed to mimic laboratory experimental evolution.
They find that when a new trait optimum is distant (i.e. when a strong selective pressure is
applied), the genomic response to selection often involves substantial increases in allele
frequencies that are consistent with selective sweeps. However, for close trait optima (weaker
selection pressure), adaptation proceeds through smaller shifts in allele frequencies. The "sweeps
versus shifts" question – in other words, the question of whether population-genetic models
focused on identifying few loci of large effect are better at explaining trait evolution than
quantitative-genetic models that assume extensive polygenicity – looms beyond the experimental
evolution literature (Höllinger et al. 2019), and we believe that selection intensity stands out as a
major factor key to resolving this question. A major goal of this study is therefore to test to
prediction that high selection intensity results in larger changes in allele frequencies at and
around target sites.
We are also interested in exploring if the same targets of selection respond to both high
and moderate selection intensities. In the simplest scenario, there is some set of "ethanol
resistance" alleles segregating in our ancestral population, and these should increase in frequency
in both selection intensity treatments. However, if the genetics of complex traits is characterized
by widespread pleiotropy as some suggest (Visscher and Yang 2016; Boyle et al. 2017) and/or
epistasis, this might not be the case. For example, due to pleotropic effects and resulting trade-
offs with other characters, alleles that are adaptive under strong ethanol stress conditions might
not be favored under weaker ethanol stress conditions and vice versa. If this prediction holds
true, it would suggest a clear need to incorporate this possibility as a factor when constructing
models of polygenic adaptation. Lastly, alleles that are beneficial at late but not early stages of
adaptation have been previously identified in evolving outcrossing *S. cerevisiae* populations,
which could indicate epistatic interactions (Phillips et al. 2020). If this is the case and generally
true, it could also drive treatment-specific responses in combination with pleiotropy; for
example, an allele that is beneficial at moderate levels of ethanol stress may not confer those
same benefits at higher ethanol stress due to negative interactions with alleles unambiguously
beneficial under high ethanol stress.
In summary, with this work we aim to address the following questions: (i) do populations
experiencing stronger selection intensities display larger changes in allele frequencies at and
around targets of selection compared to populations experiencing weaker selection intensities;
and (ii) do targets of selection vary with selection intensity? We address these questions using
genomic data from outcrossing *S. cerevisiae* populations subjected to zero, moderate, and high
ethanol stress for ~200 generations.
118
119
120

Materials and Methods

Experimental populations and selection regimes

All experimental populations used in this study were derived from the "S12" population described in detail in Phillips et al. (2021). Briefly, this population was created by combining 12 haploid strains from the barcoded SGRP yeast strain collection (Cubillos et al. 2009). The specific strains used are: Y12, YSP128, SK1, DBVPG6044, UWOPS05_217_3, L_1528, L_1374, DBVPG6765, YJM981, YJM975, BC187, and N273614. As described in Linder et al. (2020), these strains have been modified to enable easy crossing and diploid recovery – these modified strains were kindly provided by Dr. Anthony Long. The MATa and MATα strains contain *ho* deletions to prevent mating-type switching, but each contain a different drug-resistance marker inserted into a pseudogene (YCR043C) tightly linked to the mating-type locus (*MATa, hoΔ, ura3::KanMX-barcode, ycr043C::NatMX* and *MATα, hoΔ, ura3::KanMX-barcode, ycr043C::hphMX*). These genotypes enable haploids of each mating type to be recovered using media supplemented with either hygromycin B or nourseothricin sulfate, and they enable newly mated a/α diploids to be recovered in media supplemented with both drugs.

To create the S12 recombinant outbred population from these haploid strains, we used a semi-round-robin design of initial crossing, followed by 12 iterations of what we call our "weekly outcrossing protocol" to maximize genetic diversity and allow for some domestication to laboratory handling conditions (detailed in Burke et al. 2020; Phillips et al. 2021). The experimental populations featured in this study were then derived from samples of this base population: 20 control replicates (C_{1-20}), 20 moderate ethanol stress replicates (M_{1-20}), and 20 high ethanol stress replicates (H_{1-20}). Levels of ethanol used in the two stress treatments were chosen based on the results of exploratory growth rate assays (see next section for general assay methods). We found that 10% ethanol was close to the limit of what could support population growth (at least for a 48-hour period), and we thus chose this as the high stress treatment. We chose the moderate treatment to include 6% ethanol as this resulted in a doubling time approximately mid-way between the high stress and control conditions (Supplementary Figure 1).

The weekly outcrossing protocol described by Burke et al. (2020) was also used to maintain sexual reproduction in these 60 replicate populations, with minor modifications to increase throughput (i.e. smaller volumes). This protocol involves batch culture of diploids in 1mL of liquid medium in alternating wells of 24-well plates (Corning); every other well contained sterile YPD which was monitored for growth (which would indicate potential cross-contamination) throughout the experiment. After batch culture of diploids, the entire 1mL of culture was washed and resuspended in 1mL minimal sporulation media (1% potassium acetate), and incubated with shaking for 72 hours (30°C/200 rpm). After sporulation, a modified random spore isolation protocol was implemented to disrupt asci and isolate spores. This protocol involves resuspending sporulated cultures in 1mL Y-PER Yeast Protein Extraction Reagent (Thermo), followed by incubation at 50°C for 15 minutes to kill vegetative diploid cells. Cultures were then resuspended in a 1% zymolyase (Zymo Research) solution to weaken ascus

walls, and vortexed at maximum speed with 0.5 mm silica beads (BioSpec) to mechanically
agitate the asci. Following these steps, spores were transferred to YPD agar plates supplemented
with nourseothricin sulfate (100mg/L), hygromycin B (300mg/L) and G418 (200mg/L) and
incubated at 30°C for 48 hours, during which time spores mated (due to close proximity on the
plate) and diploids germinated. The resulting lawns of new diploid cells were scraped off plates
using sterile glass slides and transferred to 10mL of sterile YPD media (1/10th of this culture was
preserved in 15% glycerol and archived at -80°C for DNA extraction and sequencing). From
these super-saturated cultures, 10 μL were transferred into 1mL of treatment-specific medium
and incubated for 48 hours (30C/200rpm), with a 1/100 dilution halfway through to increase
generational turnover. For the control treatment, standard YPD (2% yeast extract, 1% peptone,
2% dextrose) was used as the culture medium. For the moderate and high ethanol stress
treatments, YPD was supplemented with either 6% or 10% ethanol (by volume), respectively.
After this period of batch culture, the next weekly outcrossing iteration would commence
(Supplementary Figure 2). All replicates were handled in parallel for a total of 15 weeks, with
one cycle of outcrossing per week.
By estimating cell density at various steps of the outcrossing protocol (via OD$_{600}$ of liquid
cultures, and/or colony counts of dilution series), we can approximate the number of asexual
generations that occurred in each treatment during the experiment. We estimate that ~10 cell
doublings occurred during the 48-hour period of batch culture – this phase of competitive growth
in treatment-specific medium is most relevant to our major questions relating to selection
strength. We infer that an additional ~4 cell doublings occur during the non-competitive phases
of the protocol (i.e. growth on agar medium). We observed that growth was slower (and cell
densities lower) in the high ethanol stress treatment compared to the others, but this did not lead
to dramatically different estimates of cell doublings among the treatments. Ultimately, we
project that over 200 asexual generations elapsed in all treatments over 15 weeks. This
generational turnover is difficult to precisely estimate for several reasons, chiefly because some
aspects of population biology were not routinely quantified throughout the experiment. For
example, our protocol development (*cf*. Burke et al. 2020) suggests that sporulation and mating
efficiencies are high in the base population, and that the mortality induced by the weekly
outcrossing protocol is low. But it was not possible to quantify these metrics in all 60
experimental replicates every week. If sporulation efficiency, mating efficiency, and/or spore
viability was low in a particular replicate during a given week, this would reduce the number of
viable cells in the population (compared to the baseline expectation) and lead to a higher
estimate of cell doublings for the week. Thus, we feel that our ballpark estimates can be
considered rather conservative.
196
Growth rate assays
Growth in liquid medium was the primary phenotype we tracked in this experiment. We
compared growth between evolved populations and the ancestor in each media type to detect

evidence of phenotypic adaptation in a given treatment. We also compared growth of evolved
populations across all media types to detect evidence of potential phenotypic trade-offs.
For each population assayed, a culture would first be grown overnight in 10 mL YPD in a
shaking incubator (30 °C/200 rpm). Cultures were then quantified via absorbance at OD_{600}
(values of 1/100 dilutions typically ranged from 0.10 to 0.15 across all treatments and
timepoints) and diluted to a standard starting concentration of $OD_{600} = 0.1$ using the appropriate
media type. These dilutions were then transferred to individual wells of a 96-well plate (Corning)
in 200 μL volumes. A Tecan Spark multimodal microplate reader measured absorbance in each
well every 30 minutes over a 48-hour period at 30°C with no shaking. Measurements were taken
at four positions within each well and the average values were used for subsequent data analysis.
To avoid edge effects, wells on the outer perimeter of the plates were not used for assays and
simply filled with sterile YPD. In a given plate reader assay, we included 9 technical replicates
of the ancestral population and 10 randomly selected replicates from each treatment. Multiple
plate reader assays were run under control, moderate ethanol stress, and high ethanol stress
conditions. Growth curves were generated using the raw readings, and the R (R CoreTeam 2021)
package "Growthcurver" (Sprouffske and Wagner 2016) used to estimate doubling times and
carrying capacity. These estimates are obtained by fitting data to the following logistic equation
that gives the number of cells N_t (as measured by absorbance) at time t:
218

$$N_t = \frac{K}{1 + \left(\dfrac{K - N_0}{N_0}\right) e^{-rt}}$$

219
where starting population size is represented by N_0 carrying capacity by K. Here carrying
capacity is simply defined as the maximum population size in a particular environment. Lastly, r
represents the growth rate that would occur if there were no limits on total population size. This
value is also used calculate the doubling time which is defined as $\frac{Ln2}{R}$. Figures shown in the
manuscript are representative of patterns seen across multiple assays. However, we note that
while patterns were consistent across assays, some specific values varied between plate reader
runs (e.g. carrying capacity estimates from cycle 15 samples varied between runs under high
ethanol conditions, but the observation that carrying capacity in H replicates was lower than
replicates from the other treatments remained consistent).
229
DNA extraction, sequencing, and read mapping
The 20 replicates of each treatment were sampled for pooled-population genome
sequencing at three distinct timepoints: after a single outcrossing cycle, after 7 outcrossing
cycles, and after 15 outcrossing cycles. To sample populations for sequencing, 1 mL of freezer
stocks were revived on YPD agar plates. After 48 hours of growth at 30ºC, the resulting lawns
were broadly sampled by wooden applicator (to capture as much genetic diversity as possible)
and these cells were inoculated into 10mL of liquid YPD culture, and growth overnight in the

shaking incubator. Genomic DNA was extracted from samples with Qiagen's Yeast/Bact Kit

following the manufacterer's protocol. After checking DNA quantity, sequencing libraries were

prepared for Illumina sequencing with the Nextera Kit DNA Sample Preparation Kit,

implementing some routine modifications to increase throughput (e.g. Baym et al. 2015).

Libraries were pooled into groups of 48 and run on at least one PE150 lane of the HiSeq3000

housed at OSU's Center for Quantitative Life Sciences (CQLS); samples with lower-than-

average coverages were re-quantified, re-pooled and re-sequenced such that high (>50X average

genome-wide) coverages were achieved across all experimental replicates.

The SNP calling pipeline we routinely use has been described previously (e.g. Phillips et

al. 2020. We used GATK v4.0 (Van der Auwera & O'Connor 2020) to align raw data to the *S*

cerevisiae S288C reference genome (R64-2-1) and create a single VCF file for all variants

identified across all populations, using standard best practices workflows and default filtering

options. This VCF file was converted into a "raw" SNP frequency table by extracting the AD

(allele depth) and DP (unfiltered depth) fields for all SNPs passing quality filters; the former

field was used as the minor allele count observed at a presumed SNP, and the latter was used as

the total coverage observed at that site. The VCF file was also used as an input for SnpEff v4.3

(Cingolani et al. 2012) to extract potential functional effects of individual SNPs. SnpEff

annotates each variant in a VCF file (e.g. by tagging whether it occurs within a protein-coding

sequence) and calculates the effect(s) each produces on known genes (e.g. amino acid changes).

We applied several filtering and quality control steps to the raw SNP table generated

from the steps above ensure only high-confidence sites were used in subsequent analysis. Mean

genome-wide coverage across the 180 samples sequenced for this study ranged from ~50X to

~300X with a median value of ~80X (Supplementary Table 1). We began our filtering process

by only considering sites where sequence coverage exceeded 20X in all populations. Next, we

removed all sites that were not expected to be polymorphic based on the previously described

sequences of the founder strains used to create the ancestral population (*cf.* Phillips et al. 2021).

We also removed sites that were not polymorphic in ancestral population itself, as for this study

our objective is to track the evolution of standing genetic variants, and not *de novo* mutations.

Finally, we only considered sites where the minor allele frequencies between 0.02 and 0.98

across the entire dataset. This removes sites where the alternate nucleotide is fixed across all

populations, and sites where errors in sequencing and/or variant calling may create the

appearance of polymorphism. These steps ultimately resulted in a SNP table with 61,281 SNPs.

<u>Identifying candidate sites and selection response categories</u>

In pursuing the goal of associating particular candidate SNPs with adaptation, we found it

useful to assign these to one of five distinct categories: (i) general adaptation to laboratory

handling, (ii) general adaptation to ethanol stress, (iii) specific adaptation to the high ethanol

treatment, (iv) specific adaptation to the moderate ethanol treatment, and (v) specific adaptation

to the control treatment. Comparing the magnitude of change at sites across these categories

allows us to assess whether greater selection intensity results in greater shifts in allele

frequencies, and comparing the location of sites observed across these categories allows us to
assess whether or not different selection intensities involve different genomic targets. Notably,
the decision to include category (v) above was made *after* observing evidence of continued and
unique adaptation in our control treatment; this was an unanticipated outcome that has
thematically influenced our interpretation of all other results. To identify and differentiate
between these types of responses we used a combination of within- and between-treatment
comparisons. For within-treatment comparisons (e.g. comparing SNP frequencies between cycle
1 and cycle 15 for the C_{1-20} populations), we relied on the Cochran-Mantel-Haenszel (CMH)
tests. For between-treatment comparisons (e.g. comparing SNP frequencies between C_{1-20} and
H_{1-20}), we used a generalized linear mixed model (GLMM) approach. Patterns of overlap
between these different comparisons were then used to isolate the different types of responses we
were interested in (see Table 1 for details on how we define each type of response). As coverage
variation can impact statistical results when using nucleotide counts, as both these methods do
we scaled coverage across the genome to a uniform level of 50X before performing these tests
(Wiberg et al. 2017). Importantly, we chose approaches best-suited to the detection of parallel
responses to selection across replicates, as we believe such sites are most likely to be true targets
of selection.
The CMH tests for within-treatment comparisons are commonly used in E&R studies,
and simulations suggest this is one of the most accurate methods available for identifying
responses to selection based on sampling populations over time (Vlachos et al. 2019). Tests were
performed using the "poolSeq" package (Taus et al. 2017) in R. For a given within-treatment
comparison, the data were paired appropriately (e.g. C_1 cycle 1 with C_1 cycle 15, C_2 cycle 1 with
C_2 cycle 15, and so on for the control treatment) and tests were performed at all 61,281 K SNPs.
Significant sites show a consistent change in frequency from cycle 1 to cycle 15, across all
replicate populations. As we sequenced populations at an intermediate timepoint (cycle 7) we
also had the option to use more sophisticated time-series approaches for this task. However,
given that simulations (Vlachos et al. 2019) and empirical comparisons (Phillips et al. 2020)
have shown that these approaches perform similarly to the CMH test using starting and ending
points in an E&R study, we favored the latter approach.
For between-treatment comparisons, as there was no meaningful way to pair replicate
populations (e.g. pairing C_1 with H_1 is no more meaningful than C_1 with H_2), the CMH test was
not appropriate. We instead opted to use the GLMM approach, which emphasizes parallel
changes across replicate populations, and is becoming popular in E&R work (e.g. Jha et al.,
2015, Kawecki et al., 2021). Three pairwise comparisons were performed using this approach:
C_{1-20} versus H_{1-20}, C_{1-20} versus M_{1-20}, and H_{1-20} versus M_{1-20} (note: these comparisons only use
cycle 15 data). For each comparison, the "lme4" package (Bates et al. 2015) was used to fit a
GLMM with binomial error structure to nucleotide counts at a given polymorphic site. The
following model statement was used:
315

$$y_i = Treatment + Replicate\,(Treatment) + e_i$$

316
where y_i is the frequency of the ith SNP, $Treatment$ is a fixed effect, and $Replicate$ is a
random factor nested within $Treatment$, and e_i refers to the binomial error term. To test for
significance, the "anova" function in R was used to compare the full model to a simplified model
without the fixed effect. Using this approach, we were able to identify SNPs that were
consistently differentiated between treatments at the end of the experiment (cycle 15).
To correct for multiple comparisons and establish significance thresholds, we used the
same general permutation approach described in Burke et al. (2014) and Graves et al. (2017).
Using C_{1-20} cycle 1 versus C_{1-20} cycle 15 as an example, the relevant replicate populations were
randomly assigned to one of two groups, the CMH tests was performed at each SNP in the
shuffled data set, and the smallest p-value generated was recorded. This was then repeated 999
times for a total of 1000 permutations. The "quantile" function in R was then applied to the
resulting list of 1000 p-values to define a genome-wide false-positive rate, per site, at 0.5%. That
is to say, a Type I error rate of 0.005 is required for a site to be considered significantly
differentiated. We used a 0.5% cutoff instead of a 5% cutoff seen in previous efforts based on
recent work making the case for 0.005 thresholds instead of 0.05 when interpreting p-values
(Benjamin et al. 2018). This same general procedure was used for between-treatment
comparisons using the GLMM approach. However, for GLMM comparisons, we only ran 200
permutations instead of 1000 as this method is more computationally intensive than the CMH.
We did perform a trial using 1000 permutations for one comparison and found that it produced a
very similar threshold.
Once SNPs belonging to each category of selection response listed in Table 1 were
identified, we compared these for potential differential enrichment of gene ontology (GO) terms.
These lists were created using the SnpEff output: for each response type, we created a list of
genes associated with at least one candidate SNP in that category (note: here we consider all
effect types as valid associations). Text files with candidate list and associated SnpEff annotation
are available through Dryad for interested individuals (see Data Availability statement for
details). GO term enrichment analyses for each gene list were then performed using Metascape
(Zhou et al. 2019) with default settings.
345
<u>Sweeps versus shifts</u>
To assess the prediction that more intense selection should be associated with greater
changes in allele frequencies, we simply compared the mean changes in SNP frequency at
candidate sites between treatments using ANOVAs. If this core idea is correct, among sites
associated with general ethanol stress, we would expect to see significantly greater changes in
SNP frequency in the high ethanol stress treatment compared to the moderate ethanol stress
treatment. Next, to evaluate the prediction that more intense selection should involve signatures
of selection more consistent with selective sweeps (larger allele frequency changes at fewer loci)
rather than polygenic shifts (smaller allele frequency changes at more loci), we generated
genome-wide heterozygosity plots for replicates in each treatment to see if there were signs of

the types of depressions in heterozygosity characteristic of selective sweeps (Burke 2012). If it is
the case that more intense selection is associated with more sweep-like dynamics, this should be
evident when comparing patterns of genetic variation in the high ethanol stress replicates to the
moderate ethanol stress and control replicates.
We defined heterozygosity as *2pq* where *p* and *q* refer to the reference (major) and
alternate (minor) nucleotide frequencies. To assess genome-wide patterns, we generated sliding-
window plots showing estimates across each chromosome. To do this we first calculated
heterozygosity at each polymorphic site in our dataset in each replicate population, then averaged
across replicates within a given treatment on a per-site basis for a given timepoint. To define
windows, we used the GenWin package (Beissinger et al. 2015) package in R. This package uses
a smoothing spline technique to define windows biased on breakpoints in the data. As a
"smoothness" parameter is user defined, we tried a range of values before settling on
"smoothness = 3000". Our final selection was made based on what most clearly represented the
data. Scripts and data to generate plots with more or less smoothness are available for interested
parties through Dryad and GitHub (see "Data availability" statement for details).
371

Results
<u>Adaptation and growth rates</u>
To characterize the long-term consequences of each experimental treatment, we assayed
the growth of evolved (cycle 15) replicates in all three media types (control, moderate ethanol,
high ethanol) and compared these to the growth of the ancestral population (Figure 1). To assess
statistical differences, we used Kruskal-Wallis tests to compare mean values across all groups,
and pairwise Wilcoxon rank sum tests to determine which treatment pairs were significantly
different (note: the Benjamini-Hochberg procedure was used to correct for multiple
comparisons). We find that in plain YPD, differences in doubling time and carrying capacity are
small (Figure 1A and B, Supplementary Figure 3A), but significant for both doubling time
(Kruskal-Wallis test p=0.0001) and carrying capacity (p= 0.001). In the case of doubling time,
there are no significant differences between the C, M, and H replicates. However, the ancestral
population has a significantly longer doubling time than evolved populations (mean of 1.3 hours
for the ancestor versus 1.25 to 1.26 for C, M, and H populations); this implies that replicates in
all treatments adapted to laboratory conditions in ways that improved growth in YPD. For
carrying capacity, the H replicates appear to stop doubling at a lower cell concentration
compared to the ancestral population, and the C and M treatments (mean of 0.9 in the H
replicates versus ~1 for the others, note: carrying capacity is represented as log(OD) as shown in
Supplementary Figure 3). While we do find some statistically significant differences here, they
are clearly quite small. As such, it appears that adaptation to ethanol stress in the moderate and
high ethanol stress has not had a large negative impact on the ability of replicates in these groups
to grow in YPD (i.e. no obvious suggestion of an adaptive trade-off).
We do observe differences in growth patterns among treatments in 6% ethanol (Figure
1C), with respect to both doubling time (Figure 1D, Kruskal-Wallis test p =1.75×10^{-6}) and

carrying capacity (Supplementary Figure 3B, Kruskal-Wallis test p=8.05×10^{-7}). Both the M and
C treatments grow significantly faster than the ancestral population, evidencing adaptation. By
contrast, doubling time in the H treatment could not be significantly distinguished from the
ancestor, and was correspondingly slower than the C and M treatments. Carrying capacity is
significantly different in all pairwise comparisons except for the ancestror vs. C treatment.
Carrying capacity is lowest in the H treatment (mean=0.98) followed by the M treatment (mean
= 1.07), and the C treatment (mean=1.1) resembled that of the ancestor (mean=1.1).
In populations assayed in 10% ethanol (Figure 1E), we also find significant differences in
both doubling time (Figure 1F, Kruskal-Wallis test p=1.16×10^{-5}) and carrying capacity
(Supplementary Figure 3C, Kruskal-Wallis test p=5.46×10^{-7}) among treatments. With respect to
doubling time, we find significant differences between all pairwise comparisons except for the M
replicates versus the H replicates. Doubling time is slowest in the ancestral population (mean =
3.64 hours), followed by the C replicates (mean = 3.26 hours), followed by the M and H
replicates (mean = 2.64 and 2.57 hours). For carrying capacity, all pairwise comparisons are
significantly different except for the ancestral population versus the C replicates. The ancestral
population has the highest carrying capacity (mean = 1.54, same for the C replicates), followed
by the M replicates (1.11) and H replicates (mean = 0.78). So, while populations adapted to
ethanol stress treatments generally have faster growth rates under high ethanol stress conditions,
they also stop doubling sooner. And adaptation to high ethanol stress specifically is associated
with the lowest carrying capacity.
While these growth rate assays are useful for ranking phenotypic differences among
ancestral and evolved populations (and they are internally consistent), we do not believe the raw
metrics reported necessarily reflect what occurred during the actual selection experiment. For
instance, in the plate reader assays, populations in 10% ethanol appear to still be in lag phase
after nearly 24 hours of growth (Figure 1E). While we observed that replicates of the H treatment
grew more slowly than the other treatments throughout the experiment, routine OD_{600} readings
indicate that all populations were in log phase at 24 hours. So, whenever cultures were
transferred to fresh liquid media during the experiment, cells were in log phase in all treatments;
we felt that it was important to avoid sampling cells at different stages as this could introduce
unwanted variation among treatments. There are many reasons why the absolute values of
growth parameters estimated during the phenotype assays differ from what was observed in the
selection experiment, but we suspect the main drivers are volume differences (200 μL in 96 well
plates versus 1mL in 24 well plates), and agitation (no shaking in the plate reader versus 200rpm
in the shaking incubator).
430
<u>Evidence for different response categories among candidate SNPs</u>
To assess the genomic response to selection, we first compared SNP frequencies between
cycles 1 and 15 for each treatment using the CMH test (Supplementary Figure 4). Here we find
clear responses to selection as evidenced by consistent changes in SNP frequencies in each
treatment, including the controls. As the number of candidate regions overlap across both ethanol

stress treatments and controls, it appears that adaptation to general laboratory handling (i.e., our
weekly outcrossing protocol) is a major feature of this study. To better parse these responses to
selection, we used a GLMM approach to identify differentiated SNPs between treatments during
cycle 15 (Supplementary Figure 5). Based on patterns of overlap between all of these
comparisons, we identified 5 major response types: (i) general adaptation to laboratory handling,
(ii) general adaptation to ethanol stress, (iii) specific adaptation to the high ethanol treatment, (iv)
specific adaptation to the moderate ethanol treatment, and (v) specific adaptation to the control
treatment. (Table 1). As seen in Figure 2 where these categories are overlaid on the CMH results
comparing cycle 1 and 15 for each treatment, almost all significant SNPs can be assigned to one
of these 5 categories.
Considering the major peaks in Figure 2, we do observe some mixing of categories
within certain peaks, but most are either made up of a single response type or have a clear
majority type. And while there are a number of regions that correspond to our three ethanol
response types (general, high, and moderate ethanol responses), our strongest signals are
observed in regions associated with general laboratory selection. So, while ethanol exposure
clearly imposes stress, laboratory conditions appear to be major selective pressures on their own.
Here it should be noted that the ancestral population sampled to establish all experimental
replicates had already experienced 12 "domestication" cycles with outcrossing prior to this
experiment. As such, there was some expectation that these populations would already be
adapted to routine culture and outcrossing protocols. A principal component analysis (PCA) of
SNP frequencies indicates general differentiation between treatments, and within treatments over
time, based on clustering patterns (Figure 3).
The existence of treatment-specific responses addresses our hypothesis that complex
genetic architectures (e.g. extensive pleiotropy and/or epistasis) could lead to different targets of
selection despite the same core selective pressure. As shown in Figure 2, treatment specific
responses can be observed in each group. For instance, there are regions of the genome showing
significant responses to selection in the H replicates (Figure 2A) and not the M replicates (Figure
2B) and vice versa despite both being exposed to ethanol stress. We also find cases where
significant regions are found in the C replicates (Figure 3A) but not in either the H or M
replicates despite all three treatments adapting to the same general laboratory conditions and
maintenance protocols.
To address our hypothesis that more intense selection would lead to greater changes in
allele frequencies, we first compared the mean change in candidate SNP frequencies per replicate
per treatment between cycle 1 and 15 for sites that fall into the general ethanol selection category
(Figure 4A). As expected, there is a greater shift in SNP frequencies in the ethanol treatments
compared to the controls at these sites. However, we do not find clear evidence of greater
changes in SNP frequencies in the high ethanol stress replicates compared to the moderate
ethanol stress replicates. We also repeated this approach for the general laboratory selection
category. Here we find that while changes in candidate SNP frequencies are not significantly
different between the control and moderate ethanol stress selection treatments, shifts are smaller

in the high ethanol stress treatment compared to both. This perhaps suggests some interaction
between adaptation for high ethanol stress conditions and general laboratory adaptation.
478   Observed changes in SNP frequencies between cycle 1 and 15 for treatment-specific
responses also suggest complex underlying dynamics. Starting with control-specific responses,
we find that these sites show significantly more change in the control treatment versus both
ethanol stress treatments (Figure 5A). And, the magnitude of change is much smaller in the high
ethanol stress treatment compared to the moderate ethanol stress treatment. Again, we suggest
this may evidence a possible interaction between adaptation to ethanol stress and laboratory
conditions. We observe the opposite pattern when we focus on the high ethanol stress-specific
candidate SNPs (Figure 5B). Here as expected, we see that the change in frequencies in the high
ethanol stress SNPs is greater than the other two. These sites appear to shift more and more
consistently in the moderate ethanol stress treatment versus the controls as well, perhaps
suggesting some weak selection in their favor under moderate ethanol stress. Finally, among the
moderate ethanol stress-specific SNPs, we observe greater changes in the moderate ethanol stress
sites compared to the other treatments as expected (Figure 5C). However, this change is not
significantly different from what was observed in the control treatment, suggesting perhaps that
this category also involves weekly selected laboratory selection responses.
493
494   <u>SNP frequency trajectories</u>
495   Given the complex genomic responses to selection observed in each experimental
treatment, we examined individual SNP trajectories in each response type in an attempt to
deepen our understanding of how selection is acting in our system. For some of the most
significant markers in peaks corresponding to our different response categories, we plotted raw
allele frequency in each replicate over the three sequenced timepoints. In representative sites
associated with general laboratory stress and general ethanol stress, we observe the expected
pattern of a clear and consistent increase in frequency in all three treatments for the former but
only the two ethanol stress treatments for the latter (Figure 6 and Supplementary Figure 6).
However, patterns are more complicated when looking at treatment specific response types.
504   In representative high ethanol stress-specific sites, we observe the greatest consistent
increase in the high ethanol stress replicates (Figure 7A and Supplementary Figure 7A).
However, the trajectories in the moderate ethanol stress treatment trend consistently upward to a
greater degree than what we see in the control treatment. This again suggests that some high
ethanol stress response sites are being weakly selected for in the moderate ethanol treatment. We
do not find such clear patterns looking at our representative moderate ethanol stress-specific
responses. Here we see the greatest consistent change in the moderate ethanol stress treatment,
but changes in the other two treatments show an upward trend as well (Figure 7B and
Supplementary Figure 7B). Here one interpretation is that this "moderate ethanol treatment"
category is capturing sites that are also beneficial in both or one of the other treatments, but more
so under moderate ethanol stress.

Finally, in representative control-specific sites, we again observe evidence of strong
directional selection in the control treatment and perhaps weaker selection in the moderate
ethanol stress treatment (Figure 7C and Supplementary Figure 7C). However, we find that in the
high ethanol stress treatment, SNP frequency changes very little over the course of the
experiment and this observation is quite consistent across replicates and independent peaks. This
is also consistent with the small changes observed across all sites in the category as shown in
Figure 5A (red boxplot). Our interpretation of this result is that these sites might provide
significant benefits in terms of adaptation to general laboratory conditions and outcrossing
protocols, but these benefits may diminish (or incur costs) as greater ethanol stress is introduced.
However, there does not appear to be strong enough selection for them to be completely
eliminated, at least on the time scale of this experiment.
526
<u>Sweeps versus shifts in response to selection</u>
The assess whether more intense selection was associated selective sweeps, we simply
visualized patterns of heterozygosity across the genome and looked for evidence of dramatic
reductions in genetic diversity (Burke 2012). Specifically, we compared genome-wide levels of
variation within each treatment at cycles 1 and 15 to evaluate the effect of 15 cycles of selection
on diversity, in a way that was consistent among replicates. Mean heterozygosity across the C,
M, and H treatments during cycle 15 does not provide any clear indication of more sweep-like
regions with depressed heterozygosity in the ethanol stress treatments versus the controls, or in
the high ethanol stress treatment versus the moderate (Supplementary Figure 8B, D, and F).
While we certainly observe genomic regions where heterozygosity is low at the end of the
experiment, it is important to contextualize this observation relative to the heterozygosity
observed at the start of the experiment; in most of these regions heterozygosity was low to begin
with (Supplementary Figure 8A vs 8B; 8C vs 8D; 8E vs 8F). This can be visualized more clearly
in Supplementary Figure 9 where we plot the absolute difference in mean heterozygosity over
time in each treatment. We do find that some of our most significant peaks from CMH
comparisons between cycles 1 and 15 are associated with pronounced reductions in
heterozygosity (e.g. the regions of chromosome 7 and 11 where heterozygosity dramatically
decreased correspond to most significant peaks in Supplementary Figure 4). However, the depth
of these depressions does not appear to be associated with intensity of selection. For instance, on
chromosome 3 there is a relatively isolated region we believe is responding to ethanol stress in
the both M and H but not the C populations (Figure 2). Here we find there is also a large change
in heterozygosity seen in the ethanol treatments that is not present in the controls (Supplementary
Figure 9). However, it is not the case this region looks like more of a complete sweep in the high
ethanol stress treatment than in the moderate ethanol stress treatment (Supplementary Figure 9).
551
<u>Candidate genes associated with different response categories</u>
To compare potential differences in genes and pathways associated with our major
response types (Table 1), we performed gene enrichment analyses using the candidate genes

associated with each category. As shown in Supplementary Table 2, we find do find enrichment
terms like "signaling" and "growth" that are shared across categories. However, we also find
distinct enriched terms within categories (both broadly and narrowly defined). This lends some
creditability to the idea that selection is targeting different biological mechanisms across
treatments. For instance, considering terms enriched in the general ethanol selection category, we
find "alcohol catabolic process" and "response to organic substance" – terms that intuitively
might underlie adaptation to long-term ethanol stress – that are not enriched in the general
laboratory selection category. We also find distinct enrichment patterns with respect to
treatment-specific categories. The most enriched terms in the high ethanol stress-specific
response category are associated with cytokinesis and asexual reproduction, while the most
enriched terms in the moderate ethanol stress-specific response category are more associated
with pseudohyphal and filamentous growth, as well as metabolic function (e.g. lipid metabolism
and sugar homeostasis). Lastly, the most enriched terms in the control-specific response category
are related to autophagy and immune function.
569
Discussion
Using ethanol stress as the focal selective pressure, our experiment was designed to test if
stronger selection was associated with greater changes in allele frequencies at target sites, and if
targets of selection varied among treatments involving different selection intensities due to
complex trait architectures. However, our genomic analysis makes it clear that despite the use of
an ancestral population that was previously subjected to a significant period of laboratory
domestication (involving a dozen cycles of outcrossing and hundreds of asexual generations),
adaptation to laboratory conditions is a major feature of our results (Figure 2 and Supplementary
Figure 4C). While this complicates our interpretations, we believe our study still has the power
to address our core questions, as the control treatment allows us to distinguish between adaptive
responses associated with laboratory domestication and those associated with ethanol stress
(Table 1). And while we cannot definitively say ethanol exposure is exerting a greater selective
pressure than general laboratory handling, it stands to reason that total selection intensity is
highest in the high ethanol stress treatment, followed by the moderate ethanol stress treatment,
and lowest in the control treatment.
585
Patterns of growth in evolved populations imply different avenues of adaptation
The observed growth-related phenotypes in our experimentally-evolved populations
suggest complex evolutionary dynamics underlying adaptation to ethanol stress. In YPD (the
control medium), replicates of the C, M, and H treatments all have similar doubling times after
15 cycles and all grow slightly faster than the ancestral population (Figure 1). This is perhaps
expected given that our genomic results indicate all populations are continuing to adapt to
laboratory conditions. However, under moderate and high ethanol stress conditions, patterns are
more complicated. Under moderate ethanol stress, replicates of the H treatment have slower
growth rates and lower carrying capacities (Figure 1C-D, and Supplementary Figure 3B) than all

other groups assayed. And while they do have higher doubling times than controls and the
ancestral population in high ethanol stress conditions (Figure 1E-5), they again have the lowest
carrying capacities (Supplementary Figure 3C). Replicates of the M treatment have similar
doubling times under high ethanol conditions, but do not display the same level of reduction in
carrying capacity (Supplementary Figure 3C). Therefore, we conclude that adaptation to ethanol
stress involves mechanisms not directly related to growth rate. Differences between the moderate
and high ethanol stress populations across media types also implicate potential trade-offs
between growth rate and final sustainable population size.
These observations of potential trade-offs support the idea that pleiotropy and genetic
interactions are general features of complex trait architecture. In the case of stress resistance,
there is already support for this idea in the literature based on the work of Kawecki et al. (2021).
Here the authors find that even though the same genes underlie starvation resistance in in
experimentally-evolved *D. melanogaster*, directionally of allele frequency shifts change
depending on whether selection is imposed at larval or adult life stages due to underlying trade-
offs. More broadly, we do not find it is simply the case that increased selection intensity results
in some clear and proportional phenotypic response. As such, selection intensity should be
carefully considered when using experimental evolution to predict how specific traits might
change in response to a given selective pressure.
613
<u>The magnitude of allele frequency change at target sites is context-dependent</u>
If increased selection intensity does result in greater changes in SNP frequency at and
around targets of selection as simulation work by Christodoulaki et al. (2019) suggest, we would
expect that at candidate sites underlying the general response to general stress, changes should be
greater in the high ethanol stress treatment compared to the moderate ethanol treatment. In
principle, this should also be true for candidate sites associated with general laboratory
adaptation; we would predict that changes in the high and moderate ethanol stress treatments
should be greater than those observed in the control treatment. With this in mind, our results are
mixed when looking across the different response types outlined in Table 1. In the general
ethanol selection category, the magnitude of change in candidate SNPs is similar in the high and
moderate ethanol stress treatment (Figure 4A). And for the general laboratory selection category,
SNP changes in the high ethanol stress treatment are actually smaller than both the control and
moderate ethanol stress treatments (Figure 4B). These observations refute the expectation that
stronger selection intensity should result in larger frequency changes at target sites. However,
our other response types suggest there is perhaps still some merit to this idea.
Among sites we have identified as control-specific and high ethanol-specific, we do
observe examples that are strongly selected for in the primary treatment, and weakly selected for
in the moderate ethanol stress treatment (Figure 5 and 7, Supplementary Figure 7). In the case of
the high ethanol-specific candidates, these perhaps represent instances where more intense
selection has in fact produced greater changes in frequencies at target sites. However,
interpretation is more complicated for control-specific candidates. While these responses are

presumably adaptative for the selective pressures imposed by experimental protocols common to
all three treatments, they appear to be less so with increasing ethanol exposure due to some
interaction. All together, we interpret these results as further evidence that different selection
intensities can in fact potentiate different trait architectures.
We considered the possibility that in this experiment, adaptation to laboratory protocols
may have been stressful enough that the inclusion of ethanol exposure had little additional
impact (i.e. all treatments are under very intense selection). This could explain why the general
laboratory selection category is accounts for so many of the most significant candidates SNPs we
identify. However, growth rate assays make it clear that the ancestral populations and controls
are indeed negatively impacted by ethanol exposure (Figure 1). As such, we generally argue that
the three experimental treatments experienced distinct and specific selection pressures, and that
our results partially support the idea of a positive correlation between selection intensity and
allele frequency change.
648
<u>Stronger selection does not obviously promote selective sweep dynamics</u>
We predicted that stronger selection intensities should produce signatures of adaptation
more consistent with selective sweeps (dramatic change observed at fewer loci), while weaker
selection intensities should produce signatures of adaptation more consistent with polygenic
adaptation (subtle shifts observed at many loci). Patterns of heterozygosity observed across the
genome in experimental populations of each treatment do not provide strong support for these
predictions. Genomic regions that could be considered most "sweep-like" (i.e. reductions in
variation where heterozygosity approaches zero) are rare in all treatments, and regions that best
fit this description were generally already low in variation prior to the start of the experiment
(Supplementary Figure 8). In the rare regions where we do observe large reductions in
heterozygosity associated with one or more of our 5 categories of candidate sites, it is not the
case that these regions look more "sweep-like" in one treatment type compared to another; in
other words, we find no clear evidence to support the idea that the high ethanol stress is more
likely to result in sweep-like patterns than the moderate ethanol or control treatment (the region
of reduced heterozygosity mid-way through chromosome 7 provides an illustrative example).
664
<u>The presence of different selection response categories reinforces that trait architectures vary</u>
<u>with selection intensity</u>
As noted previously, we believe the existence of treatment-specific responses to selection
across groups indicate that genetic architecture of adaptation varies with selection intensity. For
instance, while we do find evidence of "general ethanol selection" responses, there are also a
number responses that appear specific to either high or moderate ethanol exposure (red peaks in
Figure 2A, orange peaks in Figure 2B). Based on trajectories (Figure 7A and B, Supplementary
Figure 7A and B), it is possible that some of these sites are responding to selection in both
treatments but how strongly they are favored varies with exposure. However, even with this
interpretation, the observed context-dependence supports the idea that trait architecture varies

with selection intensity. This is further supported by the apparent phenotypic trade-offs in the M
and H replicates described above.
Surprisingly, the clearest evidence for different trait architectures at different selection
intensities comes from our control populations and control-specific responses (Figure 2C).
Unlike our moderate or high ethanol specific peaks which are often small or contain SNPs from
multiple response categories, control-specific responses include some of our most significant and
"pure" peaks. So, while these regions of the genome are strongly associated with the
maintenance and laboratory conditions that are shared by all experimental treatments, they
appear to be (at most) weakly selected under moderate ethanol stress and not selected at all under
high ethanol stress (Figure 7C and Supplementary Figure 7C). Here one possibility is while these
alleles do confer advantages under control conditions, they have costs in the presence of ethanol
and those costs increase as the level of ethanol exposure increases (i.e. context-dependence due
to pleiotropy-associated trade-offs). Alternatively, this pattern could also be explained by
negative epistatic interactions that amplify with increasing levels of ethanol stress. While we do
not have the ability to distinguish between these two scenarios, both point towards complex
relationships between selection intensity and the genetic architecture of adaptative traits.
691
<u>Selection targets different biological mechanisms across treatments</u>
As shown by our gene enrichment analysis, different biological functions appear to be
associated with adaptation across treatments. Focusing on the M and H populations, we find that
adaptation to general ethanol stress involves genes related to alcohol catabolic processes,
signaling, and responses to abiotic stimuli (Supplementary Table 2). However, we also observe
distinct enrichment terms within the M and H categories; while adaptation to high ethanol stress
is associated with GO terms relating to the integrity of cell division (e.g. cytokinesis, DNA
repair, and budding), these functions may either be unnecessary or come with some cost at lower
levels of ethanol stress. Similarly, while adaptation to moderate ethanol stress is associated with
GO terms related to metabolic function, these are not implicated at high levels of ethanol
exposure. Together, these results provide further evidence that the genetic architecture of
adaptation varies with selection intensity due to underlying differences and potential trade-offs in
the mechanisms through which adaptation occurs.
With respect to the various mechanisms associated with ethanol resistance in and across
the moderate and high ethanol stress treatments, our findings are consistent with past work
suggesting ethanol resistance in *S. cerevisiae* is complex and involves many genetic and
physiological mechanisms (reviewed in Ding et al. 2009 and Ma and Liu 2010). Signaling, lipid
metabolism, carbohydrate homeostasis all stand out as mechanisms implicated by past studies.
Work focusing on adaptation to high ethanol exposure in asexual *S. cerevisiae*, in this case
peaking at 12% of the culture medium, also specifically implicated mutations in genetic
pathways associated with cell cycle and DNA replication (Voordeckers et al. 2015). Here the
authors suggest that delayed cell cycle progression allows for greater protection for individual
cells. This sort of survival mechanism, which might more accurately be described as stress

tolerance rather than stress resistance, may drive the apparent lower carrying capacity observed in the populations adapted for high ethanol stress in the presence of ethanol. This stands out as an intuitive example of a potential context-dependent adaptative mechanism associated with ethanol stress.

Conclusions

  Our results generally support the idea that selection intensity significantly impacts evolutionary dynamics. As such, this should be a major point of consideration when synthesizing results across E&R studies even when they feature the same stress or condition as a focal selective pressure. It should also be considered when attempting to extrapolate findings from E&R studies to natural populations. For instance, candidate sites and dynamics observed in E&R studies with intense selection are perhaps unlikely translate very well to a natural population facing more moderate environmental changes and vice versa. Our observations also indicate a greater need for models of polygenic adaptation to include the possibility of extensive pleiotropy and epistasis than sometimes thought. Furthermore, while here we chose to focus on selection intensity, it is likely that other dynamic circumstances of natural environments, such as temporal heterogeneity in selection, will also affect outcomes in meaningful ways. As such, we believe our findings illuminate a clear need for studies that explicitly demonstrate how experimental and population-genetic factors shape evolutionary dynamics in E&R studies moving forward. Lastly, as with the work of Pfenninger and Foucault (2020), our results demonstrate the importance of appropriate controls even when timeseries data are available. Our controls ultimately prevented us from making a number of spurious associations between large portions of observed genomic responses to selection and ethanol stress resistance.

References

Barghi, N., Hermisson, J., Schlötterer, C. (2020). Polygenic adaptation: a unifying framework to understand positive selection.

Bates, D., Machler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, 67, 1-48.

Baym, M., Kryazhimskiy, S., Liberman, T.D., Chung, H., Desai, M.M., & Kishony, R. (2015). Inexpensive multiplexed library preparation for megabase-sizes genomes. *PLoS One*, 10**,** e0131262.

Beissinger, T.M., Rosa, G.J.M, Kaeppler, S.M., Gianola, D., & de Leon, N. (2015). Defining window-boundaries for genomic analyses using smoothing spline techniques. *Genetics Selection Evolution*, 47, 30 .

Boyle, E.A., Li, Y.I., & Pritchard, J.K. (2017) An expanded view of complex traits: from polygenic to omnigenic. *Cell*, 169, 1177-1186 .

757 Burke, M.K. (2012), How does adaptation sweep through the genome? Insights from long-term
758 selection experiments. *Proceedings of the Royal Society B,* 279, 5029-5038.
759
760 Burke, M.K., Liti, G., & Long, A.D. (2014). Long standing genetic variation drives repeatable
761 experimental evolution in outcrossing populations of *Saccharomyces cerevisiae*. *Molecular*
762 *Biology and Evolution, 32*, 3228-3239.
763
764 Burke, M.K., McHugh, K.M., & Kutch, I.C. (2020). Heat shock improves random spore analysis
765 in diverse strains of Saccharomyces cerevisiae. *Frontiers in Genetics*, 11, 1636.
766
767 Christodoulaki, E., Barghi, N, & Schlötterer, C (2019). Distance to trait optimum is a crucial
768 factor determining the genomic signature of polygenic adaptation. *bioRxiv*,
769 doi.org/10.1101/721340.
770
771 Cingolani, P., Platts, A., Wang, L., Coon, M., Nguyen, T., Wang, L., Land, S.J., Ruden, D.M., &
772 Lu, X. (2012). A program for annotating and predictiong the effects of single nucleotide
773 polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w1118; iso-2,
774 iso-3. *Fly*, 6, 80-92.
775
776 Cubillos, F.A., Louis, E.J., & Liti, G. (2009) Generation of a large set of genetically tractable
777 haploid and diploid *Saccharomyces* strains. *FEMS Yeast Research,* 9, 1217-1225.
778
779 Ding, J., Huang, X., Zhang, L., Zhao, N, Yang, D., & Zhang, K. (2009). Tolerance and stress
780 response to ethanol in the yeast Saccharomyces cerevisiae. *Applied Microbiology and*
781 *Biotechnology*, 85, 253-263.
782
783 Graves, J.L., Hertweck, K.L., Phillips, M.A., Han, M.V., Cabral, L.G., Barter, T.T., … Rose
784 MR. 2017. Genomics of parallel experimental evolution in Drosophila. *Molecular Biology and*
785 *Evolution,* 34, 831–842.
786
787 Hayward, L.K., & Sella, G. (2021). Polygenic adaptation after a sudden change in environment.
788 bioRxiv, doi.org/10.1101/792952.
789
790 Höllinger, I., Pennings, P.S., & Hermisson, J. (2019). Polygenic adaptation: from sweeps to
791 subtle frequency shifts. *PLoS Genetics*, 15, e1008035.
792
793 Jha, A.R., Miles, C.M., Lippert, N.R., Brown, C.D., White, KP, & Kretiman, M. (2015). Whole-
794 genome resequencing of experimental populations reveals polygenic basis of egg-size variation
795 in *Drosophila melanogaster. Molecular Biology and Evolution,* 32, 2616-2632.
796
797 Kawecki, T.J., Erkosar, B., Dupuis, C., Hollis, B., Stillwell, R.C., Kapun, M. (2021). Molecular
798 Biology and Evolution, 28, 2732-2749.
799
800 Kessner, D., & Novembre, J. (2015). Power analysis of artificial selection experiments using
801 efficient whole genome simulation of quantitative traits. *Genetics*, 199, 991-1005.
802

803   Linder, R.A., Zabanavar, B., Majumder, A., Shyan-Chiao Hoang, H., Delgado, V., Tran, R., La,
804   V., Leemans, S., & Long, A.D. (2021). bioRxiv: doi.org/10.1101/2021.08.27.457900.
805
806   Long AD, Liti G, Luptak A, Tenaillon O. (2015). Elucidating the molecular architecture of
807   adaptation via evolve and resequence experiments. *Nature Reveiews Genetics,* 16, 567-582.
808
809   Ma, M., & Liu, Z.L. (2010). Mechanism of ethanol tolerance in *Saccharomyces cerevisiae.*
810   *Applied Microbiology and Biotechnology*, 87, 829-845.
811
812   Pfenninger, M. &, Foucault, Q. (2020). Genomic processes underlying rapid adaptation of a
813   natural *Chironomus riparius* population to unintendedly applied experimental selection
814   pressures. *Molecular Ecology*, 29, 536-548.
815
816   Phillips, M.A., & Burke, M.K. (2021). Can laboratory evolution experiments teach us about
817   natural populations? *Molecular Ecology*, 30, 877-879.
818
819   Phillips, M.A., Kutch, I.C., Long, A.D., & Burke, M.K. (2020) Increased time sampling in an
820   eolve-and-resequence experiment with outcrossing *Saccharomyces cerevisiae* reveals multiple
821   paths of adaptive change. *Molecular Ecology*, 29, 4898-4912.
822
823   Phillips, M.A., Kutch, I.C., McHugh, K.M., Taggard, S.K., & Burke, M.K. (2021). Crossing
824   design shapes patterns of genetic variation in synthetic recombinant populations of
825   *Saccharomyces cerevisiae*. *Scientific Reports*, 11, 19551.
826
827   R Core Team. (2021). R: a language and environment for statistical computing. R foundation for
828   Statistical Computing, Vienna, Austria. www.R-project.org.
829
830   Schlötterer, C., Kofler, R., Versace, E., Tobler, R., & Franssen, S.U. (2015). Combining
831   experimental evolution with next-generation sequencing: a powerful tool to study adaptation
832   from standing genetic variation. *Heredity*, 114, 431-440.
833
834   Sprouffske, K. & Wagner, A. (2016). Growthcurver: an R package for obtaining interpretable
835   meterics from microbial growth curves. *BMC Bioinformatics,* 17, 172 (2016).
836
837   Stetter, M.G., Thornton, K., & Ross-Ibarra, J. (2018). Genetic architecture and selective sweeps
838   after polygenic adaptation to distant trait optima. *PLoS Genetics*, 14, e1007794.
839
840   Taus, T., Futschik, A., & Schlötterer, C. (2017). Quantifying selection with pool-seq time series
841   data. *Molecular Biology and Evolution,* 34, 3023-3034.
842
843   Van der Auwera, GA., & O'Connor, B.D. (2020). Genomics in the cloud: using Docker, GATK,
844   WDL, and Terra. O'reilly Media.
845
846   Visscher, P.M., & Yang, J. (2016) A plethora of pleiotropy across complex traits. *Nature*
847   *Genetics*, 48, 707-708.
848

849   Vlachos, C., & Kofler, R. (2019). Optimizng the power to identify the genetic basis of complex
850   traits with evolve and resequence studies. Molecular Biology and Evolution, 36, 2890-2905.
851
852   Vlachos, C., Burny, C., Pelizzola, M., Borges, R., Futschik, A., Kofler, R., & Schlötterer, C.
853   (2019). Benchmarking software tools for detecting and quantifying selection in evolve and
854   resequecing studies. *Genome Biology,* 20, 169.
855
856   Voordeckers, K., Komine, J., Das, A., Espinosa-Cantu, A., De Maeyer, D., Arslan, A., Van Pee,
857   M., van der Zande, E, Meert, W., Yang, Y., Zhu, B., Marchal, K., DeLuna, A., Verstrepen, K.J.
858   (2015). Adaptation to high ethanol reveals complex evolutionary pathways. *PloS Genetics*, 11,
859   e1005635.
860
861   Wiberg, R.A., Gaggiotti, O.E., Morrisssey, M.B., Ritchie M.G. (2017). Identifying consistent
862   allele frequency differences in studies of stratified populations. Methods in Ecology and
863   Evolution, 8, 1899-1909.
864

Data Availability
The raw sequence files generated over the course of this project are available through NCBI
SRA (BioProject ID: PRJNA839395) and scripts used to process raw data and perform SNP
calling are available through Github (https://github.com/mollyburke/Burke-Lab-SNP-calling-
pipeline). Core scripts necessary to reproduce our results are available through Github
(https://github.com/mphillips67/Genomics-Ethanol-Stress-Outcrossing-Yeast) and major input
and results files are available through Dryad (https://doi.org/10.5061/dryad.tqjq2bw27).
872

873   **Author Contributions**
874   M.A.P. and M.K.B. conceived of this project. M.A.P., R.K.B., M.S., and S.Z. did the lab work
875   necessary to generate the data sets used in this study. M.A.P. carried out the data analysis, and
876   M.A.P. and M.K.B. wrote the manuscript.
877

878   **Acknowledgements**
879   We thank Oregon State University's Center for Quantitative Life Sciences for use of their
880   computational and sequencing resources. This work was supported by startup funds provided to
881   M.K.B. by the College of Science at Oregon State University, and M.A.P. was supported by a
882   National Science Foundation Postdoctoral Fellowship (NSF 1906246).

Figures

893 Figure 1. Growth rates and doubling times for ancestral population and experimental populations
894 after 15 cycles of adaptation in plain YPD (A and B), 6% ethanol YPD (C and D), and 10%
895 ethanol YPD (E and F). Nine technical replicates for ancestor, and 10 randomly chosen replicates
896 from each treatment were used in these assays.

897
898
899   Figure 2. CMH results comparing SNP frequencies between cycles 1 and 15 in (A) High ethanol
900   stress, (B) Moderate ethanol stress, and (C) Control treatments. Black lines represent
901   permutation derived significance thresholds, and color coding of significantly differentiated sites
902   corresponds response categories described in Table 1.

903
904
Figure 3. PCA plot showing how different time points for replacement population in each
treatment cluster based on SNP frequencies.

908   Figure 4. Boxplots showing the absolute mean change in SNP frequency for each replicate in

909   each treatment for (A) General ethanol response candidate SNPs and (B) General Laboratory

910   Selection Candidate SNPs.

913
914 Figure 5. Boxplots showing the absolute mean change in SNP frequency for each replicate in
915 each treatment for (A) Control Specific Candidate SNPs, (B) High Ethanol Stress Candidate
916 SNPs, and (C) Moderate Ethanol Stress Candidate SNPs.

922
923
924   Figure 6. SNP frequency trajectories for representative (A) General Laboratory Selection
925   response site, and (B) General Ethanol Selection response sites. Trajectories for all 20 replicates
926   within each treatment are plotted.
927

Figure 7. SNP frequency trajectories for representative (A) High Ethanol Stress Specific response site, (B) Moderate Ethanol Stress Specific response, and (C) Control Specific response sites. Trajectories for all 20 replicates within each treatment are plotted.

940 **Tables**
941 Table 1. The five major response types used to categorize our candidate sites and how they are
942 defined based on how results overlap across within and between treatment statistical
943 comparisons.
944

RESPONSE TYPE	DEFINITION
GENERAL LABORATORY SELECTION	Significantly differentiated sites that overlap across all three treatments based on comparisons between cycles 1 and 15 for each treatment. Sites that are differentiated between Controls versus High and Moderate Ethanol Stress treatments are also removed to account for directionality.
GENERAL ETHANOL SELECTION	Significantly differentiated sites that overlap across High and Moderate Ethanol Stress treatments based on comparisons between cycles 1 and 15 for each. Any sites shared with Control cycle 1 versus 15 are removed, along with sites that are differentiated between High versus Moderate Ethanol stress during cycle 15.
HIGH ETHANOL STRESS SPECIFIC RESPONSE	Significantly differentiated sites that are unique to High Ethanol Stress treatment based on cycle 1 versus cycle 15 comparisons (i.e. removing overlap with Control and Moderate Ethanol Stress cycle 1 versus 15 comparisons).
MODERATE ETHANOL STRESS SPECIFIC RESPONSE	Significantly differentiated sites that are unique to Moderate Ethanol Stress treatment based on cycle 1 versus cycle 15 comparisons (i.e. removing overlap with Control and High Ethanol Stress cycle 1 versus 15 comparisons).
CONTROL SPECIFIC RESPONSE	Significantly differentiated sites that are unique to Control treatment based on cycle 1 versus cycle 15 comparisons (i.e. removing overlap with High and Moderate Ethanol Stress cycle 1 versus 15 comparisons).

953 **Supplementary Figures**

957 Supplementary Figure 1. Growth rates (A) and doubling times (B) for ancestral population in
958 plain YPD, 6% ethanol YPD, and 10% ethanol YPD.

964 Supplementary Figure 2. General schematic showing how the H, M, and C populations were
965 maintained over the course of this study.

979   Supplementary Figure 3. Carrying capacity estimates based on growth rate assays for ancestral
980   population and experimental populations after 15 cycles of adaptation in plain YPD (A), 6%
981   ethanol YPD (B), and 10% ethanol YPD (C). Nine technical replicates for ancestor, and 10
982   randomly chosen replicates from each treatment were used in these assays.

Supplementary Figure 4. CMH results comparing SNP frequencies between cycles 1 and 15 in (A) High ethanol stress, (B) Moderate ethanol stress, and (C) Control treatments. Black lines represent permutation derived significance thresholds.

988
989
990 Supplementary Figure 5. GLMM results from comparing SNP frequencies cycle 15 SNP
991 frequencies between (A) High and Moderate Ethanol Stress treatments, (B) High Ethanol Stress
992 and Control treatments, and (C) Moderate Ethanol Stress and Control treatments. Black lines
993 represent permutation derived significance threshold.

Supplementary Figure 6. SNP frequency trajectories for representative (A) General Laboratory Selection response site, and (B) General Ethanol Selection response sites. Trajectories for all 20 replicates within each treatment are plotted.

1011

1012

1013 Supplementary Figure 7. SNP frequency trajectories for representative (A) High Ethanol Stress
1014 Specific response site, (B) Moderate Ethanol Stress Specific response, and (C) Control Specific
1015 response sites. Trajectories for all 20 replicates within each treatment are plotted.

1016

1017

1018
1019
1020   Supplementary Figure 8. Mean heterozygosity for C_{1-20} replicates (A and B), M_{1-20} replicates (C
1021   and D), and H_{1-20} replicates (E and F) plotted across the genome during cycles 1 and 15. Panels
1022   A, C, and E were generated cycle 1 data, while B, D, and F are from cycle 15.

Supplementary Figure 9. Absolute difference in mean heterozygosity between cycles 1 and 15 for C_{1-20} replicates (A), M_{1-20} replicates (B), and H_{1-20} replicates (C) plotted across the genome. The gold square over chromosome 3 in each panel represents a region where there are candidate SNP's in both ethanol treatments, but not in the controls.

Supplementary Tables

1058

1059 Supplementary Table 1. Average sequence coverage for each timepoint of each replicate
1060 sequenced.

1061

Replicate	Cycle	Mean Coverage
Ancestor	-	88.38
C1	1	53.00
	7	78.78
	15	144.82
C2	1	57.18
	7	89.32
	15	53.21
C3	1	56.26
	7	76.21
	15	118.98
C4	1	137.77
	7	95.03
	15	87.49
C5	1	104.24
	7	138.28
	15	91.45
C6	1	62.48
	7	150.01
	15	115.99
C7	1	59.82
	7	108.10
	15	96.40
C8	1	86.30
	7	163.51
	15	130.89
C9	1	151.69
	7	108.04
	15	51.33
C10	1	121.45
	7	193.92

	15	54.42
C11	1	72.61
	7	202.69
	15	54.11
C12	1	76.27
	7	50.01
	15	88.65
C13	1	131.62
	7	93.09
	15	117.77
C14	1	97.99
	7	51.87
	15	68.71
C15	1	80.67
	7	91.39
	15	74.77
C16	1	59.72
	7	92.11
	15	65.09
C17	1	108.88
	7	69.87
	15	73.76
C18	1	58.25
	7	75.39
	15	69.35
C19	1	74.23
	7	82.11
	15	60.47
C20	1	65.85
	7	92.91
	15	73.25
M1	1	148.45
	7	75.50
	15	63.77
M2	1	151.08
	7	68.35

	15	126.24
M3	1	202.20
	7	70.62
	15	151.19
M4	1	88.88
	7	180.83
	15	48.66
M5	1	93.68
	7	143.10
	15	146.67
M6	1	97.64
	7	62.12
	15	138.83
M7	1	148.94
	7	90.99
	15	293.71
M8	1	153.92
	7	75.30
	15	118.00
M9	1	65.18
	7	94.02
	15	70.62
M10	1	161.58
	7	89.92
	15	83.01
M11	1	95.15
	7	69.57
	15	126.96
M12	1	77.64
	7	133.60
	15	106.45
M13	1	107.26
	7	72.31
	15	134.66
M14	1	118.18
	7	107.43

	15	143.51
M15	1	87.80
	7	57.45
	15	81.69
M16	1	127.69
	7	100.79
	15	170.28
M17	1	89.98
	7	108.87
	15	100.55
M18	1	71.52
	7	76.33
	15	59.61
M19	1	85.46
	7	110.75
	15	74.14
M20	1	101.33
	7	94.21
	15	58.27
H1	1	99.97
	7	137.61
	15	92.27
H2	1	102.67
	7	133.16
	15	78.03
H3	1	86.47
	7	149.63
	15	182.16
H4	1	104.77
	7	96.76
	15	197.48
H5	1	80.32
	7	78.75
	15	108.65
H6	1	82.64
	7	77.99

	15	173.67
H7	1	50.45
	7	62.33
	15	48.20
H8	1	54.53
	7	68.14
	15	55.64
H9	1	71.14
	7	71.95
	15	85.80
H10	1	88.05
	7	80.73
	15	78.34
H11	1	62.50
	7	93.30
	15	119.56
H12	1	59.88
	7	69.80
	15	96.67
H13	1	78.81
	7	92.58
	15	65.30
H14	1	98.72
	7	134.69
	15	117.62
H15	1	61.00
	7	69.39
	15	64.24
H16	1	47.84
	7	123.63
	15	81.27
H17	1	61.66
	7	161.66
	15	103.26
H18	1	102.09
	7	134.52

	15	125.07
	1	74.00
H19	7	145.69
	15	174.77
	1	73.81
H20	7	97.08
	15	175.79