1

Decoding Sequence-Structure-Function-Evolution of basic Leucine Zippers of Aureochromes from Heterokont Algae

Madhurima Khamaru[a], Debarshi Bose[a], Anwesha Deb[a] and Devrani Mitra[a*]

[a]Department of Life Sciences, Presidency University, 86/1 College Street, Kolkata – 700073, WB, India

[*]Address of correspondence - devrani.dbs@presiuniv.ac.in

Running title – Decoding bZIPs: Aureochromes versus the Rest

Keywords

Highlights

- Aureochrome is perhaps the only light-responsive transcription factor in bZIP superfamily.

- We draw a comparative between aureochromes and other bZIPs via *in-silico*/*in-vitro* methods.

- Aureochromes form a distinctly separate lineage, midway in bZIP evolution.

- The unique histidine facilitates aureochromes' interaction with cognate DNA substrates.

Abstract

The blue light photoreceptor cum transcription factors, Aureochromes (Aureos), are present exclusively in photosynthetic stramenopiles. Co-existence of Light-Oxygen-Voltage (LOV) and basic leucine zipper (bZIP) is unique to Aureos – therefore ideal to study light-dependent DNA binding/transcriptional regulation. Further, Aureos' inverse effector-sensor topology, resembling several sensory eukaryotic transcription factors, makes them prototypical optogenetic scaffolds. In absence of 3D data, this study aims for a thorough investigation of

the bZIP domains from Aureos and others, and their interaction with substrate DNA using
tools from sequence/structural bioinformatics, network theory, molecular dynamics
simulation and *in vitro* experiments. An in-depth comparison of 173 Aureo/plant/opisthokont
bZIPs reveals Aureos' uniqueness and evolutionary significance in DNA binding specificity
as well as dimer stability. An all-atom network analysis on representative bZIP-DNA co-
crystal structures, especially the measurement of eigenvector centrality, further adds
importance to hydrophobic interactions in the zipper region to stabilize bZIP dimer and
facilitate DNA binding in Aureos and other bZIPs. Perhaps the most notable finding is the
unique histidine substitution at the basic region of Aureos unlike any other bZIPs. Not only is
this residue important for DNA binding, this can serve as a potential switch point in
Aureo/bZIP evolution.

1. Introduction

Regulation of gene expression is essential for the survival of every life form. Transcription
factors (TFs), upon binding with the cognate DNA, turn genes on/off at precise spatio-
temporal resolution. Different classes of transcription factors exist. They differ on the basis of
their DNA binding structural motifs and activity. Examples include basic leucine zipper
(bZIP), basic helix-loop-helix (bHLH), Zinc-coordinating DNA-binding motifs (Zinc finger),
helix-turn-helix (HTH), beta scaffold factors like STAT (Signal transducer and activator of
transcription proteins), MADS (MCM1, AGAMOUS, DEFICIENS, and SRF) box, HMG
box (high mobility group box), TBP (TATA-binding protein), Beta barrel alpha helix etc.
Among these, bZIPs comprise one of the largest group of eukaryotic transcription factors (Ji
et al., 2018). The uniqueness of bZIPs lies in their conserved nature of structure and sequence
specificity. They possess N-X7-R/K signature motif and binds mostly to the *cis* acting DNA
elements of a promoter having an 'ACGT' core sequence (Wang et al., 2018) (Rodríguez-
Martínez et al., 2017). Interdigitation between the partner helices leads to the dimerization of
bZIP — resulting in a superimposed coiled coil structure or 'zipper' (Bader & Vogt,
2006)(Sinden, 1994). This zipper-like structure is extended by the N-terminal helical
segment, which can bind to the DNA major groove via basic amino acid residues(Glover &
Harrison, 1995). Dimerization is essential for the recognition of substrate DNA and
transcriptional regulation. The C-terminal zipper motif consists of heptads, which can be
represented by a helical wheel. The residues are conventionally marked from 'a' to 'g'
(Grigoryan & Keating, 2006), where leucines generally occupy every 'd' position, whereas
the hydrophobic residues including leucine/isoleucine dominate 'a' (Bader & Vogt, 2006).

However, presence of a conserved asparagine at 'a' position in the middle of zipper could

result in destabilization of the oligomer (Harbury et al., 1993). The 'b', 'c' and 'f' positions

are generally filled by polar residues. The charged residues at the 'e' and 'g' positions also

aid in the zipper dimer stability (Deppmann et al., 2004)(Podust et al., 2001). Leucine

residues, owing to their hydrophobic side chains, create 'knobs' which can fit into the 'holes'

generated in between the side chains of amino acids present in adjacent helices (Bader &

Vogt, 2006). At the other end, the basic amino acid residues complement well with the

negatively charged phosphate backbone of DNA — this results in favorable electrostatic

interactions. Besides, they can also participate in hydrogen bonding/polar interactions and

can even engage in non-polar interactions with the portions of extended side chains. Besides

phosphates, the nitrogenous base and deoxyribose sugar are also involved in hydrogen

bonding, van der Waal's as well as stacking interactions. The affinity of bZIP TFs towards

the cognate target DNA sequences is imparted by the sequence composition of its basic

region as well as the length of the leucine zipper (Hakoshima, 2005a). The dimerization

partner is also crucial for its sequence specific DNA binding (Pogenberg et al., 2014). Their

ability to bring in change in the binding specificity upon hetero-dimerization is indicative of

this (Busch & Sassone-Corsi, 1990). The electrostatic attraction and repulsion forces, acting

between the polar residues that flank the hydrophobic interaction surface of the zippers, are

major deciding factors behind the formation of homo/heterodimers (C. R. Vinson et al.,

1993). The oppositely charged residues located at 'g' and the following e' positions across the

hydrophobic interface, form the inter-helical ion bridges. The interactions between d-d', a-a',

g-e' are vital for the zipper formation and stabilization (Podust et al., 2001).

Widespread among the regulatory proteins of gene expression, the bZIPs play a

crucial role in cellular proliferation and differentiation in eukaryotes. It is known to be the

ancient DNA binding module found across three kingdoms of life (Hakoshima, 2005b).

Several groups of opisthokont and plant bZIPs are well defined and characterized. Few

homologs of bZIPs are also evident in fungi, algae, protists and even in bacteria. However,

the only blue light (BL) responsive photoreceptor with a bZIP TF activity has been recently

discovered (Takahashi et al., 2007) in Aureos – found exclusively in photosynthetic

stramenopiles (Rayko et al., 2010). The flavin binding Light-Oxygen-Voltage (LOV) BL

sensor at the C-terminus, makes Aureos different from any other conventional bZIPs. Aureos

perform organism specific diverse functionalities [**SI-1**] e.g. - branching in *Vaucheria sp*

(Takahashi et al., 2007)*,* cell cycle progression in *Phaeodactylum sp* (Kroth et al., 2017)*,*

lipid accumulation in *Nannochloropsis sp.* (Huang et al., 2014) etc. The broad spectrum
functions of Aureos include promotion of high light acclimation, photo protection in diatoms
etc (Huysman et al., 2013). Considering maximum penetration of BL in sea water, most of
the physiological and developmental responses are BL dependent in photosynthetic marine
algae. However, it lacks phototropins (Rayko et al., 2010), which is one of the major BL
photoreceptors in land plants. Aureos instead aid in the optimization of photosynthesis (Mann
et al., 2020) under varied light conditions. In fact, Aureos' significance in carrying out BL-
mediated gene expression is not only important to study BL signaling in photosynthetic
marine algae, they are potentially relevant for optogenetics as well. The direct DNA binding
ability of Aureo in response to BL without an intricate signaling mechanism makes Aureo an
ideal optogenetic scaffold. They are capable of bringing in faster changes in gene expression
under fluctuating light conditions (Mann et al., 2020).   Dual presence of a LOV
photoreceptor and a bZIP TF domain empowers Aureo to serve as a natural optogenetic tool
(Hepp, Trauth, Hasenjäger, et al., 2020) (Duan et al., 2018) (Jerng et al., 2021). In contrary to
commonly-sighted domain topology in most LOV photoreceptors, the inverse effector-sensor
arrangement in all Aureo is similar to that in several eukaryotic TFs (Matiiv & Chekunova,
2018) — a feature suitable for developing synthetic sensory TFs (Boral et al., 2022) as well
as optogenetic designs (S. Banerjee & Mitra, 2020). While a substantial amount of literature
is available against Aureo LOV sensors especially on their sequence-structure-function
(Mitra et al., 2012) (A. Banerjee, Herman, Kottke, et al., 2016) (Hepp, Trauth, Hasenj, et al.,
2020) (Heintz & Schlichting, 2016), structure based mechanistic insights is lacking for the
effector bZIPs. And till date, no experimental structure corresponding to full-length or even
the bZIP-linker-LOV module of Aureo is available in PDB.   Therefore, a comprehensive
sequence-structure characterization of this effector bZIP is attempted using several *in-silico*
tools e.g. for phylogenetic studies, network theory, modelling, docking and molecular
dynamics (MD) simulation to understand the formers' uniqueness and significance.   We
include all available varieties of Aureo_bZIPs and compare them with other traditional bZIP
TFs found across plants and opisthokonts to understand the uniqueness and evolutionary
significance. We further validate our *in-silico* results and assess the DNA binding ability of
Aureo_bZIPs using *in-vitro* electrophoretic mobility shift assay (EMSA) after cloning,
overexpression and protein purification. In this study, we attempt to draw an effective
comparative between the bZIP domains of Aureos and the rest *via* a holistic sequence-
structure-function characterization.

2. Materials and Methods

2.1. Retrieval of bZIP sequences from different organisms: The amino acid sequences of Aureos and other representative bZIP proteins from plants and opisthokonts were collected from the UniProt (Apweiler et al., 2018) protein sequence database. BLASTP (Altschul et al., 1990) search was also performed against the database of non-redundant protein sequences using available bZIP sequences as queries. A few sequences, especially those of Aureos from *Phaeodactylum sp*. were obtained from the available literature (Costa et al., 2013). Amino acid sequences of the 78 bZIP proteins from plants, 69 bZIPs from opisthokont species and 26 Aureo bZIP sequences were shortlisted for further analyses. The respective domain boundaries of the bZIPs were finalized either from 'conserved domain' analysis of NCBI and Pfam, or from PSI-BLAST or domain BLAST of NCBI.

2.2 Sequence comparison and molecular evolutionary analysis: A total of 173 bZIP sequences thus retrieved, were subjected to multiple sequence alignment using Clustal omega (Sievers et al., 2011) algorithm — first group-wise and then collectively. This gave us a preliminary understanding on the subtle differences in bZIP sequences that could lead to differences in DNA binding activity. The sequences were further considered for the construction of a phylogenetic tree using MEGA X programme (Kumar et al., 2018). The corresponding accession numbers of all the database entries of these bZIP proteins along with the names and source organisms are listed in the table (**SI-2**). For molecular evolutionary analysis, all 173 bZIP sequences were first aligned using MUSCLE algorithm, inbuilt in MEGA X. The most rational tree of all the selected bZIP proteins was drawn using the neighbor joining (NJ) method, which is simple and fast. Poisson model, which considers equality of substitution rates among all the sites and assumes equal frequencies of amino acids at the time of correction of the multiple substitutions at the same sites, was followed as the substitution method. The phylogeny was tested through bootstrap analysis where the number of performances of bootstrap replications for each node was kept at 1000. The redundant and partial sequences were discarded. Only the reviewed and annotated sequences were considered. The guide tree was then represented and labeled by the online phylogenetic tree manipulation tool iTOLv5. The clades in red and green represent the maximum and minimum values of bootstrap respectively. A covariance analysis on Aureo bZIPs were further performed using the GREMLIN (Kamisetty et al., 2013) server. Default parameters like 4 iterations, >75% query coverage and 10^{-10} /10^{-4} E-value cut-off were chosen as appropriate for this study.

2.3 Residue Interaction Graphs to understand bZIP-DNA interactions: The protein-DNA

complex structure can be represented as residue interaction graph (RIG) or an all-atom

network, where nodes (amino acids/nucleotide residues) are linked by edges within a given

distance cut off (e.g. 4.5 Å). To investigate further on the bZIP-DNA complexes, RIGs were

first generated from eight representative homo/heterodimeric bZIP co-crystal structures from

all the bZIP subfamilies, using the Biographs module of Biopython (Cock et al., 2009) and

NetworkX (Hagberg et al., 2008). Eight structures representing different bZIP subgroups are

- 2WTY (MAF), 1JNM (JUN-CRE complex), 1GD2 (PAP), 1YSA (GCN4), 1FOS (JUN-

FOS heterodimer), 1NWQ (CEBP), 1DH3 (CREB) and 2H7H (JUN-AP1 complex). The

RIGs were generated considering all atom-pair contacts within a given distance of 4.5 Å.

Centrality measures as simple as degree centrality, geodesic/shortest path based closeness

and betweenness centrality have been extremely helpful to identify influential nodes from

residue-level interaction data (Deb et al., 2020). However, to capture all communication at

the residue level, we calculated eigenvector centrality (EC) of the constructed RIGs. How

well connected a node is, in terms of its connectivity with other well-connected nodes of a

network, can be measured by EC. The weighted sum of the centralities of all nodes, which

are connected to node *'i'* by an edge *'A_{ij}'*, is defined as EC of node *'i'* (*C_i*) (Negre et al.,

2018)- $C_i = \varepsilon^{-1}\sum_{j=1}^{n}A_{ij}c_j$, where *'c'* denotes the eigenvector associated with eigenvalue ε

of '*A*'.

The distribution of EC was plotted next. We studied the value of Z (X) = (X-<X>)/σ, where

<X> and σ represent mean and standard deviation respectively. For subsequent analysis, we

therefore considered only those values of X which are associated with Z (X) > 1.5.

2.4 Modeling and docking of Aureo bZIPs - comparison with other bZIPs: Till date, no

experimental 3D structure corresponding to the bZIP region of Aureo is available in RCSB

PDB. Hence, the bZIP sequences, representatives from all five subtypes of Aureos sourced

from *Ectocarpus/Saccharina* sp, were individually modeled using SWISS-MODEL

(Waterhouse et al., 2018). The best models, selected on the basis of best Ramachandran

parameters, were next energy minimized using Chimera (Pettersen et al., 2004). The energy

minimized structures were next subjected to molecular docking analysis using NPDock

(Tuszynska et al., 2015). Review of literature suggests that the bZIP domain of Aureo binds

to a consensus DNA sequence TGACGT, named 'Aureo-box'(Matiiv & Chekunova, 2018)

(Takahashi et al., 2007). So, for the *in silico* docking of modeled Aureo_bZIPs with DNA,

'Aureo-box'(TGACGT) containing CRE (cAMP responsive element) DNA (A. Banerjee,

Herman, Kottke, et al., 2016) from co-crystal structure of CREB bZIP-CRE DNA [(PDB ID:

1DH3, (Schumacher et al., 2000) was used as the DNA substrate /ligand. Preliminary

knowledge on protein-DNA interaction interface, as acquired from sequence-structure

superposition with solved bZIP-DNA co-crystal structures and residue interaction graphs was

used as input in docking. The energy parameters, interface interactions as well as buried

surface area on the best scoring structures of Aureo_bZIP-DNA complexes were calculated

using PDBePISA (Evgeny, 2010). In order to perform a detailed comparative study between

aureochromes and the rest bZIPs, the hydrogen bonding and polar interactions at the basic

region of bZIP-DNA interfaces of all were analyzed using Chimera (Pettersen et al., 2004)

and Pymol (Delano, 2002). *In silico* mutagenesis was performed in few docked structures to

understand the functional as well as evolutionary significance of N-terminal basic region

residues of Aureo_bZIPs.

2.5 System Setup, Molecular Dynamics Simulations and Post-Simulation Methodologies:

Following phylogeny and docking analyses, unique placement of histidine was noticed

exclusively at the basic region of Aureos and not in others. In order to investigate the specific

contribution of this residue in DNA binding activity, Molecular Dynamics (MD) Simulations

were performed using the GROMACS (version 2020.1) (Pronk et al., 2013) (Van Der Spoel

et al., 2005). The docked structure of *Ectocarpus siliculosus* Aureo1_bZIP-DNA complex

(*Es*Aureo1_wild) and its *in-silico* mutant versions (*Es*Aureo1_HA and *Es*Aureo1_HS) were

considered as initial files for MD simulation. AMBER99SB-ILDN force field was applied.

The docked bZIP-DNA complex was solvated using TIP3P water molecules and was limited

into a cubic lattice of dimensions 1.2 x 1.2 x 1.2 nm^3. The structures had positive charges at

normal physiological pH conditions, thus the addition of chloride (Cl$^-$) ions ensured the

simulation systems' electrical neutrality. The entire molecular structure was then relaxed by

performing energy minimization, starting with the steepest descent and followed by the

conjugate gradient algorithm. To remove the solute-solute and solute-solvent steric clashes

and allow the system to adapt to its environment stably, a harmonic restraint was imposed on

the solute particles keeping the solvent particles unrestrained, for over 50000 energy

minimisation steps for both algorithms. Before moving forward, the potential energy (E$_{pot}$) of

the system was checked to be a negative value, since it is correlated with the sufficient

stability of the system required for simulation. The equilibration process was conducted in

two steps. First, the isothermal-isochoric or NVT (constant number of particles-volume-

temperature) ensemble, followed by an isothermal-isobaric or NPT (constant number of

particles-pressure-temperature) ensemble was conducted for 100 ps each, with position

restraining for the non-water residues and constraints of the H-bonds. A leapfrog integrator

was used to analyze the dynamic system numerically. The Berendsen thermostat algorithm

was used for re-scaling the particle velocities in the simulation model to relax the system and

regulate the temperature inside the box to be at around 300 K. The Particle Mesh Ewald

(PME) method was utilized for rapid stochastic relaxation of the system to equilibrium and to

calculate long-range electrostatics. The pressure fluctuation of the simulation system was

accounted for by running an average over the entire length, which was compared with the

reference pressure set as 1 bar and an isothermal compressibility allowance of 4.5 x 10-5

bar^{-1} for water. The LINCS (LINear Constraint Solver) algorithm was used for bond

(involving hydrogens) restraining. An integration time step of 2 fs was applied and the log

file/ coordinate update was done every 1 ps. The system, thus prepared, was finally exposed

to the subsequent forces and ran for 60 ns (20 ns at a stretch for 3 consecutive times), at 300

K temperature.

The data obtained post-simulation (as .xvg files) were plotted using the XMGRACE plotting

tool. PyMOL (Delano, 2002) software was used for visual assessment of the bZIP-DNA

complexes during the simulatory period. The GROMACS in-built analysis package (Van Der

Spoel et al., 2005) was extensively utilized to elucidate all the analytical data for 60 ns MD

simulation run, which were compared for the wild-type and the mutant variants.

2.6 Cloning of wild type Aureochrome-1 from Ectocarpus siliculosus: The pUC-57

plasmids harboring the codon optimized synthetic gene corresponding to *Ectocarpus*

siliculosus aureochrome1 (*Es*Aureo1; Uniprot accession no.- D7FMH8) was purchased. PCR

amplification of the bZIP-linker-LOV sequence of wild type *Es*Aureo1 (*Es*Aureo1_Wild)

DNA sequence was carried out using the following forward primer (FP) and reverse primer

(RP) respectively-5'CTAGGCTAGCATGAAAGATCTGACCGAAGAACAGCGCATTG3';

5'AGCTGAGCTCTTATTTGCTCGGGCCCGCCAGCTG3'. The *Es*Aureo1_wild was then

cloned into the pET28a expression vector at the NheI/SacI restriction sites.

2.7 Site directed mutagenesis: To uncover the role of unique histidine substitution at the

basic region of Aureos, His362 residue in *Es*Aureo1_wild was next mutated to alanine and

serine. These mutations led to the formation of *Es*AureoHA and *Es*AureoHS respectively.

Following primers were used to generate *Es*AureoHA and *Es*AureoHS constructs

respectively-

*Es*AureoHA_FP:5'CGCAACCGCGAAGCGGCGAAACGCAGC3', *Es*AureoHA_RP:5'GC

TGCGTTTCGCCGCTTCGCGGTTGCG3';*Es*AureoHS_FP:5'CGCAACCGCGAAAGCGC

GAAACGCAGC3' ; *Es*AureoHS_RP:5'GCTGCGTTTCGCGCTTTCGCGGTTGCG3'.

2.8 Overexpression and purification of WT and mutant proteins: *Es*Aureo1_wild and its mutatants - *Es*AureoHA and *Es*AureoHS were transformed into the *Escherichia coli* C43 strain to express the wild type protein along with its mutants. Colonies obtained through transformation were then inoculated into the kanamycin containing Luria-Bertani (LB) (*HiMedia*) broth cultures and grown overnight at 37˚C. In each case, 1% inoculum was transferred to the fresh media and grown at 37˚C till OD_{600} reached 0.6. Following IPTG (*Promega*) induction, the cells were grown in the dark at 21˚C overnight. Bacterial cells were then harvested by centrifugation at 8,000 rpm for 10 minutes. The cell pellet was then dissolved in lysis buffer (20 mM Tris, pH 8; 10 mM NaCl; 10% Glycerol) and incubated with protease inhibitor cocktail (*Sigma*), a pinch of lysozyme (*SRL*) and phenylmethylsulfonyl fluoride (PMSF, *Promega*). The dissolved cell pellet was then subjected to ultrasonic vibrations (*Hielscher*) and high-speed centrifugation (*Thermo Fisher Scientific*) at 12500 rpm for 1 hour at 4˚C. Following centrifugation, the supernatant was collected and subjected to Ni-NTA affinity chromatography (Qiagen). Following loading and wash, hexa-histidine tagged protein fractions were eluted using the elution buffer (20 mM Tris, pH 8; 10 mM NaCl; 100 mM Imidazole; 10% Glycerol). The eluted fractions containing the desired protein were then subjected to PD-10 desalting column (*Sigma*) for the removal of excess imidazole. The desalted protein was then concentrated using Amicon® Ultra Centrifugal Filter. The quality of protein was checked using SDS-PAGE (*Bio-Rad*). Further purification by Heparin/gel filtration chromatography resulted in extremely poor yield. Therefore, affinity purified and desalted protein fractions were used for EMSA studies, carried out at least in triplicates from multiple batches of protein purification. The concentration of the wild type as well as the mutant proteins was specifically determined with respect to flavin [at 450 nm, considering the extinction co-efficient of flavin (12500 $M^{-1}cm^{-1}$) using UV-Vis spectroscopy (*Shimadzu*)].

2.9 Electrophoretic Mobility Shift Assay: HPLC purified and lyophilized single stranded oligonucleotides of 24bp containing 'Aureo-box' sequence, 'TGACGT', (FP: 5'TGTCCTTGGCTGACGTCAGCCAAGCAC3';RP:5'GTGCTTGGCTGACGTCAGCCA AGGACA3') were purchased (*IDT*). Following resuspension in annealing buffer (10 mM Tris, pH 8.0; 20 mM NaCl) the complementary DNA strands in equimolar amounts were annealed by heating at 95˚C for 2 mins followed by gradual overnight cooling. The binding

buffer (50 mM Tris, pH 8.0; 50 mM NaCl; 1.25 mM MgCl$_2$; 0.01 mg/mL BSA; 20% glycerol) was next added to the double stranded DNA (dsDNA) to achieve a final concentration of 0.5 uM. Each protein was then serially diluted and added to the solution of dsDNA and binding buffer mixture. A total of nine samples were prepared for protein-DNA mixtures with varying protein concentrations. The substrate DNA minus protein was considered as negative control. All samples including the control were incubated on ice under BL for 45 minutes prior to the commencement of electrophoresis. The electrophoretic run was performed using Bio-Rad's Mini-PROTEAN system at 4˚C under BL-illuminated condition. The samples were resolved at 150 Volts at 4˚C for one hour. Upon the completion of electrophoresis, the native gel was stained with SYBR Gold nucleic acid stain (*Thermo Fisher Scientific*) for 45 minutes and imaged in the Bio-Rad ChemiDoc MP system.

3. Results

3.1. Sequence retrieval and analysis:

A total number of 173 bZIP sequences inclusive of 78 from plants, 69 from opisthokonts and 26 from Aureos are considered [**SI-2**] herein to assess amino acid sequence composition/conservation/variations. The conservation of leucines/non-polar amino acid residues at every a/d position in the C-terminal dimerization domain/zipper region of Aureos corroborate well with other bZIPs. However, we do observe few remarkable variations at the N-terminal basic region of Aureo_bZIPs when compared with others. The basic region responsible for DNA recognition possesses a signature sequence of nine amino acid residues [N-(X)$_7$-K/R] [**SI-3**]. A histidine is uniquely found at the 4th position [N-(X)$_2$-**H**-(X)$_4$-R] of the basic region in Aureos [**Figure-1.a**] unlike others. While in plant-bZIPs serine is the invariable occupant (except plant bZIP subfamily D, F) [**SI-3.**]; alanine is mostly present in opistokont bZIPs, excluding CREB, CREB3, ATF6 [**SI-3**]. Therefore, the basic region sequence of opisthokont and plant bZIPs can be broadly designated as [N-(X)$_2$-**A**-(X)$_4$-K/R] and [N-(X)$_2$-**S**-(X)$_4$-R] respectively. Notably in Aureo2, histidine is substituted with leucine [**Figure-1.a**]. An in-depth investigation into all the Aureo further divulged that *Sj*Aureo2 despite being designated as a type-2 Aureo, possesses histidine [**SI-3**]. This observation can be justified from the phyletic pattern [**Figure-1.b**], where *Sj*Aureo2 clubs with other Aureo paralogs rather than the Aureo2 cluster. As neither histidine nor leucine is found at the debatable 4[th] position in any other bZIP across three kingdoms of life, their persistence in aureos could bear an evolutionary significance. Again, the presence of threonine as the 8[th] residue in Aureo2 basic region [N-(X)$_2$-LA-(X)$_2$-**TR**] [**Figure-1.a**] is also notable considering

the presence of serine at the homologous position in most other Aureo bZIPs. Incidentally,

the 8th position of all bZIPs is occupied mostly by serine or cysteine. A covariation of

histidine at 4th and serine at the 8th position is further detected in the co-variation analysis

using GREMLIN (Kamisetty et al., 2013)[**Figure-1.c**]. This probably explains why both

histidine and serine are simultaneously replaced by leucine and threonine respectively in

Aureo-2 paralogs.

*3.2 Phylogeny estimation and course of bZIP evolution***:**

The evolutionary position and the phylogenetic relationship of all selected 173 bZIP proteins

from plants, opisthokonts and Aureos can be inferred from the phylogenetic tree construction

[**Figure-2**]. However, the deeply rooted divergence within the ancient nodes may not have

been detected due to the comparatively shorter length of bZIP domains. Nevertheless, three

distinct lineages from a common ancestor have been observed. The first lineage includes

BATF, PAR, and E4BP4. The second lineage comprises opisthokont bZIP families such as

CREB2, ATF2, CEBP, ATF3, FRA, and FOS along with the viral bZIP family Zta. The

remaining plant and opisthokont bZIPs are found to cluster together with Aureo bZIPs in the

third lineage. However, the last common ancestor of the third lineage further diverges into

two distinct clades. One is solely represented by all Aureo_bZIPs, while the other splits

further into two branches: the opisthokont bZIPs (e.g., PAP, GCN4, JUN, MAF, CREB3,

CREB, and ATF6) and all plant bZIPs (e.g., A, B, C, D, E, F, G, H, I, and S). However, the

most appropriate sister clade or immediate ancestor of Aureo bZIPs could not be ascertained.

The phylogenetic tree of all bZIPs illustrates the tendency of Aureo_bZIPs to club together in

the same clade, forming a distinct sub-lineage without intermingling with bZIPs either from

plants or opisthokonts. Most Aureo paralogs within the Aureo-bZIP clade align by subtype,

with a few exceptions, while the remaining ones lose their distinctiveness and cluster

randomly [**Figure- 2**]. In summary, the phylogeny estimation of bZIPs from multiple

genomes not only identifies the ancestor-descendant relationships among different bZIP

members but also highlights the evolutionary relationship of Aureo_bZIPs with the rest.

3.3. Network analysis and bZIP dimer stability towards substrate DNA interaction:

Aureo bZIPs are unique in the sense that they are BL-regulated. In fact, the LOV → bZIP

signaling relies on an allosteric pathway (A. Banerjee, Herman, Serif, et al., 2016). The

resulting conformational changes at the bZIP-substrate DNA interface further promotes DNA

binding without much alteration in DNA binding affinity. In absence of full-length

experimental structure of any Aureo at dark and light states, it would be difficult to precisely
identify the intervening residues that transduce light energy into DNA binding. Despite
significant homology observed between bZIP and LOV domains individually with other
bZIP/LOV members respectively, the existence of heterogeneous linkers with varying
sequence composition and length further complicates this problem. While bZIP dimerization
is an essential pre-requisite of DNA binding activity, Aureos contain the only bZIPs that are
connected with a BL-sensor. Therefore, one crucial question is: how does an Aureo bZIP
compare with other bZIPs? We therefore constructed all-atom networks from eight co-
crystallized bZIP-DNA complexes. While centrality measures are frequently used for
network analysis, in this study, we particularly focused on Eigenvector Centrality (EC) –
especially given their implications in allostery (Negre et al., 2018). EC that considers both the
number of connections made by a node as well as the flow of information associated with
them, EC has been successfully used to identify residues responsible for allosteric
communication in enzymes (Negre et al., 2018). We did not consider Aureo model structures
for RIG analysis because structural artifacts could affect the result. We therefore calculated
EC from networks constructed upon eight co-crystal structures of homo/heterodimeric bZIP-
DNA complexes [PDB IDs: 1DH3, 1FOS, 1GD2, 1JNM, 1NWQ, 1YSA, 2H7H and 2WTY]
[**SI-4**]. Residues with high EC values with $Z(X) > 1.5$ are only considered and marked in
green [**Figure-3**]. Additionally, two to three residues from monomeric units showing highest
EC values are color coded in red to check any further significance. The residues involved in
DNA binding are marked in blue [**Figure-3**]. In all cases, residues with high EC values are
found to congregate at the zipper region, albeit at variable distances from the basic region.
Residues with highest EC values specifically occupy 'a' and 'd' positions of a heptad in
every single case, as seen in the figure as well as multiple sequence alignment [**Figure-3.b**].
We find this result extremely interesting considering well-known hydrophobic interaction
between residues at positions 'a' and 'd' towards leucine zipper dimerization, its stability as
well as their contribution in allosteric regulation (Cruz et al., 2023). In terms of EC, after 'a'
and 'd', the next important positions happen to be 'e' and 'g'. As emphasized in our earlier
study (Khamaru et al., 2024), the electrostatic compatibility in between 'e' and 'g' can
singularly govern bZIP homo/heteromerization(C. Vinson et al., 2002)(Krylov et al., 1994).
Sequence comparison reveals that the homologous residues in Aureos corresponding to high
EC values in other bZIPs are mostly conserved (similar if not identical) except for the 'd'
position in the 4th heptad. This particular position in Aureo is occupied by polar or even
charged amino acid residues, whereas this is always leucine in other bZIPs. Overall, our EC

analysis combined with sequence alignment reveals that Aureo zippers and the rest are
generally comparable. However, precise contribution of important heptad residues (a/d/e/g)
in allosteric regulations and/or signal transduction, as predicted by EC analysis, is subjected
to further experimental validations.

3.4. Decoding bZIP-DNA interactions - insights from modeled and docked Aureo subtypes:
Following *Es*Aureo1, *Es*Aureo2, *Es*Aureo3, *Es*Aureo4 and *Sj*Aureo5 homodimer modeling
and energy minimization, each one was subjected to *in silico* docking experiments with the
substrate DNA sequence encompassing a TGACGT core which is known to be the binding
sequence for Aureo bZIPs, termed as 'Aureo-box'. The best scoring complex structures were
selected and bZIP-DNA interfaces were characterized based on hydrogen bonding/salt
bridge/van der Waals' interactions, interface area and energy. The bZIP-DNA structures of
all five Aureo-bZIP representatives are depicted in [**Figure-4**]. The accuracy of our docked
structures was further assessed by checking their RMSD when superimposed with the
modeled/docked structure obtained through alpha fold. To test the reliability of our docked
structures, we conducted PDBePISA (Protein Interfaces, Surfaces and Assemblies) analysis
and compared them with experimental data on bZIP-DNA complex [1DH3 and 1FOS
representing homo- and heterodimer-DNA interactions respectively]. The emphasis was
given on three parameters – the interface area, solvation energy and number of non-covalent
interactions, precisely hydrogen bonding and salt bridge interactions. While the interface area
between monomeric units of protein chains ranges from ~900 Å^2-1200 Å^2 (homodimer to
heterodimer respectively), that between protein-DNA ranges from ~300-400 Å^2. A negative
value of gain in solvation free energy (Δ^iG) upon interface formation in both protein-
protein/DNA complex formations indicate positive affinity between them and the
hydrophobic interface. In the case of all homodimeric Aureos [*Es*Aureo1-4, *Sj*Aureo5], the
interface area is very similar to homodimeric 1DH3/1JNM, with an average of 915 Å^2.
Aureo-DNA interface area ranges in between ~350-450 Å^2 with a very similar number of
hydrogen bonding/salt bridge interactions like in experimental structures. Further, the Δ^iG for
both protein-protein/DNA in Aureo-DNA complexes are not only negative but the values are
very similar to those of native co-crystal structures. The N-terminal basic regions being the
key determinants behind DNA binding specificity of bZIPs, they were first screened for polar
including hydrogen bonding interactions with the DNA bases. At first a detailed analysis of
the hydrogen bonding interactions involving the basic region and the substrate DNA from
experimentally derived co-crystal structures of the bZIP-DNA complexes [**SI-5**] was

performed to determine the segment of bZIP crucial for the DNA binding and to identify the crucial residues within the basic region that facilitate DNA binding. Analysis of the bZIP-DNA interaction patterns of all the co-crystal structures revealed multifarious specificities of bZIP proteins despite overall structure-sequence conservation. Since, the five subtypes of Aureo_bZIPs exhibited minor difference in residue composition of the signature motif of bZIP meant for DNA binding, all the five Aureo_bZIPs were docked with DNA [**Figure-4**] and were then screened for the hydrogen bonds and polar interactions present between the basic region residues and the substrate DNA bases to understand whether histidine takes part in the interaction with DNA. In *Es*Aureo1-DNA complex, His362 (NE2) is found to interact with dT'8. *Es*Aureo4 also exhibits interaction of His50 (NE2) with DNA. His153 (NE2) of *Sj*Aureo5 interacts with OP2 of dG'6. All the polar interactions detected at the protein-DNA interface of the docked structures are color coded and presented in tabular format [**SI-6**].

Characteristic of all bZIPs, in Aureo-DNA complexes too, the interactions between asparagine and arginine/lysine and substrate DNA in the signature $[N-(X)_7-K/R]$ sequence are conserved. Additionally, interactions from basic amino acids towards substrate binding are also prominent. Perhaps the most notable finding is the involvement of the histidine in DNA binding activity in all cases. As already mentioned, the absence of histidine at the homologous position of plant/opisthokont bZIPs led to further speculation about its possible evolutionary impact on the origin and evolution of all Aureo in the bZIP superfamily.

3.5 Exploring the importance of Histidine substitution by *in silico* mutageneis, phylogeny and co-evolution analysis:

To further understand the role of histidine at the N-terminal basic region of the Aureo bZIPs, *in-silico* mutagenesis in sequence and structure was conducted. Histidine was mutated to serine (as in plant bZIPs), alanine (as in opisthokont bZIPs) and leucine (as in Aureo2) to find out the consequences in terms of evolution as well as DNA binding respectively. No significant polar interaction or hydrogen bonding is found when His362 is mutated to alanine. When leucine replaces His362/His153 in the docked *Es*Aureo1 / *Sj*Aureo5-DNA complex*,* no interactions with any of the nucleotides can be seen. His51 in *Es*Aureo3, upon substitution by leucine, remains in close contact with dG'6 of the substrate DNA. Similar situation arises in *Es*Aureo4, where His50 upon mutation to leucine remains in close contact with dG'6. However, no polar contacts could be observed in both these cases. His362 of *Es*Aureo1 on being mutated to serine, comes in close contact with dC`7. While *Es*Aureo2's Leu197 to

serine mutation maintains dT'8 contact, His51→Ser in *Es*Aureo3, shows possible interaction with the N7 of DG'6. Incorporation of serine at the homologous histidine positions in *Es*Aureo4 and *Sj*Aureo5, fails to show any significant interaction. We must acknowledge that the experimental 3D structure of the Aureo-DNA complex could be most reliable. However, our modeled docked structures indicate an active role of His in Aureo-box DNA binding activity. Single residue substitution can influence DNA binding affinity, as noticed in one of our earlier collaborative studies – FD1/FD2 belonging to plant bZIP subfamily A, exhibited a 10-fold difference in DNA binding activity (Dutta et al., 2021). To elaborate further on single histidine substitution, not only we performed MD simulations but also conducted *in vitro* experiments with *Es*Aureo1_wild, *Es*Aureo1_HA and *Es*Aureo1_HS constructs – we discuss the results in the subsequent section of this manuscript. We also wanted to check the evolutionary significance of histidine at this specific position. Therefore, we re-constructed the phylogenetic tree after altering the histidine/leucine position in Aureo by serine (similar to plant bZIPs) and alanine (similar to opisthokont bZIPs). Very interestingly, upon serine substitution, Aureos shift closer to the plant bZIPs sharing a common evolutionary lineage with all the plant bZIP subgroups except D, F and H. All the opisthokont bZIPs are clustered in a different lineage in this case [**Figure-5.a**]. Perhaps this is important to mention here that plant bZIPs D, F and H differ from the remaining plant bZIPs at the crucial 4th/8th position of basic region signature sequence. D and F plant bZIPs possess alanine instead of serine [N(X)$_2$$\underline{A}(X)_5$], similar to opisthokont bZIPs [**SI-3**]. F and H bZIPs possess Y and A respectively at the 8th position instead of typical serine [N(X)$_6$$\underline{Y/A}$R] [**SI-3**]. Upon mutation to alanine, the bZIPs of Aureo aligned in a lineage with opisthokont and plant bZIPs, albeit differently from the wild types. [**Figure-5.b**]. All through we considered a bootstrap value of 1000 to obtain statistically confident data. And indeed, the presence of histidine in Aureo does make a difference in their evolutionary origin. Co-evolution studies conducted with *Es*Aureo1 bZIP with an E-value cut-off 10^{-10} followed by Weblogo analysis shows distinct presence of histidine [**Figure-1.c**] in Aureo subtypes. Upon increasing the E-value to 10^{-4} to accommodate more bZIP sequences however led to complete disappearance of histidine and appearance of serine/alanine at the homologous position [**Figure-5.c**].

3.6 Probing the Dynamics of bZIP-DNA Interactions through Molecular Dynamics (MD) Simulations:

MD simulations were performed to understand the consistency of interaction between *Es*Aureo1_wild, *Es*Aureo1_HA and *Es*Aureo1_HS and the substrate DNA, albeit over a short

time scale. We particularly monitored the behaviour of residue 362 (His, Ala and Ser

respectively) throughout the MD run. Thus, the post-simulation analyses conducted include

RMSD (Root Mean Square Deviation), RMSF (Root Mean Square Fluctuation), the radius of

gyration (Rg), Number of Contacts (within 3 nm cutoff) and the number of hydrogen bonds.

3.6.1. RMSD

The RMSD of the protein backbone (Cα-atoms) was analyzed from the simulation data and

plotted to compare structural stability across variants. Notably, *Es*Aureo1_wild exhibited

consistently lower deviation compared to the mutant variants (*Es*Aureo1_HA and

*Es*Aureo1_HS), with average RMSD values of 0.346 nm, 0.395 nm, and 0.414 nm,

respectively [**Figure-6.a**]. While all variants showed similar deviation trends at the outset, the

wild type demonstrated a marked decrease in deviation around 5 ns, contrasting with the

rising trends observed in the mutants. A transient surge in deviation for *Es*Aureo1_wild

occurred near 17 ns, followed by alignment with mutant trends between 25 and 30 ns. After a

minor fluctuation around 30 ns, the wild type resumed its low-deviation profile, maintaining

this trend until the end of the simulation (60 ns). Notably, the *Es*Aureo1_HS variant exhibited

a significant increase in RMSD between 50 and 60 ns, stabilizing at a higher deviation. These

observations underscore the structural resilience of *Es*Aureo1_wild compared to its mutants.

3.6.2 RMSF

To determine the flexibility of the protein residues, especially His362, the RMSF data was

elucidated and plotted against the simulation runtime. A subtle variation, commonly

suspected to arise from differences in binding dynamics and solvent interactions, is generally

expected between protein chains C and D of the homodimer; hence, they were analyzed and

plotted separately [**Figure-6.c,d**]. In both polypeptide chains of Aureo-dimer, His362 of

*Es*Aureo1_wild shows the least flexibility, followed by Ala362 of *Es*Aureo1_HA and then

Ser362 of *Es*Aureo1_HS.

3.6.3 Radius of gyration

The Rg represents the overall compactness of the protein by measuring the average distance

of its atoms from an imaginary axis passing through its center of gravity. Analysis of the Rg

curves [**Figure-6.b**] reveals that *Es*Aureo1_wild exhibits greater compactness throughout the

simulation compared to its mutant counterparts. Up to the 30 ns mark, all variants display a

similar trend, but beyond this point, the wild-type variant maintains a consistency of its Rg,

indicating a more compact structure with minimal fluctuation. In contrast, the *Es*Aureo1_HA

and *Es*Aureo1_HS mutants show increased fluctuations and less compact conformations over

time. Notably, the *Es*Aureo1_HS variant exhibits a sudden and significant drop in Rg after

the 50 ns mark, reflecting a collapse in compactness that correlates with the sharp rise in
RMSD observed in the same time frame. This behavior underscores the distinct dynamic
stability of the wild variant compared to its mutants till the end of the simulation (60 ns).

3.6.4. State of hydrogen bonds

The variation in the number of hydrogen bonds made by His362 with the DNA in both the
wild and mutant variants was obtained and was plotted against the simulation time (60000ps).
According to the graphs, the residue His362 of *Es*Aureo1_wild shows the maximum number
of hydrogen bonds, fluctuating between 0 to 4 [**Figure-7.a**]. The residue Ser362 of
*Es*Aureo1_HS shows a moderate number of H-bonds throughout the simulation, fluctuating
between 0 to 3 [**Figure-7.b**]. The residue Ala362 of EsAureo1_HA shows no H-bonds
throughout the simulation period with the DNA [**Figure-7.c**]. Hydrogen bonding interactions
were visualized from the post-MD trajectory files, using PyMOL, with a 10ns interval
between each state analyzed for an overall period of 60ns [**Figure-8**].

3.7. In vitro protein-DNA interaction study by Electrophoretic Mobility Shift Assay:

EMSA was performed with the purified and desalted samples of *Es*Aureo1_wild along with
its two mutants - *Es*Aureo_HS and *Es*Aureo_HA to compare their DNA binding ability under
BL. In all cases, the shift from unbound to substrate DNA bound state was observed with the
increase in concentrations of protein [**Figure-9**]. Presence of free DNA bands are indicative
of protein being unbound with its substrate DNA. Gradual disappearance of free DNA bands
and the resultant shift in electrophoretic mobility is indicative of protein-DNA complex
formation. The highest concentration of *Es*Aureo1_wild at which an unbound DNA band was
evident first was 0.04 μM [**Figure-9.a**]. In the case of *Es*Aureo_HS [**Figure-9.b**] and
*Es*Aureo_HA [**Figure-9.c**], the first presence of an unbound DNA band showed up at
concentrations 0.076 μM and 0.082 μM respectively. Hence, although all of the variants
manifest DNA binding, the result of EMSA indicates that *Es*Aureo1_wild has a higher DNA
binding affinity. And the unique presence of histidine at the basic region of Aureo bZIPs
seems justifiable for their specific interaction with Aureo-box substrates.

4. Discussion

The global network of TFs comprising interactions with specific substrate DNA or partner
protein molecules regulates one of the most fundamental processes of life - gene expression.
Besides the possibility of simple/auto-regulation, a TF can participate in feed forward

loops/feedback loops (FFLs/FBLs)(Alon, 2007) as homodimer or heterodimer, in complex and dynamic gene expression regulatory units [**Figure-10**]. While the orderliness and sequence composition of the basic and fork regions of bZIPs play a crucial role in DNA binding specificity, sequence organization and stability of heptad repeats in the zipper region is equally important. Even a subtle change in heptad sequence can impact dimerization specificity and stability, and the resulting manifestation could be significant. Choosing partners from the same or different group of bZIPs, respectively for homo- and hetero-dimerization can generate substantial functional diversity. The 3D structure of a protein can be considered as a network of nodes (amino acid residues) connected via edges (non-covalent interactions) and the relative importance of the individual amino acid residues can be obtained by calculating different topological parameters of RIG. Therefore, apart from characterizing protein-protein/DNA interface in terms of bonding interactions, surface area and energy, we conducted all-atom network analysis. Centrality measures like degree, closeness and betweenness effectively captures influential nodes in a network. In this study, we emphasized on measuring EC values considering its potential to detect nodes important for making connections and information flow, especially allostery. Interestingly, residues with highest EC values are always at 'a' and 'd' positions of a heptad in all eight representative co-crystal structures. Importance of hydrophobic residues at the strategic positions necessary for dimerization and consequent enhancement of DNA binding activity has been shown for c-Fos mutants(Porte et al., 1997). Incorporation of isoleucine at the 2nd heptad not only promoted homodimerization but also conferred DNA binding specificity towards TRE substrate in c-Fos mutants. While EC calculations for Aureos await determination of experimental 3D structures, sequence comparison does indicate that Aureo bZIPs are mostly similar to other bZIPs at least in these specific nodes except 'd' residue in heptad 4. Measurement of EC undoubtedly reiterates the importance of hydrophobic residues towards maintaining stable bZIP dimerization, signal transfer and thereby promoting DNA binding activity. Therefore, insights obtained from EC measurements could be helpful for designing synthetic sensory TF in the near future.

The BL-responsive sensory TFs like Aureo, the pivot of this study, comes with an additional advantage. Such an association of one major group of eukaryotic TF (here, bZIPs) with a light sensor (here, LOV) is rare and fascinating. Our study therefore attempts to draw a comparative analysis of Aureos and the other conventional bZIP proteins without LOV sensors. With distinct differences noticed, it remains to be clarified whether or not these

variations arose through evolutionary acquisition of the sensor domain to confer
physiological advantages to the host organisms. Though bZIPs are supposed to have far
reaching origin, the expansion and diversification of the bZIP superfamily has been found to
be positively correlated with the evolution of eukaryotes (Jindrich & Degnan, 2016). The
number of bZIP homologs as well as the complexity of bZIPs have increased over the time in
the process of evolution. Initially bZIPs were supposed to evolve independently in 4
eukaryotic lineages – planta, opisthokonta, amoebozoa, and heterokonta (Jindrich & Degnan,
2016). Among them 17 bZIP subfamilies within opisthokont, namely, YAP, MAF, XBP1,
CNC, BATF, CREB, CREB-2, OASIS, PAR, E4BP4, ATF-2, ATF-4, ATF-6, FOS, JUN,
C/EBP, PAP (Amoutzias et al., 2007) (Fujii et al., 2000) and ten distinct subclasses (A, C, D,
G and S) of plant bZIPs (Jakoby et al., 2002) (Corrêa et al., 2008) are evident. Whereas in
heterokontophyta (**SI-7]**, the phylum comprising of 12 families under the algal group-4
as proposed in the algal classification by Robert Lee (Lee, 2018), the most abundant bZIP
transcription factor is Aureo (Rayko et al., 2010). It has been observed in the evolutionary
analysis performed by us that all the bZIPs of different Aureo homologs together form a
different clade albeit shared the ancestor with opishtokont bZIP subfamilies- JUN, MAF,
CREB, CREB3, PAP, GCN4 and plant bZIP subfamilies. But substitution of histidine results
in an assembly of Aureo-bZIPs predominantly with plant bZIPs sharing a common ancestral
lineage. This study therefore investigates how nuances at the residue level are manifested in
the phyletic distribution of bZIP members, including the evolution of Aureos and their
subtypes.

The bZIPs of the metazoans recognize 6 different consensus DNA-binding sites- c-AMP
responsive element (CRE- TGACGTCA), TPA responsive element (TRE- ATGAGTCAT),
CAAT box (TTGCGCAA), PAR binding sites (TTACGTAAT), Maf recognition element
(MARE- CTGATCAGC) and CRE-like elements (TGACGTGC) (Amoutzias et al., 2007)
(Rodríguez-Martínez et al., 2017). Again, CRE, TRE and YAP (yeast activator protein) sites
can be recognized by fungal bZIPs. Plant bZIPs recognize G-box (CACGTG), A-box
(TACGTA), C-box (GACGTC), S-box (TGACGTG), D-box (TGACGTG) (Wang et al.,
2018) etc. We conducted *in-silico* and *in-vitro* studies, using Aureo-box containing CRE
(TGACGTCA) as the substrate DNA [sequence taken from X-ray structure of CREB bZIP-
CRE complex, PDB ID: 1DH3]. The docking, MD simulation as well as EMSA studies were
performed not only to check the DNA binding efficiency of the Aureo_bZIPs (wild type and
mutants) but also to explore the precise role of histidine in DNA binding. Both theory and

experiments indicate significance of histidine at the basic region of aureochrome. It can be
concluded that His362 plays a key role in positioning the *Es*Aureo1 bZIP to the core ACGT
sequence by favorably interacting with its flanking sequence. Changing this amino acid to
Ser (as in plants) and Ala (as in opisthokonts) leads to a significant decrease in polar
interactions [**Figure-7,8**] [**SI-8**] and therefore impacts the stability of the complex [**Figure-**
6.a] [as found from MD Simulation study]. However, at the same time, the overall binding
strength of all the constructs remains comparable [as confirmed by the EMSA studies]
[**Figure-9**]. This makes it clearer that His is very strongly related to the Aureo box, and thus
the slightest change in this recognition box (as seen in opisthokonts and plants) makes this
amino acid residue site prone to change. This provides an answer for the causality of co-
evolution. It is worth reiterating that Aureos are light-responsive TFs and therefore should
function like a photo-switch. Studies regarding the kinetics of binding and the width of
spectrum of the changes in functionality of the bZIP due to variation in substrate DNA
sequence, are still an area requiring further work. The binding specificity of a TF to DNA is
intricately governed by its interaction with the minor groove geometry and regional
electrostatics. Among the three mutants analyzed through simulation, *Es*Aureo1_wild has the
maximum DNA binding ability with the highest number of hydrogen bonds present at the
bZIP-DNA interface. *Es*Aureo1_HS ranks in between wild type and *Es*Aureo1_HA. The
alanine mutant modeled after opisthokont bZIP proteins, emerged as distinct, with Ala362
failing to interact with the flanking residues of the ACGT core. While the number of non-
covalent interactions, e.g. hydrogen bonds, determines DNA binding specificity(Lin & Guo,
2019), prompt and selective binding is eventually reflected in protein-DNA association
kinetics. Molecular docking, MD simulation as well as EMSA studies reveal that the
placement of histidine bespoke 'Aureo-box' specific interaction. However, it would be highly
interesting to see whether histidine makes any difference in association kinetics of Aureos
towards substrate recognition/binding upon BL illumination. Strength and specificity of DNA
binding interactions as well as variable association kinetics can be adopted further for
developing precision optogenetic tools.

Perhaps another important observation that we made in this study is the unique presence of
histidine in all Aureo bZIPs, except Aureo2 from several stramenopiles. Moreover, Aureo2
also possesses non-functional LOV (Takahashi et al., 2007). Out of eleven conserved amino
acids in the flavin-binding pocket, nine are conserved in Aureo2. The reason behind such
differences in Aureo2 is elusive – it is possible that the sensor domain of Aureo2 could be an

intermediate between non-light sensing PAS (Per-ARNT-Sim, ancestor to LOV) and LOV.

Our study further unveiled that histidine co-evolves with serine at the 8th position of the

signature bZIP sequence. Interestingly in Aureo2, both 4th and 8th positions are differently

occupied by leucine and threonine respectively. Despite such marked changes in the basic

region, Aureo2 clubs with the rest Aureos in the phylogenetic tree and not with

plants/opisthokonts. However, upon single substitution from leucine to alanine, Aureo2 shifts

lineage and sides with CEBPs of opisthokonts, while retaining proximity with other Aureos.

Considering the presence of non-functional sensor domain and close association with

opisthokont bZIPs just upon single amino acid substitution, we speculate that Aureo2 might

have evolved first among the Aureos. During the course of evolution, possibly with the gain

of function mutation, Aureos have finally acquired the ability to sense light and perform light

dependent transcriptional activities. And perhaps to preserve this very important property,

several other Aureo paralogs emerged through gene duplication events.

In summary, the bZIPs of aureochromes share close structural and functional similarities with

other bZIP TFs found across plants and animals, exhibiting comparable dimerization

properties and DNA-binding affinities to substrate DNA. Consequently, this bZIP integrated

with its blue light sensor, can be utilized in plants and animals either as a genetically encoded

recombinant protein driven by a synthetic promoter or as a transgene expressed under native

promoters in target cells to achieve tissue-specific gene regulation via light stimulation.

Furthermore, orthogonal manipulation through engineered gene regulatory circuits can

facilitate non-invasive control over existing regulatory networks. Our previous study

indicates that the aureochrome-bZIP monomer dimerizes with bZIP monomers within the

aureochrome family as well as with other non-aureochrome bZIPs (Khamaru et al., 2024). As

such, introducing light-sensitive aureochrome-bZIPs into genetically modified transgenic

lines or transiently transformed cells, can enable light-mediated regulation of developmental

pathways and stress responses by interacting with the inherent bZIP proteins of plants or

animals. Additionally, all the bZIP proteins regulate crucial developmental processes of

plants and animals like- growth, cell division, and differentiation can be further modulated by

adjusting the duration of gene activation or repression through fine tuning of the light sensor

module associated with these bZIPs. Interestingly, all the subclasses of plant bZIP proteins

that bind to promoter or regulator present upstream of the target genes, recognize the ACGT

sequence of DNA. As Aureo-bZIPs also recognizes ACGT sequences, it seems to be a

promising optogenetic tool for regulating plant gene expression and influencing the existing

bZIP dependent metabolic and developmental pathways within plant kingdom. Thus,
characterization of these exclusive light regulated bZIPs not only gives insight to the
transcriptional regulation pattern of the lower group of plants i.e. the heterokont algae, but
shed light into the scope of optogenetic regulation of plant genes.

Figure Legends

Figure-1:

a. Multiple sequence alignment of the bZIP domains of Aureos
b. Phylogenetic tree comprising all the aureochrome homologs.
c. Weblogo result of the co-evolution study of aureochrome1 at E-value cut-off 10^{-10}.
696
Figure-2:

Phylogenetic tree of the bZIPs from plants, opisthokonts and Aureos. Different
subgroups of bZIPs are represented by different colours. Highest to lowest bootstrap
values are colour coded as green and red respectively.
Figure-3:

a. DNA-bZIP complex where the residues identified as important from eigenvector
centrality measure are colour coded. Green colour denotes EC value higher than 1.5,
Red colour indicates knob like structure (d & a) with the highest EC value, blue
colour represents DNA binding residues.
b. Multiple sequence alignment showing DNA binding residues (coloured blue) and
residues with higher EC values (coloured green/red).

Figure-4:

Docked bZIP structures of *Ectocarpus siliculosus* (*Es*) aureochrome1 (gray),
aureochrome2 (pale green), aureochrome3 (yellow), aureochrome4 (pink) and
Saccharina japonica (*Sj*) aureochrome5 (teal) bZIPs in complex with substrate DNA
sequences containing 'Aureo-box', TGACGT motif. Interactions between amino acid
residues of the DNA recognizing the basic region of bZIPs and the corresponding
nucleotide of substrate DNA are labeled.

Figure-5:

a. Re-construction of phylogenetic tree upon mutation of Aureos' histidine to serine
(plant like)
b. Re-construction of phylogenetic tree upon mutation of Aureos' histidine to alanine
(opisthokont like)
c. Weblogo result obtained from co-evolution study at E-value cut-off 10^{-4}.
721

Figure 6:

a. Root Mean Square Deviation (RMSD) in nm plotted for *Es*Aureo1_wild (green),
*Es*Aureo1_HS (blue) and *Es*Aureo1_HA (orange) protein backbone for 60 ns
timescale of MD simulation.
b. Radius of Gyration (Rg) in nm for *Es*Aureo1_wild (green), *Es*Aureo1_HS (blue) and
*Es*Aureo1_HA (green) plotted against the simulation time of 60 ns.
c. Root Mean Square Fluctuation (RMSF) of the protein residues for *Es*Aureo1_wild
(orange), *Es*Aureo1_HS (green) and *Es*Aureo1_HA (yellow), as obtained from 60ns
simulation data, for Protein chain C.
d. Root Mean Square Fluctuation (RMSF) of the protein residues for *Es*Aureo1_wild
(orange), *Es*Aureo1_HS (green) and *Es*Aureo1_HA (yellow) for Protein Chain D as
obtained from 60 ns simulation data.
*[EsAu1 has been used as the abbreviation of EsAureo1]

Figure 7:

a. Average number of intermolecular hydrogen bonds between His362 in
*Es*Aureo1_wild and DNA during 60 ns timescale.
b. Average number of intermolecular hydrogen bonds between Ser362 in *Es*Aureo1_HS
and DNA
c. Average number of intermolecular hydrogen bonds between Ala362 in *Es*Aureo1_HA
and DNA.
*[EsAu1 has been used as the abbreviation of EsAureo1]

Figure 8:

Post-simulation visualization of Residue 362 and its hydrogen bonding interaction
with DNA (shown in yellow dashed line), at timesteps of 10, 20, 30, 40, 50 and 60ns

(from left to right). a. *Es*Aureo1_wild in green, b. *Es*Aureo1_HS in blue, c.

*Es*Aureo1_HA in orange. *[*Es*Au1 has been used as the abbreviation of *Es*Aureo1]

Figure 9:

In vitro protein DNA interaction study of Aureochrome_bZIPs through electrophoretic

mobility shift assay-a. *Es*Aureo1_wild, b. *Es*Aureo_HS, c. *Es*Aureo_HA.

Figure 10:

Cartoon and diagrammatic representation of the sensory transcription regulation network,

where Aureochrome is perceived to act as a sensory, blue-light regulated transcription factor

regulating gene expression. The concept of sensory transcription regulation network is

adopted from Network motifs: theory and experimental approaches by Alon U. [ref. 56,

doi:10.1038/nrg2102]

Acknowledgement

MK and AD acknowledge University Grants Commission (UGC) and Department of

Biotechnology (DBT), Government of India respectively, for their doctoral research

fellowships. DM acknowledges funding from DBT, Government of India

(BT/PR26435/BRB/10/1627/2017) for supporting the research work during the initial phase

and SERB, Govt of India (CRG/2023/006463) at the present phase. The authors acknowledge

the central instrumentation facility as well as the computer facility of the department

supported by DBT-BUILDER and DST-FIST programs respectively. The authors also thank

Soumen Roy, Bose Institute - Kolkata for important discussions on network analysis.

Declaration of interests

The authors declare no conflict of interest. No competing interest.

Author contribution

Madhurima Khamaru: Data curation, Investigation, Methodology, Formal analysis,

Validation, Visualization, Writing – original draft, Writing – review and editing; **Debarshi**

Bose: Investigation, Methodology, Formal analysis, Visualization, Writing – original draft;

Anwesha Deb: Investigation, Methodology; **Devrani Mitra**: Conceptualization, Supervision,

Funding acquisition, Project administration, Investigation, Methodology, Formal analysis,

Validation, Visualization, Writing – original draft, Writing – review and editing.

Data availability statement:

All data generated or analyzed during this study are included in the main text article as well as in the supplementary information. Further enquiries can be directed to the corresponding author.

References

Alon, U. (2007). Network motifs: theory and experimental approaches. *Nature Reviews Genetics 2007 8:6*, *8*(6), 450–461. https://doi.org/10.1038/nrg2102

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local alignment search tool. *Journal of Molecular Biology*, *215*(3), 403–410. https://doi.org/10.1016/S0022-2836(05)80360-2

Amoutzias, G. D., Veron, A. S., Weiner, J., Robinson-Rechavi, M., Bornberg-Bauer, E., Oliver, S. G., & Robertson, D. L. (2007). One billion years of bZIP transcription factor evolution: Conservation and change in dimerization and DNA-binding site specificity. *Molecular Biology and Evolution*, *24*(3), 827–835. https://doi.org/10.1093/molbev/msl211

Apweiler, R., Bairoch, A., Wu, C. H., Barker, W. C., Boeckmann, B., Ferro, S., Gasteiger, E., Huang, H., Lopez, R., Magrane, M., Martin, M. J., Natale, D. A., O'Donovan, C., Redaschi, N., & Yeh, L. S. L. (2018). UniProt: the universal protein knowledgebase. *Nucleic Acids Research*, *46*(5), 2699–2699. https://doi.org/10.1093/NAR/GKY092

Bader, A. G., & Vogt, P. K. (2006). Leucine Zipper Transcription Factors: bZIP Proteins. In *Encyclopedic Reference of Genomics and Proteomics in Molecular Medicine* (pp. 964–967). Springer Berlin Heidelberg. https://doi.org/10.1007/3-540-29623-9_2180

Banerjee, A., Herman, E., Kottke, T., & Essen, L. O. (2016). Structure of a Native-like Aureochrome 1a LOV Domain Dimer from Phaeodactylum tricornutum. *Structure*, *24*(1), 171–178. https://doi.org/10.1016/j.str.2015.10.022

Banerjee, A., Herman, E., Serif, M., Maestre-Reyna, M., Hepp, S., Pokorny, R., Kroth, P. G., Essen, L.-O., & Kottke, T. (2016). Allosteric communication between DNA-binding and light-responsive domains of diatom class I aureochromes. *Nucleic Acids Research*, *44*(12), 5957–5970. https://doi.org/10.1093/nar/gkw420

Banerjee, S., & Mitra, D. (2020). Structural Basis of Design and Engineering for Advanced Plant Optogenetics. *Trends in Plant Science*, *25*(1), 35–65. https://doi.org/10.1016/j.tplants.2019.10.002

Boral, A., Khamaru, M., & Mitra, D. (2022). Designing synthetic transcription factors: A structural perspective. *Advances in Protein Chemistry and Structural Biology*, *130*. https://doi.org/10.1016/BS.APCSB.2021.12.003

Busch, S. J., & Sassone-Corsi, P. (1990). Dimers, leucine zippers and DNA-binding domains. *Trends in Genetics*. https://doi.org/10.1016/0168-9525(90)90071-D

Cock, P. J. A., Antao, T., Chang, J. T., Chapman, B. A., Cox, C. J., Dalke, A., Friedberg, I., Hamelryck, T., Kauff, F., Wilczynski, B., & De Hoon, M. J. L. (2009). Biopython: freely available Python tools for computational molecular biology and bioinformatics. *Bioinformatics (Oxford, England)*, *25*(11), 1422–1423. https://doi.org/10.1093/BIOINFORMATICS/BTP163

Corrêa, L. G. G., Riaño-Pachón, D. M., Schrago, C. G., Vicentini dos Santos, R., Mueller-Roeber, B.,
& Vincentz, M. (2008). The Role of bZIP Transcription Factors in Green Plant Evolution:
Adaptive Features Emerging from Four Founder Genes. *PLoS ONE*, *3*(8), e2944.
https://doi.org/10.1371/journal.pone.0002944

Costa, B., Sachse, M., Jungandreas, A., Bartulos, C. R., & Gruber, A. (2013). Aureochrome 1a Is
Involved in the Photoacclimation of the Diatom Phaeodactylum tricornutum. *PLoS ONE*, *8*(9).
https://doi.org/10.1371/journal.pone.0074451

Cruz, P., Paredes, N., Asela, I., Kolimi, N., Molina, J. A., Ramírez-Sarmiento, C. A., Goutam, R.,
Huang, G., Medina, E., & Sanabria, H. (2023). Domain tethering impacts dimerization and
DNA-mediated allostery in the human transcription factor FoxP1. *The Journal of Chemical*
Physics, *158*(19). https://doi.org/10.1063/5.0138782

Deb, A., Grewal, R. K., Roy, S., & Mitra, D. (2020). Residue interaction dynamics in Vaucheria
aureochrome1 light-oxygen-voltage: Bridging theory and experiments. *Proteins*, *88*(12), 1660–
1674. https://doi.org/10.1002/PROT.25984

Delano, W. (2002). *The PyMOL Molecular Graphics System*.

Deppmann, C. D., Acharya, A., Rishi, V., Wobbes, B., Smeekens, S., Taparowsky, E. J., & Vinson,
C. (2004). Dimerization specificity of all 67 B-ZIP motifs in Arabidopsis thaliana : a comparison
to Homo sapiens B-ZIP motifs. *Nucleic Acids Research*, *32*(11), 3435–3445.
https://doi.org/10.1093/NAR/GKH653

Duan, L., Hope, J. M., Guo, S., Ong, Q., François, A., Kaplan, L., Scherrer, G., & Cui, B. (2018).
Optical Activation of TrkA Signaling. *ACS Synthetic Biology*, *7*(7), 1685–1693.
https://doi.org/10.1021/ACSSYNBIO.8B00126/SUPPL_FILE/SB8B00126_SI_001.PDF

Dutta, S., Deb, A., Biswas, P., Chakraborty, S., Guha, S., Mitra, D., Geist, B., Schäffner, A. R., &
Das, M. (2021). Identification and functional characterization of two bamboo FD gene homologs
having contrasting effects on shoot growth and flowering. *Scientific Reports*, *11*(1).
https://doi.org/10.1038/S41598-021-87491-6

Evgeny, K. (2010). Crystal contacts as nature's docking solutions. *Journal of Computational*
Chemistry, *31*(1), 133–143. https://doi.org/10.1002/JCC.21303

Fujii, Y., Shimizu, T., Toda, T., Yanagida, M., & Hakoshima, T. (2000). *letters Structural basis for*
the diversity of DNA recognition by bZIP transcription factors.

Glover, J. N. M., & Harrison, S. C. (1995). Crystal structure of the heterodimeric bZIP transcription
factor c-Fos–c-Jun bound to DNA. In *Nature*. https://doi.org/10.1038/373257a0

Grigoryan, G., & Keating, A. E. (2006). Structure-based Prediction of bZIP Partnering Specificity.
Journal of Molecular Biology, *355*(5), 1125–1142. https://doi.org/10.1016/j.jmb.2005.11.036

Hagberg, A. A., Schult, D. A., & Swart, P. J. (2008). Exploring network structure, dynamics, and
function using NetworkX. *7th Python in Science Conference (SciPy 2008)*, 11–15.
https://www.bibsonomy.org/bibtex/272096c6553d7057409fe78ca698eb332/mschuber

Hakoshima, T. (2005a). Leucine Zippers. In *Encyclopedia of Life Sciences*. John Wiley & Sons, Ltd.
https://doi.org/10.1038/npg.els.0005049

Hakoshima, T. (2005b). Leucine Zippers. In *Encyclopedia of Life Sciences*.
https://doi.org/10.1038/npg.els.0005049

Harbury, P. B., Zhang, T., Kim, P. S., & Alber, T. (1993). A switch between two-, three-, and four-
stranded coiled coils in GCN4 leucine zipper mutants. *Science (New York, N.Y.)*, *262*(5138),

1401–1407. https://doi.org/10.1126/SCIENCE.8248779

Heintz, U., & Schlichting, I. (2016). Blue light-induced LOV domain dimerization enhances the
affinity of aureochrome 1a for its target DNA sequence. *ELife*, *5*(JANUARY2016).
https://doi.org/10.7554/ELIFE.11860

Hepp, S., Trauth, J., Hasenj, S., Bezold, F., Essen, L.-O., & Taxis, C. (2020). *An Optogenetic Tool for*
Induced Protein Stabilization Based on the Phaeodactylum tricornutum Aureochrome 1a
LighteOxygeneVoltage Domain. https://doi.org/10.1016/j.jmb.2020.02.019

Hepp, S., Trauth, J., Hasenjäger, S., Bezold, F., Essen, L. O., & Taxis, C. (2020). An Optogenetic
Tool for Induced Protein Stabilization Based on the Phaeodactylum tricornutum Aureochrome
1a Light–Oxygen–Voltage Domain. *Journal of Molecular Biology*.
https://doi.org/10.1016/j.jmb.2020.02.019

Huang, Y., Wang, L., Zheng, M., Zheng, L., Tong, Y., & Li, Y. (2014). Overexpression of
NgAUREO1, the gene coding for aurechrome 1 from Nannochloropsis gaditana, into
Saccharomyces cerevisiae leads to a 1.6-fold increase in lipid accumulation. *Biotechnology*
Letters, *36*(3), 575–579. https://doi.org/10.1007/s10529-013-1386-0

Huysman, M. J. J., Fortunato, A. E., Matthijs, M., Costa, B. S., Vanderhaeghen, R., Van den Daele,
H., Sachse, M., Inzé, D., Bowler, C., Kroth, P. G., Wilhelm, C., Falciatore, A., Vyverman, W.,
& De Veylder, L. (2013). AUREOCHROME1a-Mediated Induction of the Diatom-Specific
Cyclin dsCYC2 Controls the Onset of Cell Division in Diatoms (Phaeodactylum tricornutum).
The Plant Cell, *25*(1), 215–228. https://doi.org/10.1105/tpc.112.106377

Jakoby, M., Weisshaar, B., Dröge-Laser, W., Vicente-Carbajosa, J., Tiedemann, J., Kroj, T., & Parcy,
F. (2002). bZIP transcription factors in Arabidopsis. *Trends in Plant Science*, *7*(3), 106–111.
https://doi.org/10.1016/S1360-1385(01)02223-3

Jerng, H. H., Patel, J. M., Khan, T. A., Arenkiel, B. R., & Pfaffinger, P. J. (2021). Light-regulated
voltage-gated potassium channels for acute interrogation of channel function in neurons and
behavior. *PLOS ONE*, *16*(3), e0248688. https://doi.org/10.1371/JOURNAL.PONE.0248688

Ji, C., Mao, X., Hao, J., Wang, X., Xue, J., Cui, H., & Li, R. (2018). Analysis of bZIP transcription
factor family and their expressions under salt stress in Chlamydomonas reinhardtii. *International*
Journal of Molecular Sciences, *19*(9). https://doi.org/10.3390/ijms19092800

Jindrich, K., & Degnan, B. M. (2016). The diversification of the basic leucine zipper family in
eukaryotes correlates with the evolution of multicellularity Genome evolution and evolutionary
systems biology. *BMC Evolutionary Biology*, *16*(1), 1–12. https://doi.org/10.1186/s12862-016-
0598-z

Kamisetty, H., Ovchinnikov, S., & Baker, D. (2013). Assessing the utility of coevolution-based
residue-residue contact predictions in a sequence- and structure-rich era. *Proceedings of the*
National Academy of Sciences of the United States of America, *110*(39), 15674–15679.
https://doi.org/10.1073/PNAS.1314045110/SUPPL_FILE/SD05.XLSX

Khamaru, M., Nath, D., Mitra, D., & Roy, S. (2024). Assessing Combinatorial Diversity of
Aureochrome Basic Leucine Zippers through Genome-Wide Screening. *Cells, Tissues, Organs*,
213(2), 133–146. https://doi.org/10.1159/000527593

Kroth, P. G., Wilhelm, C., & Kottke, T. (2017). An update on aureochromes_ Phylogeny –
mechanism – function. *Journal of Plant Physiology*, *217*, 20–26.
https://doi.org/10.1016/j.jplph.2017.06.010

Krylov, D., Mikhailenko, I., & Vinson, C. (1994). A thermodynamic scale for leucine zipper stability

and dimerization specificity: e and g interhelical interactions. *The EMBO Journal*, *13*(12), 2849. https://doi.org/10.1002/J.1460-2075.1994.TB06579.X

Kumar, S., Stecher, G., Li, M., Knyaz, C., & Tamura, K. (2018). MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. *Molecular Biology and Evolution*, *35*(6), 1547–1549. https://doi.org/10.1093/MOLBEV/MSY096

Lee, R. E. (2018). Phycology. *Phycology*. https://doi.org/10.1017/9781316407219

Lin, M., & Guo, J. T. (2019). New insights into protein–DNA binding specificity from hydrogen bond based comparative study. *Nucleic Acids Research*, *47*(21), 11103–11113. https://doi.org/10.1093/NAR/GKZ963

Mann, M., Serif, M., Wrobel, T., Eisenhut, M., Madhuri, S., Flachbart, S., Weber, A. P. M., Lepetit, B., Wilhelm, C., & Kroth, P. G. (2020). The Aureochrome Photoreceptor PtAUREO1a Is a Highly Effective Blue Light Switch in Diatoms. *IScience*, *23*(11), 101730. https://doi.org/10.1016/J.ISCI.2020.101730

Matiiv, A. B., & Chekunova, E. M. (2018). Aureochromes – Blue Light Receptors. *Biochemistry (Moscow)*, *83*(6), 662–673. https://doi.org/10.1134/S0006297918060044

Mitra, D., Yang, X., & Moffat, K. (2012). Crystal structures of aureochrome1 LOV suggest new design strategies for optogenetics. *Structure*. https://doi.org/10.1016/j.str.2012.02.016

Negre, C. F. A., Morzan, U. N., Hendrickson, H. P., Pal, R., Lisi, G. P., Patrick Loria, J., Rivalta, I., Ho, J., & Batista, V. S. (2018). Eigenvector centrality for characterization of protein allosteric pathways. *Proceedings of the National Academy of Sciences of the United States of America*, *115*(52), E12201–E12208. https://doi.org/10.1073/PNAS.1810452115

Pettersen, E. F., Goddard, T. D., Huang, C. C., Couch, G. S., Greenblatt, D. M., Meng, E. C., & Ferrin, T. E. (2004). UCSF Chimera—A visualization system for exploratory research and analysis. *Journal of Computational Chemistry*, *25*(13), 1605–1612. https://doi.org/10.1002/JCC.20084

Podust, L. M., Krezel, A. M., & Kim, Y. (2001). Crystal structure of the CCAAT box/enhancer-binding protein beta activating transcription factor-4 basic leucine zipper heterodimer in the absence of DNA. *The Journal of Biological Chemistry*, *276*(1), 505–513. https://doi.org/10.1074/JBC.M005594200

Pogenberg, V., Consani Textor, L., Vanhille, L., Holton, S. J., Sieweke, M. H., & Wilmanns, M. (2014). Design of a bZip Transcription Factor with Homo/Heterodimer-Induced DNA-Binding Preference. *Structure*, *22*(3), 466–477. https://doi.org/10.1016/j.str.2013.12.017

Porte, D., Oertel-Buchheit, P., John, M., Granger-Schnarr, M., & Schnarr, M. (1997). DNA binding and transactivation properties of Fos variants with homodimerization capacity. *Nucleic Acids Research*, *25*(15), 3026–3033. https://doi.org/10.1093/NAR/25.15.3026

Pronk, S., Páll, S., Schulz, R., Larsson, P., Bjelkmar, P., Apostolov, R., Shirts, M. R., Smith, J. C., Kasson, P. M., Van Der Spoel, D., Hess, B., & Lindahl, E. (2013). GROMACS 4.5: a high-throughput and highly parallel open source molecular simulation toolkit. *Bioinformatics*, *29*(7), 845–854. https://doi.org/10.1093/BIOINFORMATICS/BTT055

Rayko, E., Maumus, F., Maheswari, U., Jabbari, K., & Bowler, C. (2010). Transcription factor families inferred from genome sequences of photosynthetic stramenopiles. *The New Phytologist*, *188*(1), 52–66. https://doi.org/10.1111/J.1469-8137.2010.03371.X

Rodríguez-Martínez, J. A., Reinke, A. W., Bhimsaria, D., Keating, A. E., & Ansari, A. Z. (2017). Combinatorial bZIP dimers display complex DNA-binding specificity landscapes. *ELife*, *6*.

946   https://doi.org/10.7554/eLife.19272

947 Schumacher, M. A., Goodman, R. H., & Brennan, R. G. (2000). *The Structure of a CREB*
948   *bZIPSomatostatin CRE Complex Reveals the Basis for Selective Dimerization and Divalent*
949   *Cation-enhanced DNA Binding**. https://doi.org/10.1074/jbc.M007293200

950 Sievers, F., Wilm, A., Dineen, D., Gibson, T. J., Karplus, K., Li, W., Lopez, R., McWilliam, H.,
951   Remmert, M., Söding, J., Thompson, J. D., & Higgins, D. G. (2011). Fast, scalable generation of
952   high-quality protein multiple sequence alignments using Clustal Omega. *Molecular Systems*
953   *Biology*, *7*(1), 539. https://doi.org/10.1038/MSB.2011.75

954 Sinden, R. R. (1994). DNA–Protein Interactions. *DNA Structure and Function*, 287–325.
955   https://doi.org/10.1016/B978-0-08-057173-7.50013-4

956 Takahashi, F., Yamagata, D., Ishikawa, M., Fukamatsu, Y., Ogura, Y., Kasahara, M., Kiyosue, T.,
957   Kikuyama, M., Wada, M., & Kataoka, H. (2007). AUREOCHROME, a photoreceptor required
958   for photomorphogenesis in stramenopiles. *Proceedings of the National Academy of Sciences*,
959   *104*(49), 19625–19630. https://doi.org/10.1073/pnas.0707692104

960 Tuszynska, I., Magnus, M., Jonak, K., Dawson, W., & Bujnicki, J. M. (2015). NPDock: a web server
961   for protein–nucleic acid docking. *Nucleic Acids Research*, *43*(Web Server issue), W425.
962   https://doi.org/10.1093/NAR/GKV493

963 Van Der Spoel, D., Lindahl, E., Hess, B., Groenhof, G., Mark, A. E., & Berendsen, H. J. C. (2005).
964   GROMACS: Fast, flexible, and free. *Journal of Computational Chemistry*, *26*(16), 1701–1718.
965   https://doi.org/10.1002/JCC.20291

966 Vinson, C., Myakishev, M., Acharya, A., Mir, A. A., Moll, J. R., & Bonovich, M. (2002).
967   Classification of Human B-ZIP Proteins Based on Dimerization Properties. *Molecular and*
968   *Cellular Biology*, *22*(18), 6321–6335. https://doi.org/10.1128/MCB.22.18.6321-6335.2002

969 Vinson, C. R., Hai, T., & Boyd, S. M. (1993). Dimerization specificity of the leucine zipper-
970   containing bZIP motif on DNA binding: Prediction and rational design. *Genes and*
971   *Development*, *7*(6), 1047–1058. https://doi.org/10.1101/gad.7.6.1047

972 Wang, Y., Zhang, Y., Zhou, R., Dossa, K., Yu, J., Li, D., Liu, A., Mmadi, M. A., Zhang, X., & You,
973   J. (2018). Identification and characterization of the bZIP transcription factor family and its
974   expression in response to abiotic stresses in sesame. *PLOS ONE*, *13*(7), e0200850.
975   https://doi.org/10.1371/journal.pone.0200850

976 Waterhouse, A., Bertoni, M., Bienert, S., Studer, G., Tauriello, G., Gumienny, R., Heer, F. T., De
977   Beer, T. A. P., Rempfer, C., Bordoli, L., Lepore, R., & Schwede, T. (2018). SWISS-MODEL:
978   homology modelling of protein structures and complexes. *Nucleic Acids Research*, *46*(W1),
979   W296–W303. https://doi.org/10.1093/NAR/GKY427

Figure 1

Tree scale: 0.1

bootstrap
- 0
- 0.25
- 0.5
- 0.75
- 1

Colored ranges

- Aureochrome bZIP
- GCN4
- JUN
- PAP
- FOS
- FRA
- ATF3
- CEBP
- ATF2
- CREB2
- PAR
- BATF
- E4BP4
- ATF6
- CREB3
- CREB
- MAF
- CNC-bZIP
- D-type of Plants
- F-type of Plants
- E-type of Plants
- I-type of Plants
- A-type of Plants
- C-type of Plants
- S-type of Plants
- B-type of Plants
- H-type of Plants
- G-type of Plants
- Viral Zta

Figure 2

Figure 3

```
                gabcdefgabcdefgabcdefgabcdefgabcd
PAP GD2    SSKRKAQNRAAQRAFRKRKEDHLKALETQVVTLKELHSSTTLENDQLRQKVRQLEEEL
C-FOS
1FOS       RRIRRERNKMAAAKSRNRRRELTDTLQAETDQLEDEKSALQTEIANLLKEKEKLEFIL
C-JUN
1FOS       AERKRMRNRIAASKSRKRKLERIARLEEKVKTLKAQNSELASTANMLREQVAQLKQKV
GCN4
1YSA       AALKRARNTEAARRSRARKLQRMKQLEDKVEELLSKNYHLENEVARLKKLVGER----
ViralJUN
2H7H       AERKRNRIAASKSRKRKLERIARLEEKVKTLKAQNSELASTANMLREQVAQLKQKV
CEBP
1NWQ       YRVRRERNNIAVRKSRDKAKQRNVETQQKVLELTSDNDRLRKRVEQLSRELDTLRG--
CREB
1DH3       REVRLMKNREAARESRRKKKEYVKSLENRVAVLENQNKTLIEELKALKDLYSHK----
C-JUN
1JNM       AERKRMRNRIAASKSRKRKLERIARLEEKVKTLKAQNSELASTANMLREQVAQLKQKV
MAFB
2WTY       QKRRTLKNRGYAQSCRYKRVQQKHHLENEKTQLIQQVEQLKQEVSRLARERDAYKVKS
EsAureo1   RIERRERNREHAKRSRVRKKFLLDSLQRSVDAIQAENESLKGSIVGSLGERGREL---
EsAureo2   AKKRRERNRVLARRTRLRKKFFFQSLQQQVARLQRENERLKGIVTTRCPDSVGEIL--
EsAureo3   RLDRRERNREHAKRSRVRKKFLLESLQKSVTSLQEENEKLRGAIRSNLGPEEAKEL--
EsAureo4   KLERRERNREHAKRSRIRKKFMLECLQEQLLAMRKQNLALRQVVKEHMPDEASTV---
```

Heptad-1 Heptad-2 Heptad-3 Heptad-4

Figure 4

Figure 5

Plant bZIPs

Opisthokont bZIPs

Aureochrome bZIPs

a

b

c

Figure 8

Figure 9

a

b

c

bZIP

LOV   LOV

Regulation of gene expression

BLUE
LIGHT

N —— bZIP —— A' α | FMN LOV | J α —— C

Figure 10

Simple Regulation

Negative Auto-regulation

Positive Auto-regulation

Gene Z

Gene Z   Gene Z   Gene Z   Gene Z

Incoherent and Coherent FFLs   Dynamics in FFLs

Homo-dimer as Single
Input Module

Z1   Z2   Z3

Hetero-dimer as Single
Input Module

Z1   Z2   Z3

Z1   Z2   Z3

Participation in multiple-input, multiple-output module