Wnt signalling, cell fate determination and anteroposterior polarity of the skate gill arch skeleton

Jenaid M. Rees[1], Victoria A. Sleight[2], Stephen J. Clark[3], Tetsuya Nakamura[4] and J. Andrew Gillis[1,5,*]

[1]Department of Zoology, University of Cambridge, Cambridge CB2 3EJ, United Kingdom

[2]School of Biological sciences, University of Aberdeen, Aberdeen, AB24 3FX

[3]Babraham Institute, Cambridge, United Kingdom

[4]Department of Genetics, Rutgers University

[5]Marine Biological Laboratory, Woods Hole, MA, 02543

*Correspondence: jag93@cam.ac.uk

The gill arch skeleton of cartilaginous fishes (sharks, skates, rays and holocephalans) exhibits anterior–posterior polarity, with a series of appendages (branchial rays) projecting from the posterior margin of the gill arch cartilages. We previously demonstrated in the skate (*Leucoraja erinacea*) that branchial rays derive from a posterior domain of pharyngeal arch mesenchyme, which is responsive to Shh signalling from a distal gill arch epithelial ridge (GAER) signalling centre. However, how branchial ray progenitors are specified exclusively within posterior gill arch mesenchyme is not known. Here we show that the GAER of the skate is of endodermal origin, arises at the endoderm–ectoderm boundary, and is a source of Fgf and Shh signals that are transduced broadly and posteriorly, respectively, within underlying arch mesenchyme. Using RNAseq, we discover that genes encoding several Wnt ligands are expressed in the ectoderm immediately adjacent to the GAER, and that these Wnt signals are transduced largely in the anterior arch mesenchyme. These tissue origin and gene expression features are largely conserved in the hyoid arch of the chick and are therefore likely an ancestral feature of jawed vertebrates. Finally, using pharmacological manipulations in skate, we show that loss of Wnt signalling results in an anterior expansion of Shh signal transduction within pharyngeal arch epithelium, and the formation of ectopic anterior branchial rays. Our findings demonstrate that Wnt signalling restricts chondrogenesis to the posterior gill arch and highlights the importance of signalling interactions at embryonic tissue boundaries for cell fate determination in pharyngeal arches.

Introduction

The pharyngeal arches of vertebrate embryos give rise to much of the craniofacial skeleton, including the skeleton of the jaws and gills in fishes, and of the jaw, auditory ossicles and larynx in amniotes (Graham and Smith, 2001). The arches are formed through iterative outpockets of foregut endoderm that contact the overlying surface ectoderm. The meeting of endoderm and ectoderm generates a series of columns lined laterally by ectoderm, medially by endoderm, and containing a core of mesoderm and neural crest-derived mesenchyme (Graham and Smith, 2001). In fishes, endodermal outpockets fuse with the overlying surface ectoderm, giving rise to the gill slits and the respiratory surfaces of the gill arches (Gillis and Tidswell, 2017). In amniotes, the endodermal outpockets give rise to glandular tissues, such as the tonsils, parathyroid and ultimobranchial glands (Grevellec and Tucker, 2010). In all vertebrates, the largely neural crest-derived mesenchyme of the pharyngeal arches gives rise to the pharyngeal skeleton (Jiang et al., 2002, Kague et al., 2012, Sleight and Gillis, 2020, Couly and Le Douarin, 1990), and receives patterning and polarity information via signals from adjacent epithelia (Veitch et al., 1999, Gillis et al., 2009b, Couly et al., 2002, Brito et al., 2006).

All jawed vertebrates belong to one of two lineages: cartilaginous fishes (sharks, skates, rays and holocephalans) or bony fishes (ray- and lobe-fined fishes, the latter including tetrapods). While the gill arch skeleton of both cartilaginous and bony fishes ancestrally consisted proximally of two principal gill arch cartilages (the epi- and ceratobranchials), the gill arch skeleton of cartilaginous fishes additionally includes a distal series of fine cartilaginous rods called branchial rays (Fig.1A,B). These rays reflect the clear anteroposterior polarity of the gill arch skeleton of cartilaginous fishes, originating along the posterior margin of the epi- and ceratobranchial cartilages, and projecting laterally into the interbranchial septum of each arch (Gillis et al., 2009a). Elasmobranch cartilaginous fishes (sharks, skates and rays) possess five sets of branchial rays, associated with their hyoid (2^{nd} pharyngeal) and first four gill arches. Holocephalans, on the other hand, possess a single set of branchial rays supporting their hyoid arch-derived operculum.

The branchial rays of cartilaginous fishes develop under the influence of the gill arch

epithelial ridge (GAER): a *sonic hedgehog* (*Shh*)-expressing signalling centre located

within the posterior-distal epithelium of the gill arches (Fig.1C,D) (Gillis et al., 2009b).

As the gill arches undergo a prolonged phase of lateral expansion, Shh signalling from

the GAER is asymmetrically transduced within the posterior arch environment, as

evidenced by posterior expression of *Ptc2* in gill arch mesenchyme and epithelium.

This posterior transduction of Shh signalling, in turn, appears to underlie the

anteroposterior polarity of the gill arch skeleton, as branchial rays derive exclusively

from posterior–distal (Shh-responsive/GAER-adjacent) gill arch mesenchyme (Gillis

and Hall, 2016). In the skate, application of exogenous Shh protein within gill arches

is sufficient to induce ectopic branchial ray formation, and targeted or systemic

inhibition of hedgehog signalling using cyclopamine results in branchial ray deletion

(Gillis et al., 2009b). However, whether other signalling mechanisms function

alongside or in conjunction with GAER Shh signalling to establish and maintain gill

arch skeletal polarity remains unexplored.

Here, we show that the GAER of the little skate (*Leucoraja erinacea*) is of endodermal

origin and arises at the endoderm–ectoderm interface of each pharyngeal arch. We

find that the ectoderm immediately adjacent to the GAER expresses several genes

encoding Wnt ligands and that these Wnt signals are transduced in the anterior arch

environment in a pattern that is broadly complementary to the posterior transduction

of GAER Shh signalling. Finally, inhibition Wnt signalling in developing skate embryos

results in an anterior expansion of Shh signalling transduction, and in the formation of

ectopic branchial rays in the anterior gill arch. We propose that Wnt signalling from

pharyngeal arch ectoderm contributes to maintenance of anterior–posterior polarity of

the skate gill arch skeleton by repressing Shh signalling and chondrogenesis to the

posterior gill arch territory.

Figure 1. Overview of skate branchial ray anatomy and *Shh* expression.

(A) Skeletal preparation of a stage (S) 32 skate embryo showing the location of the **(A')** branchial ray and gill arch cartilages *in situ*. **(B)** A dissected gill arch in frontal view shows the articulation of branchial rays with the epi- and ceratobranchial cartilages of the gill arch. **(C)** *Shh* is expressed in the gill arch epithelial ridge (GAER) of the hyoid and first four gill arches (asterisks) of skate embryos. **(D)** mRNA *in situ* hybridisation on paraffin section showing *Shh* expression within the skate GAER (asterisk). Scale bars; A:2mm, C:500μm, D:100μm.

Methods

Embryo harvesting

Skate (*L. erinacea)* embryos were obtained from the Marine Resources Center at the Marine Biological Laboratory in Woods Hole, MA, U.S.A., were reared as described in Gillis *et al.* (2012) and staged according to Ballard *et al.* (1993) and Maxwell et al. (2008). Chick (*G. gallus*) from Brown Bovan Gold hens' eggs (Henry Stewart, UK) were incubated at 38°C, and staged according to Hamburger and Hamilton (1951). Skate embryos were euthanised with an overdose of MS-222 (1g/L in seawater), and all embryos were fixed in 4% paraformaldehyde (Thermofisher) as per Gillis *et al.* (2012).

Sectioning and histochemical staining

Embryos were embedded in paraffin and sectioned at 7μm as described in O'Neill et al. (2007). Sectioned embryos were stained with Masson's Trichrome as per Witten and Hall (2003) or hematoxylin and eosin by clearing in two rinses of Histosol (National Diganostics), rehydration through serial dilutions of ethanol, staining for 15 minutes in Mayers Haematoxylin (Sigma), rinsing in running tap water for 20 minutes, rinsing briefly in 95% ethanol, and staining in 0.1% w/v Eosin Y (Sigma) in 95% ethanol for 2 minutes. Slides were then washed briefly in 95% ethanol, washed briefly with 100% ethanol, cleared with Histosol and mounted with permount (Sigma).

Lineage tracing

Labelling in both skate and chick embryos was performed *in ovo* prior to formation of the pharyngeal arches and labelled embryos were then reared until they possessed established GAER or PEM signalling centres, respectively. For skate, labelling was performed at stage (S)18, and embryos were grown to S25–29. In chick, labelling was performed at HH11, and embryos were grown to HH19–20. In both chick and skate, endodermal labelling was performed by microinjection of CellTracker™ CM-DiI (1,1'-dioctadecyl-3,3,3'3'-tetramethylindocarbocyanine perchlorate; Thermofisher), prepared as per Gillis *et al.* (2012), into the pharyngeal cavity with a pulled glass needle. In chick, ectodermal labelling was performed by application of lipid-soluble CCFSE (5-(and-6)-Carboxy-2',7'-dichlorofluorescein diacetate, succinimidyl ester; Biotium) dye to the surface ectoderm, as per Richardson *et al.* (2012) and Shone and

Graham (2014). Labelled embryos were embedded and sectioned as described

above. Sections were imaged for lineage tracing and subsequently processed for *Shh*

mRNA *in situ* hybridisation (ISH) for identification of the GAER/PEM.

mRNA *in situ* hybridisation

Chromogenic mRNA in situ hybridisation (ISH) was performed on paraffin sections

and in wholemount as described in O'Neill *et al.* (2007), with modifications according

to Gillis *et al.* (2012). ISH probes against skate *Shh* (GenBank EF100667) and *Fgf8*

(GenBank EU574737.1), and against chick *BMP7* (GenBank XM_417496.6), *SHH*

(GenBank NM_204821.1) and *FGF8* (GenBank NM_001012767.1) were generated

by *in vitro* transcription using standard methods. A plasmids for chick *BMP7* was

generously provided by the lab of Claudio Stern (Streit et al., 1998), and for chick *SHH*

and *FGF8* by the lab of Cliff Tabin (Scherz et al., 2004). Following ISH, wholemounts

were rinsed, post-fixed in 4% paraformaldehyde and graded into 75% glycerol for

imaging, while slides were rinsed in PBS, post-fixed in 4% paraformaldehyde and

coverslipped with DAPI-Fluoromount® G (SouthernBiotech). Third-generation mRNA

ISH by hybridisation chain reaction (HCR) was performed as per the Choi *et al* (2018)

protocol for formaldehyde fixed, paraffin-embedded sections, with modifications as per

Criswell and Gillis (2020). Probes, buffers, and hairpins were purchased from

Molecular Instruments (Los Angeles, California, USA). HCR probe set lot numners are

as follows: for skate *Shh* (Lot PRA753), *Ptc2* (Lot PRA754), *Fgf8* (Lot PRA755),

Dusp6 (Lot PRA756), *Wnt2b* (Lot PRE300), *Wnt3* (Lot PRG814), *Wnt4* (Lot PRE301),

Wnt7b (Lot PRE302), *Wnt9b* (Lot PRE303), Notum (Lot PRG817), *Kremen1* (Lot

PRG816), *Axin2* (Lot PRG818), *Apcdd1* (Lot PRG815), *Col2a1* (Lot PRB574) and

Sox9 (Lot PRB571); for chick *SHH* (Lot PRB282), *FGF8* (Lot PRA997), *BMP7* (Lot

PRb283), *WNT2B* (Lot PRG820), *WNT3* (Lot PRG823), *WNT4* (Lot PRG821), *WNT7B*

(Lot PRG819), *WNT9B* (Lot PRG822), *NOTUM* (Lot PRG824), *KREMEN1* (Lot

PRG825), *AXIN2* (Lot PRG827), *APCDD1* (Lot PRG826). All mRNA ISH experiments

were replicated, at minimum, in triplicate.

Pharmacological manipulations

For systemic inhibition of canonical Wnt signalling, experimental and control skate

embryos were reared at 15°C in petri dishes containing 50μM IWR1 (Selleck

chemicals) in artificial seawater or artificial seawater containing an equivalent volume of vehicle (DMSO) only, respectively. IWR1 has been used previously to inhibit canonical Wnt signalling in skate embryos by Nakamura et al. (2015) and in shark embryos by Thiery et al. (2022). For these experiments, IWR1 was diluted to working concentration from a 25mM stock solution in DMSO. Skate embryos were reared in IWR1- or vehicle-containing seawater from S25 until S31/32, with drug or control seawater changes every 48 hours. Once embryos reached the desired stage, they were euthanised, fixed and processed for wholemount skeletal preparation according to the protocol of Gillis et al. (2009a) or paraffin histology as described above.

RNA-Seq analysis of differentially expressed genes in the GAER

GAER or GAER-adjacent regions (that we termed control) of gill arch 1 were manually dissected from skate embryos at S26 using tungsten needles and flash frozen in lysis buffer using liquid nitrogen. In all cases, GAER and control sample pairs were collected from the same arch within the same embryos, or from opposite arches within the same embryo. RNA was extracted from each sample using the RNAaqueous®-Micro Kit. cDNA was synthesised from extracted RNA according to the Smart-seq2 protocol (Picelli et al., 2014) and libraries were prepared using the Nextera XT kit (Illumina). Prior to sequencing, barcoded libraries were pooled in equal proportions and checked for quality, insert size and quantity using Qubit 2.0, and Agilent2100. Sequencing of the 20 libraries generated was conducted by Novogene on an Illumina Hi-Seq-XTen generating 150bp paired-end reads. The Cambridge Service for Data-Driven Discovery (CSD3) high performance computer was used for cleaning, normalisation, quality assessment, assembly, abundance quantification and annotation, as per (Hirschberger et al., 2021). Raw reads were cleaned to remove adapter contamination using Trim Galore v0.4.0 (parameters: --paired -q 20 –nextera), for quality (Phred score 20) and minimum read length (149bp) using ea-utils tool fastq-mcf (parameters: -q 20 -l 100) (Aronesty, 2011). Prior to assembly, all libraries were normalised using Trinity v2.6.6 *in silico* read normalisation (Haas et al., 2013) (parameters: --JM 100G --max_cov 75). Trinity v2.6.6 was used to assemble the normalised reads (parameters: --seqType fq --max_memory 250G --CPU 50 --min_contig_length 450) (Nishimura et al., 2017, Langmead et al., 2009, Marcais and Kingsford, 2011). Basic Local Alignment Search Tool (BLAST) was used to compare the assembly to the Uniprot protein database with an e-value cut-off of e-20 to search

for sequence similarity (Altschul et al., 1990). Transcript abundance was estimated
using Salmon, an alignment-free quantification method, via Trinity v2.6.6 utilities
(Haas et al., 2013, Patro et al., 2017). A matrix was built for each of the libraries of
gene-level abundance estimates (raw counts) using the Trinity
abundance_estimates_to_matrix.pl script. MDS principal component analysis was
used to check for batch effects and outliers. Individual embryo origin was a batch effect
and so was accordingly accounted for in the additive general linear model.
Differentially expressed genes between control and GAER tissue samples were
identified using the edgeR_3.28.0 package in R version 3.6.2 (Robinson et al., 2010)
using a generalised linear model likelihood ratio test (Chen, 2008) using a false
discovery rate of 5% and no log fold-change (LogFC) cut-off. See supplementary
information for assembly statistics, differential expression analysis and all R scripts
used. Differentially upregulated genes with a log-fold change over 2 were manually
screened for genes involved in known signalling pathways and transcription factors to
generate a list for candidate gene validation by ISH. Wnt family members were highly
upregulated in GAER tissue so were validated as candidate markers for the GAER
through third-generation ISH. Transcripts for probes in skate were based on the most
highly expressed isoforms.
230

Imaging and figure preparation

Images were taken on a Zeiss Axioscope.A1 compound microscope with a Zeiss
colibri 7 fluorescence LED light source using a Zeiss Axiocam 305 colour or 503 mono
camera and ZenPro software or a Leica M165FC stereomicroscope, a Leica DFC7000
T camera and LAS X software. All figures were assembled using Fiji and Adobe
creative cloud, with some images flipped or colours inverted for clarity and consistency
where needed.

Results

Embryonic origin and gene expression of the skate GAER

The epithelium of vertebrate pharyngeal arches derives from both ectoderm and endoderm, and the specific tissue origin of the GAER of cartilaginous fishes is not known. In the skate, *Shh* is initially expressed broadly in the anterior endodermal epithelium of each pharyngeal pouch, which subsequently gives rise to the posterior endodermal epithelium of each pharyngeal arch (Gillis and Tidswell, 2017). We speculated that the skate GAER may derive from this initially broad endodermal *Shh* expression domain, and to test this, we labelled the pharyngeal endoderm of early skate embryos by microinjection of the lipophilic dye CM-DiI (Fig.2A) into the pharyngeal cavity at stage (S)18. At this stage, the pharyngeal pouches have not yet fused with the overlying surface ectoderm, allowing for specific CM-DiI labelling of pharyngeal endodermal epithelium (Fig. 2B). Injected embryos were grown to S25– S29, by which time a GAER is established on the hyoid and gill arches and is detectable by expression of *Shh*. Of 29 embryos analysed, all retained CM-DiI label within the endodermal lining of the pharyngeal arches, and 8 showed labelling up to and including the *Shh*-expressing GAER with CM-DiI in one or more pharyngeal arches, indicating endodermal origin of these cells (Fig. 2C). Importantly, no CM-DiI labelling was ever observed in epithelial cells immediately anterior to the GAER, suggesting that this epithelium is of ectodermal origin, and that the GAER derives from endoderm at the ectoderm–endoderm interface during pharyngeal arch development.

Shh and *Fgf8* are both reported markers of the GAER (Gillis et al., 2009b), but how these expression patterns arise relative to one another through development has not been clearly shown, nor has their co-expression in the GAER been formally demonstrated. We performed chromogenic mRNA ISH on sectioned skate embryos to visualise expression of *Shh* and *Fgf8* across different stages of pharyngeal arch development. At S22 (Fig.3A,B), *Shh* is expressed in the posterior endodermal epithelium of the hyoid and gill arches (Fig.3C) consistent with previous reports (Gillis

Figure 2: Endodermal origin of the skate GAER. (**A**) Microinjection of CM-DiI into the pharyngeal cavity of skate embryos at S18 results in (**B**) specific labelling of the pharyngeal endoderm. (**C**) *Shh* is a marker of the GAER (dashed yellow line) in CM-DiI-labelled embryos, and (**C', C''**) Co-localisation of CM-DiI and *Shh* expression indicates that the cells of the GAER are of endodermal origin. Dashed white outline demarcates GAER-adjacent (and presumed ectodermal) epithelium that is never labelled with CM-DiI. (**C**), (**C'**), and (**C''**) are the same section, imaged sequentially for *Shh* expression and CM-DiI retention. A–P: Anterior–posterior axis, dpi: days post injection. Scale bars; A: 500μm, B–C: 20μm.

Figure 3: Expression of *Shh* and *Fgf8* in developing skate pharyngeal arches.
(**A**) At S22, (**B**) the pharyngeal arches are delineated by endodermal pouches that are contacting or fusing with overlying surface ectoderm. At this stage, (**C**) *Shh* is expressed in the posterior endodermal epithelium of the developing hyoid and gill arches, while (**D**) *Fgf8* is expressed in the anterior endodermal and lateral ectodermal epithelium of each arch. (**E**) By S26, (**F**) the hyoid and gill arches are expanding laterally, and (**G**) *Shh* and (**H**) *Fgf8* are both expressed in a ridge of epithelial cells along the leading edge of the expanding arches. (**I-L**) These gene expression features persist until S29. E: eye, Epi: epithelial, Mes: mesenchymal, S: stage. Scale bars; A,E,I: 1mm, all others: 40 μm.

and Hall, 2016, Gillis and Tidswell, 2017) whereas *Fgf8* is expressed in the anterior
endoderm and lateral ectoderm of these arches (and also weakly in the distal arch
endodermal epithelium, partially overlapping with *Shh*) (Fig. 3D). By S26 (Fig.3E,F),
Shh (Fig.3G) and *Fgf8* (Fig.3H) transcripts both resolve to a thin ridge of epithelial
cells (the putative GAER) along the leading edge of the expanding hyoid and gill
arches, and these expression patterns persist until S29 (Fig.3I–L). Using multiplexed
ISH by HCR at S26, we found that *Shh* and *Fgf8* are precisely co-expressed in the
GAER of the expanding hyoid and gill arches (Fig.4A–A''). To determine which tissues
are transducing signals from the GAER, we performed ISH by HCR for *Ptc2* and
Dusp6 – transcriptional readouts of Shh and MAP kinase-mediated FGF signalling,
respectively (Pearse et al., 2001, Kawakami et al., 2003). Consistent with previous
reports (Gillis and Hall, 2016), we observed *Ptc2* expression predominantly within the
posterior arch environment, including within the GAER and adjacent epithelia as well
as within posterior-distal arch mesenchyme and core mesoderm (Fig.4B–B''). In
contrast, *Dusp6* is expressed more specifically within the GAER and broadly
throughout underlying distal arch mesenchyme (Fig.4C–C''). Altogether, these
findings demonstrate that the GAER is a signalling centre of endodermal origin, and a
source of both Shh and Fgf8 signals, with the former transduced preferentially within
in the posterior arch environment, and the latter more broadly throughout distal arch
mesenchyme.
333

Conservation of embryonic origin and gene expression in a chick pharyngeal
arch signalling centre
In amniotes, the 2[nd] (hyoid) pharyngeal arch undergoes a prolonged phase of lateral
expansion, with this arch eventually fusing with the posterior ectoderm to close the
neck. Chick embryos possess a ridge of *SHH*-expressing epithelial cells along the
leading edge of the expanding hyoid arch, and this ridge has been termed the posterior
ectodermal margin (PEM) (Wall and Hogan, 1995). The PEM is reminiscent of the
GAER of cartilaginous fishes, and to test whether these structures share common
tissue origins and gene expression features, we conducted a parallel set of lineage
tracing and gene expression analyses of the PEM in chick embryos. To test the germ
layer origin of the PEM, we first traced the fate of the pharyngeal ectoderm by *in ovo*
topical application of the lipid-soluble dye CCFSE (Richardson et al., 2012,

346
347
Figure 4: Expression of *Shh* and *Fgf8* signalling components in skate gill

arches. ISH by HCR shows co-expression of (**A**) *Shh* and *Fgf8* in the skate GAER at

S26**.** (**B**) *Ptc2* serves as a transcriptional readout of Shh signal transduction, and is

expressed in the posterior arch environment, including in the GAER and adjacent

epithelia, and in posterior–distal mesenchyme and core mesoderm. (**C**) *Dusp6*

expression indicates MAP kinase-mediated Fgf signal transduction within the GAER

and broadly throughout distal arch mesenchyme. Scale bar: 50μm.

Shone and Graham, 2014). Of 29 embryos labelled at HH11, all showed some

labelling of the ectoderm-derived lining of the pharyngeal arches by HH19, with 9

embryos showing strong labelling of the ectoderm-derived epithelium of the hyoid arch

up to, but excluding, the *SHH*-expressing cells of the PEM (Fig. 2 Supplement 1A–C).

We next traced the fate of pharyngeal endoderm by microinjection of CM-DiI into the

pharyngeal cavity. Of 37 embryos labelled at HH11, all retained some CM-DiI labelling

within the endoderm-derived epithelium of the pharyngeal arches, with 9 embryos

showing CM-DiI labelling of the endodermal lining of the hyoid arch up to and including

the *SHH*-expressing PEM (Fig. 2 Supplement 1D–E). These findings point to an

endodermal origin of the PEM, with this signalling centre forming at the endoderm-

ectoderm boundary of the hyoid arch epithelium. *SHH, FGF8* and *BMP7* are all

reported markers of the PEM in chick (Wall and Hogan, 1995), and we confirmed that

while *SHH* and *BMP7* are initially expressed broadly in the pharyngeal endoderm and

FGF8 in the pharyngeal ectoderm and endoderm of the hyoid arch (Fig. 3 Supplement

1A–J), expression of all three genes subsequently resolves to a discrete epithelial

domain along the leading edge of the expanding arch (Fig. 3 Supplement 1K–T).

Finally, we found using ISH by HCR that *SHH, FGF8* and *BMP7* are co-expressed

within cells of the PEM (Fig. 3 Supplement 1U). These findings demonstrate that the

skate GAER and chick PEM share embryonic tissue origins and expression patterns

of signalling molecules through development and are therefore homologous and an

ancestral feature of the pharyngeal arche(s) of jawed vertebrates.

***Wnt* gene expression in the skate GAER and GAER-adjacent ectoderm**

In an attempt to discover additional gene expression features of the GAER, we

conducted a transcriptomic and differential gene expression analysis between the

GAER and non-GAER regions of the first gill arch of skate embryos. Briefly, we

manually dissected the posterior–distal GAER region and non-GAER (control) region

from the first gill arch of S26 skate embryos (n=5) (Fig.5), and we performed RNA

extraction, library preparation, and RNAseq analysis on these samples. Following *de*

novo transcriptome assembly, we compared gene expression levels between GAER

and control tissue and using a false discovery rate of 5% and no log fold-change

(LogFC) cut-off we generated a list of 401 genes that were upregulated in the GAER.

We sorted this list for genes encoding signalling pathway components and

Figure 5: RNA-seq and differential gene expression analysis of GAER and non-GAER gill arch tissues in S26 skate embryos. (**A**) DAPI stained skate embryo at S26 and (**B**) schematic of dissection from gill arch 1 of (**Bi**) non-GAER (control) tissue region (green) and posterior–distal GAER tissue region (blue), as guided by (**C**) *Shh* expression. (**D, Di**) Volcano plot illustrating differential gene expression between GAER and non-GAER tissues. Genes with >2 logFC and <0.05 false discovery rate (FDR) are highlighted and assigned functional category using colour coding as per key. Genes with >2 logFC are shown in larger point size. Signalling molecules and transcription factors with >2 logFC and <0.05 FDR are labelled (* denotes manual annotation of sequence). Differential expression was determined using edgeR with a general linear model and likelihood ratio test, corrected for multiple testing using the Benjamin-Hochberg method to control the FDR. Scale bars; A:1mm C: 50μm.

transcription factors with a log-fold change >2 (Fig.5E) and found several components
of the Wnt signalling pathway among genes upregulated in GAER tissue. These
included genes encoding the Wnt ligands Wnt2b, Wnt3, Wnt4, Wnt7b and Wnt9, the
transmembrane Wnt inhibitors Kremen1 (Mao et al., 2002) and Adenomatosis
Polyposis Coli Down-regulated 1 (APCDD1) (Shimomura et al., 2010), and the
secreted Wnt antagonist Notum (Zhang et al., 2015). We chose to further investigate
potential involvement of the Wnt pathway in GAER signalling by spatially validating
the expression of these genes, using the most highly expressed isoforms for HCR
probe set design.

To characterise expression of genes encoding Wnt ligands relative to *Shh* expression
in the GAER, we used multiplexed ISH by HCR to visualise transcript localisation in
sections of S26 skate gill arches. We found that while *Wnt7b* was co-expressed
with *Shh* in the cells of the GAER (Fig. 6A, A'), *Wnt2b*, *Wnt3*, *Wnt4* and *Wnt9b* were
all expressed predominantly in the ectoderm immediately adjacent to the GAER (Fig.
6B–E). We next examined the expression of the Wnt signalling downstream target
genes and found that *Apcdd1* was expressed in the GAER and anterior–distal
mesenchyme (Fig. 6F), *Notum* in the anterior–distal mesenchyme (Fig. 6G)
and *Kremen1* in the anterior (GAER-adjacent) ectoderm (Fig. 6H). Finally, broad
transcription of *Axin2* throughout distal gill arch tissues (Fig. 6I) indicates that Wnt
signalling within this territory is occurring through the canonical/β-catenin pathway
(Lustig et al., 2002). While these findings do not allow us to attribute expression of
specific readouts to signalling by particular Wnt ligands, our spatial expression data
nevertheless indicate that some Wnt signals emanating from the GAER or GAER-
adjacent ectoderm are transduced preferentially within the anterior arch environment,
in a pattern complimentary to the posterior epithelial and mesenchymal transduction
of Shh signals from the GAER.

Conservation of *WNT* gene expression in PEM-adjacent ectoderm in chick
To test whether the Wnt gene expression patterns described above for skate are a
unique feature of cartilaginous fishes or a general feature of jawed vertebrates, we
again used ISH by HCR to visualise expression in or around the hyoid arch PEM in

Figure 6: *Wnt* pathway genes are expressed in and around the GAER in skate.
In S26 skates, (**A**) ISH by HCR for *Shh* and DAPI staining was performed alongside
each gene of interest as a marker of the GAER and for general tissue overview,
respectively. For ease of visualisation in **A'–I**, *Shh* expression is not shown but GAER

cells are outlined green, and arch tissues are outlined in white. (**A'**) *Wnt7b* is co-expressed with *Shh* in the GAER. (**B)** *Wnt2b,* **(C)** *Wnt3*, **(D)** *Wnt4* and **(E)** *Wnt9b* are predominantly expressed in the ectodermal epithelium immediately adjacent the GAER. (**F**) *Apcdd1* is expressed in the GAER, and also in the anterior–distal mesenchyme underlying the GAER. (**G**) *Notum* is expressed in the anterior–distal arch mesenchyme. (**H**) *Kremen1* is expressed in the GAER and in GAER-adjacent ectodermal epithelium. (**I**) Broad *Axin2* expression in the vicinity of *Wnt*-expressing epithelium indicates signalling through the canonical Wnt signalling pathway. Epi: epithelial, mes: mesenchymal, mc: mesodermal core. Scale bar: 50μm.

chick embryos (Fig. 6 Supplement 1A). At HH20, we found that *WNT2B, WNT3, WNT4* and *WNT9B* are all predominantly expressed in the ectodermal epithelium directly adjacent to the PEM (Fig. 6 Supplement 1B–E). Unlike in skate, *WNT2B* expression in chick extends into the PEM, and we were unable to detect expression of *WNT7B* in the PEM. We also found that *APCDD1* is expressed in the distal arch mesenchyme (Fig. 6 Supplement 1F), *NOTUM* in the anterior–distal mesenchyme and the PEM (Fig. 6 Supplement 1G), and *KREMEN1* predominantly in the ectodermal (PEM-adjacent) epithelium (Fig. 6 Supplement 1H). Broad transcription of *AXIN2* was also seen in the distal arch tissues, indicating active canonical/β-catenin pathway Wnt signalling (Fig. 6 Supplement 1I). These findings demonstrate that Wnt signalling emanating predominantly from PEM- or GAER-adjacent ectoderm is conserved in chick and skate, respectively, and is therefore likely an ancestral feature of the developing pharyngeal arches of jawed vertebrates.

Wnt suppresses GAER Shh signal transduction and inhibits cartilage formation in the anterior gill arch in skate

To explore the function of Wnt signalling during skate pharyngeal arch development, we inhibited canonical Wnt signalling in skate embryos using IWR1, a small molecule tankyrase inhibitor that antagonises Wnt signalling by stabilising the Axin/β-catenin destruction complex (Lu et al., 2009, Kulak et al., 2015). Briefly, skate embryos were maintained in a bath of 50μM IWR1 or vehicle-only control (DMSO) in seawater from S25 for 72h to assess gene expression changes in response to drug treatment, or for 6–8 weeks (until S31/32) to assess effects of Wnt signalling inhibition on gill arch skeletal patterning, with replacement of drug- or control-seawater every 48h. IWR1-treated embryos showed a marked reduction in the expression of *Axin2* throughout their gill arches (Fig. 6 Supplement 2A–B), as well as a reduction in the expression of *Apcdd1* in the GAER and anterior–distal gill arch mesenchyme (Fig. 6 Supplement 2C–D) and a complete loss of *Notum* expression in anterior–distal gill arch mesenchyme (Fig. 6 Supplement 2E–F). These findings indicate that our drug treatment effectively supressed canonical Wnt signalling from the GAER and GAER-adjacent epithelium, including those signals transduced asymmetrically in the anterior gill arch environment.

Shh is a pro-chondrogenic signal from the GAER in skate gill arches (Gillis et al.,
2009b), and this signal is transduced asymmetrically with the posterior gill arch
environment (Gillis and Hall, 2016, and see above). To test whether anteriorly-
transduced Wnt signalling may act to restrict Shh signal transduction to the posterior
gill arch, we examined expression of *Shh* and *Ptc2* in skate embryos 72h after the
onset of IWR1 treatment. While we observed no noticeable change in the expression
of *Shh* or in the spatial distribution of *Ptc2* transcripts in gill arch mesenchyme with
IWR1 treatment, we did note a qualitative increase in *Ptc2* expression within the
normal mesenchymal domain, and a striking anterior expansion of *Ptc2* expression
within the epithelium adjacent to the GAER (Fig.7A–D). To quantify this epithelial
expansion, we compared the mean number of *Ptc2*+ cells anterior to the *Shh*-
expressing GAER across two or three sections at equivalent positions in DMSO
control (n=3) or IWR1 (n=4) treated embryos and found a significant anterior
expansion of *Ptc2* expression within the epithelium adjacent to the GAER in IWR1
treated embryos (Fig.7E–G; P= 0.036). This points to a role for GAER-adjacent Wnt
signalling in restricting the transduction of pro-chondrogenic Shh signalling in the
anterior gill arch territory in skate.

Finally, to test for an effect of Wnt inhibition on the gill arch skeleton, we examined
skeletal morphology in skate embryos reared in DMSO control seawater or IWR1 from
S25–S31/32. The skate hyoid and gill arch skeleton include branchial rays that
articulate exclusively with the posterior margin of the gill arch cartilages, and that
invariably develop on the posterior side of the mesodermally-derived interbranchial
muscle plate that extends down the middle of the arch (Daniel, 1934, Gillis et al.,
2009a). Skate embryos reared in DMSO control seawater showed no gill arch skeletal
defects (n=4 embryos; Fig. 8A–Ai), while embryos reared in IWR1 seawater
possessed conspicuous ectopic branchial rays in the anterior gill arch territory (n= 7/8
embryos; Fig. 8B–Bi). These ectopic branchial rays occurred along the entire
dorsoventral axis of the arch and ranged from short cartilage rods located in the
anterior–distal gill arch to nearly full-length branchial rays that extended much of the
way toward the gill arch cartilages. In section, normal branchial rays of skate embryos
reared in DMSO control seawater always developed on the posterior side of the
mesodermally-derived interbranchial muscle plate of each gill arch: this was evident

Figure 7: Wnt signalling restricts transduction of GAER Shh signalling to the posterior gill arch environment. ISH by HCR on sections of skate embryos treated for 72 hours with (**A,B**) DMSO or (**C,D**) IWR1 shows no difference in *Shh* expression but an expansion of *Ptc2* expression within the epithelium anterior to the *Shh*-expressing GAER. (**E**) Schematic illustration of the mean number of cells containing *Ptc2* transcripts anterior to the GAER in DMSO control (n=3; 6.3 cells) or (**F**) IWR1 treated (n=4; 12.94 cells) embryos, using the mean cell count of two or three sections at equivalent positions in the gill arch from each embryo. (**G**) A significant increase in *Ptc2*+ cells in the epithelium anterior to the GAER in IWR1 treated embryos compared to control (P= 0.036; t test). In **A–F**, *Shh* expression is indicated by green dashed line. Images in **A–D** taken using identical exposure settings. Scale bar: 20 μm.

Figure 8: Ectopic branchial ray formation upon inhibition of Wnt signalling. (A)
In control (DMSO) skate embryos, branchial rays (magnified in **Ai**) articulate
exclusively with the posterior margin of the gill arches. **(B)** Embryos reared in the
Wnt inhibitor IWR1 possessed ectopic branchial rays (magnified in **Bi**) that were
embedded within connective tissue of the anterior gill arch (ectopic rays present in
n=7/8 embryos examined; indicated by red asterisk). Scale bars: 500 μm.

566
567
Figure 9: Repression of cartilage development in the anterior gill arch by Wnt signalling. (A) Section of the branchial rays of control (DMSO) skate embryos, showing they develop exclusively on the posterior side of the interbranchial muscle plate, as is evident from histochemical staining and **(B–ii)** ISH by HCR for the chondrocyte markers *Sox9* and *Cola2a1*. **(C)** Section of the branchial rays of embryos reared in the Wnt inhibitor IWR1 show ectopic branchial rays form anterior to the interbranchial muscle plate, as evident from histological analysis and **(D–Dii)** ISH by HCR for *Sox9* and *Col2a1*. Red asterisks indicate ectopic branchial rays, and interbranchial muscle plate is outlined with a dashed line. Scale bars: 50 μm.

from histochemical staining (Fig. 9A) and from HCR for the chondrocyte markers *Sox9*

and *Col2a1* (Fig. 9B). Conversely, in skate embryos reared in IWR1 seawater, ectopic

branchial rays were consistently located anterior to the interbranchial muscle plate

(Fig. 9C–D). The induction of ectopic branchial rays with IWR1 treatment suggests

that Wnt signalling contributes to the maintenance of anteroposterior polarity of the

skate gill arch skeleton by repressing chondrogenesis in the anterior gill arch and may

do so by restricting transduction of GAER Shh signalling to the posterior gill arch

territory.

Discussion

The molecular mechanisms governing spatial regulation of cell fate decisions are

crucial to the establishment and maintenance of anatomical polarity within

developing tissues and organs. Here, we show that the GAER signalling centre that

forms on the hyoid and gill arches of cartilaginous fishes is of endodermal origin,

arising precisely at the boundary of surface ectoderm and pharyngeal endoderm,

and that pro-chondrogenic Shh signals from the GAER are transduced preferentially

within the posterior gill arch environment. We further show that the ectoderm

immediately anterior and adjacent to the GAER is a signalling centre that expresses

several Wnt family genes, and that Wnt signals from this centre are transduced

preferentially in the anterior arch environment. Pharmacological inhibition of

canonical Wnt signalling in skate embryos leads to an expansion of Shh signal

transduction within the anterior territory of the gill arch, and to the formation of

ectopic anterior branchial ray cartilage, highlighting Wnt–Shh signalling antagonism

across an ectodermal–endodermal tissue boundary as a key molecular regulator of

cell fate determination and anatomical polarity with the developing gill arch skeleton

(Fig. 10).

Previous analyses have reported the expression of several Wnt genes in developing

craniofacial tissues of bony vertebrates (Summerhurst et al., 2008), including

expression of *WNT2B* and *WNT9B* in the pharyngeal arch ectoderm of the chick

(Geetha-Loganathan et al., 2009), *Wnt9b* in the first and second pharyngeal arches in

zebrafish (Jezewski et al., 2008), and *Wnt4a* in zebrafish pharyngeal ectoderm (Choe

Figure 10: Wnt and Shh signalling and cell fate determination in skate gill arches. Shh signalling from the endodermally-derived GAER is transduced within the posterior-distal gill arch environment, where it promotes the differentiation of cartilaginous branchial rays. Wnt signals from ectoderm adjacent to the GAER are transduced within the anterior–distal gill arch environment, where they repress Shh signal transduction and inhibit cartilage formation in the anterior gill arch.

et al., 2013). While β-catenin-independent (i.e. non-canonical) Wnt signalling plays a
crucial role in epithelial remodelling during the early formation of pharyngeal
endodermal pouches in zebrafish (Choe et al., 2013), canonical Wnt signalling has
been implicated in patterning of the pharyngeal arch-derived skeleton in multiple taxa.
Blocking canonical Wnt signalling causes disrupted facial cartilage formation,
including reductions of Meckel's and ceratohyal cartilages in zebrafish (Alexander et
al., 2014) and loss of ectodermal canonical Wnt signalling in mice causes loss of facial
bones with hypoplasia of all facial prominences (Reid et al., 2011). Furthermore, in
mouse and human, cleft palate phenotypes may arise as a consequence of mutations
in *Wnt3* and *Wnt9b* (Niemann et al., 2004, Juriloff et al., 2006), and perturbation of
Wnt/β-catenin signalling has been implicated in CATSHL syndrome, a human
developmental disorder involving craniofacial bone malformation and mispatterning of
the pharyngeal arches (Sun et al., 2020). Our finding of a role for canonical Wnt
signalling in patterning the gill arch skeleton of skate may therefore reflect broadly
conserved roles for this pathway in pharyngeal skeletal development across jawed
vertebrates, though with the skate's possession of branchial rays offering a unique
anatomical readout of anterior–posterior polarity defects that arise within the
pharyngeal arch-derived skeleton in response to perturbations.

Our findings are also consistent with previously reported complex and context-
dependent roles for Wnt signalling in the regulation of cell fate determination events
during vertebrate skeletogenesis. In some instances, Wnt signalling functions to
promote skeletogenesis: for example, in chick, misexpression of *WNT5A/5B* promotes
early chondrogenesis of limb bud mesenchyme *in vitro* and delays the terminal
differentiation of growth plate chondrocytes *in vivo* (Church et al., 2002), and in mouse,
canonical WNT signalling from cranial ectoderm induces specification of osteoblast
progenitors of cranial dermal bone within underlying mesenchyme (Goodnough et al.,
2016). Conversely, in other contexts, Wnt signalling functions to inhibit chondrogenic
differentiation or homeostasis. In chick, signalling through WNT3, 4, 7A, 14 or FZ7
have all been shown to inhibit chondrogenesis *in vitro* or *in vivo* (Rudnicki and Brown,
1997), while in mouse conditionally-induced haploinsufficiency of the gene encoding
the β-catenin degradation complex component APC (i.e. activation of canonical Wnt
signalling) results in loss of resting zone chondrocytes and their clonal progeny in
developing growth plates (Hallett et al., 2021). In the limb bud, genes encoding WNT

ligands are expressed in the ectoderm (Kengaku et al., 1998, Parr et al., 1993,
Geetha-Loganathan et al., 2005), where they function synergistically with FGF
signalling from the apical ectoderm ridge to inhibit chondrogenesis of limb bud
mesenchyme and to promote soft connective tissue fates (ten Berge et al., 2008). Our
findings therefore demonstrate a striking parallel between developing skate gill arches
and tetrapod limb buds, with Wnt signals emanating from ridge-adjacent epithelia
functioning to inhibit chondrogenic differentiation, thereby ensuring differentiation of
cartilaginous appendages in a spatially-controlled manner.

We have previously shown that Shh signalling from the GAER is required for branchial
ray chondrogenesis in skate (Gillis and Hall, 2016), and that the application of
exogenous SHH protein to the anterior gill arch is sufficient to induce ectopic branchial
rays (Gillis et al., 2009b). However, it is unclear whether Shh from the GAER is a direct
inducer of chondrogenesis in the posterior arch mesenchyme, or rather induces
chondrogenesis indirectly, via a secondary signal (i.e. emanating from the Shh-
responsive epithelium or mesenchyme). We have observed that inhibition of Wnt
signalling in skate embryos has no effect on the expression of *Shh* in the GAER, but
rather results in an anterior expansion of Shh signal transduction in the gill arch
ectoderm, which in turn, correlates with ectopic chondrogenesis in the anterior gill
arch. This observation indicates that the pro-chondrogenic influence of GAER Shh
signalling may be indirect, occurring via a secondary Shh-depending epithelial signal
whose expression is typically posteriorly restricted by Wnt signalling from GAER-
adjacent ectoderm. The Shh and Wnt signalling pathways share multiple downstream
components (reviewed in; Ding and Wang (2017). For example, components of the
canonical Wnt signalling pathway can positively regulate Shh signalling via GSK3β,
by phosphorylating the downstream component SUFU and promoting the release of
SUFU from Gli (Takenaka et al., 2007), and by β-catenin affecting Gli1 transcriptional
activity via TCF/LEF (Maeda et al., 2006). It has also been found that Gli3 may function
as a downstream effector of the Wnt pathway, and that Wnt signalling represses Shh
activity in the dorsal neural tube through the regulation of *Gli3* expression (Alvarez-
Medina et al., 2008). Finally, the Shh signalling pathway may regulate Wnt signalling
through Gli1 and Gli2, with these factors positively regulating the expression of the
secreted Wnt inhibitor *frizzled-related protein-1* (*sFRP-1*) (He et al., 2006). These
mechanisms, or others, could account for the apparent cross regulation between Wnt

and Shh signalling that we have observed during growth and patterning of skate
pharyngeal arches.

Finally, conservation of *Shh*, *Fgf* and *Wnt* expression at the pharyngeal arch
endoderm–ectoderm boundary in skate and chick point to these as ancestral features
of the jawed vertebrate pharyngeal patterning programme. In bony vertebrates, the
PEM is a feature of the hyoid arch (Wall and Hogan, 1995). The hyoid arch of bony
fishes expands to form an operculum that is supported by dermal bone, and functions
as a protective cover for the gills that arise from the posterior pharyngeal arches
(Richardson et al., 2012), while the hyoid arch of amniotes expands caudally and
ultimately fuses with the posterior ectoderm to enclose the cervical sinus and close
the surface of the neck (Richardson et al., 2012). In elasmobranch cartilaginous fishes,
on the other hand, the GAER is a feature of the hyoid and first four gill arches, all of
which undergo lateral expansion to form a series of protective flaps that are supported
by cartilaginous branchial rays (Gillis et al., 2011). In a departure from the
elasmobranch condition, holocephalan cartilaginous fishes possess a GAER only on
their hyoid arch, and this correlates with the presence in this group of a single hyoid
arch-derived opercular gill cover supported by cartilaginous branchial rays, which
evolved convergently with the condition seen in bony fishes (Gillis et al., 2011). Recent
fossil data resolves the hyoid operculate condition of bony fishes as primitive for jawed
vertebrates (Dearden et al., 2019), with iterative lateral expansion of the gill arches
arising along the cartilaginous fish stem group, and with secondary loss of gill arch
expansion in holocephalans (Gillis et al., 2011). Therefore, while the number of
pharyngeal arches possessing a GAER/PEM varies across jawed vertebrate lineages,
the presence of these signalling centres correlates tightly with arches that undergo
lateral expansion, highlighting variation in GAER/PEM retention as a major
determinant of jawed vertebrate pharyngeal architecture. Although amniotes lack
homologues of the opercular series or branchial rays of bony and cartilaginous fishes,
respectively, further comparative work between bony and cartilaginous fishes with
disparate pharyngeal skeletal configurations may shed light on ancestral functions for
ectodermal Wnt and endodermal GAER/PEM signalling in governing pharyngeal
skeletal pattern and cell fate determination.

735
736

Acknowledgements

With thanks to Dr Kate Criswell and Dr Christine Hirschberger for their advice, and to

the Wellcome PhD Programme in Developmental Mechanisms. The authors were

funded by a Wellcome PhD studentship (214953/Z/18/Z) to JMR, and by a Royal

Society University Research Fellowship (UF130182 and URF\R\191007) and Royal

Society Research Grant (RG140377) to JAG.

Data Accessibility

All data and R scripts for analysis have been deposited on Figshare:

10.6084/m9.figshare.19615779

RNA sequencing data are available at NCBI-SRA under BioProject ID: PRJNA825354

Biosample accessions: SAMN27512544, SAMN27512545, SAMN27512546,

SAMN27512547, SAMN27512548, SAMN27512549, SAMN27512550,

SAMN27512551, SAMN27512552, SAMN27512553, SAMN27512554,

SAMN27512555, SAMN27512556, SAMN27512557, SAMN27512558,

SAMN27512559, SAMN27512560, SAMN27512561, SAMN27512562,

SAMN27512563

Probes were purchased from Molecular Instruments (Los Angeles, California, USA).

This included the following: for skate *Shh* (Lot PRA753), *Ptc2* (Lot PRA754), *Fgf8* (Lot

PRA755), *Dusp6* (Lot PRA756), *Wnt2b* (Lot PRE300), *Wnt3* (Lot PRG814), *Wnt4* (Lot

PRE301), *Wnt7b* (Lot PRE302), *Wnt9b* (Lot PRE303), Notum (Lot PRG817), *Kremen1*

(Lot PRG816), *Axin2* (Lot PRG818), *Apcdd1* (Lot PRG815), *Col2a1* (Lot PRB574) and

Sox9 (Lot PRB571); for chick *SHH* (Lot PRB282), *FGF8* (Lot PRA997), *BMP7* (Lot

PRb283), *WNT2B* (Lot PRG820), *WNT3* (Lot PRG823), *WNT4* (Lot PRG821), *WNT7B*

(Lot PRG819), *WNT9B* (Lot PRG822), *NOTUM* (Lot PRG824), *KREMEN1* (Lot

PRG825), *AXIN2* (Lot PRG827), *APCDD1* (Lot PRG826).

References

ALEXANDER, C., PILOTO, S., LE PABIC, P. & SCHILLING, T. F. 2014. Wnt signaling interacts with bmp and edn1 to regulate dorsal-ventral patterning and growth of the craniofacial skeleton. *PLoS Genet,* 10**,** e1004479.

ALTSCHUL, S. F., GISH, W., MILLER, W., MYERS, E. W. & LIPMAN, D. J. 1990. Basic local alignment search tool. *J Mol Biol,* 215**,** 403-10.

ALVAREZ-MEDINA, R., CAYUSO, J., OKUBO, T., TAKADA, S. & MARTI, E. 2008. Wnt canonical pathway restricts graded Shh/Gli patterning activity through the regulation of Gli3 expression. *Development,* 135**,** 237-47.

ARONESTY, E. 2011. *Command-line tools for processing biological sequencing data* [Online]. Available: https://github.com/ExpressionAnalysis/ea-utils [Accessed January 2020].

BALLARD, W. W., MELLINGER, J. & LECHENAULT, H. 1993. A series of normal stages for development of Scyliorhinus canicula, the lesser spotted dogfish (Chondrichthyes: Scyliorhinidae). *Journal of Experimental Zoology,* 267**,** 318-336.

BRITO, J. M., TEILLET, M. A. & LE DOUARIN, N. M. 2006. An early role for sonic hedgehog from foregut endoderm in jaw development: ensuring neural crest cell survival. *Proc Natl Acad Sci U S A,* 103**,** 11607-12.

CHEN, Y. 2008. *edgeR: differential analysis of sequence read count data* [Online]. Available: https://www.bioconductor.org/packages/release/bioc/vignettes/edgeR/inst/doc/edgeR UsersGuide.pdf [Accessed].

CHOE, C. P., COLLAZO, A., TRINH LE, A., PAN, L., MOENS, C. B. & CRUMP, J. G. 2013. Wnt-dependent epithelial transitions drive pharyngeal pouch formation. *Dev Cell,* 24**,** 296-309.

CHOI, H. M. T., SCHWARZKOPF, M., FORNACE, M. E., ACHARYA, A., ARTAVANIS, G., STEGMAIER, J., CUNHA, A. & PIERCE, N. A. 2018. Third-generation in situ hybridization chain reaction: multiplexed, quantitative, sensitive, versatile, robust. *Development,* 145.

CHURCH, V., NOHNO, T., LINKER, C., MARCELLE, C. & FRANCIS-WEST, P. 2002. Wnt regulation of chondrocyte differentiation. *J Cell Sci,* 115**,** 4809-18.

COULY, G., CREUZET, S., BENNACEUR, S., VINCENT, C. & LE DOUARIN, N. M. 2002. Interactions between Hox-negative cephalic neural crest cells and the foregut endoderm in patterning the facial skeleton in the vertebrate head. *Development,* 129**,** 1061-73.

COULY, G. & LE DOUARIN, N. M. 1990. Head morphogenesis in embryonic avian chimeras: evidence for a segmental pattern in the ectoderm corresponding to the neuromeres. *Development,* 108**,** 543-58.

CRISWELL, K. E. & GILLIS, J. A. 2020. Resegmentation is an ancestral feature of the gnathostome vertebral skeleton. *Elife,* 9.

DANIEL, J. F. 1934. *The elasmobranch fishes,* Berkeley,Calif., University of California press.

DEARDEN, R. P., STOCKEY, C. & BRAZEAU, M. D. 2019. The pharynx of the stem-chondrichthyan Ptomacanthus and the early evolution of the gnathostome gill skeleton. *Nat Commun,* 10**,** 2050.

DING, M. & WANG, X. 2017. Antagonism between Hedgehog and Wnt signaling pathways regulates tumorigenicity. *Oncol Lett,* 14**,** 6327-6333.

GEETHA-LOGANATHAN, P., NIMMAGADDA, S., ANTONI, L., FU, K., WHITING, C. J., FRANCIS-WEST, P. & RICHMAN, J. M. 2009. Expression of WNT signalling pathway genes during chicken craniofacial development. *Dev Dyn,* 238**,** 1150-65.

GEETHA-LOGANATHAN, P., NIMMAGADDA, S., PROLS, F., PATEL, K., SCAAL, M., HUANG, R. & CHRIST, B. 2005. Ectodermal Wnt-6 promotes Myf5-dependent avian limb myogenesis. *Dev Biol,* 288**,** 221-33.

GILLIS, J. A., DAHN, R. D. & SHUBIN, N. H. 2009a. Chondrogenesis and homology of the visceral skeleton in the little skate, Leucoraja erinacea (Chondrichthyes: Batoidea). *J Morphol,* 270**,** 628-43.

GILLIS, J. A., DAHN, R. D. & SHUBIN, N. H. 2009b. Shared developmental mechanisms pattern the vertebrate gill arch and paired fin skeletons. *Proc Natl Acad Sci U S A,* 106**,** 5720-4.

GILLIS, J. A. & HALL, B. K. 2016. A shared role for sonic hedgehog signalling in patterning chondrichthyan gill arch appendages and tetrapod limbs. *Development,* 143**,** 1313-7.

GILLIS, J. A., MODRELL, M. S., NORTHCUTT, R. G., CATANIA, K. C., LUER, C. A. & BAKER, C. V. 2012. Electrosensory ampullary organs are derived from lateral line placodes in cartilaginous fishes. *Development,* 139**,** 3142-6.

GILLIS, J. A., RAWLINSON, K. A., BELL, J., LYON, W. S., BAKER, C. V. & SHUBIN, N. H. 2011. Holocephalan embryos provide evidence for gill arch appendage reduction and opercular evolution in cartilaginous fishes. *Proc Natl Acad Sci U S A,* 108**,** 1507-12.

GILLIS, J. A. & TIDSWELL, O. R. 2017. The Origin of Vertebrate Gills. *Curr Biol,* 27**,** 729-732.

GOODNOUGH, L. H., DINUOSCIO, G. J. & ATIT, R. P. 2016. Twist1 contributes to cranial bone initiation and dermal condensation by maintaining Wnt signaling responsiveness. *Dev Dyn,* 245**,** 144-56.

GRAHAM, A. & SMITH, A. 2001. Patterning the pharyngeal arches. *Bioessays,* 23**,** 54-61.

GREVELLEC, A. & TUCKER, A. S. 2010. The pharyngeal pouches and clefts: Development, evolution, structure and derivatives. *Semin Cell Dev Biol,* 21**,** 325-32.

HAAS, B. J., PAPANICOLAOU, A., YASSOUR, M., GRABHERR, M., BLOOD, P. D., BOWDEN, J., COUGER, M. B., ECCLES, D., LI, B., LIEBER, M., MACMANES, M. D., OTT, M., ORVIS, J., POCHET, N., STROZZI, F., WEEKS, N., WESTERMAN, R., WILLIAM, T., DEWEY, C. N., HENSCHEL, R., LEDUC, R. D., FRIEDMAN, N. & REGEV, A. 2013. De novo transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nat Protoc,* 8**,** 1494-512.

HALLETT, S. A., MATSUSHITA, Y., ONO, W., SAKAGAMI, N., MIZUHASHI, K., TOKAVANICH, N., NAGATA, M., ZHOU, A., HIRAI, T., KRONENBERG, H. M. & ONO, N. 2021. Chondrocytes in the resting zone of the growth plate are maintained in a Wnt-inhibitory environment. *Elife,* 10.

HAMBURGER, V. & HAMILTON, H. L. 1951. A series of normal stages in the development of the chick embryo. *J. Morphol.,* 88**,** 49-92.

HE, J., SHENG, T., STELTER, A. A., LI, C., ZHANG, X., SINHA, M., LUXON, B. A. & XIE, J. 2006. Suppressing Wnt signaling by the hedgehog pathway through sFRP-1. *J Biol Chem,* 281**,** 35598-602.

HIRSCHBERGER, C., SLEIGHT, V. A., CRISWELL, K. E., CLARK, S. J. & GILLIS, J. A. 2021. Conserved and unique transcriptional features of pharyngeal arches in the skate (Leucoraja erinacea) and evolution of the jaw. *Mol Biol Evol,* 38**,** 4187-4204.

JEZEWSKI, P. A., FANG, P. K., PAYNE-FERREIRA, T. L. & YELICK, P. C. 2008.
Zebrafish Wnt9b synteny and expression during first and second arch, heart, and
pectoral fin bud morphogenesis. *Zebrafish,* 5**,** 169-77.
JIANG, X., ISEKI, S., MAXSON, R. E., SUCOV, H. M. & MORRISS-KAY, G. M. 2002.
Tissue origins and interactions in the mammalian skull vault. *Dev Biol,* 241**,** 106-16.
JURILOFF, D. M., HARRIS, M. J., MCMAHON, A. P., CARROLL, T. J. & LIDRAL, A. C.
2006. Wnt9b is the mutated gene involved in multifactorial nonsyndromic cleft lip
with or without cleft palate in A/WySn mice, as confirmed by a genetic
complementation test. *Birth Defects Res A Clin Mol Teratol,* 76**,** 574-9.
KAGUE, E., GALLAGHER, M., BURKE, S., PARSONS, M., FRANZ-ODENDAAL, T. &
FISHER, S. 2012. Skeletogenic fate of zebrafish cranial and trunk neural crest. *PLoS*
One, 7**,** e47394.
KAWAKAMI, Y., RODRIGUEZ-LEON, J., KOTH, C. M., BUSCHER, D., ITOH, T.,
RAYA, A., NG, J. K., ESTEBAN, C. R., TAKAHASHI, S., HENRIQUE, D.,
SCHWARZ, M. F., ASAHARA, H. & IZPISUA BELMONTE, J. C. 2003. MKP3
mediates the cellular response to FGF8 signalling in the vertebrate limb. *Nat Cell*
Biol, 5**,** 513-9.
KENGAKU, M., CAPDEVILA, J., RODRIGUEZ-ESTEBAN, C., DE LA PENA, J.,
JOHNSON, R. L., IZPISUA BELMONTE, J. C. & TABIN, C. J. 1998. Distinct WNT
pathways regulating AER formation and dorsoventral polarity in the chick limb bud.
Science, 280**,** 1274-7.
KULAK, O., CHEN, H., HOLOHAN, B., WU, X., HE, H., BOREK, D., OTWINOWSKI, Z.,
YAMAGUCHI, K., GAROFALO, L. A., MA, Z., WRIGHT, W., CHEN, C., SHAY,
J. W., ZHANG, X. & LUM, L. 2015. Disruption of Wnt/beta-Catenin Signaling and
Telomeric Shortening Are Inextricable Consequences of Tankyrase Inhibition in
Human Cells. *Mol Cell Biol,* 35**,** 2425-35.
LANGMEAD, B., TRAPNELL, C., POP, M. & SALZBERG, S. L. 2009. Ultrafast and
memory-efficient alignment of short DNA sequences to the human genome. *Genome*
Biol, 10**,** R25.
LU, J., MA, Z., HSIEH, J. C., FAN, C. W., CHEN, B., LONGGOOD, J. C., WILLIAMS, N.
S., AMATRUDA, J. F., LUM, L. & CHEN, C. 2009. Structure-activity relationship
studies of small-molecule inhibitors of Wnt response. *Bioorg Med Chem Lett,* 19**,**
3825-7.
LUSTIG, B., JERCHOW, B., SACHS, M., WEILER, S., PIETSCH, T., KARSTEN, U.,
VAN DE WETERING, M., CLEVERS, H., SCHLAG, P. M., BIRCHMEIER, W. &
BEHRENS, J. 2002. Negative feedback loop of Wnt signaling through upregulation
of conductin/axin2 in colorectal and liver tumors. *Mol Cell Biol,* 22**,** 1184-93.
MAEDA, O., KONDO, M., FUJITA, T., USAMI, N., FUKUI, T., SHIMOKATA, K.,
ANDO, T., GOTO, H. & SEKIDO, Y. 2006. Enhancement of GLI1-transcriptional
activity by beta-catenin in human cancer cells. *Oncol Rep,* 16**,** 91-6.
MAO, B., WU, W., DAVIDSON, G., MARHOLD, J., LI, M., MECHLER, B. M., DELIUS,
H., HOPPE, D., STANNEK, P., WALTER, C., GLINKA, A. & NIEHRS, C. 2002.
Kremen proteins are Dickkopf receptors that regulate Wnt/beta-catenin signalling.
Nature, 417**,** 664-7.
MARCAIS, G. & KINGSFORD, C. 2011. A fast, lock-free approach for efficient parallel
counting of occurrences of k-mers. *Bioinformatics,* 27**,** 764-70.
MARCONI, A., HANCOCK-RONEMUS, A. & GILLIS, J. A. 2020. Adult chondrogenesis
and spontaneous cartilage repair in the skate, Leucoraja erinacea. *Elife,* 9.

MAXWELL, E. E., FROBISCH, N. B. & HEPPLESTON, A. C. 2008. Variability and
conservation in late chondrichthyan development: ontogeny of the winter skate
(Leucoraja ocellata). *Anat Rec (Hoboken),* 291**,** 1079-87.
NAKAMURA, T., KLOMP, J., PIERETTI, J., SCHNEIDER, I., GEHRKE, A. R. &
SHUBIN, N. H. 2015. Molecular mechanisms underlying the exceptional adaptations
of batoid fins. *Proc Natl Acad Sci U S A,* 112**,** 15940-5.
NIEMANN, S., ZHAO, C., PASCU, F., STAHL, U., AULEPP, U., NISWANDER, L.,
WEBER, J. L. & MULLER, U. 2004. Homozygous WNT3 mutation causes tetra-
amelia in a large consanguineous family. *Am J Hum Genet,* 74**,** 558-63.
NISHIMURA, O., HARA, Y. & KURAKU, S. 2017. gVolante for standardizing
completeness assessment of genome and transcriptome assemblies. *Bioinformatics,*
33**,** 3635-3637.
O'NEILL, P., MCCOLE, R. B. & BAKER, C. V. 2007. A molecular analysis of neurogenic
placode and cranial sensory ganglion development in the shark, Scyliorhinus canicula.
Dev Biol, 304**,** 156-81.
PARR, B. A., SHEA, M. J., VASSILEVA, G. & MCMAHON, A. P. 1993. Mouse Wnt genes
exhibit discrete domains of expression in the early embryonic CNS and limb buds.
Development, 119**,** 247-61.
PATRO, R., DUGGAL, G., LOVE, M. I., IRIZARRY, R. A. & KINGSFORD, C. 2017.
Salmon provides fast and bias-aware quantification of transcript expression. *Nat*
Methods, 14**,** 417-419.
PEARSE, R. V., 2ND, VOGAN, K. J. & TABIN, C. J. 2001. Ptc1 and Ptc2 transcripts
provide distinct readouts of Hedgehog signaling activity during chick embryogenesis.
Dev Biol, 239**,** 15-29.
PICELLI, S., FARIDANI, O. R., BJORKLUND, A. K., WINBERG, G., SAGASSER, S. &
SANDBERG, R. 2014. Full-length RNA-seq from single cells using Smart-seq2. *Nat*
Protoc, 9**,** 171-81.
REID, B. S., YANG, H., MELVIN, V. S., TAKETO, M. M. & WILLIAMS, T. 2011.
Ectodermal Wnt/beta-catenin signaling shapes the mouse face. *Dev Biol,* 349**,** 261-9.
RICHARDSON, J., SHONO, T., OKABE, M. & GRAHAM, A. 2012. The presence of an
embryonic opercular flap in amniotes. *Proc Biol Sci,* 279**,** 224-9.
ROBINSON, M. D., MCCARTHY, D. J. & SMYTH, G. K. 2010. edgeR: a Bioconductor
package for differential expression analysis of digital gene expression data.
Bioinformatics, 26**,** 139-40.
RUDNICKI, J. A. & BROWN, A. M. 1997. Inhibition of chondrogenesis by Wnt gene
expression in vivo and in vitro. *Dev Biol,* 185**,** 104-18.
SCHERZ, P. J., HARFE, B. D., MCMAHON, A. P. & TABIN, C. J. 2004. The limb bud
Shh-Fgf feedback loop is terminated by expansion of former ZPA cells. *Science,* 305**,**
396-9.
SHIMOMURA, Y., AGALLIU, D., VONICA, A., LURIA, V., WAJID, M., BAUMER, A.,
BELLI, S., PETUKHOVA, L., SCHINZEL, A., BRIVANLOU, A. H., BARRES, B.
A. & CHRISTIANO, A. M. 2010. APCDD1 is a novel Wnt inhibitor mutated in
hereditary hypotrichosis simplex. *Nature,* 464**,** 1043-7.
SHONE, V. & GRAHAM, A. 2014. Endodermal/ectodermal interfaces during pharyngeal
segmentation in vertebrates. *J Anat,* 225**,** 479-91.
SLEIGHT, V. A. & GILLIS, J. A. 2020. Embryonic origin and serial homology of gill arches
and paired fins in the skate, Leucoraja erinacea. *Elife,* 9**,** 2020.07.02.183665.
STREIT, A., LEE, K. J., WOO, I., ROBERTS, C., JESSELL, T. M. & STERN, C. D. 1998.
Chordin regulates primitive streak development and the stability of induced neural

cells, but is not sufficient for neural induction in the chick embryo. *Development,* 125**,**
507-19.
SUMMERHURST, K., STARK, M., SHARPE, J., DAVIDSON, D. & MURPHY, P. 2008.
3D representation of Wnt and Frizzled gene expression patterns in the mouse embryo
at embryonic day 11.5 (Ts19). *Gene Expr Patterns,* 8**,** 331-48.
SUN, X., ZHANG, R., CHEN, H., DU, X., CHEN, S., HUANG, J., LIU, M., XU, M., LUO,
F., JIN, M., SU, N., QI, H., YANG, J., TAN, Q., ZHANG, D., NI, Z., LIANG, S.,
ZHANG, B., CHEN, D., ZHANG, X., LUO, L., CHEN, L. & XIE, Y. 2020. Fgfr3
mutation disrupts chondrogenesis and bone ossification in zebrafish model mimicking
CATSHL syndrome partially via enhanced Wnt/beta-catenin signaling. *Theranostics,*
10**,** 7111-7130.
TAKENAKA, K., KISE, Y. & MIKI, H. 2007. GSK3beta positively regulates Hedgehog
signaling through Sufu in mammalian cells. *Biochem Biophys Res Commun,* 353**,**
501-8.
TEN BERGE, D., BRUGMANN, S. A., HELMS, J. A. & NUSSE, R. 2008. Wnt and FGF
signals interact to coordinate growth with cell fate specification during limb
development. *Development,* 135**,** 3247-57.
THIERY, A.P., STANDING, A.S., COOPER, R.L. & FRASER, G.J. 2022. An epithelial
signalling centre in sharks supports homology of tooht morphogenesis in vertebrates.
eLife, 11, e73173.
VEITCH, E., BEGBIE, J., SCHILLING, T. F., SMITH, M. M. & GRAHAM, A. 1999.
Pharyngeal arch patterning in the absence of neural crest. *Curr Biol,* 9**,** 1481-4.
WALL, N. A. & HOGAN, B. L. 1995. Expression of bone morphogenetic protein-4 (BMP-
4), bone morphogenetic protein-7 (BMP-7), fibroblast growth factor-8 (FGF-8) and
sonic hedgehog (SHH) during branchial arch development in the chick. *Mech Dev,*
53**,** 383-92.
WITTEN, P. E. & HALL, B. K. 2003. Seasonal changes in the lower jaw skeleton in male
Atlantic salmon (Salmo salar L.): remodelling and regression of the kype after
spawning. *J Anat,* 203**,** 435-50.
ZHANG, X., CHEONG, S. M., AMADO, N. G., REIS, A. H., MACDONALD, B. T.,
ZEBISCH, M., JONES, E. Y., ABREU, J. G. & HE, X. 2015. Notum is required for
neural and head induction via Wnt deacylation, oxidation, and inactivation. *Dev Cell,*
32**,** 719-30.

1013

Figure 2 Supplement 1: Endodermal origin of the chick PEM. (A) Topical

application of CCFSE to chick embryos at HH11 results in specific labelling of the

surface ectoderm. Labelled embryos were incubated to HH19–20, by which time (**B**)

SHH is expressed in the PEM of the hyoid arch (yellow dashed line). Imaging of

CCFSE in the adjacent section from the same embryo as in **B** reveals (**C**) CCFSE-

labelled hyoid arch ectoderm (white dashed line) extending up to (but excluding) the

PEM**.** (**D**) Microinjection of the pharyngeal cavity with CM-DiI at HH11 results in

specific labelling of the foregut endoderm. CM-DiI-labelled embryos were grown to

HH19–20, at which point (**E***)* *SHH* is expressed in the PEM of the hyoid arch (yellow

dashed line). (**E'**) Co-labelling of cells of the PEM with *SHH* and CM-DiI indicates

endodermal origin of this signalling centre. A–P: Anterior–posterior axis, hpi: hours

post injection, NT: neural tube. Scale bars: A = 80μm, B–E = 20 μm.

1026

1027

1028

1029

1030

1031

1032

1033

1034

1035

1036

1037

1038

1039

Figure 3 Supplement 1: The chick PEM co-expresses *SHH*, *FGF8* and *BMP7*.

(**A**) At HH15, (**B**) endodermal pouches delineate the pharyngeal arches. (**C**) *SHH* is expressed in the posterior endodermal epithelium of the developing hyoid arch, while (**D**) *FGF8* is expressed in the lateral hyoid arch ectodermal arch epithelium and posterior endodermal epithelium, and (**E**) *BMP7* is expressed throughout the endodermal epithelium. (**F**) By HH17, (**G**) the hyoid arch begins to expand laterally, and (**H**) *SHH,* (**I**) *FGF8,* and (**J**) *BMP7* transcripts appear to be co-expressed in a domain of posterior endodermal epithelium (the presumptive PEM; indicated by red asterisk). (**K**) By HH19, (**L**) the hyoid arch is expanding caudally, and (**M**) *SHH*, (**N**) *FGF8* and (**O**) *BMP7* are co-expressed in the PEM. (**P**) By HH24, (**Q**) the hyoid arch is expanding posteriorly, lateral to the PEM and (**R**) *SHH*, (**S**) *FGF8* and (**T**) *BMP7*

1051   transcripts remain co-localised at the PEM. (**U**) ISH by HCR at HH20 shows co-

1052   expression of (**U'**) *SHH*, (**U''**) *FGF8* and (**U'''**) *BMP7* in the PEM in chick, with these

1053   three genes sharing a sharp anterior boundary. Epi: epithelial, Mes: mesenchymal.

1054   Scale bars: A,F,K,P = 2mm, all others = 20 μm.

1055

1056

1057

1058

1059

1060

1061

1062

Figure 6 Supplement 1: *WNT* pathway genes are expressed in and around the PEM in chick. (**A, A'**) ISH by HCR for *SHH* was performed alongside each gene of interest as a marker of the PEM. For ease of visualisation in **B–I**, *SHH* expression is not shown but rather these PEM cells are outlined green, and the arch is outlined in white for orientation. (**B**) *WNT2B,* (**C**) *WNT3,* (**D**) *WNT4* and (**E**) *WNT9B* are expressed predominantly in the ectodermal arch epithelium adjacent to the PEM. (**F**) *APCDD1* is expressed in distal arch mesenchyme and weakly in the overlying arch epithelium. (**G**) *NOTUM* is expressed in distal arch epithelium and mesenchyme, and

is expressed particularly strongly in the cells of the PEM and in the mesenchyme cells

under the PEM-adjacent ectodermal epithelium. (**H**) *KREMEN1* is expressed in

ectodermal epithelium adjacent to the PEM, and in distal arch mesenchyme. (**I**) *AXIN2*

is expressed broadly throughout the distal arch epithelium and mesenchyme, with

particularly strong expression in the anterior–distal mesenchyme and in cells of the

PEM. Scale bar: 50 μm.

1078

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

Figure 6 Supplement 2: Pharmacological inhibition of Wnt signalling in skate leads to downregulation of Wnt pathway genes. ISH by HCR was performed on sections of S26 skate embryos treated for 72hrs with either DMSO (control) or the canonical Wnt inhibitor IWR1. *Shh* was performed alongside each gene of interest as a marker of the GAER. (**A–A'**) *Axin2* is broadly expressed throughout the distal arch epithelium and mesenchyme in control embryos, with particularly strong expression in the cells of the GAER and in developing gill buds. (**B**) *Axin2* expression is substantially depressed with IWR1 treatment. (**C–C'**) *Apcdd1* is expressed in the cells of the GAER, and also within anterior–distal mesenchyme in control embryos, and (**D**) both of these expression domains show reduced expression with IWR1 treatment. (**E–E'**) *Notum* is expressed within anterior-distal arch mesenchyme in control embryos. (**F**) *Notum* expression is completely lost in the anterior-distal arch mesenchyme with IWR1 treatment. All images are representative of experiments performed in triplicate. For ease of visualisation in (**A', B, C', D, E', F**) *SHH* expression is not shown but rather these GAER cells are outlined green, and the arch is outlined in white for orientation. Scale bar: 50 μm.