Expanded Distribution and Predicted Suitable Habitat for the Critically Endangered

Yellow-tailed Woolly Monkey (*Lagothrix flavicauda*) in Peru

Melissa A. Zarate[1,2], Sam Shanee[3,4], Elvis Charpentier[5], Yeissy Sarmiento[4], and Christopher A.

Schmitt[1,2]

[1] Department of Anthropology, College of Arts and Sciences, Boston University, Boston, MA USA

[2] Department of Biology, College of Arts and Sciences, Boston University, Boston, MA USA

[3] Neotropical Primate Conservation, Windrush, Looe Hill, Seaton, Cornwall, UK

[4] Asociación Neotropical Primate Conservation Perú, Jr Andalucia #373, Moyobamba, San Martin,

Perú

[5] Equipo Primatológico del Perú, Calle Internacional M-43, Iquitos, Maynas, Perú

Corresponding authors' emails: mazarate@bu.edu; caschmit@bu.edu

Address: Department of Anthropology, Boston University

232 Bay State Road

Boston, MA 02215 USA

Submission date: 5/19/22

ABSTRACT

The Tropical Andes Biodiversity Hotspot holds a remarkable number of species at risk of

extinction due to anthropogenic habitat loss, hunting and climate change. One of these species, the

Critically Endangered yellow-tailed woolly monkey (*Lagothrix flavicauda*), was recently sighted

in Junín region, 206 kilometres south of its previously known distribution. The range extension,

combined with continued habitat loss, calls for a re-evaluation of the species' distribution and

available suitable habitat. Here, we present novel data from surveys at 53 sites in the regions of

Junín, Cerro de Pasco, Ayacucho and Cusco. We encountered *L. flavicauda* at 9 sites, all in Junín,

and the congeneric *L. l. tschudii* at 20 sites, but never in sympatry. Using these new localities along

with all previous geographic localities for the species, we made predictive Species Distribution

Models based on Ecological Niche Modelling using a generalized linear model and maximum

entropy. Each model incorporated bioclimatic variables, forest cover, vegetation measurements,

and elevation as predictor variables. Model evaluation showed >80% accuracy for all measures.

Precipitation was the strongest predictor of species presence. Habitat suitability maps illustrate

potential corridors for gene flow between the southern and northern populations, although much

of this area is inhabited by *L. l. tschudii*. An analysis of the current protected area (PA) network

showed ~47% of remaining suitable habitat is unprotected. With this, we suggest priority areas for

new protected areas or expansions to existing reserves that would conserve potential corridors

between *L. flavicauda* populations. Further surveys and characterization of the distribution in

intermediate areas, combined with studies on genetic flow, are still needed to protect this species.

Key Words (7/8 used): *Oreonax*, Maxent, Generalized Linear Modelling, Conservation, Species

Distribution Modelling (SDM), Range Expansion

Introduction

Changes in species' distributions are of major concern for conservationists due to the

continual expansion of human populations and land use (Hansen *et al.* 2001; Newbold *et al.* 2014;

Dai *et al.* 2021) that shift, contract, or fragment a species' geographic range. Such altered ranges

may isolate populations, enhancing the likelihood of inbreeding depression or local extinction

(Ewers and Didham 2005; Calkins *et al.* 2021; Solórzano-Garcia *et al.* 2021). Information on the

relative distributions of different populations of a species can reveal what types of landscapes or

species community composition—in the case of competitive exclusion—act as geographic or

ecological barriers to gene flow (Case and Taper 2000; Blair and Melnick 2012; Sales *et al.* 2019;

Pázstor *et al.* 2020). To conserve natural biodiversity in the face of expanding land use by humans,

conservation management must operate on a landscape scale, understanding species-ecosystem

relationships and species' responses to habitat change (Bellamy *et al.* 2013; Robillard *et al.* 2015;

Xioa *et al.* 2019). Informed management planning involves understanding the true current

distribution of a species and the necessary habitat components to predict how distribution changes

will occur in the future, and the best possible options for area protection.

The yellow-tailed woolly monkey (*Lagothrix flavicauda*) is considered Critically

Endangered and amongst the most threatened primate species in the world (Mittermeier *et al.* 2012;

Shanee *et al.* 2021), making it a key focus of conservation initiatives in northern Peru (Shanee *et*

al. 2018). The Tropical Andes Biodiversity Hotspot (TABH), to which the species is endemic, is

among the top five hotspots predicted to lose the most biological diversity due to continued

anthropogenic activities (Brooks *et al.* 2002). In northern Peru, the greatest threat to species and

habitats in the TABH is habitat destruction for cattle ranching, logging, and other, mainly small

scale, economic activities, fuelled by a growing human population and aided by the construction

of highways and incentivized access to lands (Oliveira *et al.* 2007; Shanee 2011; Shanee 2012;

Programa Bosques 2015; GIZ 2016; Laurance 2018; Shanee and Shanee 2016). With this migrant

economic development, roads are continuously built to access new areas, creating a cycle of

increasing deforestation for the extraction of resources and settlement (Gallice, Larrea-Gallegos

and Vázquez-Rowe 2019). Hunting of primates for bushmeat and the pet trade is also putting *L.*

flavicauda and other species at further risk (Shanee 2011; Shanee 2012; Shanee *et al.* 2017).

The geographic range of *L. flavicauda* – long thought to be restricted to the northern regions

of Amazonas and San Martín and neighbouring areas of Huánuco, La Libertad, and Loreto – has

been continuously re-evaluated through survey efforts (Mittermeier, de Macedo-Ruiz, Luscombe

1975; Graves *et al.* 1980; Leo Luna 1980; Parker and Barkley 1981; Butchart *et al.* 1995; Shanee

et al. 2007; Shanee 2016; Shanee 2011). Recent observations in the region of Huánuco have found

L. flavicauda populations as far as the south eastern border with the region of Pasco (Aquino *et al.*

2015; Aquino, Garcia and Charpentier 2016). Camera trap observations led to the discovery of a

new population of *L. flavicauda* in 2018 (McHugh *et al*. 2020). This new population, in the

Inchatoshi Kametsha Conservation Concession, Junín, is about 206 km south of the previously

known southern range limit for the species (McHugh *et al.* 2020), with no sightings in intermediate

areas.

Lagothrix flavicauda is generally restricted to montane forests between 1,400 to 2,800 m

above sea level (m.a.s.l.) along the eastern slopes of the Andes (Aquino *et al*. 2017, Shanee 2011),

with occasional occurrences below this elevation (Allgas *et al.* 2014; Paterson and Lopez Wong

2014). Their preferred habitat depends on local climate and forest composition (Shanee 2016;

Almeyda-Zambrano *et al.* 2019), and they are able to survive at least for a time in moderately

disturbed habitats when hunting pressure is low (Aquino *et al*. 2015; Shanee and Shanee 2015).

This may be, in part, due to a flexible diet consisting of flowers, leaf petioles, epiphytic roots,

vertebrates, and soil, although they are predominantly frugivorous. Fruits make up almost half of

their dietary intake, which is dominated by a handful of species and genera with large fleshy fruits

(Shanee and Shanee 2011, Shanee 2014; Fack *et al.* 2018), making food tree density and phenology

important factors for sustaining viable populations. Previous studies suggest that the protected area

(PA) network in northern Peru is insufficient to protect *L. flavicauda* from the consequences of

human population growth, habitat degradation, and climate change (Buckingham and Shanee

2009; Shanee 2016). As the number of PAs in Peru change, along with our knowledge of primate

species distributions, it is critical to re-assess their coverage with current data.

Many conservation and research initiatives use *L. flavicauda* as a flagship species in Peru

(Shanee *et al*. 2018). Although these have led to an increasing number of private, communal, and

state PAs within the species distribution (Shanee *et al.* 2017; Shanee *et al.* 2020), they have

predominantly focused on populations in Amazonas and San Martín (Shanee and Shanee 2015).

Field surveys are vital to determining the species' distribution and habitat preferences outside these

well-studied regions; however, they are often expensive, time consuming, and can be unfeasible

depending on the terrain, access, and public order limitations (Young 1996; Shanee and Shanee

2016). Predictive species distribution modelling (SDM) can highlight optimal areas for field

surveys through comparison of the ecological conditions found within a given species' known

range, with those across a wider area, showing where conditions are most favourable for the

species presence (Phillips, Anderson and Schapire 2006; Ramirez-Villegas *et al.* 2014; Guisan,

Thuiller and Zimmerman, 2017).

Species' distributions are predicated on three main conditions: the species must have the

means to disperse into and out of a habitat, the habitat must have the correct combination of

environmental variables to make it suitable, and the abiotic conditions must be able to maintain

necessary species interactions (Guisan *et al.* 2017). SDM considers the second condition by using

known habitat parameters to determine which are most important to the distribution of the focal

species, and uses them to locate potentially suitable habitat (Dong *et al*. 2019; Cianfranni *et al.*

2010; Liu *et al.* 2019). Such modeling may be used to prioritize search efforts when financial or

human resources are not available for surveying a region of interest in person. For endangered

species, analyzing correlations between these variables and known species occurrences can

highlight conditions which the species requires for its long-term survival in the region of interest,

and can be used to develop more effective conservation strategies. With this, conservationists can

determine why a species may not be present in a particular area (Liu *et al.* 2019) or increase

research and conservation effort in areas that appear to be suitable (LaRue and Nielsen 2008).

Here we compare the results of two SDM approaches—Generalized Linear Modelling

(GLM) and Maximum Entropy (MaxEnt)—using the results of our recent field surveys in central

and southern Peru. We created SDMs combining these novel localities with all previous records

to create the most complete SDM for the species to date. We use the results of these approaches to

highlight remaining suitable *L. flavicauda* habitat, particularly in its less-studied southern range.

Finally, we evaluate the current Peruvian PA network's coverage of suitable habitat within the

species' distribution, noting priority areas for new PAs, conservation corridors, and future surveys

between the northern and recently-discovered Junín populations.

Methods

Population Surveys

We conducted five surveys between May 2019 and May 2021. Due to Covid-19 related

travel restriction, no surveys were conducted between May and December 2020, with the May

2020 survey being cut short due to implementation of national quarantine measures. Surveys were

carried out in the regions of Ayacucho, Cerro de Pasco, Cusco, and Junín. Survey sites were

selected based on preliminary MaxEnt models which used previously published localities, and

updated with new sightings as the study progressed, with sites selected considering access routes,

land ownership and researcher safety.

Survey efforts followed methods of previous surveys for the species (Shanee 2011). We

gathered locality data along existing trail systems with local residents as field guides. Occasionally

new trails were opened to enter new areas; however, this was typically avoided to minimize habitat

disturbance. We visited sites for 2-9 days, with effort determined by available habitat size, and the

possibility of *L. flavicauda* being found or confirmation of its presence. We recorded localities and

points of visual or audio detection of all primates with a handheld GPS. We also gathered

secondary evidence from local informants in and around the areas visited. Using images and verbal

descriptions of *Lagothrix* spp. we cross-referenced information from multiple informants at each

site.

Species Distribution Modelling

We selected two standard SDM methods— GLM and MaxEnt. To avoid overestimation of

suitable habitat, we limited prediction outputs to between -80° W and -65° W, and -5° S and -15°

S. We filtered this to only include areas between 1,000 and 3,500 m.a.s.l., i.e., within the species'

altitudinal range, which includes the entirety of the eastern slopes of the Peruvian Andes. Models

used presence points from our own surveys combined with localities from recent published studies,

i.e., ≤10 years (Shanee 2011; Allgas *et al*. 2015; Aquino *et al.* 2016; Aquino *et al.* 2017; McHugh

et al. 2020), and an additional six localities from the Global Biodiversity Information Facility

(GBIF Secretariat 2021) (Fig. 1; Table S1).

Environmental data were all from publicly available sources (Table S2). We used the 19

bioclimatic variables from *WorldClim2*, available in the R package raster v. 3.1-5 (Hijmans *et al.*

2020) and elevation data from the NASA Shuttle Radar Topography Mission (SRTM; v. 4) at 30-

meter resolution. We used forest cover data from the Global Forest Change (v. 1.7) dataset of

percent tree cover from the year 2000, in which tree cover is given as a gradient of percentage

canopy connectivity per 30 m pixel for vegetation taller than 5 m. Forest loss and gain between

the years 2000 and 2018 were also analyzed, but were excluded from the model due to an

insufficient number of areas experiencing any recorded loss or gain (fewer than 10 pixels within

the study region, which did not coincide with the location of any presence points). We also used

the Vegetation Health Product (VHP) dataset, at 1 km resolution, from the NOAA Centre for

Satellite Applications and Research, which has been created from an algorithm providing estimates

of moisture, thermal, and wind conditions in combination with infrared imaging to achieve an

estimate of vegetation conditions in response to weather impacts. All data layers were re-sampled

to 30 m resolution.

Generalized Linear Model

All GLMs were evaluated in R (v. 3.5.2; R Core Team 2018). We modelled the ecological

niche of *L. flavicauda* as a binomial response variable of presence/pseudoabsence using a binomial

variance and logistic link function. We randomly selected pseudoabsence points to counter

sampling bias from the lack of absence data existing for the species using the *randomPoints*

function in the *dismo* package v.1.1-4 (Hijmans *et al.* 2017), generating 355 in the study area

within an elevation mask including only areas within the altitudinal range. All presence points

were excluded from the background point extraction process. We then extracted and standardized

values from all environmental layers at each pseudoabsence and presence point to be used in the

models. Models were built excluding correlated predictor variables based on Pearson's *r* value (±

0.75). Our final model was selected using the corrected Akaike Information Criterion (AICc)

procedure (Akaike 1973; Hurvich and Tsai 1993).

We evaluated the predictive ability of our model using the area under the ROC curve

(AUC), which summarizes model accuracy by giving the probability that the model ranks random

presence sites over pseudoabsence sites. We calculated the AUC by building the models from a

training set, and then applying it to the remainder of the data as a test set to evaluate predictive

performance (Fielding and Bell 1997; Hirzel *et al.* 2006). Due to the relatively small number of

presence points, we bootstrapped this process 1000 times using multiple distributions of presence

data in the training and testing sets (Hein *et al.* 2007; Guisan *et al.* 2017). We also tested model

accuracy using Cohen's kappa statistic and percent accuracy from a confusion matrix as a

threshold-dependent method of evaluation (Guisan *et al.* 2017). Using training and test data, we

set a probability threshold so that any point with a presence probability of 0.4 or higher was marked

as a presence when establishing a confusion matrix of model predictions. The kappa statistic value

ranges from -1 (complete disagreement between predicted and actual values) to 0 (predictions

equated to random chance), to +1 (complete agreement with actual values).

Maximum Entropy Model

The MaxEnt model was created and evaluated in MaxEnt Programming Software (v. 3.4.4;

Phillips 2006) and re-evaluated in R. We applied default settings to our MaxEnt run, with the

following exceptions: background predictions were written for evaluation and mapping purposes

and we selected a random test percentage of 25% of our presence points. We used the same non-

correlated variables as in the GLM. MaxEnt measures variable contribution by permutation

importance determined by randomly permuting the values of each variable among the training

points, measuring the resulting decrease in training AUC, and normalizing these values to give a

percentage (Phillips *et al.* 2006). MaxEnt evaluates the model using a ROC curve and calculating

the model AUC. We corroborated the given value by bootstrapping the AUC calculation 1000

times in R using the background prediction values.

Suitable Habitat Classification and Protected Area Assessment

To assess habitat suitability, we assigned habitat based on the predicted probability of

species presence to categories of "Good" (P(robability of species presence) between 0.25 and

0.75), "Very Good" (P>0.75), and "Low" (P<0.25), with "Good" and "Very Good" being

considered suitable habitat. To incorporate the probable impact of hunting into our results, we

considered habitat within 1 km of human settlement to have high hunting pressure for *L. flavicauda*

(Shanee 2016). We used data from Humanitarian OpenStreetMap Team (2020) to distinguish

human settlements, and deemed any suitable habitat within a 1 km buffer of settlement as low

suitability. We overlaid the suitable habitat predicted by both models to find where the predictions

intersected. Finally, we conducted an analysis of the PA coverage of suitable *L. flavicauda* habitat

by overlaying the Peruvian PA network from the World Database of Protected Areas (WDPA;

UNEP-WCMC 2020) with the habitat predictions and calculated the percent of habitat that was

considered Good or Very Good within the network. This was repeated with the network of non-

governmental conservation concessions in Peru, downloaded from the National Forestry Service

(SERFOR), as these concessions are not included in the WDPA files.

Data Accessibility

All open-source data used for modelling are openly accessible via the links provided in

the references. Presence data, prediction results, and all code can be found at

https://doi.org/10.5061/dryad.dz08kps0g.

Fig. 1 *Lagothrix flavicauda* occurrences used as presence points in habitat suitability modelling
(HSM) in this study.

Results

Population Surveys

In total we surveyed 53 new sites in central and southern Peru. We encountered *L.*
flavicauda at 9 of these sites, all to the north of the Mantaro river in Junín (Fig. 2). In surveys south
of this point we did not find any evidence of the species, from our own surveys or from local

informants, suggesting a possible southern limit for the species between ~11.6 and ~12° south

(Table 1). We recorded the congeneric Peruvian woolly monkey (*L. lagotricha tschudii*) at 20 sites,

both north and south of the southern *L. flavicauda* population. At no sites were the species

sympatric.

Fig. 2 Survey results for *L. flavicauda* and *L. lagotricha tschudii*.

Table 1. Coordinates where *L. flavicauda* and *L. l. tschudii* were found in our surveys.
256

Region	Species	Longitude	Latitude	m.a.s.l.
Junín	*L. flavicauda*	-74.7659	-11.4530	2179
Junín	*L. flavicauda*	-74.7568	-11.4378	1869
Junín	*L. flavicauda*	-74.7609	-11.4490	1866
Junín	*L. flavicauda*	-74.6467	-11.4496	1609
Junín	*L. flavicauda*	-74.6770	-11.4678	1825
Junín	*L. flavicauda*	-74.7609	-11.4490	1866
Junín	*L. flavicauda*	-74.5504	-11.6610	2012

Junín	*L. flavicauda*	-74.5793	-11.6301	2067
Junín	*L. flavicauda*	-74.7372	-11.4069	1422
Pasco	*L. l. tschudii*	-75.4709	-10.6859	1410
Pasco	*L. l. tschudii*	-75.3698	-10.7384	1257
Pasco	*L. l. tschudii*	-75.4238	-10.7223	1323
Pasco	*L. l. tschudii*	-75.5182	-10.5882	1612
Pasco	*L. l. tschudii*	-75.5358	-10.5748	2088
Pasco	*L. l. tschudii*	-75.5171	-10.6693	2349
Junín	*L. l. tschudii*	-75.4529	-10.9879	1771
Junín	*L. l. tschudii*	-75.4635	-10.9928	2490
Junín	*L. l. tschudii*	-75.4307	-10.9911	1622
Junín	*L. l. tschudii*	-75.4352	-11.0447	1672
Junín	*L. l. tschudii*	-75.3518	-11.4285	2365
Junín	*L. l. tschudii*	-74.8955	-11.2327	1609
Junín	*L. l. tschudii*	-74.9112	-11.2358	1795
Junín	*L. l. tschudii*	-74.9229	-11.2421	1080
Pasco	*L. l. tschudii*	-74.6540	-10.8546	1465
Pasco	*L. l. tschudii*	-74.6510	-10.8259	1410
Pasco	*L. l. tschudii*	-74.6600	-10.8186	1257
Pasco	*L. l. tschudii*	-75.0401	-10.6526	1323
Pasco	*L. l. tschudii*	-75.0399	-10.6866	1612
Cusco	*L. l. tschudii*	-72.8110	-12.7401	1948

257
258

Species Distribution Modelling

After the multicollinearity reduction, the GLM with the lowest AICc (242) contained elevation, mean diurnal temperature range, isothermality, precipitation of wettest month, precipitation seasonality, precipitation of coldest quarter, percent forest cover and VHP as predictor variables. These were the variables used in both model types. The suitable habitat predicted by the two models overlapped by 47.5%, with the GLM being more conservative in the southern regions (Fig. 3).

Fig 3. Suitable habitat (Good and Very Good) as indicated by the GLM, MaxEnt model, and their
overlap.

Generalized Linear Modelling

The GLM found all predictor variables to be significantly associated with the likelihood of

L. flavicauda presence with the exceptions of isothermality and forest cover. The variable most

highly associated with *L. flavicauda* habitat was precipitation seasonality, which had a negative

relationship. Precipitation of the wettest month had the strongest positive correlation (Table 2).

Table 2. Correlation coefficients and standard error of residual deviances for each standardized
predictor variable in the GLM. Estimates indicate the average change in the response variable
associate with a one unit increase in the (standardized) predictor variable. Shading indicates
significant association.
279

Predictor	Estimate	Standard Error	P value
Intercept	-6.228	1.008	<0.0005
Elevation	-0.768	0.249	0.00201
Bio2	0.983	0.466	0.0349
Bio3	0.616	0.327	0.0600
bio13	1.653	0.738	0.0251
bio15	-11.019	2.309	<0. 0005
bio19	-4.831	1.107	<0. 0005
Forest Cover	0.441	0.306	0.149
VHP	- 1.256	0.247	<0.0005

Based on the GLM, 1.4% of the study area was found to be Very Good habitat, and 11.4

% was considered Good (Table 3). Only a very small proportion of remaining habitat was within

the 1 km human settlement buffer (Table 3). Suitable habitats in the southern regions of the *L.*

flavicauda range run north-south through central Huánuco and Pasco, but coincide with areas

where the species has been found to be absent, or where *L. l. tschudii* was found instead (this study;

Aquino *et al.* 2017, 2019). In Junín, additional areas of Good habitat are present around and

immediately south of the newly discovered populations. Of all Very Good and Good habitat,

17.72% and 25.95%, respectively, were found to be within Peru's current PA network (Table 3),

most of which are in PAs in the northern regions of the study area, while much of the area between

the northern and Junín populations remains unprotected.

A single iteration of the threshold-independent AUC calculation gave an AUC of 0.93 (Fig.

S1). The bootstrapped evaluation gave a slightly lower mean AUC of 0.89 (95% CI 0.86-0.95).

Applying the GLM to the training dataset with a 0.4 threshold produced predictions with 81.76%

accuracy. The model was better able to correctly predict the pseudoabsences (86.36%) than

presences (59.26%), likely due to the larger amount of pseudoabsence points in the test dataset

(Table S3). The kappa statistic (0.47) indicated that the model produced predictions in good

agreement with the test data, though this was not significant (p=0.70).

Table 3. Percentages of the study area that the GLM and MaxEnt model considered to be Very
Good, Good and unsuitable and the amount of each habitat type within the protected area (PA)
network in Peru.
301

Modeling Approach	Habitat Suitability	Within Study Region (% of region- before human settlement)	Within Study Region (% of region- after human settlement)	Within PAs (% of habitat)
GLM	Unsuitable	87.10	87.33	14.32
	Good	11.47	11.28	25.95
	Very Good	1.43	1.39	17.72
MaxEnt	Unsuitable	86.01	86.2	13.79
	Good	11.66	11.49	21.5
	Very Good	2.33	2.31	38.1

MaxEnt Model

The MaxEnt model corroborated that precipitation seasonality had a strong negative

correlation with species presence (Table 4). There were some differences in variable importance

in comparison to the GLM results, including relatively low importance of Precipitation of Coldest

Quarter (Table 2). The AUC for the MaxEnt model was 0.95 (Fig. S2) and corroborated by the

mean AUC from bootstrap (0.95; 95% CI 0.93-0.97), indicating greater accuracy than the GLM.

The MaxEnt model suggested a higher percentage of Good and Very Good habitats

compared to the GLM (Table 3; Fig. 4a), with percentages minimally effected after removing areas

within the human settlement buffer. Suitable habitat in the southern regions of the study area

included more Good habitat running north to south in Junín (Fig. 4b). Overlaying the PA network

also showed more suitable habitat overall within the network compared to the GLM, with 21.5%

and 38.1% of Good and Very Good being within PAs, respectively (Table 3; Fig. 5a), though much

of the suitable habitat in the southern portion of the species' range remains unprotected (Fig. 5b).

Fig. 4 Habitat suitability maps for *L. flavicauda* as predicted by the MaxEnt model a.)
throughout the described study area and b.) within the regions surrounding the southernmost *L.*
flavicauda occurrences.

Fig. 5 PA network coverage over habitat suitability as predicted by the Maxent model a.) over the entire study area and b.) within the regions surrounding the southernmost *L. flavicauda* occurrences

Discussion

We carried out the first survey of *L. flavicauda* in the recently discovered southern expansion of its known range (McHugh *et al.* 2020), and our results suggest that this is a very localized population isolate of the species. Models of suitable habitat including the regions south

of Huánuco and our new occurrence records provide important information on the habitat

availability for the species, thereby providing insight into priority areas for focused conservation

efforts and placement of conservation corridors and new protected areas. Our use of free, open-

source modelling platforms will make updating and refining this analysis relatively simple with

the addition of novel presence data. Our updated suitable habitats will hopefully lead to the

discovery of more *L. flavicauda* populations, possibly connecting the northern and southern

distributions.

Our surveys discovered multiple new sightings of *L. flavicauda* in central Junín, but only

of *L. l. tschudii* in Cerro de Pasco, northern Junín and northern Cusco, and neither *Lagothrix* taxa

in Ayacucho. The lack of encounters or secondary evidence of *L. flavicauda* in areas with evidence

of *L. l. tschudii* may be due, in large part, to competitive exclusion. *L. l. tschudii* has been found

to be one of the most abundant high-altitude species in previous surveys in Cerro de Pasco and

Huánuco (Aquino *et al.* 2019), and inhabits forests with niche conditions and at elevations suitable

for *L. flavicauda* (Aquino *et al.* 2016; Serrano-Villavicencio *et al.* 2021). When comparing the

suitable habitat in our models with our observations of *L. l. tschudii*, we found that most are in

areas well-suited to the species (Fig. 6), paralleling the work of Aquino *et al.* (2016) which

indicated that *L. flavicauda* occupied similar but separate, higher-altitude habitats than *L. l.*

tschudii. While competitive exclusion may explain the replacement of either *Lagothrix* taxa with

its congener, further surveys are needed to rule out possible sympatry in other areas, and to confirm

the absence of both taxa in Ayacucho, with future models incorporating both species.

352
353 **Fig 6** *L. l. tschudii* occurrences compared to suitable habitat predicted by the MaxEnt model. *L.*
354 *flavicauda* and *L. l. tschudii* were never found occurring at the same site.
355

The lack of *Lagothrix* sightings in Ayacucho does not conclusively confirm absence,

especially in areas where our survey efforts were limited by security issues. Almost 60% of the

areas visited in this region were cultivated for coca (for cocaine production), while the remaining

were small forest patches. More continuous forests did exist, but these were beyond the local

jurisdictions, and controlled by armed guerrilla groups involved in drug trafficking, which we did

not enter, either due to lack of permission or for our safety and that of our guides. Local informants

indicated the presence of "large monkeys," in these forests, and that they are rarely seen in the

smaller patches. Informants also indicated that these monkeys were more common in previous

years, but that hunting by indigenous Ashaninka groups in the area diminished their numbers. This,

with further increases in hunting in combination with deforestation due to coca cultivation likely

contribute to the species' rarity. In the patches we were able to survey in these areas we only

recorded the smaller bodied *Saguinus*, *Saimiri*, and *Cebus*. Finally, some of those interviewed in

the Union Mantara and Vizcatán areas in Junín claimed, based on photographs, to have seen *L.*

flavicauda in high elevation areas controlled by guerrilla groups. In particular, they pointed out

the area bordered by the Río Montaro as a place where *L. flavicauda* has definitely been seen.

During interviews, local people who had previously ceased coca cultivation in favour of coffee

and cacao indicated a recent return to coca in the area due to low market prices for legal crops.

This trend could have implications for future surveys and conservation actions.

Both of our models showed a strong negative correlation between *L. flavicauda* presence

and precipitation seasonality, consistent with the results of a study by Shanee (2016) in the northern

distribution of the species. This may be due to the phenology of arboreal food sources on which *L.*

flavicauda depends. Though the species has been found to use a variety of food resources and plant

parts, studies have shown seasonal shifts in feeding behaviour and dietary components (Shanee

2014; Shanee and Shanee 2011b). Andean species of *Cercropia*, for example, whose fruits and

leaves are commonly consumed by *L. flavicauda* (Shanee 2014), experience annual reproductive

changes due to climatic seasonality (Zalamea *et al.* 2011). Moreover, *L. flavicauda* have

commonly been observed consuming *Ficus* spp. fruit (Leo Luna 1980; Shanee 2014; Almeyda-

Zambrano *et al.* 2019), which have inter- and intra-specific asynchronous fruiting; together with

abundant production, this accounts for the genus' importance for many frugivores (Bronstein *et*

al. 1990; Kattan and Valenzuela 2013). Overall, while modelling with bioclimatic variables can

aid in determining areas with suitable niche conditions, these models do not provide clear

explanations for a given species' presence or absence.

Changing environmental conditions due to current and future climate change, in

combination with increasing anthropogenic impacts in the TABH are predicted to decrease

climatic niche availability and, in turn, species richness of plants (Ramirez-Villegas *et al.* 2014).

These changes will likely lead to further habitat loss and population reductions in *L. flavicauda*

and other primate species (e.g., Shanee 2016). Research suggests that the southern Peruvian Andes

are experiencing a slight decrease in precipitation along with significant increases in surface

temperatures (Vuille *et al.* 2003). Monthly rainfall projections based on fourth report of IPCC

climate change models suggest a significant future increase in rainfall seasonality in the tropical

Andes (Lavado Casimiro *et al.* 2011). Our models suggest that precipitation and seasonality are

among the most important predictors of *L. flavicauda* presence, therefore such changes could have

dire consequences for the species. This is especially true as lower-altitude species migrate to

occupy higher elevations with climate change, increasing competition for resources (Fisher 2011).

Future scenarios are further complicated by the fragmentation of habitats and intervening barriers

to dispersal (Sales *et al.* 2019), which would limit *L flavicauda's* ability to colonize new areas.

The buffer used to estimate areas as "Low" suitability based on proximity to human

settlement is meant to reflect hunting pressure on *L. flavicauda* (Buckingham and Shanee 2009).

Biodemographic hunting models have been created comparing human population and

demographic data with primate population density to demonstrate the deleterious impact of

hunting on primate "prey" species (Levi *et al.* 2011). Although hunting is known to be one of the

greatest threats to *L. flavicauda* (Shanee 2012), the assumption that there is high hunting pressure

closer to human settlements can be problematic due to differing practices among communities.

Conversely, in areas where community conservation initiatives have been enacted, such as the

community of Yambrasbamba in Amazonas, hunting of the species has been largely eliminated

and populations of *L. flavicauda* have increased (Shanee *et al.* 2015). In addition to hunting,

different regions in Peru have varying wildlife trafficking trends and pathways in which primates

are often trafficked as pets or tourist attractions, with large-bodied primates being the most

common victims (Shanee *et al.* 2017). Given this, using pressures from human hunting and

trafficking activity as variables in models like this may require more detailed surveying of local

practices to get a more nuanced representation of localized anthropogenic pressures.

While both models showed high statistical accuracies in their predictions, our MaxEnt

model performed better. While the GLM was more conservative in the predicted amount of

suitable habitat in the southern regions, there was a large amount of overlap in predictions between

both models. As with all models, ours are only as accurate as the data used. We were limited by

the relatively low number of localities compared to many other primate habitat suitability models

(Liu *et al.* 2019). Regardless, our model predictions and survey results should encourage the

continuation of surveys in southern Peru to increase presence data and model accuracy for the

species.

Overlaying the PA network and the MaxEnt predictions showed that ~75% of suitable

habitat is currently unprotected, with most protection being in the northern regions of the species'

range (Fig. 5c). This is much lower than that of a gap analysis performed in 2009 (Buckingham

and Shanee 2009), likely due to the recent growth of private and communal protected areas, as

well as increases in state parks in the northern regions of the *L. flavicauda* range (Shanee et al.

2017; Shanee 2018; Shanee et al. 2020). Peru has 18.8% of its land covered by the current PA

network, with 91.7% being within governmental PAs and 8.3% in private or communal PAs

(Shanee *et al.* 2020). While national PAs generally cover larger areas, they operate at a regional

scale and rarely give rights or leadership roles to local people (Shanee *et al.* 2017; Horwich *et al.*

2012). The use of protected areas is key for the conservation of *L. flavicauda*, but new state

protected areas can require large areas, free of other land uses, and can take years to formally

create. Private and communal PAs, on the other hand, have proven effective for conservation in

the regions of Amazonas and San Martin where relatively high human population densities limit

opportunities for large state-run PAs (Horwich *et al.* 2012; Shanee *et al.* 2020). In these areas,

local people are the primary decision makers and enforce their own regulations, ensuring that the

methods of protection are in line with the values of their community. The promotion of

communally and privately managed protected areas, particularly in key corridor areas, and

education across landscapes can provide much needed protection (Shanee *et al*. 2015). Further,

conservation concessions cover a great amount of suitable habitat in San Martín (Fig. 5). The

inclusion of the conservation concessions caused the proportion of Very Good habitat within

protected areas to increase from 34.63% to 38.1 %.

In the southern portion of the species range, our analysis illustrated that only the small

areas of Pampa Hermosa and Pui Pui cover an area with some suitable *L. flavicauda* habitat near

the southern most populations. As such, the development of PA and community conservation

initiatives in the species' southern distribution are urgently needed, though successful conservation

actions in the areas highlighted here will depend on an understanding of the socioeconomic and

political situation to be able to garner local support (Chazdon *et al.* 2009).

Monitoring distributions, gene flow, and habitat suitability is critical to the conservation of

L. flavicauda. Future research should use genetic assessments of diversity between the southern

and northern populations of *L.* flavicauda to determine if gene flow is occurring. Patterns of

environmental variables should also be analysed between populations to better understand factors

that facilitate migration between populations, and how they could be constrained with climate

change and increasing human development (Sales *et al.* 2019). Using landscape genetic methods

could also determine what landscape and/or ecological features strongly facilitate or hinder gene

flow between populations and be used in planning new PAs or corridors (Olah *et al.* 2016).

460   Further investigation of the home ranges, and resource and land use of *L. flavicauda* will

allow for better interpretation of the importance of SDM predictor variables. Differences in these

attributes between the northern populations and the newly-discovered Junín populations will

specifically indicate the necessary factors a habitat needs to sustain viable populations. This should

involve working with local communities near and within the protected areas containing suitable

habitat in the south, with a particular focus on environmental education as our experience suggests

a lack of understanding of the importance of animals and ecosystem function within many local

immigrant (i.e., non-indigenous) communities. The presence of drug traffickers and armed groups

limited which sites we could visit, particularly in the area of Vizcatán in southern Junín, thus this

area remains un-surveyed, a situation which needs to be urgently rectified once conditions allow.

Further successful surveys will increase the number of presence and absence points for the species,

will allow for more accurate analyses of their environmental needs, and encourage the initiation

of conservation efforts in these areas to continue the growth of both state and private/community

led efforts for *L. flavicauda*. This will also increase the likelihood of finding other high-altitude

primate species localities and help us understand their distributions and conservation needs.

Acknowledgements

We thank the local communities we visited for allowing us to survey their surrounding

forests and for their guidance in finding the monkeys. This work was funded by Boston University,

and by Neotropical Primate Conservation thanks to grants from Primate Society of Great Britain,

American Society of Primatologists, International Primatological Society, and Primate

Conservation inc. Field work was carried out under Peruvian research permits (N° 173-2016-

SERFOR-DGGSPFFS, and addendums 213-2016, 350-2017, and 389-2019). We also thank Dr.

Suchi Gopal, who provided guidance in the performance of GIS analyses.

References

Akaike, H. (1973). Information theory and extension of the Maximum Likelihood principle.

Second International Symposium on Information Theory, Budapest, Hungary.

Almeyda-Zambrano, S.L. *et al.* (2019). Habitat preference in the critically endangered yellow

tailed woolly monkey (*Lagothrix flavicauda*) at La Esperanza, Peru. *Am. J. of Primatol.*

81: e23032.

Allgas, N., Shanee, S., Peralta, A. and Shanee, N. (2014). Yellow-tailed woolly monkey (*Oreonax*

flavicauda: Humboldt 1812) Altitudinal Range Extension, Uchiza, Perú. *Neotrop.*

Primates 21(2): 207-208.

Aquino, R., Zárate, R., López, L., García, G. and Charpentier, E. (2015). Current status and threats

to *Lagothrix flavicauda* and other primates in montane forest of the Región Huánuco.

Primate Conserv. 29: 1–11.

Aquino, R., Garcia-Mendoza, G. and Charpentier, E. (2016). Distribution and current status of the

Peruvian yellow-tailed woolly monkey (*Lagothrix flavicauda*) in montane forests of the

Región Huánuco, Peru. *Primate Conserv.* 30: 31-37.

Aquino, R., García, G., Charpentier, E. and Lòpez, L. (2017). Conservation status of *Lagothrix*

flavicauda and other primates in montane forests of San Martín and Huánuco, Peru.

Rev Peru. Biol. 24(1): 025-034.

Aquino, R., López, L., Falcón, R., Díaz, S. and Gálvez, H. (2019). First inventory of primates in

the montane forests of the Pasco and Ucayali regions, Peruvian Amazon. *Primate Conserv.*

33: 1-11

Bellamy, C., Scott, C. and Altringham, J. (2013). Multiscale, presence-only habitat suitability

models: fine-resolution maps for eight bat species. *J. Appl. Ecol.* 50: 892-901.

Blair, M.E. and Melnick, D.J. (2012). Scale-Dependent Effects of a Heterogeneous Landscape

on Genetic Differentiation in the Central American Squirrel Monkey (Saimiri oerstedii).

PLoS ONE 7(8): e43027.

Bronstein J.L, Gouyon, P., Gliddon, C., Kjellberg, F. and Michaloud, G. (1990). The ecological

consequences of flowering asynchrony in monoecious figs: a simulation study. *Ecology*

71: 2145-2156.

Butchart, S. H. M., Barnes R., Davies, C. W. N., Fernández, M. and Seddon, N. (1995)

Threatened mammals of the Cordillera de Colán, Peru. *Oryx* 29 (4):275-281.

doi:doi:10.1017/S003060530002127X

Brooks, T.M. *et al*. 2002. Habitat loss and extinction in the hotspots of biodiversity.

Conserv. Biol. 16(4): 909-923.

Buckingham, F. and Shanee, S. (2009). Conservation priorities for the Peruvian yellow tailed

woolly monkey (*Oreonax flavicauda*): a GIS risk assessment and gap analysis. *Primate*

Conserv. 24: 65–71.

Case, T.J. and Taper, M.L. (2000). Interspecific competition, environmental gradients, gene

flow, and the coevolution of species' borders. *Am. Nat.* 155(5), 583-605.

Calkins, M.T., Beever, E.A., Boykin, K.G., Frey, J.K. and zndersen, M.C. (2012). Not-so-

splendid isolation: modeling climate-mediated range collapse of a montane mammal

Ochotona princeps across numerous ecoregions. *Ecography* 35(9), 780-791.

Chazdon, R.L., Harvey, C.A., Komar, O., Griffith, D.M., Ferguson, B.G., Martínez-Ramos, M.,

Morales, H., Nigh, R., Soto-Pinto, L., Van Breugal, M. and Philpott, S. M. (2009).

Beyond reserves: A research agenda for conserving biodiversity in human-modified

tropical landscapes. *Biotropica* 41(2), 142-153.

Cianfrani, C., Le Lay, G., Hirzel, A.H. and Loy, A. (2010). Do habitat suitability models

reliably predict the recovery areas of threatened species? *J. Appl. Ecol.* 47: 421-430.

10.1111/j.1365-2664.2010.01781.x

Dai, Y., Peng, G., Wen, C., Zahoor, B., Ma, X., Hacker, C.E., and Xue, Y. (2021). Climate and

land use changes shift the distribution and dispersal of two umbrella species in the Hindu

Kush Himalayan region. *Sci. Total Environ.* 777, 146207.

Delêtre, M., Soengas, B., Jai Vidaurre, P., Isela Meneses, R., Delgado Vásquez, O., Oré Balbín,

I., Santayana, M., Heider, B. and Sørensen, M. (2017). Ecotypic differentiation under

farmers' selection: molecular insight into the domestication of *Pachyrhizus* Rich. ex DC.

(Fabaceae) in the Peruvian Andes. *Evol. Appl.* 10(5): 498-513.

Di Fiore, A. (2004). Diet and feeding ecology of woolly monkeys in a western Amazonian

rainforest. *Int. J. Primatol.* 25: 767-801.

Dong, X., Chu, X., Huang, Q., Zhang, J. and Bai, W. 2019. Suitable habitat prediction

Sichuan snub-nosed monlkeys (*Rhinopithecus roxellana*) and its implications for

conservation in Baihe Nature Reserve, Sichuan, China. *Environ. Sci. and Pollut. Res.* 26:

32374-32384. 10.1007/s11356-019-06369-3

Dormann, C.F. *et al*. (2013). Collinearity: A review of methods to deal with it and a simulation

study evaluating their performance. *Ecography* 36: 027–046.

Ewers, R.M. and Didham, R.K. (2006). Confounding factors in the detection of species

responses to habitat fragmentation. *Bio. Rev.* 81(1), 117-142.

Hein, S., Binzenhoefer, B., Poethke, H.J., Biedermann, R., Settele, J. and Schroeder, B. (2007).

The generality of habitat suitability models: A practical test with two insect groups. *Basic*

Appl. Ecol. 8(4), 310-320.

Fack, V., Shanee, S., Vercauteren Drubbel, R., Del Viento, M., Meunier, H. and Vercauteren, M.

(2018). Observation of snake (Colubridae) predation by yellow-tailed woolly monkeys

(Lagothrix flavicauda) at El Toro study site, Peru. *Neotrop. Primates* 24 (2):79-81

Fick, S.E. and Hijmans, R.J. (2017). Worldclim 2: New 1-km spatial resolution climate surfaces

for global land areas. *Int. J. Climatol.* 37(12): 4302-4315.

Fielding, A.H. and Bell, J.F. (1997). A Review of Methods for the Assessment of Prediction

Errors in Conservation Presence/Absence Models. *Environ. Conserv.* 24: 38-49.

Fisher, D.O. (2011). Trajectories from extinction: where are missing mammals rediscovered?

Glob. Ecol. Biogeogr. 20: 415-425.

Gallice, G.R., Larrea-Gallegos, G and Vázquez-Rowe, I. (2019). The threat of road expansion in

the Peruvian Amazon. *Onyx* 53(2): 284–292.

GIZ (2016). Cambio de uso actual de la tierra en la Amazonía peruana: Avances e

implementación en el marco de la Ley Forestal y de Fauna Silvestre 29763. Cooperación

Alemana, Lima, Perú

Graves, G.R. and O'Neill, J.P (1980). Notes on the yellow-tailed woolly monkey (Lagothrix

flavicauda) of Peru. *J. Mammal.* 61(2):345-357

Guisan, A., Thuiller, W. and Zimmermann, N.E. (2017). *Habitat Suitability and Distribution*

Models with Applications in R. Cambridge University Press, Cambridge, UK.

Hansen, A.J., Neilson, R.P., Dale, V.H., Flather, C.H., Iverson, L.R., Currie, D.J., Shafer, S.,

Cook, R. and Bartlein, P.J. (2001). Global change in forests: responses of species,

communities, and biomes: interactions between climate change and land use are projected

to cause large shifts in biodiversity. *BioScience 51*(9): 765-779.

Hansen, M. C. *et al*. (2013). High-Resolution Global Maps of 21st Century Forest Cover Change.

Science 342: 850–53. Website: http://earthenginepartners.appspot.com/science-2013-

global-forest. Downloaded February 2020.

Hijmans, R.J., Phillips, S., Leathwick, J. and Elith, J. (2017). R Package 'dismo': Species

Distribution Modeling. Website: http://rspatial.org/sdm/. Accessed: February 2020.

Hijmans, R.J. *et al*. (2020). Package 'raster': Geographic data analysis and modeling. Website:

https://cran.r-project.org/web/packages/raster/raster.pdf. Accessed: April 2020.

Hirzel, A.H., Le Lay, G., Helfer, V., Randin, C., and Guisan, A. (2006). Evaluating the ability of

habitat suitability models to predict species presences. *Ecol. Modell.* 199(2), 142-152.

Horwich, R.H., Lyon, J., Bose, A. and Jones, C.B. (2012). Preserving biodiversity and

ecosystems: catalyzing contagion. In *Deforestation Around the World*: 283–318.

Moutinho, P. (Ed.). Rijeka: InTech.

Humanitarian OpenStreetMap Team. (2020). United Nations Office for the Coordination of

Humanitarian Affairs. Website: https://data.humdata.org/dataset/hotosm_per_buildings.

Downloaded February 20, 2020.

Hurvich, C.M. and Tsai, C. (1993). A corrected Akaike information criterion for vector

592   autoregressive model selection. *J. Time Ser. Anal.* 14(3): 271-279.

593 Kattan, G.H. and Valenzuela, L.A. (2013). Phenology, abundance and consumers of figs (*Ficus*

594   spp.) in a tropical cloud forest: evaluation of a potential keystone resource. *J. Trop. Ecol.*

595   29(5): 401-407.

596 *Lagothrix flavicauda* (Humboldt, 1812) in GBIF Secretariat (2021). GBIF Backbone Taxonomy.

597   Checklist dataset https://doi.org/10.15468/dl.xmcds9 accessed via GBIF.org on 2021-09-

598   04.

599 LaRue, M.A. and Nielsen, C.K. 2008. Modelling potential dispersal corridors for cougars in

600   midwestern North America using least-cost path methods. *Ecol. Modell.* 212: 372-381.

601 Laurance, W. F. (2018). Conservation and the global infrastructure tsunami: Disclose, debate,

602   delay!. *Trends Ecol. Evol.* 33(8), 568-571.

603 Lavado Casimiro, W.S., Labat, D., Guyot, J.L. and Ardoin-Bardin, S. (2011). Assessment of

604   climate change impacts on the hydrology of the Peruvian Amazon-Andes basin. *Hydrol.*

605   *Process.* 25: 3721-3734.

606 Leo Luna, M. (1980). First field study of the yellow-tailed woolly monkey. *Oryx* 15: 386-389.

607 Levi, T., Shepard, G.H., Ohl-Schacherer, J., Wilmer, C. C., Peres, C. A. and Yu, D. W. (2011).

608   Spatial tools for modeling the sustainability of subsistence hunting in tropical forests. *Ecol.*

609   *Appl.* 21(5): 1802-1818.

610 Liu, J., Fitzgerald, M., Liao, H., Luo, Y., Jin, Y., Li, X., Yang, X., Hirata, S. and Matsuzawa, T.

611   (2019). Modeling habitat suitability for Yunnan snub-nosed monkeys in Laojun Mountain

612   National Park. *Primates* 10.1007/s10329-01900776-3

613 McHugh, S.M., F.M. Cornejo, J. McKibben, M.Zarate, T.Carlos, C.F. Jiménez and

C.A. Schmitt. (2020). First detection and updated geographic range of the Peruvian yellow-

tailed woolly monkey (*Lagothrix flavicauda*) in the region Junín, Peru. *Oryx* 54(6): 814-

818. DOI: 0.1017/S003060531900084X.

Mittermeier R.A, de Macedo-Ruiz, H, Luscombe, A (1975). A woolly monkey rediscovered in

Peru. *Oryx* 13 (1):41-46

Newbold, T. *et al.* (2014). A global model of the response of tropical and sub-tropical forest

biodiversity to anthropogenic pressures. *Proc. R. Soc. B: Biol. Sci.* 281(1792), 20141371

NOAA (National Oceanic and Atmospheric Administration). 2020. Climate Prediction Center.

Website:https://www.cpc.ncep.noaa.gov/products/assessments/assess_97/samer.html.

Accessed: June 2020.

Olah, G., Smith, A.L., Asner, G.P., Brightsmith, D.J., Heinsohn, R.G. and Peakall, R. (2017).

Exploring dispersal barriers using landscape genetic resistance modelling in scarlet

macaws of the Peruvian Amazon. *Landsc. Ecol.* 32: 445-456.

Oliveira, P.J.C., Asner, G.P., Knapp, D.E., Almeyda, A., Galván-Gildemeister, R., Keene, S.,

Raybin, R.F., Smith, R.C. (2007). Land-Use Allocation Protects the Peruvian Amazon.

Science 317 (5842):1233. doi:10.1126/science.1146324

Parker, T.A and Barkley, L.J (1981). New locality for the yellow-tailed woolly monkey. *Oryx* 16

(1):71-71

Pásztor, L., Barabás, G. and Meszéna, G. (2020). Competitive Exclusion and Evolution:

Convergence Almost Never Produces Ecologically Equivalent Species: (A Comment on

McPeek,"Limiting Similarity? The Ecological Dynamics of Natural Selection among

Resources and Consumers Caused by Both Apparent and Resource Competition"). *Am.*

Nat. 195(4), E112-E117.

Patterson, B.D. and López Wong, C. (2014) Mamíferos/Mammals. In *Peru: Cordillera Escalera.*

Rapid Biological Inventories Report 26: 154-167. Pitman, N *et al.* (Ed.). Chicago: The

Field Museum.

Phillips, S., Anderson, R.P. and Schapire, R.E. (2006). Maximum entropy modeling of species

geographic distributions. *Ecol. Modell.* 190(3-4): 231-259.

Phillips, S.J., Dudík, M., Elith, J., Graham, C. H., Lehmann, A., Leathwick, J., Ferrier, S. (2009).

Sample selection bias and presence-only distribution models: implications for background

and pseudo-absence data. *Ecol. Appl.* 19: 181–197.

Programa Bosques (2015) Bosque - no bosque y pérdida de bosque Húmedo Amazónico 2000-

2014. Programa Nacional de Conservación de Bosques para la Mitigación del Cambio

Climatico, Lima, Peru.

R Core Team. (2018). R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. Website: https://www.Rproject.org/.

Ramirez-Villegas, J., Cuesta, F., Devenish, C., Peralvo, M., Jarvis, A. and Arnillas, C.A. (2014).

Using species distribution models for designing conservation strategies of Tropical Andean

biodiversity under climate change. *J. Nat. Conserv.* 22: 391-404.

Robillard, C.M., Coristine, L.E., Soares, R.N. and Kerr, J.T. (2015). Facilitating climate-change-

induced range shifts across continental land-use barriers. *Conserv. Biol.* 29(6), 1586-

1595.

Rodriguez, E., Morris, C.S., Belz, J.E., Chapin, E.C., Martin, J.M., Daffer, W., and Hensley, S.

(2005). An assessment of the SRTM topographic products, Technical Report JPL D-31639.

Jet Propulsion Laboratory, pp 143. Pasadena, California. Website:

http://srtm.csi.cgiar.org/srtmdata/. Downloaded February 2020.

Sales, L.P., Ribeiro, B.R., Pires, M.M., Chapman, C.A. and Loyola, R. (2019). Recalculating

route: dispersal constraints will drive the redistribution of Amazon primates in the

Anthropocene. *Ecography* 42: 1789-1801.

Serrano-Villavicencio, J.E., Shanee, S. and Pacheco, V. (2021). *Lagothrix flavicauda* (Primates:

Atelidae). *Mamm. Species* 53(1010) 134-144. DOI: 10.1093/mspecies/seab013

Shanee, N., Shanee, S. and Maldonado, A.M. (2007). Conservation assessment and planning for

the yellow-tailed woolly monkey (*Oreonax flavicauda*) in Peru. *Wildlife Biol.* 3: 73-82.

Shanee, S. and Shanee, N. (2009). A new conservation NGO, neotropical primate conservation:

project experiences in Perú. *Int. NGO J.* 4(7): 329-332.

Shanee, S. (2011). Distribution survey and threat assessment of the yellow-tailed woolly

monkey (*Oreonax flavicauda*; Humboldt 1812), northeastern Peru. *Int. J. Primatol.* 32:

691-707.

Shanee, S. and Shanee N. (2011a). Population density estimates for the critically endangered

yellow-tailed woolly monkey (*Lagothrix flavicauda*) at La Esperanza, northeastern Peru.

Int. J. Primatol. 32: 691-180.

Shanee, S. and Shanee, N. (2011b). Activity budget and behavioural patterns of free-ranging

yellow-tailed woolly monkeys Oreonax flavicauda (Mammalia: Primates), at La

Esperanza, northeastern Peru. *Contrib. Zool.* 80:269–277.

Shanee, N. (2012) Trends in local wildlife hunting, trade and control in the Tropical Andes

Hotspot, Northeastern Peru. Endangered Species Research 19 (2):177-186

Shanee, S. (2014). Ranging behaviour, daily path lengths, diet and habitat use of yellow tailed

woolly monkeys (Oreonax flavicauda) at La Esperanza, Peru. In *The woolly monkey:*

behaviour, ecology, conservation and systematics: 167–186. Defler, T. R and P. R.

Stevenson (Ed.). New York: Springer Verlag.

Shanee, N., Shanee, S. and Horwich, R.H. (2015). Effectiveness of locally run conservation

initiatives in north-east Peru. *Oryx* 49(2): 239-247.

Shanee, S. and Shanee, N. (2015). Measuring success in a community conservation project:

local population increase in a critically endangered primate, the yellow-tailed woolly

monkey (*Lagothrix flavicauda*) at La Esperanza, northeastern Peru. *Trop. Conserv. Sci.*

8(1): 169-186.

Shanee, S. (2016). Predicting Future Effects of Multiple Drivers of Extinction Risk in Peru's

Endemic Primate Fauna. In *Ethnoprimatology, Developments in Primatology, Progress*

and Prospects: 315–349. Waller, M. T. (Ed.), Switzerland: Springer International

Publishing.

Shanee, N. and Shanee, S. (2016). Land trafficking, migration and conservation in the "No Man's

Land" of northeastern Peru. *Trop. Conserv. Sci.* 9(4): 1-16.

Shanee, N., Mendoza, A.P. and Shanee, S. (2017). Diagnostic overview of the illegal trade in

primates and law enforcement in Peru. *Am. J. Primatol.* 79 (11): e22516.

doi:10.1002/ajp.22516

Shanee, S., Shanee, N., Monteferri, B., Allgas, N., Pardo, A.A. and Horwich, R.H. (2017).

Protected area coverage of threatened vertebrates and ecoregions in Peru: Comparison of

communal, private and state reserves. *J. Environ. Manage.* 202, 12–20.

Shanee, S. Allgas, N. and Shanee, N. (2018). Community conservation as a tool for primate

conservation in Peru. In *Primatology, Biocultural Diversity and Sustainable*

Development in Tropical Forests: 320-329. Rommens, D. and J. Pulido Mata (Ed.).

Mexico: UNESCO.

Shanee, S., Shanee, N., Lock, W. and Espejo-Uribe, M.J. (2020). The development and growth of

non-governmental conservation in Peru: Privately and communally protected areas.

Hum. Ecol. 48: 681-693.

Shanee, S., Cornejo, F.M., Aquino, R., Mittermeier, R.A. & Vermeer, J. (2021). *Lagothrix*

flavicauda (amended version of 2019 assessment). *The IUCN Red List of Threatened*

Species 2021: e.T39924A192307818. https://dx.doi.org/10.2305/IUCN.UK.2021-

1.RLTS.T39924A192307818.en. Accessed on 31 January 2022.

Solórzano-García, B., Zubillaga, D., Piñero, D. and Vázquez-Domínguez, E. (2021).

Conservation implications of living in forest remnants: inbreeding and genetic structure

of the northernmost mantled howler monkeys. *Biotropica* 53(4), 1163-1177.

Strier, K.N. (1992). Atelinae adaptations: behavioral strategies and ecological constraints. *Am. J.*

Phys. Anthropol. 88: 515-524.

UNEP-WCMC. 2020. Protected Area Profile for Peru from the World Database of Protected

Areas, June 2020. Available at: www.protectedplanet.net

Vermote, E. (2019). NOAA CDR Program. NOAA Climate Data Record (CDR) of AVHRR

Normalized Difference Vegetation Index (NDVI), Version 5. Vegetation Health Product.

NOAA   National   Centers   for   Environmental   Information.

https://doi.org/10.7289/V5ZG6QH9. Downloaded May 20, 2020.

Vuille, M., Bradley, R.S., Werner, M. and Keimig, F. (2003). 20th century climate change in the

Tropical Andes: observations and model results. In *Variability and Change in High*

Elevation Regions: Past, Present & Future. Advances in Global Change Research, vol 15:

75-99. Diaz, H. F. (Ed.). Dordrecht: Springer.

Xiao, H., McDonald-Madden, E., Sabbadin, R., Peyrard, N., Dee, L. E. and Chadès, I. (2019).

The value of understanding feedbacks from ecosystem functions to species for managing

ecosystems. *Nat. Comm.* 10(1), 1-10.

Young, K.R. (1996). Threats to biological diversity caused by coca/cocaine deforestation in

Peru. *Environ. Conserv.* 23 (1):7-15. doi:doi:10.1017/S0376892900038200

Zalamea, P., Munoz, F., Stevenson, P.R., Paine, C.E.T., Sarmiento, C., Sabatier, D. and Heuret,

P. (2011). Continental-scale patterns of *Cecropia* reproductive phenology: evidence from

herbarium specimens. *Proceedings of the Royal Society B* 278: 2437-2445.