Root-specific activation of plasma membrane H$^+$-ATPase 1 enhances plant growth and shoot

accumulation of nutrient elements under nutrient-poor conditions in *Arabidopsis thaliana*

Kota Monden[a], Takehiro Kamiya[b], Daisuke Sugiura[c], Takamasa Suzuki[d], Tsuyoshi Nakagawa[a] and

Takushi Hachiya[a†]

[a]Department of Molecular and Functional Genomics, Interdisciplinary Center for Science Research,

Shimane University, Matsue 690-8504, Japan

[b]Graduate School of Agricultural and Life Sciences, The University of Tokyo, Tokyo 113-8657, Japan

[c]Graduate School of Bioagricultural Sciences, Nagoya University, Nagoya 464-8601, Japan

[d]College of Bioscience and Biotechnology, Chubu University, Kasugai 487-8501, Japan

[†]To whom correspondence should be addressed.

E-mail: takushi.hachiya@life.shimane-u.ac.jp

Tel.: +81-852-32-6288

Fax: +81-852-32-6109

Abbreviations: GO, Gene Ontology; NUE, Nutrient uptake efficiency; PCA, Principal component

analysis; PM, plasma membrane; RPM, Reads per million mapped reads; RT, Reverse transcription;

SA, Salicylic acid; SAR, Systemic acquired resistance; WT, Wild-type

Abstract

Plasma membrane (PM) H^+-ATPase contributes to nutrient uptake and stomatal opening by creating proton gradient across the membrane. Previous studies report that a dominant mutation in the *OPEN STOMATA2* locus (*OST2-2D*) constitutively activates *Arabidopsis* PM H^+-ATPase 1 (AHA1), which enlarges proton motive force for root nutrient uptake. However, the stomatal opening is also constitutively enhanced in the *ost2-2D*, causing the drought hypersensitivity. To develop a plant whose nutrient uptake is improved with normal stomatal movement, we generated grafted plants (scion/rootstock : Col-0 (WT)/*ost2-2D*), and compared their growth and ionome profile with those of control plants (WT/WT) under nutrient-rich or nutrient-poor conditions. WT/*ost2-2D* shoots had larger fresh weight, dry weight, rosette diameter, leaf blade area, and contents of C, N, K, Ca, S, P, Mg, Na, Mn, B, Co, and Mo under nutrient-poor conditions compared with WT/WT shoots. The root fresh and dry weights and primary root length were greater in WT/*ost2-2D* roots than in WT/WT roots under both nutrient conditions. These results suggest that root-specific activation of PM H^+-ATPase enhances root nutrient uptake, which accelerates the plant growth under nutrient-poor condition. Our study presents a novel approach to improving nutrient uptake efficiency in crops and vegetables; thus, it could contribute to the implementation of low-input sustainable agriculture.

Keywords: *Arabidopsis thaliana*; micrografting; nutrient uptake efficiency; OPEN STOMATA 2; plasma membrane H^+-ATPase

1. Introduction

As the global population increases, crop production must also increase. Nutrient availability is an important determinant of crop productivity; thus, fertilizers are often applied in large amounts to improve crop production. Intensive fertilizer use, however, puts a heavy burden on the environment, e.g., through groundwater pollution, eutrophication, and greenhouse gas emissions [1]. N, P, and K, macronutrients typically in high demand in plants, account for a major portion of applied fertilizers. The artificial synthesis of N fertilizers requires massive fossil fuel consumption; i.e., ammonia synthesis via the Harber–Bosch process consumes 1%–2% of the global energy budget currently [2]. Additionally, the P contained in fertilizers is derived from nonrenewable rock phosphate that will be depleted in the future [3]. Thus, developing plants with high nutrient uptake efficiency (NUE) is crucial to achieving sustainable agriculture.

It is widely accepted that plasma membrane (PM) H^+-ATPase provides proton motive force for root nutrient uptake mediated by specific transporters and channels [4]. Constitutive overexpression of *Oryza sativa* PM H^+-ATPase 1 (*OSA1*) under the control of the *CaMV-35S* promoter was shown to enhance root uptake of ammonium, plant growth, and productivity in both the field and laboratory [5]. Conversely, similar overexpression of PM H^+-ATPase genes failed to confer any positive effects on *Arabidopsis* and tobacco plants [6, 7]. Importantly, post-translational activation of PM H^+-ATPase often has major effects on *Arabidopsis* physiological responses, including stomatal opening and hypocotyl elongation [8, 9]. Therefore, with the aim of improving the NUE of *Arabidopsis*, we focused on the *Arabidopsis OPEN STOMATA 2* (*OST2*)-dominant mutant (*ost2-2D*), which expresses a constitutively active allele of the *Arabidopsis* PM H^+-ATPase 1 (*AHA1*) [9]. This mutant exhibits lower apoplastic pH and longer-sized root epidermal cells, corresponding to enhancements in proton motive force and acid growth in the roots, respectively [10]. However, PM hyperpolarization in the guard cells of *ost2-2D* causes drought hypersensitivity due to the prevention of stomatal closure, even with the addition of abscisic acid [9, 11, 12]. Thus, the root-specific activation of OST2/AHA1 could be an ideal design for improving NUE. To achieve this design, we grafted Col-0 (wild-type; WT) scions onto rootstock originating from WT or *ost2-2D* and analyzed

the vegetative growth, nutrient element content, and transcriptomes of the grafted plants grown under

nutrient-rich or nutrient-poor conditions. We show that *ost2-2D*-derived rootstock enhances the

growth of shoots and roots as well as the shoot accumulation of several nutrient elements under

nutrient-poor conditions.

2. Materials and methods

2.1. Plant materials and growth conditions

Seeds of Col-0 (WT) and *ost2-2D* [9] were surface sterilized and sown in a solid medium containing

50 mL of half-strength Murashige and Skoog (1/2-MS) salts including vitamins (Code M0222;

Duchefa Biochemie, Haarlem, The Netherlands) supplemented with 0.05% (w/v) MES-KOH (pH 5.7),

1% (w/v) sucrose, and 0.5% (w/v) gellan gum (Fujifilm Wako, Osaka, Japan). The seeds were placed

in the dark at 4°C for 3 days to break dormancy. Seedlings were grown vertically under a PPFD of

25 µmol m^{-2} s^{-1} (16/8-h light/dark cycle) at 22°C. Micrografting was performed aseptically using 5-

day-old seedlings [13], which were cut perpendicularly in the hypocotyl with an injection needle tip

(Code NN-2613S; TERUMO, Tokyo, Japan). The obtained scion was connected with the partner

rootstock in a silicon microtube (Code 1-8194-04; AS ONE, Osaka, Japan). The grafted plants were

incubated for 5 days under a PPDF of 50 µmol m^{-2} s^{-1} (constant light) at 27°C and further cultivated

for a few days under a PPFD of 25 µmol m^{-2} s^{-1} (16/8-h light/dark cycle) at 22°C. Successfully grafted

plants without adventitious root formation were selected for similar size; transferred to vermiculite

pots, soil pots, or solid media; and grown for use in the following experiments. For vermiculite

cultivation, the grafted plants were grown under a PPFD of 100–120 µmol m^{-2} s^{-1} (16/8-h light/dark

cycle) at 22°C–23°C. Before the cultivation, ~150 mL of vermiculite was soaked with 80 mL of 1/2-

MS or 1/50-MS salts without vitamins (Code M0221; Duchefa Biochemie) supplemented with 0.05%

(w/v) MES-KOH (pH 5.7). Four days after the onset of cultivation, 30 mL of the same compositions

of 1/2-MS or 1/50-MS salts was supplied to the vermiculite. The plants were then cultivated for an

additional 3 days; after which, they were sampled for experiments. For soil cultivation, the grafted

plants were grown for 7 days under a PPFD of 100–120 µmol m^{-2} s^{-1} (16/8-h light/dark cycle) at

22°C–23°C in the pots containing ~150 mL of nutrient-rich soil (Supermix A; Sakata Seed, Kyoto,

Japan). For an *in vitro* culture on solid media, the plants were grown for 3 days (vertically) or 7 days

(horizontally) under a PPFD of 100–120 µmol m^{-2} s^{-1} (16/8-h light/dark cycle) at 22°C–23°C in

30 mL (horizontally) or 50 mL (vertically) of 1/2-MS or 1/50-MS salts without vitamins (Duchefa

Biochemie) supplemented with 0.05% (w/v) MES-KOH (pH 5.7), 1% (w/v) sucrose, and 0.8% (w/v)

purified agar (Code 01162-15; Nacalai Tesque, Kyoto, Japan). To allow nondestructive sampling of

the roots, the surfaces of solid media were covered with cellophane sheets [14]. All other details are

described in the figure legends.

2.2. Growth analysis of plants under vermiculite and soil cultivation

Shoots were weighed on a precision balance (HR-202i; A&D, Tokyo, Japan) and scanned at a

resolution of 600 dpi (GT-X980; EPSON, Tokyo, Japan) to measure rosette diameter. Leaf blades

were harvested from the shoots and scanned at a resolution of 600 dpi to measure the area of the leaf

blades. The shoots were then oven-dried at 80°C for 2 days and recovered in a desiccator with silica

gel. Shoot dry weights were measured using the precision balance, and the rosette diameter and leaf

blade area were measured using ImageJ version 1.52p.

2.3. Growth analysis of plants under in vitro *culture on solid media*

The shoots and roots of horizontally grown plants were weighed using the precision balance, after

which they were oven-dried at 80°C for 2 days and recovered in a desiccator with silica gel. The dry

weights of the shoots and roots were measured using the precision balance. The roots of vertically

grown plants were scanned at a resolution of 600 dpi to measure the length and number of primary

and lateral roots (>0.3 mm). Root length was measured using ImageJ version 1.52p.

2.4. Determination of nutrient elements

Harvested shoots were oven-dried at 80°C for 2 days, and the dried samples were ground using a spoon

and recovered in a desiccator with silica gel. For determination of C and N, the dried powder was

weighed and encapsulated in a tin boat, and C and N concentrations were measured using a CN

analyzer (Vario EL III; Elementar Analysensysteme GmbH, Hanau, Germany). For determination of

K, P, S, Ca, Fe, Mg, Mn, Na, B, Co, Cu, Mo, and Zn, the dried samples were weighed and digested

using 2 mL of concentrated HNO_3 followed by the addition of 1 mL of H_2O_2 at 120°C until it

evaporated. The digested samples were dissolved in 0.08 M HNO_3 and the element concentrations

were measured using inductively coupled plasma–mass spectrometry (Agilent 7800 ICP-MS; Agilent

Technologies, Santa Clara, CA, USA) using 2 ppb indium as an internal standard.

2.5. Extraction of total RNA

Roots were harvested, immediately frozen using liquid N_2, and stored at −80°C until use. Frozen

samples were ground with a Multi-Beads Shocker (Yasui Kikai, Osaka, Japan) using zirconia beads

(diameter: 5 mm). Total RNA was extracted using an RNeasy Plant Mini Kit (Qiagen, Tokyo, Japan)

according to the manufacturer's instructions. For RNA-Seq library preparation, RNA was purified

using on-column DNase digestion (Qiagen).

2.6. RT-qPCR

Reverse transcription (RT) was performed using ReverTra Ace qPCR RT Master Mix with gDNA

Remover (Toyobo, Osaka Japan) according to the manufacturer's instructions. The synthesized cDNA

was diluted 10-fold with distilled water and used in quantitative PCR (qPCR). RT-qPCR was

performed on a QuantStudio 1 (Thermo Fisher Scientific) with KOD SYBR qPCR Mix (Toyobo)

according to the manufacturer's instructions. Transcript levels were quantified using a relative standard

curve with *ACTIN3* as the internal standard [19]. Primer sequences are shown in Supplementary Table

1.

2.7. RNA-Seq analysis

RNA quality was evaluated using a Qubit RNA IQ Assay Kit with a Qubit 4 Fluorometer (Thermo

Fisher Scientific, Tokyo, Japan). The RNA samples for which RNA IQs were 9.1–10.0 were used for

the following preparation. cDNA libraries were constructed using a NEBNext Ultra II RNA Library

Prep with Sample Purification Beads (Code E7775; New England Biolabs, Tokyo, Japan), a NEBNext

Poly(A) mRNA Magnetic Isolation Module (Code E7490; New England Biolabs), and a NEBNext

Multiplex Oligos for Illumina (Index Primers Set 3) (Code E7710; New England Biolabs) according

to the manufacturer's instructions. These cDNA libraries were sequenced using NextSeq 500 (Illumina,

Tokyo, Japan), and the produced bcl files were converted to fastq files using bcl2fastq (Illumina). The

obtained reads were analyzed according to the method described by Notaguchi et al. [15] and mapped

to the *Arabidopsis* reference (TAIR10) using Bowtie [16] with the following options: "--all --best --

strata." The number of reads mapped to each reference was counted, and the total read counts of splice

variants from a gene were used for further analysis via the iDEP ver. 095 integrated web application

[17] with the default settings. Pathway analysis was performed using the Metascape web application

[18]. Information about read counts is provided in Supplementary Table 2.

2.8. Statistical analysis

Unpaired two-tailed Welch's *t*-tests were conducted using R v.2.15.3.

3. Results

3.1. Confirmation of root-specific missense mutation of OST2/AHA1

To manipulate the activity of OST2/AHA1 in a root-specific manner, we grafted WT scions onto

rootstock originating from WT or *ost2-2D* (Supplementary Fig. 1A). To confirm the missense

mutation (G867S) of *OST2/AHA1* reported previously [9], we extracted RNA from the shoots and

roots of grafted plants (scion/rootstock: WT/WT and WT/*ost2-2D*). RT-PCR was performed using

primers specific to *OST2/AHA1*; after which, the PCR products were sequenced directly. The missense

mutation was detected only in the cDNA derived from the WT/*ost2-2D* roots (Supplementary Fig. 1B).

3.2. Root-specific activation of OST2/AHA1 enhances shoot growth under nutrient-poor conditions

Grafted plants were transferred to vermiculite pots to analyze the effects of root-specific activation of

OST2/AHA1 on shoot growth. Following cultivation for 7 days under nutrient-rich (1/2-MS salts) or

nutrient-poor conditions (1/50-MS salts), shoot growth was analyzed. WT/*ost2-2D* plants exhibited

increased shoot growth relative to that of WT/WT plants under 1/50-MS but not 1/2-MS conditions

(Fig. 1A). Under 1/50-MS conditions, shoot fresh and dry weights were significantly increased by

28% and 31%, respectively, in WT/*ost2-2D* plants than in WT/WT plants (Fig. 1A). Additionally,

rosette diameter and leaf blade area were significantly increased by 13% and 29%, respectively, in

WT/*ost2-2D* plants than in WT/WT plants under 1/50-MS conditions (Fig. 1A). Under 1/2-MS

conditions, however, only the shoot dry weight was significantly increased by 11% in WT/*ost2-2D*

plants (Fig. 1A). When the grafted plants were grown in pots containing nutrient-rich soil or in solid

media containing 1/2-MS or 1/50-MS, the shoot fresh and dry weights, rosette diameter, and leaf blade

area in WT/*ost2-2D* plants were slightly, albeit not always significantly, increased compared with

those in WT/WT plants (Supplementary Fig. 2A, B). Relative water contents of shoots did not

substantially decrease in WT/*ost2-2D* plants compared with those in WT/WT plants regardless of

nutritional availability and cultivation style (Supplementary Fig. 3A–C). Thus, root-specific activation

of OST2/AHA1 does not cause shoot desiccation due to excessive stomatal opening.

3.3. Root-specific activation of OST2/AHA1 enhances root growth irrespective of nutrient availability

Barbez et al. [10] reported that the epidermal cell length of the primary roots of *ost2-2D* seedlings is

significantly larger than that of WT plants, suggesting that root growth is altered because of root-

specific activation of OST2/AHA1. To confirm this, the root growth of grafted plants grown vertically

on solid media containing 1/2-MS or 1/50-MS salts was analyzed. The appearance of the roots in

WT/WT plants differed according to nutrient condition, and their fresh and dry weights and lateral

root length were generally increased under 1/50-MS conditions than under 1/2-MS conditions (Fig.

1B and Supplementary Fig. 4A). In WT/*ost2-2D* plants relative to in WT/WT plants, root fresh weight

and root dry weight were significantly increased by 30% and 41% under 1/2-MS conditions,

respectively, and by 28% and 23% under 1/50-MS conditions, respectively (Fig. 1B). The primary

root length was significantly increased in WT/*ost2-2D* plants by 22% and 32% under 1/2-MS and

1/50-MS conditions, respectively; however, no significant difference was observed in the lateral root

length of the grafted plants (Fig. 1B). In grafted plants grown horizontally on solid media, root fresh

and dry weights were increased in WT/*ost2-2D* plants than in WT/WT plants irrespective of nutrient

availability (Supplementary Fig. 4B).

3.4. Root-specific activation of OST2/AHA1 increases the shoot content of nutrient elements

Zhang et al. [5] reported that constitutive overexpression of *OSA1* significantly increases the content

of C, N, K, P, and Ca in the leaves of rice grown in nutrient-rich hydroponic solution. Thus, we

determined the content of C and the 14 nutrient elements contained in MS salts in the shoots of grafted

plants grown in vermiculite. The content of C, N, K, Ca, S, P, Mg, Na, Mn, B, Co, and Mo was

significantly increased in WT/*ost2-2D* shoots than in WT/WT shoots under 1/50-MS conditions (Fig.

2). In particular, the content of macronutrients N, K, Ca, S, and P was consistently increased by 54%,

37%, 24%, 31%, and 44%, respectively, in WT/*ost2-2D* shoots than in WT/WT shoots (Fig. 2). Under

1/2-MS conditions, however, only Na content was significantly increased in the WT/*ost2-2D* shoots

than in the WT/WT shoots, although several elements showed increasing trends in the WT/*ost2-2D*

shoots (Fig. 2). The shoot concentrations of nutrient elements were generally comparable between the

grafted plants regardless of nutrient availability (Supplementary Fig. 5).

3.5. Root-specific activation of OST2/AHA1 alters root expression of genes related to biotic responses

To explore the effects of root-specific activation of *OST2/AHA1* on the root transcriptome, we

performed RNA-Seq analysis on the roots of grafted plants grown horizontally on solid media

containing 1/2-MS or 1/50-MS salts. Hierarchical clustering and principal component analysis (PCA)

classified the RNA-Seq data into four groups according to genotype and nutrient condition

(Supplementary Fig. 6A, B). The k-means clustering analysis revealed two gene clusters for which the

expression was induced (cluster 2) or repressed (cluster 3) in WT/*ost2-2D* roots than in WT/WT roots

(Fig. 3A, B and Supplementary Table 2). Each cluster was characterized using pathway analysis;

cluster 2 was enriched in Gene Ontology (GO) terms associated with systemic acquired resistance

(SAR) and responses to fungal infection, whereas cluster 3 was enriched in the GO term "lipid X

metabolic process" (Fig. 3C). The expression of multiple genes encoding UDP-3-*O*-acyl N-

acetylglycosamine deacetylase family proteins LPXC1–5, which contribute to the biosynthesis of lipid

X, a precursor of lipid A-like molecules [20], was consistently repressed in WT/*ost2-2D* roots

(Supplementary Fig. 7A, B), although the physiological roles played by endogenous lipid A-like

molecules are unknown.

4. Discussion

The present study demonstrates that root-specific activation of OST2/AHA1 enhances shoot growth

and increases the shoot content of all the macronutrient elements and six micronutrient elements in *A.*

thaliana cultivated in nutrient-poor vermiculite (Figs. 1A and 2). The root-specific activation of

OST2/AHA1 also increased root weight and primary root length in plants grown in solid media (Fig.

1B and Supplementary Fig. 4), corresponding to root cell expansion due to apoplastic acidification,

i.e., acid growth [10]. Given that the availability of nutrient elements limits plant vegetative growth

under nutrient-poor conditions, the improvement of proton motive force and root growth could

facilitate nutrient uptake and shoot growth. In the present study, the concentrations of N, K, P, Mg, B,

and Mo in WT/WT shoots were decreased by 19%–53% when nutrient availability was reduced

(Supplementary Fig. 5), suggesting that the supply of these elements could limit shoot growth under

nutrient-poor conditions. Because the shoot content of N, K, P, Mg, B, and Mo was consistently

increased in WT/*ost2-2D* plants than in WT/WT plants (Fig. 2), root-specific activation of

OST2/AHA1 may enhance shoot growth by increasing root uptake of these elements. Interestingly,

root expression of the high-affinity nitrate uptake transporter *NRT2.1* and iron uptake transporter *IRT1*

was higher in WT/*ost2-2D* plants than in WT/WT plants under nutrient-poor conditions

(Supplementary Fig. 8A, B). IRT1 has broad specificity for divalent heavy metals other than Fe

including Mn, Zn, Co, and Cd under Fe-deficient conditions [21]. Therefore, up-regulation of *NRT2.1*

and *IRT1* might have contributed to increases in the shoot content of N, Mn, and Co under nutrient-

poor conditions (Fig. 2).

Our transcriptome analysis suggested that root-specific activation of OST2/AHA1 up-

regulates genes related to SAR and fungal infection responses in the roots (Fig. 3C). This corresponds

to an observation made previously that salicylic acid (SA) levels and expression of SA-responsive

marker genes are elevated in the leaves of *ost2-1D* mutants [9], because endogenous SA acts as a

signal that mediates local and systemic defense responses to pathogen infection [22]. Interestingly,

lowering the pH in external media *per se* is known to induce expression of genes responsive to

pathogens, pathogen elicitors, and SA in *A. thaliana* roots [23]. Therefore, root-specific activation of

OST2/AHA1 may up-regulate expression of these genes through apoplastic acidification [10].

In conclusion, we succeeded in increasing plant vegetative growth under nutrient-poor conditions via

root-specific activation of PM H^+-ATPase using a grafting strategy. Genome editing technologies are

applicable for substituting specific DNA sequences for a gene of interest in various plant species;

therefore, it may be possible to design and produce rootstocks with activated PM H^+-ATPase. Overall,

our study presents a novel approach to improving NUE in crops and vegetables; thus, it could

contribute to the implementation of low-input sustainable agriculture.

Acknowledgments

Seeds of *ost2-2D* were kindly provided by Dr. Jeffrey Leung (CNRS, France). We thank Maki Saiki

for ICP-MS analysis. The authors would like to thank Enago (www.enago.jp) for the English language

review.

Funding

This study was supported by a KAKENHI Grant from Japan Society for the Promotion of Science

[Grant-in-Aid for Scientific Research (C) No. JP20K05771 to TH] and the Ichimura Foundation for

New Technology (to TH).

Conflict of interest

288 The authors declare that this study was conducted in the absence of any commercial relationships that

289 could lead to any potential conflicts of interest.

References

[1]   S. Savci, An Agricultural Pollutant: Chemical Fertilizer, Int. J. Environ. Sci. Dev. 3 (2012) 77-
80. https://doi.org/10.7763/IJESD.2012.V3.191.

[2]   V. Kyriakou, I. Garagounis, A. Vourros, et al., An Electrochemical Haber-Bosch Process,
Joule 277 (2020) 142-158. https://doi.org/10.1016/j.joule.2019.10.006.

[3]   J. Cooper, R. Lombardi, D. Boardman, et al., The future distribution and production of global
phosphate rock reserves, Resour. Conserv. Recycl. 57 (2011) 78–86.
https://doi.org/10.1016/j.resconrec.2011.09.009.

[4]   M. G. Palmgren, Plant plasma membrane H^+-ATPases: powerhouses for nutrient uptake, Annu.
Rev.Plant Biol. 52 (2001) 817–845. https://doi.org/10.1146/annurev.arplant.52.1.817.

[5]   M. Zhang, Y. Wang, X. Chen, et al., Plasma membrane H^+-ATPase overexpression increases
rice yield via simultaneous enhancement of nutrient uptake and photosynthesis, Nat. Commun.
12 (2021) 735. https://doi.org/10.1038/s41467-021-20964-4.

[6]   M. Hashimoto-Sugimoto, T. Higaki, T. Yaeno, et al., A Munc13-like protein in *Arabidopsis*
mediates H^+-ATPase translocation that is essential for stomatal responses, Nat. Commun. 4
(2013) 2215. https://doi.org/10.1038/ncomms3215.

[7]   F. Gévaudant, G. Duby, E. von Stedingk, et al., Expression of a Constitutively Activated
Plasma Membrane H^+-ATPase Alters Plant Development and Increases Salt Tolerance, Plant
Physiol. 144 (2007) 1763–1776. https://doi.org/10.1104/pp.107.103762.

[8]   J. Falhof, J. T. Pedersen, A. T. Fuglsang, et al., Plasma Membrane H^+-ATPase Regulation in
the Center of Plant Physiology, Mol. Plant 9 (2016) 323–337.

https://doi.org/10.1016/j.molp.2015.11.002.

[9]   S. Merlot, N. Leonhardt, F. Fenzi, et al., Constitutive activation of a plasma membrane H^+-

ATPase prevents abscisic acid-mediated stomatal closure, EMBO J. 26 (2007) 3216–3226.

https://doi.org/10.1038/sj.emboj.7601750.

[10]   E. Barbez, K. Dünser, A. Gaidora, et al., Auxin steers root cell expansion via apoplastic pH

regulation in *Arabidopsis thaliana*, Proc. Natl. Acad. Sci. USA. 114 (2017) E4884–E4893.

https://doi.org/10.1073/pnas.1613499114.

[11]   S. D'Alessandro, Y. Mizokami, B. Légeret, et al., The Apocarotenoid β-Cyclocitric Acid

Elicits Drought Tolerance in Plants, iScience 19 (2019) 461–473.

https://doi.org/10.1016/j.isci.2019.08.003.

[12]   J. H. Wong, M. Klejchová, S. A. Snipes, et al., SAUR proteins and PP2C.D phosphatases

regulate H^+-ATPases and K^+ channels to control stomatal movements, Plant Physiol. 185

(2021) 256–273. https://doi.org/10.1093/plphys/kiaa023.

[13]   K. Monden, M. Kojima, Y. Takebayashi, et al., Root-specific Reduction of Cytokinin

Perception Enhances Shoot Growth in *Arabidopsis thaliana*, Plant Cell Physiol. (2022) 256–

273. https://doi.org/10.1093/pcp/pcac013.

[14]   T. Hachiya, T. Oya, K. Monden, et al., A cellophane-supported *Arabidopsis* culture for

seamless transfer between different media is useful for studying various nitrogen responses,

Soil Sci. Plant Nutr. 67 (2021) 277–282. https://doi.org/10.1080/00380768.2021.1908094.

[15]   M. Notaguchi, T. Higashiyama, T. Suzuki, Identification of mRNAs that move over long

distances using an RNA-Seq analysis of *Arabidopsis/Nicotiana* benthamiana heterografts,

Plant Cell Physiol. 56 (2014) 311–321. https://doi.org/10.1093/pcp/pcu210.

[16]   B. Langmead, C. Trapnell, M. Pop, et al., Ultrafast and memory-efficient alignment of short

DNA sequences to the human genome, Genome Biol. 10 (2009) R25.

https://doi.org/10.1186/gb-2009-10-3-r25.

[17]   S. X. Ge, E. W. Son, R. Yao, iDEP: an integrated web application for differential expression

and pathway analysis of RNA-Seq data, BMC Bioinform. 19 (2018) 534.

https://doi.org/10.1186/s12859-018-2486-6.

[18]   Y. Zhou, B. Zhou, L. Pache, et al., Metascape provides a biologist-oriented resource for the

analysis of systems-level datasets, Nat. Commun. 10 (2019) 1523.

https://doi.org/10.1038/s41467-019-09234-6.

[19]   T. Hachiya, J. Inaba, M. Wakazaki, et al., Excessive ammonium assimilation by plastidic

glutamine synthetase causes ammonium toxicity in *Arabidopsis thaliana*, Nat. Commun. 12

(2021) 4944. https://doi.org/10.1038/s41467-021-25238-7.

[20] C. Li, Z. Guan, D. Liu, et al., Pathway for lipid A biosynthesis in *Arabidopsis thaliana*

resembling that of *Escherichia coli*, Proc. Natl. Acad. Sci. USA. 108 (2011) 11387–11392.

https://doi.org/10.1073/pnas.1108840108.

[21] A. Jogawat, B. Yadav, Chhaya, et al., Metal transporters in organelles and their roles in heavy

metaltransportation and sequestration mechanisms in plants, Physiol. Plant 173 (2021) 259–

275. https://doi.org/10.1111/ppl.13370.

[22] D. F. Klessig, H. W. Choi, D. A. Dempsey, et al., Systemic Acquired Resistance and Salicylic

Acid:Past, Present, and Future, Mol. Plant Microbe Interact. 31 (2018) 871–888.

https://doi.org/10.1094/MPMI-03-18-0067-CR.

[23] I. D. A. Lager, O. Andréasson, T. L. Dunbar, et al., Changes in external pH rapidly alter plant

gene expression and modulate auxin and elicitor responses, Plant Cell Environ. 33

(2010)1513–1528. https://doi.org/10.1111/j.1365-3040.2010.02161.x.

Figure legends

Fig. 1. Effects of root-specific activation of OST2/AHA1 on the growth of shoots and roots. (A)

Shoot appearance, shoot fresh and dry weights, rosette diameter, and leaf blade area of the grafted

plants. Grafted plants were transferred to vermiculite pots and grown for 7 days with a supply of 1/2-

MS or 1/50-MS salts. All data pooled from six independent grafting experiments are presented as

means ± SD (n = 43–44). (B) Root fresh and dry weights and primary and lateral root lengths of

grafted plants. Grafted plants were transferred to 1/2-MS or 1/50-MS media and grown vertically for

3 days. All data pooled from four independent grafting experiments are presented as means ± SD

(n = 30). WT/WT (W/W) and WT/*ost2-2D* (W/*o*) denote plants corresponding to the WT scion grafted

on the WT rootstock and the WT scion grafted on the *ost2-2D* rootstock, respectively. *$P < 0.05$;

***$P < 0.001$; NS, not significant at $P > 0.05$ (unpaired two-tailed Welch's *t*-test). Scale bar: 0.5 cm.

Fig. 2. Effects of root-specific activation of OST2/AHA1 on the shoot content of nutrient

elements. The shoot content of 15 nutrient elements in the grafted plants, which were transferred to

vermiculite pots and grown for 7 days with a supply of 1/2-MS or 1/50-MS salts. Shoot samples

harvested from three (for C and N determination) and five (for determination of the other elements)

independent grafting experiments were subjected to element analysis. Shoots from 4 to 11 plants in

each experiment were pooled as one biological replicate. Data are presented as means ± SD (n = 3 or

5). WT/WT (W/W) and WT/*ost2-2D* (W/*o*) denote the plants corresponding to the WT scion grafted

on the WT rootstock and the WT scion grafted on the *ost2-2D* rootstock, respectively. *$P < 0.05$;

**$P < 0.01$; NS[†] $P < 0.1$; NS, not significant at $P > 0.05$ (unpaired two-tailed Welch's *t*-test).

Fig. 3. Effects of root-specific activation of OST2/AHA1 on the root transcriptome. (A) Heat map

from k-means clustering, (B) violin plots of normalized transcript levels in clusters 2 and 3, and (C)

pathway analyses of the genes in clusters 2 and 3. Grafted plants were transferred to 1/2-MS or 1/50-

MS media and grown horizontally for 7 days. Root samples harvested from two independent grafting

experiments were subjected to RNA-Seq analysis. Roots from four plants in each experiment were

pooled as one biological replicate. Genes for which the standard deviation was ranked in the top 2000

were used for k-means clustering with four clusters, which included 769, 288, 219, and 724 genes

(clusters 1–4), respectively. Purple and green represent higher and lower expressions, respectively.

Information about genes included in each cluster is provided in Supplementary Table 3. Pathway

analyses were performed using the obtained clusters 2 and 3, and significantly enriched terms are

shown. WT/WT (W/W) and WT/*ost2-2D* (W/*o*) denote plants corresponding to the WT scion grafted

on the WT rootstock and the WT scion grafted on the *ost2-2D* rootstock, respectively.

Figure 1

A

1/2-MS   1/50-MS

WT/WT   WT/*ost2-2D*   WT/WT   WT/*ost2-2D*

Shoot fresh weight

NS   ***

mg

Shoot dry weight

*   ***

mg

Rosette diameter

NS   ***

cm

W/W W/o   W/W W/o
1/2-MS   1/50-MS

Leaf blade area

NS   ***

cm²

W/W W/o   W/W W/o
1/2-MS   1/50-MS

B

Root fresh weight

***   ***

mg

Root dry weight

***   ***

mg

Primary root length

***   ***

cm

W/W W/o   W/W W/o
1/2-MS   1/50-MS

Lateral root length

NS   NS

cm

W/W W/o   W/W W/o
1/2-MS   1/50-MS

Figure 2

Figure 3