Environmental DNA reveals invasive crayfish microbial associates and ecosystem-wide biodiversity before and after eradication

Running Title: eDNA reveals crayfish microbial associates

Kimberly M. Ballare[a], Anna Worth[a], Tyler Goodearly[b,c], Dannise V. Ruiz-Ramos[d], Eric Beraut[e], Hailey Nava[a], Colin Fairbairn[a], Robert K. Wayne[e], Beth Shapiro[a,f], Ginny Short[b], Rachel S. Meyer[a]

[a] University of California Santa Cruz, Dept. of Ecology and Evolutionary Biology, Santa Cruz, CA 95064, USA
[b] Center for Natural Lands Management, Temecula, CA 92590, USA
[c] Current Affiliation: University of Hawai i at Mānoa, Dept. of Human Nutrition, Food, and Animal Sciences, Honolulu, HI, 96822, USA
[d] University of California Merced, Dept. of Life and Environmental Sciences, Merced, CA 95343, USA
[e] University of California Los Angeles, Dept. of Ecology and Evolutionary Biology, Los Angeles, CA 90095, USA
[f] Howard Hughes Medical Institute, University of California Santa Cruz, Santa Cruz, CA 95064, USA

Author Contributions: K.M.B., G.S., T.G., and R.S.M. conceived and designed experiments and

sampling scheme. G.S. coordinated site access and supervised field work. T.G. maintained and sampled

aquaria. K.M.B., G.S., H.N., D.V.R-R., T.G., and R.S.M collected field samples and coordinated

volunteer field sampling. K.M.B., A.W., D.V.R-R, E.B., H.N., and C.F. conducted molecular laboratory

work. K.M.B., A.W., H.N., C.F, and R.S.M. conducted literature review. K.M.B. and R.S.M. curated,

analyzed, and interpreted data. K.M.B. and T.G. generated the map figure and K.M.B. generated the

data figures. R.K.W., B.S., G.S., and R.S.M. provided supervision, funding, materials, and analysis tools.

K.M.B. and R.S.M. wrote the manuscript with input from all co-authors.

Corresponding Author: Rachel Meyer, rameyer@ucsc.edu, 1-206-351-7997

Keywords: restoration; holobiome; Procambarus clarkii; bioindicators; eDNA; conservation

management

Abstract

Biodiversity monitoring in conservation projects is essential to understand environmental status

and recovery. However, traditional field surveys can be expensive, time-consuming, biased

towards visual detection, and focused on measuring a limited set of taxa. Environmental DNA

(eDNA) methods provide a new approach to biodiversity monitoring that has the potential to

sample a taxonomically broader set of organisms with similar effort, but many of these

approaches are still in the early stages of development and testing. Here, we use multilocus

eDNA metabarcoding to understand how the removal of invasive red swamp crayfish impacts

local biodiversity of a desert oasis ecosystem, as well as to detect crayfish both directly and

indirectly. We tracked crayfish DNA signatures, microbial DNA associated with crayfish, and

biodiversity of plant, fungal, animal, and bacterial communities through time. We were

unsuccessful in detecting crayfish directly in either control tanks or oases using targeted

metabarcoding primers for invertebrates and eukaryotes, similar to previous studies which have

shown variable levels of success in detecting crayfish from environmental samples. However,

we were successful in discerning a suite of 90 crayfish-associated taxa to serve as candidate

bioindicators of invasive presence using 16S and Fungal ITS2 metabarcoding. Ranking these

90 taxa by their geographic distribution in eDNA surveys and by evidence of crayfish-

associations in the literature, we support 9 taxa to be high-ranking, and suggest they be

prioritized in future biomonitoring. Biodiversity analyses from five metabarcode loci including

plants, animals, and both prokaryotic and eukaryotic microbes showed that communities

differed but that species richness remained relatively similar between oases through time. Our

results reveal that, while there are limitations of eDNA approaches to detect crayfish and other

invasive species, microbial bioindicators offer a largely untapped biomonitoring opportunity for

invasive species management, adding a valuable resource to a conservation manager's toolkit.

Keywords: restoration; holobiome; Procambarus clarkii; bioindicators; eDNA; conservation
63
64   management
65

1. Introduction

1. Introduction

Environmental DNA (eDNA) is an increasingly popular tool for biodiversity monitoring across

landscapes and taxa. Using eDNA, reliable biodiversity surveys can be quickly and easily

obtained by land managers and community scientists without need for expertise in field methods

or taxonomy (Meyer et al., 2021; Suarez-Menendez et al., 2020). These approaches are

particularly promising for invasive species monitoring and conservation projects (Ficetola et al.,

2008; Sepulveda et al., 2020) because they make it possible for managers to simultaneously

monitor biodiversity and detect invasive species using a single technique. Biomonitoring during

restoration and conservation management projects is also an essential part of understanding

whether restoration interventions have been successful, e.g. whether taxa of interest are

present in an ecosystem. Both traditional and eDNA surveys can be useful for direct invasive

species detection and monitoring, as long as the taxon in question can be reliably detected (e.g.

Crane et al., 2021; Halstead et al., 2017; Ratsch et al., 2020) However, some taxa are

challenging to detect and quantify using either method even when they are known to be present:

including cryptic or elusive species, species persisting at low densities, or with variable

detection during their life cycles.

One set of taxa that can be challenging to detect with either traditional or molecular approaches

are crayfish (Astocoidea L.). When abundant, crayfish are easily trapped or detected with visual

surveys (e.g. Short, 2021), but crayfish traps also risk mortality of non-target organisms

(Tréguier et al., 2014). When rare, crayfish can escape detection through visual surveys or

trapping because they can burrow in sediment and find refugia for up to several months

(Pritchard et al., 2021), potentially leading to false negative detection in early stages of invasion,

or false successes of eradication programs for invasive species (e.g. Nunes et al., 2017).

Molecular detection approaches also have variable levels of success (Chucholl et al., 2021;

Dougherty et al., 2016; Harper et al., 2018; Ikeda et al., 2016; Troth et al., 2020). For the

globally invasive red swamp crayfish (*Procambarus clarkii* Giard, 1852) in particular, targeted

detection assays from environmental samples have not proven to be a stable monitoring

approach (Cai et al., 2017; Mauvisseau et al., 2018; Thalinger et al., 2021). For example, in a

2014 study, *P. clarkii* was not reliably detected from environmental samples even when using

species-specific primers tested and designed for the study (Tréguier et al., 2014). Additionally,

some assays fail when researchers modify protocols such as changing filters, extraction

volumes, and substrates (Geerts et al., 2018; Tréguier et al., 2014), or when attempting to

detect crayfish at different points in their life cycle (Dunn et al., 2017). One potential explanation

is that crayfish do not shed DNA reliably into the environment (Curtis and Larson, 2020), which

is also a known issue with detecting other crustacean and invertebrate species in marine or

aquatic ecosystems (e.g. green crab, Crane et al., 2021).

To mitigate problems with direct detection of a particular species, an emerging eDNA method is

to use microbial proxies as bioindicators of sentinel species for tracking invasion and other

disturbances (Astudillo-García et al., 2019). The characterization of microbial communities can

also provide vital information about the biology of the target species (Ficetola et al., 2019), as

well as characterize the community ecology of the site. Microbial proxies used for indirect

detection can include both infectious agents (Wittwer et al., 2018), as well as more benign

associates such as gut microbes, diet species, and symbionts (Ficetola et al., 2019; Skelton et

al., 2017). While the use of eDNA metabarcoding is appealing because it can detect many

species simultaneously and be used on a variety of substrates (Jarman et al., 2018), few

studies use metabarcoding profiles to generate bioindicator proxies for target species detection.

It may be difficult to detect "true" microbial bioindicators, that is, a taxon may appear driven by a

target species, but in fact that microbe is common in many environments and has broad

ecological associations and roles. However, because *P. clarkii* and other crayfish have multiple

known symbionts, pathogens, and other microbial associates verified across many studies

(Chen et al., 2021; Dragičević et al., 2021), this makes it an ideal taxon to test detection

success through microbial proxies. Because of the commonness of invasive red swamp crayfish

across the globe (Geiger et al., 2005), the outsized negative impacts that they have on local

biodiversity and ecosystem function (Oficialdegui et al., 2020; Souty-Grosset et al., 2016), and

the known detection challenges, multiple methods of detection are clearly needed.

Here, we explore how environmental DNA can be used to track the invasive red swamp

crayfish, as well as reveal novel components of the biology of the invaded ecosystem. We

tested multiple existing published assays for direct detection of *P. clarkii*, and used universal

metabarcoding to 1) detect signals of crayfish presence, 2) generate a reliable list of microbial

crayfish bioindicators, and 3) track the biotic effects of crayfish presence and pesticide

eradication methods on the overall ecological community. Because the red swamp crayfish has

such a large known effect on aquatic biodiversity, consuming living plants, detritus, and animals

as an opportunistic omnivore (Alcorlo et al., 2004; Bucciarelli et al., 2019), we expect to see

discernable correlations with the surrounding biotic community including changes in community

composition and species richness over time.

2. Methods

2.1 Study System

We conducted our study in two palm oases in the Thousand Palms Oasis Preserve (Thousand

Palms Oasis and Simone Pond, Figure 1) managed by the Center for Natural Lands

Management (CNLM), as well as in an experimental tank system (see section 2.3 below).

Located in the Colorado Desert, California, USA, the Thousand Palms Oasis Preserve contains

numerous distinct oases fed by a common underground aquifer, with the vegetative community

dominated by the native California Fan Palm (*Washingtonia filifera)*. Simone Pond is distinct

from other oases in the preserve as it is characterized by an open habitat and more consistent

water depth throughout the year. Thousand Palms Oasis is located approximately one km south

of Simone Pond. Surrounded by the same dominant vegetation community, it differs from

Simone Pond in that it has less light that permeates through the fan palm canopy, and a shallow

water depth. Other than Western mosquitofish *Gambusia affinis* which is an invasive species

that persists in both oases, Thousand Palms Oasis has no fish species. There is no record of

crayfish existing in Thousand Palms Oasis.

Simone Pond has had multiple invasive species introductions, including red swamp crayfish

(*Procambarus clarkii*) in the 1950's, and tilapia (*Oreochromis* sp.) in 2015, both of which caused

significant biodiversity loss in Simone Pond (Short, 2021). Importantly, Simone Pond has

unique geophysical characteristics including an open habitat that allows for light permeation,

high water quality, and submergent and emergent vegetation, which make it a candidate

location for the introduction of the rare and endangered desert pupfish (*Cyprinodon macularius*

Baird & Girard 1853, Martin and Saiki, 2005; Varela-Romero et al., 2002). Unfortunately,

previous attempts to introduce desert pupfish to Simone Pond were unsuccessful due to

competition with and predation by red swamp crayfish. CNLM Staff used several methods to try

to remove the invasive species and found cypermethrin (Cyper TC®, Control Solutions Inc.,

EPA Reg. No. 53883–92 [(±)α-cyano-(3-phenoxyphenyl)methyl(±) - cis/trans-3-(2,2-

dichloroethenyl)- 2,2-dimethylcyclopropanecarboxylate]) to be the most effective at controlling

crayfish (Short, 2021).

Cypermethrin is an insecticide that kills target organisms by blocking voltage-gated sodium

channels to affect the peripheral nervous system (Shafer et al., 2005). Derived from the

insecticidal compounds found in chrysanthemum, cypermethrin is known be more toxic to

crustaceans than vertebrate fish species (Morolli et al., 2006; Sandodden and Johnsen, 2010;

Soares et al., 2019), with mild toxicity present in amphibians and mammals (Edwards et al.,

1986; Khan et al., 2009). Cypermethrin and related compounds have been shown to be

effective at eradicating invasive crayfish populations at low doses (e.g. Sandodden and

Johnsen, 2010), although it is also toxic to other aquatic organisms including benthic

invertebrates (Zhang et al., 2018). Prior to applying the pesticide, preserve staff drained water to

a temporary reservoir, and manually filtered all visible tilapia and crayfish. After the initial

cypermethrin application to the drained Simone Pond, they allowed Simone Pond to refill

naturally. Pesticide application in Simone Pond began in July 2019 and continued monthly until

July 2020 (Pesticide research authorization permit #1905023, California Department of

Pesticide Regulation). During and after the application of the pesticide CNLM staff conducted

regular monitoring including trapping and visual surveys, which revealed a substantial

population of crayfish in Simone Pond. Prior to pesticide application, over 6,300 crayfish were

removed via baited minnow traps between January and June 2019, with an estimated 26,000

individuals present in January 2019 (Leslie Depletion Model; Leslie and Davis, 1939). No

crayfish or tilapia were observed for 100 days after initial application of the pesticide, but one

dead crayfish was found 101 days after the initial application. As of May 2022, no others have

been observed since October 2019 (Short, 2021).

2.2 Environmental Sample Collection

We used environmental DNA sampling to detect crayfish and monitor changes in biodiversity in

Simone Pond and Thousand Palms Oasis before, during, and after the cypermethrin application

program. Specifically, we collected environmental samples at four time points (Table 1,

Supplemental Table S1): 1) June 2019, immediately before cypermethrin application began; 2)

December 2019, after 5 monthly applications of cypermethrin; 3) June 2020, after the last

monthly cypermethrin application; and 4) Dec 2020, 5 months after the last application of

cypermethrin. We collected sediment (fully water submerged to rarely water submerged

surfaces of the oasis, containing soil usually mixed with algal and siliceous matter) and soil

(terrestrial surface, never water submerged) samples during each sampling event, and water

samples during the June 2019 sampling event. Water samples were collected using a

randomized transect approach in each oasis. One liter of water was collected in an IV fluid bag

and passively drip filtered through a 0.22 um Sterivex filter (Millipore, Burlington,

Massachusetts, USA). Filters were immediately refrigerated, and DNA was extracted the

following day. Sediment or soil samples were collected at regular spatial intervals around the

water edge of the oasis, with sampling locations selected so as to maximize environmental

heterogeneity by taking samples from sites that were rarely, frequently, or fully submerged, from

under palm trees, in aquatic grass beds, in dense algal masses, and along oasis perimeter

fencing. At each site, we collected a sample of three 2ml cryotubes filled with soil or sediment.

We obtained these substrates by scraping off the surface with the tube itself (soil) or by

obtaining an underwater surface sediment sample using a 2-inch diameter PVC pipe and

scraping sediment off of it with the tube itself. Water and substrate samples were immediately

frozen at -20˚C and transferred to -80˚C for archiving. Immediately before DNA extraction, we

thawed soils and sediments on ice, homogenized them by vortexing, and pooled an aliquot from

each tube together to represent the single site sample. This mixture was again homogenized

and 0.25g was taken for DNA extraction.

2.3. Aquarium eDNA Sampling

In June 2019, we established thirteen four-gallon aquarium tanks using unfiltered well water

from the preserve source aquifer. The aquaria contained four treatments: four crayfish-only

tanks (three containing live crayfish, and one with a crayfish carcass), three tilapia-only tanks,

three crayfish and tilapia tanks, and three no-animal (blank) tanks. Crayfish were not fed for the

duration of their time in the tanks to avoid introducing a foreign food microbiome. After two

weeks, we collected a single one-liter water sample from each tank and processed samples for

DNA extraction as described in section 2.2 above. Non-invasive sampling of these tanks was

compliant with NRC guidelines of animal treatment, and was an approved component of CNLM

Project G2023 (Short, 2021).

2.4. DNA Extraction, Library Preparation, and Sequencing

<u>2.4.1 Metabarcoding</u>

We extracted soil and sediment samples using a standard 0.25g input into the Qiagen DNeasy

PowerSoil Kit (Qiagen Inc., Germantown, MD) and followed manufacturer's instructions. We

extracted water DNA from the Sterivex filter cartridge using the Qiagen DNeasy Blood and

Tissue Kit following the protocol reported in Spens et al. (2017). From soil, sediment, and water

DNA extracts, we amplified 5 metabarcode markers to detect different taxonomic groups: 16S

targeting Bacteria and Archaea (515F 806R; Caporaso et al., 2012), 18S targeting Eukaryota

(Euk_1391f and EukBr; Amaral-Zettler et al., 2009), fungal ITS1 targeting Fungi (abbreviated

hereafter as "FITS", ITS5 and 5.8S; White et al., 1990) and plant ITS2, targeting plants and

green algae (abbreviated hereafter as 'PITS', ITS-S2F and ITS-S3R; Gu et al., 2013), and CO1,

initially targeting the full region (LCO1490 and HCO219; Hebert et al., 2003) to search for

crayfish DNA in the June 2019 sample. After preliminary sequence data indicated high

proteobacteria swamping in the initial CO1 samples, we switched to the partial CO1 primers for

subsequent samples targeting both invertebrates and protists (mlCOlintF and Fol-degen-rev;

Leray et al., 2013).

For each marker, we performed three replicate PCRs to account for stochastic amplification. We

then pooled PCR replicates by marker, quantified the amount of DNA in each pool using the

Qubit^{TM} dsDNA Broad Range assay (Invitrogen^{TM}), and finally pooled all reactions per sample of

five markers together using equimolar concentrations of each reaction. We indexed the

amplicons in each sample pool using dual unique Nextera indices (UMI), then combined all

metabarcoding libraries for sequencing on either an Illumina MiSeq with the v3 kit for 2x300bp

reads (18S, CO1, PITS (all sampling points), and 16S and FITS for the June 2019 samples) or a

NextSeq 550 for 2x150bp reads (for 16S and FITS in Dec 2019, June 2020, and Dec 2020).

Sequencing was conducted at UCLA or UCSC sequencing facilities to produce fastq files of

DNA libraries.

To provide Preserve staff with preliminary results after each sampling event we processed and

sequenced each of the four sample batches separately. To test for batch effects, we randomly

selected 10 samples from each sample date (five samples per oasis), re-extracted DNA,

amplified two markers (16S and 18S), and sequenced them together on the NextSeq as

described above. We co-analyzed these with the original samples to detect whether community

composition of replicates fell within their respective batch groups in ordinations and

permutational ANOVA.

We analyzed metabarcoding sequence data using the *Anacapa Toolkit* (Curd et al., 2019),

which uses CutAdapt (Martin, 2011) and FastX-Toolkit (Gordon and Hannon, 2010) for Fastq

quality control and filtering, and DADA2 (Callahan et al., 2016) for amplicon sequence variant

(ASV) assignment and chimera detection and removal. Taxonomic assignment for each ASV

was made using Bowtie2 (Langmead and Salzberg, 2012) and the Bayesian Lowest Common

Ancestor (BLCA; Gao et al., 2017) algorithm with our metabarcode-specific CRUX reference

databases (version 2; made from October 2019 NCBI data; doi:10.5068/D1QX0W), of which the

18S and CO1 databases included sequences from *P. clarkii*. We assumed the default

parameters except the minimum quality score was set to 32. Taxonomic assignments at each

classification level were retained when the bootstrap confidence cutoff score was over 70 for

16S and FITS or over 80 for CO1, 18S, and PITS. The ASVs with the same resulting taxonomic

entry from domain to species were summed to be a single entry as the species-equivalent. We

converted metadata and ASV tables (Supplemental Table S1) to *Phyloseq* (McMurdie and

Holmes, 2012) objects for analysis using *Ranacapa* (Kandlikar et al., 2018). We filtered data for

analysis first by decontaminating tables with the decontam package (Davis et al., 2018) and

prevalence set to 0.1, and then removing any taxa represented by fewer than 10 total reads

across all samples.

2.4.2 Metagenome

After cypermethrin crayfish control began in Simone Pond, hundreds to thousands of killed

crayfish carcasses degraded in Simone Pond, and pieces of the exoskeletons were occasionally

found intact. In December 2019, we found a piece of crayfish carapace exoskeleton in the

Simone Pond surface sediment. To examine the metagenome signatures on this exoskeleton

and test for any retained crayfish DNA in decomposing carcasses, we washed the sample with

pure water and ground 0.25g with a Qiagen PowerLyzer Homogenizer and extracted DNA. A

Tn5 library was prepared according to Russell et al. (2020) and sequenced using an Illumina

NextSeq 550 (Illumina, San Diego, California). We used bbduk from BBTools (Bushnell, 2014)

to trim low quality (q<25) reads and remove adapters. Then, we merged reads with bbmerge

and mapped them to the *P. clarkii* genome (ASM2042438v2) using BWA (v 0.7.17-r1188; Li

and Durbin, 2009). Low entropy mapped reads were removed using bbduk with entropy=0.9

entropywindow=25 entropyk=10 entropymask=f. Merged reads were matched to taxa through

MG-RAST (Meyer et al., 2008).

2.4.3 Quantitative PCR

In an attempt to have more sensitive direct detection of crayfish than metabarcoding, we also

used published quantitative PCR (qPCR) assays to amplify *P. clarkii* from samples where

crayfish were present. We selected two primer sets that reported reliable detection of red

swamp crayfish from freshwater DNA extracts: Mauvisseau et al. (2018) [CO1-Pc-03-F 5'-

GGAGTTGGAACAGGATGGACT-3', CO1-Pc-03-R 5'-AATCTACAGATGCTCCCGCA-3']; and

Knudsen et al. (2019) [Procla_co1_F04 5'-GCGGGAGCATCTGTAGATTT-3', Procla_co1_R04

5'-ATAGCTCCTGCCAACACAGG-3']. The templates we screened included known crayfish

positive and known crayfish negative samples from the June 2019 DNA extracts: three sediment

samples and two water samples from Simone Pond (which contained an estimated 26,000

crayfish), one water sample and two sediment samples from Thousand Palms Oasis (which had

no crayfish), and eight water samples from the tank experiments (5 samples from crayfish

positive and 3 samples from crayfish negative tanks). Reactions were performed in triplicate and

included an 18S reaction (same 18S primer as used for metabarcoding) as a positive control.

Melt curve temperatures were also recorded. The published assays used different annealing

temperatures: 60 C° (Knudsen et al. 2019) and 59 C° (Mauvisseau et al. 2018). Using iTaq

Universal SYBR Green Supermix (BioRad, Hercules, California, USA), we ran these assays

twice, once at each annealing temperature, and found results were similar, so we report the

assay at 59 C°. We note we did not use FAM probes as in original publications for these assays,

as the probes were solely designed to screen out other related crayfish in multi-species

systems; we had no crayfish present in our system other than *P. clarkii*.

2.5 Statistical Analysis

2.5.1 Bioindicator Analyses

We calculated microbial bioindicators of crayfish presence using community similarity

percentage analysis (SIMPER; Clarke, 1993) in the R package *vegan* (Oksanen et al., 2015).

SIMPER performs pairwise comparisons of species within each sample category and finds the

average contributions of each species to overall Bray-Curtis dissimilarity, as well as the

likelihood of each species occurring in one site type over another. From the June 2019 (pre-

eradication) sample batch, we grouped together all tank and oasis samples (sediment, soil, and

water) containing crayfish, and grouped together all tank and oasis samples without crayfish.

We calculated the contribution of each taxon to the Bray-Curtis dissimilarity between groups

containing crayfish and those without. We performed this twice: both the log (x+1) sequence

read abundance and presence-absence data of sample sets rarefied to 3000 reads with 1000

permutations. To mitigate the tendency of SIMPER to preferentially detect more variable taxa as

having between-group effects (Warton et al., 2012), we only considered taxa as significant

drivers of community difference if they were significant in both the presence-absence and log

(x+1) read abundance transformations (as per Ballare et al., 2019). We repeated the SIMPER

analysis for each time point post-eradication.

We considered the taxa that were found to be significant drivers of community differences in the

June 2019 SIMPER analyses to be 'candidate bioindicators' of crayfish presence. We vetted

these candidate bioindicator taxa in multiple ways. 1) We scored geographic distribution by

cross referencing each taxon within 1000 eDNA samples collected across California that used

the same assay (CALeDNA publicly available data, ucedna.com; Meyer et al., 2021).

Bioindicators that were present in most of these samples were given a low score because they

are clearly not likely to be crayfish-specific based on distribution. Bioindicators with regional or

restricted distribution, such as only in watersheds, or restricted to the oases or Coachella Valley

were given high scores. 2) We scored bioindicators by information on their ecological roles and

any reported associations with crayfish species. To review these bioindicator roles, we

conducted a literature review using GLOBI API Plus (Poelen et al., 2014), and also reviewed

literature in Google Scholar searches for the bioindicator taxon at the species, genus, or family

level and 'crayfish'. From these, we developed a composite score to and assigned each taxon a

bioindicator strength weight from 0-6 (0 having no evidence based on geography or role that

there is a relationship with crayfish, and 6 being tightly related to both crayfish presence based

on role and specificity in our ecosystem), using the following scoring logic: 1 point for being

found in under 200 samples out of the 1000 CALeDNA samples, 1 point for geographic range

more specific than all of California (e.g. southern California or Coachella Valley), 2 points if most

occurrences are in Simone Pond, and 2 points for having a published direct interaction with

crayfish (any genus). If there was no direct crayfish interaction found in the manual literature

review or GLOBI search, 1 point was added for co-occurring with crayfish or any interaction with

Arthropoda. All points were taken away if found mostly in Thousand Palms Oasis.

2.5.2 Ecosystem Biodiversity Analyses

We used R packages *Phyloseq* (McMurdie and Holmes, 2012) and *vegan* (Oksanen et al.

2015), for community biodiversity analyses visualized plots using *ggplot2* (Wickham et al.,

2016). For these analyses we removed repeat samples and conducted tests only on sediment

samples to avoid differences in community due to sample type (soil, sediment, or water), as

variable compositions of clay, sand, and silt between soil and sediment samples may cause

differing rates of DNA absorption and retention (Barnes et al., 2014; Cai et al., 2006). Sediment

also tends to be a more reliable substrate to detect biodiversity over time as it retains a much

longer record of DNA than water samples (days to weeks or even months, Turner et al., 2015;

Wei et al., 2018). We examined alpha and beta diversity patterns for each metabarcode dataset

separately (16S, 18S, FITS, PITS, and CO1), comparing biodiversity between oases, both

within and between timepoints. To evaluate the changes in the Arthropoda community that we

suspected was sensitive to cypermethrin treatment, we merged the CO1 and 18S data and

subsetted it to Arthropoda only, and then conducted the same biodiversity analyses to test for

differences in alpha and beta diversity of arthropods in both Simone Pond and Thousand Palms

Oasis over the four time points. We conducted Analysis of Variance (ANOVA) and post-hoc

Tukey tests to test for differences in observed alpha diversity over time, and permutational

ANOVA (PERMANOVA) to test for significant differences in community composition at different

time points in the different oases, using the Jaccard distance metric. Although PERMANOVA is

more robust than other permutation methods (e.g. ANOSIM, Mantel test) to detect differences in

community composition in the presence of uneven dispersion (Anderson and Walsh, 2013), we

also conducted post-hoc beta dispersion tests (PERMDISP2) to test whether significant

community differences detected by PERMANOVA were due to differences in dispersion rather

than true biological difference.

Our analysis to look for batch effects indicated that the repeated samples did not separate from

the other batches in NMDS ordination space, indicating little effect of sequencing batch on

community composition (Supplemental Figure S1). However, PERMANOVA analyses of 18S

and 16S data sets with the repeat samples included indicated some small but significant

community differences between sample processing dates (PERMANOVA: 18S F=2.90,

$p<0.001$; 16S F=10.864, $p<0.001$). To remain conservative, subsequent PERMANOVA for all

markers included the processing batch as a stratum to correct for possible differences in

community due to batch effects only. We visualized overall community similarity between time

points and oases using NMDS ordination.

3. Results

3.1. Direct Detection of Crayfish DNA Signatures

<u>3.1.1. Metabarcoding</u>

We expected to find crayfish DNA signatures in metabarcoding results from COI and 18S loci,

but from the 84 total samples analyzed from July 2019 that included 21 sediment samples from

Simone Pond (Simone Pond), 4 water samples from Simone Pond, and the 13 experiment tanks

of which 7 contained crayfish, we only detected a signal of Decapoda (the order for crayfish) in

18S from 3 samples. While in all cases *Procambarus clarkii* was the most congruent sequence,

it was not assigned to high confidence (ASVs required a bootstrap confidence score above 80

to be included in our final dataset). These positive samples were all from water samples: from

two of the experimental tanks with crayfish and tilapia, and one from Simone Pond. Given that

water retains a shorter temporal record of DNA compared to sediment, and only one natural

oasis sample yielded a positive result even when an estimated ~26,000 crayfish were in the

oasis, we determined metabarcoding for the purpose of direct detection would not be useful for

crayfish monitoring.

3.1.2. Crayfish qPCR

We found that qPCR using both sets of published *CO1* primers for *P. clarkii* were also unstable

at detecting crayfish, producing false positive amplification signals from two negative controls.

While some known crayfish positive samples did show qPCR amplification with both published

primer sets, most failed to amplify crayfish, such as water samples from Simone Pond and the

tank with the crayfish carcass (Supplemental Table S2).

3.1.3. Crayfish DNA in exoskeleton material

Because our qPCR results did not detect crayfish DNA from water containing a rotting carcass,

this led us to ask how much crayfish DNA persists in carcass material. Only 121 out of 602,844

merged reads mapped the carapace reads to the *Procambarus clarkii* genome (0.02%),

indicating severe DNA degradation. MG-RAST metagenome analysis of the carapace reads

revealed over 99% of the reads were microbial and dominated by Proteobacteria (Supplemental

Table S3).

3.2. Bioindicator Analysis

3.2.1. Candidate Crayfish Bioindicator taxa from SIMPER analysis

From the samples collected in July 2019, using 16S and FITS datasets, we detected 90

microbial taxa that were significantly more likely to occur in samples containing crayfish

(crayfish tanks and Simone Pond) than samples where crayfish were known to be absent (blank

tank, tilapia-only tanks, and Thousand Palms Oasis), hereafter referred to as candidate crayfish

bioindicators. Candidate bioindicator taxa were from diverse microbial lineages, originating from

three kingdoms, 43 orders, and 56 families (Figure 2a, Supplemental Table S4). Each individual

bioindicator taxon was found in a minimum of two samples and a minimum of 30 total sequence

reads. After conducting the GLOBI search, literature review, and applying qualitative scoring, 60

of these taxa had a weight above 0 but less than 5 (indicating weak to moderate crayfish

association), and 9 taxa had a weight of 5 or above, indicating tight crayfish association (Table

2, Supplemental Table S4).

Heatmap visualizations of Euclidean distance of composition and frequency of 90 candidate

bioindicator taxa between oases and throughout time indicate that composition of bioindicator

taxa differed between and within oases in both 2019 timepoints and in June 2020 (Figure 2b),

indicating strong community turnover of bioindicator taxa during the study period, especially

between oases when crayfish were present and when cypermethrin was applied. The same

heatmap analysis conducted on 90 random taxa (averaged frequency from 20 permutations)

from the total microbial data set (merged 16S and FITS) showed a substantially different pattern

than the heatmap made with only bioindicator taxa, with fewer community differences between

oases and timepoints (Figure 2c).

3.2.2. SIMPER Analysis of Simone Pond and Thousand Palms Oasis After Crayfish Removal

The SIMPER analyses for the remaining sampling points had fewer taxa return as significant

drivers of beta diversity between the two oases, indicating that communities between the two

oases may be becoming more similar over time post-crayfish eradication. In the December

2019 samples, taken 5 months into the eradication program, we only detected four taxa that

were significantly more likely to occur in Simone Pond as compared to Thousand Palms Oasis.

We found no taxa more significantly likely to occur in Simone Pond in June 2020, and one

significant taxon in Dec 2020 (Table 3). In December 2019, two of the original bioindicator taxa

(crayfish association rank of 0) were significantly more likely to occur in Simone Pond in both

June 2019 and Dec 2019 samples (Table 3, bolded text).

3.2.3. Discovery of candidate bioindicator taxa in crayfish carapace sample

We looked for the presence of candidate bioindicator taxa in the microbial metagenome of the

crayfish carcass (Section 3.1.3) and found strong overlap (Supplemental Table S3). At the

family level, we found 24 overlapping families with the 90 bioindicators, encompassing 32

bioindicator taxa. At the genus level, we found 17 overlapping genera encompassing 18

bioindicator taxa including top ranked bioindicators (rank 5 or 6): *Cyanobium sp, Aspergillus*,

Talaromyces, and *Scenedesmus*. *Shewanella*, a taxon that we did not rank highly in our

qualitative score (rank 2) but that is documented to make up 20% of the *P. clarkii* crayfish gut

microbiome (Shui et al., 2020), was the dominant of the overlapping genera with 20,106 reads

and the third highest in overall relative abundance in the carcass metagenome. These

bioindicators were found at no such level in the sample periods following crayfish removal

(Supplemental Table S1).

3.3. Individual Metabarcode Biodiversity Analysis

3.3.1 Metabarcode Sequencing Summary

We generated 47.0 million MiSeq paired reads and 52.6 million NextSeq paired reads. Negative

controls including field blanks, extraction blanks, and PCR blanks made up 2 million of those

MiSeq reads and 3.7 million of the NextSeq reads (NCBI bioproject, will provide reference after

manuscript acceptance). The *16S* marker targeting Prokaryota recovered 2280 taxa in June

2019 samples from MiSeq sequencing and in subsequent sampling windows of December

2019, June 2020, and December 2020, which used the NextSeq for less costly sequencing, we

recovered 2979, 3899, and 2614 taxa, respectively. 91% of these taxa were Bacteria, and 5%

were Archaea and 4% Eukaryota. In the same sample date order, the *18S* marker targeting

Eukaryota recovered 1668, 1213, 1041, and 1049 taxa and all were assigned to Eukaryota. The

FITS marker targeting Fungi recovered only Eukaryota, and included 636, 605, 967, and 649

taxa. The full *CO1* marker targeting animals and protists used in June 2019 recovered 427 taxa

and contained 76% Bacteria, 0.7% Archaea, and 23% Eukaryota. The partial CO1 marker used

in subsequent sampling amplified 916, 760, and 513 taxa consisting of over 99% Eukaryota and

0.3% Bacteria. The *PITS* marker targeting Chlorophyta (land plants and green algae) recovered

125, 141, 145, and 100 taxa and all were Eukaryota, of which 66% belonged to Streptophyta

and the remainder to Chlorophyta.

3.3.2 Beta Diversity Analyses

For four of the marker datasets (18S - Eukaryotes, FITS - Fungi, 16S - Bacteria and Archaea,

and PITS - Plants), Simone Pond and Thousand Palms Oasis had significantly different

community composition from each other at each sample time point (PERMANOVA p<0.05,

Figure 3a, Supplemental Figure S2a-c, Supplemental Table S5). PERMANOVA analyses for

the CO1 marker set indicated that communities were significantly different at the latter three

sample periods, but not in June 2019 (Figure S2d), which was due to Proteobacteria swamping,

an unintended effect of using the full-length COI primers for the June 2019 dataset.

Pairwise PERMANOVA tests also indicated differences in community composition within oases

between different sample time periods. However, the time periods that significantly differed

within an oasis was variable within and between datasets (Supplemental Table S5). For

instance, Simone Pond 18S eukaryotic communities in Jun 2020 and Dec 2020 were

significantly different, but in Thousand Palms Oasis they were not. The test for differences in

beta dispersion for Simone Pond 18S was significant when comparing Jun 2019 and Dec 2019

samples (F=4.06, p=0.012) indicating that significant differences in these two sampling points

shown by PERMANOVA could be due to differences in dispersion. All other 18S community

comparisons from Simone Pond and Thousand Palms Oasis were not significantly different in

beta dispersion, indicating that PERMANOVA results are accurately reflecting a true difference

in community structure between remaining time points. Full PERMANOVA and Beta Dispersion

Test results for all marker sets are reported in Supplemental Table S5.

3.3.3 Overall Alpha Diversity Analyses

Alpha diversity estimates were more variable between individual marker datasets, with different

taxonomic groups showing different levels of species richness between oases and timepoints.

In general, ANOVA with post-hoc Tukey tests indicated that extrapolated species richness

(Chao1) at individual time points within each dataset did not significantly differ between Oases

(Figure 3b, Supplemental Figure S3). There were two exceptions to these results, where

Bacteria and Archaea (16S) in Simone Pond 16S showed significantly lower species richness

than Thousand Palms Oasis in Jun 2019 (Supplemental Figure S3b, Supplemental Table S6),

and Plants (PITS) in Simone Pond having significantly higher species richness than Thousand

Palms Oasis in Dec 2019 (Supplemental Figure S3c).

When species richness of both Oases was considered together, each marker dataset showed

significant differences between timepoints, although there was not any time point that was

consistently more species-rich. For 18S Eukaryotes, Jun 2019 and Dec 2019 had the highest

species richness for both oases (Figure 3b); for 16S, PITS, and CO1, Dec 2019 showed the

highest levels of species richness (Supplemental Figure S3b-d); and Jun 2020 showed the

highest species richness for FITS (Supplemental Figure S3a).

3.3.4. Arthropod Biodiversity Analyses

We conducted alpha and beta diversity analyses of only Arthropoda taxa by merging 18S

Eukaryotes and CO1 Invertebrates and Protists datasets and removing all non-arthropod taxa.

We observed a significant effect of sample date on arthropod species richness in Simone Pond

(ANOVA, F=13.85, p<0.001). Post-hoc Tukey tests indicate that all samples at time points after

crayfish eradication in Simone Pond had lower observed alpha diversity than the Jun 2019

sample pre-eradication (p<0.001, Figure 2d). In Thousand Palms Oasis, arthropod diversity

was also significantly affected by sample period (ANOVA, F=3.84, p=0.019), but Tukey tests

reveal that arthropod species richness in Thousand Palms Oasis remained more stable

between sample points, with only the June 2020 sample significantly lower than the December

2019 sample (p=0.023, Figure 3d). We found no other significant difference in species richness

between any sample point in Thousand Palms Oasis (see Supplemental Table S6 for full

ANOVA and post-hoc test results).

PERMANOVA analyses indicate significant arthropod community turnover across time in both

Simone Pond and Thousand Palms Oasis (Table S5, Figure 3c). In Thousand Palms Oasis, all

sample periods had significantly different communities from each other except for June 2020

and December 2020, similar to results for the single marker datasets described above. All

sample points had significantly different communities in Simone Pond and higher F values than

comparisons between Thousand Palms Oasis, indicating stronger separation of arthropod

communities between sample points in Simone Pond than Thousand Palms Oasis.

4. Discussion

4.1 Candidate bioindicators of crayfish presence are valuable to monitor over time

Our analyses show individual taxa as significant drivers of community difference between

crayfish positive and negative samples, which allows for flagging of potential bioindicator

species at different times and between invaded and non-invaded sites. We found that several

microbial taxa identified through eDNA metabarcoding are reliable bioindicators for invasive

crayfish. SIMPER analyses of the two oases and experimental tanks identified 90 taxa that were

significantly more likely to occur when crayfish were present. After conducting literature review

and qualitative scoring on these taxa, we found 9 microbial taxa that are highly linked to

crayfish. We propose these 9 taxa as candidate crayfish bioindicators that can be used as

potential warning signs of invasive crayfish presence in our study site as well as other systems

where crayfish are suspected or known to be present. We found that many of these taxa are

directly linked to crayfish as a specialized host, and others that are likely associated with

crayfish due to shared habitat requirements. Our highest ranked candidate bioindicators, with

rank values of 6 (*Cynobium* sp., *Aspergillus flavus*, and *Talaromyces helicus*) were previously

identified as key associated taxa in invasive *P. clarkii* or *C. chelax* in Italy and Australia (Dörr et

al., 2012; Foysal et al., 2019; Garzoli et al., 2014). In our study, out of 1003 total eDNA samples

from across the state of California (Meyer et al. 2021), *Cynobium sp.* and *Talaromyces helicus*

were only found in samples in the Coachella Valley, suggesting that they may thrive in desert

environments where crayfish are present. *Aspergillius flavus, Phoma herbarum* and several of

the mid-ranked biodindicators in our study have been previously classified as fungal plant

pathogens (Dörr et al., 2012; e.g. Stagnospora sp., Liu et al., 2012; Ramimonilia apicalis, Ricks,

2019). Garzoli et al. (2014) suggested that these and other fungal species are common in the

crayfish gut to help break down plant detritus; a primary source of food for *P. clarkii,* and thus

are likely to be tightly associated with *P. clarkii*. Overall, 4 out of 9 candidate bioindicator taxa

were fungal species in the Phylum Ascomycota. This finding is consistent with the hypothesis

that invasive crayfish may act as vectors for fungal plant pathogens (Dörr et al. 2012; Garzoli et

al. 2014). The remaining four taxa in the 9 candidate bioindicators are all green algae in the

family Scenedesmaceae. Congeners of these taxa have been previously found to co-occur with

crayfish, and have been shown to be sensitive to crayfish pesticides (Buřič et al., 2013;

Sarıkaya et al., 2011), suggesting that they may constitute important habitat for *P. clarkii*. We

note that invasive tilapia were also present in our study system, potentially inflating the number

of taxa significantly associated with crayfish in the pre-eradication samples. However, given

that we included experimental tanks containing only tilapia as non-crayfish samples, we are

confident that any tilapia-specific taxa were filtered out of our final data set of crayfish-

associated taxa. We suggest that the microbial taxa reported here constitute a proxy indicator

community for crayfish presence, which we propose as a valuable addition to a multi-pronged

detection approach along with visual, trapping, and/or direct molecular surveys, as per Tregulier

et al. (2014).

It is unclear why multiple methods of direct molecular detection of crayfish failed in our study. It

is possible that the timing of our sample did not coincide with a period when crayfish shed the

most DNA (e.g. during molting or gravidity, Dunn et al. 2017), and so the levels of DNA in both

the oasis and tank samples were not high enough to be detected at high confidence.

Additionally, the published primer studies noting successful detection of *P. clarkii* from

environmental samples used crayfish samples in Europe, so it is also possible that there are

mutations in primer binding sites between these geographically disparate populations. Genome

sequencing of multiple *P. clarkii* populations around the globe would be useful to understand if

sequence differences are contributing to unstable molecular detection, but this was not within

the scope of our study. Regardless, the variability of detection success of *P. clarkii* both within

and between studies suggest that a multi-pronged approach including some combination of

visual surveys, trapping, and molecular detection, including direct (qPCR, targeted PCR) and/or

indirect (metabarcoding, microbial proxies) is necessary for the management of this species.

We suggest that for managers using molecular surveys, microbial indicators can be tracked

during the same analysis. If either crayfish or microbial indicators are found in the sequencing

data, managers can be on alert for potential invasion or crayfish refugia.

4.2 Community composition but not species richness varies between oases through time

Biodiversity analyses of five metabarcode loci (18S, eukaryotes; FITS, fungi; 16S, Bacteria and

Archaea; CO1, animals; and PITS, plants) showed that community composition between oases

in all taxa was significantly different at each sample period (with the exception of CO1 in Jun

2019 due to proteobacteria contamination). Communities continued to show significant

differences even in the final sample period (December 2020), which was 17 months post-

eradication and 5 months post-pesticide application in Simone Pond. This result is somewhat

surprising as both oases are fed from the same water source, are located less than 0.5 km

apart, and are characterized by the same broad vegetation type (California Fan Palm,

Washingtonia filifera). Due to the legacy of invasive crayfish and tilapia in Simone Pond, we

expected that community composition would be different between Simone Pond and Thousand

Palms Oasis in the pre-eradication sample but that the two sites would become more similar

over time. However, our results indicate that Simone Pond may still be recovering from invasive

presence and eradication. Indeed, it is possible that complete ecological recovery in Simone

Pond may never occur to the extent that communities between Thousand Palms Oasis and

Simone Pond become similar (Prior et al., 2018). It is also possible that Thousand Palms Oasis

and Simone Pond support inherently different biological communities, currently and

historically. Thousand Palms Oasis has a more closed canopy and shallower depth than

Simone Pond, which is why only Simone Pond and not Thousand Palms Oasis is targeted as a

re-introduction site for the desert pupfish (Marsh and Sada, 1993; Varela-Romero et al., 2002).

While community composition differed between oases, species richness remained relatively

consistent between them. Different datasets showed different peaks of species richness at

different time points, but richness rarely differed between oases at a single time point. Again, we

were surprised by this result as we expected that invasive presence and eradication would

differentially affect overall species richness between invaded and non-invaded oases. This was

only true when analyzing the species richness of Arthropoda over time. Other studies have

similarly shown that invasive species may not decrease overall species richness, especially

when considering native and non-native species (Fork, 2010; Forys and Allen, 2002; Matthews

et al., 2009). Tracking both native and overall community composition and richness between

oases, along with invasive species bioindicators, will continue to be of interest as the next

phases of restoration at the preserve continue (Short, 2017).

4.3 Cypermethrin negatively impacted arthropod biodiversity in Simone Pond

Although the pesticide cypermethrin was deemed necessary to eradicate crayfish from Simone

Pond, managers were concerned about the potential for off-target effects of the pesticides on

other arthropod taxa. Our analyses of taxa in Arthropoda in CO1 and 18S datasets reveal that

all three sample periods post-eradication had significantly lower arthropod species richness than

the pre-eradication sample in Simone Pond. For example, taxa in the classes Malacostraca

(which also include crayfish), and Hexanuplia (containing mainly Copepods) were found in the

June 2019 sample in Simone Pond, but were either eliminated or found in lower sequence read

abundance in subsequent samples. The final sample in December 2020, where pesticide had

not been applied for 5 months, had levels of arthropod species richness that were similar to the

earlier post-eradication samples. Analysis of Thousand Palms Oasis did not show the same

change in arthropod species richness through time: post-eradication samples had similar or

higher levels of arthropod species richness compared to December 2019. As Thousand Palms

Oasis was effectively a control sample with no pesticide applied, these results suggest that,

while the cypermethrin negatively impacted arthropod species richness in Simone Pond, it did

not affect arthropod communities in the other oases in the preserve. Previous studies have

highlighted the persistent toxicity of pyrethroid pesticides in sediments on benthic invertebrates

(Li et al., 2017; Zhang et al., 2018). Further monitoring of sites treated with cypermethrin for the

purposes of crayfish eradication will be necessary to understand the long-term effects of

cypermethrin treatment on both aquatic and terrestrial arthropod communities.

5. Conclusions and Conservation Applications

We used multi-locus metabarcoding of environmental DNA in an aquatic ecosystem undergoing

conservation management to simultaneously identify microbial bioindicators for an invasive

crayfish and track biodiversity changes. We were unable to reliably detect crayfish DNA in

environmental samples using either metabarcoding or qPCR, which is consistent with previous

studies of crayfish (Treguier et al. 2014) and other crustaceans (e.g. Crane et al. 2021).

However, our community-level analyses of bacterial and fungal markers in habitats where

crayfish were known to be present (in tanks with crayfish and the invaded oasis pre-eradication)

revealed a community of microbial taxa that are strongly associated with crayfish

presence. Analyses of post-eradication environmental samples from the invaded pond revealed

few of these bioindicator taxa, suggesting that crayfish are no longer present, consistent with

the traditional survey data. Our biodiversity analyses of five metabarcode datasets also

revealed that biological communities differed between oases through time, although species

richness estimates tended to be similar between oases at individual time points. Finally, we

found that arthropod diversity declined after crayfish pesticide treatment but did not decline in

the oasis that was not treated with pesticide, suggesting that cypermethrin use may have

detrimental off-target effects to other species. Overall, we recommend conducting ongoing

monitoring using multiple methods including eDNA for native and invasive species as a powerful

approach to track biodiversity and invasive status throughout a conservation project. As

managers of invaded systems look for solutions for restoration and monitoring, the use of

metabarcoding can be an invaluable tool to track cryptic species with multiple approaches and

monitor biodiversity.

Acknowledgments

The authors wish to acknowledge Deborah Rogers, Center for Natural Lands Management (CNLM)

Director of Conservation Science and Stewardship, for providing invaluable feedback on manuscript

drafts. We thank C. Orland, H. Bregoff, C.W. Schaal, A. Monzón, E. De Garcas, R. Henderson, and Y.

Xiao for assistance with environmental DNA sampling. We finally wish to thank the members of the

CALeDNA team and eDNA working group in the Paleogenomics Lab at the University of California

Santa Cruz for overall support, as well as continual feedback on data analysis and visualization

throughout the project. This work was funded by the HHMI Professors Program grant no. GT10483 as

well as the State of California Proposition 1 and USFWS Strategic Habitat Conservation Grants.

References

Alcorlo, P., Geiger, W., Otero, M., 2004. Feeding Preferences and Food Selection of the Red
Swamp Crayfish, Procambarus clarkii, in Habitats Differing in Food Item Diversity.
Crustaceana 77, 435–453.
Amaral-Zettler, L.A., McCliment, E.A., Ducklow, H.W., Huse, S.M., 2009. A Method for Studying
Protistan Diversity Using Massively Parallel Sequencing of V9 Hypervariable Regions of
Small-Subunit Ribosomal RNA Genes. PLOS ONE 4, e6372.
https://doi.org/10.1371/journal.pone.0006372
Anderson, M.J., Walsh, D.C.I., 2013. PERMANOVA, ANOSIM, and the Mantel test in the face of
heterogeneous dispersions: What null hypothesis are you testing? Ecological
Monographs 83, 557–574. https://doi.org/10.1111/j.2041-210X.2011.00127.x
Astudillo-García, C., Hermans, S.M., Stevenson, B., Buckley, H.L., Lear, G., 2019. Microbial
assemblages and bioindicators as proxies for ecosystem health status: potential and
limitations. Appl Microbiol Biotechnol 103, 6407–6421. https://doi.org/10.1007/s00253-
019-09963-0
Ballare, K.M., Neff, J.L., Ruppel, R., Jha, S., 2019. Multi-scalar drivers of biodiversity: local
management mediates wild bee community response to regional urbanization.
Ecological Applications 29, e01869.

Barnes, M.A., Turner, C.R., Jerde, C.L., Renshaw, M.A., Chadderton, W.L., Lodge, D.M., 2014.
720   Environmental Conditions Influence eDNA Persistence in Aquatic Systems. Environ. Sci.
721   Technol. 48, 1819–1827. https://doi.org/10.1021/es404734p
722   Bucciarelli, G.M., Suh, D., Lamb, A.D., Roberts, D., Sharpton, D., Shaffer, H.B., Fisher, R.N.,
723   Kats, L.B., 2019. Assessing effects of non-native crayfish on mosquito survival.
724   Conservation Biology 33, 122–131. https://doi.org/10.1111/cobi.13198
725   Buřič, M., Kouba, A., Máchová, J., Mahovská, I., Kozák, P., 2013. Toxicity of the
726   organophosphate pesticide diazinon to crayfish of differing age. Int. J. Environ. Sci.
727   Technol. 10, 607–610. https://doi.org/10.1007/s13762-013-0185-4
728   Bushnell, B., 2014. BBTools. BBMap short read aligner, and other bioinformatic tools. Available
729   online at: https://sourceforge. net/projects/bbmap.
730   Cai, P., Huang, Q., Zhang, X., Chen, H., 2006. Adsorption of DNA on clay minerals and various
731   colloidal particles from an Alfisol. Soil Biology and Biochemistry 38, 471–476.
732   https://doi.org/10.1016/j.soilbio.2005.05.019
733   Cai, W., Ma, Z., Yang, C., Wang, L., Wang, W., Zhao, G., Geng, Y., Yu, D.W., 2017. Using
734   eDNA to detect the distribution and density of invasive crayfish in the Honghe-Hani rice
735   terrace World Heritage site. PLOS ONE 12, e0177724.
736   https://doi.org/10.1371/journal.pone.0177724
737   Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., Holmes, S.P., 2016.
738   DADA2: High-resolution sample inference from Illumina amplicon data. Nature methods
739   13, 581–583.
740   Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Huntley, J., Fierer, N., Owens,
741   S.M., Betley, J., Fraser, L., Bauer, M., Gormley, N., Gilbert, J.A., Smith, G., Knight, R.,
742   2012. Ultra-high-throughput microbial community analysis on the Illumina HiSeq and
743   MiSeq platforms. ISME J 6, 1621–1624. https://doi.org/10.1038/ismej.2012.8
744   Chen, X., Fan, L., Qiu, L., Dong, X., Wang, Q., Hu, G., Meng, S., Li, D., Chen, J., 2021.
745   Metagenomics Analysis Reveals Compositional and Functional Differences in the Gut
746   Microbiota of Red Swamp Crayfish, Procambarus clarkii, Grown on Two Different
747   Culture Environments. Frontiers in Microbiology 12.
748   Chucholl, F., Fiolka, F., Segelbacher, G., Epp, L.S., 2021. eDNA Detection of Native and
749   Invasive Crayfish Species Allows for Year-Round Monitoring and Large-Scale Screening
750   of Lotic Systems. Frontiers in Environmental Science 9.
751   Clarke, K.R., 1993. Non-parametric multivariate analyses of changes in community structure.
752   Australian Journal of Ecology 18, 117–143.
753   Crane, L.C., Goldstein, J.S., Thomas, D.W., Rexroth, K.S., Watts, A.W., 2021. Effects of life
754   stage on eDNA detection of the invasive European green crab (Carcinus maenas) in
755   estuarine systems. Ecological Indicators 124, 107412.
756   https://doi.org/10.1016/j.ecolind.2021.107412
757   Curd, E.E., Gold, Z., Kandlikar, G.S., Gomer, J., Ogden, M., O'Connell, T., Pipes, L., Schweizer,
758   T.M., Rabichow, L., Lin, M., Shi, B., Barber, P.H., Kraft, N., Wayne, R., Meyer, R.S.,
759   2019. Anacapa Toolkit: An environmental DNA toolkit for processing multilocus
760   metabarcode datasets. Methods in Ecology and Evolution 10, 1469–1475.
761   Curtis, A.N., Larson, E.R., 2020. No evidence that crayfish carcasses produce detectable
762   environmental DNA (eDNA) in a stream enclosure experiment. PeerJ 8, e9333.
763   https://doi.org/10.7717/peerj.9333
764   Davis, N.M., Proctor, D.M., Holmes, S.P., Relman, D.A., Callahan, B.J., 2018. Simple statistical
765   identification and removal of contaminant sequences in marker-gene and metagenomics
766   data. Microbiome 6, 1–14.
767   Dörr, A.J.M., Rodolfi, M., Elia, A.C., Scalici, M., Garzoli, L., Picco, A.M., 2012. Mycoflora on the
768   cuticle of the invasive crayfish Procambarus clarkii. Fundamental and Applied Limnology
769   180, 77–84.

Dougherty, M.M., Larson, E.R., Renshaw, M.A., Gantz, C.A., Egan, S.P., Erickson, D.M.,
Lodge, D.M., 2016. Environmental DNA (eDNA) detects the invasive rusty crayfish
Orconectes rusticus at low abundances. Journal of Applied Ecology 53, 722–732.
https://doi.org/10.1111/1365-2664.12621
Dragičević, P., Bielen, A., Petrić, I., Hudina, S., 2021. Microbial pathogens of freshwater
crayfish: A critical review and systematization of the existing data with directions for
future research. Journal of Fish Diseases 44, 221–247. https://doi.org/10.1111/jfd.13314
Dunn, N., Priestley, V., Herraiz, A., Arnold, R., Savolainen, V., 2017. Behavior and season
affect crayfish detection and density inference using environmental DNA. Ecology and
Evolution 7, 7777–7785. https://doi.org/10.1002/ece3.3316
Edwards, R., Millburn, P., Hutson, D.H., 1986. Comparative toxicity of cis-cypermethrin in
rainbow trout, frog, mouse, and quail. Toxicology and Applied Pharmacology 84, 512–
522. https://doi.org/10.1016/0041-008X(86)90256-5
Ficetola, G.F., Manenti, R., Taberlet, P., 2019. Environmental DNA and metabarcoding for the
study of amphibians and reptiles: species distribution, the microbiome, and much more.
Amphibia-Reptilia 40, 129–148. https://doi.org/10.1163/15685381-20191194
Ficetola, G.F., Miaud, C., Pompanon, F., Taberlet, P., 2008. Species detection using
environmental DNA from water samples. Biology Letters 4, 423–425.
https://doi.org/10.1098/rsbl.2008.0118
Fork, S.K., 2010. Arthropod Assemblages on Native and Nonnative Plant Species of a Coastal
Reserve in California. Environ Entomol 39, 753–762. https://doi.org/10.1603/EN09185
Forys, E.A., Allen, C.R., 2002. Functional Group Change within and across Scales following
Invasions and Extinctions in the Everglades Ecosystem. Ecosystems 5, 339–347.
https://doi.org/10.1007/s10021-001-0078-0
Foster, Z.S.L., Sharpton, T.J., Grünwald, N.J., 2017. Metacoder: An R package for visualization
and manipulation of community taxonomic diversity data. PLOS Computational Biology
13, e1005404. https://doi.org/10.1371/journal.pcbi.1005404
Foysal, M.J., Nguyen, T.T.T., Chaklader, M.R., Siddik, M.A.B., Tay, C.-Y., Fotedar, R., Gupta,
S.K., 2019. Marked variations in gut microbiota and some innate immune responses of
fresh water crayfish, marron (Cherax cainii, Austin 2002) fed dietary supplementation of
Clostridium butyricum. PeerJ 7, e7553. https://doi.org/10.7717/peerj.7553
Gao, X., Lin, H., Revanna, K., Dong, Q., 2017. A Bayesian taxonomic classification method for
16S rRNA gene sequences with improved species-level accuracy. BMC bioinformatics
18, 1–10.
Garzoli, L., Paganelli, D., Rodolfi, M., Savini, D., Moretto, M., Occhipinti, A., Picco, A., 2014.
First evidence of microfungal "extra oomph" in the invasive red swamp crayfish
Procambarus clarki. Aquatic Invasions 9, 47–58. https://doi.org/10.3391/ai.2014.9.1.04
Geerts, A.N., Boets, P., Van den Heede, S., Goethals, P., Van der heyden, C., 2018. A search
for standardized protocols to detect alien invasive crayfish based on environmental DNA
(eDNA): A lab and field evaluation. Ecological Indicators 84, 564–572.
https://doi.org/10.1016/j.ecolind.2017.08.068
Geiger, W., Alcorlo, P., Baltanás, A., Montes, C., 2005. Impact of an introduced Crustacean on
the trophic webs of Mediterranean wetlands. Biological Invasions 7, 49–73.
https://doi.org/10.1007/s10530-004-9635-8
Gordon, A., Hannon, G., 2010. Fastx-toolkit. FASTQ/A short-reads preprocessing tools
(unpublished) http://hannonlab. cshl. edu/fastx_toolkit 5.
Gu, W., Song, J., Cao, Y., Sun, Q., Yao, H., Wu, Q., Chao, J., Zhou, J., Xue, W., Duan, J.,
2013. Application of the ITS2 Region for Barcoding Medicinal Plants of Selaginellaceae
in Pteridophyta. PLOS ONE 8, e67818. https://doi.org/10.1371/journal.pone.0067818
Halstead, B.J., Wood, D.A., Bowen, L., Waters, S.C., Vandergast, A.G., Ersan, J.S., Skalos,
S.M., Casazza, M.L., 2017. An evaluation of the efficacy of using environmental DNA

(eDNA) to detect giant gartersnakes (Thamnophis gigas) (No. 2017–1123), Open-File
Report. U.S. Geological Survey. https://doi.org/10.3133/ofr20171123

Harper, K.J., Anucha, P., Turnbull, J.F., Bean, C.W., Leaver, M.J., 2018. Searching for a signal:
environmental DNA (eDNA) for the detection of invasive signal crayfish, Pacifasticus
leniusculus (Dana, 1852). Management of Biological Invasions 9, 137–148.
https://doi.org/10.3391/mbi.2018.9.2.07

Hebert, P.D.N., Cywinska, A., Ball, S.L., deWaard, J.R., 2003. Biological Identifications through
DNA Barcodes. Proceedings: Biological Sciences 270, 313–321.

Ikeda, K., Doi, H., Tanaka, K., Kawai, T., Negishi, J.N., 2016. Using environmental DNA to
detect an endangered crayfish Cambaroides japonicus in streams. Conservation Genet
Resour 8, 231–234. https://doi.org/10.1007/s12686-016-0541-z

Jarman, S.N., Berry, O., Bunce, M., 2018. The value of environmental DNA biobanking for long-
term biomonitoring. Nature Ecology & Evolution 2, 1192–1193.
https://doi.org/10.1038/s41559-018-0614-3

Kandlikar, G.S., Gold, Z.J., Cowen, M.C., Meyer, R.S., Freise, A.C., Kraft, N.J., Moberg-Parker,
J., Sprague, J., Kushner, D.J., Curd, E.E., 2018. ranacapa: An R package and Shiny
web app to explore environmental DNA data with exploratory statistics and interactive
visualizations. F1000Research 7.

Khan, A., Faridi, H.A.M., Ali, M., Khan, M.Z., Siddique, M., Hussain, I., Ahmad, M., 2009. Effects
of cypermethrin on some clinico-hemato-biochemical and pathological parameters in
male dwarf goats (Capra hircus). Experimental and Toxicologic Pathology 61, 151–160.
https://doi.org/10.1016/j.etp.2008.07.001

Knudsen, S.W., Agersnap, S., Møller, P.R., Andersen, J.H., 2019. Development of species-
specific eDNA-based test systems for monitoring of freshwater crayfish. NIVA-rapport.

Langmead, B., Salzberg, S.L., 2012. Fast gapped-read alignment with Bowtie 2. Nature
methods 9, 357–359.

Leray, M., Yang, J.Y., Meyer, C.P., Mills, S.C., Agudelo, N., Ranwez, V., Boehm, J.T., Machida,
R.J., 2013. A new versatile primer set targeting a short fragment of the mitochondrial
COI region for metabarcoding metazoan diversity: application for characterizing coral
reef fish gut contents. Frontiers in Zoology 10, 34. https://doi.org/10.1186/1742-9994-10-
34

Leslie, P.H., Davis, D.H.S., 1939. An Attempt to Determine the Absolute Number of Rats on a
Given Area. Journal of Animal Ecology 8, 94–113. https://doi.org/10.2307/1255

Li, H., Cheng, F., Wei, Y., Lydy, M.J., You, J., 2017. Global occurrence of pyrethroid
insecticides in sediment and the associated toxicological effects on benthic
invertebrates: An overview. Journal of Hazardous Materials 324, 258–271.
https://doi.org/10.1016/j.jhazmat.2016.10.056

Li, H., Durbin, R., 2009. Fast and accurate short read alignment with Burrows–Wheeler
transform. bioinformatics 25, 1754–1760.

Liu, Z., Zhang, Z., Faris, J.D., Oliver, R.P., Syme, R., McDonald, M.C., McDonald, B.A.,
Solomon, P.S., Lu, S., Shelver, W.L., Xu, S., Friesen, T.L., 2012. The Cysteine Rich
Necrotrophic Effector SnTox1 Produced by Stagonospora nodorum Triggers
Susceptibility of Wheat Lines Harboring Snn1. PLOS Pathogens 8, e1002467.
https://doi.org/10.1371/journal.ppat.1002467

Marsh, P.C., Sada, D.W., 1993. Draft Desert Pupfish RP Amendment.pdf. U.S. Fish and Wildlife
Service.

Martin, B.A., Saiki, M.K., 2005. Relation of desert pupfish abundance to selected environmental
variables in natural and manmade habitats in the Salton Sea basin. Environ Biol Fish 73,
97–107. https://doi.org/10.1007/s10641-004-5569-3

Martin, M., 2011. Cutadapt removes adapter sequences from high-throughput sequencing
reads. EMBnet.journal 17, 10–12. https://doi.org/10.14806/ej.17.1.200

Matthews, J.W., Peralta, A.L., Soni, A., Baldwin, P., Kent, A.D., Endress, A.G., 2009. Local and
landscape correlates of non-native species invasion in restored wetlands. Ecography 32,
1031–1039. https://doi.org/10.1111/j.1600-0587.2009.05863.x
Mauvisseau, Q., Coignet, A., Delaunay, C., Pinet, F., Bouchon, D., Souty-Grosset, C., 2018.
Environmental DNA as an efficient tool for detecting invasive crayfishes in freshwater
ponds. Hydrobiologia 805, 163–175. https://doi.org/10.1007/s10750-017-3288-y
McMurdie, P.J., Holmes, S., 2012. Phyloseq: a bioconductor package for handling and analysis
of high-throughput phylogenetic sequence data, in: Biocomputing 2012. World Scientific,
pp. 235–246.
Meyer, F., Paarmann, D., D'Souza, M., Olson, R., Glass, E., Kubal, M., Paczian, T., Rodriguez,
A., Stevens, R., Wilke, A., Wilkening, J., Edwards, R., 2008. The metagenomics RAST
server – a public resource for the automatic phylogenetic and functional analysis of
metagenomes. BMC Bioinformatics 9, 386. https://doi.org/10.1186/1471-2105-9-386
Meyer, R.S., Ramos, M.M., Lin, M., Schweizer, T.M., Gold, Z., Ramos, D.R., Shirazi, S.,
Kandlikar, G., Kwan, W.-Y., Curd, E.E., Freise, A., Parker, J.M., Sexton, J.P., Wetzer,
R., Pentcheff, N.D., Wall, A.R., Pipes, L., Garcia-Vedrenne, A., Mejia, M.P., Moore, T.,
Orland, C., Ballare, K.M., Worth, A., Beraut, E., Aronson, E., Nielson, R., Lewin, H.A.,
Barber, P.H., Wall, J., Kraft, N., Shapiro, B., Wayne, R.K., 2021. The CALeDNA
program: Citizen scientists and researchers inventory California's biodiversity. California
Agriculture 75. https://doi.org/10.3733/ca.2021a0001
Morolli, C., Quaglio, F., Della Rocca, G., Malvisi, J., Di Salvo, A., 2006. EVALUATION OF THE
TOXICITY OF SYNTHETIC PYRETHROIDS TO RED SWAMP CRAYFISH
(PROCAMBARUS CLARKII, GIRARD 1852) AND COMMON CARP (CYPRINUS
CARPIO, L. 1758). Bull. Fr. Pêche Piscic. 1381–1394.
https://doi.org/10.1051/kmae:2006042
Nunes, A.L., Hoffman, A.C., Zengeya, T.A., Measey, G.J., Weyl, O.L., 2017. Red swamp
crayfish, Procambarus clarkii, found in South Africa 22 years after attempted eradication.
Aquatic Conservation: Marine and Freshwater Ecosystems 27, 1334–1340.
https://doi.org/10.1002/aqc.2741
Oficialdegui, F.J., Sánchez, M.I., Clavero, M., 2020. One century away from home: how the red
swamp crayfish took over the world. Rev Fish Biol Fisheries 30, 121–135.
https://doi.org/10.1007/s11160-020-09594-z
Oksanen, J., Blanchet, F.G., Kindt, R., Legendre, P., Minchin, P.R., O'hara, R., Simpson, G.L.,
Solymos, P., Stevens, M.H.H., Wagner, H., 2015. Vegan: community ecology package.
2015. R package version 2.
Poelen, J.H., Simons, J.D., Mungall, C.J., 2014. Global biotic interactions: An open
infrastructure to share and analyze species-interaction datasets. Ecological Informatics
24, 148–159. https://doi.org/10.1016/j.ecoinf.2014.08.005
Prior, K.M., Adams, D.C., Klepzig, K.D., Hulcr, J., 2018. When does invasive species removal
lead to ecological recovery? Implications for management success. Biol Invasions 20,
267–283. https://doi.org/10.1007/s10530-017-1542-x
Pritchard, E.G., Chadwick, D.D.A., Patmore, I.R., Chadwick, M.A., Bradley, P., Sayer, C.D.,
Axmacher, J.C., 2021. The 'Pritchard Trap': A novel quantitative survey method for
crayfish. Ecological Solutions and Evidence 2, e12070. https://doi.org/10.1002/2688-
8319.12070
Ratsch, R., Kingsbury, B.A., Jordan, M.A., 2020. Exploration of Environmental DNA (eDNA) to
Detect Kirtland's Snake (Clonophis kirtlandii). Animals 10, 1057.
https://doi.org/10.3390/ani10061057
Ricks, N., 2019. A Metagenomic Approach to Understand Stand Failure in Bromus tectorum.
Theses and Dissertations.

Russell, S.L., Pepper-Tunick, E., Svedberg, J., Byrne, A., Castillo, J.R., Vollmers, C., Beinart,
R.A., Corbett-Detig, R., 2020. Horizontal transmission and recombination maintain
forever young bacterial symbiont genomes. PLOS Genetics 16, e1008935.
https://doi.org/10.1371/journal.pgen.1008935
Sandodden, R., Johnsen, S.I., 2010. Eradication of introduced signal crayfish Pasifastacus
leniusculus using the pharmaceutical BETAMAX VET.®. AI 5, 75–81.
https://doi.org/10.3391/ai.2010.5.1.9
Sarıkaya, R., Sepici-Dinçel, A., Ça□lan Karasu Benli, A., Selvi, M., Erkoç, F., 2011. The acute
toxicity of fenitrothion on narrow-clawed crayfish (Astacus leptodactylus Eschscholtz,
1823) in association with biomarkers of lipid peroxidation. Journal of Biochemical and
Molecular Toxicology 25, 169–174. https://doi.org/10.1002/jbt.20373
Sepulveda, A.J., Nelson, N.M., Jerde, C.L., Luikart, G., 2020. Are Environmental DNA Methods
Ready for Aquatic Invasive Species Management? Trends in Ecology & Evolution 35,
668–678. https://doi.org/10.1016/j.tree.2020.03.011
Shafer, T.J., Meyer, D.A., Crofton, K.M., 2005. Developmental neurotoxicity of pyrethroid
insecticides: critical review and future research needs. Environ Health Perspect 113,
123–136. https://doi.org/10.1289/ehp.7254
Short, G., 2021. Desert Pupfish Refuge Habitat Restoration Project. Center for Natural Lands
Management.
Short, G., 2017. Ten year management plan for the Thousand Palms Oasis and McCallum
Grove Preserve (S004 & S050) and Whitewater Mitigation Area (S049) Thousand
Palms, CA. Center for Natural Lands Management.
Shui, Y., Guan, Z.-B., Liu, G.-F., Fan, L.-M., 2020. Gut microbiota of red swamp crayfish
Procambarus clarkii in integrated crayfish-rice cultivation model. AMB Express 10, 5.
https://doi.org/10.1186/s13568-019-0944-9
Skelton, J., Geyer, K.M., Lennon, J.T., Creed, R.P., Brown, B.L., 2017. Multi-scale ecological
filters shape the crayfish microbiome. Symbiosis 72, 159–170.
https://doi.org/10.1007/s13199-016-0469-9
Soares, M.P., Oliveira, N., Rebelo, D., Marcondes, S.F., Fernandes, C.E., Domingues, I.,
Soares, A., Hayd, L., 2019. Cypermethrin-based formulation Barrage® induces
histological changes in gills of the Pantanal endemic shrimp Macrobrachium
pantanalense. Environmental Toxicology and Pharmacology 67, 66–72.
https://doi.org/10.1016/j.etap.2019.01.014
Souty-Grosset, C., Anastácio, P.M., Aquiloni, L., Banha, F., Choquer, J., Chucholl, C., Tricarico,
E., 2016. The red swamp crayfish Procambarus clarkii in Europe: Impacts on aquatic
ecosystems and human well-being. Limnologica 58, 78–93.
https://doi.org/10.1016/j.limno.2016.03.003
Suarez-Menendez, M., Planes, S., Garcia-Vazquez, E., Ardura, A., 2020. Early Alert of
Biological Risk in a Coastal Lagoon Through eDNA Metabarcoding. Frontiers in Ecology
and Evolution 8.
Thalinger, B., Deiner, K., Harper, L.R., Rees, H.C., Blackman, R.C., Sint, D., Traugott, M.,
Goldberg, C.S., Bruce, K., 2021. A validation scale to determine the readiness of
environmental DNA assays for routine species monitoring. Environmental DNA 3, 823–
836. https://doi.org/10.1002/edn3.189
Tréguier, A., Paillisson, J.-M., Dejean, T., Valentini, A., Schlaepfer, M.A., Roussel, J.-M., 2014.
Environmental DNA surveillance for invertebrate species: advantages and technical
limitations to detect invasive crayfish Procambarus clarkii in freshwater ponds. Journal of
Applied Ecology 51, 871–879. https://doi.org/10.1111/1365-2664.12262
Troth, C.R., Burian, A., Mauvisseau, Q., Bulling, M., Nightingale, J., Mauvisseau, C., Sweet,
M.J., 2020. Development and application of eDNA-based tools for the conservation of

972   white-clawed crayfish. Science of The Total Environment 748, 141394.
973   https://doi.org/10.1016/j.scitotenv.2020.141394
974 Turner, C.R., Uy, K.L., Everhart, R.C., 2015. Fish environmental DNA is more concentrated in
975   aquatic sediments than surface water. Biological Conservation, Special Issue:
976   Environmental DNA: A powerful new tool for biological conservation 183, 93–102.
977   https://doi.org/10.1016/j.biocon.2014.11.017
978 Varela-Romero, A., Ruiz-Campos, G., Yépiz-Velázquez, L.M., Alaníz-García, J., 2002.
979   Distribution, habitat and conservation status of desert pupfish (Cyprinodon macularius)
980   in the Lower Colorado River Basin, Mexico. Reviews in Fish Biology and Fisheries 12,
981   157–165. https://doi.org/10.1023/A:1025006920052
982 Warton, D.I., Wright, S.T., Wang, Y., 2012. Distance-based multivariate analyses confound
983   location and dispersion effects. Methods in Ecology and Evolution 3, 89–101.
984   https://doi.org/10.18637/jss.v032.i10
985 Wei, N., Nakajima, F., Tobino, T., 2018. A Microcosm Study of Surface Sediment Environmental
986   DNA: Decay Observation, Abundance Estimation, and Fragment Length Comparison.
987   Environ. Sci. Technol. 52, 12428–12435. https://doi.org/10.1021/acs.est.8b04956
988 White, T., Bruns, T., Lee, S., Taylor, J., 1990. Amplification and direct sequencing of fungal
989   ribosomal RNA genes for phylogenetics, in: PCR Protocols: A Guide to Methods and
990   Applications. Academic Press, New York, pp. 315–322.
991 Wickham, H., Chang, W., Wickham, M.H., 2016. Package 'ggplot2.' Create elegant data
992   visualisations using the grammar of graphics. Version 2, 1–189.
993 Wittwer, C., Stoll, S., Strand, D., Vrålstad, T., Nowak, C., Thines, M., 2018. eDNA-based
994   crayfish plague monitoring is superior to conventional trap-based assessments in year-
995   round detection probability. Hydrobiologia 807, 87–97. https://doi.org/10.1007/s10750-
996   017-3408-8
997 Zhang, J., You, J., Li, H., Tyler Mehler, W., Zeng, E.Y., 2018. Particle-scale understanding of
998   cypermethrin in sediment: Desorption, bioavailability, and bioaccumulation in benthic
999   invertebrate Lumbriculus variegatus. Science of The Total Environment 642, 638–645.
1000   https://doi.org/10.1016/j.scitotenv.2018.06.098

1001

1002

1003 **Tables**

1004 **Table 1.** Sampling Summary

Sample Date	Simone Pond Condition	N Samples Simone Pond				N Samples Thousand Palms Oasis				N Samples Tanks
		Water	Sediment	Soil	Total	Water	Sediment	Soil	Total	Water Only
Jun 2019	Invasives Present	4	21	0	25	10	27	6	43	13
Dec 2019	Cypermethrin applied monthly since July 2019	0	31	3	34	0	14	2	16	0
Jun 2020	Cypermethrin applied monthly since July 2019	0	31	6	37	0	15	5	20	0
Dec 2020	No mitigation for 5 months	0	28	3	31	0	8	8	16	0

1005

1006 **Table 2.** Highly ranked crayfish associates (N=9) ordered by qualitative rank value

Domain	Phylum	Class	Order	Family	Genus	Species	Rank Value

Bacteria	Cyanobacteria	Cyanophyceae	Synechococcales	Synechococcaceae	Cyanobium	unknown	6
Eukaryota	Ascomycota	Eurotiomycetes	Eurotiales	Aspergillaceae	Aspergillus	*Aspergillus flavus*	6
Eukaryota	Ascomycota	Eurotiomycetes	Eurotiales	Trichocomaceae	Talaromyces	*Talaromyces helicus*	6
Eukaryota	Ascomycota	Dothideomycetes	Pleosporales	Didymellaceae	Phoma	*Phoma herbarum*	5
Eukaryota	Ascomycota	Dothideomycetes	Pleosporales	Massarinaceae	Stagonospora	unknown	5
Eukaryota	Chlorophyta	Chlorophyceae	Sphaeropleales	Scenedesmaceae	Desmodesmus	*Desmodesmus intermedius*	5
Eukaryota	Chlorophyta	Chlorophyceae	Sphaeropleales	Scenedesmaceae	Desmodesmus	*Desmodesmus pseudocommunis*	5
Eukaryota	Chlorophyta	Chlorophyceae	Sphaeropleales	Scenedesmaceae	Desmodesmus	*Desmodesmus serratus*	5
Eukaryota	Chlorophyta	Chlorophyceae	Sphaeropleales	Scenedesmaceae	Scenedesmus	*Scenedesmus bijugus*	5

1007 *Table 2 Note: List was obtained from the original 90 candidate bioindicators that were significantly more*
1008 *likely to occur in Jun 2019 crayfish invaded samples (see Supplemental Table S4), after literature review*
1009 *and qualitative scoring (possible values 0-6).*
1010

1011 **Table 3. SIMPER Significant Taxa After Crayfish Removal**

Sample Date	Phylum	Class	Order	Family	Genus	Species	Rank Value
Dec 2019	Spirochaetes	Spirochaetia	Leptospirales	Leptospiraceae	Leptospira	Leptospira sp. enrichment culture clone B4-33	NA
Dec 2019	**Proteobacteria**	**Deltaproteobacteria**	**Syntrophobacterales**	**Syntrophobacteraceae**	**unknown**	**unknown**	**0**
Dec 2019	**Proteobacteria**	**Betaproteobacteria**	**Nitrosomonadales**	**unknown**	**unknown**	**unknown**	**0**
Dec 2019	Chloroflexi	Anaerolinealyae	Anaerolineales	Anaerolineaceae	Thermomarinilinea	Thermomarinilinea sp.	NA
June 2020	NA						
Dec 2020	Acidobacteria	Vicinamibacteria	unknown	Vicinamibacteraceae	unknown		NA

1012

Table 3 Note: Taxa that are significantly more likely to occur in invaded Simone Pond as compared to
non-invaded Thousand Palms Oasis in the post-eradication samples. Two taxa from the Dec 2019
sample were also significantly associated with crayfish-positive samples collected in July 2019, these are
indicated in bold. However, both of these had a rank value of 0, indicating little to no true association with
crayfish. We observed no taxa significantly associated with Simone Pond in the June 2020 samples, and
one taxon was significant in the December 2020 sample.

1019

1020 **Figure Legends**

1021 **Figure 1**. Map of study area, located in Colorado desert, Southern California, USA. Left, Aerial view of

1022 Thousand Palms Oasis Preserve. Top right, Simone Pond, oasis invaded with *Procambarus clarkii.*

1023 Bottom right, Thousand Palms Oasis, not invaded. Shaded portions indicate inundated areas.

1024

Figure 2. a) Phylogenetic heat tree of 90 crayfish bioindicator species, made using the *Metacoder* R

package (Foster et al., 2017). Nodes and branches with higher thickness had higher numbers of

sequence reads, darker teal were found in more samples. b,c) Heat maps of frequency of b) 90

bioindicator taxa and c) 90 taxa sampled randomly from 16S and FITS datasets with 20 permutations.

Color indicates Euclidean distance in frequency and composition of taxa between sample points, with

blue and purple indicating more similarity in composition and green and yellow indicating more

differences. Square areas within the dashed lines are comparisons between Simone Pond and

Thousand Palms Oasis. Areas outside the dashed lines are comparisons within the same oasis.

1033

Figure 3. Biodiversity within a, b) the full 18S (Eukaryote) dataset and c, d) Arthropoda only from 18S

and CO1 combined datasets. Thousand Palms Oasis (TPO) is represented in blue and Simone Pond

(SP) is represented in orange for panels a, b, and d . a) Axes 1 and 2 of NMDS ordination faceted by

sample date. b) Violin plot of Chao1 richness at each sample point. Internal white points and bars

indicate mean value with 95% confidence interval generated from 1000 bootstraps. Letters indicate

significant group differences in species richness by sample date (group a or b). There was no difference

in species richness between oases within a timepoint. c) Barplot showing relative sequence read

abundance of detected classes of Arthropoda in each sample and oasis. d) Violin plot of Chao1 richness

of Arthropoda at each sample point, faceted by oasis.   Internal white points and bars indicate mean value

with 95% confidence interval generated from 1000 bootstraps. Letters indicate significant group

differences in species richness by oasis and sample date (group a or b, within an oasis).

1045

1046

1047

Simone Pond
Thousand Palms Oasis
Colorado Desert
California State Outline

a)

b)

c)

Euclidean Distance