Metabolic shift toward ketosis in asocial cavefish increases social-like collective behavior.

Motoko Iwashita[1], Amity Tran[1], Miki Wong[2,3], Ryan Lee[4,5], and Masato Yoshizawa[1]*

[1] School of Life Sciences, the University of Hawai'i at Mānoa, Honolulu, HI 96822, USA
[2] Nā Pu'uwai Native Hawaiian Healthcare System, Kaunakakai, HI 96748, USA
[3] Nutrition Services Department, Shriners Hospitals for Children, Honolulu, HI 96826, USA
[4] Medical Staff Department, Shriners Hospitals for Children, Honolulu, HI 96826, USA
[5] Department of Pediatrics, University of Hawai'i at Mānoa, Honolulu, HI 96813, USA

Corresponding author:
Masato Yoshizawa
yoshizaw@hawaii.edu
2538 McCarthy Mall, EDM 216
Honolulu, HI 96822, USA

Keywords: Ketogenic diet, Social affinity, Collective behavior, Astyanax, Plasticity, Metabolism

Conflict of Interest: The authors declare no competing financial interests.

Acknowledgement: We thank C Balaan for constructive comments regarding insights on social-like interactions in cavefish. We are grateful to V Crystal, J Choi, L Lu, J Nguyen, VFL Fernandes, K Lactaoen, M Worsham, H Hernandez, N Doeden, J Kato, M Ito, R Balmilero-Unciano, E Doy, A Martinez, and D Mones for fish care assistance. We gratefully acknowledge supports from the National Institute of Health (P20GM125508) to MY, Hawaii Community Foundation (18CON-90818) to MY.

Abstract

Social affinity and collective behavior are nearly ubiquitous in the animal kingdom, but many lineages feature evolutionarily asocial species. These solitary species may have evolved to conserve energy in food-sparse environments. However, the mechanism by which metabolic changes regulate social affinity is not well investigated. In this study, we used the Mexican tetra (*Astyanax mexicanus*), which features riverine sighted surface (surface fish) and cave-dwelling populations (cavefish), to address the impact of metabolic shifts on asociality and other cave-associated behaviors in cavefish, including repetitive turning, sleeplessness, swimming longer distances, and enhanced foraging behavior. After 1 month of ketosis-inducing ketogenic diet feeding, asocial cavefish exhibited significantly higher social affinity, whereas social affinity regressed in cavefish fed the standard diet. The ketogenic diet also reduced repetitive turning and swimming in cavefish. No detectable behavioral shifts were found regarding sleeplessness and foraging behavior, suggesting that other evolved behaviors are not regulated by ketosis. We further examined the effects of the ketogenic diet via supplementation with extragenic ketone bodies, revealing that ketone bodies are pivotal molecules associated with social affinity. Our study indicated that fish that evolved to be asocial remain capable of exhibiting social affinity under ketosis.

Introduction

Wild animals experience frequent fasting because of daily, seasonal, and yearly changes in food availability. Physiologically, fasting can increase the secretion of appetite-related hormones (e.g., ghrelin, peptide Y, orexin) and induce a metabolic shift into nutritional ketosis (McCue, 2010). Concerning behavioral outputs, fasting also induces shifts including boldness in foraging involving risk-taking (Padilla et al., 2016) and a shift from avoiding to approaching prey (Filosa et al., 2016). Interestingly, fasting also induces non-foraging–related behaviors including aggression toward cohorts (Fokidis et al., 2013; Solianik et al., 2016) and engagement in social dominance (Nakajo et al., 2020). These non-foraging behaviors could be evoked by the metabolic changes that occur in a state of nutritional ketosis instead of the increased production of appetite-related hormones. However, it is not fully understood whether ketosis itself in the absence of hunger drives these non-foraging behaviors. Such knowledge will open a path to understanding the effects of different dietary intakes on changing environments, such as switching from ketosis-inducing very low-carbohydrate diets to glycolysis-inducing carbohydrate-rich diets or vice versa.

Recently, the ketosis-inducing ketogenic diet (KD), which contains a high amount of fat, sufficient protein, and very low amount of carbohydrates, gained popularity among humans because of its neuroprotective and anti-inflammatory effects without effects on appetite-related hormone levels (Deemer et al., 2020; Ludwig, 2020; Sumithran et al., 2013). The KD is an effective treatment for refractory seizures, and there is some evidence that it may be beneficial for other nervous system-based disorders such as Alzheimer's disease, Perkinson's disease, and autism (Lee et al., 2018; McDonald and Cervenka, 2018; Phillips et al., 2018; Ruskin and Masino, 2012). Because modern humans evolved to acquire resistance to starvation (Bellisari, 2008), our body physiology and behavioral tendencies possibly evolved to accommodate drastic metabolic changes. However, the major molecular mechanisms for these positive are largely unknown (Ludwig, 2020; Qin et al., 2021). We were therefore motivated to explore the effects of metabolic shifts, particularly from glycolysis to ketosis, on behavioral

outputs such as social affinity using a single species consisting of typical and starvation-resistant

populations.

A suitable model system for this purpose is the Mexican cavefish (*Astyanax mexicanus*). *A.*

mexicanus has emerged as a model system of diverse aspects of evolution and development, including

those with relevance to human medicine, e.g., cataract formation, diabetes, albinism-related syndrome,

and insomnia (Aspiras et al., 2015; Bilandžija et al., 2018, 2013; Duboué et al., 2012, 2011; Jaggard et

al., 2017; Keene et al., 2016; Ma et al., 2014; McGaugh et al., 2014; Riddle et al., 2018; Rohner et al.,

2013; Strickler et al., 2007). *A. mexicanus* consists of surface riverine epigean (surface fish) and cave-

dwelling hypogean (cavefish) populations. Cavefish diverged from their surface-dwelling relatives

20,000–200,000 years ago (Fumey et al., 2018; Herman et al., 2018), and they have subsequently

evolved many distinct morphological and behavioral phenotypes in the food-sparse cave environment,

including eye regression/loss, pigment reduction, increased mechanosensory lateral line activity,

adherence to vibration stimuli, sleeplessness, hyperactivity, repetitive circling, and reduced social

affinity (Iwashita and Yoshizawa, 2021; Keene et al., 2016; Yoshizawa, 2015; Yoshizawa et al., 2018).

Compared to cavefish, surface fish exhibit typical teleost phenotypes, including typical eyed and

pigmented morphologies, no strong adherence to vibration stimuli, nocturnal sleep patterns, and social

affinity. Many cavefish traits are believed to have evolved to adapt to food-sparse dark environments.

Indeed, wild cavefish were estimated to be exposed to approximately 6 months of food-sparse conditions

annually (Espinasa et al., 2021), and they are likely to have the ability to withstand starvation via

increased fat storage (Aspiras et al., 2015), slower weight loss during starvation (Huppop, 1986),

reduced energy-costing circadian activities, and lack of eyes (Moran et al., 2015, 2014).

Concerning social-like behavior, cavefish exhibit no detectable schooling behavior (Kowalko et

al., 2013; Patch et al., 2020; Pierre et al., 2020) or hierarchal dominance (Elipot et al., 2013). By

contrast, surface fish school/shoal with cohorts and model fish (Kowalko et al., 2013) exhibit group

hierarchical dominance (Elipot et al., 2013). Because social behaviors in many fish (*e.g.*, zebrafish) are

promoted by visual stimuli, blind cavefish might not express social-like activities because of the absence

of visual acuity. However, a recent detailed study illustrated that surface fish exhibit a high level of

social-like nearby interactions (one-by-one affinity) in the dark, and these of which were promoted by

mechanosensory lateral line inputs (Iwashita and Yoshizawa, 2021). Interestingly, blind cavefish

displayed much lower levels, albeit significant, of nearby interactions than surface fish (Iwashita and

Yoshizawa, 2021). Further, cavefish exhibited plasticity in the level of nearby interactions in which they

increased plasticity in a familiar environment in comparison with an unfamiliar environment (Iwashita

and Yoshizawa, 2021), which is similar to the findings in patients with autism (Helt et al., 2020; Runco

et al., 1986).

Thus far, similarities between cavefish and patients with autism have been investigated in terms

of gene regulation- and innate behavior-profiles. First, the cavefish gene expression profile is closer to

that of patients with autism than to that of other known model systems (cavefish and surface fish

transcriptomes exhibited the same directional gene expression changes observed in the brains of patients

with autism [>58.5% of 3152 cavefish orthologs]). Conversely, other proxy systems (BTBR mice

[classic autism model] and *shank3* knockout mice) exhibit much less overlap (<11%) (Lee et al., 2019;

Provenzano et al., 2016; Yoshizawa et al., 2018). Second, cavefish's evolved behaviors—asociality,

repetitive behavior, sleeplessness, higher swimming activity, adherence to a particular vibration

stimulus, and higher anxiety-related plasma cortisol levels—are similar to those in patients with autism

(Yoshizawa et al., 2018). Third, cavefish and human ancestors are starvation-resistant, and they could

share some metabolic pathways (Aspiras et al., 2015; Bellisari, 2008; Huppop, 1986; Riddle et al.,

2018). These similarities and a fact that KD feeding increases sociality in patients with autism

(Evangeliou et al., 2003; Lee et al., 2018; Li et al., 2021; Napoli et al., 2014) prompted us to study the

effects of ketosis on social affinity in asocial cavefish.

In this study, we assessed the effects of the KD on an evolutionarily asocial cave population of *A.*

mexicanus. The time-course experiment revealed that 1 month of KD feeding promoted and sustained

the juvenile level of nearby interactions, whereas control diet (CD)-fed cavefish exhibited diminished

nearby interactions. KD feeding also reduced repetitive turning and swimming activity, which are the

hallmarks of the autism-like condition. However, the effects of the KD were limited. For example,

sleeplessness and high adherence to a particular vibrating stimulus were not detectably changed under

the 1-month KD treatment. To reveal the molecular basis of the effects of the KD, we provided

supplementation with a major ketone body, beta-hydroxybutyrate (BHB). This experiment indicated that

BHB supplementation promoted nearby interactions and reduced repetitive turning, covering the major

effect of the KD.

Overall, ketosis appears to be capable of significantly shifting the asociality of evolved cavefish

toward the surface fish phenotype, providing new insights into the contribution of the diet to the

evolution of behavior.

Results

From our observation in their wild habitat (the Mexican cave Pachón, Supplemental Movie 1), cavefish swam slower and remained near each other more frequently than the lab population. Because the cave environment has a limited diet compared to that of the surface, we predicted that cavefish experience frequent ketosis induced by fasting.

To avoid appetite-related behavior, we developed a ketogenic diet (KD) based on a human milk formula (KetoCal3:1® with Zeigler zebrafish standard irradiated diet at a 5:1 weight ratio [nutritionally complete, ketogenic medical food; Nutricia North America, Inc. Gaithersburg, MD, USA]; Table 1; Materials and Methods). To test whether our KD could induce a shift in the balance of ketone body and glucose levels, we measured blood ketone and glucose levels after chronic dietary treatment. Three-month-old fish (juvenile–young adult stage) were used in this study because many adult-type behaviors of cavefish emerge in this stage, including higher adherence to a vibration stimulus (vibration attraction behavior [VAB]) (Yoshizawa et al., 2010), less social affinity, and longer swimming distances compared to the findings in surface fish. After 5 weeks of KD feeding, both ketone and glucose concentrations were decreased compared to the findings in CD-fed fish (KetoCal3:1 and Zeigler zebrafish diet at a 1:5 weight ratio; Table 1; Fig. 1A–C). For both diets, surface fish exhibited a significantly higher ketone body level than cavefish (Fig. 1B), whereas cavefish exhibited a higher glucose level than surface fish (Fig. 1C). The glucose ketone index (GKI $= \frac{glucose\ (mM)}{ketone\ (mM)}$), a medical index proposed for brain cancer diagnosis (Meidenbauer et al., 2015), was lower in surface fish than in cavefish, and the value was reduced under KD feeding in both surface fish and cavefish compared to that in their CD-fed counterparts (Fig. 1D). This result indicated that KD feeding more strongly reduced glucose levels than ketone body levels, resulting in a lower GKI in KD-fed fish than in CD-fed fish (Fig. 1D) and suggesting that KD feeding could shift the metabolic state from glycolysis toward ketosis.

161   Table 1. Nutrient composition of each diet used in the study

%	Brine shrimp	Zeigler zebrafish standard diet	KetoCal3:1	Control diet (CD)	Ketogenic diet (KD)
Protein	58.4	54.0	15.3	47.6	21.8
Lipid/fat	14.7	14.4	67.7	23.3	58.8
Carbohydrate	5.2	11.6	7.2	10.9	7.9
Ash	7.2	15.5	NA	NA	NA
Calories (kcal/g)	5.9	3.9	7.0	4.4 (20% w/v in agar)	6.5 (20% w/v in agar)

(Paffenhöfer, 1967; Panda, 2016)

Regarding this dietary treatment, we first examined its ontogenic (developmental) effects on

collective social-like behavior (Iwashita and Yoshizawa, 2021). Many adult behaviors emerge in the

transition from juvenile to young adult (adolescent) in 3–4-month-old *A. mexicanus* fish, including

foraging behavior, VAB (Yoshizawa, 2015; Yoshizawa et al., 2010), adult-type regulation of sleep

(independent from catecholamine) (Duboué et al., 2012; Jaggard et al., 2020, 2017; Yoshizawa et al.,

2015), and the collective behavior in young adults (under higher Reynold's number; Iwashita and

Yoshizawa, 2021). We therefore investigated the shift of collective behavior in 3–4-month-old fish using

criteria based on the vicinity of two fish (≤5 cm) and duration of nearby interactions (≥4 s) during

tracking in a four-fish group (Iwashita and Yoshizawa, 2021) (Fig. 2B). At 3 months old, ('Pre-

treatment' in Fig. 2), surface fish exhibited social-like nearby interactions for 17.0 ± 4.4 s (Fig. 2C) and

3.1 ± 0.4 bout number of nearby interactions (Fig. 2D) during the 5-min assay. Conversely, cavefish

exhibited an approximately 50% shorter interaction duration (8.3 ± 1.5 s; Fig. 2C) and a smaller bout

number of interactions (1.8 ± 0.3; Fig. 2D). To track the effect of the KD treatment, fish were fed the

KD for 5 weeks, followed by CD feeding during weeks 6–9 to assess the persistence of the effects of the

KD (Fig. 2A, C, and D).

The nearby interactions of surface fish did not differ between CD and KD feeding (Fig. 2C and

D). By contrast, the nearby interactions of cavefish were significantly decreased by CD feeding

compared to the effects of KD feeding in weeks 4 and 5 (Fig. 2C and D), and interactions remained

depressed through week 9 by CD feeding. This effect of the KD diet on nearby interactions did not

persist. After KD deprivation and CD feeding, the nearby interactions of KD-fed cavefish were

indistinguishable from those of CD-fed fish (6–9 weeks, Fig. 2C, and D), suggesting that KD has a

promotive/supportive effect on collective behavior in genetically asocial cavefish.

To investigate in detail whether KD-promoted nearby interactions have social-like properties, we

explored swimming speed, which is an indicator of the movement of fish in the vicinity of each other.

Fish are more likely to have an opportunity to express affinity toward each other at a slower swimming

speed (Iwashita and Yoshizawa, 2021). Indeed, surface fish moved slower during nearby interactions

than during the non-nearby interaction period (Iwashita and Yoshizawa, 2021) (Fig. 3A). Consistently,

KD-fed cavefish swam slower during the nearby interaction period than during the non-nearby

interaction period ('5 weeks;' Fig. 3B). Overall swimming speed was also slower in the KD group than

in the CD group ('5 weeks;' Fig. 3B). These findings indicated that KD-fed cavefish exhibited more

social-like nearby interactions with a similar speed profile as surface fish. In surface fish, there was no

major difference in swimming speed profiles between CD and KD feeding ('5 weeks;' Fig. 3A). To

address whether the slower swimming speed was sufficient to increase nearby interactions, we tracked

the total swimming distance within 5 min from pre-treatment to week 9 of feeding (Supplementary Fig.

1). KD-fed cavefish exhibited a significantly shorter swimming distance (slower swimming speeds)

from the first week of feeding (Supplementary Fig. 1), which was much earlier than when the higher

level of nearby interactions emerged (weeks 4–5). This result suggests that although a slower swimming

speed is associated with nearby interactions (Fig. 3) and KD feeding reduced swimming speeds in

cavefish, a slower speed itself is not sufficient to induce nearby interactions.

Repetitive turning is frequently observed in an antagonistic relationship with nearby interactions

in cavefish (Iwashita and Yoshizawa, 2021). That is, individuals with few nearby interactions frequently

exhibit a high level of turning bias or 'repetitive turning.' Accordingly, CD-fed cavefish with few nearby

interactions (4–5 weeks) exhibited significantly higher turning bias than KD-fed cavefish (Fig. 4A, B).

KD-fed cavefish displayed a comparable level of balanced turning as surface fish (close to a score of '1'

in Fig. 4B). In summary, these results suggest that KD feeding could reduce repetitive turning while

maintaining longer nearby interactions.

Tracking behavioral changes each week (Fig. 2) may result in confounding factors such as fish

remembering the recording environment. To clarify whether our results captured the genuine effects of

KD feeding, we repeated 4–5-week dietary treatment in a new set of fish (Supplementary Fig. 2).

Similarly, surface fish did not exhibit a detectable change in the duration and number of nearby

interactions between CD and KD feeding (Supplementary Fig. 2A, B). By contrast, CD-fed cavefish

displayed fewer nearby interactions, whereas the level of nearby interactions was retained in KD-fed

cavefish, resulting in a higher level of nearby interactions in KD-fed cavefish (Supplementary Fig. 2A,

B). In this repeated experiment, the results for swimming distance and repetitive turning were also

similar to those in the previous experiment; specifically, CD-fed cavefish swam longer distances than

KD-fed cavefish (Supplementary Fig. 2C), whereas surface fish did not exhibit a detectable shift in

swimming distance between the CD and KD groups. Additionally, CD-fed cavefish displayed a high

level of turning bias, whereas KD-fed cavefish exhibited balanced turning (Supplementary Fig. 2D).

We then explored other changes induced by KD feeding, including changes in sleep, 24-h

swimming distance, and adherence to a vibrating stimulus, which are distinct between surface fish and

cavefish. Cavefish largely exhibit reduced sleep duration and swim almost all day, perhaps to find

nutrients in the food-sparse environment (Duboué et al., 2012, 2011; Jaggard et al., 2017; Yoshizawa et

al., 2015). After 5 weeks of dietary treatment on the 3 – 4 months old fish, both surface fish and cavefish

exhibited a shorter sleep duration than observed before treatment regardless of the diet (Fig. 5A,

particularly at night), suggesting growth between 3 – 4 and 4 – 5 months old exerted a negative effect on

the sleep duration. However, there was no detectable difference between CD and KD feeding.

Animals' sleep is usually fragmented, involving repeated sleep/awake cycles during the night

(diurnal animals) or day (nocturnal animals) (Campbell and Tobler, 1984). Then, the structure and

regulation of sleep are typically analyzed according to the number of events (bout). Our detailed sleep

analysis illustrated that KD-fed cavefish displayed fewer sleep bouts during the night than CD-fed

cavefish (Fig. 5B). However, the number of sleep bouts did not differ between CD and KD feeding (5

weeks; Supplementary Fig. 3). Overall, the sleep phenotype was not dramatically changed by KD

feeding, and cavefish exhibited a shortened sleep duration.

The sleep duration is negatively correlated with the 24-h swimming distance (Yoshizawa et al.,

2015). Cavefish displayed overall higher activities, which was consistent with previous findings

(Duboué et al., 2011; Yoshizawa et al., 2015) and consistent with the findings of longer swimming

distance in the nearby interaction assay (Fig. 5C). CD-fed cavefish swam longer distances than KD-fed

cavefish. Surface fish, in contrast, did not exhibit a detectable difference in swimming distance between

KD and CD feeding. Overall, the KD treatment induced little changes in sleep behaviors in both surface

and cavefish.

In general, the KD is assumed to induce ketosis without increasing appetite. We then checked the

shift of foraging behavior under KD feeding. Cavefish evolutionarily exhibit increased foraging

behavior, termed VAB, in which fish adhere to a particular vibration stimulus (35–40 Hz) in the dark

(Yoshizawa et al., 2010). VAB is advantageous for prey capture in the dark. Cavefish and surface fish

did not exhibit a detectable difference in VAB between CD and KD feeding, whereas VAB was

significantly increased during 1 month of growth (pre-treatment vs. 5 weeks; Supplementary Fig. 4A).

The swimming distance in 3-min VAB assays was decreased in KD-fed cavefish, consistent with the

nearby interaction and sleep studies (5 weeks; Supplementary Fig. 4B). In summary, the VAB analysis

indicated that KD feeding did not increase a foraging behavior.

Although the KD diet induced significant changes in some behavioral outputs, it suppressed

growth during treatment. The average weights of KD-fed surface fish and cavefish were 55.5 % and

69.9 % of those in their CD-fed counterparts, respectively (5 weeks; Fig. 6B). The standard length of

KD-fed surface fish was also significantly reduced (5 weeks; Fig. 6A).

Are these behavioral and growth changes induced by ketosis? The KD contains high amounts of

fat and other ingredients. This question motivated us to test the molecular basis of the effects of the KD

by adding major ketosis metabolites to the standard diet.

In humans, the KD induces ketosis, in which the liver releases the major ketone body, beta-

hydroxybutyrate (BHB), via beta-oxidation of fat (Evans et al., 2017). Instead of supplying a massive

amount of fat using the KD, BHB (sodium salt form of racemic BHB: 50% L-form and 50% D-form;

only the D-form is considered to be biologically active) might be responsible for the majority of effects

observed after KD feeding. With this idea, sodium salt BHB was provided to both surface fish and

cavefish for 4 weeks. The result indicated that the BHB supplemental diet promoted nearby interactions

in cavefish (Fig. 7A and B) and reduced the duration of nearby interactions in surface fish (Fig. 7A).

Turning bias tended to be reduced by BHB supplementation in cavefish, although significance was not

achieved (Fig. 7C). Swimming distance was not reduced in surface fish or cavefish (Fig. 7D). The body

growth of BHB-treated surface fish and cavefish was comparable to that in control fish (standard length

and weight; Supplementary Fig 5A and B, respectively). The night-time sleep and VAB did not exhibit

detectable differences between control and BHB treatment (Supplementary Fig. 6A and B, respectively)

In summary, BHB treatment covered many effects of KD treatment did, including changes in

social interactions and repetitive turning. BHB had fewer negative effects on growth and swimming

activities, suggesting that ketone bodies are responsible for the 'positive' effects of KD feeding.

Discussion

In this study, we examined the behavioral shifts induced by KD feeding and BHB supplementation. Ketosis is expected to occur frequently in wild animals because of a failure to find food (fasting) or an absence of carbohydrate inputs/synthesis (available nutrients). Certain levels of socialness can be beneficial to animal species for mating and finding food. Under KD feeding, cavefish maintained their juvenile level of nearby interactions until the treatment ended (5 weeks). Nearby interactions were then reduced to an indistinguishable level from the control levels within 1 month after stopping KD feeding. Surface fish exhibited a higher number of nearby interactions than cavefish, and no detectable difference was observed in nearby interaction levels between CD- and KD-fed surface fish. KD feeding also effectively reduced repetitive turning in cavefish, whereas CD-fed treated cavefish exhibited a high level of repetitive turning. Under KD feeding, both surface fish and cavefish significantly reduced swimming distances during development, during which the swimming distance typically becomes longer. There were no detectable changes in sleep and foraging behavior (VAB) under 1 month of KD feeding. These patterns in behaviors and growth were not changed in two replicated experiments (social affinity and repetitive turning), supporting the consistency of the observed effects under KD feeding. Finally, the major KD metabolite, BHB, could cover the KD effect, indicating the ketone body has the pivotal role on this treatment.

Effects of the KD on blood ketone levels and body growth

Under 4–5 weeks of KD feeding, blood ketone and glucose levels were reduced compared to the effects of the CD in both surface fish and cavefish, contradicting our expectation that ketone levels would be higher in the KD group. However, GKI (Meidenbauer et al., 2015) was significantly lower under KD feeding than under CD feeding. These significant changes in GKI in both surface fish and cavefish suggest that the metabolic condition was shifted toward ketosis by KD feeding. In general, cavefish had a higher GDI than surface fish under both diets, suggesting that the cavefish physiology

was constitutively biased toward glycolysis. For example, blood glucose levels in cavefish under KD

feeding were similar to those in surface fish under CD feeding, whereas cavefish had 3-fold lower

ketone levels than surface fish under CD feeding, resulting in a higher GKI even under KD feeding.

KD feeding for 4–5 weeks also resulted in slowed body growth. This growth retardation has been

observed in patients with epilepsy chronically fed a KD (Coppola et al., 2010; Napoli et al., 2014), and

these results were consistent with our observations in KD-fed fish. The detailed molecular/physiological

mechanisms are largely unknown, but this study identified BHB as a potential candidate mediating KD-

associated phenotypes (see below).

Effects of ketones in the TCA cycle and epigenetics in the brain

In mammals, KD feeding causes a 'starvation'-like state, causing the liver to release ketone

bodies into the bloodstream. BHB is the major ketone body produced by the liver through beta-

oxidation. The gut epithelia also absorb and circulate ketone bodies from the diet and/or gut microbiota.

Both liver- and gut-derived ketone bodies can cross the blood–brain barrier and exert two functions: (i)

inhibit histone deacetylase, which influences epigenetic regulation and induces gene expression in

neurons; and (ii) act as a general energy source that is converted into acetyl-CoA to fuel the aerobic

TCA cycle in neurons. Both pathways have the potential to alter brain function. The facts that cavefish

easily tolerate high blood glucose levels, at which surface fish was paralyzed (Riddle et al., 2018), and

Wnt signaling is upregulated in cavefish, potentially resulting in high glycolytic activity as humans do

(Vallée and Vallée, 2018; Yoshizawa et al., 2018), support the aforementioned hypothesis that cavefish

exhibit high blood glucose levels and generate energy via glycolysis. Also, ketone bodies can also

promote behavioral shifts by changing the epigenetic state by inhibiting histone deacetylase

(Krautkramer et al., 2017; Szyf, 2015). Histone deacetylase inhibition increases gene expression in

general. This possibility is supported by the fact that cavefish have more downregulated genes (2913

genes, $\log_2 < -1.0$) than upregulated genes (1643 genes, $\log_2 > 1.0$) in the transcriptome at 72 h post-

fertilization (Gross et al., 2013; Yoshizawa et al., 2018). In addition, more methylated loci are found in

the eye genes of cavefish than in surface fish, which could also be true for other tissues (Gore et al.,

2018), and most of these methylated gene loci were downregulated. The brains of patients with autism

are also expected to be hypermethylated, resulting in a transcription-less condition (Zhu et al., 2014).

Therefore, these two pathways, namely metabolism and epigenetics, were highlighted as possible targets

of ketone bodies during behavioral shifts under ketosis. Future research should address these

possibilities to clarify the metabolism-based evolution of behavior (*cf.* (Qin et al., 2021)

Ontogeny of nearby interactions and the KD

In this study, 3-4 months-old cavefish exhibited a weak but detectable level of nearby

interactions (social affinity), and this social affinity decayed under CD feeding. Interestingly, KD-fed

cavefish and surface fish fed either diet maintained a similar level of nearby interactions during 5 weeks

of dietary feeding. The reduction of nearby interactions in CD-treated cavefish can be explained by (1)

quicker exhaustion under CD feeding (aerobic ketosis produces more adenosine triphosphate than

anaerobic glycolysis), (2) greater anxiety in the recording environment (Iwashita and Yoshizawa, 2021),

and (3) less social motivation. The first explanation is unlikely because CD-fed cavefish swam longer

distances than KD-fed cavefish. The higher level of anxiety could explain the findings because cavefish

exhibited increased repetitive turning, which is related to higher anxiety in mammals (Langen et al.,

2011). In addition, cavefish displayed fewer nearby interactions in an anxiety-associated unfamiliar

environment in prior research (Iwashita and Yoshizawa, 2021). In the future, the anxiety level should be

monitored using plasma cortisol levels in the future (Gallo and Jeffery, 2012). Less motivation regarding

social affinity is also a possible cause, and this variable can be monitored by assessing activities in social

decision-making networks including the preoptic area, nucleus accumbens, and striatum (O'Connell and

Hofmann, 2012, 2011). Explanations (2) and (3) are not mutually exclusive, and co-occurrence is

possible. These possibilities will be assessed in our future study.

Possible target system for ketosis

Under KD feeding and BHB treatment, we observed increased social affinity and reduced

repetitive turning and swimming distances. However, we detected no changes in sleep and VAB. From

the knowledge of neurotransmitters and their associated behaviors, these behavioral phenotypes

indicated the possible involvement of the dopaminergic and serotonergic systems but less involvement

of the cholinergic, orexin/hypocretinergic, histaminergic, or adrenergic system. The mechanism by

which ketosis or ketone bodies more strongly affect the dopaminergic and serotonergic systems than the

other systems is unclear. However, the findings that the dopaminergic and serotonergic systems are

sensitive to ketosis in terms of evolution are extremely interesting in its mechanism and for future

therapy applications.

Ketosis in the cave environment

Cave-dwelling animals usually experience less temperature fluctuation and fewer dietary inputs

(Culver and Pipan, 2009), but these features can vary. The diets of cave-dwelling animals in the dry

season (approximately 6 months/year) could be organic matter in the pool bottoms, bat guano (larger

adults), or small crustaceans (smaller fish), whereas food is sparse in the rainy season (approximately 6

months/year) (Espinasa et al., 2021, 2017). These available diets contain extremely low amounts of

carbohydrates, and they can be high in protein and fat (e.g., crustaceans). Although some amino acids,

lactate, and glycerol can be used for glucose synthesis in fish (Polakof et al., 2012), cavefish are

expected to be exposed to carbohydrate-deprived diets or frequent fasting and therefore frequent ketosis.

In prior research, wild cavefish swam slower and exhibited similar social affinity as observed in KD-fed

cavefish in this study (Movie 1). Although these observations and dietary inputs suggested that wild

cavefish may be under frequent ketosis, recent multiple reports indicated that cavefish may be under

anaerobic glycolysis to adapt to anaerobic cave water conditions because of the approximately 20%

lower oxygen level in cave pools (Boggs et al., 2022; van der Weele and Jeffery, 2022). In addition,

cavefish tend to store lipids instead of using them through the enhanced PPARγ pathway (Xiong et al.,

2022). These expectations of low ketosis appear to contradict expectations in the wild—starved ketosis

conditions. However, it appears to fit well with the findings in cavefish, namely high blood glucose and

low ketone levels even under KD feeding in this study, as well as higher GKI in cavefish than in surface

fish in both the CD and KD groups (Fig. 1D). Cavefish appear to have evolved to maintain a high GKI

(high blood glucose and low ketone levels); therefore, the physiology of cavefish may allow them to

survive in the low-oxygen condition by using anaerobic glycolysis. KD-fed cavefish behave similarly as

wild cavefish because the balance between ketosis and glycolysis could reach a similar level as that in

the wild after KD feeding. By contrast, if cavefish are fed a typical carbohydrate-rich lab fish diet, it

may overactivate glycolysis and result in a higher GKI, which may lead to reduced social affinity and

increased repetitive circling and swimming distance. The future use of a pharmacological glycolysis

inhibitor (e.g., 2-deoxy-D-glucose; Yao et al., 2011) can reveal the relationship between GKI and

cavefish behaviors.

Summary statement

Solitary animals surprisingly share a set of dysregulated genes and behavioral outputs. In this

study, we demonstrated that a diet that induces ketosis shifts these behaviors toward the surface fish

phenotype regardless of the presence of many dysregulated genes. In addition to the gene therapy

approach, ketone body-based treatment may open a path for sustainable and less toxic therapy for

multigenic psychiatric disorders, including autism, although the target pathways remain unclear.

Concerning the genetics of behavior, differentially expressed metabolic genes have been largely

overlooked because it was difficult to interpret. Because mitochondria-based disorders are highlighted in

neuroscience (Chauhan et al., 2012; Rajasekaran et al., 2015), the balance between glycolysis and

ketosis could be the starting point for identifying a therapeutic target. The known evolved behaviors

should be also revisited by investigating whether the metabolic shift promotes the variations of

behaviors.

Materials and Methods

Fish maintenance and rearing in the lab

A. mexicanus surface fish used in this study were the laboratory-raised descendants of original collections created in Balmorhea Springs State Park in Texas in 1999 by Dr. William R. Jeffery. Cavefish were laboratory-raised descendants originally collected from Cueva de El Pachón (Pachón cavefish) in Tamaulipas, Mexico in 2013 by Dr. Richard Borowsky.

Fish (surface fish and Pachón cave populations) were housed in the University of Hawaiʻi at Mānoa *Astyanax* facility with temperatures set at $21 \pm 0.5°C$ for rearing, $24 \pm 0.5°C$ for behavior experiments, and $25 \pm 0.5°C$ for breeding (Elipot et al., 2014; Yoshizawa et al., 2015). Lights were maintained on a 12-h:12-h light:dark cycle (Elipot et al., 2014; Yoshizawa et al., 2015). For rearing and behavior experiments, the light intensity was maintained at 30–100 Lux. Fish husbandry was performed as previously described (Elipot et al., 2014; Keene et al., 2016; Yoshizawa et al., 2015). Fish were raised to adulthood and maintained in standard 42-L tanks in a custom-made water-flow tank system. Adult fish were fed a mixed diet to satiation twice daily starting 3 h after the lights were turned on (Zeitgeber time 3 [ZT3] and ZT9; Zeigler Adult zebrafish irradiated diet, Zeigler Bros, Inc, Gardners, PA; TetraColor Tropical Fish Food Granules, Tetra, Blacksburg, VA, USA; Jumbo Mysis Shrimp, Hikari Sales USA, Inc., Hayward, CA, USA). All fish in the behavioral experiments were between 2.5 and 5 cm in standard length and between 3 and 12 months old. Fish ages were stated in each experiment. All fish care and experimental protocols were approved under IACUC (17-2560) at the University of Hawaiʻi at Mānoa.

Ketogenic diet

To prepare the KD, we used a mixture of a human KD (KetoCal3:1) and zebrafish standard diet (adult zebrafish irradiated diet) in a 5:1 ratio. The gross caloric amounts were 6.99 kcal/g for KetoCal3:1 and 3.89 kcal/g for the zebrafish diet. Regarding the CD, we used the same KetoCal3:1 and zebrafish

irradiated diet but mixed at a 1:5 ratio. The KetoCal3:1 powder and ground zebrafish irradiated diet were

mixed in the aforementioned ratios and solidified with 1% agar at a final concentration of 20% w/v (2 g

of mixture in 10 mL of 1% agar). After solidification, both KD and CD agar was cut into 3-mm^3 cubes,

and each four-fish group was given 1–2 pieces every morning (ZT 0:00–3:30) and afternoon (ZT 8:00–

12:00). The fish were fed ad libitum in each feeding and the remaining amount was removed 1 h after

feeding using a pipette.

<u>Behavior assays</u>

The protocol for social-like nearby interactions was described previously (Iwashita and

Yoshizawa, 2021). Briefly, four fish raised in a home tray (15.6 × 15.6 × 5.7 cm^3 Ziploc containers, S.

C. Johnson & Sons, Inc, Racine, WI, USA) were released in a recording arena (49.5 × 24.2 × 6.5 cm^3)

with a water depth of 3 cm on the stage of a custom-made infrared (IR) back-light system within a

custom-built black box (75 × 50 × 155 cm, assembled with polyvinyl chloride pipe frame and covered

by shading film). The IR back-light system was composed of bounce lighting of IR LED strips

(SMD3528 850 nm strip: LightingWill, Guang Dong, China). The video was recorded at 20 frame/s

using VirtualDub2 software (build 44282; http://virtualdub2.com/) with the x264vfw codec for 6 min,

and the last 5 min were used for the analysis. After the recording, the fish were returned to the home

tray. The X-Y coordinates of each fish were calculated using idTracker software (Pérez-Escudero et al.,

2014) after the video image was processed for background subtraction using ImageJ (Iwashita and

Yoshizawa, 2021). This X-Y coordinate was also used for the turning bias analysis. The duration and

number of nearby interactions and swimming speed during and after nearby interaction events were

calculated using custom-made MATLAB script (MathWorks Inc., Natick, MA, USA) (Iwashita and

Yoshizawa, 2021).

The turning bias rate was calculated as $\frac{Nl}{Ns}$, where Ns and Nl represent a smaller (Ns) or larger

(Nl) number of left or right turns. This turning bias rate indicates the extent to which fish turning is

biased to the left or right, and ranging from '1' (L-R balanced) to exponential (L or R biased). The

numbers of left or right turns were calculated as changes in the angles of swimming directions in every

five frame window (0.25 s) as described previously (Iwashita and Yoshizawa, 2021). An automatic

calculation of the total number of the left or right turns is implemented in the aforementioned homemade

MATLAB script.

Analyses of sleep and swimming distance were described previously (Yoshizawa et al., 2018,

2015). Briefly, fish were recorded in a custom-designed 10.0-L acrylic recording chamber (457.2 ×

177.8 × 177.8 mm^3 and 6.4 mm thick) with opaque partitions that permit five individually housed fish

per tank (each individual chamber was 88.9 × 177.8 × 177.8 mm^3). The recording chamber was

illuminated with a custom-designed IR LED source the light-controlled room on a 12-h:12-h cycle. The

room light was turned on at 7:00 am and turned off at 7:00 pm each day. Behavior was recorded for 24 h

after overnight (18–20 h) acclimation beginning 1–2 h after turning the light on (ZT1–2). Videos were

recorded at 15 frames/s using a USB webcam with an IR high-pass filter. Videos were captured by

VirtualDub2 software with the x264vfw codec and subsequently processed using Ethovision XT

(Version 16, Noldus Information Technology, Wageningen, Netherlands). Water temperature was

monitored throughout the recordings, and no detectable differences were observed during the light and

dark periods (24.0 ± 0.5°C). The visible light during behavior recordings was approximately 30–100

Lux.

The tracking parameters for detection were as follows: the detection was set to 'subject brighter

than background' and brightness contrast was set from 20 to 255; the current frame weight was set to 15;

the video sample rate was set to 15 frames/s; and pixel smoothing was turned off. We monitored sleep,

activity, and arousal thresholds via protocols previously established for *A. mexicanus* (Yoshizawa et al.,

2015). The X-Y coordinates of each fish were subsequently processed using custom-written Perl

(v5.23.0, www.perl.org) and Python scripts (3.8).

We assayed VAB as described previously (Yoshizawa et al., 2015, 2012, 2010). Briefly, fish

were permitted to acclimate for 4–5 days prior to the assay in a cylindrical assay chamber (325-mL glass

dish, 10 cm in diameter and 5 cm in height, VWR, Radnor, PA, USA) filled with conditioned water (pH

6.8–7.0; conductivity 600–800 μS). During the assays, vibration stimuli were created using a glass rod

that vibrated at 40 Hz. The number of approaches to the vibrating rod was video recorded during a 3-min

period under infrared illumination. The number of fish approaches in a 1.3-mm radius from the vibrating

glass rod were analyzed using the X-Y coordinate of each fish head detected by the trained DeepLabCut

model (Mathis et al., 2018; Fernandez et al., *submitted*)

<u>Blood ketone and glucose measurements</u>

The fish used to measure ketone and glucose levels were the siblings of the fish used for the

behavioral assay. Eight fish were used for each treatment group (CD vs. KD, surface fish vs. cavefish;

four groups in total). All blood samples were collected 2 h after feeding. Fish were then deeply

anesthetized in ice-cold water, and blood was collected from the tail artery. Blood ketone and glucose

levels were measured using an Abbott Precision Xtra blood glucose & ketone monitoring system

(Abbott Laboratories, Abbott Park, IL, USA)

<u>Measurement of tissue structure</u>

Fish were anesthetized with 0.2 mg/mL ethyl 3-aminobenzoate methanesulfonate (MS-222:

MilliporeSigma, Burlington, MA, USA) in ice-cold conditioned water (pH: 7.0; conductivity: 700 μS),

and weight was measured after taking pictures with a standard camera (Pentax K-1 DSLR with 35-70

mm zoom lens, Ricoh, Tokyo, Japan). The standard body length and body depth were measured using

ImageJ.

<u>Statistical analysis</u>

Regarding the power analysis, we designed our experiments based on three-way repeated-

measures ANOVA with a moderate effect size (f = 0.25), alpha-error probability of 0.05, and power of

0.80, and the number of groups was eight (surface fish vs. cavefish × non-treated vs. treated × pre-

treatment vs. post-treatment). G*Power software (Erdfelder et al., 2009; Faul et al., 2007) estimated that

the sample size needed for this experiment was nine per group. We thus aimed to use at least 12 fish in

each group for all experiments in this study.

For statistical comparisons of our data, we performed tests including Student's *t*-test and two- or

three-way generalized linear model analyses to compare surface and cavefish, treatment and non-

treatment, and pre-treatment and post-treatment. We calculated Akaike's information criterion (AIC) for

each linear and generalized linear model and chose the model with the lowest AIC. Holm's post hoc

correction was used to understand which contrasts were significant (Holm, 1979).

Regarding replicates of experiments, we used different individuals for the replicates, namely two

biological replicates, using different individuals in each trial (*e.g.*, Fig. 2 and Supplementary Fig. 2).

There was no repeated usage of individual fish excluding the time-course experiment (Fig. 2). For the

experiments measuring sleep and VAB in addition to nearby interactions and turning bias (Figs. 5–6,

Supplementary Figs. 2–4), we used two biological replicates and confirmed that the averages of

experimental data did not largely differ from each other. We then merged the data acquired in two

biological replicates and presented the data as a single set of results.

The aforementioned calculations were performed using R version 4.0.4 software (packages of

car, lme4, and lmerTest) (Bates et al., 2015; Fox and Weisberg, 2019; Kuznetsova et al., 2017), and all

statistical scores are available in Supplementary Table 1, the figure legends, or the text.

527
528
529

References

Aspiras AC, Rohner N, Martineau B, Borowsky RL, Tabin CJ. 2015. Melanocortin 4 receptor mutations contribute to the adaptation of cavefish to nutrient-poor conditions. *Proc Natl Acad Sci U S A* **112**:9668–9673. doi:10.1073/pnas.1510802112

Bellisari A. 2008. Evolutionary origins of obesity. *Obes Rev*. doi:10.1111/j.1467-789X.2007.00392.x

Bilandžija H, Abraham L, Ma L, Renner KJ, Jeffery WR. 2018. Behavioural changes controlled by catecholaminergic systems explain recurrent loss of pigmentation in cavefish. *Proc R Soc B Biol Sci* **285**:20180243. doi:10.1098/rspb.2018.0243

Bilandžija H, Ma L, Parkhurst A, Jeffery WR. 2013. A potential benefit of albinism in Astyanax cavefish: Downregulation of the oca2 gene increases tyrosine and catecholamine levels as an alternative to melanin synthesis. *PLoS One* **8**:e80823. doi:10.1371/journal.pone.0080823

Boggs TE, Friedman JS, Gross JB. 2022. Alterations to cavefish red blood cells provide evidence of adaptation to reduced subterranean oxygen. *Sci Rep* **12**:1–10. doi:10.1038/s41598-022-07619-0

Campbell SS, Tobler I. 1984. Animal sleep: A review of sleep duration across phylogeny. *Neurosci Biobehav Rev* **8**:269–300. doi:10.1016/0149-7634(84)90054-X

Chauhan A, Gu F, Chauhan V. 2012. Mitochondrial respiratory chain defects in autism and other neurodevelopmental disorders. *J Pediatr Biochem* **2**:213–223. doi:10.3233/JPB-120063

Coppola G, Verrotti A, Ammendola E, Operto FF, Corte R della, Signoriello G, Pascotto A. 2010. Ketogenic diet for the treatment of catastrophic epileptic encephalopathies in childhood. *Eur J Paediatr Neurol* **14**:229–234. doi:10.1016/j.ejpn.2009.06.006

Culver DC, Pipan T. 2009. The biology of caves and other subterranean habitats, The biology of habitats series. Oxford: Oxford University Press.

Deemer SE, Plaisance EP, Martins C. 2020. Impact of ketosis on appetite regulation—a review. *Nutr Res* **77**:1–11. doi:10.1016/j.nutres.2020.02.010

Duboué ER, Borowsky RL, Keene AC. 2012. β-adrenergic signaling regulates evolutionarily derived sleep loss in the Mexican cavefish. *Brain Behav Evol* **80**:233–43. doi:10.1159/000341403

Duboué ER, Keene AC, Borowsky RL. 2011. Evolutionary convergence on sleep loss in cavefish populations. *Curr Biol* **21**:671–676. doi:10.1016/j.cub.2011.03.020

Elipot Y, Hinaux H, Callebert J, Rétaux S. 2013. Evolutionary shift from fighting to foraging in blind cavefish through changes in the serotonin network. *Curr Biol* **23**:1–10.

Elipot Y, Legendre L, Père S, Sohm F, Rétaux S. 2014. *Astyanax* transgenesis and husbandry: how cavefish enters the laboratory. *Zebrafish* **11**:291–9. doi:10.1089/zeb.2014.1005

Erdfelder E, FAul F, Buchner A, Lang AG. 2009. Statistical power analyses using G*Power 3.1: Tests for correlation and regression analyses. *Behav Res Methods* **41**:1149–1160. doi:10.3758/BRM.41.4.1149

Espinasa L, Bonaroti N, Wong J, Pottin K, Queinnec E, Rétaux S. 2017. Contrasting feeding habits of post-larval and adult *Astyanax* cavefish. *Subterr Biol* **21**:1–17. doi:10.3897/subtbiol.21.11046

Espinasa L, Heintz C, Rétaux S, Yoshisawa M, Agnès F, Ornelas-Garcia P, Balogh-Robinson R. 2021. Vibration attraction response is a plastic trait in blind Mexican tetra (Astyanax mexicanus), variable within subpopulations inhabiting the same cave. *J Fish Biol* **98**:304–316. doi:10.1111/jfb.14586

Evangeliou A, Vlanchonikolis I, Mihailidou H, Vlachonikolis I, Mihailidou H, Spilioti M, Skarpalezou A, Makaronas N, Prokopiou A, Christodoulou P, Liapi-Adamidou G, Helidonis E, Sbyrakis S, Smeitink J. 2003. Application of a Ketogenic Diet in Children With Autistic Behavior: Pilot Study. *J Child Neurol* **18**:113–118. doi:10.1177/08830738030180020501

Evans M, Cogan KE, Egan B. 2017. Metabolism of ketone bodies during exercise and training: physiological basis for exogenous supplementation. *J Physiol* **595**:2857–2871. doi:10.1113/JP273185

Faul F, Erdfelder E, Lang AG, Buchner A. 2007. G*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciencesBehavior Research Methods. Springer.

pp. 175–191. doi:10.3758/BF03193146

Filosa A, Barker AJ, Dal Maschio M, Baier H. 2016. Feeding State Modulates Behavioral Choice and
Processing of Prey Stimuli in the Zebrafish Tectum. *Neuron* **90**:596–608.
doi:10.1016/j.neuron.2016.03.014

Fokidis HB, Prior NH, Soma KK. 2013. Fasting increases aggression and differentially modulates local
and systemic steroid levels in male zebra finches. *Endocrinology* **154**:4328–4339.
doi:10.1210/en.2013-1171

Fumey J, Hinaux H, Noirot C, Thermes C, Rétaux S, Casane D. 2018. Evidence for late Pleistocene
origin of Astyanax mexicanus cavefish. *BMC Evol Biol* **18**:43. doi:10.1186/s12862-018-1156-7

Gallo ND, Jeffery WR. 2012. Evolution of space dependent growth in the teleost Astyanax mexicanus.
PLoS One **7**:e41443. doi:10.1371/journal.pone.0041443

Gore A V., Tomins KA, Iben J, Ma L, Castranova D, Davis AE, Parkhurst A, Jeffery WR, Weinstein
BM. 2018. An epigenetic mechanism for cavefish eye degeneration. *Nat Ecol Evol* **2**:1155–1160.
doi:10.1038/s41559-018-0569-4

Gross JB, Furterer A, Carlson BM, Stahl BA. 2013. An integrated transcriptome-wide analysis of cave
and surface dwelling *Astyanax mexicanus*. *PLoS One* **8**:e55659. doi:10.1371/journal.pone.0055659

Helt MS, Fein DA, Vargas JE. 2020. Emotional contagion in children with autism spectrum disorder
varies with stimulus familiarity and task instructions. *Dev Psychopathol* **32**:383–393.
doi:10.1017/S0954579419000154

Herman A, Brandvain Y, Weagley J, Jeffery WR, Keene AC, Kono TJY, Bilandžija H, Borowsky R,
Espinasa L, O'Quin K, Ornelas-García CP, Yoshizawa M, Carlson B, Maldonado E, Gross JB,
Cartwright RA, Rohner N, Warren WC, McGaugh SE. 2018. The role of gene flow in rapid and
repeated evolution of cave-related traits in Mexican tetra, Astyanax mexicanus. *Mol Ecol* **27**:4397–
4416. doi:10.1111/mec.14877

Holm S. 1979. A simple sequentially rejective multiple test procedure. *Scand J Stat*.

Huppop K. 1986. Oxygen-Consumption of Astyanax-Fasciatus (Characidae, Pisces) - a Comparison of
Epigean and Hypogean Populations. *Environ Biol Fishes* **17**:299–308.

Iwashita M, Yoshizawa M. 2021. Social-like responses are inducible in asocial Mexican cavefish despite
the exhibition of strong repetitive behavior. *Elife* **10**:e72463. doi:10.7554/eLife.72463

Jaggard JB, Lloyd E, Yuiska A, Patch A, Fily Y, Kowalko JE, Appelbaum L, Duboue ER, Keene AC.
2020. Cavefish brain atlases reveal functional and anatomical convergence across independently
evolved populations. *Sci Adv* **6**:3126–3142. doi:10.1126/sciadv.aba3126

Jaggard JB, Stahl BA, Lloyd E, Duboue ER, Keene AC, Prober DA, Duboue ER, Keene AC. 2017.
Hypocretin underlies the evolution of sleep loss in the Mexican cavefish. *Elife* **7**.
doi:10.7554/eLife.32637

Keene AC, Yoshizawa M, McGaugh SE. 2016. Biology and Evolution of the Mexican Cavefish,
Biology and Evolution of the Mexican Cavefish. Amsterdam: Elsevier Inc. doi:10.1016/C2014-0-
01426-8

Kowalko JE, Rohner N, Rompani SB, Peterson BK, Linden TA, Yoshizawa M, Kay EH, Weber J,
Hoekstra HE, Jeffery WR, Borowsky R, Tabin CJ. 2013. Loss of schooling behavior in cavefish
through sight-dependent and sight-independent mechanisms. *Curr Biol* **23**:1874–83.
doi:10.1016/j.cub.2013.07.056

Krautkramer KA, Dhillon RS, Denu JM, Carey H V. 2017. Metabolic programming of the epigenome:
Host and gut microbial metabolite interactions with host chromatin. *Transl Res* **189**:30–50.
doi:10.1016/j.trsl.2017.08.005

Langen M, Durston S, Kas MJH, van Engeland H, Staal WG. 2011. The neurobiology of repetitive
behavior: ...and men. *Neurosci Biobehav Rev* **35**:356–365. doi:10.1016/j.neubiorev.2010.02.005

Lee RWY, Corley MJ, Pang A, Arakaki G, Abbott L, Nishimoto M, Miyamoto R, Lee E, Yamamoto S,
Maunakea AK, Lum-Jones A, Wong M. 2018. A modified ketogenic gluten-free diet with MCT

improves behavior in children with autism spectrum disorder. *Physiol Behav* **188**:205–211. doi:10.1016/j.physbeh.2018.02.006

Lee Y, Kang H, Jin C, Zhang Y, Kim Y, Han K. 2019. Transcriptome analyses suggest minimal effects of Shank3 dosage on directional gene expression changes in the mouse striatum. *Animal Cells Syst (Seoul)* **23**:270–274. doi:10.1080/19768354.2019.1595142

Li Q, Liang J, Fu N, Han Y, Qin J. 2021. A Ketogenic Diet and the Treatment of Autism Spectrum Disorder. *Front Pediatr* **9**:1–7. doi:10.3389/fped.2021.650624

Ludwig DS. 2020. The Ketogenic Diet: Evidence for Optimism but High-Quality Research Needed. *J Nutr* **150**:1354–1359. doi:10.1093/JN/NXZ308

Ma L, Parkhurst A, Jeffery WR. 2014. The role of a lens survival pathway including sox2 and αA-crystallin in the evolution of cavefish eye degeneration. *Evodevo* **5**:28. doi:10.1186/2041-9139-5-28

Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, Mathis MW, Bethge M. 2018. DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. *Nat Neurosci* **21**:1281–1289. doi:10.1038/s41593-018-0209-y

McCue MD. 2010. Starvation physiology: Reviewing the different strategies animals use to survive a common challenge. *Comp Biochem Physiol - A Mol Integr Physiol* **156**:1–18. doi:10.1016/j.cbpa.2010.01.002

McDonald TJW, Cervenka MC. 2018. Ketogenic Diets for Adult Neurological Disorders. *Neurotherapeutics* **15**:1018–1031. doi:10.1007/s13311-018-0666-8

McGaugh SE, Gross JB, Aken B, Blin M, Borowsky R, Chalopin D, Hinaux H, Jeffery WR, Keene A, Ma L, Minx P, Murphy D, O'Quin KE, Rétaux S, Rohner N, Searle SMJ, Stahl BA, Tabin C, Volff JN, Yoshizawa M, Warren WC. 2014. The cavefish genome reveals candidate genes for eye loss. *Nat Commun* **5**:5307. doi:10.1038/ncomms6307

Meidenbauer JJ, Mukherjee P, Seyfried TN. 2015. The glucose ketone index calculator: A simple tool to monitor therapeutic efficacy for metabolic management of brain cancer. *Nutr Metab* **12**:1–7. doi:10.1186/s12986-015-0009-2

Moran D, Softley R, Warrant EJ. 2015. The energetic cost of vision and the evolution of eyeless Mexican cavefish. *Sci Adv* **1**:e1500363. doi:10.1126/sciadv.1500363

Moran D, Softley R, Warrant EJ. 2014. Eyeless Mexican cavefish save energy by eliminating the circadian rhythm in metabolism. *PLoS One* **9**:e107877. doi:10.1371/journal.pone.0107877

Nakajo H, Chou MY, Kinoshita M, Appelbaum L, Shimazaki H, Tsuboi T, Okamoto H. 2020. Hunger Potentiates the Habenular Winner Pathway for Social Conflict by Orexin-Promoted Biased Alternative Splicing of the AMPA Receptor Gene. *Cell Rep* **31**:107790. doi:10.1016/j.celrep.2020.107790

Napoli E, Dueñas N, Giulivi C. 2014. Potential therapeutic use of the ketogenic diet in autism spectrum disorders. *Front Pediatr* **2**:1–9. doi:10.3389/fped.2014.00069

O'Connell LA, Hofmann HA. 2012. Evolution of a vertebrate social decision-making network. *Science (80-)* **336**:1154–1157. doi:10.1126/science.1218889

O'Connell LA, Hofmann HA. 2011. Genes, hormones, and circuits: An integrative approach to study the evolution of social behavior. *Front Neuroendocrinol* **32**:320–335. doi:10.1016/j.yfrne.2010.12.004

Padilla SL, Qiu J, Soden ME, Sanz E, Nestor CC, Barker FD, Quintana A, Zweifel LS, Rønnekleiv OK, Kelly MJ, Palmiter RD. 2016. Agouti-related peptide neural circuits mediate adaptive behaviors in the starved state. *Nat Neurosci* **19**:734–741. doi:10.1038/nn.4274

Paffenhöfer G-A. 1967. Caloric content of larvae of the brine shrimpArtemia salina. *Helgoländer Wissenschaftliche Meeresuntersuchungen* **16**:130–135. doi:10.1007/BF01620694

Panda S. 2016. Circadian physiology of metabolism. *Science (80-)* **354**:1008–1015. doi:10.1126/science.aah4967

Patch A, Paz A, Holt K, Duboue E, Kowalko J, Keene A, Fily Y. 2020. Analysis of social swimming dynamics in the Mexican cavefish. *bioRxiv* 2020.01.31.929323. doi:10.1101/2020.01.31.929323

Pérez-Escudero A, Vicente-Page J, Hinz RC, Arganda S, De Polavieja GG. 2014. IdTracker: Tracking
individuals in a group by automatic identification of unmarked animals. *Nat Methods* **11**:743–748.
doi:10.1038/nmeth.2994
Phillips MCL, Murtagh DKJ, Gilbertson LJ, Asztely FJS, Lynch CDP. 2018. Low-fat versus ketogenic
diet in Parkinson's disease: A pilot randomized controlled trial. *Mov Disord* **33**:1306–1314.
doi:10.1002/mds.27390
Pierre C, Pradère N, Froc C, Ornelas-García P, Callebert J, Rétaux S. 2020. A mutation in monoamine
oxidase (MAO) affects the evolution of stress behavior in the blind cavefish Astyanax mexicanus.
bioRxiv 2020.04.09.033266. doi:10.1101/2020.04.09.033266
Polakof S, Panserat S, Soengas JL, Moon TW. 2012. Glucose metabolism in fish: A review. *J Comp
Physiol B Biochem Syst Environ Physiol* **182**:1015–1045. doi:10.1007/s00360-012-0658-7
Provenzano G, Corradi Z, Monsorno K, Fedrizzi T, Ricceri L, Scattoni ML, Bozzi Y. 2016.
Comparative gene expression analysis of two mouse models of autism: Transcriptome profiling of
the BTBR and En2-/- Hippocampus. *Front Neurosci* **10**:396. doi:10.3389/fnins.2016.00396
Qin L, Ma K, Yan Z. 2021. Rescue of histone hypoacetylation and social deficits by ketogenic diet in a
shank3 mouse model of autism. *Neuropsychopharmacology* 1–9. doi:10.1038/s41386-021-01212-1
Rajasekaran A, Venkatasubramanian G, Berk M, Debnath M. 2015. Mitochondrial dysfunction in
schizophrenia: Pathways, mechanisms and implications. *Neurosci Biobehav Rev* **48**:10–21.
doi:10.1016/j.neubiorev.2014.11.005
Riddle MR, Aspiras AC, Gaudenz K, Peuß R, Sung JY, Martineau B, Peavey M, Box AC, Tabin JA,
McGaugh S, Borowsky R, Tabin CJ, Rohner N. 2018. Insulin resistance in cavefish as an
adaptation to a nutrient-limited environment. *Nature* **555**:647–651. doi:10.1038/nature26136
Rohner N, Jarosz DFDF, Kowalko JEJE, Yoshizawa M, Jeffery WRWR, Borowsky RLRL, Lindquist S,
Tabin CJCJ. 2013. Cryptic variation in morphological evolution: HSP90 as a capacitor for loss of
eyes in cavefish. *Science (80-)* **342**:1372–1375. doi:10.1126/science.1240276
Runco MA, Charlop MH, Schreibman L. 1986. The occurrence of autistic children's self-stimulation as
a function of familiar versus unfamiliar stimulus conditions. *J Autism Dev Disord* **16**:31–44.
doi:10.1007/BF01531576
Ruskin DN, Masino SA. 2012. The nervous system and metabolic dysregulation: Emerging evidence
converges on ketogenic diet therapy. *Front Neurosci* **6**:1–12. doi:10.3389/fnins.2012.00033
Solianik R, Sujeta A, Terentjevienė A, Skurvydas A. 2016. Effect of 48 h Fasting on Autonomic
Function, Brain Activity, Cognition, and Mood in Amateur Weight Lifters. *Biomed Res Int* **2016**:1–
8. doi:10.1155/2016/1503956
Strickler AG, Byerly MS, Jeffery WR. 2007. Lens gene expression analysis reveals downregulation of
the anti-apoptotic chaperone *alpha A-crystallin* during cavefish eye degeneration. *Dev Genes Evol*
217:771–782.
Sumithran P, Prendergast LA, Delbridge E, Purcell K, Shulkes A, Kriketos A, Proietto J. 2013. Ketosis
and appetite-mediating nutrients and hormones after weight loss. *Eur J Clin Nutr* **67**:759–764.
doi:10.1038/ejcn.2013.90
Szyf M. 2015. Epigenetics, a key for unlocking complex CNS disorders? Therapeutic implications. *Eur
Neuropsychopharmacol* **25**:682–702. doi:10.1016/j.euroneuro.2014.01.009
Vallée A, Vallée JN. 2018. Warburg effect hypothesis in autism Spectrum disorders. *Mol Brain* **11**:1–7.
doi:10.1186/s13041-017-0343-6
van der Weele CM, Jeffery WR. 2022. Cavefish cope with environmental hypoxia by developing more
erythrocytes and overexpression of hypoxia inducible genes. *Elife* **11**. doi:10.7554/eLife.69109
Xiong S, Wang W, Kenzior A, Olsen L, Krishnan J, Persons J, Medley K, Peuß R, Wang Y, Chen S,
Zhang N, Thomas N, Miles JM, Alvarado AS, Rohner N. 2022. Enhanced lipogenesis through
Pparγ helps cavefish adapt to food scarcity. *Curr Biol* 1–9. doi:10.1016/j.cub.2022.03.038
Yao J, Chen S, Mao Z, Cadenas E, Brinton RD. 2011. 2-deoxy-D-glucose treatment induces

ketogenesis, sustains mitochondrial function, and reduces pathology in female mouse model of
Alzheimer's disease. *PLoS One* **6**. doi:10.1371/journal.pone.0021788
Yoshizawa M. 2015. Behaviors of cavefish offer insight into developmental evolution. *Mol Reprod Dev*
82:268–280. doi:10.1002/mrd.22471
Yoshizawa M, Gorički S, Soares D, Jeffery WR. 2010. Evolution of a behavioral shift mediated by
superficial neuromasts helps cavefish find food in darkness. *Curr Biol* **20**:1631–1636.
doi:10.1016/j.cub.2010.07.017
Yoshizawa M, Robinson BG, Duboué ER, Masek P, Jaggard JB, O'Quin KE, Borowsky RL, Jeffery
WR, Keene AC. 2015. Distinct genetic architecture underlies the emergence of sleep loss and prey-
seeking behavior in the Mexican cavefish. *BMC Biol* **13**:15. doi:10.1186/s12915-015-0119-3
Yoshizawa M, Settle A, Hermosura MCM, Tuttle LJL, Cetraro N, Passow CNCN, McGaugh SESE.
2018. The Evolution of a Series of Behavioral Traits is associated with Autism-Risk Genes in
Cavefish. *BMC Evol Biol* **18**:89. doi:10.1186/s12862-018-1199-9
Yoshizawa M, Yamamoto Y, O'Quin KE, Jeffery WR. 2012. Evolution of an adaptive behavior and its
sensory receptors promotes eye regression in blind cavefish. *BMC Biol* **10**:108. doi:10.1186/1741-
7007-11-82
Zhu L, Wang X, Li XL, Towers A, Cao X, Wang P, Bowman R, Yang H, Goldstein J, Li YJ, Jiang YH.
2014. Epigenetic dysregulation of SHANK3 in brain tissues from individuals with autism spectrum
disorders. *Hum Mol Genet* **23**:1563–1578. doi:10.1093/hmg/ddt547
745
746

Figure Legends

Figure 1. Blood glucose and ketone levels under the control diet (CD) or ketogenic diet (KD).
(A) Experimental procedure. After fish were raised for 3–4 months on a brine shrimp larva diet, fish were fed the CD or KD for 5 weeks. After 5 weeks, blood glucose and ketone levels were measured. (B) Blood ketone level (mmol/L). Ketone levels were significantly reduced by KD feeding in both surface fish (SF) and cavefish (CF). Data are presented as the mean ± standard error of the mean. Dots indicate individual data. (C) Blood glucose level (mg/dL). Glucose levels were significantly reduced by KD feeding in both SF and CF. (D) The glucose ketone index (GKI) indicated that the ratio of glucose to ketone was lowered by KD feeding in both SF and CF, suggesting that this diet altered the balance between glucose and ketone.
SF: N = 13 for CD feeding, N = 8 for KD feeding. CF: N = 13 for CD feeding, N = 11 for KD feeding.
*: $P < 0.05$, **: $P < 0.01$, ***: $P < 0.001$. All detailed statistical data are available in Supplementary Table 1.

Figure 2. Time course of nearby interaction changes during 9 weeks of control diet (CD) or ketogenic diet (KD) feeding.
(A) Experimental procedure. After rearing fish for 3–4 months on a brine shrimp larva diet, the pre-treatment recording was performed, followed by CD or KD feeding for 5 weeks. Nearby interactions were recorded every week until week 6 of feeding. Subsequently, all groups including KD-fed fish were given the CD until week 9. (B) An example of nearby interaction events among surface fish (SF). The left panel presents an example frame of the video. The colored lines indicate the trajectories of individual fish. A red-labeled fish was followed by a blue-labeled fish. Each nearby event that met the detection criteria, namely a distance of ≤ 5 cm between two fish that was maintained for more than 4 s, was counted as a nearby interaction event. The right panel presents an example of the detected events presented in a raster plot (each yellow bar indicates a nearby interaction event). Each pair of fish (six pairs among four fish) is presented in the rows. (C) Duration of nearby interactions. Although SF did not exhibit any differences in the duration of nearby interactions (s) between CD (green) and KD (blue) feeding, differences were detected among cavefish (CF) in week 5. However, the nearby interaction duration was indistinguishable from that of the CD group starting in week 6 when the KD was withdrawn from the experimental group. (D) Number of nearby interactions. Whereas SF exhibited no differences between CD and KD feeding, differences were observed in CF in weeks 4–6. After the KD was withdrawn in week 6, the number of events decreased to the level observed with CD feeding. Data are presented as the mean ± standard error of the mean. Dots indicate individual data. N = 20 for each group. *: $P < 0.05$, **: $P < 0.01$, ***: $P < 0.001$. All detailed statistical data are available in Supplementary Table 1.

Figure 3. Ketogenic diet (KD) feeding induced surface fish (SF)-like speed profiles during nearby interactions in cavefish (CF).
Swimming speed changes before, during, and after nearby interaction events in SF (A) and CF (B). The mean swimming speeds (i) for 4 s before the nearby interaction event, (ii) during the event, (iii) during 4 s after the event, and (iv) during the out-of-event period were plotted (see the top-left inset of A). (A) Swimming speed was reduced during nearby interactions in SF in both the CD and KD groups. This profile was clearer in the fifth week (right panel). (B) Swimming speed was reduced during nearby interactions only in the KD group in week 5 (right panel). The bars indicate the 25th percentiles, medians, and 75th of the data points. Dots indicate individual data. SF: N = 11 for CD, N = 20 for KD. CF: N = 16 for CD, N = 15 for KD. *: $P < 0.05$, **: $P < 0.01$, ***: $P < 0.001$. All detailed statistical data are available in Supplementary Table 1.

Supplementary Figure 1. Swimming distance of cavefish was suppressed by ketogenic diet (KD) feeding.

Surface fish exhibited an increase in the swimming distance over time under both the control diet (CD) and KD. By contrast, in cavefish, the swimming distance and activity were suppressed by KD feeding starting in week 1, and these values remained smaller than those in CD-fed fish until week 6 when the KD was withdrawn. KD-fed cavefish subsequently exhibited an increased swimming distance (week 0). Data are presented as the mean ± standard error of the mean. Dots indicate individual data. N = 20 in all groups. *: $P < 0.05$, **: $P < 0.01$, ***: $P < 0.001$. All detailed statistical data are available in Supplementary Table 1.

Figure 4. Biased turning was attenuated by the ketogenic diet (KD).

(A) Diagram and the calculation formula for the turning bias index. The changes in the left or right traveling directions were calculated every five frames (every 0.25 s) across all trajectories and expressed as radians. Positive radian values represent left (anticlockwise) turning, and negative values indicate right turning. The ratio between the numbers of clockwise and anticlockwise turns was used as the turning rate (1 – infinity, positive value). (B) Turning biases of surface fish (left) and cavefish (right). There was no difference between CD and KD feeding in surface fish, whereas the turning index in CD-fed cavefish than in KD-fed cavefish (see week 6).

Data are presented as the mean ± standard error of the mean. Dots indicate individual data. N = 20 for all groups. *: $P < 0.05$, **: $P < 0.01$, ***: $P < 0.001$. All detailed statistical data are available in Supplementary Table 1.

Supplementary Figure 2. Consistent results were obtained in the repeated experiment for the duration and number of nearby interactions, swimming distance, and turning bias under control diet (CD) or ketogenic diet (KD) feeding.

(A) The duration of nearby interactions, (B) the number of nearby interactions, (C) swimming distance, and (D) turning bias are presented. The overall tendencies were the same as those observed in the original experiments (Figs. 2 and 4, Supplementary Fig. 1). Surface fish did not exhibit any significant differences regarding the duration (A) or number of nearby interactions (B), the swimming distance (C) or the turning bias (D). The duration (A) and number of nearby interactions were maintained in KD-fed cavefish, (B) whereas they were reduced in CD-fed cavefish, which also exhibited a higher level of turning bias (D). The swimming distance was not significantly reduced in KD-fed cavefish compared to that in CD-fed controls in this repeated experiment (C).

Data are presented as the mean ± standard error of the mean. Dots indicate individual data. Surface fish: N = 28 for CD, N = 32 for KD. Cavefish: N = 28 for CD, N = 32 for KD. *: $P < 0.05$, **: $P < 0.01$, ***: $P < 0.001$. All detailed statistical data are available in Supplementary Table 1.

Figure 5. Day and night sleeping durations and swimming distances were not altered by ketogenic diet (KD) feeding.

(A) Sleep duration (min/h) during the day (left) and night (right). During 5 weeks of growth, the sleep duration decreased in surface fish and cavefish regardless of the diet. (B) Number of events (min/10 min) during the day (left) and night (right). During 5 weeks of growth, the number of sleep events was lower in surface fish under both dietary conditions and in KD-fed cavefish. (C) Swimming distance during the day (left) and night (right). Control diet (CD)-fed cavefish exhibited a longer swimming distance during the day and night. Conversely, surface fish fed either diet and cavefish fed the KD exhibited a significantly increased swimming distance only at night.

842 Data are presented as the mean ± standard error of the mean. Dots indicate individual data. Surface fish:
843 N = 28 for CD, N = 32 for KD. Cavefish: N = 28 for CD, N = 32 for KD. *: P < 0.05, **: P < 0.01, ***:
844 P < 0.001. All detailed statistical data are available in Supplementary Table 1.
845
Supplementary Figure 3. Daytime and nighttime number of sleeping events under control diet
(CD) or ketogenic diet (KD) feeding.
After the 5 weeks of growth, surface fish exhibited a reduced number of sleeping events during the day
under both diets. CD-fed cavefish exhibited reduced numbers of sleeping events during the day and
night. However, the number of sleeping events did not differ according to the diet in cavefish or surface
fish.
Data are presented as the mean ± standard error of the mean. Dots indicate individual data. Surface fish:
N = 28 for CD, N = 32 for KD. Cavefish: N = 28 for CD, N = 32 for KD. *: P < 0.05, **: P < 0.01, ***:
P < 0.001. All detailed statistical data are available in Supplementary Table 1.
855
Supplementary Figure 4. Vibration attraction behavior (VAB) and swimming distance during
VAB under control diet (CD) or ketogenic diet (KD) feeding.
(A) Number of approaches to the vibration rod in the 3-min assays. After 5 weeks of growth, the number
of approaches was increased in CD- and KD-fed cavefish, but no difference according to the diet was
detected in either surface fish or cavefish. (B) Swimming distance during VAB. KD-fed cavefish swam
significantly shorter distances than CD-fed cavefish.
Data are presented as the mean ± standard error of the mean. Dots indicate individual data. Surface fish:
N = 28 for CD, N = 32 for KD. Cavefish: N = 28 for CD, N = 32 for KD. *: P < 0.05, **: P < 0.01, ***:
P < 0.001. All detailed statistical data are available in Supplementary Table 1.
865
Figure 6. Body length and weight under control diet (CD) or ketogenic diet (KD) feeding.
(A) Standard length (cm). KD-fed surface fish and cavefish were significantly smaller than their CD-fed
counterparts. (B) Body weight (g). KD-fed surface fish and cavefish weighed less than their CD-fed
counterparts.
Data are presented as the mean ± standard error of the mean. Dots indicate individual data. Surface fish:
N = 28 for CD, N = 32 for KD. Cavefish: N = 28 for CD, N = 32 for KD. *: P < 0.05, **: P < 0.01, ***:
P < 0.001. All detailed statistical data are available in Supplementary Table 1.
873
Figure 7. Nearby interactions and other behaviors under control diet (CD) or beta-
hydroxybutyrate–supplemented diet (BHB) feeding.
(A) Duration of nearby interactions (s). After 4 weeks, the duration of nearby interactions was decreased
in BHB-treated surface fish and increased in BHB-treated cavefish. (B) Number of nearby interactions.
The number of nearby interactions was increased in BHB-treated cavefish. (C) Turning bias ratio. BHB-
treated cavefish tend to exhibit decreased biased turning, although this reduction was not significant. (D)
Swimming distance. No difference was detected between the CD and BHB groups.
Data are presented as the mean ± standard error of the mean. Dots indicate individual data. N = 20 for all
groups. *: P < 0.05, **: P < 0.01, ***: P < 0.001. All detailed statistical data are available in
Supplementary Table 1.
884
885 **Supplementary Figure 5. Body size and weight under control diet (CD) or beta-hydroxybutyrate-**
886 **supplemented diet (BHB) feeding.**
887 (A) Standard length (cm). No statistical difference was found for either diet. (B) Body weight (g). BHB-
888 treated surface fish exhibited significantly reduced weight, whereas the weight of cavefish did not differ
889 according to the diet.

Data are presented as the mean ± standard error of the mean. Dots indicate individual data. N = 20 for all groups. *: P < 0.05, **: P < 0.01, ***: P < 0.001. All detailed statistical data are available in Supplementary Table 1.

Supplementary Figure 6. Day and night sleeping durations and vibration attraction behavior (VAB) were not drastically changed by beta-hydroxybutyrate-supplemented diet (BHB) feeding. (A) Sleep duration (min/h) during the day (left) and night (right). During 4 weeks of treatment, the daytime sleep duration in surface fish and cavefish increased in BHB feeding but not in the nighttime. (B) Number of approaches per the 3 min assay (VAB level). During 4 weeks of treatment, the VAB level did not shift in surface fish or cavefish regardless of diets. Data are presented as the mean ± standard error of the mean. Dots indicate individual data. N = 20 for all groups. *: P < 0.05, **: P < 0.01, ***: P < 0.001. All detailed statistical data are available in Supplementary Table 1.

Figure 1

A   Surface fish

B   Cavefish

Figure 4

A

radian < 0 (Right turn)

radian > 0 (Left turn)

$$\text{Turning Rate} = \text{Maximum of } \left(\frac{\# \text{ of Right turn}}{\# \text{ of Left turn}}, \frac{\# \text{ of Left turn}}{\# \text{ of Right turn}} \right)$$

B

Figure 5

Supplementary Figure 3

Supplementary Figure 4