Integration of metabolism and regulation reveals rapid adaptability to growth on non-native substrates.

Vikas D. Trivedi[1,2], Sean F. Sullivan[1], Debika Choudhury[1], Venkatesh Endalur Gopinarayanan[1], Taylor Hart[1], Nikhil U. Nair[1*]

[1] Department of Chemical & Biological Engineering, Tufts University, Medford, MA
[2] Current: Department of Structural Biology and Center for Data Driven Discovery, St. Jude Children's Research Hospital, Memphis, TN

* Corresponding author: nikhil.nair@tufts.edu; @nair_lab; @nair_lab@mastodon.social

ABSTRACT

Engineering synthetic heterotrophy (i.e., growth on non-native substrates) is key to the efficient bio-based valorization of renewable and waste substrates. Among these, engineering hemicellulosic pentose utilization has been well-explored in *Saccharomyces cerevisiae* (yeast) over several decades – yet the answer to what makes their utilization inherently recalcitrant remains elusive. Through implementation of a semi-synthetic regulon, we find that harmonizing cellular and engineering objectives are key to obtaining highest growth rates and yields with minimal metabolic engineering effort. Concurrently, results indicate that "extrinsic" factors – specifically, upstream genes that direct flux of pentoses into central carbon metabolism – are rate-limiting. We also reveal that yeast metabolism is innately highly adaptable to rapid growth on non-native substrates and that systems metabolic engineering (i.e., flux balancing, directed evolution, functional genomics, and network modeling) is largely unnecessary. We posit that the need for extensive engineering espoused by prior works is a consequence of unfortunate (albeit avoidable) antagonism between engineering and cellular objectives. We also found that deletion of endogenous genes to promote growth demonstrate inconsistent outcomes that are genetic-context- and condition-dependent. For the most part, these knockouts also lead to deleterious pleiotropic effects that decrease the robustness of strains against inhibitors and stressors associated with bioprocessing. Thus, at best, perturbation of "intrinsic" factors (e.g., native metabolic, regulatory genes) results in incremental and inconsistent benefits. At worst, they are detrimental. Overall, this work provides insight into the limitations and pitfalls to realizing efficient synthetic heterotrophy using traditional/systems metabolic engineering approaches, demonstrates the innate adaptability of yeast for metabolism of non-native substrates, and provides an alternate, novel, holistic (and yet minimalistic) approach based on integrating non-native metabolic genes with a native regulon system.

KEYWORDS: synthetic biology, systems biology, metabolic engineering, sustainability, biomanufacturing

INTRODUCTION

Engineering metabolism for growth on non-native substrates (i.e., synthetic heterotrophy) has been an outstanding challenge for several decades in various microbial species (*1-4*). The traditional approach is to constitutively overexpress catabolic genes that input substrate to central carbon metabolism with the expectation that the native cellular systems can direct subsequent steps required for growth. In case initial designs do not lead to rapid growth, flux balancing (*5*), adaptive laboratory evolution (ALE) (*6, 7*), functional genomics (*8, 9*), and directed evolution (*6, 10*) approaches are used. More recently, systems-level analysis of regulatory structures in engineered strains have been used to identify dysregulated pathways and corrective interventions have provided avenues to improve engineering outcomes (*11-13*). Unifying to these approaches is that they are reactive interventions that try to rectify inefficiencies introduced due to strain engineering but are not proactive in circumventing undesirable outcomes by harmonizing recombinant activity with native cellular objectives. Typical of this approach is engineering the yeast *Saccharomyces cerevisiae* for complete and rapid utilization of lignocellulosic pentoses (*14-17*). In this work, we engineer strains of yeast for rapid growth and pentose (xylose, arabinose) utilization and assess intrinsic and extrinsic constraints and pitfalls that control desirable phenotypic outcomes. Specifically, we answer the longstanding unresolved question: *what are the inherent/intrinsic limitations within this yeast that prevent it from metabolizing non-native pentose substrates at rates equivalent to native substrates?* Surprisingly, we find that this yeast is intrinsically highly adaptable to rapid growth on non-native substrates and that it can achieve maximum growth rates with minimal engineering – which is in direct contrast with the currently accepted paradigm.

To achieve this, we first modified our semi-synthetic regulon (*18*) – a system based on synthetically activating the galactose (GAL) response system using a non-native substrate – for efficient utilization of arabinose through the isomerase pathway (*19*). We demonstrate that while initial outcomes – like with xylose, previously – are superior to constitutively overexpressing the same genes, growth rates were modest ($\mu = 0.14 - 0.17$ h^{-1}). To identify factors that constrain growth on either pentose, we performed pathway balancing, directed evolution, and systems biology-driven functional genomics. To our surprise, we found that when cells can coordinate global growth responses with substrate use through the semi-synthetic GAL regulon (REG), growth is largely extrinsically controlled – i.e., limited by the upstream/heterologous pathway that controls non-native substrate uptake and flux to central carbon metabolism. We also found through parallel investigations that genetic interventions identified through traditional/systems metabolic engineering that prune cellular metabolic and/or regulatory networks to improve strain performance often led to pleiotropic defects with mid-to-severe fitness costs. Thus, our findings suggest that an approach that synergizes activation of the GAL regulon with an optimized upstream heterologous metabolic module reveals the hidden metabolic adaptability of yeast.

78 **RESULTS**
79
A semi-synthetic regulon design outperforms a constitutive overexpression strategy.
Previously, we demonstrated that coupling activation of galactose (GAL)-responsive regulon to
catabolism and growth on non-native substrate, xylose, enabled faster growth and more complete
utilization when compared to the constitutive overexpression of the same genes ([18]). This is because
GAL regulon activation upregulates growth responsive genes and suppresses starvation responses,
in contrast to the traditional constitutive overexpression approaches where the opposite holds true
(**Figure 1A**) ([18]). We wanted to assess whether the benefit was unique to xylose or if growth on other
substrates, like arabinose, could also benefit from activation of the GAL regulon. Since Gal3p$^{Syn4.1}$
was engineered to activate the GAL regulon in a xylose-dependent manner, we wondered if
arabinose could do so as well since the two sugars are structurally similar. We quantified activation of
GAL1p-EGFP by wild-type Gal3p or engineered Gal3p$^{Syn4.1}$ in the presence of native Gal2p and/or
engineered Gal2p$^{2.1}$ permease ([18]). We observed that a Gal3p$^{Syn4.1}$ and Gal2p$^{2.1}$ co-expressing strain
showed activation on arabinose with low background and high dynamic range (**Figure 1B**).
Interestingly, arabinose showed higher activation of *GAL1p-EGFP* than xylose, even though both
Gal3p$^{Syn4.1}$ and Gal2p$^{2.1}$ were engineered for activity on xylose (**Figure 1C**). Given robust activation,
we constructed a semi-integrant strain by integrating accessory genes including, sensor (*GAL3$^{Syn4.1}$*),
transporter (*GAL2$^{2.1}$*), and transaldolase (*TAL1*) – expressed under GAL promoters – to generate a
"REG" (regulon) strain background (**Figure 1D**). Similarly, as control, and to compare the with the
traditional engineering approach, we integrated *GAL2$^{2.1}$* and *TAL1* under strong constitutive
promoters to generate the "CONS" parental strain. We then assessed the growth of both strains on
arabinose or xylose after transformation with plasmid-encoded *araBAD* or *XYLA*3-XKS1* catabolic
genes, respectively. In ARA-REG and XYL-REG all genes were expressed under GAL-responsive
promoters whereas in ARA-CONS and XYL-CONS all genes were under constitutive promoters
(**Figure 1D**). On both carbon sources, strains that used the GAL regulon to coordinate substrate
assimilation with global metabolism demonstrated higher growth rates (μ: ARA-REG = 0.14 ± 0.04 h^{-1},
XYL-REG = 0.17 ± 0.01 h^{-1}) compared to those that constitutively overexpressed the same genes (μ:
ARA-CONS = 0.06 ± 0.01 h^{-1}, XYL-CONS = 0.11 ± 0.01 h^{-1}). They also reached higher final cell
densities (OD$_{600}$: ARA-REG = 6.01 ± 1.62, ARA-CONS = 1.31 ± 0.04, XYL-REG = 10.8 ± 0.62, XYL-
CONS = 5.3 ± 0.49) (**Figure 1E-F**). We then wanted to assess what factors prevented these REG
strains from achieving maximum aerobic growth rate (μ = $0.22 - 0.29$ h^{-1}) ([13, 20]). We explored two
potential limitations – i) intrinsic, i.e., native metabolic or regulatory systems downstream of substrate
uptake, and ii) extrinsic, i.e., upstream metabolic modules (primarily heterologous genes) that direct
non-native substrates to central carbon metabolism.
113
114 **Perturbation of intrinsic (downstream) factors result in modest and inconsistent**
115 **improvements.**
116 In traditional/systems metabolic engineering strategies, adaptive laboratory evolution (ALE) or rational
117 data-driven approaches are often used to improve growth rates of initial strain designs, since growth
118 rates and biomass yields are most often suboptimal. We decided to use a (data-driven) systems

 biology approach to identify "intrinsic" genetic targets to modify to improve the growth rates of our

120   REG strains

121
Figure 1. Implementation of a semi-synthetic regulon design for synthetic heterotrophy. A) The GAL
regulon, synthetically activated by Gal3p$^{Syn4.1}$, regulates GAL-responsive genes as well as hundreds of
downstream genes to promote growth. The traditional approach of constitutive overexpression cannot activate
downstream genes, which leads to starvation response and slow growth (*18*). **A)** Dose-dependent *GAL1p-EGFP*
activation by arabinose of strain expressing Gal3p^{WT} or Gal3p$^{Syn4.1}$ in and Gal2p^{WT} or Gal2p$^{2.1}$. **B)** Comparison of
GAL1p activation by 2 % of native sugar, galactose, and non-native sugars – xylose and arabinose. **C)** Genetic
structure of semi-integrant REG and CONS strain backgrounds. Growth curve of REG and CONS strains on **D)**

arabinose and **E)** xylose when transformed with respective arabinose and xylose isomerase pathway genes, respectively. REG and CONS strains use GAL-responsive and constitutive promoters, respectively, for all genes indicated (**Table S1 and S2**).

Figure 2. Systems analysis to identify intrinsic limitations in yeast for growth on pentoses. A) Differential expression volcano plot for transcriptomic comparison of strains grown individually on galactose or pentose sugars (xylose or arabinose) with the number of genes relatively upregulated in each condition overlaid. **B)** Plots of Betweenness Centrality (BC) (arbitrary units) vs. significance ($-\log_{10}$P) for the subset of significantly ($p < 0.05$) differentially expressed genes demonstrate how native gene targets for deletion were identified using three yeast gene regulatory networks (GRNs): EGRIN (top), YEASTRACT (center), and CLR (bottom). Genes were categorized as either, scoring highest in BC in a given network and picked for deletion (red), not picked for deletion (gold), identified as deletion target in another network (purple), or all remaining significant nodes (grey). **C)** 24 highly connected genes identified via differential gene expression and gene regulatory network analyses (with $p < 0.05$) as targets for deletion to improve growth on pentoses.

 on xylose and arabinose (i.e., pentoses). By "intrinsic" factors, we mean native yeast genes that are

 far downstream of substrate uptake and catabolism (e.g., cell wall biosynthesis, DNA repair).

 As the GAL regulon has evolved to function harmoniously during growth on a native substrate, we

 hypothesized that identifying expression differences between galactose and the two non-native

 substrates and closing the gap between them may aid in better synchronizing GAL regulation with

 pentose metabolism. To do so, we performed RNA-seq on REG strains grown on arabinose, xylose,

 and galactose to characterize their respective expression profiles. A differential gene expression

 (DGE) analysis between individual sugars revealed that a total of 838 genes were differentially

 regulated between arabinose and galactose and 1,430 genes between xylose and galactose strains

 (**Figure S1**) (p-value < 0.05 after Benjamini–Hochberg correction). However, rather than pursue the

 individual substrate comparisons, we chose to repeat the DGE analysis to compare pentose REG vs.

 galactose utilization to identify "intrinsic" factors that differentiate growth on the two pentoses vs. the

 hexose galactose. Upon doing so, we found 1,845 genes that were relatively upregulated on pentoses

 and 1,678 genes that were relatively upregulated during growth on galactose (**Figure 2A**). Given the

 large number of differentially expressed genes (3,523) between the two conditions (pentoses vs.

galactose), we sought to use gene regulatory networks (GRN) to identify highly-connected, regulatory

elements that could serve as major contributors to the observed divergence in transcriptional profiles

and therefore growth rates. To do so, we first identified three different yeast GRNs that were

previously validated for their ability to improve the accuracy of yeast growth phenotypes when

integrated with existing metabolic models (*21*). These GRNs – EGRIN, YEASTRACT, and CLR – all

sought to capture influences on expression between yeast genes but differed in the data types (gene

expression, DNA-binding, protein-protein interactions, etc.) used to infer these connections. The

resulting networks were represented as graphs with nodes corresponding to genes and edges

between nodes corresponding to an inferred connection between genes. To utilize these networks,

we began by associating each node with the expression fold-change and (Benjamini-Hochberg

corrected) significance values from the differential expression comparison between the pentose- and

galactose-grown REG strains. We then eliminated all nodes with corrected p-values >0.05 along with

any connections to those nodes to isolate significantly perturbed sections of the network. We next

utilized the network visualization and analysis software tool Cytoscape (*22*) to assign a betweenness

centrality (BC), a measure of the centrality of a given node in a network, value to the remaining genes

in each network. Plotting BC vs significance in gene expression change helped narrow down a subset

of regulatory genes as targets for deletion (**Figure 2B**). Finally, we manually curated a subset of 24

genes among these that have roles in regulation, metabolism, and stress response for deletion to test

their effect of growth and phenotype (**Figure 2C**).

We generated barcoded deletion (KO) libraries of these "intrinsic" factors in wild-type and REG strain

backgrounds and performed growth enrichments on glucose, galactose, xylose, and arabinose

(**Figure 3A**). Comparing population shifts using barcodes, we identified eight genes that displayed

positive fitness on both xylose and arabinose but not on galactose – indicating a role in controlling

growth primarily on pentoses. We then tested their growth individually on galactose, xylose, and

arabinose (**Figure 3B-D**) and found that only Δ*GLN3* was either neutral or beneficial on both

pentoses – all other deletions were detrimental for growth on xylose. Further, the growth rates of all

knockout strains were still lower than that of the parental strain on galactose ($\mu = 0.24 \pm 0.03$ h^{-1} when

GAL1-7-10 are expressed from a plasmid, **Figure S2**), indicating additional limitations. Overall, the

systems metabolic engineering approach failed to identify genetic factors broadly beneficial to growth

on pentoses. In contrast, the genome-wide expression changes effected by GAL regulon activation

are beneficial for all three substrates.

Pentose metabolism is largely extrinsically (upstream) controlled with a regulon approach.

Our studies have so far focused on identifying "intrinsic" factors that may be controlling/bottlenecking

growth on pentoses and given the inconsistent benefits, we wondered whether the limitations were, in

fact, "extrinsic". The former implies that the native metabolic and/or regulatory capacity of this yeast is

inherently limited for effective pentose metabolism and improving growth rate requires vast

restructuring of associated (intrinsic) metabolic and/or regulatory networks. The latter implies that

there are no inherent limitations in this yeast and that it is already poised for rapid growth on

pentoses, but the observed low growth rates are due to suboptimal design of the upstream metabolic

module that includes the heterologous (extrinsic) genes. To assess whether the "extrinsic limitation"
paradigm has any merits, we needed to optimize the design of the upstream metabolic module
responsible for substrate uptake and flux into central carbon metabolism (i.e., glycolysis).
203

204
205   **Figure 3. Effect of gene knockouts on growth phenotype. A)** Fitness map of knockout library of 24 genes
206   identified from the mutual information network. The scale represents fitness ranging from blue (negative) to
207   orange (positive). Growth rates of KOs with positive fitness on **B)** galactose, **C)** xylose (*GAL1p- XYLA*3 GAL10p
208   -XKS1*), **D)** arabinose (*GAL1p-araB GAL10p-araA GAL7p-araD*). Glc = glucose, Gal = galactose, Xyl = xylose,
209   Ara = arabinose. The growth rates were normalized from the average of at least two biological replicates.
210

First, we looked at the effect of plasmid copy number. For arabinose, there was no difference in
growth rate when *araA-araB-araD* genes were expressed on high- or low-copy plasmids, whereas for
xylose, we only observed growth when the *XYLA*3-XKS1* gene dose was high (**Figure S3**). In either
case, changing plasmids backbone did not improve growth rate. Next, we hypothesized that balancing
expression of these heterologous genes may be required to enhance growth rate. For arabinose, we
created all six combinations of gene-promoter pairings and assessed their performance. While such
promoter swaps have been previously demonstrated to improve strain performance (*23*), we were
surprised at the marked improvement in growth rate – from 0.14 ± 0.04 h^{-1} in the original design to
0.27 ± 0.03 h^{-1} in the best re-design (**Figure 4A**). To understand the cause of this behavioral change,
we compared the relative expression levels of the *araA*, *araB,* and *araD* using quantitative reverse
transcription PCR (qPCR) (**Figure 4B**). We observed that in the poor performing combinations,
expression of *araB* (ribulokinase) was the highest, whereas, in high performing combinations,
expression level followed this pattern *araA > araB > araD*. In addition, we found a strong positive
correlation between growth rate and relative expression of *araA*:*araB* and *araA*:*araD*, respectively
(**Figure 4C-E**). The success of this approach encouraged us to attempt the same on xylose and we
found similar improvements – from 0.17 ± 0.01 h^{-1} to 0.24 ± 0.01 h^{-1} (**Figure S4**). These are

comparable to the growth rate of yeast on galactose when *GAL1-7-10* are expressed from a plasmid

(0.24 ± 0.03 h^{-1}) (**Figure S2**).

230
231 **Figure 4. Optimizing extrinsic (upstream) factors to enhance growth of yeast on pentoses. A)** Comparison
232 of growth performance of different promoter-gene combination. **B)** Gene expression analysis of the six strains by
233 qPCR. Pearson correlation coefficient between growth rate and gene expression levels of **C)** *araA*:*araB*, **D)**
234 *araB:araD,* and **E)** *araA:araD*, respectively. **G)** Enrichment of mutagenic *araBAD* library in SC+Ara. **H)** Fitness

Figure 5. **Effect of gene deletions on growth and stress-response phenotypes. A)** Growth rate of the optimized plasmids in KO-strains. Fitness of KOs in presence of individual stressors when grown on **B)** sucrose and **C)** arabinose or in the mixture of stressors on **D)** sucrose and **E)** arabinose. The relative fitness is expressed as growth of deletion strains over the fitness of parental strain under the same conditions. NaCl = sodium chloride; NaAc = sodium acetate; Fur = furfural; 5HMF = 5-hydroxymethylfurfural; Vald = veratraldehyde. 1/2x, 1/4x, 1/8x = serial dilutions of the five stressors combined. Xyl* = *GAL10p-XYLA*3 GAL7p-XKS1*; Ara* = *GAL1p-araA GAL10p-araB GAL7p-araD* in respective REG backgrounds.

Finally, we used directed evolution to improve the growth rate on arabinose further (**Figure 4F**). We randomly mutagenized the six arabinose pathway combinations, adding barcodes to track the lineages and enriched this library size of 10^8 variants in minimal (SC+Ara) and complex (2YP+Ara) arabinose media (**Figure 4F**). Over the course of 15 subcultures, the growth rate over subcultures increased from 0.12 h^{-1} to 0.22 h^{-1} and 0.18 h^{-1} to 0.26 h^{-1} in SC+Ara and 2YP+Ara, respectively (**Figure 4G, S6**). Using barcodes, we tracked the performance of promoter-gene combinations (**Figure 4H**). We observed that initially all the six plasmids start at similar abundance, but the *araA-B-D* (under *GAL1-10-7p*, respectively) was the most abundant at the end of enrichment in SC+Ara (**Figure 4H**). We picked single colonies from each condition and calculated their growth rates and

identified 4 variants that showed the highest growth rates (0.35 ± 0.04 h^{-1}) from the SC+Ara enriched

culture (**Figure S5**). We sequenced the barcodes to identify the lineage and the whole cassette to

identify the mutations and found that three of the six initial designs were represented in the four best

variants (named N-3, N-6, N-12, and N-16) (**Table S3**). Re-transformation into parent background

strain indicated that the four variants attained the same maximum growth rate as that on galactose

(0.29 ± 0.01 h^{-1}), which is the highest reported growth rate on any pentose reported in the literature for

this yeast (**Figure 4I**). Since the mutations were distributed throughout the cassettes, we quantified

the expression levels of *araA*, *araB*, and *araD* in the four strains (**Figure 4J**) and found patterns that

differed significantly than parental strains (**Figure 4B**). We saw a strong positive correlation between

growth rate and relative expression of *araA*:*araB* and negative correlation between *araB*:*araD*,

indicating that the directed evolution campaign likely altered both activity and expression in each

strain differently (**Figure S6**). These results highlight a key insight about yeast: it's ability to utilize

pentoses is largely limited only by "extrinsic" factors (upstream pathway, especially heterologous

enzyme activity) and minimally by any "intrinsic" factor (i.e., native regulation or metabolic pathway).

Engineering intrinsic (downstream) genes lead to pleiotropic fitness trade-offs.

Next, we assessed whether the global regulatory elements identified through our network analysis

would benefit our best designs (*araA-B-D* under *GAL1-10-7p* and *XYLA*3-XKS1* under *GAL1-7p*) but

found that deletion of each of the 8 genes did not result in any improvements over the parental strains

(**Figure 5A**). We considered exploring combinatorial deletions and ALE to attempt to further enhance

growth rates on these two substrates. However, we expected that significant improvements were

unlikely as both strains had already attained rates comparable to the empirically determined

maximum aerobic growth rates of this yeast (0.22 h^{-1} – 0.29 h^{-1}) (*13, 20*) with high biomass yields and

short lag, we expect that improvements would be insignificant.

One consideration not yet assessed is the suitability of these strains for bioprocessing and their

resilience to growth inhibitors associated with bioprocessing applications. Given that all deletion

targets are highly-connected genes that control key cellular processes, we were concerned that

dysregulation major networks may lead to undesirable pleiotropic effects. Since resilience against

stress is a complex phenotype, often requiring concerted response from gene networks, we tested the

fitness of all single knockout (KO) strains on several stressors (sodium chloride, sodium acetate,

furfural, 5-hydroxymethylfurfural, and veratraldehyde) relevant to bioprocessing individually and in

combination (**Figure 5B-E**). Using parental strain as reference, we observed that performance of

strains under stress was highly context-dependent. For example, in sucrose medium, all deletions lost

fitness in single or mixed inhibitor cultures (**Figure 5C, E**). Conversely, on arabinose (with GAL

regulon activated), certain KO strains had improved tolerance to stressors (e.g., *ΔTEC1, ΔMET28*)

(**Figure 5D, F**). *ΔGLN3* has previously been shown to improve fitness under isobutanol stress (*24*);

however, in our study, it was less fit than the parental strain under all stress conditions. Interestingly, it

did demonstrate improved growth rate in a sub-optimal upstream metabolic design in arabinose

(**Figure 3E**) but lost that benefit in a more optimized design (**Figure 5C, E**). Collectively, these results

highlight that deleting genes to remodel expression profiles to enhance a single phenotype (e.g.,
growth rate on a non-native substrate) can lead to some improvements, but they are often
accompanied by negative pleiotropic effects that make the strain less suitable for eventual
bioprocessing applications where growth conditions are often non-ideal.

DISCUSSION

Despite decades of effort engineering synthetic heterotrophy with *S. cerevisiae*, there is no formalized
understanding of what limits the metabolic adaptability of this yeast for growth on pentoses. It is
important to differentiate between substrate utilization/uptake from growth since the former can be
readily achieved by diverting flux to unwanted or dead-end byproducts (e.g., Crabtree metabolites,
organic acids, pentitols) often with native substrate supplementation to support biomass generation.
However, valorization to high-value products requires efficient catabolism to central metabolic
products whose concentrations and fluxes are regulated and associated with growth. Initial studies in
literature focused on upstream ("extrinsic") elements (e.g., heterologous gene expression/activity,
transporter engineering, etc. (*25-29*)) whereas recent focus has been more "intrinsic", focusing on
functional genomics, adaptively laboratory evolution (ALE), and often network/systems analysis, to
identify downstream limitations (*5-13*). Successes have been abundantly reported with improvements
in growth along with a series of deletion and overexpression targets (*PHO13, ALD6, ASK10, YPR1,*
SNF6, RGT1, CAT8, MSN4, GPD1, CCH1, ADH6, BUD21, ALP1, ISC1, RPL20B, COX4, ISU1,
SSK2, YLR042c, CYC8, PHD1, TEC1, ARR1, etc. (*13, 30-36*)). Significantly, many targets are unique
to individual studies. This is perhaps unsurprising as the impact of such interventions, and even the
performance of identical heterologous pathways, can depend on genetic context (*37, 38*). Typifying
the traditional approach is a recent report where all the aforementioned extensive systems metabolic
engineering approaches were used to develop a strain of yeast with high growth aerobic growth rate
(0.26 h^{-1}) and short lag phase (9-15 h) on xylose (*13*). Despite this, it is not clear if the insights were
translatable to engineering growth on any other substrate.

Through our work, we posit that the requirement for extensive engineering to identify intrinsic
limitations is only required when the initial strain design is sub-optimal. Indeed, constitutive
overexpression of upstream metabolic genes results in significant stress (*18*) and low growth rates
that must be compensated for through systems approaches focused on preventing cellular detection
of stress rather than activating growth promoting systems. We demonstrated that with an optimal
upstream module, strains can attain superior growth profiles (fast specific growth rate, $\mu \geq 0.24$ h^{-1};
short lag phase, <10 h; high final biomass, OD$_{600}$ >10) if a growth-associated regulon – the GAL
regulon – is activated. This builds on our prior observation that synthetic activation of the GAL regulon
potentiates cells for rapid growth on non-native substrates by upregulating growth responsive genes
and suppressing starvation responses (*18*). Importantly, using this approach our final strain designs
had minimal modifications – we deleted only one gene (other than the Leloir pathway genes) – *GRE3*
– to minimize oxidation of substrate pentoses to pentitols (*39*). Many genes that have previously been
identified as important inactivation targets to improve growth on pentoses are intact. We also

overexpressed only a minimal upstream metabolic module (*TAL1* and *GAL2*[2.1]) along with the specific

pentose metabolic genes (*araBAD* or *XYLA*3-XKS1*). Our work strongly argues that growth on

pentoses is largely extrinsically controlled and there are no major intrinsic regulatory or metabolic

limitations in this yeast. This insight presents a paradigm shift in engineering synthetic heterotrophy.

An additional advantage of this approach is the preservation of native regulatory systems that are

otherwise dysregulated in the traditional engineering approach. We found that while deletion of

endogenous genes could improve growth in strains with sub-optimal upstream modules under certain

conditions, they are associated with pleiotropic defects and are less robust in the presence of growth

inhibitors. This is not surprising since most genetic interventions aim to suppress cellular responses to

stress rather than promote growth. Thus, strains engineered for synthetic heterotrophy through

extensive gene inactivation are less robust for eventual bioprocessing applications. This highlights a

major drawback of traditional/systems metabolic engineering approaches like ALE. In contrast, our

regulon-based approach is minimalistic and holistic – our strains maintain native regulatory systems

and even exploit them toward the engineering goal (i.e., the GAL regulon). On the one hand, our

utilization of a single strain background (W303-1a) and catabolic pathway for each substrate in this

study could potentially limit the broad applicability of our findings. For example, the use of the

oxidoreductase xylose pathway has been demonstrated to improve resistance to lignocellulosic

inhibitors relative to the isomerase pathway (*40*). Also, quantitative measures of strain fitness (i.e.,

growth rates) are known to be influenced by seemingly confounding factors like the choice of

selection marker and the copy number of the plasmid bearing the auxotrophy-complementing gene

(*41*). However, the regulon approach we are advocating is not an isolated perturbation to the cell but

interfaces with a native regulatory system that coordinates the expression of hundreds of genes.

Thus, we believe it is more likely to be transferable to different strain backgrounds and target

substrates. Furthermore, this approach significantly simplifies and expedites the design-build-test

cycle for synthetic heterotrophy and demonstrates the intrinsic adaptability of yeast toward growth on

non-native substrates. We expect that this insight and approach will expand the utility of this yeast for

valorizing current and emerging waste and/abundant substrates.

DATA AVAILABILITY

RNA-seq sequencing data has been submitted to NCBI SRA and is available under accession

PRJNA837644. R script for mutual information network is available at https://github.com/nair-

lab/yeast-MINet.

ACKNOWLEDGEMENTS

The authors would like to thank current and former Nair lab members, Dr. Todd C. Chappell, and Dr.

Karishma Mohan for helpful discussions. This work was supported by NIH grant #DP2HD91798, NSF

grant #1935354, and Tufts Launchpad | Accelerator to N.U.N.

AUTHOR CONTRIBUTIONS

N.U.N., V.D.T., and D.C. conceived and designed the research project. V.D.T., S.F.S., D.C., and

N.U.N. co-wrote the manuscript. V.D.T., S.F.S., V.E.G., and T.H. performed the experiments. V.D.T.,

S.F.S., and N.U.N. analyzed the data. All the authors have reviewed the manuscript and approved it

for submission.

COMPETING INTERESTS

None.

MATERIALS AND METHODS

Strains and plasmids.

The list of plasmids and strains used are listed in Supplementary **Table S1 and S2**. Strain W303-1a (*MATa leu2-3,112 trp1-1 can1-100 ura3-1 ade2-1 his3-11,15*) and plasmids pIS374, pIS376, pIS385 were obtained from Euroscarf (Germany). The plasmids were constructed in the present using NEB-HiFi DNA assembly master mix from NEB (Beverly, MA).

Growth studies.

Overnight inoculums were grown in the required dropout SC medium (6.7 g/L yeast nitrogen base without amino acids, 2 g/L dropout mix) with sucrose (2 %). The culture was washed twice in the growth medium and resuspended at an initial OD_{600} of 0.1 with appropriate sugar (2 %) in 250 mL shake flasks containing 20 mL of media. OD_{600} measurement was checked at frequent time intervals (3-6 h) on SpectraMax M3 spectrophotometer (Molecular Devices). Growth rate was determined by plotting the values in GraphPad Prism following non-linear regression and using exponential growth equation, $Y = Y_0 exp(kX)$.

Directed evolution.

Error prone PCR libraries of arabinose cassettes (pVDT14, pVDT38-42) were generated as described earlier (*18*). Six-barcoded primers (**Table S4**) were used to track the proportion of the lineage during the enrichment. The amplicons from error prone PCR were assembled into pRS413 plasmids using yeast gap-repair cloning. To attain high library size, the linear fragments were transformed into yeast by electroporation as described previously (*42*). We attained library size of ~10^8 CFUs which was then subjected to enrichment on arabinose in 2YP (20 g/L yeast extract, 40 g/L peptone, 100 mg/L adenine hemisulfate) as well SC media. The cell pellets from each passage were frozen for quantifying barcodes using amplicon sequencing (Genewiz, Cambridge, MA). At the end of enrichment, the library pool was plated on 2YP+Ara and SC+Ara and 18 colonies were randomly picked for growth rate determination. The plasmids from best variants were isolated from strain, transformed into *E. coli*. The corresponding plasmid isolated from *E. coli* was sequenced and re-transformed into REG-strain for growth rate determination.

Genomics integrations.

Accessory cassettes (pVDT29, 30, 35, **Table S1**) were integrated into VEG16 (W303-1a, Δ*GRE3*, Δ*GAL1*, Δ*GAL3*, Δ*GAL7*, Δ*GAL10*) (*18*) strain via disintegrator plasmid (*43*). Counterselection was performed on SC medium with 1 g/L 5-FOA to remove the *URA3* marker and the locus was amplified, and sequence confirmed.

Deletion library.

To generate the deletion library, the cassettes for the required targets were amplified from the genomic DNA of the appropriate strains from the KO collection (*44*). Since the cassettes contained *KANMX* marker, the library was selected on YP+G418 (400 µg/mL). The pool was stored as a stock

and used for studying fitness by transforming it with plasmids for galactose, xylose, and arabinose
utilizing cassettes.

qPCR expression analysis.
Total yeast RNA isolation was performed on OD ~ 1 of yeast cells collected from mid-log phase
growth using the GeneJet RNA Purification Kit (Thermofisher Scientific Catalog #: K0731) according
to the instructions for isolating yeast RNA. Two samples were collected for each growth condition.
The concentration of total RNA isolated was obtained by measuring 10-fold dilutions from each
sample on a spectrophotometer. Using these concentrations, 2.5 µg of total RNA was subjected to the
'routine' DNase treatment protocol using the Invitrogen Turbo DNA-free Kit (Thermofisher Scientific
Catalog #: AM1907). Next, 20 % of the final reaction (10-50 µL, targeting 500 ng RNA) was used to
synthesize cDNA for the sample using the Invitrogen SuperScript IV First-Strand Synthesis System
(Thermofisher Scientific Catalog #: 18091050). The random hexamers supplied by the kit were used
to prime the reverse transcriptase enzyme. cDNA samples were diluted 100-fold (2 µL into 198 µL
dH$_2$O), of which 2 µL was used as the template for a 15 µL qPCR reaction using the Applied
Biosystems PowerUp SYBR Green Master Mix (Thermofisher Scientific Catalog #: A25741). Each
sample was tested for the expression level of five genes: the three arabinose catabolic genes (*araA,*
araB, and *araD*) as well as two housekeeping genes (*TFC1* and *UBC6*) using the primers listed in
Table S5. Reactions were run on an Applied Biosystems Quantstudio 5 Real-Time PCR Instrument.
The relative expression level of each catabolic genes was calculated by subtracting the geometric
mean of the cycle thresholds (Ct) of the two housekeeping genes from the Ct of that catabolic gene,
followed by conversion of the log-base two Ct into a non-log value that can be compared across
genes.

RNA-seq.
Transcriptomics of strains WT, XYL-REG, ARA-REG were performed on the mid-log phase cultures
grown on their respective carbon source (galactose, xylose, or arabinose). Cells pellets were washed
twice in water and stored at –80 °C and outsourced to Genewiz Inc. for RNA extraction and
sequencing. RNA-seq was performed on Illumina HiSeq. Raw FASTQ files were processed for
differential expression analysis using GeneiousPrime. The possible adapter sequences and low-
quality short-read (less than 50 bp) trimming were performed using BB-Trim package. The reads
were aligned to the reference genome W303 obtained from Saccharomyces Genome Database
(http://www.yeastgenome.org) using Bow-Tie2 package. The edgeR package was used to normalize
the gene count based on library size and was converted to cpm (counts per million) using. DESEQ2
package was sued for differential gene expression analysis. Genes with p-values < 0.05 and fold
change of ≥ 2 were considered as differentially expressed.

Amplicon sequencing analysis.
The total DNA was extracted and quantified using a spectrophotometer and approximately 100 ng
was used as template for PCRs. Unique, barcoded primers flanked by Illumina sequencing adaptors

were used to generate the amplicons after 15-20 cycles of PCR for sequencing (**Table S6**). The

barcoded samples were pooled and sent for amplicon sequencing (2×250 bp, Genewiz, New Jersey,

USA). For each pooled sample, we received approximately 100,000 sequencing reads. These data

were processed according to a previously described bioinformatic workflow using Geneious Prime®

2020.2.4 (*45*). Briefly, the reads were paired and merged using the BBMerge package and filtered for

poor-quality reads using the BBDuk package. The reads were mapped using BowTie2 and gene

expression was calculated and differential expression was determined using DESeq2 (*46*).

Network analysis.

The *S. cerevisiae* genes that we consider as the 'gold standard' list of transcription factors is the

unique set of genes identified in the file supplemental data **File S1**. Our analysis utilizes three yeast

genetic regulatory networks (EGRIN, YEASTRACT, and CLR) (*21*) in the context of integrating

regulatory and metabolic networks to enable more accurate prediction of yeast phenotypes in different

growth conditions (*47*). The Environment and Gene Regulatory Influence Network (EGRIN) consists

of 92 regulators and 2,588 interactions and was constructed using the cMonkey and Inferelator

computational tools trained on expression data from 2,929 microarray experiments to identify gene

clusters and potential regulatory genes. The YEASTRACT gene regulatory network was extracted

directly from the YEASTRACT database2 and consists of 177 regulators and 31,075 regulatory

associations based on both 'direct evidence' – chromatin immunoprecipitation (ChIP), ChIP-on-chip,

electrophoretic mobility shift assay, or examination of the effect of TF binding site mutations on target

gene expression – as well as 'indirect evidence' provided by gene expression changes in response to

deletion, mutation, or overexpression of a given TF3.

The EGRIN, YEASTRACT, and CLR yeast gene regulatory networks (GRN) were downloaded as

Microsoft Excel files from the supplementary information of a recent study by Wang et al. (*21*). These

excel files captured the GRN as two-column lists of genes with each row representing an inferred

connection between the genes. We imported these files as networks in the open-source network

visualization and analysis software Cytoscape (*22*). For each network, we created spreadsheets that

associates the fold-change in expression and corrected p-value from the differential gene expression

analysis between the pentose (REG) and galactose grown strains with each unique yeast gene found

in that network. By importing these files into Cytoscape, we were able to map this additional

information onto the nodes in the networks. Next, each network was filtered by removing all nodes

corresponding to genes with Benjamini-Hochberg corrected p-values > 0.05. The "Analyze Network"

function was used to generate betweenness centrality (BC), a quantitative measure of the centrality of

a given node in a network, values for each of the remaining nodes. The final list of 24 deletion targets

was manually curated from a combined list of the top 15 genes by BC from each of the three networks

based on which were most likely to be directly related to the metabolism of pentoses. The R

programming language was used for importing and manipulating data. R scripts can be found in

https://github.com/nair-lab/yeast-MINet.

Stressor studies.

We determined fitness of the strains in presence of various stressors encountered during

biomanufacturing using plate assay. Briefly, we tested the growth of by performing spot dilution on

2YP supplemented with 2 % sucrose or arabinose and various concentration of the stressors (**Table**

S7). The plates after incubation were imaged and the colonies were counted to determine the CFUs.

The CFUs in the presence of the stressor was divided by the CFUs in absence of the stressor to

determine the fitness of that strain. This value was normalized to the fitness of the parental strain and

was expressed as the relative fitness.

REFERENCES

1. J. Sun, W. Sun, G. Zhang, B. Lv, C. Li, High efficient production of plant flavonoids by microbial cell factories: Challenges and opportunities. *Metabolic Engineering*, (2022).
2. M. A. Franden *et al.*, Engineering Pseudomonas putida KT2440 for efficient ethylene glycol utilization. *Metabolic engineering* **48**, 197-207 (2018).
3. S. Pontrelli *et al.*, Escherichia coli as a host for metabolic engineering. *Metabolic engineering* **50**, 16-46 (2018).
4. T. Sun, Y. Yu, K. Wang, R. Ledesma-Amaro, X.-J. Ji, Engineering Yarrowia lipolytica to produce fuels and chemicals from xylose: A review. *Bioresource Technology* **337**, 125484 (2021).
5. T. J. Hanly, M. A. Henson, Dynamic flux balance modeling of microbial co-cultures for efficient batch fermentation of glucose and xylose mixtures. *Biotechnology and bioengineering* **108**, 376-385 (2011).
6. S.-M. Lee, T. Jellison, H. S. Alper, Directed evolution of xylose isomerase for improved xylose catabolism and fermentation in the yeast Saccharomyces cerevisiae. *Applied and Environmental Microbiology* **78**, 5708-5716 (2012).
7. L. Diao *et al.*, Construction of fast xylose-fermenting yeast based on industrial ethanol-producing diploid Saccharomyces cerevisiaeby rational design and adaptive evolution. *BMC biotechnology* **13**, 1-9 (2013).
8. D. J. Wohlbach *et al.*, Comparative genomics of xylose-fermenting fungi for enhanced biofuel production. *Proceedings of the National Academy of Sciences* **108**, 13212-13217 (2011).
9. A. Matsushika, H. Inoue, T. Kodaki, S. Sawayama, Ethanol production from xylose in engineered Saccharomyces cerevisiae strains: current state and perspectives. *Applied microbiology and biotechnology* **84**, 37-53 (2009).
10. M. Lee, H. J. Rozeboom, E. Keuning, P. de Waal, D. B. Janssen, Structure-based directed evolution improves S. cerevisiae growth on xylose by influencing in vivo enzyme performance. *Biotechnology for Biofuels* **13**, 1-18 (2020).
11. Z. Dai, J. Nielsen, Advancing metabolic engineering through systems biology of industrial microorganisms. *Current opinion in biotechnology* **36**, 8-15 (2015).
12. K. V. Presnell, H. S. Alper, Systems metabolic engineering meets machine learning: A new era for data-driven metabolic engineering. *Biotechnology journal* **14**, 1800416 (2019).
13. X. Li *et al.*, Metabolic network remodelling enhances yeast's fitness on xylose using aerobic glycolysis. *Nature Catalysis* **4**, 783-796 (2021).
14. L. Sun, Y. S. Jin, Xylose assimilation for the efficient production of biofuels and chemicals by engineered Saccharomyces cerevisiae. *Biotechnology Journal* **16**, 2000142 (2021).
15. X. Wang, J. Yang, S. Yang, Y. Jiang, Unraveling the genetic basis of fast l-arabinose consumption on top of recombinant xylose-fermenting Saccharomyces cerevisiae. *Biotechnology and Bioengineering* **116**, 283-293 (2019).
16. R. Garcia Sanchez *et al.*, Improved xylose and arabinose utilization by an industrial recombinant Saccharomyces cerevisiae strain using evolutionary engineering. *Biotechnology for biofuels* **3**, 1-11 (2010).
17. H. W. Wisselink, M. J. Toirkens, Q. Wu, J. T. Pronk, A. J. van Maris, Novel evolutionary engineering approach for accelerated utilization of glucose, xylose, and arabinose mixtures by engineered Saccharomyces cerevisiae strains. *Applied and environmental microbiology* **75**, 907-914 (2009).
18. V. Endalur Gopinarayanan, N. U. Nair, A semi-synthetic regulon enables rapid growth of yeast on xylose. *Nature communications* **9**, 1-12 (2018).
19. H. W. Wisselink *et al.*, Engineering of Saccharomyces cerevisiae for efficient anaerobic alcoholic fermentation of L-arabinose. *Applied and Environmental Microbiology* **73**, 4881-4891 (2007).
20. Z. Dai, M. Huang, Y. Chen, V. Siewers, J. Nielsen, Global rewiring of cellular metabolism renders Saccharomyces cerevisiae Crabtree negative. *Nature communications* **9**, 1-8 (2018).
21. Z. Wang *et al.*, Combining inferred regulatory and reconstructed metabolic networks enhances phenotype prediction in yeast. *PLoS computational biology* **13**, e1005489 (2017).
22. P. Shannon, Markiel a, Ozier O, et al. cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res* **13**, 1 (2003).
23. C. E. Costa, P. Carvalho, L. Domingues, Strategic combination of different promoters in lactose metabolisation and host chassis selection for high bioethanol titres from dairy wastes. *Bioresource Technology Reports* **19**, 101131 (2022).

24. K. Kuroda *et al.*, Critical roles of the pentose phosphate pathway and GLN3 in isobutanol-
specific tolerance in yeast. *Cell Systems* **9**, 534-547. e535 (2019).
25. E. Young, S.-M. Lee, H. Alper, Optimizing pentose utilization in yeast: the need for novel tools
and approaches. *Biotechnology for biofuels* **3**, 1-12 (2010).
26. E. M. Young, A. D. Comer, H. Huang, H. S. Alper, A molecular transporter engineering
approach to improving xylose catabolism in Saccharomyces cerevisiae. *Metabolic*
engineering **14**, 401-411 (2012).
27. J. G. Nijland, A. J. Driessen, Engineering of pentose transport in Saccharomyces cerevisiae
for biotechnological applications. *Frontiers in Bioengineering and Biotechnology*, 464 (2020).
28. B. Wiedemann, E. Boles, Codon-optimized bacterial genes improve L-arabinose fermentation
in recombinant Saccharomyces cerevisiae. *Applied and environmental microbiology* **74**, 2043-
2050 (2008).
29. V. Endalur Gopinarayanan, N. U. Nair, Pentose metabolism in Saccharomyces cerevisiae: the
need to engineer global regulatory systems. *Biotechnology journal* **14**, 1800364 (2019).
30. S. R. Kim *et al.*, Rational and evolutionary engineering approaches uncover a small set of
genetic changes efficient for rapid xylose fermentation in Saccharomyces cerevisiae. *PloS*
one **8**, e57048 (2013).
31. J. Usher *et al.*, Chemical and Synthetic Genetic Array Analysis Identifies Genes that
Suppress Xylose Utilization. (2011).
32. Y. Shen *et al.*, An efficient xylose-fermenting recombinant Saccharomyces cerevisiae strain
obtained through adaptive evolution and its global transcription profile. *Applied Microbiology*
and Biotechnology **96**, 1079-1091 (2012).
33. L. V. Dos Santos *et al.*, Unraveling the genetic basis of xylose consumption in engineered
Saccharomyces cerevisiae strains. *Scientific Reports* **6**, 1-14 (2016).
34. N. S. Parachin, O. Bengtsson, B. Hahn‐Hägerdal, M. F. Gorwa‐Grauslund, The deletion of
YLR042c improves ethanolic xylose fermentation by recombinant Saccharomyces cerevisiae.
Yeast **27**, 741-751 (2010).
35. J. Zha, M. Shen, M. Hu, H. Song, Y. Yuan, Enhanced expression of genes involved in initial
xylose metabolism and the oxidative pentose phosphate pathway in the improved xylose-
utilizing Saccharomyces cerevisiae through evolutionary engineering. *Journal of Industrial*
Microbiology and Biotechnology **41**, 27-39 (2014).
36. C. F. Wahlbom, R. R. Cordero Otero, W. H. van Zyl, B. r. Hahn-Ha☐gerdal, L. J. Jo☐nsson,
Molecular analysis of a Saccharomyces cerevisiae mutant with improved ability to utilize
xylose shows enhanced expression of proteins involved in transport, initial xylose metabolism,
and the pentose phosphate pathway. *Applied and environmental microbiology* **69**, 740-746
(2003).
37. J. T. Cunha, T. Q. Aguiar, A. Romaní, C. Oliveira, L. Domingues, Contribution of PRS3, RPB4
and ZWF1 to the resistance of industrial Saccharomyces cerevisiae CCUG53310 and PE-2
strains to lignocellulosic hydrolysate-derived inhibitors. *Bioresource Technology* **191**, 7-16
(2015).
38. A. Romaní, F. Pereira, B. Johansson, L. Domingues, Metabolic engineering of
Saccharomyces cerevisiae ethanol strains PE-2 and CAT-1 for efficient lignocellulosic
fermentation. *Bioresource technology* **179**, 150-158 (2015).
39. K. Traff, R. Otero Cordero, W. Van Zyl, B. Hahn-Hagerdal, Deletion of the GRE3 aldose
reductase gene and its influence on xylose metabolism in recombinant strains of
Saccharomyces cerevisiae expressing the xylA and XKS1 genes. *Applied and environmental*
microbiology **67**, 5668-5674 (2001).
40. J. T. Cunha, P. O. Soares, A. Romaní, J. M. Thevelein, L. Domingues, Xylose fermentation
efficiency of industrial Saccharomyces cerevisiae yeast with separate or combined xylose
reductase/xylitol dehydrogenase and xylose isomerase pathways. *Biotechnology for Biofuels*
12, 1-14 (2019).
41. J. T. Pronk, Auxotrophic yeast strains in fundamental and applied research. *Applied and*
environmental microbiology **68**, 2095-2100 (2002).
42. J. R. Thompson, E. Register, J. Curotto, M. Kurtz, R. Kelly, An improved protocol for the
preparation of yeast cells for transformation by electroporation. *Yeast* **14**, 565-571 (1998).
43. I. Sadowski, T. C. Su, J. Parent, Disintegrator vectors for single‐copy yeast chromosomal
integration. *Yeast* **24**, 447-455 (2007).
44. G. Giaever *et al.*, Functional profiling of the Saccharomyces cerevisiae genome. *nature* **418**,
387-391 (2002).

45.   V. D. Trivedi *et al.*, In-Depth Sequence–Function Characterization Reveals Multiple Pathways
636   to Enhance Enzymatic Activity. *ACS Catalysis* **12**, 2381-2396 (2022).
637   46.   M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for
638   RNA-seq data with DESeq2. *Genome biology* **15**, 1-21 (2014).
639   47.   M. J. Herrgård, B.-S. Lee, V. Portnoy, B. Ø. Palsson, Integrated analysis of regulatory and
640   metabolic networks reveals novel regulatory mechanisms in Saccharomyces cerevisiae.
641   *Genome research* **16**, 627-635 (2006).
642