1   **Title: Vitamin D constrains inflammation by modulating the expression of key genes on**

2   **Chr17q12-21.1**

Authors: Ayşe Kılıç[1*], Arda Halu[1*#], Margherita De Marzio[1,2], Enrico Maiorino[1], Melody G.

Duvall[3], Thayse Brueggemann[3], Joselyn J. Rojas Quintero[3], Robert Chase[1], Hooman

Mirzakhani[1], Ayse Özge Sungur[4, 5], Janine Koepke[5], Taiji Nakano[6], Hong Yong Peh[3], Nandini

Krishnamoorthy[3], Raja-Elie Abdulnour[3], Katia Georgopoulos[7], Augusto A. Litonjua[8], Marie B.

Demay[9], Harald Renz[10,11], Bruce D. Levy[3] and Scott T. Weiss[1*#]

Affiliations:

[1] Channing Division of Network Medicine, Department of Medicine, Brigham and Women's

Hospital and Harvard Medical School; Boston, MA, USA.

[2] Department of Environmental Health, Harvard TH Chan School of Public Health; Boston, MA,

USA.

[3] Division of Pulmonary and Critical Care Medicine, Department of Medicine, Brigham and

Women's Hospital and Harvard Medical School; Boston, MA, USA.

[4] Behavioral Neuroscience, Experimental and Biological Psychology, Philipps-University

Marburg; Marburg, Germany.

[5] Excellence Cluster Cardio-Pulmonary System (ECCPS), Justus Liebig University Giessen;

Giessen, Germany.

[6] Department of Pediatrics, Graduate School of Medicine, Chiba University; Chiba, Japan.

[7] Cutaneous Biology Research Center, Massachusetts General Hospital and Harvard Medical

School; Charlestown, Massachusetts, USA.

[8] Division of Pediatric Pulmonary Medicine, Golisano Children's Hospital at Strong, University

of Rochester Medical Center; Rochester, NY, USA.

[9] Endocrine Unit, Massachusetts General Hospital and Harvard Medical School; Boston, MA,

USA.

[10] Institute of Laboratory Medicine and Pathobiochemistry, Molecular Diagnostics, Philipps

University of Marburg and German Center for Lung Research (DZL); Marburg, Germany.

[11] Department of Clinical Immunology and Allergology, Laboratory of Immunopathology

Sechenov University; Moscow, Russia.

*These authors contributed equally to the manuscript.

#Corresponding author. Arda Halu (rehal@channing.harvard.edu), Scott T. Weiss

(restw@channing.harvard.edu)

38   **Abstract:** Vitamin D possesses immunomodulatory functions and vitamin D deficiency has been

39   associated with the rise in chronic inflammatory diseases, including asthma (1). Vitamin D

40   supplementation studies do not provide insight into the molecular genetic mechanisms of vitamin

41   D mediated immunoregulation. Here we provide evidence for vitamin D regulation of two human

chromosomal loci, Chr17q12-21.1 and Chr17q21.2, reliably associated with autoimmune and chronic inflammatory diseases (2–4). We demonstrate increased vitamin D receptor (VDR) expression in mouse lung CD4+ Th2 cells, differential expression of Chr17q12-21.1 and Chr17q21.2 genes in Th2 cells based on vitamin D status and identify the IL-2/Stat5 pathway as a target of vitamin D signaling. Vitamin D deficiency caused severe lung inflammation after allergen challenge in mice that was prevented by long term prenatal vitamin D supplementation. Mechanistically, vitamin D induced the expression of the *Ikzf3* encoded protein Aiolos to suppress IL-2- signaling and ameliorate cytokine production in Th2 cells. These translational findings demonstrate mechanisms for the immune protective effect of vitamin D in allergic lung inflammation with a strong molecular genetic link to the regulation of both Chr17q12-21.1 and Chr17q21.2 genes and suggest further functional studies and interventional strategies for long-term prevention of asthma and other autoimmune disorders.

One Sentence Summary: Vitamin D regulates two human chromosomal loci, Chr17q12-21.1 and Chr17q21.2, that are associated with autoimmune and chronic inflammatory diseases.

INTRODUCTION

The vitamin D receptor was genetically associated with asthma in 2004 (5) and high intake of vitamin D by pregnant women is associated with about a 50% reduction in asthma risk in the mother's offspring (6,7). Together, these findings led to a comprehensive theory about how vitamin D deficiency could influence asthma occurrence through its effects on lung and immune system development and that progressive decreases in vitamin D intake from 1946 onward could be contributing to the epidemic of allergic and autoimmune diseases (1). Vitamin D deficiency is

the most common vitamin deficiency in the world today and is particularly prevalent in pregnant

women where the fetal lung and immune system are developing in utero (8–10).

To determine the impact of vitamin D supplementation, we have previously performed a clinical

trial, the Vitamin D Antenatal Asthma Trial (VDAART) in pregnant women who either had

allergies or asthma or had family members with these conditions. The participants were given

either 4400 IU of vitamin D3 or 400 IU and were followed throughout their pregnancy and for the

first six years of the life of the child (11). The results of the trial were not statistically significant

using conventional intent to treat analysis (12,13) but the reasons were complex, as is often seen

when the effects of nutrients are studied. Most importantly, unlike conventional drug trials where

one compares drug to no drug or alternative drug, in nutrient trials there is nutrient already present

in the placebo group. This creates misclassification that can reduce trial power. When we

performed a meta-analysis of the two pregnancy related trials of vitamin D, we got a statistically

significant reduction in asthma in the offspring of women who had the higher vitamin D intake

during pregnancy and, when we adjusted for the baseline level of vitamin D in the meta-analysis,

we got a reduction in asthma risk of 50 percent, exactly what we saw in the observational studies

performed previously (14,15).

To link the results of the VDAART trial directly to the chr17q12 locus, our group then genotyped

the single nucleotide polymorphism (SNP) rs12936231, located in ZPBP2, the gene adjacent to

ORMDL3, and stratified the VDAART trial results by maternal genotype at this locus. This study

found that the vitamin D effect in the trial was significantly influenced by genotype, with the GG

genotype exhibiting a protective effect on asthma risk in the child and the CC or GC genotype not

being responsive to vitamin D; additionally this result was related to sphingolipid production with

the children protected from asthma having higher sphingolipid production (3).

This SNP (rs12936231) is one of two SNPs (the other being rs4065275) that alter the chromatin

state of a regulatory domain, specifically two CTCF binding sites, in the chr17q12-21.1 locus that

are correlated with the expression of ORMDL3 (4,16). 4C-seq assays previously demonstrated that

the ORMDL3 promoter interacts with a long-range enhancer in IKZF3 that promotes (or represses)

transcription of ORMDL3 in cells expressing both genes and the binding of CTCF per the G allele

of rs12936231 in ZPBP2 blocks this interaction, resulting in reduced transcription of ORMDL3

on haplotypes with the rs12936231-G allele (4). If rs12936231 is not activated, then there is

activation of the other CTCF binding site (rs4065275) intronic to ORMDL3, thus favoring the

expression of ORMDL3 and increased asthma risk (4).

Mouse models have contributed to our understanding of ORMDL3 function as mice expressing

the ORMDL3 transgene exhibit spontaneous increases in airway hyper responsiveness (AHR), the

essential feature of asthma. This increase in AHR was associated with airway remodeling and peri

bronchial fibrosis without airway inflammation (17). ORMDL3 is pleotropic, influencing Ca^{++}

signaling, and the unfolded protein response. Most importantly, increased expression of ORMDL3

inhibits the enzyme serine palmityl-transferase, the rate limiting step in the production of

sphingolipids, and ORMDL3 transgenic mice have reduced levels of sphingolipids (18). What has

been unclear so far is how ORMDL3 is controlled.

In the current manuscript we provide evidence for vitamin D regulation of two human

chromosomal loci, Chr17q12-21.1 and Chr17q21.2, reliably associated with autoimmune and

chronic inflammatory diseases. We demonstrate increased vitamin D receptor (VDR) expression

in mouse lung CD4+ Th2 cells, differential expression of these Chr17q12-21.1 and Chr17q21.2

genes in Th2 cells based on vitamin D status and identify the IL-2/Stat5 pathway as a target of

vitamin D signaling. Vitamin D induced the expression of the *Ikzf3* encoded protein Aiolos to

suppress IL-2- signaling and ameliorate cytokine production in Th2 cells thus demonstrating a

mechanism for how vitamin D influences autoimmunity.

MATERIALS AND METHODS

In silico analysis of VDR binding sites on Chr17q12-21

For our *in-silico* analysis, we focused on two genomic regions that are known to harbor variants

associated with asthma and autoimmune disease susceptibility. The first one is a 240 kb region

that covers parts of the 17q12 and 17q21.1 loci and contains IKZF3, ZPBP2, ORMDL3, GSDMA

and GSDMB (chr17: 37,899,254–38,139,253; hg19) and the second one is a 420 kb region in the

17q21.2 locus that includes STAT3, STAT5A and STAT5B (chr17: 40,330,000-40,750,000;

hg19). When selecting these genomic ranges, our criterion was to make the window inclusive of

any potential enhancer regions targeting the above genes of interest. We used the GeneHancer (19)

track in the UCSC Genome Browser (http://genome.ucsc.edu) to determine predicted enhancer-

target gene interactions. We obtained VDR, IKZF3, CTCF and Vitamin D-sensitive CTCF binding

sites from (20–23), converted their genomic coordinates to hg19 if a different genome build was

used in the original publication, and added them as custom tracks on the Genome Browser. We

retrieved the known disease and trait associations in these loci from the NHGRI-EBI GWAS

Catalog (downloaded on 7/26/2021) (24) and retained all hits that were deemed significant ($p<10^{-5}$

). We downloaded significant single-tissue eQTLs for IKZF3 and PSMCIP3 in all tissues from

the Common Fund (CF) Genotype-Tissue Expression Project (GTEx) Analysis Release V8

(dbGaP   Accession   phs000424.v8.p2),   through   the   GTEx   Portal

(https://www.gtexportal.org/home/) in July 2021. We calculated linkage disequilibrium (LD)

using the NCI LDmatrix Tool (https://ldlink.nci.nih.gov/?tab=ldmatrix), and by selecting all

available populations.

The Vitamin D Antenatal Asthma Trial (VDAART)

The design, eligibility and initial results of the trial have been published (11–17). Pregnant

women were recruited from 3 clinical sites across the United States – Boston Medical Center,

Boston, MA; Washington University at St. Louis, St. Louis, MO; and Kaiser Permanente

Southern California Region, San Diego, CA. The Data Coordinating Center was based in the

Channing Division of Network Medicine, Brigham and Women's Hospital, Boston, MA.

Eligible participants were women between the ages of 18 and 39 years, who presented between

the estimated gestational ages of 10 and 18 weeks; who had a history of asthma, eczema, or

allergic rhinitis, or whose partner (biologic father of the child) had a history of asthma, eczema,

or allergic rhinitis; who was a non-smoker; and who was English- or Spanish- speaking, with

intent to participate for 4 years (up to the 3rd birthday of the child). The VDAART study protocol

was approved by the institutional review boards at each participating institution and at the

Brigham and Women's Hospital. All women provided a written informed consent. The trial was

registered with clinical trials.gov: clinicaltrials.gov Identifier: NCT00920621.

Gene expression analysis in VDAART cohort

To understand the putative mechanism linking IKZF3 expression and Vitamin D to asthma risk

we considered umbilical cord blood (CB) RNA-seq data from the VDAART trial. **Figure 1-figure**

supplement 1C shows the study design. Total RNA was isolated from cord blood using the

QIAGEN PAXgene Blood RNA Kit (Qiagen) according to the manufacturer's protocol. Quality

was assessed using the Nanodrop 8000 spectrophotometer (ThermoFisher Scientific). Sequencing

libraries were constructed with the TruSeq® Stranded Total RNA Library Prep Globin Kit

(Illumina) and the NEXTflex Small RNA sequencing Kit v3 (Boo Scientific). Sequencing was

performed using a NextSeq 550 instrument (Illumina). Trimmed reads were mapped to The

GRCh38 reference genome using STAR (25). Read counts were computed with htseq (26). The

data includes 443 expression profiles of pregnant women with and without asthma undergoing

measurements of Vitamin D levels in their CB. Each sample is matched with the observed patterns

of wheezing and asthma in the newborn. We filtered all the samples where information about

Vitamin D levels, asthma/wheezing patterns or other covariates were missing, obtaining 393

expression profiles. Data was normalized with the voom function of the edgeR R package (27),

and Vitamin D levels were binarized using 30 ng/ml as threshold. We performed logistic regression

of asthma status as a function of the expression levels of IKZF1 and IKZF3, including their

interaction term and the interaction between IKZF3 and Vitamin D. Since IKZF3 forms

homodimers and heterodimers with IKZF1, we included IKZF1 in the model. We adjusted the

model for maternal asthma status and child race.

Mice

All animal studies were approved by the institutional animal care committee. Mice were

maintained in a virus- and parasite-free animal facility under a 12-h light, 12-h dark cycle.

C57BL6/J wild type (WT) mice were bred and housed at the MGH animal facility. Vitamin D-

deficient mice (C57BL6/J) were bred and maintained in a UV-free environment and weaned onto

a similar diet lacking vitamin D metabolites (TD97340; Harlan Teklad) that results in undetectable

circulating 25-hydroxyvitamin D levels (28). VDR$^{-/-}$ mice (Vd$^{rtm1Mbd/J}$), on C57BL6/J background,

were bred and housed at the MGH animal facility, weaned at 18 d on a diet that maintains normal

mineral ion homeostasis in the absence of VDR signaling (2% calcium, 1.25% phosphorus, 20%

lactose supplemented diet, TD96348; Harlan Teklad, Madison WI) (29). BALB/c male and female

mice were purchased from Charles River Laboratories at 6–8 weeks of age and fed with diets

supplemented with varying amounts of vitamin-D3 (TD97340 was used as base diet and

supplemented with either 400 IU vitamin D/kg, 1000 IU vitamin D/kg or 4000 IU vitamin D/kg).

Mice were bred on the specific diets for two generations to assure stabilization of vitamin D levels.

Resulting F2 litters were used for the experiments. Mice were housed under standard conditions

with free access to rodent chow and water. All animal experiments were carried out in accordance

with NIH Guidelines for the Care and Use of Laboratory Animals as well as guidelines prescribed

by the Institutional Animal Care and Use Committee (IACUC) at Brigham and Women's Hospital

(2016N000357), Massachusetts General Hospital (2004N000113) and Harvard Medical Area

(05115) (AAALAC 1729).

House dust mite induced-airway inflammation

Mice were exposed to D. pteronyssinus extracts (Greer Laboratories, 10.52 EU/mg endotoxin) via

the intranasal application of HDM (25 μg in saline) on days 1,2,3, 8 to 15. Control mice received

equal volumes of saline (Hospira Inc., Lake Forest, IL, USA). Twenty-four hours after the final

challenge, mice were sacrificed, and specimens sampled.

Bronchoalveolar Lavage

Bronchoalveolar Lavage (BAL) was performed with 2 × 1 mL of PBS/0.02% bovine serum

albumin (BSA). The total number of leukocytes was determined using a TC20 Automated Cell

Counter (BioRad Systems). To differentiate cell types, cytospins were prepared and stained with

Kwik-Diff solutions as per manufacturer's instructions (Fisher Scientific) and 200 immune cells

per slide were counted.

Histological staining

Lungs were perfusion fixed with zinc fixative (BD Pharmingen, San Diego, CA) Tissues were paraffin embedded, divided into sections, and stained with hematoxylin and eosin by MGH DF/HCC Specialized Histopathology Services Core Histology Core. After paraffin embedment, 3-μm sections were stained with hematoxylin and eosin.

Measurements of serum immunoglobulins

Blood samples were collected 24 hours after the final aerosol challenge and serum was prepared. Total IgE levels were measured by ELISA (Chondrex) according to manufacturer's instructions.

Dissociation of mouse tissues, in vitro polarization of CD4 Th-subsets and flow cytometry analyses

Lungs were perfused via the right heart ventricle with PBS, excised, and digested with collagenase D (2 mg/ml; Roche) in RPM 1640 (Sigma-Aldrich) for 30 minutes at 37°C. The cell suspension was washed twice with PBS. Erythrocytes were removed by hypotonic lysis. Cells were washed twice with PBS and sieved with 30-μm cell strainers (Miltenyi Biotec). The resulting single cells suspension was used to detect leukocyte populations (CD45 BV421 [clone: 104; Biolegend], CD4 PerCPeFluor710 [RM4-5; eBioscience], CD8a APC-Cy7 [53-6.7; Biolegend], CD19 PE-Cy7 [6D5; Biolegend]) and CD4+ Foxp3+ Treg (CD45 BV421, CD4 PerCPeFluor710, Foxp3 PE [FJK-16s; Thermo Fisher]), VDR AlexaFluor647 (D2K6W; Cell Signaling Technologies), Aiolos PE (8B2; Biolegend) and cytokine production by Th-cells (see below). Spleens were excised and mononuclear cells were isolated using a syringe plunger and 100-μm cell strainers. Red blood cells were removed by hypotonic lysis. Tonicity was restored by the addition of RPMI 1640 (Sigma Aldrich). The resulting cell suspension was washed twice with PBS and used for the isolation of

naïve CD4+ T cells according to the manufacturer's instructions (Naive T Cell Isolation Kit II;

Miltenyi Biotec). Naive CD4+ T cell purity was assessed by staining for CD4 PerCPeFluor710

(clone: RM4-5), CD44 FITC (IM7), and CD62L PE (mel-14). Naïve T cells were activated with

plate bound anti-CD3ε (5 µg/ml, clone: 145-2C11, BD Biosciences) and soluble anti-CD28 (0.5

µg/ml, clone: 37.51, BD Biosciences). Th-subset polarization was induced by exposing naïve

CD4+Tcells to following culture conditions for 3 days. Th1 cells: recombinant murine IL-12

(rmIL-12) (10 ng/mL; Peprotech), anti–IL-4 (5 µg/mL; BD Biosciences); Th2 cells: rmIL-4 (20

ng/mL; Peprotech), anti–IFN-γ (5 µg/mL; BD Biosciences), and anti–IL-12 (5 µg/mL; BD

Biosciences); Treg: rmTGFβ1 (2 ng/mL; Peprotech), anti–IL-4 (5 µg/mL), anti–IFN-γ (5 µg/mL)

for 3 days. Proliferation of Th1 and Th2 cells was induced using rmIL-2 (20 ng/mL; Peprotech) in

the presence of Th-polarizing reagents for an additional 2 days. Polarization efficiency was

verified by staining for intracellular cytokines (please see below) or Foxp3 expression (see above).

For intracellular cytokine staining, cells were stimulated with phorbol 12-myristate 13-acetate

(PMA)/ionomycin and brefeldin A (Cell Stimulation Cocktail; Tonbo Biosiences) for 4 hours.

Cells were fixed with Fixation Buffer (Biolegend). Cells were permeabilized using 0.3% saponin

in FACS buffer (0.5% FBS in PBS). Antibody staining was performed for 30 minutes in the dark

using CD4, IFN-γ (XMG1.2-FITC), IL-10 (JES5-16E3-PE), IL-13 (eBio13A-eFluor660), IL-17A

(TC11-18H10.1-PE-Cy7). Cells were washed twice with permeabilization buffer and suspended

in FACS buffer. For analysis, 30,000 counts (in the CD4+ gate) were recorded on a FACSFortessa

(BD Biosciences) and analyzed with FlowJo software (v10; Tree Star).

Western blot analyses

Cells were lysed in 50 mmol/L Tris/HCl (pH 7.5), 150 mmol/L NaCl, 1% NP-40, and 1× protease

and phosphatase inhibitor mix (both Roche) for 15 minutes on ice. Lysates were clarified, and total

protein concentrations were measured with BCA (Pierce). Proteins (20 µg) were separated on gradient gels (Mini-PROTEAN® TGX; BioRad) and transferred on a Immun-Blot PVDF membrane (BioRad). Unspecific binding was blocked using 5 % milk powder in TBST for 30 min. Anti-VDR (D2K6W, 1:1000; Cell Signaling Technologies, CST) was incubated in 0.5 % milk powder in TBST overnight at 4° C. Membranes were washed 3 times with TBST and incubated with secondary anti-rabbit IgG, HRP-linked Antibody (#7074; CST) for 2 hours. To control for equal loading, membranes was probed for β-actin (AC-15; 1:10,000; Sigma-Aldrich) expression for 2 hrs. at RT. Protein bands were visualized with Luminol reagent (Pierce) using the ChemiDoc XRS+system (BioRad).

Immunofluorescence staining for confocal microscopy

In vitro polarized Th2 cells (see above) were stimulated with 100 nm calcitriol (in ethanol) or ethanol [equal v/v] for the final 24 hours of culture. Cells were washed twice with PBS and spun onto poly-L-lysine coated glass slides. After drying, cells were fixed with 4 % PFA in PBS for 15 min. Cells were permeabilized using 0.2% Triton X-100 in PBS for 20 min. Cells were washed 3 times for 5 min with PBS and incubated with Image-iT FX Signal Enhancer (Thermo Fisher Scientific) for 30 min at RT. After three consecutive washes with PBS, unspecific binding was blocked using blocking buffer (PBS/5 % goat serum/0.3 % Triton X-100) for 60 min at RT. Primary antibodies directed against CD4-FITC (clone; 1:100, BD Biosciences), VDR (D2K6W; 1: 200, CST) were incubated in a humidified chamber over night at 4° C. After 3 washes with PBS, slides were incubated with the secondary reagent, anti-rabbit-Cy3 (1:500; Thermo) for 30 min at RT in the dark. After three more washes with PBS, slides were mounted with ProLong Gold with DAPI according to manufacturer's instructions (Invitrogen). Images were acquired in a single z-plane using a Zeiss immunofluorescence microscope with a 40x objective.

Total RNA isolation, quality control, and RNA-sequencing analysis

Total RNA was isolated using TRIzol™ Reagent (Invitrogen) according to the manufacturer's instructions. Quality was assessed using the Nanodrop 8000 spectrophotometer. Sequencing libraries were constructed with the TruSeq® Stranded Total RNA Library Prep Globin Kit (Illumina). Sequencing was performed using a HiSeq. 2500 instrument (Illumina). Trimmed reads were mapped to the GRCm38 reference genome using STAR (25). Read counts were computed with htseq (26). Data were normalized and differential expression was analyzed with the R package DESeq2 (v.1.34.0)(30) using a false discovery rate (FDR) < 0.05.

Pathway enrichment, Overrepresentation and Gene Set Enrichment analysis

KEGG pathway, overrepresentation and gene set enrichment analyses were performed with the "enrichKEGG", "enrichr" and "GSEA" function in the R package "clusterProfiler" (v.3.10.1) with default settings(31). P-values were adjusted using the BH-FDR correction and enriched terms with a p-adj.<0.05 selected.

Network analysis in the mice

RNA-Sequencing data extracted from murine Th2 cells were analyzed using the R package DESeq2 (v.1.22.2) (30). Raw counts were normalized and transformed for variance stabilization using the DESeq2 function rlog (regularized-logarithm transformation). Batch effects were removed using the function removeBatchEffect of the R package LIMMA (32). Gene with counts lower than 5 in at least one sample were removed for downstream analyses. Differential expression analysis was performed between control Th2 cells versus calcitriol stimulated Th2 cells using DeSeq2. The design matrix was built to consider batch effects derived from the RNA isolation

process. DESeq2-independent filtering was applied (30) and P values were corrected for multiple

hypothesis testing by using the Benjamini-Hochberg (BH)-FDR adjustment (33).

The PPI network used in our network analysis was developed by Silverbush et al. (34). As

described in the original paper, this human PPI network was reconstructed by integrating large

scale PPI databases, drug response repositories, and cancer genomics data. Only the largest

connected component of the network was considered, resulting in an interactome of 15,500

proteins and 234,585 consensus-oriented interactions. Conversion between murine and human

gene IDs was performed using the BioMart conversion tool (35). We also assumed a one-to-one

correspondence between genes and their protein products in the entire analysis.

To identify the active signaling paths in control and calcitriol stimulated Th2 cells, we modelled

the transmission of the molecular signal as a sequential path on the PPI network (called a network

path) that starts from a receptor, connects the receptor to multiple intermediate proteins, and leads

to a transcription factor. Receptors and transcription factors were selected from a predefined list

based on their role in Th2 cell biology and IL-2 expression (**Figure 4-table supplement 1**). CD3,

CD28, IL-4R, IL-13R were selected based on their role in Th2 differentiation. The IL-2R complex

was included due to being a target of vitamin D signaling. TGFβRs were selected based on the

immunomodulatory activity of this signaling path in CD4+ T cells. role in mediating

immunoregulation. Using this subset of receptors and transcription factors as input for the

implementation of our network analysis, we computed all the shortest paths connecting each

receptor-transcription factor pair on the PPI network. For both control and calcitriol stimulated

Th2 cells, we ranked each network path based on the average expression of the path's nodes. The

top selected paths represent the most active downstream processes in Th2 cells in each condition.

mRNA isolation, reverse transcription, and real-time RT-PCR

Total RNA from in vitro polarized Th2 cells was isolated with the RNeasy Mini Kit (Qiagen).

First-strand cDNA synthesis was performed with the Omniscript kit (Qiagen). Quantitative real-

time PCR was performed with the SsoFast EvaGreen Supermix Kit (BioRad) on a AriaMx Real-

time PCR System (Agilent) as per manufacturer's instructions. Primers sequences used in this

study are: Gsdma (sense: 5'-TGC TGT CAA AGG ACG CTA TG-3'; antisense: 5'-TCA ACC

AGC TTC AGT TGC AC-3'), IL-2 (sense: 5'-CCC ACT TCA AGC TCC ACT TC-3'; antisense:

5'-ATC CTG GGG AGT TTC AGG TT-3'), Ikfz3 (5'-TGT CAC CTC TGC AAC TAC GC-3'; 5'-

TTC CGC AGA ACT CAC ACT TG-3'), Ormdl3 (sense: 5'-CAC ACG GGT GAT GAA C AG

TC-3'; antisense: 5'-CTT TGC CTT GGT CTG GAG TC-3'), Stat3 (sense: 5'-GAG GAG CTG

CAG AAA GT-3'; antisense: 5'-TCG TGG TAA ACT GGA CAC CA-3'), Stat5a (sense: 5'-TAC

ATG GAC CAG GCT CCT TC-3'; antisense: 5'-GTC AAA CTC GCC ATC TTG GT-3'), Stat5b

(sense: 5'-TGT GGA TAC AGG CTC AGC AG-3'; antisense: 5'-TGG GTG GCC TTA ATG TTC

TC-3'). Relative gene expression levels were calculated with the $2^{-\Delta\Delta Ct}$ method, with normalization

to the murine ribosomal L32 as the housekeeping gene. Data are presented as fold induction over

the respective control groups (vehicle or WT). All standard procedures were performed according

to the manufacturer's instructions.

Quantification and statistical analysis

For each figure the number of replicates per experiment is indicated in the corresponding figure

legend. In the figures, mean and standard error of the mean (s.e.m.) are presented and error bars

represent the mean ± s.e.m. Two group comparisons were analyzed using unpaired two-tailed

Student's t-tests (parametric data) and Mann–Whitney U-test (non-parametric data). Multiple

group comparisons were analyzed with Kruskal-Wallis and One-way ANOVA test with Tukey's

post hoc analysis. *In vivo* experiments were analyzed using Mixed-effect analysis or Two-way

ANOVA test with Holm-Šidák's post-hoc analysis (factors: genotype & exposure, diet &

exposure). Prism 9.2 (GraphPad Software) was used to calculate statistics. P values of less than

0.05 were considered statistically significant.

RESULTS

Integrative bioinformatics identifies the potential control of key asthma and autoimmune

loci in 17q12-21.1 and 17q21.2 by vitamin D

To address the genetic and molecular mechanisms by which vitamin D can influence asthma,

allergic and autoimmune disease risk, we studied the genetic loci of two chromosome regions of

interest: 1) the 17q12-21.1 region, that includes the well-known asthma-associated *ORMDL3* gene

(36), and 2) the 17q21.2 region, that includes the *STAT5* gene, known to be extensively associated

with autoimmune diseases (37,38). We first examined the entire set of significant disease and trait

associations in the Chr17q12-21.1 (181 associations) and Chr17q21.2 (100 associations) loci using

the NHGRI-EBI GWAS Catalog (24). Both regions had multiple Th1 and Th2 diseases

associations with at least one significant SNP. Chr17q12-21.1 had the highest number of

associations related to Th2 diseases (**Figure 1***A*) and 17q21.2 had the highest number of

associations related to Th1 diseases (**Figure 1-figure supplement 1***A*). To address the possible

link to vitamin D intake, we next investigated a potential overlap of these SNPs with VDR binding

sites. Out of the 169 VDR binding sites on Chr17, seven were in the 17q12-21.1 locus,

concentrated in two bands on or near *IKZF3* and *ZPBP2* (**Figure 1***B*). These VDR binding sites

also overlapped with RXRA binding sites (**Figure 1***B*). To assess their functional relevance, we

searched for expression quantitative trait loci (eQTLs) in these VDR binding regions. We found

that four cis-eQTLs control *IKZF3* expression in whole blood and EBV-transformed lymphocytes:

rs2941522 and rs12946510 in the enhancer region targeting *ORMDL3* and *IKZF3*, and rs1453559

and rs35564481 in the enhancer region targeting *ORMDL3*, *GSDMA* and *GSDMB* (**Figure 1*B***).

Rs1453559 and rs2941522 have previous GWAS associations with both asthma and autoimmune

diseases (39–41), rs12946510 has previous GWAS associations with autoimmune diseases

(42,43), and rs35564481 has no known GWAS associations. For the 17q21.2 locus, the only VDR

binding site resides near the gene *PSMC3IP* in an enhancer site targeting *STAT5A* (**Figure 1-figure**

supplement 1*B*). PSMC3IP expression is controlled by three eQTLs overlapping with this VDR

binding site (rs4793244, rs62078362 and rs111708606), none of which have any known disease

associations. These results suggest that vitamin D binding is critical in this genomic region.

We next examined the linkage disequilibrium (LD) pattern between the Th1/Th2-associated SNPs

and the SNPs in the VDR binding sites (**Figure 1*C***). Out of the seven SNPs with VDR binding

sites identified in 17q12-21.1 and 17q21.2, the first four in 17q12-21.1 (rs2941522, rs12946510,

rs35564481, rs1453559) were in a high-LD block with many other Th1-Th2 associated SNPs in

the same region. These same four SNPs in 17q12-21.1 were also in high-LD with two SNPs in

ZPBP2 and *ORMDL3* (rs12936231 and rs4065275, respectively) that were previously shown to be

functionally relevant to asthma (3,4,16). Additionally, two of the three SNPs with VDR binding

sites in 17q21.2 (rs4793244 and rs62078362) were in strong-LD with three other Th1-Th2

associated SNPs in the17q21.2 region (rs11871801, rs2006141, rs4793090). Overall, these results

highlight strong statistical association between vitamin D binding sites and multiple genotypic

variations linked to asthma, autoimmune and allergic diseases.

Two SNPs in *ZPBP2* and *ORMDL3* (rs12936231 and rs4065275, respectively) that were

previously shown to be functionally relevant to asthma (3,4,16), were also in high linkage

disequilibrium with the four SNPs that we identified in the VDR binding regions of the 17q12-

21.1 loci (**Figure 1*C***). Importantly, in both the 17q12-21.1 and 17q21.2 regions, the VDR binding

sites coincide with CTCF and IKZF3 binding sites, as well as H3K27ac peaks indicating active

enhancer sites (**Figure 1*B*, Figure 1-figure supplement 1*B***). In particular, the two enhancer sites

in 17q12-21.1 (GH17J039753 and GH17J039859) were both predicted to interact with the

ORMDL3 promoter region (**Figure 1*B***), and the enhancer site in 17q21.2 (GH17J042576) was

predicted to interact with the *STAT5A* promoter region (**Figure 1-figure supplement 1*B***). This

suggests a similar mechanism to the one described in (4), whereby SNPs in LD with each other in

these enhancer sites affect VDR and CTCF binding, the co-dependence of which in turn could

modulate *ORMDL3* and *STAT5A* expression. Although it has been shown that CTCF binding sites

can be vitamin D-sensitive (7), no such sites were found in our genomic regions of interest (**Figure**

1*B*, Figure 1-figure supplement 1*B*).

Next, we investigated cord blood gene expression data in the Vitamin D Antenatal Asthma

Reduction Trial (VDAART) (see Methods) and looked at the genes of interest in the 17q12 and

17q21 regions and their interaction with cord blood vitamin D levels as predictors of

asthma/wheeze risk at age 3 year of age. After adjusting for relevant covariates (see Methods), the

baseline expression term of *IKZF3* was not found to be significantly associated with asthma risk,

while the interaction between vitamin D and *IKZF3* was statistically significant (p-val. < 0.05),

suggesting that the association of *IKZF3* expression with asthma risk was mediated by vitamin D

levels in the cord blood (**Figure 1- table supplement 1; Figure 1-figure supplement 1*C***).

Based on our bioinformatic analysis of the Chr17q12 and 17q21 regions, we hypothesize that low

vitamin D tissue levels is associated with decreased *IKFZ3* expression and increased expression

of *ORMDL3*, *STAT3*, *STAT5A*, *STAT5B*, and *IL2*.

Vitamin-D receptor (VDR) is expressed in Th2 cells in allergic airway inflammation

To test our hypothesis of vitamin D activity at the Chr17q12-21 locus we utilized a mouse model

of allergic airway inflammation and identified vitamin D responsive leukocyte subsets in house

dust mite (HDM)-sensitized and challenged mice using flow cytometry (**Figure 2***A*, **Figure 2-**

figure supplement 1*A-C*). VDR expression was significantly induced in HDM sensitized and

challenged mice and almost absent in leukocytes isolated from control (vehicle) lungs (**Figure 2-**

figure supplement 2*C*). Within the CD45+ population, VDR expression was high in CD4+ T cells

(p= 0.0357), while only very few CD8+ T cells and CD19+ B cells expressed VDR (**Figure 2***B*,

Figure 2-figure supplement 1*B* **HDM**). Within the CD4+ T cell population, highest VDR

expression was detected in Gata3+ Th2 cells (p= 0.002 vs Tbet+ and p< 0.0001 vs Foxp3+ T cells).

VDR expression was very low in Tbet+ Th1 and CD4+ Foxp3+ Treg cells (**Figure 2-figure**

supplement 1*C* **HDM**). The differential expression of VDR in Th2 cells was further validated by

western blot analysis using naïve CD4+ T cells immediately after isolation (0hrs), after activation

with CD3/CD28 for 48 hrs and in *in vitro* polarized Th1, Th2 and iTreg cells with well-defined

culture conditions (**Figure 2-figure supplement 2***A*, see Methods section). VDR expression

appeared as early as day 1 of Th2 culture conditions and gradually increased during the

polarization process (**Figure 2-figure supplement 2***B*). In the absence of the VDR ligand vitamin

D, VDR expression localized to the cytoplasm of Th2 cells. Addition of the VDR ligand calcitriol

led to a strong translocation of VDR into the nucleus, suggesting functionality of the expressed

receptor in the presence of its ligand (**Figure 2-figure supplement 2***C*). These results identify Th2

cells as vitamin D targets in allergic airway inflammation.

Vitamin D status regulates the expression of key genes on Chromosome17q12-21.1 and

Chr17q21.2

Taking cell type specific gene expression and selective expression of VDR in Th2 cells into

account, we next analyzed the expression of genes encoded on Chr17q12-21.1 and 17q21.2 loci in

WT and vitamin D deficient Th2 cultures as well as in Th2 cells post calcitriol stimulation.

Vitamin D deficiency differentially regulated the expression levels of several Chr17q12-21.1

encoded genes, with lower expression of *Ikzf3*, *Ormdl3* and *Gsdma* (all p=0.0286) (**Figure 2D**).

Conversely, stimulation with calcitriol induced the expression of the same genes (all p=0.043)

(**Figure 2F**). Expression of the Chr17q21.2 genes *Stat3*, *Stat5a* and *Stat5b* was not affected by

vitamin D deficiency (**Figure 2E**). However, after calcitriol stimulation *Stat5a* (p=0.0082) and

Stat5b (p=0.003) genes were strongly suppressed and *Stat3* (p= 0.0048) expression was increased

in Th2 cells. (**Figure 2G**). These results hint to a protective role of vitamin D in allergic airway

inflammation, by regulating the expression of the IL-2 downstream signaling molecules *Stat5a*

and *Stat5b*.

Deficiency in vitamin D signaling augments allergic airway inflammation

To decipher the protective role of vitamin D signaling in allergic airway inflammation, we

employed vitamin D deficient and VDR$^{-/-}$ mice (**Figure 3A**) (28). HDM sensitization of vitamin

D deficient mice led to a stronger reaction to the allergen resulting in more prominent peri-

bronchial and perivascular leukocytic infiltrates (**Figure 3B**). This phenotype was accompanied

by lower VDR expression in CD4+ T cells isolated from HDM-exposed lungs compared to WT

(**Figure 3C**). Vitamin D deficiency led to a robust increase in total immunoglobulin (Ig)E levels

(**Figure 3D**). Histological findings were associated with higher total BAL leukocyte (p= 0.0008)

(**Figure 3E**), eosinophil (p= 0.0002) (**Figure 3F**) and lymphocyte (p= 0.0023) (**Figure 3G**)

numbers in vitamin D deficient mice, while neutrophil numbers were not affected **Figure 3-figure**

supplement 1*A)*. Th2 cell numbers (p= 0.0127) were also higher in the HDM exposed mice

(**Figure 3*H***) in a manner that was independent of T helper cell skewing (**Figure 3-figure**

supplement 1*B, C*). Differentiation and recruitment of Th17 and IL-10 producing CD4+ T cells

was not affected by vitamin D deficiency (**Figure 3-figure supplement 1*A, B***). To validate these

findings from vitamin D deficiency, we next employed VDR$^{-/-}$ mice (see Methods). As with the

vitamin D deficient mice, HDM sensitization and challenge of VDR$^{-/-}$ mice led to an increased Th2

immune response. The lung phenotype in the VDR$^{-/-}$ mice was marked by dense leukocytic

infiltrates around and mucus plugs in the airways (**Figure 3*B***). The augmented systemic and local

immune response increased total IgE titers (p= 0.0489), and total BAL (p< 0.0001 vs WT HDM

and p= 0.003 vs Vit D deficient HDM), eosinophil (p< 0.0001 vs WT HDM and Vit D deficient

HDM), total lymphocytes (p=0.0194 vs WT HDM and p= 0.0421vs Vit D deficient HDM) and

Th2 cells (p= 0.0009 vs WT HDM) (**Figure 3*D-H***). VDR$^{-/-}$ did not significantly affect recruitment

of neutrophils, Treg and Th17 cells (**Figure 3-figure supplement 1*A, B, C***).

To explore the mechanisms underlying the augmented Th2-phenotype in the lungs of vitamin D

deficient and VDR$^{-/-}$ mice after HDM allergen challenge, we analyzed the transcriptional profiles

of in vitro polarized Th2 cells from WT and vitamin D deficient mice (**Figure 3*I***). Differential

expression and enrichment analysis revealed dysregulation of several pathways implicated in Th2-

cell activation, cytokine production, proliferation, and survival with prominent changes in the IL-

2/ STAT5 - pathway in vitamin D deficient Th2 cells (p= 0.0014) (**Figure 3*J*, Figure 3-table**

supplements 1, 2). The effects of impaired vitamin D signaling on these pathways was confirmed

by qRT-PCR and flow cytometry. Baseline expression of IL-2 was markedly increased at both the

RNA (p= 0.0286)) and protein levels (p= 0.0079) in vitamin D deficient and VDR$^{-/-}$ Th2 cells

(**Figure 3*K, L***). In Vitamin D deficient and VDR$^{-/-}$ Th2 cell *in vitro* incubation, the elevated IL-2

levels were accompanied by a sharp increase in IL-13 production (VitD deficient: p< 0.0001; VDR$^-$

$^{/-}$: p= 0.0141) (**Figure 3-figure supplement 2**).

Taken together, these results indicate a critical role for vitamin D/VDR signaling in regulating type

2 inflammation and identify the IL-2/ Stat5 pathway as a downstream target of vitamin D in Th2

cells.

Vitamin D suppresses the activation of the IL-2/Stat5 pathway and cytokine production in

Th2 cells

To ascertain the vitamin D dependent regulation of the IL-2/Stat5 pathway and the impact on the

effector program of Th2 cells, we first analyzed the transcriptional profile of WT Th2 cells exposed

to calcitriol during differentiation (**Figure 4***A*). Gene ontology enrichment analysis of biological

processes revealed regulation of several processes impacting inflammatory responses, including

chemotaxis and activation of cells (**Figure 4-table supplement 1**). Calcitriol stimulation of Th2

cells affected several immune related disease pathways, including asthma (**Figure 4-figure**

supplement 1, Figure 4-table supplement 2). Gene set enrichment analysis (GSEA), performed

on the differentially expressed genes, highlighted the negative regulation of IL-2/Stat5 pathway in

Th2 cells by calcitriol (**Figure 4B**, **Figure 4-table supplement 3**). Calcitriol stimulation

suppressed the expression of *Il2* (p<0.0001), *Stat5a* (p<0.0008) and *Stat5b* (p<0.0008) genes

(**Figure 4C**). Concordantly, levels of activated Itk (phospho-Itk) (p= 0.029) and expression of the

IL2Rβ (CD25) (p= 0.0441) were reduced on a per cell basis (**Figure 4D**).

Next, the impact of calcitriol stimulation on the effector function of Th2 cells was assessed by

flow cytometric analysis for the T2 cytokine IL-13. Activation of the vitamin D/VDR pathway led

to significant suppression of IL-13 production in CD4+ T cells (p= 0.0012) (**Figure 4*E***).

To further delineate the molecular impact of vitamin D on Th2 cell biology, we integrated the

murine transcriptional profile with a recently developed human Protein-Protein Interaction (PPI)

database (34). By modeling the signal transduction process as a sequential path on the PPI network

that starts from a receptor and propagates downstream to a transcription factor via multiple

intermediate proteins (for details see Methods section, **Figure 4-figure supplement 2*A***), we

inferred the active signaling flow in Th2 cells for control (vehicle) and calcitriol stimulated Th2

cells. As input of our network analysis, we selected a subset of receptors and transcription factors

that are known to be involved in CD4+ T cell activation, Th2 cell differentiation, and cytokine

production (**Figure 4-table supplement 4**). For clarity of the figure, we display the gene names.

Corresponding protein names are attached in **Figure 4-table supplement 5**. In control Th2 cells,

our network analysis revealed the engagement of a variety of receptors including IL4R, IL2RA,

CD28, and TNF-receptors. Downstream to these receptors, the most active signaling paths

highlighted the activation of STAT3, STAT5A/B, NFkB1, and IKZF1 transcription factors

(**Figure 4-figure supplement 2*B***) via multiple adapter proteins of the TRAF and JAK family.

Interestingly, these transcription factors are known to be involved in differentiation and activation

pathways in Th2 cells Conversely, after calcitriol stimulation, signal from IL4R, CD28 and

TNFSF8 propagated over different interaction partners (**Figure 4*F***). Calcitriol specific edges are

highlighted in orange color. Molecular mediators included the cytoskeletal proteins VAV1 and

RACK1 and emphasized activation of the transcription factors AP-1 (FOS/JUND dimer) and

NFATC1.

Our network analysis highlighted a central role of the transcriptional repressor IKZF1 (IKAROS)

upon calcitriol stimulation, which was targeted by multiple downstream mediators including

IKZF3 (AIOLOS). Notably, the IKZF3 gene encodes for the AIOLOS protein, a direct interaction

partner of the IKZF1-encoded IKAROS. Both transcription factors, as homo- and heterodimers,

suppress IL-2 expression on the transcriptional levels in mouse and human lymphocytes (44).

Calcitriol increased the number of edges connecting intermediate proteins to the transcriptional

repressor IKAROS. Among the connecting proteins, AIOLOS interaction with IKAROS was

implicated in repression of IL-2 expression in Th2 cells by vitamin D.

Absence of STAT5A/B mediators in the active signaling paths of calcitriol stimulated Th2 cells

further suggested repression of the Th2 cell STAT5 pathway with calcitriol compared to control

(**Figure 4-figure supplement 2*B***). To ensure that the above findings were not restricted to the

C57BL/6 mouse strain, the inverse experiment was performed in Balb/c mice. This mouse strain

is commonly used for type 2 driven inflammation.

Taken together, these results indicate that the specific suppression of the IL-2/Stat5 pathway by

vitamin D results in reduced cytokine production by Th2 cells. Our findings suggest that this

suppression was mediated by the vitamin D dependent regulation of IKAROS (IKZF1) through

the induction of AIOLOS (IKZF3).

Vitamin D supplementation alleviates the allergic phenotype in the lung by suppressing type

2 cytokine production in a dose dependent manner

To test if vitamin D supplementation prevents the development of T2 driven allergic inflammation

in the lung, custom rodent diets with select vitamin D doses were used: 400 IU/kg chow (400 IU;

low), 1000 IU/kg chow (1000 IU; regular) and 4000 IU/kg chow (4000 IU; high) (**Figure 5- figure**

supplement 1*A,* see Methods section). The 1000 IU group was used as the control group, as it

contains the average amount of vitamin D fortified in standard rodent chow. HDM exposure led

to recruitment of leukocytes into the lung tissue and BAL (**Figure 5***A, B;* **Figure 5- figure**

supplement 1*B*), with a predominant eosinophilic and lymphocytic infiltrate as well as high

numbers of IL-13+ Th2 cells (**Figure 5***C-E*). There were trends for higher cellular infiltration in

the airways when mice were supplemented with the low vitamin D doses (400 IU), but these

inflammatory responses were not significantly different from the phenotype observed with 1000

IU. Lung and airway pathology were similar between these dosing groups, with similar numbers

and frequencies of the different leukocyte populations. In sharp contrast, dietary supplementation

with the higher vitamin D dose (4000 IU) decreased the inflammatory phenotype in HDM exposed

mice compared to the 400 IU and 1000 IU dosed animals. Concordant with the histological

findings, BAL leukocytes ($p< 0.0001$) were decreased with marked reductions in BAL eosinophil

($p=0.0016$), lymphocyte ($p=0.026$) and neutrophils numbers ($p=0.0005$) (**Figure 5***B, D;* **Figure 5-**

figure supplement 1*C*), and the frequency of IL-13+ Th2 cells was significantly reduced in the

lungs with 4000 IU of vitamin D ($p= 0.0066$ vs 400 IU HDM; $p=0.0011$ vs 1000 IU HDM) (**Figure**

5*E*). Of note, dietary supplementation with high vitamin D levels, increased VDR expression in

CD4+ T cells from HDM exposed lungs ($p< 0.0001$) (**Figure 5- figure supplement 1***C*).

To ascertain that dietary supplementation with vitamin D specifically altered mouse Th2 cell

differentiation and the effector program, splenic naïve CD4+ T cells from 400 IU and 4000 IU

mice were studied. Of interest, naïve CD4+ T cells did not express VDR until they were exposed

to Th2 polarizing conditions. Gata3 expression on per cell basis was lower in 4000 IU Th2 cells (p= 0.0286) (**Figure 5- figure supplement 2***A*). This was accompanied by higher VDR protein expression (p= 0.0286) (**Figure 5- figure supplement 2***B*). The expression of IL-2 (p= 0.0004) and the Th2 cytokines IL-13 (p= 0.0159) and IL-5 (p= 0.0159) were lower in 4000 IU Th2 cultures (**Figure 5- figure supplement 2***C-E*).

To explore the specific induction of Aiolos expression for vitamin D mediated suppression of IL-2 production in Th2 cells, Aiolos expression was monitored throughout Th2 differentiation. Aiolos expression gradually increased under Th2 polarizing conditions in CD4+ T cells, starting as early as day 1 of in vitro culture (**Figure 5***F*). Simultaneous exposure to calcitriol further increased Aiolos protein levels compared to vehicle in Th2 cells (p= 0.0079) (**Figure 5***G*).

To test if vitamin D exerts its inhibitory effect on IL-2 production via Aiolos induction, Ikzf3$^{-/-}$ mice were employed, exposed to Th2 polarizing conditions and assessed for IL-2 production after calcitriol exposure. IL-2 production in control Ikzf3$^{-/-}$ Th2 cultures was significantly higher compared to WT Th2 cells. Calcitriol significantly reduced IL-2 levels in both, WT (p= 0.0001) and Ikzf3$^{-/-}$ Th2 cells (p= 0.0001) (**Figure 5***H*); however, the suppressive effect of calcitriol on IL-2 production was significantly higher in WT Th2 cells ($\sim$ 65% inhibition) compared to Ikzf3$^{-/-}$ Th2 cells ($\sim$40% inhibition) (p= 0.0001) (**Figure 5***H*).

Taken together, these results indicated that vitamin D can regulate Aiolos expression and suggested a role for Aiolos as a downstream effector for selectively fine-tuning Th2 cell IL-2 production by vitamin D.

DISCUSSION

We utilized human genetics to identify SNPs on chromosome 17q12-21.1 and 17q21.2 that were

within GWAS regions for Th1 and Th2 autoimmune diseases. Of note, many of these SNPs were

within VDR binding sites. We then determined by eQTL analysis of the genes in these two

genomic regions that IKZF3 expression was controlled by four cis-eQTLs in the enhancer region

targeting ORMDL3 and IKZF3, and in the enhancer region targeting ORMDL3, GSDMA and

GSDMB. Next, we demonstrated that two SNPs in CTCF binding sites were in strong LD with

these 4 eQTL SNPs and that the two enhancer sites in 17q12-21.1 were both predicted to interact

with the ORMDL3 promoter region, and the enhancer site in 17q21.2 was predicted to interact

with the STAT5A promoter region thus, suggesting a plausible human molecular genetic

mechanism by which vitamin D could activate IKZF3 to repress ORMDL3 and immune system

development. Finally, we leveraged bulk RNAseq in the cord blood of the VDAART trial to show

that an interaction for vitamin D level and IKZF3 expression that was associated with reduced

asthma risk at 3 years of age. **Figure 5- figure supplement 3** depicts the influence of vitamin D

binding sites on the genes in the 17q12-21 region.

To confirm our human molecular genetic findings, we chose to utilize mouse models of allergic

lung inflammation. We show strong VDR expression in CD4+ Th2 cells and alterations in

chr17q12-21.1 and 17q21.2 gene expression induced by vitamin D status. Expression of VDR has

been reported for airway epithelial cells and different immune cell types present in the lung (45).

In lymphocytes, *ex vivo* and *in vitro* studies describe low baseline VDR transcript expression levels

that were transiently increased by activating signals and calcitriol stimulation (46). Here, we report

a differential VDR expression pattern in CD4+ T cell subsets from allergic inflamed lungs and

post lineage specific cytokine stimulation *in vitro*, with strongest expression in Th2 cells.

Activation of VDR suppressed IL-2, IL-5, and IL-13 production by Th2 cells. The expression of

VDR in Gata3+Th2 cells in the lung might represent an endogenous control mechanism to curb

the pathogenic cell activity and cytokine production after repetitive allergen contact. *Ex vivo* and

in vitro Th1, nTreg and iTreg cells expressed low VDR protein levels at baseline. It remains to be

determined whether other Th-subsets described in allergic inflammation express VDR at baseline

and if either calcitriol or other secondary signals, specific for each subset, could induce VDR

expression during inflammation.

Vitamin D deficiency that had been introduced over two generations, impaired vitamin D

signaling and led to an augmented allergic phenotype in the lung after allergen exposure with a

dominating Th2-signature locally and systemically. This was likely due to specific deregulation of

Th2 cells instead of Th cell skewing, since neither Th17 nor Treg development and recruitment

were significantly affected. In contrast, in mice supplemented with high vitamin D doses over two

generations, VDR expression and consequently vitamin D responsiveness was higher in CD4+ T

cells from the animals' lungs, almost completely abrogating the inflammatory process in the lung

after allergen challenge. While the vitamin D dose in the mice was chosen to replicate the

VDAART trial dose in the mouse model, the duration of supplementation was adapted to the

protocol for generating vitamin D deficient mice. Both factors might influence the outcome here

compared to prior rodent studies with inconclusive results. We provide a discussion of vitamin D

toxicity in our response to the reviewers. More work to determine the exact levels of vitamin D

that confer this protective effect in the mice would also be worthy of further investigation.

Differentiation of Th2 cells is driven by IL-4 induced Gata3 expression in naïve CD4+ T cells.

This process depends on a positive feedforward loop by IL-2 and its downstream STAT5 signaling

pathway for the effective expression of IL-4 (47). Therefore, either interfering with Gata3

expression or with the IL-2 pathway represents a promising strategy to finetune Th2-driven

immunity. Our data highlight the specific regulation of the IL-2/STAT5 pathway by vitamin D.

Vitamin D deficiency and impaired VDR signaling caused elevated IL-2 production by Th2 cells,

leading to high IL-13 production. Elevated IL-13 levels *in vivo*, could amplify the local T2

responses in the lung. In sharp contrast, exposure of polarizing Th2 cells to calcitriol was able to

suppress IL-2 production, reduce expression and activation of STAT5A/B and therefore reduce

IL-13 production by Th2 cells. The transcription factor AIOLOS has previously been shown to

suppress IL-2 expression at the transcriptional level (44). We report here a vitamin D mediated

induction of Aiolos expression in Th2 cells, that is impaired by vitamin D deficiency. High Aiolos

expression coincided with lower IL-2 production, ameliorated STAT5A/B expression, and reduced

IL-13 production in Th2 cells. *In vitro* Th2 polarization of naïve CD4+ T cells isolated from mice

supplemented with higher vitamin D levels, without further exogenous calcitriol stimulation,

developed a weakened Th2 phenotype, characterized by lower IL-2, IL-5, and IL-13 production.

Since naïve CD4+ T cells do not express VDR, these results, together with our human data,

implicate changes in chromosomal accessibility by high vitamin D in the developing immune

system, which needs further investigation.

Experiments here with Ikzf3$^{-/-}$ mice demonstrated significant reductions in vitamin D mediated

control of Th2 cell cytokine production, but the vitamin D response was not completely

impaired, suggesting additional mechanisms for vitamin D regulation of Th2 cells – several

possibilities were uncovered by our PPI analyses. These will be the subject of future research. In

addition, we only investigated a Th2 mouse model of asthma and have not extended our mouse

work to Th1 models of autoimmunity. We acknowledge that the impact of vitamin D on Th2

biology is conflicting in the literature. While several groups report Th2 promoting activity, we,

and others, show inhibition of type 2 cytokine production (48–51). These discrepancies could be

due to the model system studied, e.g., PBMC and purified CD4+ T cells, or the dose of vitamin

D or the mouse strain.

We have still to consider other asthma genes that might be linked to these two genetic loci as there

are likely other VDR binding sites of importance in both asthma and Th1 autoimmunity. Our

preliminary results confirm that three asthma susceptibility loci: 2q12.1 (IL1RL1), 6p21.32 (HLA-

DQA1/B1/A2/B2) and 22q12.3 (IL2RB) each have VDR and IKZF3 binding sites either in

enhancers predicted by GeneHancer to target these genes or within these genes themselves. It is

important to consider other data that might have enhanced the results that we present here. For our

human genetic data, additional use of ChIP or co-IP to establish STAT induction and activation

would have been of potential value. For our mouse studies, additional cytokine measurements in

the mice as well as measurement of airway resistance would have added to the pathophysiological

data linking IKFZ3 expression to TH2 response. Finally, quantification of histology and confocal

images could provide an overview of VDR expression in the lungs**.**

In summary, we have leveraged clinical trial data, molecular genetic bioinformatic data and mouse

model data to outline for the first time, a comprehensive molecular genetic mechanism for how

vitamin D influences not only asthma development, but both Th1 and Th2 autoimmune disease.

The significance of these findings relates to the high prevalence of vitamin D deficiency, especially

during pregnancy, and the strong possibility that the epidemic of asthma and autoimmunity might

be significantly reduced if serum vitamin D levels were elevated worldwide.

List of Supplementary Materials

Fig S1 to S11

Tables S1 to S9

Acknowledgments: We wish to thank the VDAART trial participants and clinical staff for their critical work on the VDAART study. We wish to thank also our Channing colleagues for their comments.

Funding:

National Heart, Lung, and Blood Institute grant 5P01HL132825

National Heart, Lung, and Blood Institute grant 5UH3OD023268

National Heart, Lung, and Blood Institute grant 5K25HL150336

German Society for Clinical Chemistry and Laboratory Medicine (DGKL) and the

German Research Foundation (DFG) grant KI 1868/3-1

Author contributions:

A.K. and S.T.W. jointly supervised this work. A.K., A.H. and S.T.W. designed the experiments, analyzed data and wrote the manuscript. A.K., M.G.D., T.R.B., J.R.Q., A.O.S, J.K., T.N., H.Y.P., N.K. and R.E.A performed experiments. A.H., E.M., M.D.M. and R. C. analyzed data. M.D. and K.G. provided critical mouse lines and scientific input. A.A.L. assisted on VDAART related analyses. H.R. and B.D.L. supervised experiments and provided critical scientific input. All authors critically reviewed the manuscript.

Competing interests: Dr. Weiss receives royalties from UpToDate and is an investor in Histolix. All other authors declare that they have no competing interests.

Data and materials availability: All data are available in the main text or the supplementary

materials.

Figure legends

Figure 1. VDR binding sites on Chr17q12 and q21.1 overlap with open chromatin signatures.

A, Significant disease and trait associations in 17q12-21.1 (y-axis), retrieved from the NHGRI EBI

GWAS Catalog, with at least one significant SNP (x-axis) in Chr17q12-21.1. Colored bars denote

autoimmune diseases mapped to Th1 (red bars) or Th2 (blue bars) driven immunity. *B*, University

of California Santa Cruz (UCSC) Genome Browser tracks showing chromosomal location,

common dbSNPs, VDR-, RXR-, CTCF- and IKZF3-binding sites and H3K27Ac marks present in

the Chr17q12-21.1 locus. VDR binding sites overlapping with active regulatory elements are

highlighted by colored boxes. *C*, Linkage disequilibrium (LD) between the significant SNPs in the

Chr17q12-21.1 (magenta) and 21.2 (green) region, highlighting Th1/Th2 associated SNPs (red and

blue, respectively), eQTLs in the VDR binding regions in Chr17q12 21.1 and 21.2 (yellow), and

functional asthma SNPs from the literature (cyan).

Figure 1-figure supplement 1. *A***,** Significant disease and trait associations in 17q21.2 (y-axis),

retrieved from the NHGRI-EBI GWAS Catalog, with at least one significant SNPs (x-axis) in

Chr17q21.2. Colored bars denote autoimmune diseases mapped to Th1 (red bars) or Th2 (blue

bars) driven immunity. *B***,** University of California Santa Cruz (UCSC) Genome Browser tracks

showing chromosomal location, common dbSNPs, VDR-, RXR-, CTCF- and IKZF3-binding sites

and H3K27Ac marks present in the Ch17q21.2 locus. VDR binding sites overlapping with active

regulatory elements are highlighted by colored boxes. **C,** Overview of the analyses conducted on

the VDAART transcriptomic and clinical data.

Figure 2. VDR expression is elevated in Th2 cells and vitamin D regulated expression of genes

on Chr1q12 and Chr17q21. A, Scheme of the house dust mite (HDM) induced airway

inflammation protocol. **B,** Flow cytometric analysis and cell frequencies of VDR expression in

CD45+ CD4+ T cells in the respective groups (n=4-5). **C,** Flow cytometric analyses and cell

frequencies of VDR expression in Th1 (Tbet+), Th2 (Gata3+) and Treg (Foxp3+) cells in the

respective groups (n=4-7). Quantitative RT-PCR analysis of relative mRNA expression levels of

genes encoded on Chr17q12-21.1 in (**D**) Th2 cultures of WT and vitamin D deficient mice and (**E**)

control and calcitriol stimulated cultures. Quantitative RT-PCR analysis of relative mRNA

expression levels of genes encoded on Chr17q21.2 in (**F**) Th2 cultures of WT and vitamin D

deficient mice and (**G**) control and calcitriol stimulated cultures. Gene expression levels were

normalized to the house keeping gene L32 and are expressed relative to the expression level in

WT (**D, F**) or WT-vehicle (**E, G**) Th2 cells. (n ≥ 4 per group). Each symbol represents one mouse.

Numbers in flow plots indicate percentages. Error bars indicate the s.e.m. Statistical tests: two-

tailed Mann-Whitney U test (**B, C**). *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$. Data summarize results

from two independent experiments.

Figure 2-figure supplement 1. Baseline VDR expression is confined to the CD4+ Foxp3- Teff

cell population. A, Gating strategy for the analysis of VDR expression in different leukocyte

populations in the lung. **B,** Flow cytometric analysis and cell frequencies of VDR expression in

CD45+, CD4+, CD8+ and CD19+ in the respective groups (n=4-5). **C,** Representative flow plots

of lung CD45+ CD4+ T cells isolated from saline and HDM- immunized and -challenged mice for the co-expression of Foxp3 and VDR. Each symbol represents one mouse. Numbers in flow plots indicate percentages. Error bars indicate the s.e.m. Statistical tests: two-tailed Mann-Whitney U test. *P < 0.05. Data summarize results from two independent experiments.

Figure 2-figure supplement 2. VDR expression is induced during CD4+ Th2 cell differentiation. *A,* Representative immunoblot for VDR expression in freshly isolated (0 hrs) and activated CD4+ CD62L+ CD44- naïve T cells (48 hrs.) and in vitro polarized Th1, Th2 and iTreg cells. Kidney samples of WT and VDR-/- mice were used as controls. *B,* Representative immunoblot for VDR expression in polarizing Th2 at the indicated days of in vitro culture. *C,* Representative image of VDR localization in in vitro polarized Th2 cells stimulated with vehicle (EtOH, [% v/v]) or calcitriol.

Figure 3. Vitamin D deficiency augments allergic airway inflammation. *A,* Schematic presentation of the different mouse strains used. *B,* Representative H&E-stained lung sections from saline and HDM-exposed wild type (C57/BL6; WT), vitamin D deficient and VDR-/- mice (x 10 magnification). Black arrows point to the peri bronchial infiltrate. *C,* Representative flow plots assessing VDR expression in lung CD4+ T cells from the indicated groups. *D,* Total IgE levels detected in the sera of respective groups. *E-G,* Absolute numbers of (*E*) leukocytes, (*F*) eosinophils, and (*G*) lymphocytes in the airways of the respective groups. *H,* Frequencies of IL-13 producing Th2 in the lung. Data summarize results from two or three independent experiments with n>8 per group. *I,* Heat map showing differentially expressed genes between WT and vitamin

D deficient Th2 cultures identified by RNA-Sequencing analysis. Analysis and visualization were

performed with DESeq2. *J,* Overrepresentation analysis of Hallmark gene sets in the differentially

expression gene set. The x-axis represents the gene ratio in the respective gene set. Dot sizes denote

the number of genes in the respective pathway. The dot color indicates the FDR-adjusted p-value.

Analysis and visualization were performed with the clusterProfiler package. *K,* Quantitative RT-

PCR analysis of Il-2 mRNA expression in Th2 cultures from the respective groups (n=4 per group).

Gene expression levels were normalized to the house keeping gene L32 and are expressed as fold-

induction compared to WT Th2 cells. *L,* Flow cytometric analysis and cell frequencies of IL-2

production in WT and VDR-/- Th2 cultures (n=5 per group). Each symbol represents one mouse.

Error bars indicate the s.e.m. Statistical significance was determined with: Mixed-effect analysis

with Holm-Šidák's post-hoc analysis (*D –H*), two-tailed Mann-Whitney U test (*K, L*). *P < 0.05,

**P < 0.01, ***P < 0.001, ****P < 0.0001. Data summarize results from two or three independent

experiments with n>8 per group.

Figure 3-figure supplement 1. Vitamin D deficiency does not alter CD4+ IL-10 and IL-17A

expression allergic airway inflammation. Frequencies of BAL neutrophils (*A*), IL-10 (*B*) and

IL-17A (*C*) CD4+ T cells. Each symbol represents one mouse. Error bars indicate the s.e.m. Data

summarize results from two or three independent experiments with n>8 per group.

Figure 3-figure supplement 2. Vitamin D deficiency increases IL-13 expression in Th2 cells.

Flow cytometric analysis and frequencies of IL-13 expression in WT, vitamin D deficient and

VDR-/- Th2 cultures (n=5-9 per group). Each symbol represents one individual sample and data

summarize the results from two or three independent experiments. Error bars indicate the s.e.m.

Statistical tests: Kruskal- Wallis Test with Dunn's post-hoc analysis. *P < 0.05, **P < 0.01, ****P

< 0.0001.

Figure 4. Vitamin D stimulation of polarizing Th2 cells suppresses the IL-2/Stat5 pathway.

A, Heat map of differentially expressed genes between vehicle (EtOH [% v/v]) and calcitriol

treated WT Th2 cells. Analysis and visualization were performed with DESeq2. *B*, Gene set

enrichment analysis (GSEA) of differentially expressed genes. Analysis and visualization were

performed with the clusterProfiler package. *C*, Quantitative RT-PCR for Il-2, Stat5a and Stat5b

mRNA expression in Th2 cells exposed to either vehicle or calcitriol. Gene expression levels were

normalized to the house keeping gene L32 and are expressed as fold-induction compared to control

cells. *D*, Flow cytometric analysis and mean fluorescence intensities of Itk and Il2rβ expression in

indicated groups. *E*, Flow cytometric analysis and quantification of IL-13 expression in respective

groups. *F*, Network analysis to visualize vitamin D induced molecular changes in the molecular

interactome of calcitriol stimulated CD4+ Th2. Predefined set of cell membrane receptors were

set as starting points (orange nodes) and annotated transcription factors were set as ending points

(blue). All intermediate proteins on the PPI are visualized as grey nodes. Molecular interactions

only overrepresented in calcitriol stimulated cells are shown as orange arrows. Each symbol in the

bar graphs represents one individual sample and data summarize the results from two or three

independent experiments with n≥ 5. Error bars indicate the s.e.m. Statistical test: two-tailed

Student's t-test (***C, E***) and two-tailed Mann-Whitney U-test (D). *P < 0.05, **P < 0.01, ***P <

0.001.

Figure 4-figure supplement 1. KEGG pathways analysis post vitamin D stimulation. List of

KEGG pathways enriched (p-adj. < 0.05) in DE genes after calcitriol stimulation. The x-axis

indicates the number of genes annotated to each pathway. The bar colors indicate the FDR-adjusted

p-value based on the colormap on the right side of the figure. The R package clusterProfiler was

used for the enrichment analysis and visualization.

Figure 4-figure supplement 2. Active signaling pathways in Th2 cells. *A,* Schematic

representation of the network analysis employed. ***B***, Network analysis to visualize active signaling

paths in control stimulated Th2 cultures. Predefined set of cell membrane receptors were set as

starting points (orange nodes) and annotated transcription factors were set as ending points (blue).

All intermediate proteins on the PPI are visualized as grey nodes. Molecular interactions only

overrepresented in control stimulated cells are shown as purple arrows.

Figure 5. Prenatal vitamin D supplementation protects from asthma development. *A*, H&E

staining of representative lung sections of indicated experimental groups. Arrowheads point to peri

bronchial infiltrates. ***B-E,*** Absolute numbers of total BAL (***B***) leukocytes, (***C***) eosinophils, (***D***)

lymphocytes, and (***E***) lung IL-13+ Th2 cells. ***F,*** Flow cytometric analysis of Aiolos expression in

differentiating Th2 cells at indicated time points. ***G***, QRT-PCR analysis of Ikzf3 transcripts in

respective groups. Expression levels were normalized to the house keeping gene L32 and are

expressed as fold change expression over the control samples WT-vehicle/ WT. **H**, Flow

cytometric analysis of Aiolos expression in control and calcitriol stimulated Th2 cells. **I,** Flow

cytometric analysis and frequencies of IL-2+ cells in WT and Ikzf3-/- cultures post indicated

treatment. Each symbol represents and independent sample (n= 9 per group). Flow cytometric

analysis and quantification of VDR expression in lung CD4+ T cells of the respective groups. Each

symbol represents one individual mouse and data summarize the results from two to three

independent experiments with n>8 per group. Error bars indicate the s.e.m. Statistical significance

was determined with Mixed-effect analysis with Holm-Šidák's post-hoc analysis (**B-E**), two-tailed

Mann-Whitney U-test (**G, H**) and One-way ANOVA with Holm-Šidák's post-hoc analysis (**G**). *P

< 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Figure 5-figure supplement 1. Prenatal vitamin D supplementation increases VDR

expression in CD4+ T cells. *A*, Schematic presentation of the dietary intervention protocol and

breeding scheme with vitamin D. ***B, Absolute neutrophil numbers in BAL. C***, Flow cytometric

analysis and quantification of VDR expression in lung CD4+ T cells of the respective groups. Each

symbol represents one individual mouse and data summarize the results from two to three

independent experiments with n>8 per group. Error bars indicate the s.e.m. Statistical significance

was determined with One-way ANOVA with Holm-Šidák's post-hoc analysis. *P < 0.05, **P <

0.01, ***P < 0.001, ****P < 0.0001.

Figure 5-figure supplement 2. Prenatal vitamin D supplementation mitigates Th2

development and cytokine production in vitro. *A-E*, Representative flow plots and cell

frequencies of (***A***) Gata3, (***B***) VDR, (***C***), IL-2, (***D***) IL-13, (***E***) IL-5 expression in indicated groups.

Each symbol represents one individual sample and data summarize the results from two to three

independent experiments (n= 4-10 per group). Error bars indicate the s.e.m. Statistical test: two-

tailed Mann-Whitney U-test (**a – e**). *P < 0.05, **P < 0.01, ***P < 0.001.

Figure 5-figure supplement 3. Schematic summary of vitamin D molecular genetics in the

17q12-21 region.

1.   Litonjua AA, Weiss ST. Is vitamin D deficiency to blame for the asthma epidemic? J

Allergy Clin Immunol [Internet]. 2007 Nov [cited 2023 Sep 25];120(5):1031–5. Available

from: https://pubmed.ncbi.nlm.nih.gov/17919705/

2.   Bouzigon E, Corda E, Aschard H, Dizier M-H, Boland A, Bousquet J, et al. Effect of

17q21 variants and smoking exposure in early-onset asthma. N Engl J Med [Internet].

2008 Nov 6 [cited 2023 Sep 25];359(19):1985–94. Available from:

https://pubmed.ncbi.nlm.nih.gov/18923164/

3.   Kelly RS, Chawes BL, Guo F, Zhang L, Blighe K, Litonjua AA, et al. The role of the

17q21 genotype in the prevention of early childhood asthma and recurrent wheeze by

vitamin D. Eur Respir J [Internet]. 2019 Oct 1 [cited 2022 Dec 8];54(4). Available from:

https://pubmed.ncbi.nlm.nih.gov/31439681/

4.   Schmiedel BJ, Seumois G, Samaniego-Castruita D, Cayford J, Schulten V, Chavez L, et

al. 17q21 asthma-risk variants switch CTCF binding and regulate IL-2 production by T

cells. Nat Commun [Internet]. 2016 Nov 16 [cited 2022 Dec 8];7. Available from:

https://pubmed.ncbi.nlm.nih.gov/27848966/

5.   Raby BA, Lazarus R, Silverman EK, Lake S, Lange C, Wjst M, et al. Association of

vitamin D receptor gene polymorphisms with childhood and adult asthma. Am J Respir

Crit Care Med [Internet]. 2004 Nov 15 [cited 2023 Sep 25];170(10):1057–65. Available

from: https://pubmed.ncbi.nlm.nih.gov/15282200/

6.   Camargo CA, Rifas-Shiman SL, Litonjua AA, Rich-Edwards JW, Weiss ST, Gold DR, et

al. Maternal intake of vitamin D during pregnancy and risk of recurrent wheeze in

children at 3 y of age. Am J Clin Nutr [Internet]. 2007 Mar 1 [cited 2023 Sep

25];85(3):788–95. Available from: https://pubmed.ncbi.nlm.nih.gov/17344501/

7.   Devereux G, Litonjua AA, Turner SW, Craig LCA, McNeill G, Martindale S, et al.

Maternal vitamin D intake during pregnancy and early childhood wheezing. Am J Clin

Nutr [Internet]. 2007 Mar 1 [cited 2023 Sep 25];85(3):853–9. Available from:

https://pubmed.ncbi.nlm.nih.gov/17344509/

8.   van der Pligt P, Willcox J, Szymlek-Gay EA, Murray E, Worsley A, Daly RM.

Associations of Maternal Vitamin D Deficiency with Pregnancy and Neonatal

Complications in Developing Countries: A Systematic Review. Nutrients [Internet]. 2018

May 18 [cited 2023 Sep 25];10(5). Available from:

https://pubmed.ncbi.nlm.nih.gov/29783717/

9.   Roth DE, Abrams SA, Aloia J, Bergeron G, Bourassa MW, Brown KH, et al. Global

prevalence and disease burden of vitamin D deficiency: a roadmap for action in low- and

middle-income countries. Ann N Y Acad Sci [Internet]. 2018 Oct 1 [cited 2023 Sep

25];1430(1):44–79. Available from: https://pubmed.ncbi.nlm.nih.gov/30225965/

10.   Cashman KD. Vitamin D Deficiency: Defining, Prevalence, Causes, and Strategies of

Addressing. Calcif Tissue Int [Internet]. 2020 Jan 1 [cited 2023 Sep 25];106(1):14–29.

Available from: https://pubmed.ncbi.nlm.nih.gov/31069443/

11.   Litonjua AA, Lange NE, Carey VJ, Brown S, Laranjo N, Harshfield BJ, et al. The Vitamin

D Antenatal Asthma Reduction Trial (VDAART): rationale, design, and methods of a

randomized, controlled trial of vitamin D supplementation in pregnancy for the primary

prevention of asthma and allergies in children. Contemp Clin Trials [Internet]. 2014 [cited

2023 Sep 25];38(1):37–50. Available from: https://pubmed.ncbi.nlm.nih.gov/24614387/

12.   Litonjua AA, Carey VJ, Laranjo N, Harshfield BJ, McElrath TF, O'Connor GT, et al.

Effect of Prenatal Supplementation With Vitamin D on Asthma or Recurrent Wheezing in

Offspring by Age 3 Years: The VDAART Randomized Clinical Trial. JAMA [Internet].

2016 Jan 26 [cited 2023 Sep 25];315(4):362–70. Available from:

https://pubmed.ncbi.nlm.nih.gov/26813209/

13.   Litonjua AA, Carey VJ, Laranjo N, Stubbs BJ, Mirzakhani H, O'Connor GT, et al. Six-

Year Follow-up of a Trial of Antenatal Vitamin D for Asthma Reduction. N Engl J Med

[Internet]. 2020 Feb 6 [cited 2023 Jan 19];382(6):525–33. Available from:

https://www.nejm.org/doi/full/10.1056/NEJMoa1906137

14.   Wolsk HM, Chawes BL, Litonjua AA, Hollis BW, Waage J, Stokholm J, et al. Prenatal

vitamin D supplementation reduces risk of asthma/recurrent wheeze in early childhood: A

combined analysis of two randomized controlled trials. PLoS One [Internet]. 2017 Oct 1

[cited 2023 Sep 25];12(10). Available from: https://pubmed.ncbi.nlm.nih.gov/29077711/

15.   Wolsk HM, Harshfield BJ, Laranjo N, Carey VJ, O'Connor G, Sandel M, et al. Vitamin D

supplementation in pregnancy, prenatal 25(OH)D levels, race, and subsequent asthma or

recurrent wheeze in offspring: Secondary analyses from the Vitamin D Antenatal Asthma

Reduction Trial. J Allergy Clin Immunol [Internet]. 2017 Nov 1 [cited 2023 Sep

25];140(5):1423-1429.e5. Available from: https://pubmed.ncbi.nlm.nih.gov/28285844/

16. Verlaan DJ, Berlivet S, Hunninghake GM, Madore AM, Larivière M, Moussette S, et al.

Allele-specific chromatin remodeling in the ZPBP2/GSDMB/ORMDL3 locus associated

with the risk of asthma and autoimmune disease. Am J Hum Genet [Internet]. 2009 Sep 11

[cited 2022 Dec 13];85(3):377–93. Available from:

https://pubmed.ncbi.nlm.nih.gov/19732864/

17. Miller M, Rosenthal P, Beppu A, Mueller JL, Hoffman HM, Tam AB, et al. ORMDL3

transgenic mice have increased airway remodeling and airway responsiveness

characteristic of asthma. J Immunol [Internet]. 2014 Apr 15 [cited 2022 Dec

8];192(8):3475–87. Available from: https://pubmed.ncbi.nlm.nih.gov/24623133/

18. Zhang Y, Willis-Owen SAG, Spiegel S, Lloyd CM, Moffatt MF, Cookson WOCM. The

ORMDL3 Asthma Gene Regulates ICAM1 and Has Multiple Effects on Cellular

Inflammation. Am J Respir Crit Care Med [Internet]. 2019 Feb 15 [cited 2023 Sep

25];199(4):478–88. Available from: https://pubmed.ncbi.nlm.nih.gov/30339462/

19. Fishilevich S, Nudel R, Rappaport N, Hadar R, Plaschkes I, Stein TI, et al. GeneHancer:

genome-wide integration of enhancers and target genes in GeneCards. Database (Oxford)

[Internet]. 2017 [cited 2023 Sep 25];2017. Available from:

https://pubmed.ncbi.nlm.nih.gov/28605766/

20. Neme A, Seuter S, Carlberg C. Vitamin D-dependent chromatin association of CTCF in

human monocytes. Biochim Biophys Acta [Internet]. 2016 Nov 1 [cited 2023 Sep

25];1859(11):1380–8. Available from: https://pubmed.ncbi.nlm.nih.gov/27569350/

21. Neme A, Seuter S, Carlberg C. Selective regulation of biological processes by vitamin D

based on the spatio-temporal cistrome of its receptor. Biochim Biophys acta Gene Regul

Mech [Internet]. 2017 Sep 1 [cited 2023 Sep 25];1860(9):952–61. Available from:

https://pubmed.ncbi.nlm.nih.gov/28712921/

22. Ramagopalan S V., Heger A, Berlanga AJ, Maugeri NJ, Lincoln MR, Burrell A, et al. A

ChIP-seq defined genome-wide map of vitamin D receptor binding: associations with

disease and evolution. Genome Res [Internet]. 2010 Oct [cited 2023 Sep 25];20(10):1352–

60. Available from: https://pubmed.ncbi.nlm.nih.gov/20736230/

23. Lu X, Chu CS, Fang T, Rayon-Estrada V, Fang F, Patke A, et al. MTA2/NuRD Regulates

B Cell Development and Cooperates with OCA-B in Controlling the Pre-B to Immature B

Cell Transition. Cell Rep [Internet]. 2019 Jul 9 [cited 2023 Sep 25];28(2):472-485.e5.

Available from: https://pubmed.ncbi.nlm.nih.gov/31291582/

24. Buniello A, Macarthur JAL, Cerezo M, Harris LW, Hayhurst J, Malangone C, et al. The

NHGRI-EBI GWAS Catalog of published genome-wide association studies, targeted

arrays and summary statistics 2019. Nucleic Acids Res [Internet]. 2019 Jan 8 [cited 2023

Sep 25];47(D1):D1005–12. Available from: https://pubmed.ncbi.nlm.nih.gov/30445434/

25. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast

universal RNA-seq aligner. Bioinformatics [Internet]. 2013 Jan [cited 2023 Sep

25];29(1):15–21. Available from: https://pubmed.ncbi.nlm.nih.gov/23104886/

26. Anders S, Pyl PT, Huber W. HTSeq—a Python framework to work with high-throughput

sequencing data. Bioinformatics [Internet]. 2015 Jan 15 [cited 2023 Sep 25];31(2):166–9.

Available from: https://dx.doi.org/10.1093/bioinformatics/btu638

27.   McCarthy DJ, Chen Y, Smyth GK. Differential expression analysis of multifactor RNA-

Seq experiments with respect to biological variation. Nucleic Acids Res [Internet]. 2012

May [cited 2023 Sep 25];40(10):4288. Available from: /pmc/articles/PMC3378882/

28.   Sakai Y, Kishimoto J, Demay MB. Metabolic and cellular analysis of alopecia in vitamin

D receptor knockout mice. J Clin Invest [Internet]. 2001 [cited 2023 Sep 25];107(8):961–

6. Available from: https://pubmed.ncbi.nlm.nih.gov/11306599/

29.   Li YC, Amling M, Pirro AE, Priemel M, Meuse J, Baron R, et al. Normalization of

mineral ion homeostasis by dietary means prevents hyperparathyroidism, rickets, and

osteomalacia, but not alopecia in vitamin D receptor-ablated mice. Endocrinology

[Internet]. 1998 [cited 2023 Sep 25];139(10):4391–6. Available from:

https://pubmed.ncbi.nlm.nih.gov/9751523/

30.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome Biol [Internet]. 2014 Dec 5 [cited 2023 Sep

25];15(12):1–21. Available from:

https://genomebiology.biomedcentral.com/articles/10.1186/s13059-014-0550-8

31.   Yu G, Wang LG, Han Y, He QY. clusterProfiler: an R package for comparing biological

themes among gene clusters. OMICS [Internet]. 2012 May 1 [cited 2023 Sep

25];16(5):284–7. Available from: https://pubmed.ncbi.nlm.nih.gov/22455463/

32.   Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers differential

expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Res

[Internet]. 2015 Apr 20 [cited 2018 Feb 18];43(7):e47–e47. Available from:

http://academic.oup.com/nar/article/43/7/e47/2414268/limma-powers-differential-

expression-analyses-for

33.   Benjamini Y, Drai D, Elmer G, Kafkafi N, Golani I. Controlling the false discovery rate in

behavior genetics research. Behav Brain Res [Internet]. 2001 Nov 8 [cited 2023 Sep

25];125(1–2):279–84. Available from: https://pubmed.ncbi.nlm.nih.gov/11682119/

34.   Silverbush D, Sharan R. A systematic approach to orient the human protein-protein

interaction network. Nat Commun [Internet]. 2019 Dec 1 [cited 2022 Mar 15];10(1).

Available from: https://pubmed-ncbi-nlm-nih-gov.ezp-prod1.hul.harvard.edu/31289271/

35.   Smedley D, Haider S, Durinck S, Pandini L, Provero P, Allen J, et al. The BioMart

community portal: an innovative alternative to large, centralized data repositories. Nucleic

Acids Res [Internet]. 2015 [cited 2023 Sep 25];43(W1):W589–98. Available from:

https://pubmed.ncbi.nlm.nih.gov/25897122/

36.   Moffatt MF, Kabesch M, Liang L, Dixon AL, Strachan D, Heath S, et al. Genetic variants

regulating ORMDL3 expression contribute to the risk of childhood asthma. Nature

[Internet]. 2007 Jul 26 [cited 2022 Dec 8];448(7152):470–3. Available from:

https://pubmed.ncbi.nlm.nih.gov/17611496/

37.   Litherland SA, Xie TX, Grebe KM, Davoodi-Semiromi A, Elf J, Belkin NS, et al. Signal

transduction activator of transcription 5 (STAT5) dysfunction in autoimmune monocytes

and macrophages. J Autoimmun [Internet]. 2005 Jun [cited 2023 Sep 25];24(4):297.

Available from: /pmc/articles/PMC2605968/

38.   Kara S, Pirela-Morillo GA, Gilliam CT, Wilson GD. Identification of novel susceptibility

genes associated with seven autoimmune disorders using whole genome molecular

interaction networks. J Autoimmun [Internet]. 2019 Feb 1 [cited 2023 Sep 25];97:48–58.

Available from: https://pubmed.ncbi.nlm.nih.gov/30391024/

39.   Syreeni A, Sandholm N, Sidore C, Cucca F, Haukka J, Harjutsalo V, et al. Genome-wide

search for genes affecting the age at diagnosis of type 1 diabetes. J Intern Med [Internet].

2021 May 1 [cited 2023 Sep 25];289(5):662–74. Available from:

https://pubmed.ncbi.nlm.nih.gov/33179336/

40.   Shrine N, Portelli MA, John C, Soler Artigas M, Bennett N, Hall R, et al. Moderate-to-

severe asthma in individuals of European ancestry: a genome-wide association study.

Lancet Respir Med [Internet]. 2019 Jan 1 [cited 2023 Sep 25];7(1):20. Available from:

/pmc/articles/PMC6314966/

41.   Marinho S, Custovic A, Marsden P, Smith JA, Simpson A. 17q12-21 variants are

associated with asthma and interact with active smoking in an adult population from the

United Kingdom. Ann Allergy Asthma Immunol [Internet]. 2012 [cited 2022 Dec

8];108(6). Available from: https://pubmed.ncbi.nlm.nih.gov/22626592/

42.   Hitomi Y, Kojima K, Kawashima M, Kawai Y, Nishida N, Aiba Y, et al. Identification of

the functional variant driving ORMDL3 and GSDMB expression in human chromosome

17q12-21 in primary biliary cholangitis. Sci Rep [Internet]. 2017 Dec 1 [cited 2023 Sep

25];7(1). Available from: /pmc/articles/PMC5460198/

43.   Keshari PK, Harbo HF, Myhr KM, Aarseth JH, Bos SD, Berge T. Allelic imbalance of

multiple sclerosis susceptibility genes IKZF3 and IQGAP1 in human peripheral blood.

BMC Genet [Internet]. 2016 Apr 14 [cited 2023 Sep 25];17(1). Available from:

https://pubmed.ncbi.nlm.nih.gov/27080863/

44.   Quintana FJ, Jin H, Burns EJ, Nadeau M, Yeste A, Kumar D, et al. Aiolos promotes TH17

differentiation by directly silencing Il2 expression. Nat Immunol [Internet]. 2012 Aug

[cited 2023 Sep 25];13(8):770–7. Available from:

https://pubmed.ncbi.nlm.nih.gov/22751139/

45.   Pfeffer PE, Hawrylowicz CM. Vitamin D in Asthma: Mechanisms of Action and

Considerations for Clinical Trials. Chest [Internet]. 2018 May 1 [cited 2023 Sep

25];153(5):1229–39. Available from: https://pubmed.ncbi.nlm.nih.gov/28923762/

46.   Baeke F, Korf H, Overbergh L, van Etten E, Verstuyf A, Gysemans C, et al. Human T

lymphocytes are direct targets of 1,25-dihydroxyvitamin D3 in the immune system. J

Steroid Biochem Mol Biol [Internet]. 2010 Jul [cited 2023 Sep 25];121(1–2):221–7.

Available from: https://pubmed.ncbi.nlm.nih.gov/20302932/

47.   Zhu J, Cote-Sierra J, Guo L, Paul WE. Stat5 activation plays a critical role in Th2

differentiation. Immunity [Internet]. 2003 [cited 2023 Sep 25];19(5):739–48. Available

from: https://pubmed.ncbi.nlm.nih.gov/14614860/

48.   Boonstra A, Barrat FJ, Crain C, Heath VL, Savelkoul HFJ, O'Garra A. 1alpha,25-

Dihydroxyvitamin d3 has a direct effect on naive CD4(+) T cells to enhance the

development of Th2 cells. J Immunol [Internet]. 2001 Nov 1 [cited 2023 Sep

25];167(9):4974–80. Available from: https://pubmed.ncbi.nlm.nih.gov/11673504/

49.   Cantorna MT, Snyder L, Lin YD, Yang L. Vitamin D and 1,25(OH)2D regulation of T

cells. Nutrients [Internet]. 2015 Apr 22 [cited 2023 Sep 25];7(4):3011–21. Available from:

https://pubmed.ncbi.nlm.nih.gov/25912039/

50.   Pichler J, Gerstmayr M, Szépfalusi Z, Urbanek R, Peterlik M, Willheim M. 1

alpha,25(OH)2D3 inhibits not only Th1 but also Th2 differentiation in human cord blood

T cells. Pediatr Res [Internet]. 2002 Jul [cited 2023 Sep 25];52(1):12–8. Available from:

https://pubmed.ncbi.nlm.nih.gov/12084841/

51.   Staeva-Vieira TP, Freedman LP. 1,25-dihydroxyvitamin D3 inhibits IFN-gamma and IL-4

levels during in vitro polarization of primary murine CD4+ T cells. J Immunol [Internet].

2002 Feb 1 [cited 2023 Sep 25];168(3):1181–9. Available from:

https://pubmed.ncbi.nlm.nih.gov/11801653/

1055

1056

1057

1058   **Appendix 1**

1059   **Primer sequences**

Gene	species	sequence
Gsdma antisense	mouse	5'-TCA ACC AGC TTC AGT TGC AC-3'
Gsdma sense	mouse	5'-TGC TGT CAA AGG ACG CTA TG-3'
Ikzf3 sense	mouse	5'-TGT CAC CTC TGC AAC TAC GC-3'
Ikzsf3 antisense	mouse	5'-TTC CGC AGA ACT CAC ACT TG-3'
IL-2 antisense	mouse	5'-ATC CTG GGG AGT TTC AGG TT-3'
IL-2 sense	mouse	5'-CCC ACT TCA AGC TCC ACT TC-3'
Ormdl3 antisense	mouse	5'-CTT TGC CTT GGT CTG GAG TC-3'
Ormdl3 sense	mouse	5'-CAC ACG GGT GAT GAA C AG TC-3'
Stat3 antisense	mouse	5'-TCG TGG TAA ACT GGA CAC CA-3'

Stat3 sense	mouse	5'-GAG GAG CTG CAG CAG AAA GT-3'
Stat5a antisense	mouse	5'-GTC AAA CTC GCC ATC TTG GT-3'
Stat5a sense	mouse	5'-TAC ATG GAC CAG GCT CCT TC-3'
Stat5b antisense	mouse	5'-TGG GTG GCC TTA ATG TTC TC-3'
Stat5b sense	mouse	5'-TGT GGA TAC AGG CTC AGC AG-3'

1060

1061

Appendix 2

Antibodies

Antibody	clone	Vendor	catalog #
Aiolos PE	8B2	Biolegend	653204
CD122 (IL2Rβ) APC	5H4	Biolegend	5H4
CD19 PE-Cy7	6D5	Biolegend	115520
CD4 FITC	RM4-5	Biolegend	100510
CD4 PerCPeFluor710	RM4-5	eBioscience	46-0042-82
CD44 FITC	IM7	Biolegend	103006
CD45 BV421	104	Biolegend	109832
CD62L PE	MEL-14	Biolegend	104408
CD8a APC-Cy7	53-6.7	Biolegend	100714
Foxp3 PE	FJK-16s	Invitrogen	12-5773-82
Gata3 PE	16E10A23	Biolegend	653804
IFNg FITC	XMG1.2	Biolegend	505806
IL-10 Pacific Blue	JES5-16E3	Biolegend	505020

IL-13 eFluor660	eBio13A	eBioscience	50-7133-82
IL-13 PE	eBio13A	molecular probes	A18492
IL17A PE-Cy7	TC11-18H10.1	Biolegend	506922
IL-2 PE-TexasRed	JES6-5H4	Biolegend	503840
monoclonal b-actin-Peroxidase	AC-15	Sigma-Aldrich	A3854
Phospho-BTK/ITK	M4G3LN	eBioscience	12-9015-42
Tbet PE	4B10	Biolegend	644809
VDR	D2K6W	Cell Signaling Technology	12550S
VDR AlexaFluor647	D2K6W	Cell Signaling Technology	custom order

1064

1065 **Appendix 3**

Reagent	Vendor	catalog #
anti-mouse CD28 (37.51)	BD Biosciences	553294
anti-mouse CD3e (145-2C11)	BD Biosciences	553057
b-mercaptoethanol	gibco	21985-032
calcitriol	Sigma-Aldrich	D1530
collagenase D	Roche	11088866001
Kwik-Diff soultion	Shandon	9990700
DMSO	Sigma-Aldrich	D2438
ProLong gold Anitfade Mountant with DAPI	ThermoFisher Scietific	P36935
rat anti-mouse anti-IFNg	BD Biosciences	554430
Anti-Mouse CD16/CD32 (Mouse BD Fc Block	BD Biosciences	553141
hamster anti-mouse anti-IL-4 (11B11)	BD Biosciences	554433

L-Glutamin	Sigma- Aldrich	G7513
LS columns	Miltenyi	130-042-401
microRNeasy	Qiagen	74004
naive CD4+ T cell isolation kit	Miltenyi	130-104-453
MEM non-essential amino acid solution (100x)	Sigma- Aldrich	M7145
Penicillin/Streptomycin	Sigma- Aldrich	P4333
Cell stimulation Cocktail (500x) phorbol myristate acetate (PMA)/Ionomycin	Tocris	TNB-4975
phosphate buffered saline	Gibco	14190-144
recombinant human TGFb1	Peprotech	100-21C
recombinant mouse IL-12	Peprotech	210-12
recombinant mouse IL-13	Peprotech	210-13
recombinant mouse IL-2	Peprotech	212-12
recombinant mouse IL-33	Peprotech	210-33
recombinant mouse IL-4	Peprotech	214-14
recombinant mouse IL-6	Peprotech	216-16
recombinant human IL-2	Peprotech	200-02
RPMI1640	Sigma-Aldrich	R8758
STAT3 inhibitor III WP1066	Sigma-Aldrich	573129
STAT5 inhibitor	EMD Millipore	573106
AS1517499 (STAT6 inhibitor)	Millipore-Sigma	SML190
U0126	CellSignaling Technologies	9903S

LY294002	CellSignaling Technologies	9901S
SsoFast Eva Green	Bio-Rad	1725201
10x zink fixative	BD Pharmingen	552658
Tris-Glycine-SDS Running Buffer	Boston Bioproducts	BP-150
Transfer buffer	Boston Bioproducts	BP-190
Tris Buffered Saline with Tween 20	CellSignaling Technologies	9997S

1066

Vitamin D constrains inflammation by modulating the expression of key genes on Chr17q12-21.1

Authors:
Ayşe Kılıç[1*], Arda Halu[1*], Margherita De Marzio[1,2], Enrico Maiorino[1], Melody G. Duvall[3], Thayse Brueggemann[3], Joselyn J. Rojas Quintero[3], Robert Chase[1], Hooman Mirzakhani[1], Ayse Özge Sungur[4, 5], Janine Koepke[5], Taiji Nakano[6], Hong Yong Peh[3], Nandini Krishnamoorthy[3], Raja-Elie Abdulnour[3], Katia Georgopoulos[7], Augusto A. Litonjua[8], Marie B. Demay[9], Harald Renz[10,11], Bruce D. Levy[3] and Scott T. Weiss[1*]

Affiliations:
Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School; Boston, MA, USA.
Department of Environmental Health, Harvard TH Chan School of Public Health; Boston, MA, USA.
Division of Pulmonary and Critical Care Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School; Boston, MA, USA.
Behavioral Neuroscience, Experimental and Biological Psychology, Philipps-University Marburg; Marburg, Germany.
Excellence Cluster Cardio-Pulmonary System (ECCPS), Justus Liebig University Giessen; Giessen, Germany.
Department of Pediatrics, Graduate School of Medicine, Chiba University; Chiba, Japan.
Cutaneous Biology Research Center, Massachusetts General Hospital and Harvard Medical School; Charlestown, Massachusetts, USA.
Division of Pediatric Pulmonary Medicine, Golisano Children's Hospital at Strong, University of Rochester Medical Center; Rochester, NY, USA.
Endocrine Unit, Massachusetts General Hospital and Harvard Medical School; Boston, MA, USA.
Institute of Laboratory Medicine and Pathobiochemistry, Molecular Diagnostics, Philipps University of Marburg and German Center for Lung Research (DZL); Marburg, Germany.
Department of Clinical Immunology and Allergology, Laboratory of Immunopathology Sechenov University; Moscow, Russia.

Figure 1

Figure 1-figure supplement 1

A HDM 25 µg i.n. HDM 25 µg i.n. sacrifice

6-8 wks 1 3 8 15 16 days

B saline HDM

CD4 PerCP-eF710

VDR- AF647

0.84 16.1

CD4+ VDR+ (%)

☐ saline
■ HDM

*

C in CD4+ and transcription factor + gate

Tbet-PE 0.2
Gata3-PE 26.8
Foxp3-PE 0.95

VDR- AF647

VDR⁺ (%)

☐ *Tbet+*
■ *Gata3+*
▨ *Foxp3+*

** ***

☐ WT/ WT- vehicle   ■ Vitamin D def   ▨ WT-calcitriol

D *Ikzf3* *Ormdl3* *Gsdma*   **f** *Ikzf3* *Ormdl3* *Gsdma*

rel. mRNA expression

* * *   ** ** **

E *Stat5a* *Stat5b* *Stat3*   **g** *Stat5a* *Stat5b* *Stat3*

rel. mRNA expression

ns ns ns   ** *** **

Figure 2

Figure 2 - figure supplement 1

A

naive CD4+

WT VDR⁻/⁻ 0h 48h Th1 Th2 iTreg

VDR

β-actin

B

days

Th0 1 2 3 5

VDR

β - actin

C

EtOH

DAPI CD4 VDR

calcitriol

DAPI CD4 VDR

Figure 2 - figure supplement 2

Figure 3

Figure 4

Figure 4 - figure supplement 1

A

Receptors

TFs

Mapping to PPI

Ranking

<Expression>	
Ctrl	*Calcitriol*
6063.77	8755.65
5184.54	6565.09
5053.26	5612
4837.35	4973.34
...	...
2005.85	2156.38

B

Figure 4 - figure supplement 2

Figure 5

Figure 5 - figure supplement 2

Cartoon summarizing Vitamin D molecular genetics at 17q12-21

1. VitD -> asthma through sphingolipids
2. VitD -> asthma through ORMDL3
3. VitD -> asthma through IKZF3
4. VitD -> asthma through CTCF

Loading order:
1- marker
2- empty
3- WT kidney sample: positive control
4- VDR-/- kidney sample: negative control
5- CD4+ CD62L+CD44- naïve T cells fresh
6- Th0 cells (24 h CD3 & CD28 stimulation)
7- in vitro Th1
8- in vitro Th2
9- in vitro iTreg
10- WT kidney sample: positive control
11- VDR-/- kidney sample: negative control

Source data-Figure 2 - figure supplement 2A

Loading order:
1- marker
2- empty
3- CD4+ CD62L+CD44- naïve T cells fresh
4- in vitro differentiating Th2 day 1
5- in vitro differentiating Th2 day 2
6- in vitro differentiating Th2 day 3
7- in vitro differentiating Th2 day 5
8- empty
9- WT lung sample: positive control
10- VDR-/- lung sample: negative control

Source data-Figure 2 - figure supplement 2B