Identification of novel basil downy mildew resistance genes using *de novo*

comparative transcriptomics

Kelly S. Allen[1*], Gregory A. DeIulio[1*], Robert Pyne[2], Jacob Maman[1], Li Guo[1], Robert L.

Wick[3], James Simon[2], Anne Gershenson[1], Li-Jun Ma[1]

*These authors contributed equally to this work.

1- University of Massachusetts Amherst Department of Biochemistry and Molecular

Biology, Amherst, MA, USA

2- Rutgers University Department of Plant Biology, New Brunswick, NJ, USA

3- University of Massachusetts Amherst Stockbridge School of Agriculture, Amherst, MA,

USA

Corresponding Author:

Li-Jun Ma

Telephone: (413) 545-1205

E-mail: lijun@biochem.umass.edu

Total word count (excluding summary, references and legends):	7610	No. of figures:	8
Summary:	196	No. of Tables:	2
Introduction:	950	No of Supporting Information files:	5 (2 figures, 3 tables)
Materials and Methods:	1674		
Results:	3588		
Discussion:	1145		
Acknowledgements:	124		

Summary

- Sweet basil (*Ocimum basilicum* L.) production is threatened by the oomycete pathogen *Peronospora belbahrii* causing basil downy mildew (BDM); BDM resistant cultivar 'Mrihani' (MRI) was identified in a germplasm screen, and fertile progeny were produced through a breeding program with BDM-susceptible 'Newton' (SB22), but the molecular mechanisms conferring resistance in MRI and progeny remained unknown

- Comparative transcriptomics was performed to identify candidate resistance genes and potential mechanisms for BDM resistance; RNA samples from BDM-infected MRI and SB22 plants were harvested at 4 time points during the first 3 days of infection to differentiate interactions in resistant and susceptible plants.

- Three categories of genes uniquely induced in resistant MRI upon pathogen challenge were identified: nucleotide-binding leucine rich repeat proteins (NLRs), multi-functional receptor-like kinases (RLKs), and secondary metabolic enzymes; validation of the top resistance candidate NLR gene confirmed its unique presence in MRI as well as in two of four resistant MRIxSB22 F_2 progeny.

- In MRI, pathogen challenge also upregulated transcripts in the salicylic acid synthesis pathway, suggesting its role in BDM resistance, and demonstrating the application of using comparative transcriptomics to identify resistance genes and mechanisms in non-model crops for marker-assisted breeding approaches.

Keywords: *Ocimum basilicum* **(sweet basil),** *Peronospora belbahrii*, **resistance, downy mildew, salicylic acid, transcriptomics**

INTRODUCTION

Basil (genus *Ocimum*) is a major herb crop with diverse species and cultivars possessing distinct phenotypes in plant size, leaf shape, aroma, and flavor (Vieira & Simon, 2006). Sweet basil (*Ocimum basilicum* L.) is the most popular basil and is cultivated for culinary use and essential oil production for applications including medicine, health care products, and food additives. In 2019, revenue generated in the US from sweet basil and other culinary herbs grown for dry processing and fresh market sales was estimated to be $165 million dollars, and other estimates have even valued the retail market above $300 million dollars (Wyenandt *et al.*, 2015; *(dataset) USDA National Agricultural Statistics Service (2017).*)

Basil Downy Mildew (BDM), caused by the biotrophic oomycete *Peronospora belbahrii*, has become the most important disease posing a serious threat to global basil production since its introduction and movement in Europe in 2001 and the US in 2007 (Belbahri *et al.*, 2005; Wyenandt *et al.*, 2015). The pathogen enters plant tissue through open stomata or direct penetration of the upper cuticle, and colonizes leaf tissue intercellularly, causing characteristic symptoms of interveinal chlorosis several days after infection (Wyenandt *et al.*, 2015; Cohen *et al.*, 2017). Following sustained periods of high relative humidity (>85%), the pathogen produces sporangiophores bearing infective sporangia, which emerge through stomata and create a gray to dark-gray discoloration corresponding to

interveinal chlorosis (Wyenandt *et al.*, 2015; Cohen *et al.*, 2017). The sporangia are

aerially dispersed and cause polycyclic infections throughout large areas of production

(Wyenandt *et al.*, 2015; Cohen *et al.*, 2017). After prolonged infection, the leaves

desiccate and are abscised from the plant (Wyenandt *et al.*, 2015; Cohen *et al.*, 2017).

Symptoms and signs of BDM disease render plants unfit for commercial sale (Wyenandt

et al., 2015).

First reported in Uganda in 1930, BDM began to attract attention in 2001 when disease

instances were increasingly reported across the world with reports from the Americas,

Asia, and Europe (Garibaldi *et al.*, 2004, 2005; McLeod *et al.*, 2006; Khateri *et al.*, 2007;

Roberts *et al.*, 2009; Ronco *et al.*, 2009; Martínez de la Parte *et al.*, 2010; Nagy & Horváth,

2011; Kanetis *et al.*, 2014; Šafránková & Holková, 2014; Choi *et al.*, 2016). BDM was first

reported in the US in 2007 in Florida, and then in the Northeast the following year (Roberts

et al., 2009; Wyenandt *et al.*, 2015). As of 2021, BDM has been reported in 44 US states,

including Hawaii and the District of Columbia (Wyenandt *et al.*, 2015; McGrath, 2021,

2022a), threatening productivity in every growing region.

Chemical and cultural control methods to prevent infection and reduce disease spread

include conventional fungicides (Homa *et al.*, 2014; McGrath, 2020b), nocturnal fanning

(Cohen & Ben-Naim, 2016), nocturnal illumination (Cohen *et al.*, 2013), and daytime solar

heating (Cohen & Rubin, 2015). However, these measures can be prohibitively time

consuming and costly, necessitating the development of improved basil cultivars with

BDM resistance. Initial screening of basil germplasm identified the *O. basilicum* 'Mrihani'

(MRI) (Horizon Seed Co., Williams, OR) with significant resistance to BDM (Pyne *et al.*,

2015). This cultivar has a unique anise/fennel aroma and flavor profile, and a distinctive

phenotype as compared to other *O. basilicum* cultivars selected for culinary use. For

example, the MRI leaves are smaller than other basil cultivars and have serrated leaves

compared to the large downward cupped smooth leaves of commercial sweet basil

(Figure 1A). Most importantly, the taste and smell of MRI due to its unique methyl chavicol

chemotype differs considerably from eugenol-enriched sweet basil (Pyne *et al.*, 2015).

This cultivar was used in a parental cross with BDM-susceptible and the Fusarium-

resistant 'Newton' (also referred to as Rutgers breeding line SB22), and successfully

produced fertile offspring. From the six-generation breeding design, four BDM resistant

cultivars ('Devotion', 'Obsession', 'Passion', and 'Thunderstruck') were selected from the

backcrossed population progeny with improved downy mildew resistance and desirable

phenotypes and chemotypes (Simon *et al.*, 2018).

Screening of the full-sibling family offspring of the MRI x SB22 cross revealed additive

and dominant gene effects of the MRI-conferred resistance, leading to the hypothesis that

dominant alleles are involved in resistance (Pyne *et al.*, 2015). A quantitative trait locus

(QTL) analysis of the F_2 mapping population from the cross between MRI and SB22

identified a single locus *(dm11.1)* that accounts for 20%-28% of the variance observed

among the F_2 population (Pyne *et al.*, 2017). In addition, this study also identified two

minor loci *(dm9.1 and dm14.1)* that respectively contributed 5-16% and 4-18% of the F_2

population's phenotypic variation. Resistance (R) genes involved in quantitative disease

resistance map to QTLs (Nelson *et al.*, 2018), and the results of the linkage mapping

performed in the 'MRI' X SB22 F2 mapping population suggest that these loci may contain

genes conferring quantitative disease resistance in 'MRI'. Without genome assemblies of

the basil cultivars, the underlying causative genes remained unknown.

A comparative transcriptomic analysis was designed to identify unique genes involved in

the interactions of the resistant cultivar MRI and the susceptible commercial cultivar SB22

with the pathogen *P. belbahrii.* RNA was extracted at 12, 24, 48, and 72 hours post

inoculation (hpi) representing roughly germination, penetration, and intercellular growth

stages. Global transcription expression profiles identified three categories of genes

uniquely induced in the MRI cultivar upon pathogen challenge, including R genes

encoding nucleotide-binding leucine rich repeat proteins (NLRs), receptor-like kinases

(RLKs) that sense conserved microbe-associated molecular patterns, and secondary

metabolic enzymes. Validation of the top candidate resistance NLR protein-encoding

gene confirmed its unique presence in the MRI cultivar as well as two out of the four

resistance hybrids. Unique upregulation of the salicylic acid synthesis pathway in MRI

suggests the perturbation of this important hormone signaling pathway in conferring BDM

resistance.

MATERIALS AND METHODS

Sample preparation

Inbred *O. basilicum* genotypes SB22 (*P. belbahrii* susceptible) and MRI (*P. belbahrii*

resistant) plants were grown from seed. Previously infected sweet basil leaves with fresh

P. belbahrii sporulation were harvested and agitated for 5 minutes in sterile distilled water

(diH$_2$0). The inoculum mixture was filtered with 40 μm nylon mesh. A 1 mL subsample

from the filtered inoculum was pipetted into an Eppendorf tube and frozen at -80°C to

serve as a pathogen control. The remaining inoculum was centrifuged at 1,000g for 1 min

and diH$_2$0 decanted. The resulting sporangia pellet was resuspended in diH$_2$O, and the

inoculum concentration was adjusted to 1 x 10^5 sporangia/mL. Four-to-six-week-old MRI

and SB22 plants were spray-inoculated at the 6-leaf (3 true leaf set) growth stage with

approximately 1 mL/leaf and plants were incubated at 100% relative humidity for 24

hours. A set of MRI and SB22 plants were sprayed with diH2O in triplicate to serve as the

mock inoculated control.

Four disks per true leaf were sampled from both genotypes at 12, 24, 48 and 72 hpi and

immediately flash frozen in liquid nitrogen. The water control leaves were harvested at 12

hpi only. Total RNA was extracted from freshly ground tissue using the Spectrum™ Plant

Total RNA Kit (Sigma Aldrich). RNA samples were used to generate sequence libraries

using a library prep kit from New England Biolabs (NEB #E7530). Paired-end sequence

reads of 75 bp were generated at the TUFTs genomic center at the Tufts University

School of Medicine using the Hi-Seq Illumina platform. Higher coverage analyses were

specifically designed for inoculated samples that contain both the pathogen and the host

due to the increased complexity of these samples, allowing for further study of pathogen

expression.

Generating transcript assemblies and FPKM expression

FASTQC version 0.11.5 (Andrews, 2010) was used to assess average read quality. Paired-end reads (fastq files) were provided to Trinity version 2.4.0 and assembled using default parameters (Grabherr *et al.*, 2011). Datasets were assembled including single sets using either all MRI datasets and the sporangia control (MRI Combined Assembly) or all SB22 datasets including the sporangia control (SB22 Combined Assembly). Separately assembled control data provided organism specific databases of genes and transcripts.

The resulting output files served as the references for expression quantification. RSEM version 1.2.29 (Li & Dewey, 2011) and bowtie version 1.0.0 (Langmead *et al.*, 2009) were used to calculate FPKM (Fragments per Kilobase exon per Million mapped reads) values for assembled contigs while tracking replicate information. RNAseq datasets from both MRI and SB22 were mapped to the infected MRI Combined Assembly to standardize the reference which allowed us to use previously generated gene annotations and to cluster genes from both cultivars together. In all cases the standard settings were used for assembly and transcript quantification. Additionally, edgeR (Robinson *et al.*, 2010) was used to calculate differential gene expression. Expression data from both MRI and SB22 data mapped to the MRI Combined Assembly using Trinity and edgeR was used to assess differential expression between all timepoints within a single cultivar. Trinity DEG output data was filtered for genes with a p-value less than 0.05 and FDR less than 0.01.

Reciprocal BLAST hits and reference gene phylogeny

To explore the overall sequence conservation between MRI and SB22, we performed a reciprocal BLAST using all sequences within the MRI and SB22 water control assemblies. Briefly, all MRI sequences were compared to the SB22 transcriptome, and all SB22 sequences were compared to the MRI transcriptome. The hit with the highest BLAST score for each gene was chosen. Results were compared and pairs of top scoring genes were considered reciprocal best BLAST hits (i.e., MRI gene X BLASTs to SB22 gene Z, and SB22 gene Z BLASTs to MRI gene X).

Sequence conservation between MRI and SB22 was further assessed by performing a phylogenetic analysis using 9 protein-coding chloroplast genome genes based on a prior analysis (Rastogi *et al.*, 2015). The MRI and SB22 chloroplast genes were identified using BLAST against the MRI and SB22 water control assemblies, and the top hits with the highest bit scores were chosen and translated into coding sequences. Sequence alignment of the MRI and SB22 coding sequences was performed against sequences from 14 asterid lineage plants downloaded from NCBI Organelle Genome Resources database, with *Spinacia oleracea* L. and *Arabidopsis thaliana* L. set as outgroups (Rastogi *et al.*, 2015). The gene sequences were aligned using MAFFT (Madeira *et al.*, 2019), and the tree was generated using IQ-TREE (Minh *et al.*, 2020).

Sequence Translation and Annotation

We generated a database of sequence annotations for MRI genes. All MRI genes with an expression of FPKM >1 in at least one time point were chosen and the longest transcript

associated with that gene was compared to the NCBI non-redundant database using

cloud BLAST through Blast2GO. Annotations were saved as a searchable database in

text format. Genes were filtered by taxonomic hit to verify their species of origin as

needed.

To facilitate easier searches for gene families of interest, we translated the longest

nucleotide sequence associated with each gene into six-frame translated protein

sequences using EMBOSS, searching for only those translated sequences between

START and STOP codons longer than 30 amino acids. In many cases to verify protein

domain structure, the nucleotide or protein sequence was analyzed using either the NCBI

conserved domain finder (Marchler-Bauer *et al.*, 2015), PFAM (Finn *et al.*, 2016), or

InterProScan (Jones *et al.*, 2014). Sequences were aligned using MEGA 6 for visual

inspection (Tamura *et al.*, 2013).

Expression clustering toward candidate resistance gene identification

Genes were clustered using Trinity version 2.2.0 based on read counts following the steps

outlined in the Trinity manual (Robinson *et al.*, 2010). Expression data generated by

mapping all datasets to the MRI Combined Assembly were used for clustering. A matrix

of gene expression at all timepoints and replicates was used to define clusters with the

edgeR function associated with Trinity, using p=50 and p=20 (a grouping parameter for

cluster creation, with higher numbers forming larger and broader clusters). The resulting

clusters, available in pdf format, were visually examined for clusters which displayed the

target expression profile.

BLAST search for secondary metabolite enzyme and resistance genes

To analyze MRI unique gene families and defense hormone signaling genes, we performed BLASTp search against MRI and SB22 translated nucleotide sequences using *A. thaliana*, or in some cases sweet basil, protein sequences retrieved from NCBI. Generally, the hit with the highest bit score was chosen as the top hit for each sequence. In cases of short alignment length or low sequence identity, the recovered MRI or SB22 hit was compared to the green plant database on NCBI. BLAST version 2.2.22 was used in all cases to compare protein sequences (Altschul *et al.*, 1990) at the Massachusetts green-energy high performance computing center (MGHPCC).

PCR screen of parent and cultivar genomic DNA for unique genes

Genomic DNA was prepared from approximately 80mg of newly emerging leaf tissue of MRI, SB22, 'Devotion', 'Obsession', 'Passion' and 'Thunderstruck' cultivars using the E.Z.N.A. SP Plant DNA Kit (Omega BioTek, Norcross, GA) (Pyne *et al.*, 2017). Primers amplifying transcript sequences were designed for MRI and SB22 shared genes as well as MRI unique genes including the comp160460c0 transcript, and were ordered from IDT (Coralville, IA). The primers were either external primers designed to amplify the whole gene (MRI_134-F 5'-CCGAGAAAATCGATCTAGAGAG-3', MRI_2869-R 5'-CTAGCTTGATCTTTTAATTGGTGGAAAAAT-3') or internal primers for specific regions of interest (Supporting Information Table S1). Primers amplifying a 198bp fragment of the *O. basilicum* Actin gene (ObActin_2-F 5'-GTTATGCACTTCCCCATGCT-3', ObActin_2-R 5'-GAGCTGTTCTTTGCGGTCTC-3') were used in positive control reactions for all cultivars. PCR was performed with Q5 High-Fidelity DNA Polymerase (New England

Biolabs, Ipswich, MA) using manufacturer-recommended cycling conditions with a 30

second denaturation cycle to ensure full denaturation of genomic DNA, a 30 second

extension time to amplify the 198bp ObActin region and a 120 second extension time to

amplify the ~3.5kb comp160460-encoded gene on a Mastercycler proS (Eppendorf,

Hamburg, Germany). Water was used as a negative control template in all reaction sets.

Amplicons were visualized on 1.5% agarose gels stained with SYBR Safe DNA Gel Stain

(Invitrogen, Waltham, MA), and imaged under UV light.

NLR allele analysis

Successfully amplified comp160460c0 products from MRI, 'Devotion' and 'Obsession'

were purified using a QIAquick PCR purification kit (QIAGEN, Hilden, Germany) and

single amplicon copies were ligated into the pMiniT 2.0 vector using the NEB PCR Cloning

Kit (New England Biolabs, Ipswich, MA). Individual clones were selected and confirmed

via colony PCR using Q5 High-Fidelity DNA Polymerase (New England Biolabs, Ipswich,

MA) with primers 1BF and 8R which were designed to amplify the coiled coil and NB-ARC

domain coding sequences of MRI-R1. Plasmid DNA was prepared from confirmed clones

using the Zyppy Plasmid Miniprep Kit (Zymo Research, Irvine, CA).

Individual comp160460c0 clones from MRI, 'Devotion' and 'Obsession' were sequenced

using NEB PCR Cloning Kit Cloning Analysis Forward and Reverse primers to flank the

insert, as well as internal primers (Supporting Information Table S1). Twelve clones were

sequenced from MRI, and six clones were sequenced from both 'Devotion' and

'Obsession'. Consensus sequences for each clone were assembled and annotated to

identify intron regions by alignment to transcript sequences; coding regions were

confirmed and annotated using InterProScan (Jones *et al.*, 2014). Assembled sequences

were identified as individual alleles, nucleotide and predicted protein sequences were

aligned using the EMBL-EBI Clustal Omega multiple sequence alignment tool (Madeira

et al., 2019), and alignments were visualized using Jalview 2.11.1.4 (Waterhouse *et al.*,

2009).

Protein structures were predicted using RoseTTAFold (Baek *et al.*, 2021), and the

predicted secondary structure was added to the Clustal Omega allele sequence

alignment using ESPript 3.0 (Robert & Gouet, 2014). Allele structures were used as a

query search against the Protein Data Bank in DALI, and structure pairwise comparison

was performed (Holm, 2020). Allele structures were further analyzed and aligned using

UCSF Chimera (Pettersen *et al.*, 2004). Allele expression was analyzed by mapping the

variable coding regions to the RNA-seq data using Burrows-Wheeler Aligner software

package (BWA-MEM) (Li & Durbin, 2009), and results were visually examined using the

Integrated Genomics Viewer (Robinson *et al.*, 2011).

RESULTS

Sequencing data reflect phylogenetic relatedness of MRI and SB22

We generated 12.8 million (MRI), 14.3 million (SB22), and 9.9 million (Sporangia) high

quality paired-end reads per replicate for three controls (Table 1, Supporting Information

Figure S1). Considering the increased complexity of infected samples, we doubled the

sequence coverage and generated an average 24.8 million and 27.6 million high quality

Illumina paired-end reads per replicate per infection sample for MRI and SB22,

respectively (Table 1, Supporting Information Figure S1). All sequence data were

deposited at NCBI under GEO NUMBER: GSE111387.

Dataset (organism)	Read pairs (millions)	Mapping Percentage	Average base quality	Number of Genes	Average gene length	Average coverage
Sporangia (*P. belbahrii*)	29.7	84.1	37.28	10144	1603	230.4
MRI Water (*O. basilicum* MRI)	42.7	62.0	37.20	66930	893	66.4
MRI 12 hpi (*O. basilicum* MRI, *P. belbahrii*)	75.9	63.3	37.21	81873	897	98.1
MRI 24 hpi (*O. basilicum* MRI, *P. belbahrii*)	75.3	63.3	37.22	89302	864	92.6
MRI 48 hpi (*O. basilicum* MRI, *P. belbahrii*)	77.0	63.1	37.23	85832	860	98.7
MRI 72 hpi (*O. basilicum* MRI, *P. belbahrii*)	69.8	63.4	37.22	78061	909	93.5
SB22 Water (*O. basilicum* SB22)	38.3	62.9	37.18	52844	977	69.9
SB22 12 hpi (*O. basilicum* SB22, *P. belbahrii*)	85.9	63.9	37.18	74918	934	117.6
SB22 24 hpi (*O. basilicum* SB22, *P. belbahrii*)	83.2	63.3	37.19	72699	949	114.5
SB22 48 hpi (*O. basilicum* SB22, *P. belbahrii*)	75.9	65.2	37.15	84478	870	100.9
SB22 72 hpi (*O. basilicum* SB22, *P. belbahrii*)	86.6	66.6	37.20	80675	903	117.7
MRI Combined Assembly	370.4	64.8	37.22	133,441	765	352.6
SB22 Combined Assembly	399.6	66.0	37.19	118,296	692	477.7

Table 1. Summary of assembled transcripts by timepoint.

In total, 240.2 million paired-end reads were used to generate the MRI Combined

Assembly, containing 341,633 unique transcripts corresponding to 133,441 genes called

by Trinity. The SB22 Combined Assembly was generated using 263.9 million paired-end

reads and contained 118,296 genes and a total of 322,696 unique transcripts. The MRI

and SB22 plant-only control assemblies contained 66,930 and 52,844 genes respectively,

and the sporangia control contained 10,144 assembled genes. As expected, more genes

were assembled in infected samples, representing both host and pathogen transcripts,

genes expressed only during infection, and assembly errors (fragmented sequences)

introduced as transcriptome complexity increased.

Though fewer reads were sequenced for both water and sporangia control samples, the

sporangia control produced the highest sequence coverage (>90x) and longest average

gene length (1,603 bp) for assembled genes, likely due to the smaller

genome/transcriptome size of the (inactive) pathogen. The average assembled gene

length of *O. basilicum* transcripts from a previously published transcriptome was 1,363

bp (Rastogi *et al.*, 2014), larger than our average assembly size. The average gene size

of the oomycete *Phytophthora infestans* was 1,523 bp (Haas *et al.*, 2009), roughly

equivalent to the sporangia control assembly.

To assess the genetic diversity between MRI and SB22, we performed a BLAST search

between MRI and SB22 water control assemblies and identified 20,943 reciprocal hits,

likely representing orthologs between these two plants. These orthologs are highly similar

with an average pairwise sequence identity of 98.66% (Figure 1B). We further assessed

the genetic similarity between the cultivars by conducting a phylogenetic analysis of nine

protein-coding chloroplast genome orthologs across members of the asterid clade, to

which *O. basilicum* belongs, and with *A. thaliana* and *S. oleracea* set as outgroups

(Rastogi *et al.*, 2015). The analysis showed high sequence conservation among the

Ocimum spp. with MRI and SB22 grouped together (Figure 1C). We anticipate that unique

genes or differentially regulated genes are likely to contribute to the MRI and SB22

phenotypic variations.

Figure 1: Basil cultivar phenotypic and genetic diversity. (a) BDM susceptible 'Newton' (SB22) and resistant 'Mrihani' (MRI) cultivars. (b) Bi-directional blast hit among 20,943 genes shared between the SB22 and MRI water references. The nucleotide identity for each top BLAST hit is graphed here with a bin size of 0.25%. (c) Multigene phylogeny of chloroplast genome orthologs across members of the asterid clade.

Using expression profiles in the individual assemblies, MRI transcripts can be divided into

36,414 predicted plant transcripts (present in the MRI water control), 9,988 predicted

pathogen transcripts (present in the sporangia control), and 29,502 infection unique

transcripts (absent in both plant and pathogen controls) (Table 2). Similarly, the SB22

transcripts include 31,702 transcripts with a plant origin, 9,426 transcripts of pathogen

origin, and 26,486 transcripts uniquely present in the infection samples. Consistent with

SB22 susceptibility, we saw a significant increase in the number of expressed pathogen

genes in the susceptible host, observing an almost 20-fold increase from 12 to 72 hpi

compared to only a two-fold increase for MRI (Figure 2). Measuring total pathogen mRNA

abundance (the number of reads mapping to roughly 4,553 pathogen genes), we detected

a 43-fold increase in mapped pathogen reads from the SB22 72 hpi samples relative to

the MRI 72 hpi samples (Figure S2).

359 **Table 2: Expressed genes across timepoints**

Dataset	Total Genes	Number of Genes FPKM > 1		
		Plant Control	Sporangia Control	Infection Unique
MRI Water	36,528	36,414*	0	0
MRI 12 hours	39,349	29,247	472	9,534
MRI 24 hours	40,171	29,074	389	10,606
MRI 48 hours	40,973	28,852	179	11,840
MRI 72 hours	39,279	28,365	825	9,987
Sporangia	10,102	0	9,988*	0
Total unique	**76,018**	**36,414**	**9,988**[+]	**29,502**
SB22 Water	31,794	31,702*	0	0
SB22 12 hours	35,561	25,770	274	9,431
SB22 24 hours	36,732	25,584	699	10,364
SB22 48 hours	40,216	25,746	1,788	12,598
SB22 72 hours	40,838	25,414	5,375	9,962
Sporangia	9,518	0	9,426*	0
Total unique	**67,706**	**31,702**	**9,426**[+]	**26,486**

*- When the data was split into 3 sets 114 MRI points and 92 SB22 points were removed due to ambuiguity
+- The numbers displayed in in the chart above are total genes in each category as split by FPKM based separation after mapping to each master assembly. If the gene had a non-zero FPKM in the sporangia samples it was included in the "sporangia" sample. The difference in gene count is a reflection of mapping to two seperate assemblies.

Figure 2. Plant and pathogen gene expression profiles from clustering of the MRI
Combined Assembly. Clustering was done using all datasets mapped to the MRI
combined assembly. Cluster numbers are displayed above each cluster image. Dataset
and replicate are listed on the bottom x-axis. Sporangia expressed gene expression is
in red, infection unique gene expression is in gray and gene expression in the mock
infected water controls are in black for both MRI and SB22.
376
Clustering analysis highlights transcripts with potential functions involved in

host - pathogen interactions

Based on overall ortholog identity, we were confident that SB22 reads from orthologous

genes would align to the MRI assembly. Using the MRI Combined Assembly as a

reference, we mapped all the SB22 and MRI reads using RSEM with default parameters.

Initial coarse clustering using all genes from MRI and SB22 mapped data resulted in 12

clusters with on average 2,810 genes per cluster (Supporting Information Table S2).

Three clusters that lacked consistency among biological replicates were not included in

further analyses. Three clusters (A1, A2 and A3) characterized transcripts primarily

belonging to the pathogen (Figure 3 panel A), as all transcripts showed significant

expression in sporangia pathogen control samples (Figure 3 grey bar in the middle), but

no expression in both water-inoculated plant control samples (Figure 3 two blue bars

representing MRI-water only and SB22-water only).

Six clusters containing primarily plant transcripts were identified. All transcripts showing

significant expression in plant and infected plant samples that were absent from

sporangia pathogen control samples were filtered (Panels B and C). The remaining genes

from the six plant gene clusters represent plant transcripts during pathogen challenge.

Transcripts within three clusters, B-1, B-2 and B-3, had comparable expression profiles

between MRI and SB22, indicating conserved functions between the two different plant

hosts. Both B-1 and B-2 clusters show a pattern consistent with a 12-hour shift in

photoperiod, but these clusters respond in opposite directions. Cluster B-1 was

upregulated at 24, 48, and 72 hpi and was enriched for metabolism, oxidation-reduction,

and photosynthesis functions. No GO terms were significantly enriched in cluster B-2

which showed downregulation at 24, 48, and 72 hpi; however, of those GO terms

annotated by Blast2Go metal ion functions were predominant. The largest cluster, B-3,

roughly represents stably expressed plant genes. Cluster B-3 is enriched for many

categories including various metabolic processes, protein modification, and protein

localization, among others.

Transcripts in three clusters, C-1, C-2, and C-3 displayed differential regulation responses

between MRI and SB22 during pathogen challenge (Figure 3B). In Cluster C-1, MRI

expression is high but almost completely absent from the SB22 transcriptome. In Cluster

C-2, MRI expression was higher than SB22 and was enriched for genes related to

defense response, response to stress, response to stimulus, and DNA integration. In

Cluster C-3, the expression of MRI genes is instead lower than SB22. No GO terms were

enriched in cluster C-3.

Figure 3. Clusters with expressed plant genes. Nine gene clusters produced by coarse clustering are displayed here. Expression value, y-axis, is on a log2 scale. Three replicates for each dataset are represented by tick marks on the bottom x-axis, with mean data plotted as dark blue points and gray shading representing the standard deviation. Datasets from MRI are highlighted in orange with 3 replicates from 12 to 72 hpi, SB22 datasets are colored pink with 3 replicates from 12 to 72 hpi, both water controls are labeled blue (MRI left, SB22 right), and the pathogen sporangia control data is colored gray. Column A, pathogen gene clusters as all transcripts showed significant expression in sporangia pathogen control samples but were absent from plant and infected plant samples. Columns B and C gene clusters contain primarily plant transcripts as all transcripts showed significant expression in plant and infected plant samples but were absent from sporangia pathogen control samples. Column B, plant genes expressed similarly in both cultivars. Column C, plant genes with different profiles between cultivars.

MRI unique expressed genes include NLR, RLK and secondary metabolic enzymes

To understand potential mechanisms underlying the resistance, we repeated the

clustering with higher stringency (p=20, see methods for details) resulting in 188 clusters.

Eight clusters were chosen as they were expressed in MRI, minimally expressed in SB22,

and showed no expression in the sporangia control. A comprehensive filtering process of

the eight clusters resulted in a total 369 MRI unique candidate genes. These MRI unique

candidate genes can be grouped into secondary metabolic enzymes (22 genes),

immunity related genes including 22 nucleotide-binding site leucine-rich repeat (NLR)

genes and 25 receptor-like kinases (RLK) or receptor-like proteins and others.

Detecting secondary metabolic enzymes as MRI unique genes is expected as these two

basil plants produce distinct secondary compounds. For instance, SB22 accumulates a

significant amount of eugenol, while MRI predominantly accumulates methylchavicol

(Rob Pyne, unpublished data). This distinct chemotype prevents MRI from immediate

commercial use. Examining twenty-five secondary metabolite related genes predicted as

MRI unique genes, we found enzymes related to secondary metabolites which specifically

differentiate the MRI and SB22 chemotypes, including cinnamate p-coumarate carboxyl

methyltransferase, enzymes involved in anthocyanin biosynthesis, and chavicol/eugenol

O-methyltransferase, the enzyme that catalyzes the conversion of chavicol to

methylchavicol, as would be predicted from the chemotypes.

Unique expression of RLKs and NLRs, both immunity related protein families, in MRI is

of particular interest. Basic plant immunity consists of pattern-triggered immunity (PTI)

and effector-triggered immunity (ETI) (Jones & Dangl, 2006; Cui *et al.*, 2015). Plant PTI

uses receptor-like proteins/kinases (RLPs/RLKs), a large gene family (Chern *et al.*, 2016;

Mendy *et al.*, 2017) that have roles as sensors of microbe-associated molecular patterns

(MAMPs) and induce downstream defense reactions (Jones & Dangl, 2006; Dodds &

Rathjen, 2010; Bigeard *et al.*, 2015; Zhou & Zhang, 2020). Plant ETI employs an

intracellular nucleotide-binding site and leucine-rich repeat domain receptors (NLRs) that

play roles in sensing effector proteins secreted by pathogens and regulating downstream

defense signaling (Jones & Dangl, 2006; Cui *et al.*, 2015; Cesari, 2018; Monteiro &

Nishimura, 2018; Wang & Chai, 2020).

To investigate basil PTI, we further characterized the RLK BLAST hits. The best RLK

candidate is transcript mri_comp170662, which encodes a full-length malectin-like RLK.

Some predicted receptor-like kinases (RLK) as truncated fragments, such as three

transcripts containing only the leucine-rich repeat (LRR) domain, two containing only

protein kinase domains, and two that were too short to have domain affiliation. Two

transcripts contained an RLK-like domain involved in antifungal and salt tolerance

(Sawano *et al.*, 2007; Zhang *et al.*, 2009). Using the full-length top candidate RLK

transcript mri_comp170662, we identified 12 and 10 homologs in the MRI and SB22

assemblies, respectively. Ten SB22/MRI orthologous pairs can be readily identified

between these two sister cultivars based on a phylogeny (three pair members lacking a

malectin domain were excluded from the tree) (Figure 4a). Three MRI orphan transcripts

including this top RLK candidate mri_comp170662 showed increased expression at 12

and 24 hpi compared to the water control (Figure 4b). This RLK candidate contains LRR,

Malectin and kinase domains (Figure 4c).

Figure 4. Malectin-like RLK proteins in MRI and SB22. (a) Alignment of 19 of 22 RLK proteins by their conserved malectin domain. Gray bars with numbers represent orthologous pairs and boxes indicate sequences absent from the SB22 cultivar. (b) Fold change compared to water for MRI and SB22 RLKs across infected plant samples. (c) Protein domain structure of 22 RLK hits generated from translated nucleotide sequences. Transcript IDs are those of the adjacent fold change row.

NLRs upregulation in MRI during infection

To identify unique MRI NLR resistance genes involved in ETI, we focused on Cluster C-1 where all 17 transcripts are highly expressed in MRI upon infection but almost absent from the SB22 transcriptome. Two transcripts, comp_178221_c0 and comp_160460_c0, are putative NLR resistance genes encoding a late blight resistance protein homolog R1A (gi:848916018 and gi:848932751) from spotted monkey flower (*Erythranthe guttata*), which belongs to the order Lamiales including basil.

Members of the plant NLR protein family have been characterized as sensors,
recognizing specific microbial effectors in response to ongoing host-pathogen
coevolution, or as helpers involved in signal transduction (Wu *et al.*, 2017). These proteins
are highly conserved in eukaryotic immune responses (Wu *et al.*, 2017). NLRs contain a
central nucleotide-binding domain and a C-terminal leucine-rich repeat region that confers
specificity to the receptor (Wu *et al.*, 2017; Prigozhin & Krasileva, 2020). There are three
subfamilies of NLRs defined by presence of one of three functional N-terminal domains:
Resistance To Powdery Mildew 8 (RPW8), Coiled-Coil (CC), or Toll/Interleukin-1
Receptor homology (TIR) (Prigozhin & Krasileva).

The coiled-coil domain NLR subfamily (CC-NLRs) comprises several well-characterized
intracellular receptors, including Recognition of Peronospora Parasitica 1 (RPP1) genes
conferring resistance to downy mildew in *A. thaliana* (Krasileva *et al.*, 2010). CC-NLRs
are characterized by an N-terminal CC domain, which has been associated with
oligomerization in characterized CC-NLRs including barley mildew locus A MLA10 and *A.*
thaliana ZAR1, an NLR that polymerizes to form a plant "resistosome" during the immune
response (Maekawa *et al.*, 2011; Adachi *et al.*, 2019).

We focused on validating our computational prediction and further examined our top
candidate NLR transcript comp160460_c0 identified in Cluster C-1. Transcript
comp_160460_c0, MRI Resistance gene 1 (MRI-R1), encodes a full-length CC-NLR
protein of 887 aa with all three functional domains. This NLR candidate is uniquely
expressed and differentially upregulated in MRI in the presence of the pathogen (Figure

5A). Based on mapping results and expression analyses, while MRI-R1 is expressed in

the control samples, its expression is significantly increased in the pathogen-inoculated

samples throughout the time-course of infection. Specifically, MRI-R1 was 2-fold

upregulated between 12 and 24 hpi and upregulated expression was maintained 48- and

72-hpi. Mapping the SB22 infected assemblies against the MRI-R1 transcript sequence

as a reference confirmed that there was no detectable expression of MRI-R1 detected in

susceptible samples.

To determine whether MRI-R1 activity could be attributed to presence/absence

polymorphism between MRI and SB22 or the differential regulation at transcriptional level,

we designed primers to amplify the coding region of the full-length MRI-R1 transcript

using PCR. Forward primer 1BF flanks the 5' end of the coding sequence, and reverse

primer 14R targets the 3' end of the gene. This primer pair produced an amplicon of the

MRI-R1 gene from MR1, but not from SB22 or the water template negative control (Figure

4B). Internal primers were designed to further examine MRI-R1 like genes in MRI and

SB22 (Supporting Information Table S1). PCR using internal forward primer 9F, which

flanks the NB-ARC domain, paired with 3' end primer 14R, resulted in gene amplification

from MRI and potentially off-target or ortholog amplification from SB22 (Figure 5B). These

results suggest that the full MRI-R1 gene is unique to MRI, but that there is some partial

sequence conservation in a similar gene of unknown functional status in SB22. Thus,

MRI-R1 represents another case of R gene polymorphism among closely related

organisms.

The MRI-R1 amplicon detected in the MRI gDNA appeared to be approximately 800bp

larger than predicted from the RNA transcripts, indicating the potential presence of a non-

coding intronic region (Figure 5B). We also observed sequence polymorphisms among

MRI-R1 transcripts with 6 different isoforms predicted for the same gene (2 with full coding

sequences, comp160460_c0_seq2 and comp160460_c0_seq5). To investigate transcript

polymorphisms we isolated and sequenced individual clones from the PCR products. A

total of 12 full-length amplicons were cloned from three MRI plants. Due to the length of

the transcript and to help validate the sequences, between 12 to 25 overlapping

sequences were generated. Assembly and annotation of these MRI-R1 sequences

revealed 4 separate alleles of the gene, supporting the allelic polymorphisms observed in

the transcriptomic data (Figure 4C). The pairwise nucleotide sequence identity among the

four MRI-R1 alleles range from 80% to 96.45%. Alleles 1 and 2 share an identical 827

nucleotide intron sequence while alleles 3 and 4 share an identical 376 nucleotide intron

sequences, which accounts for the size discrepancy in the agarose gel electrophoresis

result (Figure 5B).

Analysis of the protein sequences using InterProScan (Jones *et al.*, 2014) showed that

all four alleles contain the CC, NB-ARC, and LRR domains with canonical functional

motifs. The N-terminus of all four alleles begins with an identical MADA motif, a functional

motif conserved in approximately 20% of CC-NLR immune receptors across distantly

related plant species (Bentham *et al.*, 2018; Adachi *et al.*, 2019). The MADA motif has

been shown to be necessary for *Nicotiana benthamiana* NRC4 cell death like the ZAR1

resistosome (Adachi *et al.*, 2019). Similarly, the EDVID motif known to be involved in self-

association, direct interactions with cofactors and, in some cases, cell death signaling

resulting in a hypersensitive response (Bentham *et al.*, 2018), is also present in all 4

alleles. The NB-ARC domain of all four alleles also contains the Walker A (P-loop) motif

(GMFGLGKT) (Ramakrishnan *et al.*, 2002), which is critical for nucleotide binding (Steele

et al., 2019). Also present in the four alleles is the MHD-type motif IHD, which has been

shown to be involved in inhibition of autoactivation of R proteins in the absence of a

pathogen (van Ooijen *et al.*, 2008). The MHD motif is proposed to act as a molecular

switch for R protein activation, and the histidine and aspartate residues are the most

highly conserved across R proteins, with the histidine occupying a critical position in an

ADP-binding pocket (van Ooijen *et al.*, 2008). The most variable regions among these 4

alleles are in and between the CC and NB-ARC domains (Figure S3).

Figure 5. MRI-R1 unique presence, expression, and alleles in MRI. A) Expression of MRI-R1 in *P. belbahrii*-inoculated MRI and SB22 cultivars expressed in TPM (transcripts per million). B) Unique amplification of MRI-R1 from MRI (3673bp) using the external 1BF forward and 14R reverse primers. Internal primer, 9F, paired with 14R produces amplicons of expected size 1206bp in both MRI and SB22. The negative control (NEG) is water in place of gDNA template. C) Gene models of the 4 MRI-R1 alleles with domains and subdomains predicted by InterProScan colored as: coiled coil (gold), NB-ARC (blue), and leucine-rich repeat (green). Introns between the coiled coil and NB-ARC domains are represented as a single line.

Protein structural models of all four alleles were generated in the Robetta server utilizing RoseTTAFold, a top-ranked deep-learning based protein structure prediction method (Baek *et al.*, 2021; Du *et al.*, 2021). The resulting allele protein structures were queried using a distance matrix alignment in the DALI server to search for the closest structural homologs (Holm, 2020).

The resulting top hits for MRI-R1 included plant proteins such as NB-ARC domain from

the tomato immune receptor NRC1 (6S2P), LRR receptor-like serine/threonine-protein

kinase FLS2 (4MN8), and LRR receptor-like serine/threonine-protein kinase GSO1

(6S6Q). The top structural homologs also included human and animal LRR proteins such

as Leucine-rich repeat transmembrane neuronal protein 2 (5Z8X), Dimeric bovine tissue-

extracted decorin (1XCD), and human osteomodulin (5YQ5). The top ten structural

homologs ranged in Z scores of 18-22.2 (with Z scores above 20 indicating definite

homology, and above 8 indicating probable homology), and average deviation in distance

between aligned Cα atoms in 3-D superimpositions, indicated by root mean-squared

deviation (RMSD) ranged from 2.2-8.5 angstroms (Supporting Information Table S3)

(Holm, 2020).

Structural modeling and homology comparison of each allele were used to refine the initial

sequence-based boundary predictions of CC, NB-ARC, and LRR domains (Figure 6A).

All 4 allele structural models were aligned, revealing conservation of predicted functional

domain structures and active sites. As expected, based on high sequence identity, all four

alleles shared high structural similarity, with slight differences in domain boundaries

predicted by the models (Figure 5B). Allele expression analysis showed a clear bias for

expression of MRI-R1 Allele 1 by MRI in response to basil downy mildew infection.

A   B

Figure 6. MRI-R1 protein structural modeling and alignment. a) Predicted protein structure of MRI-R1 allele 1 with domains and motifs colored from N to C terminus as follows: coiled coil domain in yellow, NB-ARC domain in blue, Walker A motif in red, Walker B motif in purple, and LRR domain in green. b) Alignment of predicted structures for all 4 MRI-R1 alleles.

The presence of MRI-R1 in two Backcross progenies 'Devotion' and 'Obsession'

To assess the contribution of MRI-R1 in the four new downy mildew resistant (DMR) basil cultivars that integrated MRI resistance genes, we tested each for the presence of the MRI-R1 gene. A partial sequence of ObActin, the ubiquitous positive control, was successfully amplified from both BDM-susceptible SB22, BDM-resistant MRI, and all four DMR cultivars, while no amplification was detected from the negative water template control (Figure 7). Interestingly, MRI-R1 was detected in only two, 'Devotion' and 'Obsession', out of the four new DMR cultivars using primers designed to amplify the full coding region of the gene (Figure 7). The two amplicons are estimated to be 3064 bp and 3515 bp, corresponding to allele 4 and allele 1 for 'Devotion' and

Page **32** of **55**

'Obsession', respectively. These amplicon sizes include 24bp of the 5' UTR (beginning

with primer 1BF), and the 827bp intron (alleles 1 and 2) or the 376 bp intron (alleles 3

and 4). The six individual clones selected and sequenced from 'Devotion' and

'Obsession' were uniform, indicating that only one allele of MRI-R1 was passed from

MRI to these offspring.

These four new DMR Cultivars were selected from genetic breeding efforts. Briefly, F1

progeny were generated through the MRI (female)×SB22 (male) cross, exhibiting

dominant gene action (Pyne *et al.*, 2015). An F2 family was generated after F1 self-

pollination and a single resistant individual RUMS469-11 was selected for hybridization

with elite sweet basil inbred line 'SB13', which demonstrates downy mildew and

Fusarium wilt tolerance. Twenty individuals achieving the highest category of reduced

disease severity from the RUMS469-11 (female)×SB13 (male) cross were self-

pollinated to generate full sibling families evaluated for response to downy mildew. The

DMR 'Devotion', 'Obsession', 'Passion' and 'Thunderstruck' were selected from these

inbred lines. We anticipate these four new selected DMR cultivars should inherit genetic

resistance genes from both MRI and SB13.

Figure 7. MRI-R1 detection in DMR offspring. Gel electrophoresis showing amplification of ObActin fragment positive control from all basil cultivars tested on the left, MRI-R1 amplification from 'Mrihani' (MRI), 'Devotion', and 'Obsession' on the right using the external 1BF forward and 14R reverse primers. The negative control ('control') is water in place of gDNA template.

Differential upregulation of salicylic acid biosynthesis pathways in MRI

Both NLR and RLK genes interact with hormone signaling pathways to activate host defense (McHale *et al.*, 2006). To understand the involvement of plant hormone signaling pathways involved in susceptibility and resistance responses, we examined the expression of genes involved in ethylene, jasmonic acid (JA), abscisic acid (ABA), jndole-3-acetic acid (Auxin), gibberellic acid (GA), and salicylic acid (SA) based on *A. thaliana* annotation (1e-20, sequence similarity >40%). No significant differences were observed between the MRI and SB22 pattern of expression in ethylene, jasmonic acid, abscisic acid, or gibberellic acid pathway genes (Figure S4, Table S2). We saw a difference in expression profile for two of five auxin genes; YUC1 and TAA1 both were upregulated at early timepoints in MRI, however they were not statistically significantly differentially expressed (p value > 0.05).

Figure 8. **MRI and SB22 diverge transcriptionally at salicylic acid synthesis.** Bar graphs are the average and standard deviation of three replicates. White bars indicate MRI data and gray bars indicate SB22 for each timepoint. Infection releases the repression WRKY70 has on SARD1 expression. Double arrows leading from SID2 and PAL indicate more than one step to the SA molecule.

The most striking difference was observed among genes required for the synthesis of

salicylic acid (Figure 8). Plants possess two biosynthesis pathways to synthesize SA,

both starting from chorismate, but subsequent steps involve either isochorismate

synthase (ICS) or SALICYLIC ACID INDUCTION DEFICIENT 2 (SID2) (Wildermuth *et al.*, 2001) and phenylalanine ammonia-lyase (PAL) (Olsen *et al.*, 2008). There is no

significant difference in PAL expression between the two cultivars, but we did observe a

drastic induction of ICS 24 hpi in MRI. For the water control and 12 hpi samples, there

was no significant difference in ICS expression between MRI and SB22. Compared to

SB22, ICS expression in the MRI cultivar is 15, 6 and 5-fold higher at 24, 48 and 72 hpi,

respectively.

SARD1 is upstream of SID2 in the SA synthesis pathway, and the average SARD1

expression across all time points is 7-fold higher in MRI compared to SB22 and has

roughly 2-, 2.5-, 0- and 3.6-fold increases at 12, 24, 48 and 72 hpi compared to the water

control. Similarly, PR1, a commonly used marker gene downstream of SA synthesis, is

significantly induced in MRI upon pathogen challenge starting at 24 hpi, confirming the

unique increase of SA in MRI upon pathogen challenge.

Together, our data suggest that MRI-derived BDM resistance involves SA. The most

significant change occurred 24 hpi, one day after the encounter of the plant with the

pathogen. WRKY70 is known to repress SARD1 expression in the absence of pathogens

and is required for the activation of some defense genes (Li *et al.*, 2004; Zhou *et al.*,

2018). In MRI, we saw a strong induction in the transcription of WRKY70 at 12 hpi,

increasing expression of SARD1 from water to 24 hpi, and a 23-fold rise in the relative

normalized expression of SID2 at 24 hpi relative to 12 hpi. In SB22, we observed only a

slight increase in WRKY70 expression which was delayed until 24 hpi, and SID2

expression rose only 2.5-fold by 24 hpi and remained effectively stable at later time points.

Similarly, we saw a roughly 2-fold induction in the expression of PR1 in MRI between 12

hpi and 24 hpi, consistent with the upregulation of SID2, while at the same time there was

no increase in PR1 expression in SB22 between 12 and 72 hpi.

DISCUSSION

Here we report the results of transcriptomic sequencing of basil cultivars infected with

Peronospora belbahrii with the goals of identifying genes conferring resistance and

understanding what pathways are involved in susceptibility. We sequenced 33 datasets

to a reasonable coverage, likely capturing most transcripts, although total gene length

was shorter than a previous study in sweet basil (Rastogi *et al.*, 2014), and Trinity's

estimate of gene count in the plant samples was greater than expected. Nonetheless, we

identified roughly 21,000 orthologous genes between cultivars with overall high sequence

conservation consistent with closely related individuals. Phylogenetic analysis supports

the hypothesis that SB22 and MRI are closely related, interspecific genotypes, which was

previously shown using SSRs to confirm conservation of the orthologs in the two breeding

parents (Pyne *et al.*, 2018). These results underscore the close relationship between

cultivars and their sexual compatibility, likely providing us with a pool of true genes shared

in MRI and SB22. At the time of this study, no sweet basil genome was available for use

in a reference-based assembly, though a draft genome for Genovese-type cultivar 'Perrie'

has since been published (Gonda *et al.*, 2020). Furthermore, several unique MRI gene

candidate sequences were used to search the *O. basilicum* draft genome and no

significant BLAST hits were retrieved.

Gene expression clustering identified patterns consistent with MRI unique infection-

expressed genes. Annotated MRI unique candidates were found to be enriched for NLR,

RLK, and secondary metabolic proteins. Analysis of the NLR and malectin-like RLK genes

in both MRI and SB22 identified orthologous pairs and cultivar-unique genes. Members

of the NLR and RLK families are known to act alongside or upstream of disease signaling

pathways and serve as good candidate resistance genes. Secondary metabolite genes

identified in the two cultivars correlate well with known chemotypic characteristics

differentiating MRI and SB22. The top candidate NLR prediction is supported by PCR

screening revealing its presence in resistant MRI and absence in susceptible SB22. This

result was further strengthened with the cloning and sequencing-based confirmation of

MRI-R1 alleles, as well as predicted amino acid sequence analysis and protein structure

modeling supporting the hypothesis that this is a unique NLR likely involved in immune

responses to pathogen infection.

The presence of MRI-R1 was detected in only two out of the four resistant cultivars,

'Devotion' and 'Obsession'. However, recent quantitative trait loci (QTL) analysis

detected at least two major genomic regions (LOD>4.0) that control DM resistance in

the MRI x SB22 F2 mapping population (Pyne *et al.*, 2017), suggesting that the

predicted involvement of MRI-R1 in quantitative disease resistance may be redundant

or have shared function(s) with other NLRs and/or RLKs. We cannot conclusively

determine the functionality of the alleles simply based on the bias for MRI-R1 allele

expression in MRI. The presence of Allele 1 in 'Obsession' and Allele 4 in 'Devotion'

indicates that there may be functional redundancy, and further understanding of the

conserved motifs in these alleles suggests that there may be interacting partners that

have an impact on the activity of these proteins. Nevertheless, we are confident that our

transcriptomic pipeline is powerful in detecting resistant genes involved in the host-

pathogen interactions.

In addition to prediction of specific genes likely conferring resistance, this comparative

transcriptomic approach was also valuable in revealing physiological mechanisms

involved in basil downy mildew resistance. Salicylic acid signaling is an integral part of

plant defense responses and has been demonstrated to be involved in defense against

downy mildews and other biotrophic pathogens (Delaney Terrence P. *et al.*, 1994; Mohr

et al., 2010). Pathogen virulence strategies have developed to overcome and inhibit

salicylic acid defenses, thus enhancing susceptibility to biotrophic pathogens such as

downy mildew organisms (Caillaud *et al.*, 2013, 2016). This suggests a likely role of

salicylic acid signaling in MRI BDM resistance, though the specific mechanisms of

signaling induction remain unknown. We hypothesize that multiple NLRs and RLKs are

active and have interacting and/or redundant roles in mediating the SA signaling pathway

in MRI. If this hypothesis is correct, utilizing multiple targets in a marker-assisted selective

breeding program will be more effective and robust to pass resistance from parents to

progeny in order to slow the evolution of new pathogen races.

This study has utilized mRNA sequencing over an infection time course to provide

strong evidence linking susceptibility to known mechanisms which control defense

responses, and prediction of genes regulating the resistant phenotype. Transcriptomics

without the need for a reference genome is a powerful tool for comparative analyses

given the availability of methods for data annotation and pattern identification. The

strong resistance phenotype of MRI compared to SB22 likely led to the strength of the

visible signal between the cultivars. NLRs that confer resistance to biotrophic pathogens

have increased genetic diversity, and here we observe that, despite high genetic

conservation between orthologs (Fig. 1), there are distinct genetic differences in the

NLR repertoire (Van de Weyer *et al.*, 2019).

Breeding cultivars with quantitative resistance has been shown to produce durable

resistance, typically through the activity of multiple minor-effect genes (Brown, 2015; Niks

et al., 2015). Other BDM resistant cultivars have been produced through interspecific

hybridization of *O. basilicum* with *O. americanum* var. *pilosum*, leading to dominant

resistance against two races of *P. belbahrii* (Ben-Naim & Weitman, 2021). We have

observed that the downy mildew-resistant basil cultivars succumb to infection in

production systems in different regions, consistent with the report of emerging pathogen

races (Ben-Naim & Weitman, 2021). Therefore, understanding the physiological and

molecular bases of host-pathogen interactions is critical to rapidly developing improved

cultivars, monitoring strategies, and management practices. Identification of suitable

molecular markers conferring multiple sources of resistance will improve breeding for

basil downy mildew resistance and advance the understanding of molecular mechanisms

of resistance and pathogenicity. These developments in approaches and knowledge can

be broadly utilized by plant breeders and pathologists working with downy mildew

pathogens on many different crops. After further validation, the genes predicted here can

likely serve as molecular markers for the selection of downy mildew resistant sweet basils,

and investigation and validation of physiological responses to infection may aid in

developing more robust phenotyping assays. This comparative transcriptomics approach

not only revealed candidate resistance genes, but also offered us new insights into

differential infection responses in resistant and susceptible cultivars, which will open new

avenues of investigation to further combat basil downy mildew.

This method of analysis is broadly applicable to two-organism biological systems where

the identification of genes involved in any specific interaction is desired. Although

genomes for holy basil (*O. tenuiflorum*) had been published prior to this study and sweet

basil (*O. basilicum*) and *P. belbahrii* genomes have been published since, the methods

described here worked exclusively from RNA sequencing data and did not require the

whole genome sequence. Any system utilizing two organisms from different kingdoms

could be examined using these methods as DNA sequences are differentiable down to

reasonable taxonomic levels.

Acknowledgements

We thank the MGHPCC for providing high-performance computing capacity for the

RNAseq data analysis. This work was supported by the United States Department of

Agriculture, Specialty Crop Research Initiative, National Institute of Food and Agriculture

(USDA/SCRI/NIFA 2018-51181-28383) and USDA Hatch grant (MASR-2009-04374,

MAS-2018-00496). L.-J.M. is also supported by an Investigator Award in Infectious

Diseases and Pathogenesis by the Burroughs Wellcome Fund (BWF-1014893), the

National Eye Institute of the National Institutes of Health (R01EY030150) and the National

Science Foundation (IOS-1652641).

We thank Chris Joyner and David O'Neil of the College of Natural Sciences Greenhouse

at University of Massachusetts for continued support with plant propagation. We also

thank Erin Patterson for assistance with phylogenetic analysis and Dr. Dilay Hazal Ayhan

for help with allele expression analysis.

Author Contribution

JS, L-JM, RP, and LG participated in the designing of the experiment. RP prepared all

materials for the RNAseq, GAD performed all RNA-seq data analysis. KSA performed

phylogenetic analysis and R-gene cloning, sequencing, and protein structural analyses.

AG directed protein structural and sequence analysis. JM provided initial protein structural

modeling analysis. KSA, GAD and L-JM wrote the manuscript. KSA and GAD prepared

the figures, and all authors edited the paper.

Disclaimer: This article was prepared while Anne Gershenson was employed at the

University of Massachusetts Amherst. The opinions expressed in this article are the

author's own and do not reflect the view of the National Institutes of Health, the

Department of Health and Human Services, or the United States government.

References

References

Adachi H, Contreras MP, Harant A, Wu C, Derevnina L, Sakai T, Duggan C,

Moratto E, Bozkurt TO, Maqbool A, *et al.* 2019. An N-terminal motif in NLR immune

receptors is functionally conserved across distantly related plant species (J-M Zhou, D

Weigel, and J-M Zhou, Eds.). *eLife* **8**: e49956.

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. **1990**. Basic local alignment

search tool. *Journal of Molecular Biology* **215**: 403–410.

Andrews S. **2010**. *FASTQC. A quality control tool for high throuhput sequence data*.

Baek M, DiMaio F, Anishchenko I, Dauparas J, Ovchinnikov S, Lee GR, Wang J,

Cong Q, Kinch LN, Schaeffer RD, *et al.* 2021. Accurate prediction of protein structures

and interactions using a three-track neural network. *Science* **373**: 871–876.

Belbahri L, Calmin G, Pawlowski J, Lefort F. **2005**. Phylogenetic analysis and Real

Time PCR detection of a presumably undescribed Peronospora species on sweet basil

and sage. *Mycological Research* **109**: 1276–1287.

Ben-Naim Y, Weitman M. **2021**. Joint action of Pb1 and Pb2 provide dominant

complementary resistance against new races of Peronospora belbahrii (Basil Downy

Mildew). *Phytopathology®*: PHYTO-02-21-0065-R.

Bentham AR, Zdrzałek R, De la Concepcion JC, Banfield MJ. **2018**. Uncoiling CNLs:

Structure/Function Approaches to Understanding CC Domain Function in Plant NLRs.

Plant and Cell Physiology **59**: 2398–2408.

862 **Bigeard J, Colcombet J, Hirt H**. **2015**. Signaling Mechanisms in Pattern-Triggered

863 Immunity (PTI). *Cell Signaling* **8**: 521–539.

864 **Brown JKM**. **2015**. Durable Resistance of Crops to Disease: A Darwinian Perspective.

865 *Annual Review of Phytopathology* **53**: 513–539.

866 **Caillaud M-C, Asai S, Rallapalli G, Piquerez S, Fabro G, Jones JDG**. **2013**. A downy

867 mildew effector attenuates salicylic acid-triggered immunity in Arabidopsis by interacting

868 with the host mediator complex. *PLoS biology* **11**: e1001732–e1001732.

869 **Caillaud M-C, Asai S, Rallapalli G, Piquerez S, Fabro G, Jones JDG**. **2016**.

870 Correction: A Downy Mildew Effector Attenuates Salicylic Acid-Triggered Immunity in

871 Arabidopsis by Interacting with the Host Mediator Complex. *PLOS Biology* **14**:

872 e1002408.

873 **Cesari S**. **2018**. Multiple strategies for pathogen perception by plant immune receptors.

874 *New Phytologist* **219**: 17–24.

875 **Chern M, Xu Q, Bart RS, Bai W, Ruan D, Sze-To WH, Canlas PE, Jain R, Chen X,**

876 **Ronald PC**. **2016**. A Genetic Screen Identifies a Requirement for Cysteine-Rich–

877 Receptor-Like Kinases in Rice NH1 (OsNPR1)-Mediated Immunity. *PLOS Genetics* **12**:

878 e1006049.

879 **Choi YJ, Choi IY, Lee KJ, Shin HD**. **2016**. First Report of Downy Mildew Caused by

880 Peronospora belbahrii on Sweet Basil (Ocimum basilicum) in Korea. *Plant Disease* **100**:

881 2335–2335.

Cohen Y, Ben Naim Y, Falach L, Rubin AE. **2017**. Epidemiology of Basil Downy

Mildew. *Phytopathology®* **107**: 1149–1160.

Cohen Y, Ben-Naim Y. **2016**. Nocturnal Fanning Suppresses Downy Mildew Epidemics

in Sweet Basil. *PLOS ONE* **11**: e0155330.

Cohen Y, Rubin AE. **2015**. Daytime Solar Heating Controls Downy Mildew

Peronospora belbahrii in Sweet Basil. *PLOS ONE* **10**: e0126103.

Cohen Y, Vaknin M, Ben-Naim Y, Rubin AE. **2013**. Light Suppresses Sporulation and

Epidemics of Peronospora belbahrii. *PLOS ONE* **8**: e81282.

Cui H, Tsuda K, Parker JE. **2015**. Effector-Triggered Immunity: From Pathogen

Perception to Robust Defense. *Annual Review of Plant Biology* **66**: 487–511.

(dataset) USDA National Agricultural Statistics Service (2017). USDA National

Agricultural Statistics Service.

Delaney Terrence P., Uknes Scott, Vernooij Bernard, Friedrich Leslie, Weymann

Kris, Negrotto David, Gaffney Thomas, Gut-Rella Manuela, Kessmann Helmut,

Ward Eric, *et al.* **1994**. A Central Role of Salicylic Acid in Plant Disease Resistance.

Science **266**: 1247–1250.

Dodds PN, Rathjen JP. **2010**. Plant immunity: towards an integrated view of plant–

pathogen interactions. *Nature Reviews Genetics* **11**: 539–548.

Du Z, Su H, Wang W, Ye L, Wei H, Peng Z, Anishchenko I, Baker D, Yang J. **2021**.

The trRosetta server for fast and accurate protein structure prediction. *Nature Protocols*

16: 5634–5651.

Finn RD, Coggill P, Eberhardt RY, Eddy SR, Mistry J, Mitchell AL, Potter SC,

Punta M, Qureshi M, Sangrador-Vegas A, *et al.* **2016**. The Pfam protein families

database: towards a more sustainable future. *Nucleic Acids Research* **44**: D279–D285.

Garibaldi A, Minuto A, Gullino ML. **2005**. First Report of Downy Mildew Caused by

Peronospora sp. on Basil (Ocimum basilicum) in France. *Plant Disease* **89**: 683–683.

Garibaldi A, Minuto A, Minuto G, Gullino ML. **2004**. First Report of Downy Mildew on

Basil (Ocimum basilicum) in Italy. *Plant Disease* **88**: 312–312.

Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X,

Fan L, Raychowdhury R, Zeng Q, *et al.* **2011**. Full-length transcriptome assembly

from RNA-Seq data without a reference genome. *Nature Biotechnology* **29**: 644–652.

Haas BJ, Kamoun S, Zody MC, Jiang RHY, Handsaker RE, Cano LM, Grabherr M,

Kodira CD, Raffaele S, Torto-Alalibo T, *et al.* **2009**. Genome sequence and analysis

of the Irish potato famine pathogen Phytophthora infestans. *Nature* **461**: 393–398.

Holm L. **2020**. Using Dali for Protein Structure Comparison. In: Methods in Molecular

Biology. Structural Bioinformatics. New York, NY: Humana, 29–42.

Homa K, Barney WP, Ward DL, Wyenandt CA, Simon JE. **2014**. Evaluation of

Fungicides for the Control of Peronospora belbahrii on Sweet Basil in New Jersey. *Plant*

Disease **98**: 1561–1566.

Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J,

Mitchell A, Nuka G, *et al.* 2014. InterProScan 5: genome-scale protein function

classification. *Bioinformatics* **30**: 1236–1240.

Jones JDG, Dangl JL. **2006**. The plant immune system. *Nature* **444**: 323–329.

Kanetis L, Vasiliou A, Neophytou G, Samouel S, Tsaltas D. **2014**. First Report of

Downy Mildew Caused by Peronospora belbahrii on Sweet Basil (Ocimum basilicum) in

Cyprus. *Plant Disease* **98**: 283–283.

Khateri H, Calmin G, Moarrefzadeh, Lassaad B, Lefort F. **2007**. First report of downy

mildew caused by Peronospora sp. on basil in Northern Iran. *JOURNAL OF PLANT*

PATHOLOGY **89**: S70.

Krasileva KV, Dahlbeck D, Staskawicz BJ. **2010**. Activation of an *Arabidopsis*

Resistance Protein Is Specified by the in Planta Association of Its Leucine-Rich Repeat

Domain with the Cognate Oomycete Effector. *The Plant Cell* **22**: 2444–2458.

Langmead B, Trapnell C, Pop M, Salzberg SL. **2009**. Ultrafast and memory-efficient

alignment of short DNA sequences to the human genome. *Genome Biology* **10**: R25.

Li B, Dewey CN. **2011**. RSEM: accurate transcript quantification from RNA-Seq data

with or without a reference genome. *BMC Bioinformatics* **12**: 323.

Li J, Brader G, Palva ET. **2004**. The WRKY70 Transcription Factor: A Node of

Convergence for Jasmonate-Mediated and Salicylate-Mediated Signals in Plant

Defense[W]. *The Plant Cell* **16**: 319–331.

Madeira F, Park YM, Lee J, Buso N, Gur T, Madhusoodanan N, Basutkar P, Tivey

ARN, Potter SC, Finn RD, *et al.* 2019. The EMBL-EBI search and sequence analysis

tools APIs in 2019. *Nucleic Acids Res* **47**: W636–W641.

Maekawa T, Cheng W, Spiridon LN, Töller A, Lukasik E, Saijo Y, Liu P, Shen Q-H,

Micluta MA, Somssich IE, *et al.* 2011. Coiled-Coil Domain-Dependent

Homodimerization of Intracellular Barley Immune Receptors Defines a Minimal

Functional Module for Triggering Cell Death. *Cell Host & Microbe* **9**: 187–199.

Marchler-Bauer A, Derbyshire MK, Gonzales NR, Lu S, Chitsaz F, Geer LY, Geer

RC, He J, Gwadz M, Hurwitz DI, *et al.* 2015. CDD: NCBI's conserved domain

database. *Nucleic Acids Research* **43**: D222–D226.

Martínez de la Parte E, Pérez-Vicente L, Bernal B, García D. **2010**. First report of

Peronospora sp. on sweet basil (Ocimum basilicum) in Cuba. *Plant Pathology* **59**: 800–

800.

McGrath MT. **2020b**. Efficacy of Conventional Fungicides for Downy Mildew in Field-

Grown Sweet Basil in the United States. *Plant Disease* **104**: 2967–2972.

McGrath MT. **2021**. Where in the USA is Basil Downy Mildew?

McGrath MT. **2022a**. Basil Downy Mildew. *Cornell Vegetables*.

McHale L, Tan X, Koehl P, Michelmore RW. **2006**. Plant NBS-LRR proteins:

adaptable guards. *Genome Biology* **7**: 212.

McLeod A, Coertze S, Mostert L. **2006**. First Report of a Peronospora Species on

Sweet Basil in South Africa. *Plant Disease* **90**: 1115–1115.

Mendy B, Wang'ombe MW, Radakovic ZS, Holbein J, Ilyas M, Chopra D, Holton N,

Zipfel C, Grundler FMW, Siddique S. **2017**. Arabidopsis leucine-rich repeat receptor–

like kinase NILR1 is required for induction of innate immunity to parasitic nematodes.

PLOS Pathogens **13**: e1006284.

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler

A, Lanfear R. **2020**. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic

Inference in the Genomic Era. *Molecular Biology and Evolution* **37**: 1530–1534.

Mohr TJ, Mammarella ND, Hoff T, Woffenden BJ, Jelesko JG, McDowell JM. **2010**.

The Arabidopsis Downy Mildew Resistance Gene RPP8 Is Induced by Pathogens and

Salicylic Acid and Is Regulated by W Box cis Elements. *Molecular Plant-Microbe*

Interactions® **23**: 1303–1315.

Monteiro F, Nishimura MT. **2018**. Structural, Functional, and Genomic Diversity of

Plant NLR Proteins: An Evolved Resource for Rational Engineering of Plant Immunity.

Annual Review of Phytopathology **56**: 243–267.

Nagy G, Horváth A. **2011**. Occurrence of Downy Mildew Caused by Peronospora

belbahrii on Sweet Basil in Hungary. *Plant Disease* **95**: 1034–1034.

Nelson R, Wiesner-Hanks T, Wisser R, Balint-Kurti P. **2018**. Navigating complexity to

breed disease-resistant crops. *Nature Reviews Genetics* **19**: 21–33.

Niks RE, Qi X, Marcel TC. **2015**. Quantitative Resistance to Biotrophic Filamentous

Plant Pathogens: Concepts, Misconceptions, and Mechanisms. *Annual Review of*

Phytopathology **53**: 445–470.

van Ooijen G, Mayr G, Kasiem MMA, Albrecht M, Cornelissen BJC, Takken FLW.

2008. Structure–function analysis of the NB-ARC domain of plant disease resistance

proteins. *Journal of Experimental Botany* **59**: 1383–1397.

Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin

TE. **2004**. UCSF Chimera—A visualization system for exploratory research and

analysis. *Journal of Computational Chemistry* **25**: 1605–1612.

Prigozhin DM, Krasileva KV. **2020**. *Intraspecies diversity reveals a subset of highly*

variable plant immune receptors and predicts their binding sites. Plant Biology.

Prigozhin DM, Krasileva KV. Analysis of intraspecies diversity reveals a subset of

highly variable plant immune receptors and predicts their binding sites. : 35.

Pyne R, Honig J, Vaiciunas J, Koroch A, Wyenandt C, Bonos S, Simon J. **2017**. A

first linkage map and downy mildew resistance QTL discovery for sweet basil (Ocimum

basilicum) facilitated by double digestion restriction site associated DNA sequencing

(ddRADseq) (T Yin, Ed.). *PLOS ONE* **12**: e0184319.

Pyne RM, Honig JA, Vaiciunas J, Wyenandt CA, Simon JE. **2018**. Population

structure, genetic diversity and downy mildew resistance among Ocimum species

germplasm. *BMC Plant Biology* **18**: 69.

Pyne RM, Koroch AR, Wyenandt CA, Simon JE. **2015**. Inheritance of Resistance to

Downy Mildew in Sweet Basil. *Journal of the American Society for Horticultural Science*

140: 396–403.

Ramakrishnan C, Dani VS, Ramasarma T. **2002**. A conformational analysis of Walker

motif A [GXXXXGKT (S)] in nucleotide-binding and other proteins. *Protein Engineering,*

Design and Selection **15**: 783–798.

Rastogi S, Kalra A, Gupta V, Khan F, Lal RK, Tripathi AK, Parameswaran S,

Gopalakrishnan C, Ramaswamy G, Shasany AK. **2015**. Unravelling the genome of

Holy basil: an "incomparable" "elixir of life" of traditional Indian medicine. *BMC*

Genomics **16**: 413.

Rastogi S, Meena S, Bhattacharya A, Ghosh S, Shukla RK, Sangwan NS, Lal RK,

Gupta MM, Lavania UC, Gupta V, *et al.* **2014**. De novo sequencing and comparative

analysis of holy and sweet basil transcriptomes. *BMC Genomics* **15**: 588.

Robert X, Gouet P. **2014**. Deciphering key features in protein structures with the new

ENDscript server. *Nucleic Acids Research* **42**: W320–W324.

Roberts PD, Raid RN, Harmon PF, Jordan SA, Palmateer AJ. **2009**. First Report of

Downy Mildew Caused by a Peronospora sp. on Basil in Florida and the United States.

Plant Disease **93**: 199–199.

Robinson MD, McCarthy DJ, Smyth GK. **2010**. edgeR: a Bioconductor package for

differential expression analysis of digital gene expression data. *Bioinformatics* **26**: 139–

140.

Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G,

Mesirov JP. **2011**. Integrative genomics viewer. *Nature Biotechnology* **29**: 24–26.

Ronco L, Rollán C, Choi YJ, Shin HD. **2009**. Downy mildew of sweet basil (Ocimum

basilicum) caused by Peronospora sp. in Argentina. *Plant Pathology* **58**: 395–395.

Šafránková I, Holková L. **2014**. The First Report of Downy Mildew Caused by

Peronospora belbahrii on Sweet Basil in Greenhouses in the Czech Republic. *Plant*

Disease **98**: 1579–1579.

Sawano Y, Miyakawa T, Yamazaki H, Tanokura M, Hatano K. **2007**. Purification,

characterization, and molecular gene cloning of an antifungal protein from Ginkgo biloba

seeds. **388**: 273–280.

Simon JE, Pyne RM, Wyenandt CA. **2018**. Downy mildew resistant/tolerant sweet

basil varieties. 10159212, filed November 14, 2017, and issued December 25, 2018.

https://patents.justia.com/patent/10159212.

Steele JFC, Hughes RK, Banfield MJ. **2019**. Structural and biochemical studies of an

NB-ARC domain from a plant NLR immune receptor. *PLOS ONE* **14**: e0221226.

Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. **2013**. MEGA6: Molecular

Evolutionary Genetics Analysis version 6.0. *Molecular biology and evolution* **30**: 2725–

2729.

Van de Weyer A-L, Monteiro F, Furzer OJ, Nishimura MT, Cevik V, Witek K, Jones

JDG, Dangl JL, Weigel D, Bemm F. **2019**. A Species-Wide Inventory of NLR Genes

and Alleles in Arabidopsis thaliana. *Cell* **178**: 1260-1272.e14.

Vieira RF, Simon JE. **2006**. Chemical characterization of basil (Ocimum spp.) based on

volatile oils. *Flavour and Fragrance Journal* **21**: 214–221.

Wang J, Chai J. **2020**. Structural Insights into the Plant Immune Receptors PRRs and

NLRs. *Plant physiology* **182**: 1566–1581.

Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ. **2009**. Jalview

Version 2—a multiple sequence alignment editor and analysis workbench.

Bioinformatics **25**: 1189–1191.

Wildermuth MC, Dewdney J, Wu G, Ausubel FM. **2001**. Isochorismate synthase is

required to synthesize salicylic acid for plant defence. *Nature* **414**: 562–565.

Wu C-H, Abd-El-Haliem A, Bozkurt TO, Belhaj K, Terauchi R, Vossen JH, Kamoun

S. **2017**. NLR network mediates immunity to diverse plant pathogens. *Proceedings of*

the National Academy of Sciences **114**: 8113–8118.

Wyenandt CA, Simon JE, Pyne RM, Homa K, McGrath MT, Zhang S, Raid RN, Ma

L-J, Wick R, Guo L, *et al.* 2015. Basil Downy Mildew (*Peronospora belbahrii*):

Discoveries and Challenges Relative to Its Control. *Phytopathology®* **105**: 885–894.

Zhang L, Tian L-H, Zhao J-F, Song Y, Zhang C-J, Guo Y. **2009**. Identification of an

apoplastic protein involved in the initial phase of salt stress response in rice root by two-

dimensional electrophoresis. *Plant physiology* **149**: 916–928.

Zhou M, Lu Y, Bethke G, Harrison BT, Hatsugai N, Katagiri F, Glazebrook J. **2018**.

WRKY70 prevents axenic activation of plant immunity by direct repression of SARD1.

New Phytologist **217**: 700–712.

Zhou J-M, Zhang Y. **2020**. Plant Immunity: Danger Perception and Signaling. *Cell* **181**:

978–989.

1066

1067

1068

1069

1070

1071

1072

1073

1074

1075

1076

Supporting Information

Additional supporting information may be found in the online version of this article.

Fig. S1 Average Quality Scores. Average quality scores plotted on the Y-axis by base number on

the X-axis for all paired-end reads.

Fig. S2 Pathogen Read Percentage. Percentage of reads mapped to the *Peronospora belbahrii*

sporangia data in MRI (black bars) and SB22 (grey bars) combined data at each timepoint

plotted on the X-axis.

Table S1 Primers Used In This Study

Table S2 Coarse Clustering Results

Table S3 Top ten structural homologs of MRI-R1 identified by DALI PDB Search