Regulation by cyclic-di-GMP attenuates dynamics and enhances robustness of bimodal curli gene activation in *Escherichia coli*

Olga Lamprecht[1#†], Maryia Ratnikava[1#], Paulina Jacek[1], Eugen Kaganovitch[1], Nina Buettner[1], Kirstin Fritz[1], Ina Biazruchka[1], Robin Köhler[1], Julian Pietsch[1], and Victor Sourjik[1]*

[1]Max Planck Institute for Terrestrial Microbiology and Center for Synthetic Microbiology (SYNMIKRO), Karl-von-Frisch Strasse 14, 35043 Marburg, Germany

[#]These authors contributed equally to this work

[†]Present address: Eawag: Swiss Federal Institute of Aquatic Science and Technology, Dübendorf, Switzerland

*Corresponding author: Victor Sourjik, victor.sourjik@synmikro.mpi-marburg.mpg.de

Abstract

Curli amyloid fibers are a major constituent of the extracellular biofilm matrix formed by bacteria of the Enterobacteriaceae family. Within *Escherichia coli* biofilms, curli gene expression is limited to a subpopulation of bacteria, leading to heterogeneity of extracellular matrix synthesis. Here we show that bimodal activation of curli expression occurs not only in submerged and macrocolony biofilms, but also in well-mixed planktonic cultures of *E. coli*, resulting in all-or-none stochastic differentiation into distinct subpopulations of curli-positive and curli-negative cells at the entry into the stationary phase of growth. Stochastic curli activation in individual *E. coli* cells could further be observed during continuous growth in a conditioned medium in a microfluidic device, which further revealed that the curli-positive state is only metastable. In agreement with previous reports, regulation of curli gene expression by c-di-GMP via two pairs of diguanylate cyclase and phosphodiesterase enzymes, DgcE/PdeH and DgcM/PdeR, modulates the fraction of curli-positive cells under all tested growth conditions. Unexpectedly, removal of this regulatory network does not abolish the bimodality of curli gene expression, although it affects dynamics of activation and increases heterogeneity of expression levels among individual cells. Moreover, the fraction of curli-positive cells within an *E. coli* population shows stronger dependence on growth conditions in the absence of c-di-GMP regulation. We thus conclude that, while not required for the emergence of bimodal curli gene expression in *E. coli*, this c-di-GMP regulatory network attenuates the frequency and dynamics of gene activation and increases its robustness to cellular heterogeneity and environmental variation.

Keywords: Gene expression, bacteria, biofilm, bistability, differentiation, amyloid fibers

Introduction

Curli amyloid fibers are the key component of the extracellular matrix produced during biofilm formation by *Escherichia coli*, *Salmonella enterica*, and other Enterobacteriaceae [1-9]. In *E. coli* and *S. enterica* serovar Typhimurium, curli genes are organized in two divergently transcribed *csgBAC* and *csgDEFG* operons that share a common intergenic regulatory region [10]. Expression of these operons is under regulation of the stationary phase sigma factor σ^S (RpoS) and thus becomes activated during the entry into the stationary phase of growth [4, 11-14]. This activation is achieved by the σ^S-dependent induction of the transcriptional regulator CsgD, which then controls the expression of the *csgBAC* operon that encodes the major curli subunit CsgA along with the curli nucleator CsgB and the chaperone CsgC [7, 8, 15]. In turn, *csgD* expression in *E. coli* and *S.* Typhimurium is either directly or indirectly regulated by multiple cellular factors that mediate responses to diverse environmental changes, including both global and specific transcriptional regulators, small regulatory RNAs and second messengers (reviewed in [16-19]).

One of the key regulators of *csgD* is the transcription factor MlrA [13, 14, 20, 21]. The activity of MlrA depends on cellular levels of bacterial second messenger bis-(3'-5')-cyclic dimeric guanosine monophosphate (c-di-GMP), and in *E. coli* this control is known to be mediated by a pair of the interacting diguanylate cyclase (DGC) and phosphodiesterase (PDE) enzymes, DgcM and PdeR, that form a ternary complex with MlrA [12, 14, 22]. MlrA is kept inactive by binding PdeR, and this interaction is relieved when the latter becomes active as a PDE thus acting as the trigger enzyme [22, 23]. This inhibition is counteracted both by DgcM, which locally produces c-di-GMP to engage PdeR, as well as by the global pool of c-di-GMP. Besides its enzymatic activity, DgcM might also activate MlrA through direct protein interaction. Another DGC-PDH pair, DgcE and PdeH, provides global regulatory input into the local DgcM-PdeR-MlrA regulation [12, 24]. At least under typical conditions used to study *E. coli* biofilms, this regulatory network appears to be the only c-di-GMP-dependent input that controls *csgD* expression [24].

Previous studies of *E. coli* macrocolony biofilms formed on agar plates showed that curli expression occurs in the upper layer of the colony, but even in this layer its expression remained heterogeneous [25-27], indicating an interplay between global regulation of curli gene expression by microenvironmental gradients within biofilms and its inherent stochasticity. Differentiation of *E. coli* into distinct subpopulations of cells either expressing or not expressing curli was also observed in submerged biofilms formed in liquid cultures, whereby curli expression was associated with cellular aggregation [28]. Furthermore, bi- or multimodality of *csgD* reporter activity was also observed in the early stationary phase among planktonic cells in *S.* Typhimurium [29, 30] and *E. coli* [27]. Given established c-di-GMP-dependent regulation of CsgD activity, it was proposed that bistable curli expression originates from a toggle switch

created by mutual inhibition between DgcM and PdeR, which could act as a bistable switch on
csgD gene expression [27, 31].

In this study we demonstrate that differentiation of *E. coli csgBAC* operon expression, leading
to formation of distinct subpopulations of curli-positive and -negative cells, occurs not only in
submerged and macrocolony biofilms but also stochastically in a well-stirred planktonic culture
and thus in the absence of any environmental gradients. Similar stochastic and reversible
differentiation could be further observed among cells growing in conditioned medium in the
microfluidic device. Although the c-di-GMP regulatory network consisting of the DgcE/PdeH
and DgcM/PdeR pairs modulates the fraction of curli-positive cells in the population and the
dynamics and uniformity of curli gene activation, it is not required to establish the bimodality of
their expression.

Materials and methods

Bacterial strains and plasmids

All strains and plasmids used in this study are listed in Table S1. A derivative of *E. coli* W3110
[26] that was engineered to encode a chromosomal transcriptional sfGFP reporter downstream
of the *csgA* gene [28] (VS1146) was used here as the wild-type strain. Gene deletions were
obtained with the help of P1 phage transduction using strains of the Keio collection [32] as
donors, and the kanamycin resistance cassette was removed using FLP recombinase [33].
For expression, *dgcE* and *pdeH* genes were cloned into the pTrc99A vector [34].

Growth conditions for planktonic cultures

Planktonic *E. coli* cultures were grown in tryptone broth (TB) medium (10 g tryptone, 5 g NaCl
per liter, pH 7.0) or lysogeny broth (LB) medium without salt (10 g tryptone, 5 g yeast extract,
per liter, pH 7.0), supplemented with antibiotics where necessary. Overnight cultures were
inoculated from LB agar plates, grown at 30°C and diluted 1:100, unless indicated otherwise,
in 5-10 ml of fresh TB and grown at 30°C at 200 rpm in 100 ml flasks in a rotary shaker until
the indicated OD_{600} or overnight (18-25 h; OD_{600} ~ 1.3-1.8). Alternatively, cultures were grown
in TB in 96-well plates with orbital shaking in a plate reader, with 200 µl culture per well. Where
indicated, bacterial cultures were supplemented with either 1 mM L-serine (after 6 h of growth)
or 0.1–10 mg/l DL-serine hydroxamate at inoculation.

Growth and quantification of submerged biofilms

Submerged biofilms were grown and quantified as described previously [28], with minor
modifications. Overnight bacterial cultures grown in TB were diluted 1:100 in fresh TB medium
and grown at 200 rpm and 30°C in a rotary shaker to OD_{600} of 0.5. The cultures were then
diluted in fresh TB medium to a final OD_{600} of 0.05, and 300 µl was loaded into a 96-well plate

(Corning Costar, flat bottom; Sigma-Aldrich, Germany) and incubated without shaking at 30°C

for 46 h.

For quantification of biofilm formation, the non-attached cells were removed and the wells were

washed once with phosphate-buffered saline (PBS; 8 g NaCl, 0.2 g KCl, 1.44 g Na_2HPO_4, 0.24

g KH_2PO_4). Attached cells were fixed for 20 min with 300 µl of 96% ethanol, allowed to dry for

40 min, and stained with 300 µl of 0.1% crystal violet (CV) solution for 15 min at room

temperature. The wells were subsequently washed twice with 1x PBS, incubated with 300 µl

of 96% ethanol for 35 min and the CV absorption was measured at OD_{595} using INFINITE M

NANO$^+$ plate reader (Tecan Group Ltd., Switzerland). Obtained CV values were normalized to

the OD_{600} values of the respective biofilm cultures.

Macrocolony biofilm assay

123   Macrocolony biofilms were grown as described previously [26]. Briefly, 5 µl of the overnight

124   liquid culture grown at 37°C in LB medium (10 g tryptone, 5 g NaCl, and 5 g yeast extract per

125   liter) was spotted on salt-free LB agar plates supplemented with Congo red (40 µg/ml). Plates

126   were incubated for 8 days at 28°C.

Fluorescence measurements

Measurements of GFP expression in an INFINITE M1000 PRO plate reader (Tecan Group

Ltd., Switzerland) were done using fluorescence excitation at 483 nm and emission at 535 nm.

Relative fluorescence was calculated by normalizing to corresponding OD_{600} values of the

culture.

For fluorescence measurements using flow cytometry, aliquots of 40-300 µl of liquid bacterial

cultures were mixed with 2 ml of tethering buffer (10 mM KH_2PO_4, 10 mM K_2HPO_4, 0.1 mM

EDTA, 1 µM L-methionine, 10 mM lactic acid, pH 7.0). Macrocolonies were collected from the

plate, resuspended in 10 ml of tethering buffer and then aliquots of 40 µl were mixed with 2 ml

of fresh tethering buffer. Samples were vigorously vortexed and then immediately subjected to

flow cytometric analysis using BD LSRFortessa Sorp cell analyzer (BD Biosciences, Germany)

using 488-nm laser. In each experimental run, 50,000 individual cells were analyzed. Absence

of cell aggregation was confirmed by using forward scatter (FSC) and side scatter (SSC)

parameters. Data were analyzed using FlowJo software version v10.7.1 (FlowJo LLC,

Ashland, OR, US).

Microfluidics

145   Conditioned medium was prepared by cultivating wild-type *E. coli* in TB in a rotary shaker at

146   30°C for 20 h, after which the cell suspension was centrifuged at 4000 rpm for 10 min, medium

147   was filter-sterilized and stored at 4°C. Mother machine [35] microfluidics devices were

designed, fabricated and operated as described in Supporting protocols. *E. coli* cells from the overnight culture in TB were loaded into the mother machine by manual infusion of the cell suspension through one of the two inlets using a 1-ml syringe. Cells were first allowed to grow at 30 °C for 4 h in fresh TB, then switched to the conditioned TB and cultivated for up to 26 h. Phase contrast and GFP fluorescence images were acquired using a Nikon Eclipse Ti-E inverted microscope with a time interval of 10 min. Details of image analysis are described in Supporting protocols.

Results

Bimodal curli gene expression is induced in planktonic culture

In order to characterize curli expression in planktonic culture of *E. coli*, we followed the induction of a chromosomal transcriptional reporter for the *csgBAC* operon, where the gene encoding for stable green fluorescent protein (GFP) was cloned downstream of *csgA* with a strong ribosome binding site as part of the same polycistronic RNA [28]. In our previous study of submerged *E. coli* biofilms, this reporter showed bimodal expression both in the surface-attached biofilm and in the pellicle at the liquid-air interface, and its activity correlated with the recruitment of individual cells into multicellular aggregates [28]. When *E. coli* culture was grown at 30°C in tryptone broth (TB) liquid medium, this reporter became induced during the transition to stationary phase (Figure 1A), which is consistent with previous reports [12, 14, 27]. The GFP reporter level remains high in the stationary phase, which also explains its initially high fluorescence in our plate reader experiments, where the culture is inoculated using stationary-phase cells. The observed induction of curli expression occurred at similar densities in the cultures with different initial inoculum size. In both cases the onset of induction apparently coincided with the reduction of the growth rate, which might occur due to depletion of amino acids in the medium and induction of the stringent response [36, 37], consistent with the proposed role of the stringent response in the regulatory cascade leading to curli gene expression [18, 23]. In agreement with that, curli expression was strongly reduced when *E. coli* cultures were grown in a concentrated TB medium (Figure S1A) or when TB medium was supplemented with serine (Figure S2A). Moreover, curli reporter induction was enhanced by the addition of serine hydroxamate (SHX), which is known to mimic amino acid starvation and induce the stringent response [38] (Figure S2A).

In order to investigate whether curli expression was uniform or heterogeneous within planktonic *E. coli* populations, we next measured curli reporter activity in individual cells using flow cytometry. The reporter was induced only in a fraction of cells, and this bimodality of curli expression became increasingly more pronounced at later stages of culture growth, reaching its maximum in the overnight culture (Figure 1B). Thus, the bimodal induction of curli gene expression is observed not only in biofilms but also in well-mixed planktonic cultures. Upon

inspection of the flask culture by microscopy, no cell aggregation could be observed,
suggesting that curli gene induction in a subpopulation of bacteria is not due to the formation
of suspended biofilm-like aggregates [39]. While curli activation was more pronounced in a cell
culture growing in an orbital shaker in a flask (Figure 1B), bimodality was also observed for
cultures grown in a plate reader (Figure S1B and Figure S2B). Notably, stimulation of curli
expression by SHX, or its suppression by additional nutrients, affected the fraction of positive
cells rather than their expression levels (Figure S1B and Figure S2B).
192

Bimodality of curli gene expression does not require the known c-di-GMP regulatory
network
Subsequently, we investigated dependence of bimodal activation of curli gene expression in
planktonic culture on the known regulation of curli expression in *E. coli* by the global (DgcE
and PdeH) and local (DgcM and PdeR) pairs of DGC/PDE enzymes [12, 14, 22] (Figure 2A).
In agreement with this model of c-di-GMP-dependent regulation, activation of curli reporter
was completely abolished in the absence of MlrA (Figure 2B), and it was strongly reduced by
deletions of *dgcE* and *dgcM* and enhanced by deletions of *pdeH* and *pdeR* genes (Figure 2C).
Notably, under these conditions a small fraction of curli-positive cells could still be detected in
cultures of both DGC gene deletion strains, and a small fraction of curli-negative cells was
observed for both PDE gene deletion strains, indicating that deletions of these genes do not
eliminate the bimodality of curli gene expression or its levels in curli-positive cells but rather
change the probability of curli genes to become activated in individual cells.
Consistently, bimodality was also retained upon combined deletions of pairs of DGC/PDE
genes. Removal of the entire global level of c-di-GMP regulation led to a bimodal pattern of
curli reporter activation within the population of Δ*pdeH* Δ*dgcE* cells (Figure 2D) that was similar
to that observed in the wild-type culture, despite a smaller fraction of curli-positive cells in the
deletion strain. Even more surprisingly, the distribution of curli reporter expression remained
bimodal upon removal of the local level of c-di-GMP regulation in Δ*pdeR* Δ*dgcM* strain,
although the fraction of curli-positive cells was reduced in this background, too.
We observed that level of curli gene expression in Δ*pdeR* Δ*dgcM* strain was lower than in
Δ*pdeR* strain, which confirms that DgcM can promote curli expression independently of PdeR
[12, 14, 22] (Figure 2A). This conclusion is further supported by the comparison between the
triple deletion strain Δ*pdeH* Δ*dgcE* Δ*pdeR* that expresses only DgcM (Figure 2E) and the
quadruple deletion strain Δ*pdeH* Δ*dgcE* Δ*pdeR* Δ*dgcM* that lacks the entire regulatory network
(Figure 2F), with much higher fraction of culri-positive cells observed in the former background.
Most importantly, the bimodality of curli expression was still observed in the quadruple deletion
strain (Figure 2F).

We thus conclude that, whereas the known c-di-GMP-dependent regulation of MlrA activity
does clearly affect the fraction of curli-positive cells, it is not required to establish bimodality of
curli gene expression in the planktonic cell population. The network has further only little impact
of the level of curli reporter activity in individual positive cells (i.e., on position of the positive
peak in the flow cytometry data), although reporter intensity in positive cells appeared to be
slightly reduced in the strains lacking DgcM.
Vibrio cholerae transcriptional regulator VpsT, a close homologue of CsgD, has been shown
to be directly regulated by binding to c-di-GMP [40]. Furthermore, in both *S.* Typhimurium and
E. coli, curli gene expression might also be regulated by c-di-GMP independently of *csgD*
transcription [41-43]. We thus aimed to examine whether *E. coli* curli gene expression was no
longer sensitive to the global cellular level of c-di-GMP in the absence of the local PdeR/DgcM
regulatory module. Indeed, whereas the overexpression of DgcE or of PdeH, which both affect
the global pool of c-di-GMP in *E. coli*, had strong impacts on the fraction of curli-positive cells
in the wild type, the quadruple mutant was insensitive to such overexpression (Figure 3). This
result suggests that in this background, the expression of the *csgBAC* reporter is indeed no
longer affected by global c-di-GMP levels. This further excludes regulation by other DGCs that
contribute to the common c-di-GMP pool, although it does not rule out hypothetical regulation
by a local DGC/PDE pair that is insensitive to global c-di-GMP levels.
239
Curli gene activation shows higher growth-condition dependent variability in the
absence of the c-di-GMP regulatory network
We next explored how the fraction of curli-positive cells in the population depends on the
conditions of culture growth, with or without regulation by c-di-GMP. As noted above, when
the wild-type *E. coli* culture was grown in multi-well plates, both the fraction of curli-positive
cells and the level of reporter activity in positive cells were moderately reduced compared to
incubation in a flask in an orbital shaker (Figure 2B, Figure 4A and Figure S1B). The reduction
in the number of curli-positive cells under these growth conditions was even more pronounced
for Δ*pdeR* Δ*dgcM* or Δ*pdeH* Δ*dgcE* Δ*pdeR* Δ*dgcM* strains (Figure 4A and Figure S3), where
only a small fraction of cells (12%) became positive, compared to 40-50% in the flask culture
(Figure 2D,F). Interestingly, this difference was less for the Δ*pdeH* Δ*dgcE* strain (24% vs 30%)*.*
Deletions of individual DGC or PDE genes generally showed expected phenotypes, but
fractions of curli-positive cells in Δ*pdeH* and Δ*pdeR* strains were again reduced compared with
flask cultures (Figure 2C). Another notable difference between flask and multi-well plate
cultures was that the low-fluorescence peak of the wild-type culture was not fully negative but
apparently contained a large fraction of cells with incompletely activated curli reporter, which
could also be seen in the Δ*pdeH* or Δ*pdeR* strains but not in the Δ*pdeH* Δ*dgcE* Δ*pdeR* Δ*dgcM,*
Δ*pdeR* Δ*dgcM* or Δ*pdeH* Δ*dgcE* strains. Similar results were obtained even upon prolonged

incubation in the plate reader (Figure S4), confirming that the observed difference with the overnight flask culture was not because of the different growth stage but rather due to differences in growth conditions.

We further tested reporter activation under growth conditions that favor biofilm formation. During formation of static submerged biofilms in multi-well plates, where cultures are grown without shaking, the overall curli activation pattern in cell populations of individual or double and quadruple deletion strains (Figure 4B) was comparable to that in the flask culture grown in the orbital shaker (Figure 2B-D, F), although effects of DGC and PDE gene deletions on the fraction of curli-positive cells were apparently reduced. Curli gene activation in individual mutant strains correlated well with the levels of submerged biofilm formation (Figure S5), but the lack of regulation by c-di-GMP resulted in stronger reduction of the biofilm biomass than simply expected from its effect on curli gene expression, consistent with additional roles of c-di-GMP in biofilm formation.

All strains were also grown in the form of macrocolony biofilms on agar plates containing the salt-free LB medium, as done previously [26] (Figure 4C), as well as in the liquid salt-free LB medium as a control (Figure 4D). Expression of curli reporter in individual cells within the macrocolony was analyzed by flow cytometry immediately after colony resuspension (see Materials and Methods), so that the level of stable GFP reporter should reflect cell-specific curli expression within the macrocolony. The overall dependency of reporter activation on the deletions of individual DGC and PDE genes was again qualitatively similar to other growth conditions (Figure 4C). However, the fraction of curli-positive cells in the macrocolony biofilms was less sensitive to deletions of individual *dgc* or *pdh* genes than in planktonic cultures, regardless whether the latter were grown in TB (Figure 2C) or in the salt-free LB (Figure 4D). This agrees with stronger Congo red staining of macrocolonies of Δ*dgcE* and Δ*dgcM* strains compared to the Δ*mlrA* negative control (Figure S6). Thus, during macrocolony biofilm formation, regulation by c-di-GMP appears to be less important for the activation of curli gene expression, possibly because other regulators contribute to this activation in a highly structured environment. Importantly, the curli-positive cell fraction in both macrocolony and planktonic cultures of Δ*pdeR* Δ*dgcM* and Δ*pdeH* Δ*dgcE* Δ*pdeR* Δ*dgcM* strains growing on the salt-free LB was much higher than under other growth conditions.

To summarize, under all of the tested growth conditions, regulation by the known c-di-GMP regulatory network was not required for bimodal curli activation, although it indeed affected the fraction of curli-positive cells. Nevertheless, the absence of this c-di-GMP control apparently makes the fraction of curli-positive cells in the population more sensitive to environmental conditions, since this fraction exhibited much larger variation in the quadruple deletion compared to the wild-type strain among different growth conditions tested here.

Dynamics and variability of stochastic curli gene activation in individual cells is controlled by the c-di-GMP regulatory network

In order to investigate the dynamics of curli gene activation, and the impact of c-di-GMP regulation, at the single-cell level, we utilized a "mother machine" device — a microfluidic chip where growth of individual bacterial cell lineages could be followed in a highly parallelized manner over multiple generations [35] (Figure 5 and Supporting protocols). Since our design of the mother machine allows rapid switching of the medium supplied to the cells (Figure 5D), we could mimic nutrient depletion in order to activate curli expression in continuously growing single cells. For that, wild-type, individual DGC or PDE, or quadruple deletion strains were first loaded into the mother machine from overnight cultures, allowed to grow in fresh TB medium for several generations, and then shifted to the TB medium that was pre-conditioned by growing a batch culture (see Materials and Methods and Supporting protocols). A fraction of curli-positive cells was observed at the beginning of these experiments, which was strain-specific and consistent with the expected fraction of positive cells in the overnight cultures of each respective strain. After resuming exponential growth in the fresh medium all cells turned off curli expression (Figure 6A,C, Figure S7 and Movie S1, Movie S2 and Movie S3). Following the shift to conditioned medium, cell growth rate was strongly reduced (Figure 6E and Figure S8), and after several generations of slow growth, individual cells of all strains activated curli expression, while other cells remained in the curli-off state (Figure 6A,C,F, Figure S7 and Movie S1, Movie S2 and Movie S3). Importantly, the frequencies of curli-positive cells in individual mutants were consistent with those of the planktonic cultures, confirming that the overall control of curli gene expression by the c-di-GMP regulatory network is similar for growth in conditioned medium in the mother machine.

Although the overall frequency of curli reporter activation was comparable for the wild-type and $\Delta pdeH \Delta dgcE \Delta pdeR \Delta dgcM$ cells lacking c-di-GMP regulatory network, we further compared the dynamics of single-cell curli activation in both backgrounds. First, we observed that, in both strains, curli reporter activation was transient — and after several generations in the curli-on state individual cells turned curli expression off again during continuous growth in the conditioned medium (Figure 6B,D, Figure S8 and Figure S9). In some cases, the initial transient pulse of activation was even followed by a second activation event. Although such transient curli reporter activation was observed in both backgrounds, with a similar delay after exposure to conditioned medium (Figure 6H), its dynamics showed several important differences between strains. Most clearly, the rate of curli activation in individual cells was higher in the absence of the c-di-GMP-regulatory network (Figure 6G,I and Figure S10A,B), which led to stronger but more transient increase in their fluorescence (Figure 6J,K and Figure S10B). Additionally, the level of reporter induction showed greater intercellular heterogeneity in the $\Delta pdeH \Delta dgcE \Delta pdeR \Delta dgcM$ strain (Figure 6,D,J, Figure S9). Thus, the control of curli

expression by c-di-GMP reduces the rate but also the cell-to-cell variability of curli induction

within population.

Discussion

Expression of the curli biofilm matrix genes is known to be heterogeneous or even bistable in

communities of *E. coli* [25-28] and *S.* Typhimurium [29, 30], which might have important

functional consequences for the biomechanics of bacterial biofilms [27] and for stress

resistance and virulence of bacterial populations [29, 30]. In the well-structured macrocolony

biofilms, differentiation into subpopulations with different levels of curli matrix production is

largely deterministic and driven by gradients of nutrients and oxygen [18], but bimodality of

curli expression might also emerge stochastically, in a well-mixed population or between cells

within the same layer of a macrocolony [27, 29, 30]. How this bimodality originates within the

extremely complex regulatory network of curli genes [17, 19, 23, 27] remains a matter of

debate. Although earlier work proposed that bistable expression of curli might be a

consequence of positive transcriptional feedback in *csgD* regulation [30, 44], the most recently

suggested model attributed bistability to the properties of the c-di-GMP regulatory switch

formed by DgcM, PdeR and MlrA [27, 31]. Furthermore, although this bimodal pattern of curli

gene expression has typically been referred to as bistable, temporal dynamics and stability of

its induction in individual cells has not been directly investigated.

Here, we studied the expression of the major curli *csgBAC* operon in well-stirred planktonic *E.*

coli cultures as well as in submerged and macrocolony biofilms and during growth in a

microfluidics device. Consistent with previous studies [12, 14, 27], the induction of curli gene

expression in growing *E. coli* cell populations was observed during entry into the stationary

phase of growth. Curli activation was apparently dependent on depletion of nutrients, most

likely amino acids, since it could be suppressed by increasing the levels of nutrients or, more

specifically, by addition of serine, and conversely, it could be enhanced by the SHX-mediated

stimulation of the stringent response. We further observed that activation of the *csgBAC*

operon exhibited strongly pronounced bimodality under all tested conditions, resulting in two

distinct, curli-positive and -negative, subpopulations of cells even in a well-stirred planktonic

culture. Stochastic bimodality of curli reporter activation was confirmed in a microfluidic device,

where, following shift to conditioned medium, only a fraction of the cell population turned on

curli gene expression.

Such differentiation is apparently consistent with previous reports of bimodal *csgD* expression

in the stationary phase of *S.* Typhimurium [29, 30] and *E. coli* [27] culture growth. However, in

contrast with the previous interpretation of such a *csgD* expression pattern as bistability, we

observed that activation of curli gene expression during continuous cell growth in the

microfluidic device was only transient and unstable. Moreover, our data indicate that the

bimodality of the *csgBAC* expression is unlikely to be (fully) explained by the bimodal
expression of *csgD*. Firstly, compared to the bimodality of the *csgBAC* reporter, the previously
reported bimodality of *csgD* expression in *E. coli* [27] was much less pronounced and became
apparent only in later stages of planktonic culture growth. Moreover, in that study, all wild-type
cells showed activation of the *csgD* reporter, differing only in the level of this activation, which
itself cannot explain the on/off bimodality of the *csgBAC* reporter activation.

Secondly, and most importantly, whereas the bimodality of *csgD* expression was observed to
be dependent on regulation by the DgcE/PdeH/DgcM/PdeR network [27], this was clearly not
the case for *csgBAC* reporter expression. Under all tested conditions, including planktonic
cultures, submerged and macrocolony biofilms, and growth in the microfluidic device, the
differentiation into distinct subpopulations of positive and negative cells in our experiments
occurred in the absence of this entire c-di-GMP regulatory network. Since removal of the
DgcM/PdeR module further made curli gene expression insensitive to global levels of c-di-
GMP, contributions of other *E. coli* DGCs and PDEs to curli regulation in this deletion
background are unlikely, even if these enzymes might in principle affect curli gene expression
in wild-type *E. coli* cells via the common c-di-GMP pool. We therefore conclude that c-di-GMP
regulation might be dispensable for activation or bimodality of curli gene expression in *E. coli*,
although we could not rule out regulation of curli gene expression by a hypothetical local
DGC/PDE module that is insensitive to global c-di-GMP levels.

Nevertheless, c-di-GMP control clearly plays an important role during the establishment of
bimodality. Consistent with previous studies, we observed that the DGC and PDE proteins
determine the fraction of curli-positive cells [22, 23, 27], although they have no or only little
effect on the average level of *csgBAC* expression in individual cells. Our results further suggest
that the DgcE/PdeH/DgcM/PdeR regulatory network might make curli expression in *E. coli*
populations more robust, helping to ensure that the fraction of curli-positive cells is less
variable under different culture growth conditions. This might be related to the observed effect
of this network on the temporal expression dynamics in a continuously growing culture, with
faster but more heterogeneous activation of curli expression in the absence of the c-di-GMP
control, although whether and how there two different kinds of variability and their suppression
by are connected remains to be investigated. It is further possible that the increased
robustness provided by c-di-GMP regulation is due to the network-induced bimodality of *csgD*
expression discussed above [27], which might provide an additional stabilizing feedback.

How might the observed pulsatile activation of the curli-positive state originate at the single-
cell level and lead to the differentiation into distinct subpopulations in *E. coli* culture? Pulsing
in expression was proposed to be common to many gene regulatory circuits [45, 46, 47]. It was
recently described in *E. coli* for the upstream regulator of curli, σ^S [48], as well as for the
flagellar regulon [49, 50] that is anti-regulated with curli [28]. However, in neither of these cases

did pulsing lead to bimodality of expression, and their relation to the observed pulses in curli expression thus remains to be seen.

Regardless of the origin of pulsatile expression, the timing and duration of curli activation pulses can explain how such transient activation leads to differentiation into two distinct subpopulations in the planktonic culture. Since the curli expression is only turned on during transition to the stationary phase, individual cells stochastically activate curli genes just before the culture growth ceases, thus making subsequent reversion to the curli-negative state impossible. Although curli gene regulation in biofilms might be more complex, spatial stratification of gene expression in the macrocolony biofilms was proposed to resemble the temporal regulation of expression in planktonic cultures [18, 26]. Thus, also in this case, curli activation would happen in a cell layer just before it ceases to grow, fixing the differentiation between curli-positive and curli-negative cells within this colony layer.

Acknowledgements

We thank Sarah Hoch, Silvia González Sierra and Gabriele Malengo for technical help and advice.

References

1.   Arnqvist A, Olsen A, Pfeifer J, Russell DG, Normark S. The Crl protein activates cryptic genes for curli formation and fibronectin binding in *Escherichia coli* HB101. Mol Microbiol. 1992;6(17):2443-52. Epub 1992/09/01. doi: 10.1111/j.1365-2958.1992.tb01420.x. PubMed PMID: 1357528.

2.   Hufnagel DA, Depas WH, Chapman MR. The Biology of the *Escherichia coli* Extracellular Matrix. Microbiol Spectr. 2015;3(3). Epub 2015/07/18. doi: 10.1128/microbiolspec.MB-0014-2014. PubMed PMID: 26185090; PubMed Central PMCID: PMCPMC4507285.

3.   Larsen P, Nielsen JL, Dueholm MS, Wetzel R, Otzen D, Nielsen PH. Amyloid adhesins are abundant in natural biofilms. Environmental Microbiology. 2007. doi: 10.1111/j.1462-2920.2007.01418.x.

4.   Romling U, Bian Z, Hammar M, Sierralta WD, Normark S. Curli fibers are highly conserved between *Salmonella typhimurium* and *Escherichia coli* with respect to operon structure and regulation. J Bacteriol. 1998;180(3):722-31. Epub 1998/02/11. doi: 10.1128/JB.180.3.722-731.1998. PubMed PMID: 9457880; PubMed Central PMCID: PMCPMC106944.

5.   Vidal O, Longin R, Prigent-Combaret C, Dorel C, Hooreman M, Lejeune P. Isolation of an *Escherichia coli* K-12 mutant strain able to form biofilms on inert surfaces: involvement of a new ompR allele that increases curli expression. J Bacteriol. 1998;180(9):2442-9. Epub 1998/05/09. doi: 10.1128/JB.180.9.2442-2449.1998. PubMed PMID: 9573197; PubMed Central PMCID: PMCPMC107187.

6.   Zogaj X, Bokranz W, Nimtz M, Romling U. Production of cellulose and curli fimbriae by members of the family Enterobacteriaceae isolated from the human gastrointestinal tract.

Infect Immun. 2003;71(7):4151-8. Epub 2003/06/24. doi: 10.1128/IAI.71.7.4151-4158.2003.
PubMed PMID: 12819107; PubMed Central PMCID: PMCPMC162016.
7.   Tursi SA, Tukel C. Curli-Containing Enteric Biofilms Inside and Out: Matrix Composition,
Immune Recognition, and Disease Implications. Microbiol Mol Biol Rev. 2018;82(4). Epub
2018/10/12. doi: 10.1128/MMBR.00028-18. PubMed PMID: 30305312; PubMed Central
PMCID: PMCPMC6298610.
8.   Chapman MR, Robinson LS, Pinkner JS, Roth R, Heuser J, Hammar M, et al. Role of
Escherichia coli curli operons in directing amyloid fiber formation. Science.
2002;295(5556):851-5. Epub 2002/02/02. doi: 10.1126/science.1067484. PubMed PMID:
11823641; PubMed Central PMCID: PMCPMC2838482.
9.   Dueholm MS, Albertsen M, Otzen D, Nielsen PH. Curli Functional Amyloid Systems Are
Phylogenetically Widespread and Display Large Diversity in Operon and Protein Structure.
PLoS ONE. 2012. doi: 10.1371/journal.pone.0051274.
10.   Gerstel U, Römling U. The csgD promoter, a control unit for biofilm formation in
Salmonella typhimurium. Research in Microbiology2003.
11.   Brombacher E, Baratto A, Dorel C, Landini P. Gene expression regulation by the Curli
activator CsgD protein: modulation of cellulose biosynthesis and control of negative
determinants for microbial adhesion. J Bacteriol. 2006;188(6):2027-37. Epub 2006/03/04. doi:
10.1128/JB.188.6.2027-2037.2006. PubMed PMID: 16513732; PubMed Central PMCID:
PMCPMC1428138.
12.   Pesavento C, Becker G, Sommerfeldt N, Possling A, Tschowri N, Mehlis A, et al. Inverse
regulatory coordination of motility and curli-mediated adhesion in *Escherichia coli*. Genes Dev.
2008;22(17):2434-46. Epub 2008/09/04. doi: 10.1101/gad.475808. PubMed PMID: 18765794;
PubMed Central PMCID: PMCPMC2532929.
13.   Romling U, Rohde M, Olsen A, Normark S, Reinkoster J. AgfD, the checkpoint of
multicellular and aggregative behaviour in *Salmonella typhimurium* regulates at least two
independent pathways. Mol Microbiol. 2000;36(1):10-23. Epub 2000/04/12. doi:
10.1046/j.1365-2958.2000.01822.x. PubMed PMID: 10760159.
14.   Weber H, Pesavento C, Possling A, Tischendorf G, Hengge R. Cyclic-di-GMP-mediated
signalling within the sigma network of *Escherichia coli*. Mol Microbiol. 2006;62(4):1014-34.
Epub 2006/10/03. doi: 10.1111/j.1365-2958.2006.05440.x. PubMed PMID: 17010156.
15.   Barnhart MM, Chapman MR. Curli biogenesis and function. Annu Rev Microbiol.
2006;60:131-47. Epub 2006/05/18. doi: 10.1146/annurev.micro.60.080805.142106. PubMed
PMID: 16704339; PubMed Central PMCID: PMCPMC2838481.
16.   Ishihama A. Prokaryotic genome regulation: Multifactor promoters, multitarget
regulators and hierarchic networks. FEMS Microbiology Reviews2010.
17.   Rossi E, Cimdins A, Luthje P, Brauner A, Sjoling A, Landini P, et al. "It's a gut feeling" -
Escherichia coli biofilm formation in the gastrointestinal tract environment. Crit Rev Microbiol.
2018;44(1):1-30. Epub 2017/05/10. doi: 10.1080/1040841X.2017.1303660. PubMed PMID:
28485690.
18.   Serra DO, Hengge R. Bacterial Multicellularity: The Biology of *Escherichia coli* Building
Large-Scale Biofilm Communities. Annu Rev Microbiol. 2021;75:269-90. Epub 2021/08/04.
doi: 10.1146/annurev-micro-031921-055801. PubMed PMID: 34343018.
19.   Simm R, Ahmad I, Rhen M, Le Guyon S, Romling U. Regulation of biofilm formation in
Salmonella enterica serovar Typhimurium. Future Microbiol. 2014;9(11):1261-82. Epub
2014/12/02. doi: 10.2217/fmb.14.88. PubMed PMID: 25437188.

20.   Brown PK, Dozois CM, Nickerson CA, Zuppardo A, Terlonge J, Curtiss R. MIrA, a novel
regulator of curli (AgF) and extracellular matrix synthesis by *Escherichia coli* and *Salmonella*
enterica serovar Typhimurium. Molecular Microbiology. 2001. doi: 10.1046/j.1365-
2958.2001.02529.x.
21.   Ogasawara H, Yamada K, Kori A, Yamamoto K, Ishihama A. Regulation of the
Escherichia coli csgD promoter: Interplay between five transcription factors. Microbiology.
2010. doi: 10.1099/mic.0.039131-0.
22.   Lindenberg S, Klauck G, Pesavento C, Klauck E, Hengge R. The EAL domain protein YciR
acts as a trigger enzyme in a c-di-GMP signalling cascade in *E. coli* biofilm control. EMBO J.
2013;32(14):2001-14. Epub 2013/05/28. doi: 10.1038/emboj.2013.120. PubMed PMID:
23708798; PubMed Central PMCID: PMCPMC3715855.
23.   Hengge R. Linking bacterial growth, survival, and multicellularity - small signaling
molecules as triggers and drivers. Curr Opin Microbiol. 2020;55:57-66. Epub 2020/04/04. doi:
10.1016/j.mib.2020.02.007. PubMed PMID: 32244175.
24.   Sarenko O, Klauck G, Wilke FM, Pfiffer V, Richter AM, Herbst S, et al. More than
Enzymes That Make or Break Cyclic Di-GMP-Local Signaling in the Interactome of GGDEF/EAL
Domain Proteins of *Escherichia coli*. mBio. 2017;8(5). Epub 2017/10/12. doi:
10.1128/mBio.01639-17. PubMed PMID: 29018125; PubMed Central PMCID:
PMCPMC5635695.
25.   DePas WH, Hufnagel DA, Lee JS, Blanco LP, Bernstein HC, Fisher ST, et al. Iron induces
bimodal population development by *Escherichia coli*. Proc Natl Acad Sci U S A.
2013;110(7):2629-34. Epub 2013/01/30. doi: 10.1073/pnas.1218703110. PubMed PMID:
23359678; PubMed Central PMCID: PMCPMC3574911.
26.   Serra DO, Richter AM, Klauck G, Mika F, Hengge R. Microanatomy at cellular resolution
and spatial order of physiological differentiation in a bacterial biofilm. mBio.
2013;4(2):e00103-13. Epub 2013/03/21. doi: 10.1128/mBio.00103-13. PubMed PMID:
23512962; PubMed Central PMCID: PMCPMC3604763.
27.   Serra DO, Hengge R. A c-di-GMP-Based Switch Controls Local Heterogeneity of
Extracellular Matrix Synthesis which Is Crucial for Integrity and Morphogenesis of *Escherichia*
coli Macrocolony Biofilms. J Mol Biol. 2019;431(23):4775-93. Epub 2019/04/08. doi:
10.1016/j.jmb.2019.04.001. PubMed PMID: 30954572.
28.   Besharova O, Suchanek VM, Hartmann R, Drescher K, Sourjik V. Diversification of gene
expression during formation of static submerged biofilms by *Escherichia coli*. Frontiers in
Microbiology. 2016. doi: 10.3389/fmicb.2016.01568.
29.   Grantcharova N, Peters V, Monteiro C, Zakikhany K, Römling U. Bistable expression of
CsgD in biofilm development of *Salmonella enterica* serovar Typhimurium. Journal of
Bacteriology. 2010. doi: 10.1128/JB.01826-08.
30.   MacKenzie KD, Wang Y, Shivak DJ, Wong CS, Hoffman LJ, Lam S, et al. Bistable
expression of CsgD in *Salmonella enterica* serovar Typhimurium connects virulence to
persistence. Infect Immun. 2015;83(6):2312-26. Epub 2015/04/01. doi: 10.1128/IAI.00137-15.
PubMed PMID: 25824832; PubMed Central PMCID: PMCPMC4432751.
31.   Yousef KP, Streck A, Schütte C, Siebert H, Hengge R, von Kleist M. Logical-continuous
modelling of post-translationally regulated bistability of curli fiber expression in *Escherichia*
coli. BMC Systems Biology. 2015. doi: 10.1186/s12918-015-0183-x.
32.   Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, et al. Construction of
Escherichia coli K-12 in-frame, single-gene knockout mutants: the Keio collection. Mol Syst

Biol. 2006;2:2006 0008. Epub 2006/06/02. doi: 10.1038/msb4100050. PubMed PMID: 16738554; PubMed Central PMCID: PMCPMC1681482.

33. Cherepanov PP, Wackernagel W. Gene disruption in *Escherichia coli*: TcR and KmR cassettes with the option of Flp-catalyzed excision of the antibiotic-resistance determinant. Gene. 1995;158(1):9-14. Epub 1995/05/26. doi: 10.1016/0378-1119(95)00193-a. PubMed PMID: 7789817.

34. Amann E, Ochs B, Abel K-J. Tightly regulated tac promoter vectors useful for the expression of unfused and fused proteins in *Escherichia coli*. Gene. 1988;69(2):301-15. doi: https://doi.org/10.1016/0378-1119(88)90440-4.

35. Wang P, Robert L, Pelletier J, Dang WL, Taddei F, Wright A, et al. Robust growth of *Escherichia coli*. Curr Biol. 2010;20(12):1099-103. Epub 2010/06/12. doi: 10.1016/j.cub.2010.04.045. PubMed PMID: 20537537; PubMed Central PMCID: PMCPMC2902570.

36. Potrykus K, Cashel M. (p)ppGpp: still magical? Annu Rev Microbiol. 2008;62:35-51. Epub 2008/05/06. doi: 10.1146/annurev.micro.62.081307.162903. PubMed PMID: 18454629.

37. Varik V, Oliveira SRA, Hauryliuk V, Tenson T. HPLC-based quantification of bacterial housekeeping nucleotides and alarmone messengers ppGpp and pppGpp. Sci Rep. 2017;7(1):11022. Epub 2017/09/10. doi: 10.1038/s41598-017-10988-6. PubMed PMID: 28887466; PubMed Central PMCID: PMCPMC5591245.

38. Patacq C, Chaudet N, Letisse F. Crucial Role of ppGpp in the Resilience of *Escherichia coli* to Growth Disruption. mSphere. 2020;5(6). Epub 2020/12/29. doi: 10.1128/mSphere.01132-20. PubMed PMID: 33361126; PubMed Central PMCID: PMCPMC7763551.

39. Krasteva PV, Fong JC, Shikuma NJ, Beyhan S, Navarro MV, Yildiz FH, et al. *Vibrio cholerae* VpsT regulates matrix production and motility by directly sensing cyclic di-GMP. Science. 2010;327(5967):866-8. Epub 2010/02/13. doi: 10.1126/science.1181185. PubMed PMID: 20150502; PubMed Central PMCID: PMCPMC2828054.

40. Ahmad I, Cimdins A, Beske T, Romling U. Detailed analysis of c-di-GMP mediated regulation of csgD expression in *Salmonella typhimurium*. BMC Microbiol. 2017;17(1):27. Epub 2017/02/06. doi: 10.1186/s12866-017-0934-5. PubMed PMID: 28148244; PubMed Central PMCID: PMCPMC5289004.

41. Gualdi L, Tagliabue L, Landini P. Biofilm formation-gene expression relay system in *Escherichia coli*: modulation of sigmaS-dependent gene expression by the CsgD regulatory protein via sigmaS protein stabilization. J Bacteriol. 2007;189(22):8034-43. Epub 2007/09/18. doi: 10.1128/JB.00900-07. PubMed PMID: 17873038; PubMed Central PMCID: PMCPMC2168689.

42. Levine JH, Lin Y, Elowitz MB. Functional roles of pulsing in genetic circuits. Science. 2013;342(6163):1193-200. Epub 2013/12/07. doi: 10.1126/science.1239999. PubMed PMID: 24311681; PubMed Central PMCID: PMCPMC4100686.

43. Locke JC, Young JW, Fontes M, Hernandez Jimenez MJ, Elowitz MB. Stochastic pulse regulation in bacterial stress response. Science. 2011;334(6054):366-9. Epub 2011/10/08. doi: 10.1126/science.1208144. PubMed PMID: 21979936; PubMed Central PMCID: PMCPMC4100694.

44. Levine JH, Fontes ME, Dworkin J, Elowitz MB. Pulsed feedback defers cellular differentiation. PLoS Biol. 2012;10(1):e1001252. Epub 2012/02/04. doi: 10.1371/journal.pbio.1001252. PubMed PMID: 22303282; PubMed Central PMCID: PMCPMC3269414.

45.   Patange O, Schwall C, Jones M, Villava C, Griffith DA, Phillips A, et al. *Escherichia coli*
can survive stress by noisy growth modulation. Nat Commun. 2018;9(1):5333. Epub
2018/12/19. doi: 10.1038/s41467-018-07702-z. PubMed PMID: 30559445; PubMed Central
PMCID: PMCPMC6297224.
46.   Kim JM, Garcia-Alcala M, Balleza E, Cluzel P. Stochastic transcriptional pulses
orchestrate flagellar biosynthesis in *Escherichia coli*. Sci Adv. 2020;6(6):eaax0947. Epub
2020/02/23. doi: 10.1126/sciadv.aax0947. PubMed PMID: 32076637; PubMed Central
PMCID: PMCPMC7002133.
47.   Sassi AS, Garcia-Alcala M, Kim MJ, Cluzel P, Tu Y. Filtering input fluctuations in intensity
and in time underlies stochastic transcriptional pulses without feedback. Proc Natl Acad Sci U
S A. 2020;117(43):26608-15. Epub 2020/10/14. doi: 10.1073/pnas.2010849117. PubMed
PMID: 33046652; PubMed Central PMCID: PMCPMC7604450.
598

Figure 1: Bimodal activation of curli gene expression in *E. coli* planktonic cultures. *E. coli* cells carrying genomic transcriptional reporter of *csgBAC* operon were grown in liquid tryptone broth (TB) medium at 30 °C under constant shaking. **(A)** Optical density (OD_{600}) and relative fluorescence (fluorescence/OD_{600}; AU, arbitrary units) of the culture during growth in a plate reader, starting from two different dilutions of the overnight culture. Error bars indicate standard error of the mean (SEM) of 10 technical replicates. **(B)** Distribution of single-cell fluorescence levels, measured by flow cytometry, of cultures grown from overnight culture diluted 1:1000 to indicated OD_{600} or overnight (ON; 25 h) in flasks in an orbital shaker.

Figure 2: Regulation of curli gene expression by c-di-GMP. (A) Current model of regulation of curli

gene expression by c-di-GMP in *E. coli*, adapted from [27]. The regulation is mediated by two pairs of

diguanylate cyclases (DGCs; blue) and phosphodiesterases (PDEs; orange). PdeH and DgcE control

global levels of c-di-GMP, whereas PdeR and DgcM mediate local c-di-GMP-dependent regulation of

curli gene expression by controlling activity of transcription factor MlrA, which activates another curli-

specific transcription factor CsgD. **(B-F)** Flow cytometry measurements of curli gene expression in *E.*

coli planktonic cultures grown in TB overnight in flasks in an orbital shaker, shown for the wild-type (WT)

and Δ*mlrA* strain (B), and individual (C), double (D), triple (E) and quadruple (F) deletions of DGC or

PDE enzymes, as indicated. Fraction of positive cells in the population (mean of three biological

replicates ± SEM) is indicated for each strain. Note that the scale in the *y* axes is different for individual

strains to improve readability.

Figure 3: Decoupling of curli gene expression from c-di-GMP regulation in the absence of PdeR/ DgcM regulatory module. *E. coli* wild-type (WT) cells or cells lacking c-di-GMP regulatory enzymes (Δ*pdeH* Δ*dgcE* Δ*pdeR* Δ*dgcM*) were transformed with either empty pTrc99a plasmid (control) or with pTrc99a plasmid carrying *dgcE* (pTrc99a::dgcE) or *pdeH* (pTrc99a::pdeH) genes. Expression from the vector was induced with 1 µM IPTG. Bacteria were grown in TB overnight in flasks with shaking and cultures were subjected to the flow cytometry analysis. Fraction of positive cells in the population (mean of three biological replicates ± SEM) is indicated for each strain.

Figure 4: Dependence of curli gene expression on c-di-GMP regulation under different growth conditions. (A-D) Flow cytometry measurements of curli expression in the wildtype (WT) and indicated gene deletion strains either after 24 h of growth in liquid TB in a plate reader (A) or as submerged biofilms without agitation (B), as macrocolony biofilms on salt-free LB agar (C), or as overnight planktonic cultures in salt-free LB in flasks in an orbital shaker. Fraction of positive cells in the population (mean of three biological replicates ± SEM) is indicated for each strain.

Figure 5: Design and operation of the microfluidic mother machine device. (A) Schematic overview

of the chip layout, featuring four independent supply channels (green) for cell inoculation and media

supply. **(B)** Detailed view of the area marked by a rectangle in (A), showing the switching junction and

a part of the mother machine cultivation sites (blue). The junction is formed by two inlets, leading to one

central supply channel. Control of pressure at each inlet allows for prioritization of one medium over the

other through the supply channel, and, ultimately, the mother machine cultivation sites. Residual

medium flows out through waste channels located to each side of the central supply channel. Medium

flowing through the supply channel exits the chip through one outlet. **(C)** Detailed view of the area

marked in **(B)** by a rectangle, showing the mother machine cultivation sites. Each of the four channels

contains 57 mother machine cultivation sites, each of which contains 30 mother machine traps with

widths of either 0.9, 1, or 1.1 µm. The mother machine traps feature a 0.3 µm wide constriction on the

bottom, preventing the mother cell from exiting the trap while allowing perfusion of the medium. The

supply channels (green) are 8 µm in depth, the mother machine traps (blue) are 0.8 µm in depth. **(D)**

On-chip medium switching visualized by merged phase contrast and mCherry images of the channel

junction. Media are supplied through separate inlets (top and bottom), which are separated in the center

of the channel by a PDMS barrier. The direction of flow is indicated by white arrows. Water was supplied

through the top inlet, while a 0.2 µM sulforhodamine B solution was supplied through the bottom inlet,

visualizing the flow pattern in the junction. The pressure at the top inlet was kept constant at 200 mbar.

Depending on the pressure set at the bottom inlet, it is possible to select which one of the two media

flows into the central supply channel to the mother machine growth sites.

Figure 6: Impact of c-di-GMP regulation on dynamics of curli gene induction in individual cells.
E. coli cells in a microfluidic device (mother machine) were shifted from a fresh to conditioned TB medium after 4 h of growth to induce curli expression. **(A-D)** Examples of image time series and single-cell fluorescence traces for the wild-type (WT) (A,B) and for the Δ*pdeH* Δ*dgcE* Δ*pdeR* Δ*dgcM* strain disabled in the c-di-GMP regulation (C,D), growing in a mother machine in one experiment. Expression of the curli reporter is indicated by the green overlay on the phase contrast image. **(E)** Median instantaneous growth rate — fold rate of change in length — of cells grown in microfluidics experiments as described in (A). The switch to conditioned medium is at time zero, as indicated. Shaded area is the interquartile range. The number of cells in the device varies with time, but is on average n = 296 for WT and n = 522 for the quadruple deletion strain. **(F)** Distributions of curli expression at different time points

in the microfluidics experiment. Shown are kernel density estimates of curli expression in WT and in the

quadruple deletion strain at selected time points. **(G)** Median curli expression profile for single-cell traces

aligned by the time at which they exceed a threshold of 10^3 fluorescence units. Shaded area is

interquartile range; n = 128 for WT and 230 for the quadruple deletion strain. **(H-K)** Distributions of curli

induction parameters for the wild-type and for the c-di-GMP-regulation disabled strain, in two

independent experiments, r1 (same experiment as in A-G) and r2. Shown are histograms of the times

at which a threshold of 10^3 fluorescence units were crossed **(H)**, the maximum rates of increase in

fluorescence **(I)**, the fluorescence amplitudes at the first peak **(J)** and the half-time of fluorescence decay

after reaching the peak **(K)** for cell traces from both microfluidics experiments. **$p < 10^{-2}$ in an unpaired

two-sample t-test assuming normal distributions with unknown and unequal variances. ns, non-

significant.

Supporting information

A

Figure S1: Dependence of curli gene expression on nutrient levels. Wild-type *E. coli* cultures were grown as in Figure 1A but with different indicated concentrations of TB. **(A)** Bacterial growth and activity of transcriptional curli reporter. Error bars indicate SEM of 6 technical replicates. **(B)** Distribution of single-cell fluorescence levels after 24 h of growth in a plate reader measured by flow cytometry. Note that the scale in the *y* axes is different for individual conditions to improve readability.

Figure S2: Stimulation of curli gene expression by stringent response. Wild-type *E. coli* cultures

were grown as in Figure 1A but with addition of either indicated concentrations of serine hydroxamate

(SHX) at inoculation point or 1 mM serine after 6 h of growth. **(A)** Bacterial growth and activity of

transcriptional curli reporter. Error bars indicate SEM of 6 technical replicates. **(B)** Distribution of single-

cell fluorescence levels after 24 h of growth in a plate reader measured by flow cytometry.

Figure S3: Relative fluorescence of curli reporter in planktonic culture grown in TB medium in a

plate reader. Error bars indicate SEM of 5 technical replicates. *E. coli* planktonic cultures of the wild-

type (WT), Δ*mlrA* strain, and individual, double and quadruple deletions of DGC or PDE enzymes were

grown in TB in a plate reader.

Figure S4: Curli gene expression upon prolonged cultivation in a plate reader. Wild-type (WT) and
ΔpdeH ΔdgcE ΔpdeR ΔdgcM strains were grown in TB in a plate reader as in Figure S3, but during
36 h. **(A)** Induction of transcriptional curli reporter. Error bars indicate SEM of 10 technical replicates.
(B) Distribution of single-cell fluorescence levels in populations of both strains after 36 h of growth
measured by flow cytometry.

Figure S5: Curli gene expression in submerged biofilm cultures. Biofilm formation by indicated

strains grown in multi-well plates in TB medium, quantified using crystal violet (CV) staining. Error bars

indicate SEM of 3 independent replicates.

Figure S6: Curli expression in macrocolony biofilms. Images of macrocolonies of indicated strains

after 8 days of growth on salt-free LB agar plates.

Figure S7: Distributions of curli expression at different time points in the microfluidics
experiment. Shown are kernel density estimates of curli expression in the wild type WT, individual and
quadruple deletions of DGC or PDE enzymes at selected time points. Cells in stationary phase were
loaded into mother machine chip supplied with fresh medium, and then switched to the conditioned
media at the ''0 hours'' time point. Note that the scale in the *y* axes is different for individual conditions
to improve readability.

Figure S8: Growth rates and fraction of curli expressing cells over time. Stationary phase cells

were introduced into mother machine devices, supplied with fresh medium and then switched to

conditioned medium after 4 h of growth, as in Figure 6. **(A)** Median instantaneous growth rates for the

wild-type and for the Δ*pdeH* Δ*dgcE* Δ*pdeR* Δ*dgcM* strain disabled in c-di-GMP regulation. Growth rate

drops rapidly and cells switch on curli expression after a switch to conditioned medium. Shaded area is

interquartile range. **(B)** Fraction of cells with fluorescence exceeding 1000 units. **(C)** Number of detected

cells. Two biological replicates (r1 and r2) were performed for each strain; data for the r1 replicate are

also shown in Figure 6. Note that in the r2 experiment for the quadruple deletion strain, cells were only

imaged after medium switching.

Figure S9: Single-cell traces of cell fluorescence for all cells for the wild-type and for the c-di-
GMP-regulation disabled strain. Data are from the same biological replicates (r1 and r2) as in Figure
S8.

737 Figure S10: The rate and variability of curli induction for the wild-type and for the c-di-GMP-
738 regulation disabled strain. (A) Median curli expression, **(B)** production rate, and **(C)** number of cells

739 for traces from both microfluidics experiments (r1 and r2) aligned by the time at which they exceeded a

740 threshold of 10^3 fluorescence units. Shaded area is interquartile range. Compared to the WT, the rate

741 of curli induction is faster but traces show more variability in a mutant without the global or local c-di-

742 GMP regulatory modules.

744 **Table S1. *E. coli* strains and plasmids used in this study.**

Strains	Relevant genotype	Reference
W3110	W3110 derivative with functional RpoS	[1]
VS1146	W3110 *csgA*::*csgA*_RBS_*sfgfp*	[2]
VS1857	VS1146 Δ*mlrA*	This work
VS1732	VS1146 Δ*pdeH*	This work
VS1720	VS1146 Δ*dgcE*	This work
VS1258	VS1146 Δ*pdeR*	This work
VS1257	VS1146 Δ*dgcM*	This work
VS1717	VS1146 Δ*pdeH* Δ*dgcE*	This work
VS1713	VS1146 Δ*pdeR* Δ*dgcM*	This work
VS1885	VS1146 Δ*pdeH* Δ*dgcE* Δ*pdeR*	This work
VS1729	VS1146 Δ*pdeH* Δ*dgcE* Δ*pdeR* Δ*dgcM*	This work
Plasmids		
pTrc99a	Expression vector; *Ptrc* promoter inducible by isopropyl-β-D-thiogalactopyranoside (IPTG); pBR ori; Ap^R	[3]
pVS2689	pTrc99a::*pdeH*	This work
pVS1644	pTrc99a::*dgcE*	This work

1. Serra DO, Richter AM, Klauck G, Mika F, Hengge R. Microanatomy at cellular resolution and
spatial order of physiological differentiation in a bacterial biofilm. mBio. 2013;4(2):e00103-13. Epub
2013/03/21. doi: 10.1128/mBio.00103-13. PubMed PMID: 23512962; PubMed Central PMCID:
PMCPMC3604763.
2. Besharova O, Suchanek VM, Hartmann R, Drescher K, Sourjik V. Diversification of gene
expression during formation of static submerged biofilms by *Escherichia coli*. Frontiers in Microbiology.
2016. doi: 10.3389/fmicb.2016.01568.
3. Amann E, Ochs B, Abel K-J. Tightly regulated tac promoter vectors useful for the expression of
unfused and fused proteins in *Escherichia coli*. Gene. 1988;69(2):301-15. doi:
https://doi.org/10.1016/0378-1119(88)90440-4.

Supporting protocols

Design and fabrication of the mother machine. A microfluidic device was used for time-resolved studies of *E. coli* growth and curli expression, which features mother machines traps for the observation of one-dimensional cell growth. The design of the microfluidic device is shown in Figure 5. The channel structure of the microfluidic chip features four independent channels with a depth of 8 µm, allowing the realization of four conditions, e.g. different media compositions, in one experiment. Each channel contains two inlets enabling on-chip switching between two different media, which are supplied at defined pressures. By controlling the pressure ratio between the two inlets, the medium supplied at the higher pressure flows through the central channel following the junction, while the other medium is pushed away from the junction and flows out through the waste channel (Figure 5D). The medium which flows through the central channel reaches the mother machine cultivation sites before exiting the chip through a single outlet.

The device was produced using a two-layer soft lithography method as described previously [1]. Based on our in-house made design of the channel layout, a 100 mm silicon wafer was produced by e-beam lithography (ConScience, Sweden). The wafer contains the channel layout as a positive relief. The mother machine traps, which are shown in blue in Figure 5A-C, were structured by etching the wafer by 0.8 µm, giving the mother machine the appropriate vertical dimension for cell trapping. The supply channels, which are shown in green in Figure 5A-C, are implemented as photoresist structures with a height of 8 µm on the wafer. The wafer served as a master mold for liquid polydimethylsiloxane (Sylgard 184 PDMS, VwR International GmbH, Germany), which was mixed at volumetric ratio of 7:1 with a cross-linking agent, degassed in a desiccator for 30 min and poured over the wafer to a height of approximately 4 mm and thermally cured at 80 °C overnight. The cured PDMS was peeled off from the wafer and manually cut into separate chips. Inlet and outlet holes were punched with a 0.75 mm punching tool (Robbins True-Cut Disposable Biopsy Punch 0.75mm with Plunger, Robbins Instruments, USA). The surface of the chip was cleaned by a rinse with isopropanol and the application of adhesive tape (tesafilm, Germany) prior to bonding. The chip was irreversibly bonded to a glass substrate by applying oxygen plasma to both chip and glass surfaces (Diener Femto, Diener GmbH, Germany) and bringing the treated surfaces together. The bond was strengthened by storing the bonded device in the oven at 80 °C for 2 min. The device was mounted on an inverted fluorescence microscope (Nikon Eclipse Ti, Nikon Corporation, Japan) equipped with an incubator. The microscope setup included an Andor Zyla 4.2 sCMOS camera (Oxford Instruments, UK), an objective with 100x magnification (Plan Apochromat λ Oil, NA=1.45, WD=170 µm; Nikon, Japan) and a perfect focus system (Nikon Corporation, Japan) for focus drift compensation.

Growth experiments. *E. coli* cells were allowed to grow in TB medium until the stationary phase prior to inoculation into the device. Conditioned medium was prepared by cultivating the wild-type *E. coli* cells in TB medium for 20 h (batch culture), after which the cell suspension was centrifuged at 4000 rpm for 10 min. The medium then was filter sterilized and stored at 4°C. The mother machine growth sites were loaded with the undiluted cell suspension by manual infusion of the cell suspension through one of the two inlets using a 1-ml syringe. The connection between the syringe and the chip was realized by tygon tubing (Tygon S-54-HL, inner diameter = 0.51 mm, outer diameter = 1.52 mm, VwR International GmbH, Germany) in combination with blunt dispensing needles (general purpose tips, inner diameter = 0.41 mm, outer diameter = 0.72 mm, Nordson EFD, USA). Medium flow was controlled by programmable pressure regulators (LineUP FlowEZ, FLUIGENT, France), which generated flow by applying pressure on 50-ml medium reservoirs (P-CAP series, FLUIGENT, France). Fresh and conditioned TB medium were respectively filled in separate reservoirs, and each one was pressurized by one module of the pressure regulator. After the cell inoculation both media were connected to the inlets of the channel via tygon tubing and blunt dispensing needles. The pressure at the inlet of the fresh TB medium was set to 200 mbar and remained constant throughout the experiment. During the selection of the positions for imaging the pressure at the inlet of the conditioned medium was set to 250 mbar, allowing the conditioned medium to flow through the junction to the mother machine growth sites and thereby maintaining the stationary state of the cells. At the beginning of imaging, the pressure at the inlet of the conditioned medium was reduced to 150 mbar and programmed to increase back to 250 mbar after 4 h of on-chip cultivation, thereby activating a medium switch from fresh to conditioned TB medium. Phase contrast and GFP fluorescence images were acquired with a time interval of 10 min.

Analysis of microfluidics data. Cells were segmented from phase contrast images by making use of a fully convolutional neural network based on the U-net architecture [2]. A set of manually curated cell outlines was prepared for training (1105 outlines) and validation (346 outlines) of the network. The training set was augmented by scaling, rotation, flipping and addition of white noise. A U-net of depth three, with 8, 16 and 32 filters along the contracting path, was trained to predict cell interiors from phase contrast images. Phase contrast images were normalized by subtracting the median and scaling to intensities expected between the 2^{nd} and 98^{th} percentiles. Cell interiors were defined from the curated outlines by filling each outline and then subjecting it to two rounds of morphological erosion. The erosion step ensured that neighbouring cells predicted by the network were well separated, such that distinct cell instances could be clearly identified simply by thresholding the prediction and labelling

connected regions. After instance identification, two rounds of morphological dilation restored
each mask to its original size. Finally, a smooth outline for each cell was obtained as a two-
dimensional spline defined by equidistant knots placed on the mask edge.

To track cells between time points, we applied a length conservation strategy for the cells along
each trench. At each time point, we ordered cell outlines by their depth in the trench, with
deepest cells first. We then attempted to match, in order, a cell outline in time point t with one
or more cell outlines in time point $t+1$, chosen such that the sum of their cell lengths would be
conserved within some threshold tolerance. In the trivial case, the length of the first outline at
time point t would match that of the first outline at time point $t+1$. In the event of cell division,
the cell length at time point t would match the sum of the first two cell lengths at time point $t+1$.
Since cells may grow in length between time points, we also initialised a growth rate parameter
for each cell that biased the expected cell lengths for time point $t+1$ as a fold-increase in length.
To enable adaptation to the true growth rate, the growth rate parameter was updated by a
lagging average over 20 time points. To increase robustness to errors in segmentation, we
additionally allowed state transitions from one to many and many to one, and built a proposal
tree, which branched for all valid assignments lying within the length thresholds. We searched
for the proposal with the lowest average fold-change in matched lengths, but limited branching
by retaining only the 10 best proposals for subsequent nodes (cell outlines) in the tree. The
length thresholds were deliberately set loosely such that the (sum of) cell length(s) at $t+1$ could
decrease at most five-fold or increase at most two-fold relative to the (sum of) cell length(s) at
t. This increased the number of valid proposals, but was important in cases where the growth
rate estimate was poor. For transitions where one cell outline split into more than two, or
transitions where multiple outlines merged into one cell, a new label was generated for the
corresponding cells at $t+1$. We made one exception to this labelling strategy to account for
occasional ambiguity in segmentation near division events, where a cell segmented as two
sister cells could later be segmented as a single mother cell. Specifically, when two sister cells
— i.e., cells that were previously involved in a division event — merged into one, the label was
set back to that of the mother; at the next division event, the labels of the sister cells were also
retained. Finally, note that any outlines below a minimum size threshold of 50 pixels were
ignored. All errors in tracking were manually curated.

Cell length was estimated from cell regions as the 'major axis length' of the Matlab regionprops
function — the major axis of the ellipse with same normalised second central moment as the
region. Instantaneous growth rates were estimated from the derivative of a smoothing spline
fitted to the logarithm of cell length over each cell division cycle. Knots for the spline were
placed at intervals of at most 15 time points. Single-cell fluorescence traces were quantified
from the median fluorescence within each outline. Background fluorescence varied as a
function of time due to the accumulation of cells at some trench exits, so we corrected for

background fluorescence in each trench at each time point using the median value of all non-
cell pixels. Fluorescence traces were characterised along branching lineages and were
smoothed with a Savitzky-Golay filter of order 3 and window length 21. The derivative of the
filter was used to obtain the maximum rate of fluorescence increase. Peaks in fluorescence
were identified as points for which the second derivative was less than -100 and either the first
derivative was zero, or its sign changed at the next time point. In cases where multiple
descendants shared a common peak event before branching, we counted that event only once.
The half-life of decay after a peak in fluorescence was determined by fitting an exponential
decay function with offset term to fluorescence data after the peak. Given the occasional
occurrence of a second peak in fluorescence, the decay function was only fit to time points
after the peak up to the point where the first derivative exceeded zero.
879

880 1.   Qin D, Xia Y, Whitesides GM. Soft lithography for micro- and nanoscale patterning. Nat
881 Protoc. 2010;5(3):491-502. Epub 2010/03/06. doi: 10.1038/nprot.2009.234. PubMed PMID:
882 20203666.
883 2.   Ronneberger O, Fischer P, Brox T. U-net: Convolutional networks for biomedical image
884 segmentation. arXiv. 2015;[1505.04597].
885