Three-way relationships between gut microbiota, helminth assemblages and bacterial

infections in wild rodent populations

Marie Bouilloud[1,2], Maxime Galan[1], Adélaïde Dubois[1], Christophe Diagne[3], Philippe

Marianneau[4], Benjamin Roche*[2], Nathalie Charbonnel*[1*]

[1]CBGP, INRAE, CIRAD, Institut Agro, IRD, University of Montpellier, Montpellier, France

[2]MIVEGEC, IRD, CNRS, University of Montpellier, Montpellier, France

[3]CBGP, IRD, INRAE, CIRAD, Institut Agro, University of Montpellier, Montpellier, France

[4]INRAE, Lyon, France

[*]Corresponding author

[*]Equal contribution of the coauthors

Orcid

Marie Bouilloud : marie.bouilloud@ird.fr, 0000-0001-8740-3205
17

18   Maxime Galan : maxime.galan@inrae.fr, **0000-0001-6981-5732**

20   Christophe Diagne : christophe.diagne@ird.fr, **0000-0002-6406-1270**

22   Philippe Marianneau : philippe.marianneau@inrae.fr

24   Benjamin Roche : roche.ben@gmail.com, **0000-0001-7975-4232**

26   Nathalie Charbonnel : nathalie.charbonnel@inrae.fr**, 0000-0002-7907-6539**

Abstract

30   **Abstract**

Background

Despite its central role in host fitness, the gut microbiota may differ greatly between

individuals. This variability is often mediated by environmental or host factors such as diet,

genetics, and infections. Recently, a particular attention has been given to the interactions

between gut bacteriota and helminths, as these latter could affect host susceptibility to other

infections. Further studies are still required to better understand the three-way interactions

between gut bacteriota, helminths and other parasites, especially because previous findings

have been very variable, even for comparable host-parasite systems.

Methods

In our study, we used the V4 region of the 16S rRNA gene to assess the variability of gut

bacteriota diversity and composition in wild populations of a small mammal, the bank vole

Myodes glareolus. Four sites were sampled at a regional geographical scale (100 km) along a

North-South transect in Eastern France. We applied analyses of community and microbial

ecology to evaluate the interactions between the gut bacteriota, the gastro-intestinal helminths

and the pathogenic bacteria detected in the spleen.

Results

We identified important variations of the gut bacteriota composition and diversity among

bank voles. They were mainly explained by sampling localities and reflected the North/South

sampling transect. In addition, we detected two main enterotypes, that might correspond to

contrasted diets. We found geographic variations of the Firmicutes/Bacteroidetes ratio, that

correlated positively with body mass index. We found positive correlations between the

specific richness of the gut bacteriota and of the helminth community, as well as between the

composition of these two communities, even when accounting for the influence of

geographical distance. The helminths *Aonchotheca murissylvatici, Heligmosomum mixtum*

and the bacteria *Bartonella* sp were the main taxa associated with the whole gut bacteriota

composition. Besides, changes in relative abundance of particular gut bacteriota taxa were

specifically associated with other helminths (*Mastophorus muris*, *Catenotaenia henttoneni*,

Paranoplocephala omphalodes and *Trichuris arvicolae*) or pathogenic bacteria. Especially,

infections with *Neoehrlichia mikurensis*, *Orientia* sp, *Rickettsia* sp and *P. omphalodes* were

associated with lower relative abundance of the family Erysipelotrichaceae (Firmicutes),

while coinfections with higher number of bacterial infections were associated with lower

relative abundance of a Bacteroidales family (Bacteroidetes).

Conclusions

These results emphasize complex interlinkages between gut bacteriota and infections in wild

animal populations. They remain difficult to generalize due to the strong impact of

environment on these interactions, even at regional geographical scales. Abiotic features, as

well as small mammal community composition and within host parasite coinfections, should

now be considered to better understand the spatial variations observed in the relationships

between gut bacteriota, gastro-intestinal helminths and bacterial infections.

Keywords: bank voles, co-infections, interactions, microbial community ecology, zoonoses

Introduction

Vertebrate gut microbiota plays key roles in host fitness through functions including nutrient acquisition, immunity and defence against infectious exogenous agents (hereafter called 'parasites' and including micro- and macroparasites) or proliferating indigenous organisms (Belkaid & Hand, 2014; Kamada *et al.*, 2013; Round & Mazmanian, 2009) among others. Nonetheless, the gut microbiota may differ greatly in natural environments between individuals, populations and species (Vujkovic-Cvijin *et al.*, 2020). Its composition is even subject to high temporal variation for a given individual.

These variations are shaped by ecological and/or evolutionary processes, among which stochasticity, migration and/or adaptive differences in microbes (McDonald *et al.*, 2020; Kolodny & Schulenburg, 2020). They are mediated by environmental features (e.g. acquisition of microorganisms from the environment, potentially through diet Ley *et al.*, 2008; Moran *et al.*, 2019), host factors (notably phylogeny, genetics or vertical transmission from mother to offspring) and interactions between hosts and their environment across space and time. For example, disruption of host-associated gut microbiota (termed "dysbiosis") may occur as a result of environmental change and stress affecting the host. This has been shown in the context of anthropogenic pressures (e.g. chemical exposures, Rosenfeld, 2017; urbanisation, Stothart *et al.*, 2019) or parasite infections (Trevelline *et al.*, 2019).

Understanding the relationships between gut microbiota and parasites is crucial regarding their potential impacts on human and animal health (Clemente *et al.*, 2012). Among the numerous studies of vertebrate microbiota, some of them have put an emphasis on the gut bacterial microbiota (called hereafter 'gut bacteriota') and their interactions with gastro-intestinal helminth parasites. On one hand, the gut bacteriota may act as an innate immune barrier to intestinal infections and influence the local colonisation and growth of eukaryotic parasites, including helminths, through competitive metabolic interactions or induction of host immune responses (Leung *et al.*, 2018). On the other hand, helminth infections may also directly or indirectly affect the composition of the gut bacteriota via physical contact, competition for resources or host immunoregulation (see Kreisinger *et al.*, 2015).

Interactions between helminths and the gut bacteriota may be positive or negative (Loke & Lim, 2015). They may lead to potentially local but also systemic physiological changes affecting host health. For example, helminth infections can lead to malnutrition and weight loss through the dysfunction of microbial metabolism that could result from negative impacts on fermentative gut bacteria (Leung *et al.*, 2018). Besides, some helminth infections promote

higher abundance of gut bacteria that produce short-chain fatty acids from dietary fiber (Zaiss
et al., 2015). These metabolites circulate throughout the body and are important regulators of
host physiology (glucose and fat metabolism) and immune system (Honda & Littman, 2016;
Kim, 2021). Interactions between these helminths and gut bacteria may here increase the host
anti-inflammatory and regulatory T cell suppressor responses, what may in turn affect host
susceptibility to other infections as well as the outcomes of infections (Glendinning *et al.*,
2014).
The gut microbiota may also influence microparasite infections through their immune
function against local pathogenic bacteria colonization and their role in maintaining the
intestinal epithelium integrity (Khosravi & Mazmanian, 2013). There is also strong evidence
for systemic interactions between the gut microbiota and extra-intestinal
microbiota communities, at least in laboratory mice (e.g. Rosshart *et al.*, 2017;). This
systemic impact of gut microbiota is mediated by host immunity (Zheng et al., 2020). As
such, the gut microbiota produces metabolites (e.g., bacteriocins, short-chain fatty acids,
microbial amino-acids) that translocate from the intestinal lumen to various organs (e.g., liver,
brain, lung) through the circulatory system. This may induce tissue-specific immune
responses, and affect the host's susceptibility/resistance to (non enteric) pathogens (Winckler
& Thackray, 2019; Pfeiffer & Sonnenburg, 2011). Most of these studies have focused on
viruses (e.g., influenza A, coronaviruses, Karst & Wobus, 2019) and not yet on pathogenic
bacteria (but see Rolhion & Chassaing, 2015). The systemic impact of gut bacteriota on
microparasite infections still represents a fundamental knowledge gap in wild animals (Pascoe
et al., 2017).
The three-way interactions between host's gut bacteriota, gastro-intestinal helminths and
microparasites have been scarcely investigated in a single system, despite clearly becoming
pivotal in disease ecology. Yet, the growing interest on gut bacteriota/parasitism relationships
in recent literature (P. T. Johnson *et al.*, 2015) highlights the critical need for further empirical
works. One main reason is the relatively low concordance of findings between previous
studies – even for comparable host-parasite systems (e.g. for *Trichuris* sp and the gut
microbiota, see Cortes et al., 2019; Lawson et al., 2021). Up to now, most of the research on
this topic have been conducted on model species in laboratory settings. Although experiments
under controlled conditions may help deciphering the mechanisms underlying these
interactions between gut bacteriota and parasites in vertebrates (Pascoe *et al.*, 2017), they also
have inherent limitations. On the one hand, they only included a restricted number of targeted

parasites (usually helminths and/or microparasites). Consequently, they often omitted the potential effects of species interactions between and within parasite communities at the intra-host level (Telfer *et al.*, 2010). Co-infections by helminths species have been noticed by parasitologists for decades (Montgomery & Montgomery, 1989; Haukisalmi & Henttonen, 1993). Yet, co-infections between highly divergent micro- and macroparasites are also recognized to be the rule in most hosts in natural environments. Indeed, wild animals may carry simultaneously a large number of bacteria, helminths, viruses (Hoarau *et al.*, 2020). On the other hand, they are unable to include – and then capture the complexity of – the environmental conditions as drivers of the composition of gut bacteriota and of the exposure or sensibility to these latter (Adair & Douglas, 2017). From there, studies in natural contexts deserve strong consideration because these environmental factors may impact deeply the relationships between gut bacteriota and parasitism. Empirical studies should enable to highlight associations, what is a critical pre-requisite before defining hypotheses about potential interactions and their underlying mechanisms.

Here, we strived to bridge these gaps by assessing the variability of gut bacteriota diversity and composition in wild populations of the bank vole *Myodes glareolus*, which is a small mammal reservoir of a large number of infectious agents (e.g., Abbate *et al.*, In_revision). We studied the relationships between its gut bacteriota, parasite infracommunities (focusing on gastro-intestinal helminths and pathogenic bacterial infections) and host and environmental factors that may either influence or indicate the health status of the host (e.g., proxies such as the body mass index (BMI)). The current study therefore addressed two main questions: (1) how is the structure (composition and diversity) of the gut bacteriota influenced by host and environmental factors? (2) does the structure of gut bacteriota also reflect associations with gastro-intestinal helminth and pathogenic bacterial communities?

Material and methods

Data collection

Bank vole sampling

Bank voles (*Myodes glareolus*) were trapped in summer, between late June and early September 2014 in forests located in four French localities (Table 1, Figure S1) distributed along a North-South transect in Eastern France. These localities are separated by 40 to 120 km from one another. The standardized trapping protocol used here was described in details in Dubois et al. (2018).

Rodents were euthanized using isofluorane and cervical dislocation, as recommended by

Mills (1995). A similar set of morphological measures was systematically recorded for each

individual. Age groups (juveniles and adults) were defined according to body mass and sexual

maturity. This latter was inferred using testes length and position, and seminal vesicle

development for males, or uterus size for females. Body condition was estimated using the

body mass index (BMI = weight/length^2). The digestive tract and the spleen were removed

and stored respectively in 96% ethanol and RNA later solution (-20°C).

Ethical statements: Animal capture and handling have been conducted according to the

French and European regulations on care and protection of laboratory animals (French Law

2001-486 issued on June 6, 2001 and Directive 2010/63/EU issued on September 22, 2010).

The CBGP laboratory has approval (D-34-169-003) from the Departmental Direction of

Population Protection (DDPP, Hérault, France), for the sampling of rodents and the storage

and use of their tissues.

Table 1- Number of bank voles analysed and prevalence of potentially pathogenic bacteria and gastro-intestinal

helminths for each sampling locality. N is the number of bank voles analysed (Grey cells). N_T represents the

number of individuals with data available for the three intra-host communities (gut bacteriota, pathogenic

bacteria and gastro-intestinal helminths). N_{GB}, N_{PB} and N_{GIH} respectively represent the number of individuals

with data available for each of these intra-host communities. '*Uninfected*' corresponds to the number of

uninfected bank voles for a given intra-host community. '*Co-infection*' corresponds to the number of bank voles

infected with at least two parasites of a given intra-host community. Prevalence is provided for each pathogenic

bacteria detected from the spleen, and each gastro-intestinal helminth. The red color gradient illustrates

variations in prevalence (0% = light red to 100% = dark red).

	Full data	Sampling localities			
		Mont-sous-Vaudrey	Chaux-des-Crotenay	Chatillon	Cormaranche
Full data					
N_T	124	22	34	36	32
Gut bacteriota					
N_{Gb}	161	37	42	41	41
Pathogenic bacteria					
N_{Pb}	138	22	34	38	37
Uninfected$_{Pb}$	11	3	1	3	4
Co-infection$_{Pb}$	67	10	21	16	20
Bartonella sp	0.73	0.80	0.81	0.69	0.66
Mycoplasma haemomuris	0.36	0.28	0.39	0.36	0.39
Anaplasma phagocytophilum	0.14	0.04	0.33	0.08	0.11
Neoehrlichia mikurensis	0.13	0.16	0.00	0.08	0.29
Orientia tsutsugamushi	0.08	0.00	0.08	0.21	0.00
Rickettsia sp	0.05	0.00	0.08	0.05	0.05
Spiroplasma sp	0.04	0.04	0.00	0.00	0.11
Borrelia miyamotoi	0.02	0.04	0.00	0.03	0.03
Borreliella afzelii	0.02	0.08	0.00	0.00	0.03
Mycoplasma coccoides	0.01	0.04	0.00	0.00	0.00
Gastro-intestinal helminths					
N_{Gih}	153	37	42	39	35
Uninfected$_{Gih}$	29	16	4	6	3
Co-infection$_{Gih}$	64	2	22	13	27
Heligmosomoides glareoli	0.48	0.49	0.36	0.54	0.54
Heligmosomum mixtum	0.41	0.00	0.76	0.00	0.86
Aonchotheca murissylvatici	0.35	0.03	0.26	0.59	0.54
Catenotaenia henttoneni	0.10	0.03	0.19	0.10	0.06
Paranoplocephala omphalodes	0.06	0.00	0.14	0.00	0.09
Arostrilepis horrida	0.03	0.00	0.00	0.00	0.14
Mastophorus muris	0.01	0.05	0.00	0.00	0.00
Trichuris arvicolae	0.01	0.03	0.00	0.00	0.00

199
200

Characterization of gut bacteriota

We first characterized the gut bacteriota of bank voles. We focused on the colon as rodent gut microbiota exhibits the highest level of bacterial diversity in the lower segments of the digestive tract (Suzuki & Nachman, 2016). DNA was extracted in 2016 from a 5 mm piece of colon tissue (taken about 1 cm far from the caecum - lumen was removed) of each bank vole using the ZymoBiomics 96 DNA Kit (Zymo) following the manufacturer's instructions. We amplified a 251-bp portion of the V4 region of the 16S rRNA gene (16S-V4F

[GTGCCAGCMGCCGCGGTAA] and 16S-V4R [GGACTACHVGGGTWTCTAATCC]),

following Kozich et al. (2013) and as described in Galan et al. (2016). Samples were

multiplexed using dual-indexes (index i5 in the forward primer and index i7 in the reverse

primer). Negative controls for extraction (whole reagents without DNA), for PCR (PCR mix

without DNA), and for indexing (wells without reagents corresponding to particular dual-

indexes combinations). All DNA extractions were analysed twice using two independent

technical replicates of amplicon libraries. PCR products were pooled, migrated and excised on

a low melting agarose gel (1.25%) then purified using the NucleoSpin Gel and PCR Clean-Up

kit (Macherey-Nagel) and quantified using the KAPA library quantification kit (KAPA

Biosystems) standardized to 4nM by qPCR spectrophotometry (assay). Sequencing was

performed on a 251-bp paired-end Illumina MiSeq run. The raw sequence reads (.fastq

format) have been deposited in the Zenodo Repository.

Sequence data were processed as described in Galan et al. (2016) using the pipelines

implemented in FROGS (Find Rapidly OTU with Galaxy Solution, Escudié *et al.*, 2018).

Briefly, the paired-end sequences were trimmed with CUTADAPT (Martin, 2011), merged with

FLASH (MAGOC & SALZBERG, 2011), and clustered into fine-scale molecular operational

taxonomy OTU units at 97% identity using the SWARM algorithm (Mahe *et al.*, 2014)

executed with aggregation parameter distance d=1 and a second pass performed on the seeds

of previous clusters with d=3. As such, OTUs do not correspond to a fixed clustering

threshold. Putative chimeras were removed using VSEARCH tools with de novo VUCHIME and

the cross-validation method. Taxonomy was assigned with BLASTN+ (Camacho *et al.*, 2009)

using the SILVA SSU Ref NR 128 database as a reference

(http://www.arb175silva.de/projects/ssu-ref-nr/). Filtering for false positives was carried out

as proposed by Galan et al. (2016). In short, we discarded positive results associated with

sequence counts below two OTU-specific thresholds, which checked respectively for cross-

contamination between samples (using the negative controls for extraction and PCR) and

incorrect assignment due to the generation of mixed clusters on the flow cell during Illumina

sequencing, using a false index-pairing rate for each PCR product of 0.02%, based on

estimates from Galan et al. (2016). For each sample, only OTUs found in the two technical

replicates were considered as positive, and OTUs found in only one of the two replicates were

removed. The number of sequences obtained for each technical replicate from a sample were

summed.

Lastly, we discarded OTUs and samples containing less than 500 reads in the dataset, as well

as OTUs considered to be contaminants, following (Salter *et al.*, 2014). Number of reads per

OTU were finally normalised to proportional abundance within each rodent (McKnight *et al.*,

2019). We only considered the family taxonomic rank for further analyses, but analyses at the

phylum level provided similar results (not shown).

Detection of pathogenic bacteria and gastro-intestinal helminths

We described the presence/absence of pathogenic bacteria from the spleen of each bank vole.

This lymphoid organ filters microbial cells in mammals and as such, enables to recover recent

infections (Abbate *et al.*, In_revision; Diagne *et al.*, 2017). Molecular protocols,

bioinformatics pipelines and data filtering were similar to those described above (gut

bacteriota), except for the DNA extraction from splenic tissue using DNeasy 96 Tissue Kit

(Qiagen). The potential pathogenicity of each bacterial OTU was assessed based on published

literature and on the Gideon database (https://www.gideononline.com/). Opportunistic

pathogens (*i.e.* commensal agents in healthy hosts, that become pathogenic when the balance

of the immune system is disrupted) were discarded from the dataset. Only the information of

the presence / absence of pathogenic OTUs was considered. For each bank vole, helminths

were carefully extracted and counted from the different sections of the digestive tract

(stomach, small intestine, large intestine and caecum), and classified by morphotype then

stored in 95% ethanol for further accurate identification. The latter was based on

unambiguous morphological criteria using conventional microscopy and generalist

identification keys or specific literature when available (Anderson *et al.*, 2009 ; Khalil *et al.*,

1994 ; Ribas Salvador *et al.*, 2011).

Statistical analyses

All statistical analyses were implemented in R v4.0.3 (team, 2020). For more convenience,

gut bacteriota, pathogenic bacteria and gastro-intestinal helminths were further described as

'intra-host communities'.

Gut microbiota diversity and composition

Description and analyses of bacterial communities were performed using the PHYLOSEQ

package (McMurdie & Holmes, 2013). We considered three features to analyse within hosts'

gut microbiota. i) We looked for enterotypes, *i.e.* distinct community composition types of gut

bacteriota, as found in humans (Arumugam *et al.*, 2011; Holmes *et al.*, 2012), using the

Dirichlet Multinomial Mixtures DMM (Morgan, 2021). ii) We analysed the Firmicutes

/Bacteroidetes (F/B) log-ratio, as it is often used as a proxy of health or metabolism in humans

and mice (Ley *et al.*, 2005; Toumi *et al.,* 2022). We calculated this ratio with the MICROBIOTA

package (Lahti & Shetty, 2017). iii) We characterized the alpha diversity using two metrics,

the specific richness (*i.e.* number of taxa within the host individual) and the Shannon index as

recommended in (Haegeman *et al.*, 2013).

We estimated the beta diversity, *i.e.,* the dissimilarity between host individuals in their gut

bacteriota using Bray-Curtis distances. We considered the relative abundance of OTUs

(family).

Influence of host and environmental factors on gut bacteriota diversity and composition

We tested the influence of individual characteristics (age class, gender, BMI) and localities,

independently on the F/B log-ratio and on the alpha diversity using generalized linear models

(GLM). We considered a negative binomial error distribution for the F/B ratio and the specific

richness, and a gaussian distribution for the Shannon index. Best model selection was

performed considering models with all possible combinations of factors and the DREDGE

function of the MUMIN package. The best model was selected using the Akaike information

criterion corrected for small sample size AICc, (J. B. Johnson & Omland, 2004). We assessed

the effect of each factor in the best model with the ΔAICc index. When the factor locality was

significant, Tukey's post-hoc tests were applied to evaluate pairwise differences between

localities, using the MULTCOMP package (Hothorn *et al.*, 2008). Residuals were checked to

graphically to ensure that all assumptions regarding normality, independence and the

homogeneity of variance were satisfied.

We evaluated the influence of geographic distance on the dissimilarities in gut bacteriota by

performing Mantel tests and using Pearson correlation (10,000 permutations). These tests

have less statistical power to address questions related to the variation in community

composition data among sites. Therefore, we also analysed the factors shaping the

dissimilarities in gut microbiota composition using several functions of the VEGAN package

(Oksanen *et al.*, 2020). Distance-based redundancy analyses (db-RDA) were performed to

analyse the effect of individual explanatory factors (age class, gender, BMI) and sampling

localities on dissimilarities in gut microbiota composition. Redundancy analyses are

appropriate to test hypotheses about the origin and maintenance of the variation in β diversity

(Legendre et al. 2005). We used the CAPSCALE function, followed by permutational

multivariate analyses of variance (PERMANOVA). We selected the best model, *i.e.*, the most

parsimonious one, using the ORDIR2STEP function (*P*-value adjusted and R^2 adjusted). For

each factor, we evaluated the intra-group dispersion using the BETADISPER function as

PERMANOVA analyses are sensitive to differences in dispersion among groups. A Tukey's

test was done to see if and which groups differed in relation to their variances. Lastly, we used

DESEQ2 package (Love et al., 2014) to identify the changes in bacteria taxa that best

explained gut bacteriome dissimilarities between individuals and localities. We performed

GLMs with negative binomial error (NBINOMWALDTEST method) and significant differences

were obtained after Benjamini & Hochberg corrections. They were visualised using the

METACODER package (Foster et al., 2017).

Relationships between gut bacteriota and pathogenic communities

We estimated the alpha diversity of the gastro-intestinal helminhth and pathogenic bacteria

community using the richness index (presence/absence data). We used GLMS and model

selection process described above to analyse whether the alpha diversity of each intra-host

community (gut bacteriota and pathogenic communities) was influenced by the alpha

diversity of the two other ones.

We estimated the beta diversity of the gastro-intestinal helminth and pathogenic bacteria

community using the Jaccard index (presence/absence data). The relationships between intra-

host community dissimilarities were investigated using three approaches. i) We applied

partial Mantel tests using MULTI.MANTEL (phytools package Revell, 2012) to analyse the

correlation between two matrices of dissimilarities (corresponding to two different

communities), while controlling for the effect of a third dissimilarity matrix (third

community). ii) We used db-RDA to analyse more deeply the relationships between the gut

bacteriota and the pathogenic (bacteria and helminths) communities. We included the alpha

diversity indices (richness specific) and infectious status as presence / absence) of pathogens

with prevalence greater than 10% in at least one locality as explanatory variables in these

analyses. We selected the best model using the ORDIR2STEP method. iii) We used DESEQ2 to

determine the gut bacteria taxa whose relative abundances changed with significant

explanatory variables.

Results

A total of 186 bank voles were trapped during the fieldwork campaign over the four targeted localities. For technical reasons (e.g., poor sample preservation, missing data), we could study the three intra-host communities for 124 rodents only.

Characterization of the gut bacteriota: taxa and enterotypes

Once the quality control steps were applied, the gut bacteriota dataset included 161 bank voles. We detected 10 phyla and 61 families of bacteria. At the phylum level, we found six predominant taxa that represented 99% of the gut bacteria relative abundance (Figure S2). At the family level, 11 families represented 93% of the relative abundance of the gut bacteriota (Figure 1A).

Figure 1- Composition of the intestinal bacteriota. A) The bar plot shows the individual variations of 11 bacterial families (F= Unknown family) belonging to 6 phyla and representing 93% of the total composition. Individuals (bars) are grouped by sampling localities, which are ordered from North to South. Each color represents a taxa. B) The composition of the two enterotypes identified using Dirichlet multinomial mixtures (DMMs), at family rank, is shown. Bacterial families are represented using the same colors as in A. C) the ratio (Firmicutes / Bacteroidetes) is shown for each sampling locality. Box and whisker plots represent the median and interquartile values. Black dots correspond to the mean value, and colored dots correspond to individuals. Different letters indicate statistically significant differences at $P < 0.05$, with pairwise Tukey post hoc adjustments.

We distinguished two enterotypes from the DMM approach. One (enterotype 1) was mainly

composed of the families *Helicobacteraceae*, *Lachnospiraceae*, *Muribaculaceae and*

Desulfovibrionaceae, while the other (enterotype 2) mainly included *Lactobacillaceae*,

Desulfovibrionaceae and *Eggerthellaceae* (Figure 1B).

Diversity of the gut bacteriota : the influence of sampling locality and host condition

We found that the *Firmicutes / Bateroidetes* ratio varied significantly between localities

(Figure 1C). Overall, northern localities exhibited lower F/B ratio than southern ones, with all

pairwise comparisons being significant except Chatillon *versus* Chaux-des-Crotenay.

Individual characteristics did not influence this ratio (Table S1).

The sampling locality had a significant global effect on the alpha diversity of the gut

bacteriota (*GLMs*. Specific richness: $F = 8.49$, $P < 10^{-3}$; Figure 2A; Shannon index: $F = 4.74$,

$P = 3 \times 10^{-3}$; Figure 2B; Table S2A). The locality Cormaranche exhibited a higher specific

richness than all other localities (*Tukey post hoc test*. Mont-sous-Vaudrey: $Z = 5.13$, $P_{adj} < 10^{-3}$,

Chaux-des-Crotenay: $Z = 4.57$, $P_{adj} < 10^{-3}$ and Chatillon: $Z = 3.62$, $P_{adj} = 1.7 \times 10^{-3}$) but a

lower level of diversity than Mont-sous-Vaudrey when considering taxa relative abundance

(*Tukey post hoc test*. Shannon index: $Z = -3.64$, $P_{adj} = 1.5 \times 10^{-3}$, Figure 2B). Body condition

(BMI) was also found to have a significant effect, but only when considering specific richness

($t = 2.91$; $P = 4 \times 10^{-3}$) – with higher values of BMI associated with increasing species

richness. All these results are detailed in Table S2A and Figures S3).

Figure 2- Variations of the gut bacteriota alpha diversity between localities. Alpha diversity is represented using
A) the specific richness of the gut bacteriota, and B) the Shannon index of the gut bacteriota, Results are shown
per locality, ordered from North to South. Each colored point represents an individual. Black points indicate the
average alpha diversity per locality. Box-and-whisker plots represent the median and interquartile values.
Different letters denote statistically significant differences at *P* < 0.05, with pairwise post-hoc Tukey
adjustments. C) Distance-based redundancy analysis (db-RDA) of the gut bacteriota (family) based on Bray–
Curtis dissimilarities. Dots represent individuals. The colors and shapes of the dots are associated with different
factors : Localities from North to South (Mont-sous-Vaudrey: Mont, Chaux-des-Crotenay: Chau, Chatillon:
Chat and Cormaranche: Corm) and gender (females: F and males: M). Significative factors based on the

ordiR2step analysis are shown as arrows. Ellipses represent a 80% confidence interval around the centroid of the

clusters, for each locality.

Composition of the gut bacteriota: sampling locality as the main factor of variation

We found a significant positive relationship between the dissimilarities in gut bacteriota

composition and the geographic distance (*Mantel test*. $r = 0.25$, $P = 10^{-4}$, Table S3A).

We found a significant effect of the sampling localities (*db-RDA*. $R^2_{adj} = 0.16$, $P = 1 \times 10^{-3}$)

and host gender (*db-RDA*. $R^2_{adj} = 0.01$, $P = 0.027$) on gut microbiota composition. The CAP1

axis discriminated Mont-sous-Vaudrey and Cormaranche localities (12.7% of the total

variance, Figure 2C). However, this result has to be taken cautiously as significant differences

of data dispersion were detected between localities (*betadisper*. $P = 1 \times 10^{-3}$). The locality

Cormaranche showed a lower dispersion compared to all other localities (Tukey multiple

comparisons, Table S3B).

We detected significant differences in the relative abundance of specific taxa using DeSeq2

(Table 2; Table S3C). The main changes (Log2 fold values higher than 20) were detected

between the northern (Mont-sous-Vaudrey) and southern localities. The northern population

was involved in 75% of all significant pairwise differences (Log2 fold change in composition

> 10). The gut bacteriota of these bank voles includes less Clostridiales (one unknown family;

Firmicutes), Bifidobacteriaceae (Actinobacteria) and Desulfovibrionales (two unknown

families; Proteobacteria), but more Erysipelotrichaceae (Firmicutes) than in all the three other

southern populations. The gut bacteriota of bank voles from Cormaranche (South) is

characterized by less Erysipelotrichaceae (Firmicutes), than in the three northern populations.

Table 2- Pairwise comparisons of the relative abundance of the gut bacteriota between sampling localities. Mont

= Mont-sous-Vaudrey, Chau = Chaux-des-Croteay, Chat = Chatillon, Corm = Cormaranche. The Log² fold value

is indicated for significant changes in abundance between two localities. Blue and red colors respectively

correspond to negative and positive values. Higher absolute changes in Log² fold are emphasized with darker

colors. The notation "order_fx" or "class_fx" is used when there was no assignation at the family level with the

SILVA database. Phylum is indicated in bold.

Taxonomy	Mont vs Chau	Mont vs Chat	Mont vs Corm	Chau vs Chat	Chau vs Corm	Chat vs Corm
Epsilonbacteraeota						
Helicobacteraceae	0	0	1.91	0	0	0
Proteobacteria						
Desulfovibrionaceae	−1.3	−1.63	−1.54	0	0	0
Firmicutes						
Ruminococcaceae	0	0	1.48	0	1.5	1.08
Lactobacillaceae	0	0	−1.81	0	0	0
Clostridiales	0	0	2.16	0	2.27	1.43
Clostridiales_F2	−26.15	−26.31	−26.54	0	0	0
Spirochaetes						
Spirochaetaceae	0	0	2.39	0	0	0
Actinobacteria						
Eggerthellaceae	0	−1.31	−2.42	0	−1.74	0
Bacteroidetes						
Bacteroidales_F4	0	0	0	0	0	−3.59
Muribaculaceae	0	0	1.74	0	1.93	1.64

422
423

Differences in the composition of the gut bacteriota between males and females bank voles

were driven by the phylum *Firmicutes*, with males exhibiting higher relative abundance of

this taxa than females (Table S3C).

Relationships between the diversity of the three intra-host communities

We found a significant relationship between the specific richness of the gut bacteriota and the

richness of the helminth community. A more diverse gut bacteriota was associated with a

greater number of helminth species infecting bank voles (*GLM. F* = 14.09, *P* < 10^{-3}; Figure

3A; Table S2B). We also found a positive relationship between the specific richness of the

pathogenic bacteria and the richness of the gastro-intestinal helminth community (*GLM. F*=

6.99, *P* = 9 x 10^{-3}; Figure 3A; Table S2B). Lastly, we found a significant effect of the specific

richness of both the gut bacteriota and of pathogenic bacteria on the richness of the gastro-

intestinal helminth community (*GLM.* Gut bacteriota. *t* = 3.50, *P* < 10^{-3}; pathogenic bacteria *t*

= 2.38, *P* = 0.019; Figure 3A, Table S2B).

Figure 3- Associations between the diversity and composition of the three intra-host communities. The two upper diagrams show the relationships between A) the specific richness and B) the composition of the three communities, while taking into account the influence of the sampling localities or distance geographic. The effect size and the direction of the relationship between communities are represented using an arrow, its width corresponding to the estimate (β) x 10 or the correlation index R x 10. Only significant effects are represented. C) This db-RDA triplot shows the structure of the gut bacteriota and the correlations with the pathogen communities. The arrows correspond to the significant explanatory variables. Each point corresponds to an individual, and the colors correspond to the different sampling localities. A.mur: *Aonchotheca murissylvatici*, Bar: *Bartonella* sp, H.mix: *Heligmosomum mixtum*.; Helicobacter.: Helicobacteraceae, Spiroch.: Spirochaetaceae, Clostri.F2: Clostridiales_f2, Clostri.F1: Clostridiales_f1, Lactob.:Lactobacillaceae,

Eggerthe.: Eggerthellaceae, Desulfov.: Desulfovibrionaceae , Bacter.F1: Bacteroidales_f1,
Lachno.:Lachnospiraceae; Murib.: Muribaculaceae, Ruminoco.: Ruminococcaceae

Relationships between the composition of the three intra-host communities

We found a positive relationship between dissimilarities in the gut bacteriota and

dissimilarities in the gastro-intestinal helminth community composition (*partial Mantel test. r*

$= 0.16$, $P = 1 \times 10^{-4}$; Figure 3B), but not with dissimilarities in the pathogenic bacteria

community composition (*partial Mantel test. r* $= 0.02$, $P = 0.28$). After controlling for

geographic distances, dissimilarities in gut bacteriota composition remained significantly

correlated with dissimilarities in helminth community composition (*partial Mantel test. r*

$=0.12$; $P = 0.001$). Further details are provided in Table S3A.

We detected significant associations between the whole composition of the gut bacteriota and

the presence / absence of three pathogens: *Aonchotheca murissylvatici* (*ordistep db-RDA.*

$R^2_{adj} = 0.0$, $P = 2 \times 10^{-3}$), *Bartonella* sp (*ordistep db-RDA. $R^2_{adj} = 0.01$, $P = 0.04$*) and

Heligmosomum mixtum (*ordistep db-RDA. $R^2_{adj} = 0.06$, $P = 2 \times 10^{-3}$*). The db-RDA triplot

based on the two first axes only represented 10.2% of the total variance (Figure 3C; Figure

S4). It showed that individuals infected with *Aonchotheca murissylvatici*

or *Heligmosomum mixtum*, but not infected with *Bartonella sp.,* had

more *Lactobacillaceae* (*Firmicutes*), *Desulfovibrionaceae* (*Proteobacteria*)

and *Eggerthellaceae* (*Actinobacteria).* These individuals also had less *Spirochaetaceae*

(*Spirochaeta*), *Muribaculaceae (Bacteroidetes)*, *Helicobacteraceae* (*Epsilonbacteraeota*)

and *Ruminococcaceae* (*Firmicutes*). This pattern is correlated with the sampling localities.

Individuals from northern localities are distributed on the left side of the CAP1 axis, and

southern ones on the right side of it (Figure 3C). Neither the specific richness of pathogenic

bacteria nor the specific richness of the gastro-intestinal helminth community had a

significant effect on the global composition of the gut bacteriota (Table S3D).

The specific richness of the gastro-intestinal helminth community, as well as infections with

A. murissylvatici and *H. mixtum,* were only slightly associated with different relative

abundance of particular gut bacteria taxa (*DeSeq2.* Log2 fold changes did not exceed 3.5).

These changes concerned four main families. Rhizobiaceae and Spirochaetaceae showed

negative associations with gastro-helminth specific richness and *A. murissylvatici*. Mollicutes

(undetermined family) and Saccharimonadaceae showed positive associations with *A.*

murissylvatici and *H. mixtum* (Table 3). More details are provided in Table S3E.

485   **Table 3**- Changes in relative abundance of the gut bacteriota with regard to infectious status (helminths and
486   pathogenic bacteria) and specific richness. The Log^2 fold change in relative abundance is indicated for
487   significant values only. Negative values are represented with blue colors, positive values with red colors.
488   Higher absolute changes in Log^2 fold are emphasized with darker colors.

Phylum	Family	log2FoldChange
Richness of GI helminths		
Actinobacteria	Eggerthellaceae	0.67
Proteobacteria	Rhizobiaceae	−3.79
Epsilonbacteraeota	Helicobacteraceae	−0.89
Aonchotheca murissylvatici		
Proteobacteria	Desulfovibrionaceae	1.13
Tenericutes	Mollicutes_f1	3.07
Patescibacteria	Saccharimonadaceae	3.03
Spirochaetes	Spirochaetaceae	−1.75
Actinobacteria	Eggerthellaceae	1.4
Heligmosomum mixtum		
Tenericutes	Mollicutes1	3.34
Mastophorus muris		
Proteobacteria	Desulfovibrionales3	−29.98
Firmicutes	Clostridiales_f1	−21.41
Firmicutes	Clostridiales_f2	−23.73
Catenotaenia henttoneni		
Proteobacteria	Rhizobiaceae	−24.71
Proteobacteria	Burkholderiaceae	−23.96
Firmicutes	Erysipelotrichaceae	−20.6
Paranoplocephala omphalodes		
Firmicutes	Clostridiales_f2	−25.37
Proteobacteria	Desulfovibrionales_f3	−23.74
Proteobacteria	Rhizobiaceae	−24.86
Proteobacteria	Burkholderiaceae	−22.69
Firmicutes	Erysipelotrichaceae	−22.4
Firmicutes	Christensenellaceae	−5.24
Trichuris arvicolae		
Firmicutes	Christensenellaceae	−20.69
Firmicutes	Clostridiales_f1	−18.62
Bacteroidetes	Bacteroidales_f4	−18.38
Richness of Pathogenic bacteria		
Bacteroidetes	Bacteroidales_f3	−28.51
Neoehrlichia mikurensis		
Firmicutes	Erysipelotrichaceae	−23.42
Firmicutes	Clostridiales	−2.72
Bacteroidetes	Muribaculaceae	−1.76
Orientia tsutsugamushi		
Firmicutes	Erysipelotrichaceae	−22.95
Proteobacteria	Desulfomicrobiaceae	−5.71
Bacteroidetes	Muribaculaceae	−2.45
Anaplasma phagocytophilum		
Firmicutes	Ruminococcaceae	−1.75
Rickettsia sp		
Firmicutes	Erysipelotrichaceae	−21.81

In the opposite, we found strong changes in relative abundance of gut bacteriota families with

other gastro-intestinal helminths and pathogenic bacteria infections (*DeSeq2.* Log2 fold

higher than 18, Table 3). It concerned infections with the helminth species *Mastophorus*

muris, Catenotaenia henttoneni, Paranoplocephala omphalodes and *Trichuris arvicolae.*

These associations were all negative and mostly involved the same bacterial families (Table

S3E), namely undetermined families of Bacteroidales (Bacteroidetes), Desulfovibrionales

(Proteobacteria) or Clostridiales (Firmicutes), Erysipelotrichaceae (Firmicutes), Rhizobiaceae

(Proteobacteria) and Burkholderiaceae (Proteobacteria).

Considering pathogenic bacteria, we found that higher levels of specific richness were

associated with lower relative abundance of an undetermined Bacteroidales family

(Bacteroidetes), and that *Neoehrlichia mikurensis, Orientia tsutsugamushi* and *Rickettsia* sp

infections were associated with strong decreases in relative abundance of *Erysipelotrichaceae*

(Firmicutes) (Table 3; Table S3E). Other associations between bacterial infections and

changes in relative abundance of specific gut bacteriota taxa were detected, but with little size

effect (*DeSeq2.* log2 fold changes lower than 5).

507   **Discussion**

Understanding the complex interlinkages between host microbiota, host-pathogen interactions

and health in wild animal populations has become a key topic in disease ecology. Such

understanding is instrumental for deciphering population dynamics, and designing strategies

for zoonotic risk management or biodiversity conservation. Here, we use a combination of

metabarcoding and community ecology approaches to *(i)* describe the gut microbiota of wild

rodent populations and their variations at a regional geographical scale, and *(ii)* explore the

three-way relationships between the gut bacteriota and communities of gastro-intestinal

helminths and pathogenic bacteria.

Spatial variations of gut bacteriota and their potential causes

The gut microbiota of bank voles has been mainly examined in the context of exposure to

radioactive pollutants (e.g., Lavrinienko (2018), but see Knowles et al. (2019)). In this study,

we focused on localities sampled at a regional scale (100 km) along a North-South gradient in

Eastern France.

We found significant inter-individual variations in the gut bacteriota composition although
host factors such as gender and age played little role. Interestingly, we found that all
individuals were clustered within two groups, which could be distinct enterotypes if
longitudinal surveys confirmed this clustering through time (Arumugam *et al.*, 2011).
Enterotypes have already been described in wild rodents (Goertz *et al.*, 2019; Li *et al.*, 2016),
and they might reflect distinct ways of generating energy from substrates available in the
digestive tract, as well as differences in diet (Rinninella *et al.*, 2019; Wang *et al.*, 2014).
These enterotypes could be associated with different diets in bank voles: one oriented toward
seeds and plants and another one toward insects and berries. Indeed, enterotype 1 is
characterized by families (namely, Helicobacteraceae, Lachnospiraceae and Muribaculaceae)
that are involved in the breakdown of carbohydrates, fermentation of plant saccharid and
degradation of glycan (see refs in Goertz *et al.*, 2019). These families are also predictive
signals of a high-fat diet in mice (Bowerman *et al.*, 2021; Rodriguez-Daza *et al.*, 2020).
Conversely, enterotype 2 is characterized by families (namely, Lactobacillae and
Eggerthellaceae) that can be involved in the digestion of fermented food (e.g., rodent food
store over winter), and insect skeleton (see refs in Maurice *et al.*, 2015) or in the degradation
of polyphenol (Rodriguez-Daza *et al.*, 2020). All these aliments have varying nutritional and
chemical composition and may be part of bank vole diet (Ecke *et al.*, 2018). The fraction of
these different types of resources in bank vole diet may vary with resource preference or
availabilities, reproductive status, sampling date and location (e.g., Maurice *et al.*, 2015). It
would be interesting to develop semi-natural experiments to survey rodent diet and gut
microbiome through time, and analyse the link with enterotypes in bank voles (Wang *et al.*,
2014).
There are now many evidence that the environment in which hosts evolve (through abiotic
and biotic factors) is likely to shape variations in gut bacteriota composition between
localities sampled and studies. Previous works have already shown that the structure of rodent
gut microbiota varied between localities at large spatial scales due to biogeographic or genetic
factors (Linnenbrink *et al.*, 2013). Geographic variability has also been found at smaller
spatial scales (e.g., few km Goertz *et al.*, 2019). Here, our results provide significant evidence
for spatial structure of gut bacteriota between bank vole populations that are between 50 and
130 km away, with no clear barrier to dispersal or gene flow (Dubois *et al.*, 2018).
We observed gradual changes between bank voles from the northern and southern populations
in terms of gut bacteriota richness, evenness, composition and particularly

Firmicutes/Bacteroidetes ratio. Although the links between the diversity and functional
capacity of the gut bacteriota still need to be better understood (Worsley *et al.*, 2021), it is
largely assumed that changes in diversity are associated with shifts in metabolism (Reese &
Dunn, 2018). Bank voles from southern populations exhibit higher specific richness and lower
evenness of the gut bacteriota, as well as lower dispersion of gut bacteriota composition. They
have higher levels of body condition and F/B ratio, which are indicative of an optimisation of
calorie intake and absorption, weight gain and fat storage (see refs in Wolf *et al.*, 2021).
Altogether, these results could suggest strong constraints on gut bacteriota function to
maximise energy extraction. The northern populations show the opposite patterns. Lower
BMI and lower levels of F/B ratio might reflect energy production and conversion, amino acid
transport and metabolism. Diversity patterns (higher evenness and lower specific richness of
the gut bacteriota) could suggest lower stochasticity and/or directional selection. Further
studies are required to investigate the eco-evolutionary processes driving these changes in gut
bacteriota.
Lastly, these differences in gut microbiota composition between the northern and southern
populations might also reflect physiological variations related to physiology, health and
(potentially) immunity. Indeed, Clostridiales and Bifidobacteriaceae participate in the
maintenance of intestinal homeostasis, and in the regulation of inflammation or in the gut
barrier function (Arboleya *et al.*, 2016; Hakansson & Molin, 2011; Lopetuso *et al.*, 2013).
Specific taxa within Erysipelotrichaceae may be correlated with inflammation or have
immunogenic potential (Kaakoush, 2015; Zhai *et al.*, 2019). Desulfovibrionales activities
result in the production of H_2S, which in turn negatively affects the gut barrier, production of
endotoxins and pro-inflammatory cytokines (Hu *et al.*, 2022). Our previous work revealed
that bank voles from these southern populations had lower basal level of *Tnf-a* (a pro-
inflammatory cytokine) and higher level of *Mx2* antiviral gene expression than those from
these northern populations (Dubois *et al.*, 2018). Future studies should assess *(i)* the potential
relationships between variations in gut bacteriota composition and *(ii)* the capacities to
regulate or mount immune responses and inflammation in these bank vole populations.
583
Three-way relationships between intra-host communities
At the community level, we have not found strong evidence for three-way relationships
between the gut microbiota (diversity or composition), the gastro-intestinal helminth and
pathogenic bacteria communities. Neither the specific richness nor the global composition of

a given community are related to the richness or composition of the two other ones. By

contrast, particular taxa seem to be involved in these three-way relationships.

First, some infections are significantly associated with the global composition of the gut

bacteriota, but have only little impact on specific gut bacterial taxa. This result concerns two

helminths *Heligmosomum mixtum*, *Aonchotheca murissylvatici* and the hemotrophic bacteria

Bartonella. Opposite patterns are observed for the helminths and for the bacteria. They could

reflect the antagonistic impacts of these infections on the gut bacteriota, or negative

interactions between these pathogens. On one hand, some evidence suggests that *Bartonella*

may be acting as a symbiont more than a pathogen (McKee *et al.*, 2021). Significant

coevolutionary congruence has been found between *Bartonella* species and their rodent hosts,

and *Bartonella* infections in rodents lead to an asymptomatic long lasting intra-erythrocytic

bacteraemia (Deng *et al.*, 2012; Lei & Olival, 2014). It would be interesting to test whether

associations between *Bartonella* and gut bacteriota could corroborate the hypothesis of

coadaptation between these bacteria and their rodent hosts (Hayman et al., 2013). On the other

hand, some hookworms have been shown to induce changes in rodent gut bacteriota (review

in Mutapi, 2015). Infection of mice with the nematode *Heligmosomoides polygyrus* (which is

phylogenetically close from *Heligmosomum mixtum*) lead to an increased abundance of

Lactobacillaceae in the gut microbiome (Reynolds *et al.*, 2014), as suggested by the

correlation detected in our study. Lastly, negative interactions between *H. mixtum or A.*

murissylvatici and *Bartonella* are probable. Indeed gastro-intestinal hookworms are known to

induce anaemia (Seguel & Gottdenker, 2017), while *Bartonella* invades and replicates in red

blood cells. This resource limitation driven by helminths on erythrocyte-dependent infectious

agents is an important driver of helminth-microparasite coinfection (Graham, 2008).

Therefore, the negative associations detected here between *Bartonella* and *H. mixtum or A.*

murissylvatici – and their respective links with gut bacteriota composition – likely seem to be

driven by potential complex antagonistic, synergistic and symbiotic interactions that need to

be further explored.

Second, other infections are strongly associated with large changes in the relative abundance

of one or few specific taxa from the gut bacteriota, but not with the global composition. Three

bacterial infections (*Neoehrlichia* sp, *Orientia* sp and *Rickettsia* sp) as well as *P. omphalodes*

infections exhibited the same pattern: they were associated with a lower relative abundance of

Erysipelotrichaceae. It is striking to find such common associations for these infectious agents

because observed changes in the gut microbiota during infection are rarely consistent, even

for single pathogens (Sabey *et al.*, 2021). The most obvious features shared between these

infectious agents is that they are transmitted by arthropods. Unfortunately, knowledge on

Erysipelotrichaceae and its links with infection or dysbiosis is still deficient to explain the

pattern observed. To our knowledge, such associations have been investigated in humans

only. They have shown that increased abundance of Erysipelotrichaceae could be associated

with a number of diseases such as tuberculosis, HIV and norovirus infections, inflammation-

related intestinal disease and metabolic disorders (Kaakoush, 2015). The reasons why

Neoehrlichia sp, *Orientia* sp, *Rickettsia* sp or *P. omphalodes* infections are associated with a

decreased abundance of Erysipelotrichaceae in bank voles remain to be investigated.

Marked relationships between gut bacteriota and gastro-intestinal helminth

communities

While we do not detect three-way relationships between the whole composition of gut

bacteriota, helminth and pathogenic bacteria communities, we highlight strong pairwise

associations between helminth community and gut bacteriota. The former pattern might be

explained by the fact that the pathogenic bacteria detected here do not constitute a functional

community. The main reason is that their ecological niche can be very different. For example,

hemotrophic *Mycoplasma* parasitizes erythrocytes (Alabi *et al.*, 2020) while *Borrelia*

disseminates through the bloodstream and/or lymphatic system to invade and colonize various

tissue (Zeidner *et al.*, 2001).

The strong associations between gastro-intestinal helminths and gut bacteriota may be

interpreted under two perspectives. First, the strong positive associations between the

diversity of helminth community and gut bacteriota might corroborate the hypothesis.

Besides, experimental evidence showed that helminths have the capacity to maintain higher

gut microbiota diversity and may represent gut homoeostasis (Kreisinger *et al.*, 2015). Indeed,

low-intensity, chronic helminth infections are commonly linked to high microbial diversity

and predominance of bacteria typically associated with gut health (Peachey *et al.*, 2017).

Nevertheless, this interpretation has to be taken cautiously as the diversity of both

communities was strongly influenced by the localities of sampling. The environment might

therefore shape similarly gut bacteriota and helminth community diversity.

Second, significant associations between helminth community and gut bacteriota composition

– which remain significant even when potential geographic confounding effects were

removed – may be linked to the fact that both communities reside in the same environmental

niche (host intestines). From there, they likely experience similar selection pressure (e.g., host

immune responses) which could shape their composition (Glendinning *et al.*, 2014). One

could expect therefore potentially strong interactions and reciprocal influence between them,

Unfortunately, the causal processes behind these gut microbiota and helminths interactions

are complex, multifaceted and difficult to assess. This intricacy is amplified by the fact that

experimental studies mostly focus on single helminth infections while interactions

between/within community are the rule within host organisms. The field of microbiota

research would thus benefit from taking into account the whole composition of gastro

intestinal helminth community rather than single helminth infections only.

In this study, we also highlight a large number of species-specific associations between

helminths infections and members of the gut bacteriota. High-intensity, acute helminth

infections may correlate with changes in hosts gut microbiota, through direct and indirect

interactions (e.g., immune or other processes such as malnutrition; Peachey *et al.*, 2017).

Nevertheless, the patterns of shifts in gut bacteriota associated with helminth infections

remain hardly predictable so far. As such, research works addressing this issue with

laboratory or wild animals have provided variable, and sometimes even contradictory

conclusions. Most surprising is that these inconsistent patterns are also found when focusing

on single host-helminth models. A potential explanation is that these infection-associated

microbiota shifts could depend on the presence of other helminths and the duration of

infection (Sabey et al., 2021). Local interactions between helminths and between helminths

and gut bacteria could mediate changes in infection outcomes as well as the gut bacteria and

helminth populations themselves (Glendinning *et al.*, 2014).

677   **Conclusion**

Altogether, these results emphasize complex interlinkages between gut bacteriota, gastro-

intestinal helminths and bacterial infections in wild animal populations. We emphasize the

strong impact of environment, even at fine geographical scales, on these interactions. Shifts in

diet or host genetics could mediate the spatial changes observed in gut bacteriota. However,

the processes shaping gut bacteriota diversity and composition are many and complex, and

further investigations are required to decipher the relative importance of drift, dispersal or

selection on bank vole gut bacteriota in the populations studied here. Besides, we find a

diverse array of associations between gut bacteriota and gastro-intestinal helminths or

pathogenic bacteria, some being significant at the scale of the whole community and other

being species-specific only. Whether these patterns reflect coadaptation, dysbiosis or indirect

interactions with host immunity and coinfections should now be considered to better

understand the spatial variations observed in the relationships between gut bacteriota and

health.

Acknowledgement

Data used in this work were partly produced through the genotyping and sequencing facilities

of ISEM (Institut des Sciences de l'Evolution-Montpellier) and Labex CeMEB (Centre

Méditerranéen Environnement Biodiversité).

Funding

This research was funded through the INRAE metaprogram MEM Hantagulumic and the

2018-2019 BiodivERsA joint call for research proposals, under the BiodivERsA3 ERA-Net

COFUND programme, and with the funding organization ANR. A. Dubois PhD was funded

by an INRA-EFPA/ANSES fellowship.

Conflict of interest disclosure

The authors declare that they have no financial conflict of interest with the content of this

article. N. C. is one of the PCI Inf recommenders. B.R. is part of the managing board of PCI

Inf.

Data and script availability

Raw data and scripts are available on zenodo: https://doi.org/10.5281/zenodo.7433800

Supplementary information

All supplementary materials are available on zenodo :

https://doi.org/10.5281/zenodo.7431842

Supplementary Figure S1. Maps showing the sampling area (left) and localities (right) in

France. Forests are indicated in green and water in blue. The four sampling localities are

represented with a colored polygon. The arrow indicates the North.

Supplementary Figure S2. Composition of the gut bacteriota. The relative abundance of six phyla representing 99% of the total composition is represented. Individuals are grouped by sampling localities, which are ordered from North to South. (A) Bar graph shows individual variation in phyla composition (phylum=color). (B) Box and whisker plots represent median and interquartile values for each phylum. Black dots correspond to mean values, and colored dots correspond to individuals.

Supplementary Figure S3. Variations of alpha diversity with individual factors, for the gut bacteriota (family level), pathogenic bacteria and gastro-intestinal helminths of bank voles. Alpha diversity is estimated using the specific richness (A, B and C) and the Shannon index (D, E and F). In graphs C and F, the blue line corresponds to the linear regression line.

Supplementary Figure S4. Relationships between the composition of the gut bacteriota, pathogenic bacteria and gastro-intestinal helminth communities: The db-RDA triplot shows the structure of the gut bacteriota at the phylum level and the correlations with the intra-host parasite communities. The arrows correspond to the significant explanatory variables. Each point corresponds to an individual, and the colors correspond to the different sampling localities.

Supplementary Table S1. Variation of the Firmicutes/Bacteroidetes ratio with localities and individual factors.

Supplementary Table S2. Alpha diversity metrics and statistics for the gut bacteriota, pathogenic bacteria and helminth communities of bank voles.

Supplementary Table S3. Beta diversity metrics and statistics for the gut bacteriota, pathogenic bacteria and helminth communities of bank voles.

AUTHOR CONTRIBUTIONS

M.B.: Data curation; Formal analysis; Methodology; Writing – original draft

M.G.: Conceptualization; Data curation; Formal analysis; Methodology; Supervision; Writing – review and editing

A.D.: Conceptualization; Data curation

C.A.D.: Data curation; Writing – review and editing

P.M.: Conceptualization; Funding acquisition; Investigation; Supervision; Writing – review

and editing

B.R.: Conceptualization; Methodology; Supervision; Writing – review and editing

N.C.: Conceptualization; Funding acquisition; Investigation; Methodology; Project

administration; Resources; Supervision; Writing – original draft

REFERENCES

Abbate, J., Galan, M., Razzauti, M*., et al.* (In_revision). Pathogen community composition and co-
infection patterns in a wild community of rodents. *PCI Ecology*. doi:10.1101/2020.02.09.940494

Adair, K. L.,Douglas, A. E. (2017). Making a microbiome: the many determinants of host-associated
microbial community composition. *Current Opinion in Microbiology, 35*, 23-29.
doi:10.1016/j.mib.2016.11.002

Alabi, A. S., Monti, G., Otth, C*., et al.* (2020). Molecular survey and genetic diversity of
Hemoplasmas in rodents from Chile. *Microorganisms, 8*(10). doi:10.3390/microorganisms8101493

Anderson, R. C., Chabaud, A. G.,Willmott, S. (2009). *Keys to the nematode parasites of vertebrates:*
archival volume. CABI, Wallingford, Oxon, UK.

Arboleya, S., Watkins, C., Stanton, C*., et al.* (2016). Gut Bifidobacteria populations in human health
and aging. *Frontiers in Microbiology, 7*. doi:10.3389/fmicb.2016.01204

Arumugam, M., Raes, J., Pelletier, E*., et al.* (2011). Enterotypes of the human gut microbiome.
Nature, 473(7346), 174-180. doi:10.1038/nature09944

Belkaid, Y.,Hand, T. W. (2014). Role of the microbiota in immunity and inflammation. *Cell, 157*(1),
121-141. doi:10.1016/j.cell.2014.03.011

Bowerman, K. L., Knowles, S. C. L., Bradley, J. E*., et al.* (2021). Effects of laboratory domestication
on the rodent gut microbiome. *ISME Communications, 1*(1), 49. doi:10.1038/s43705-021-00053-9

Camacho, C., Coulouris, G., Avagyan, V*., et al.* (2009). BLAST+: architecture and applications. *BMC*
Bioinformatics, 10, 421. doi:10.1186/1471-2105-10-421

Clemente, J. C., Ursell, L. K., Parfrey, L. W*., et al.* (2012). The impact of the gut microbiota on
human health: an integrative view. *Cell, 148*(6), 1258-1270. doi:10.1016/j.cell.2012.01.035

Cortes, A., Peachey, L.E., Jenkins, T.P. et al. (2019). Helminths and microbes within the vertebrate
gut - not all studies are created equal. *Parasitology* 146,1371-1378.
doi:10.1017/s003118201900088x

Deng, H. K., Le Rhun, D., Buffet, J. P. R*., et al.* (2012). Strategies of exploitation of mammalian
reservoirs by *Bartonella* species. *Veterinary Research, 43*. doi:10.1186/1297-9716-43-15

Diagne, C. A., Charbonnel, N., Henttonen, H*., et al.* (2017). Serological survey of zoonotic viruses in
invasive and native commensal rodents in Senegal, West Africa. *Vector Borne and Zoonotic*
Diseases, 17(10), 730-733. doi:10.1089/vbz.2017.2135

Dubois, A., Castel, G., Murri, S*., et al.* (2018). Bank vole immunoheterogeneity may limit nephropatia
epidemica emergence in a French non-endemic region. *Parasitology, 145*, 393–407.
doi:10.1017/S0031182017001548

Ecke, F., Berglund, A. M., Rodushkin, I*., et al.* (2018). Seasonal shift of diet in bank voles explains
trophic fate of anthropogenic osmium? *Science of the Total Environment, 624*, 1634-1639.
doi:10.1016/j.scitotenv.2017.10.056

Escudié, F., Auer, L., Bernard, M*., et al.* (2018). FROGS: Find Rapidly OTU with Galaxy Solution.
Bioinformatics, 34, 1287–1294.

Galan, M., Razzauti, M., Bard, E*., et al.* (2016). 16S metagenomics for epidemiological survey of
bacteria in wildlife. *mSystem, 1*(4), e00032-00016. doi:10.1128/mSystems.00032-16

Glendinning, L., Nausch, N., Free, A*., et al.* (2014). The microbiota and helminths: sharing the same
niche in the human host. *Parasitology, 141*(10), 1255-1271. doi:10.1017/S0031182014000699

Goertz, S., de Menezes, A. B., Birtles, R. J*., et al.* (2019). Geographical location influences the
composition of the gut microbiota in wild house mice (*Mus musculus domesticus*) at a fine spatial
scale. *Plos One, 14*(9). doi:10.1371/journal.pone.0222501
Graham, A. L. (2008). Ecological rules governing helminth-microparasite coinfection. *Proceedings of*
the National Academy of Sciences of the United States of America, 105(2), 566-570.
doi:10.1073/pnas.0707221105
Haegeman, B., Hamelin, J., Moriarty, J*., et al.* (2013). Robust estimation of microbial diversity in
theory and in practice. *Isme Journal, 7*(6), 1092-1101. doi:10.1038/ismej.2013.10
Hakansson, A.,Molin, G. (2011). Gut microbiota and inflammation. *Nutrients, 3*(6), 637-682.
doi:10.3390/nu3060637
Haukisalmi, V., Henttonen, H. Coexistence in helminths of the bank vole *Clethrionomys glareolus*. I.
Patterns of co-occurrence. *Journal of Animal Ecology* 62,221-229
Hoarau, A. O. G., Mavingui, P.,Lebarbenchon, C. (2020). Coinfections in wildlife: Focus on a
neglected aspect of infectious disease epidemiology. *Plos Pathogens, 16*(9).
doi:10.1371/journal.ppat.1008790
Holmes, I., Harris, K.,Quince, C. (2012). Dirichlet multinomial mixtures: Generative models for
microbial metagenomics. *Plos One, 7*(2). doi:10.1371/journal.pone.0030126
Honda, K.,Littman, D. R. (2016). The microbiota in adaptive immune homeostasis and disease.
Nature, 535(7610), 75-84. doi:10.1038/nature18848
Hothorn, T., Bretz, F.,Westfall, P. (2008). Simultaneous inference in general parametric models.
Biometrical Journal, 50(3), 346-363. doi:10.1002/bimj.200810425
Hu, H., Shao, W. T., Liu, Q*., et al.* (2022). Gut microbiota promotes cholesterol gallstone formation
by modulating bile acid composition and biliary cholesterol secretion. *Nature Communications,*
13(1). doi:10.1038/s41467-021-27758-8
Johnson, J. B.,Omland, K. S. (2004). Model selection in ecology and evolution. *Trends in Ecology and*
Evolution, 19, 101–108.
Johnson, P. T., de Roode, J. C.,Fenton, A. (2015). Why infectious disease research needs community
ecology. *Science, 349*(6252), 1259504. doi:10.1126/science.1259504
Kaakoush, N. O. (2015). Insights into the role of Erysipelotrichaceae in the human host. *Frontiers in*
Cellular and Infection Microbiology, 5. doi:10.3309/fcimb.2015.00004
Kamada, N., Chen, G. Y., Inohara, N*., et al.* (2013). Control of pathogens and pathobionts by the gut
microbiota. *Nature Immunology, 14*(7), 685-690. doi:10.1038/ni.2608
Karst, S.M., Wobus, C.E. (2019). Viruses and the microbiome. Current Opinion in Virology 37,III-VI.
doi:10.1016/j.coviro.2019.06.012
Khalil, L. F., Jones, A.,Bray, R. A. (1994). *Keys to the cestode parasites of Vertebrates.* CAB
International, Wallingford, Oxon, UK.
Khosravi, A.,Mazmanian, S. K. (2013). Disruption of the gut microbiome as a risk factor for microbial
infections. *Current Opinion in Microbiology, 16*(2), 221-227. doi:10.1016/j.mib.2013.03.009
Kim, C. H. (2021). Control of lymphocyte functions by gut microbiota-derived short-chain fatty acids.
Cellular & Molecular Immunology, 18(5), 1161-1171. doi:10.1038/s41423-020-00625-0
Knowles, S. C. L., Eccles, R. M.,Baltrunaite, L. (2019). Species identity dominates over environment
in shaping the microbiota of small mammals. *Ecology Letters, 22*(5), 826-837.
doi:10.1111/ele.13240
Kolodny, O.,Schulenburg, H. (2020). Microbiome-mediated plasticity directs host evolution along
several distinct time scales. *Philosophical Transactions of the Royal Society B-Biological Sciences,*
375(1808). doi:10.1098/rstb.2019.0589
Kozich, J. J., Westcott, S. L., Baxter, N. T*., et al.* (2013). Development of a dual-index sequencing
strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina
sequencing platform. *Applied and Environmental Microbiology, 79*(17), 5112-5120.
doi:10.1128/aem.01043-13
Kreisinger, J., Bastien, G., Hauffe, H. C*., et al.* (2015). Interactions between multiple helminths and
the gut microbiota in wild rodents. *Philosophical Transactions of the Royal Society B-Biological*
Sciences, 370(1675). doi:10.1098/rstb.2014.0295
Lahti, L.,Shetty, S. (2017). microbiome R package. Bioconductor. doi: 10.18129/B9.bioc.microbiome

Lawson, M.A.E., Roberts, I.S., Grencis, R.K. (2021). The interplay between *Trichuris* and the
microbiota. *Parasitology* 148,1806-1813. doi:10.1017/s0031182021000834

Lavrinienko, A., Mappes, T., Tukalenko, E*., et al.* (2018). Environmental radiation alters the gut
microbiome of the bank vole *Myodes glareolus*. *Isme Journal, 12*(11), 2801-2806.
doi:10.1038/s41396-018-0214-x

Lei, B. N. R.,Olival, K. J. (2014). Contrasting patterns in mammal-bacteria coevolution: *Bartonella*
and *Leptospira* in bats and rodents. *Plos Neglected Tropical Diseases, 8*(3).
doi:10.1371/journal.pntd.0002738

Leung, J. M., Graham, A. L.,Knowles, S. C. L. (2018). Parasite-microbiota interactions with the
vertebrate gut: Synthesis through an ecological lens. *Frontiers in Microbiology, 9*.
doi:10.3389/fmicb.2018.00843

Ley, R. E., Hamady, M., Lozupone, C*., et al.* (2008). Evolution of mammals and their gut microbes.
Science, 320(5883), 1647-1651. doi:10.1126/science.1155725

Li, H., Li, T. T., Beasley, D. E*., et al.* (2016). Diet diversity is associated with beta but not alpha
diversity of pika gut microbiota. *Frontiers in Microbiology, 7*. doi:10.3389/fmicb.2016.01169

Linnenbrink, M., Wang, J., Hardouin, E. A*., et al.* (2013). The role of biogeography in shaping
diversity of the intestinal microbiota in house mice. *Molecular Ecology, 22*(7), 1904-1916.
doi:10.1111/mec.12206

Loke, P., Lim, Y. A. L. (2015). Helminths and the microbiota: parts of the hygiene hypothesis.
Parasite Immunology, 37(6), 314-323. doi:10.1111/pim.12193

Lopetuso, L. R., Scaldaferri, F., Petito, V*., et al.* (2013). Commensal Clostridia: leading players in the
maintenance of gut homeostasis. *Gut Pathogens, 5*. doi:10.1186/1757-4749-5-23

Magoc, T.,Salzberg, S. L. (2011). FLASH: fast length adjustment of short reads to improve genome
assemblies. *Bioinformatics, 27*(21), 2957-2963. doi:10.1093/bioinformatics/btr507

Mahe, F., Rognes, T., Quince, C*., et al.* (2014). Swarm: robust and fast clustering method for
amplicon-based studies. *Peerj, 2*. doi:10.7717/peerj.593

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal, 17, 10-12. doi:10.14806/ej.17.1.200

Maurice, C. F., Knowles, S. C. L., Ladau, J*., et al.* (2015). Marked seasonal variation in the wild
mouse gut microbiota. *Isme Journal, 9*(11), 2423-2434. doi:10.1038/ismej.2015.53

McDonald, J.E., Marchesi, J.R., Koskella, B. (2020). Application of ecological and evolutionary
theory to microbiome community dynamics across systems. *Proceedings of the Royal Society B-*
Biological Sciences 287. doi:10.1098/rspb.2020.2886

McKee, C.D., Bai, Y., Webb, C.T. et al. (2021). Bats are key hosts in the radiation of mammal-
associated *Bartonella* bacteria. *Infection Genetics and Evolution* 89.
doi:10.1016/j.meegid.2021.104719

McKnight, D. T., Huerlimann, R., Bower, D. S*., et al.* (2019). Methods for normalizing microbiome
data: An ecological perspective. *Methods in Ecology and Evolution, 10*(3), 389-400.
doi:10.1111/2041-210x.13115

McMurdie, P. J.,Holmes, S. (2013). phyloseq: an R package for reproducible interactive analysis and
graphics of microbiome census data. *Plos One, 8*(4), e61217. doi:10.1371/journal.pone.0061217

Mills, J. N., Childs, J., Ksiazek, T. G*., et al.* (1995). *Methods for trapping and sampling small*
mammals for virologic testing. Atlanta: Centers for Disease Control and Prevention.

Montgomery, S.S.J., Montgomery, W.I. (1989). Spatial and temporal variation in the infracommunity
structure of helminths of *Apodemus sylvaticus* (Rodentia: Muridae). *Parasitology* 98,145–150

Moran, N. A., Ochman, H.,Hammer, T. J. (2019). Evolutionary and ecological consequences of gut
microbial communities. *Annual Review of Ecology Evolution Systematics, 50*(1), 451-475.
doi:10.1146/annurev-ecolsys-110617-062453

Morgan, M. (2021). Dirichlet-multinomial mixture model machine learning for microbiome data.

Mutapi, F. (2015). The gut microbiome in the helminth infected host. *Trends in Parasitology, 31*(9),
405-406. doi:10.1016/j.pt.2015.06.003

Oksanen, J., Blanchet, F. G., Friendly, M*., et al.* (2020). vegan: Community Ecology Package.

Pascoe, E. L., Hauffe, H. C., Marchesi, J. R*., et al.* (2017). Network analysis of gut microbiota
literature: an overview of the research landscape in non-human animal studies. *Isme Journal,*
11(12), 2644-2651. doi:10.1038/ismej.2017.133
Peachey, L. E., Jenkins, T. P.,Cantacessi, C. (2017). This gut ain't big enough for both of us. Or is it?
Helminth-microbiota interactions in veterinary species. *Trends in Parasitology, 33*(8), 619-632.
doi:10.1016/j.pt.2017.04.004
Pfeiffer, J.K., Sonnenburg, J.L. (2011). The intestinal microbiota and viral susceptibility. *Frontiers in*
Microbiology 2. doi:10.3389/fmicb.2011.00092
Reese, A. T.,Dunn, R. R. (2018). Drivers of microbiome biodiversity: A review of general rules, feces,
and ignorance. *Mbio, 9*(4). doi:10.1128/mBio.01294-18
Revell, L. J. (2012). phytools: an R package for phylogenetic comparative biology (and other things).
Methods in Ecology and Evolution, 3(2), 217-223. doi:10.1111/j.2041-210X.2011.00169.x
Reynolds, L. A., Smith, K. A., Filbey, K. J*., et al.* (2014). Commensal-pathogen interactions in the
intestinal tract Lactobacilli promote infection with, and are promoted by, helminth parasites. *Gut*
Microbes, 5(4), 522-532. doi:10.4161/gmic.32155
Ribas Salvador, A., Guivier, E., Chaval, Y*., et al.* (2011). Concomitant influence of helminth infection
and landscape on the distribution of Puumala hantavirus in its reservoir, *Myodes glareolus. BMC*
Microbiology, 11(30), 1-13. doi:http://www.biomedcentral.com/1471-2180/11/30
Rinninella, E., Raoul, P., Cintoni, M*., et al.* (2019). What is the healthy gut microbiota composition?
A changing ecosystem across age, environment, diet, and diseases. *Microorganisms, 7*(1).
doi:10.3390/microorganisms7010014
Rodriguez-Daza, M. C., Roquim, M., Dudonne, S*., et al.* (2020). Berry polyphenols and fibers
modulate distinct microbial metabolic functions and gut microbiota enterotype-like clustering in
obese mice. *Frontiers in Microbiology, 11*. doi:10.3389/fmicb.2020.02032
Rolhion, N.,Chassaing, B. (2016). When pathogenic bacteria meet the intestinal microbiota.
Philosophical Transactions of the Royal Society B-Biological Sciences, 371(1707).
doi:10.1098/rstb.2015.0504
Rosenfeld, C. S. (2017). Gut dysbiosis in animals due to environmental chemical exposures. *Frontiers*
in Cellular and Infection Microbiology, 7. doi:10.3389/fcimb.2017.00396
Rosshart, S.P., Vassallo, B.G., Angeletti, D. et al. (2017). Wild mouse gut microbiota promotes host
fitness and improves disease resistance. Cell 171,1015. doi:10.1016/j.cell.2017.09.016
Round, J. L.,Mazmanian, S. K. (2009). The gut microbiota shapes intestinal immune responses during
health and disease. *Nature Reviews Immunology, 9*(5), 313-323. doi:10.1038/nri2515
Sabey, K. A., Song, S. J., Jolles, A*., et al.* (2021). Coinfection and infection duration shape how
pathogens affect the African buffalo gut microbiota. *Isme Journal, 15*(5), 1359-1371.
doi:10.1038/s41396-020-00855-0
Salter, S. J., Cox, M. J., Turek, E. M*., et al.* (2014). Reagent and laboratory contamination can
critically impact sequence-based microbiome analyses. *Bmc Biology, 12*, 87. doi:10.1186/s12915-
014-0087-z
Seguel, M.,Gottdenker, N. (2017). The diversity and impact of hookworm infections in wildlife.
International Journal for Parasitology-Parasites and Wildlife, 6(3), 177-194.
doi:10.1016/j.ijppaw.2017.03.007
Stothart, M.R., Palme, R., Newman, A.E.M. (2019). It's what's on the inside that counts: stress
physiology and the bacterial microbiome of a wild urban mammal. *Proceedings of the Royal*
Society B-Biological Sciences 286. doi:10.1098/rspb.2019.2111
Suzuki, T. A.,Nachman, M. W. (2016). Spatial heterogeneity of gut microbial composition
along the gastrointestinal tract in natural populations of house mice. *Plos One, 11*(9).
doi:10.1371/journal.pone.0163720
team, R. c. (2020). A language and environment for statistical computing.: Austria, Vienna.
Telfer, S., Lambin, X., Birtles, R*., et al.* (2010). Species interactions in a parasite community drive
infection risk in a wildlife population. *Science, 330*(6001), 243-246. doi:10.1126/science.1190333
Trevelline, B. K., Fontaine, S. S., Hartup, B. K*., et al.* (2019). Conservation biology needs a microbial
renaissance: a call for the consideration of host-associated microbiota in wildlife management

practices. *Proceedings of the Royal Society B-Biological Sciences, 286*(1895).
doi:10.1098/rspb.2018.2448
Vujkovic-Cvijin, I., Sklar, J., Jiang, L.*, et al.* (2020). Host variables confound gut microbiota studies
of human disease. *Nature, 587*(7834), 448-454. doi:10.1038/s41586-020-2881-9
Wang, J., Linnenbrink, M., Kunzel, S.*, et al.* (2014). Dietary history contributes to enterotype-like
clustering and functional metagenomic content in the intestinal microbiome of wild mice.
Proceedings of the National Academy of Sciences of the United States of America, 111(26), E2703-
E2710. doi:10.1073/pnas.1402342111
Winkler, E.S., Thackray, L.B. (2019). A long-distance relationship: the commensal gut microbiota and
systemic viruses. *Current Opinion in Virology* 37,44-51. doi:10.1016/j.coviro.2019.05.009
Wolf, J. F., Kriss, K. D., MacAulay, K. M.*, et al.* (2021). Gut microbiome composition predicts
summer core range size in two divergent ungulates. *FEMS Microbiology Ecology, 97*(5).
doi:10.1093/femsec/fiab048
Worsley, S. F., Davies, C. S., Mannarelli, M. E.*, et al.* (2021). Gut microbiome composition, not alpha
diversity, is associated with survival in a natural vertebrate population. *Animal Microbiome, 3*(1).
doi:10.1186/s42523-021-00149-6
Zaiss, M.M., Rapin, A., Lebon, L. et al. (2015). The intestinal microbiota contributes to the ability of
helminths to modulate allergic inflammation. *Immunity* 43,998.
doi:1010.10.1016/j.immuni.2015.09.012
Zeidner, N. S., Schneider, B. S., Dolan, M. C.*, et al.* (2001). An analysis of spirochete load, strain, and
pathology in a model of tick-transmitted Lyme borreliosis. *Vector-Borne and Zoonotic Diseases,*
1(1), 35-+. doi:10.1089/153036601750137642
Zhai, S., Qin, S., Li, L.*, et al.* (2019). Dietary butyrate suppresses inflammation through modulating
gut microbiota in high-fat diet-fed mice. *FEMS Microbiology Letters, 366*(13).
doi:10.1093/femsle/fnz153
Zheng, D.P., Liwinski, T., Elinav, E. (2020). Interaction between microbiota and immunity in health
and disease. *Cell Research* 30,492-506. doi:10.1038/s41422-020-0332-7
987