Small scale geographical variation of three-way relationships between gut microbiota,

helminth assemblages and bacterial infections in wild rodent populations

Marie Bouilloud[1,2], Maxime Galan[1], Adélaïde Dubois[1], Christophe Diagne[3], Philippe Marianneau[4],

Benjamin Roche*[2], Nathalie Charbonnel*[1]*

[1]CBGP, INRAE, CIRAD, Institut Agro, IRD, University of Montpellier, Montpellier, France

[2]MIVEGEC, IRD, CNRS, University of Montpellier, Montpellier, France

[3]CBGP, IRD, INRAE, CIRAD, Institut Agro, University of Montpellier, Montpellier, France

[4]INRAE, Lyon, France

*Corresponding author

*Equal contribution of the coauthors

Orcid

Marie Bouilloud : marie.bouilloud@ird.fr, 0000-0001-8740-3205
17

18 Maxime Galan : maxime.galan@inrae.fr, 0000-0001-6981-5732

20 Christophe Diagne : christophe.diagne@ird.fr, 0000-0002-6406-1270

22 Philippe Marianneau : philippe.marianneau@inrae.fr

24 Benjamin Roche : roche.ben@gmail.com, 0000-0001-7975-4232

26 Nathalie Charbonnel : nathalie.charbonnel@inrae.fr, 0000-0002-7907-6539

Abstract

Background

Despite its central role in host fitness, the gut microbiota may differ greatly between individuals. This variability is often mediated by environmental or host factors such as diet, genetics, and infections. Recently, a particular attention has been given to the interactions between gut bacteriome and helminths, as these latter could affect host susceptibility to other infections. Further studies are still required to better understand the three-way interactions between gut bacteriome, helminths and other parasites, especially because previous findings have been very variable, even for comparable host-parasite systems.

Methods

In our study, we used the V4 region of the 16S rRNA gene to assess the variability of gut bacteriome diversity and composition in wild populations of a small mammal, the bank vole *Myodes glareolus*. Four sites were sampled at a fine geographical scale (100 km) along a North-South transect in Eastern France. We applied analyses and concepts of community and microbial ecology to evaluate the interactions between the gut bacteriome, the gastro-intestinal helminths and the pathogenic bacteria detected in the spleen.

Results

Regarding the gut bacteriome composition and diversity among bank voles, we identified important variations that were mainly explained by sampling localities and reflected the North/South sampling transect. In addition, we detected two main enterotypes, that might correspond to contrasted diets. We found geographic variations of the gut bacteriome richness and Firmicutes/Bacteroidetes ratio, both correlating positively with body mass index. We found positive correlations between the specific richness of the gut bacteriome and of the helminth community, as well as between the composition of these two communities, even when accounting for the influence of geographical distance. For the pathogenic bacteria community, no broad patterns have been detected. The helminths *Aonchothecha murissylvatici, Heligmosomum mixtum* and the bacteria *Bartonella* sp were the main taxa associated with the whole gut bacteriome composition. Besides, changes in relative abundance of particular gut bacteriome taxa were specifically associated with other helminths (*Mastophorus muris*, *Catenotaenia henttoneni*, *Paranoplocephala omphaloides* and *Trichuris arvicolae*) or pathogenic bacteria. Especially, infections with *Neoehrlichia mikurensis*, *Orientia* sp or *Rickettsia* sp were associated with lower relative abundance of the family

Erysipelotrichaceae (Firmicutes), while coinfections with higher number of bacterial infections

were associated with lower relative abundance of a Bacteroidales family (Bacteroidetes).

Conclusions

These results emphasize complex interlinkages between gut bacteriome and infections in wild

animal populations. They remain difficult to generalize due to the strong impact of environment,

even at fine geographical scales, on these interactions. Abiotic features, as well as small mammal

community composition and within host parasite coinfections, should now be considered to better

understand the spatial variations observed in the relationships between gut bacteriome, gastro-

intestinal helminths and bacterial infections.

Keywords : bank voles, co-infections, interactions, microbial community ecology, zoonoses

Introduction

Vertebrate gut microbiome plays key roles in host fitness, through functions including, among others, nutrient acquisition, immunity and defence against infectious exogenous agents (hereafter called 'parasites' and including micro- and macroparasites) or proliferating indigenous organisms (Belkaid & Hand, 2014; Kamada *et al.*, 2013; Round & Mazmanian, 2009). Nonetheless, the gut microbiome may differ greatly in natural environments between individuals, populations and species (Vujkovic-Cvijin *et al.*, 2020). This variability reflects hosts intrinsic factors (notably phylogeny, genetics or vertical transmission from mother to offspring) as well as extrinsic features (acquisition of microorganisms from the environment, potentially through diet Ley *et al.*, 2008; Moran *et al.*, 2019). It might also reveal disruption of host-associated gut microbiome (termed "dysbiosis") caused by environmental factors, among which are anthropogenic pressures (e.g., chemical exposures, Rosenfeld, 2017) or parasite infections (Trevelline *et al.*, 2019).

Today, it is obvious that understanding the relationships between gut microbiome and parasite transmission is crucial regarding their potential impacts on human and animal health (Clemente *et al.*, 2012). To this extent, vertebrate microbiome studies have placed particular emphasis on the gut bacterial microbiota (called hereafter 'gut bacteriome') and their interactions with gastro-intestinal helminth parasites. On one hand, the gut bacteriome may act as an innate immune barrier to intestinal infections and influence the colonisation and growth of eukaryotic parasites, including helminths, through competitive metabolic interactions or induction of host immune responses (Leung *et al.*, 2018). On the other hand, helminth infections may also affect, directly or indirectly through similar ecological and/or evolutionary processes, the composition of the gut bacteriome, via physical contact, competition for resources or host immunoregulation (see Kreisinger *et al.*, 2015). Helminths and the gut bacteriome interactions may thus lead to positive and negative associations (Loke & Lim, 2015), with potentially local but also systemic physiological changes affecting host health. For example, helminth infections can lead to malnutrition and weight loss, through the dysfunction of microbial metabolism that could result from negative impacts on fermentative gut bacteria (Leung *et al.*, 2018). Besides, some helminth infections favour gut bacteria that produce short-chain fatty acids from dietary fiber. These metabolites circulate throughout the body and are important regulators of host physiology (glucose and fat metabolism) and immune system (Honda & Littman, 2016; Kim, 2021). Interactions between these helminths and gut bacteria may here increase the host anti-inflammatory and regulatory T cell suppressor responses, what may in turn affect host susceptibility to other infections as well as the outcomes of infections (Glendinning *et al.*, 2014).

Besides, the gut microbiota may also influence microparasite infections, through their immune

function against pathogenic bacteria colonization and their role in maintaining the intestinal

epithelium integrity (Khosravi & Mazmanian, 2013). However, up to now, the systemic impact of

gut bacteriome on microparasite infections and vice-versa, represents a fundamental knowledge gap

in wild animal populations (Pascoe *et al.*, 2017; Rolhion & Chassaing, 2016).

Moreover, the three-way interactions between host's gut bacteriome, gastro-intestinal helminths and

microparasites have been scarcely investigated in a single system, despite clearly becoming pivotal

in disease ecology. Yet, the growing interest on gut bacteriome/parasitism relationships in recent

literature (Johnson *et al.*, 2015) highlights the critical need for further empirical works, given the

relatively low concordance of findings between previous studies – even for comparable host-

parasite systems. Up to now, most of the research on this topic have been conducted on model

species in laboratory settings. Although experiments under controlled conditions may help

emphasizing general patterns and deciphering the mechanisms underlying these interactions

between gut bacteriome and parasites in vertebrates (Pascoe *et al.*, 2017), they also have inherent

limitations. On the one hand, they only included a restricted number of targeted parasites (usually

helminths and/or microparasites), omitting the potential effects of species interactions between and

within parasite communities at the intra-host level (Telfer *et al.*, 2010). Yet, co-infections are now

recognized to be the rule in most hosts in natural environments, with wild animals that may carry

simultaneously a large number of parasites (Hoarau *et al.*, 2020). On the other hand, they are

intrinsically unable to include – and then capture the complexity of – the environmental conditions

as drivers of the composition of gut bacteriome and of the exposure or sensibility to these latter

(Adair & Douglas, 2017). From there, studies in natural contexts therefore deserve strong

consideration because these extrinsic factors may impact deeply the relationships between gut

bacteriome and parasitism.

Here, we strived to bridge these gaps by assessing the variability of gut bacteriome diversity and

composition in wild populations of the bank vole *Myodes glareolus*, a small mammal reservoir of a

large number of zoonotic agents (e.g., Abbate *et al.*, In_revision). We studied the relationships

between its gut bacteriome, parasite infracommunities – focusing on gastro-intestinal helminths and

pathogenic bacterial infections –, and host and environmental factors that may either influence or

indicate the health status (e.g., proxies such as the body mass index (BMI)). The current study

therefore addressed two main questions: (1) how is the structure (composition and diversity) of the

gut bacteriome influenced by host and environmental factors? (2) does the structure of gut

bacteriome reflect interactions with gastro-intestinal helminth and pathogenic bacterial

communities? For this purpose, we *(i)* characterized the composition of each intra-host community (gut bacteriome, pathogenic bacteria, gastro-intestinal helminths) across bank vole populations, *(ii)* investigated the alpha and beta diversity of each intra-host community, and the pairwise relationships between these communities for each diversity metrics, and *(iii)* assessed if the structure of gut bacteriome was mediated either by factors similar to those influencing the structure of gastro-intestinal helminths and pathogenic bacteria community, or by potential interactions between intra-host communities.

Material and methods

Sample collection

Bank voles (*Myodes glareolus*) were trapped between June and September 2014 in forests located in four French localities (Table 1, Figure S1) distributed along a North-South transect in Eastern France, and separated by 40 to 120 km from one another. The standardized trapping protocol used here was described in details in Dubois et al. (2018). Previous studies have revealed that these populations exhibit some variability in bank vole susceptibility to zoonotic viral infections and immune gene expression (Dubois *et al.*, 2017; Dubois *et al.*, 2018).

Rodents were euthanized using isofluorane and cervical dislocation, as recommended by Mills (1995). Morphological measures were taken. Age groups (juveniles and adults) were defined according to body mass and sexual maturity. This latter was inferred using testes length and position, and seminal vesicle development for males, or uterus size for females. Body condition was estimated using the body mass index (BMI = weight/length^2). The digestive tract and the spleen were removed and stored respectively in 96% ethanol and RNA later solution (-20°C).

Ethical statements: Animal capture and handling have been conducted according to the French and European regulations on care and protection of laboratory animals (French Law 2001-486 issued on June 6, 2001 and Directive 2010/63/EU issued on September 22, 2010). The CBGP laboratory has approval (D-34-169-003) from the Departmental Direction of Population Protection (DDPP, Hérault, France), for the sampling of rodents and the storage and use of their tissues.

Table 1- Prevalence of potentially pathogenic bacteria and gastro-intestinal helminths for each sampling locality. *N* is the number of bank voles analysed. N_T represents the number of individuals with data available for the three intra-host communities (gut bacteriome, pathogenic bacteria and gastro-intestinal helminths). N_{GB}, N_{PB} and N_{GIH} respectively represent the number of individuals with data available for each of these intra-host communities. '*Uninfected*' corresponds to the number of uninfected bank voles for a given intra-host community. '*Co-infection*' corresponds to the number of bank voles infected with at least two parasites of a given intra-host community. The color gradient illustrates variations in prevalence (0% = light red to 100% = dark red).

	Total	Sampling localities			
		Mont-sous-Vaudrey	Chaux-des-Crotenay	Chatillon	Cormaranche
Total					
N_T	124	22	34	36	32
Gut bacteriome					
N_{Gb}	161	37	42	41	41
Pathogenic bacteria					
N_{Pb}	138	22	34	38	37
Uninfected$_{Pb}$	11	3	1	3	4
Co-infection$_{Pb}$	67	10	21	16	20
Bartonella sp	0.73	0.80	0.81	0.69	0.66
Mycoplasma haemomuris	0.36	0.28	0.39	0.36	0.39
Anaplasma phagocytophilum	0.14	0.04	0.33	0.08	0.11
Neoehrlichia mikurensis	0.13	0.16	0.00	0.08	0.29
Orientia tsutsugamushi	0.08	0.00	0.08	0.21	0.00
Rickettsia sp	0.05	0.00	0.08	0.05	0.05
Spiroplasma sp	0.04	0.04	0.00	0.00	0.11
Borrelia miyamotoi	0.02	0.04	0.00	0.03	0.03
Borreliella afzelii	0.02	0.08	0.00	0.00	0.03
Mycoplasma coccoides	0.01	0.04	0.00	0.00	0.00
Gastro-intestinal helminths					
N_{Gih}	153	37	42	39	35
Uninfected$_{Gih}$	29	16	4	6	3
Co-infection$_{Gih}$	64	2	22	13	27
Heligmosomoides glareoli	0.48	0.49	0.36	0.54	0.54
Heligmosomum mixtum	0.41	0.00	0.76	0.00	0.86
Aonchotheca murissylvatici	0.35	0.03	0.26	0.59	0.54
Catenotaenia henttoneni	0.10	0.03	0.19	0.10	0.06
Paranoplocephala omphaloides	0.06	0.00	0.14	0.00	0.09
Arostrilepis horrida	0.03	0.00	0.00	0.00	0.14
Mastophorus muris	0.01	0.05	0.00	0.00	0.00
Trichuris arvicolae	0.01	0.03	0.00	0.00	0.00

Characterisation of intra-host bacterial and helminth communities

Gut bacteriome

We first characterized the gut bacteriome of bank voles. We focused on the colon as rodent gut microbiome exhibits the highest level of bacterial diversity in this part of the digestive tract (Suzuki & Nachman, 2016). DNA was extracted from a 5 mm piece of colon (taken about 1 cm far from the caecum) of each bank vole using the ZymoBiomics 96 DNA Kit (Zymo) following the manufacturer's instructions. We amplified a 251-bp portion of the V4 region of the 16S rRNA gene (16S-V4F [GTGCCAGCMGCCGCGGTAA] and 16S-V4R [GGACTACHVGGGTWTCTAATCC]), following Kozich et al. (2013) and as described in Galan et al. (2016). Samples were multiplexed using dual-indexes (index i5 in the forward primer and index i7 in the reverse primer). Negative controls for extraction (whole reagents without DNA), for PCR (PCR mix without DNA), and for indexing (wells without reagents corresponding to particular dual-indexes combinations). All DNA extractions were analysed twice using two independent technical replicates of amplicon libraries. PCR products were pooled, migrated and excised on a low melting agarose gel (1.25%) then purified using the NucleoSpin Gel and PCR Clean-Up kit (Macherey-Nagel) and quantified using the KAPA library quantification kit (KAPA Biosystems) standardized to 4nM by qPCR spectrophotometry (assay). Sequencing was performed on a 251-bp paired-end Illumina MiSeq run. The raw sequence reads (.fastq format) have been deposited in the Zenodo Repository.

Sequence data were processed as described in Galan et al. (2016) using the pipelines implemented in FROGS (Find Rapidly OTU with Galaxy Solution, Escudié *et al.*, 2018). Briefly, the paired-end sequences were trimmed with CUTADAPT (Martin, 2011), merged with FLASH (MAGOC & SALZBERG, 2011), and clustered into operational taxonomy OTU units at 97% identity using the SWARM algorithm (Mahe *et al.*, 2014) executed with aggregation parameter distance *d*=1 and a second pass performed on the seeds of previous clusters with *d*=3. Putative chimeras were removed using VSEARCH tools with de novo VUCHIME and the cross-validation method. Taxonomy was assigned with BLASTN+ (Camacho *et al.*, 2009) using the SILVA SSU Ref NR 119 database as a reference (http://www.arb175silva.de/projects/ssu-ref-nr/). Filtering for false positives was carried out as proposed by Galan et al. (2016). In short, we discarded positive results associated with sequence counts below two OTU-specific thresholds, which checked respectively for cross-contamination between samples (using the negative controls for extraction and PCR) and incorrect assignment due to the generation of mixed clusters on the flow cell during Illumina sequencing, using a false index-pairing rate for each PCR product of 0.02%, based on estimates from Galan et al. (2016). For each sample, only OTUs found in the two technical replicates were considered as positive, and OTUs found in only one of the two replicates were removed. The number of sequences obtained for each technical replicate from a sample were summed.

Lastly, we discarded OTUs and samples containing less than 500 reads in the dataset, as well as

OTUs considered to be contaminants, following (Salter *et al.*, 2014). OTUs number of reads were

finally normalised to proportional abundance within each rodent (McKnight *et al.*, 2019). Two

taxonomic ranks were considered for further analyses, phylum and family.

Further description and analyses of bacterial communities were performed using the PHYLOSEQ

package (McMurdie & Holmes, 2013). We looked for enterotypes, *i.e.* distinct community

composition types of gut bacteriome, as found in humans (Arumugam *et al.*, 2011; Holmes *et al.*,

2012), using the Dirichlet Multinomial Mixtures DMM (Morgan, 2021). We next analysed the

Firmicutes /Bacteroidetes (F/B) log-ratio, as it is often used as a proxy of health or metabolism in

humans and mice, see refs in (Lavrinienko *et al.*, 2018). We calculated this ratio with the

MICROBIOME package (Lahti & Shetty, 2017) and we tested the influence of individual

characteristics (age class, gender, BMI) and localities on this log-ratio using generalized linear

models.

Pathogenic bacteria

We next described the presence/absence of pathogenic and zoonotic bacteria from the spleen of

each bank vole. This lymphoid organ filters microbial cells in mammals and as such, enables to

recover recent infections (Abbate *et al.*, In_revision; Diagne *et al.*, 2017). Molecular protocols,

bioinformatics pipelines and data filtering were similar to those described above (gut bacteriome),

except for the DNA extraction from splenic tissue using DNeasy 96 Tissue Kit (Qiagen). The

potential pathogenicity of each bacterial OTU was assessed based on published literature and on the

Gideon database (https://www.gideononline.com/). Opportunistic pathogens (*i.e.* commensal agents

in healthy hosts, that become pathogenic when the balance of the immune system is disrupted) were

discarded from the dataset. Only the information of the presence / absence of pathogenic/zoonotic

OTUs was considered.

Gastro-intestinal helminths

For each bank vole, helminths were carefully extracted and counted from the different sections of

the digestive tract (stomach, small intestine, large intestine and caecum), and classified by

morphotype then stored in 95% ethanol for further accurate identification. The latter was based on

unambiguous morphological criteria using conventional microscopy and generalist identification

keys or specific literature when available (Anderson *et al.*, 2009; Khalil *et al.*, 1994; Ribas Salvador

et al., 2011).

Statistical analyses

All statistical analyses were implemented in R v4.0.3 (team, 2020). For more convenience, gut bacteriome, pathogenic bacteria and gastro-intestinal helminths were further described as 'intra-host communities'.

Assessment of the alpha diversity within intra-host communities

First, we characterized the alpha diversity, *i.e.* the diversity within intra-host communities (the gut bacteriome, pathogenic bacteria and gastro-intestinal helminth communities) using two metrics, the specific richness (*i.e.* number of taxa within the host individual) and the Shannon index as recommended in (Haegeman *et al.*, 2013).

Second, we tested the influence of rodent characteristics (age class, gender, BMI), localities and the two-ways interactions between these factors, on the alpha diversity of the three intra-host communities using generalized linear models (GLMs). We considered a negative binomial error distribution for the specific richness and a gamma distribution for the Shannon index. Best model selection was performed considering models with all possible combinations of factors and the DREDGE function of the MuMIn package. The best model was selected using the Akaike information criterion corrected for small sample size AICc, (Johnson & Omland, 2004). We assessed the effect of each factor in the best model with the ΔAICc index. When the factor locality was significant, Tukey's post-hoc tests were applied to evaluate pairwise differences between localities, using the MULTCOMP package (Hothorn *et al.*, 2008). Residuals were checked to graphically to ensure that all assumptions regarding normality, independence and the homogeneity of variance were satisfied.

Third, we analysed whether the alpha diversity of each intra-host community was influenced by the alpha diversity of the two other ones. We performed GLMs and followed a similar model selection process than the one explained above.

Assessment of the beta diversity of intra-host communities

We analysed the beta diversity, *i.e.,* the dissimilarity between host individuals in their intra-host communities (gut and pathogenic bacteria, and helminth communities) composition, using Bray-Curtis distances. We considered the relative abundance of OTUs for the gut bacteriome (phylum and family), the occurrence (presence/absence) of pathogenic bacteria and the abundance of gastro-intestinal helminths.

To assess the impact of the spatial dimension on beta diversity, we performed Mantel tests using
Pearson correlation (10,000 permutations) and evaluated the influence of geographic distances on
the dissimilarities in intra-host gut, pathogenic bacterial and helminth communities.
We then analysed the factors shaping the dissimilarities in intra-host community composition using
several functions of the VEGAN package (Oksanen *et al.*, 2020). Rodents that were not infected
(pathogenic bacteria or helminths) could not be included in these analyses. Distance-based
redundancy analyses (db-RDA) were performed to analyse the effect of individual explanatory
factors (age class, gender, BMI) and sampling localities on dissimilarities in intra-host community
composition, using the CAPSCALE function, followed by permutational multivariate analyses of
variance (PERMANOVA). We selected the best model, *i.e.*, the most parsimonious one, using the
ORDIR2STEP function (*P*-value adjusted and R^2 adjusted). For each factor, we evaluated the intra-
group dispersion using the BETADISPER function as PERMANOVA analyses are sensitive to
differences in dispersion among groups.
As this study focused on the variability of the gut bacteriome, we also used DESEQ2 package (Love
et al., 2014), that is designed to perform differential analysis of high-dimensional count data, to
identify the changes in bacteria taxa (phyla or families) that best explained gut bacteriome
dissimilarities between individuals and localities. We performed GLMs with negative binomial
error (NBINOMWALDTEST method) and significant differences were obtained after Benjamini &
Hochberg corrections. They were visualised using the METACODER package (Foster *et al.*, 2017).
302
Interactions between intra-host gut bacteria, pathogenic bacteria and gastro-intestinal helminth
communities
We applied partial Mantel tests using MULTI.MANTEL (phytools package Revell, 2012) to analyse
the correlation between two matrices of dissimilarities (corresponding to two different
communities), while controlling for the effect of a third dissimilarity matrix (third community).
Then, we used db-RDA to analyse more deeply the relationships between the gut bacteriome and
the pathogenic (bacteria and helminths) communities. We included the alpha diversity indices
(richness specific and Shannon index) and the infectious status (presence / abundance) of pathogens
with prevalence greater than 10% in at least one locality as explanatory variables in these analyses.
We selected the best model using the ORDIR2STEP method. As described above, we used DESEQ2 to
determine the phyla and families of the gut bacteriome whose relative abundances changed with
significant explanatory variables.
315
316   **Results**

A total of 186 bank voles were trapped during the fieldwork campaign over the four targeted

localities. For technical reasons (e.g., poor sample preservation, missing data), we could describe

the three intra-host communities for 124 rodents only.

Composition of the intra host communities

Gut bacteriome

Once the filtration steps were applied, the gut bacteriome dataset included 161 bank voles. We

detected 10 phyla and 61 families of bacteria. At the phylum level, we found six predominant taxa

that represented 99% of the gut bacteria relative abundance (Figures S2): *Firmicutes* (41% of

reads), *Proteobacteria* (20% of reads), *Epsilonbacteraeota* (12% of reads)*, Actinobacteria* (12% of

reads), *Bacteroidetes* (11% of reads) and *Spirochaetes* (3% of reads). At the family level, 11

families represented 93% of the relative abundance of the gut bacteriome (Figure 1A):

Lactobacillaceae (21% of reads), *Lachnospiraceae* (11% of reads), *Ruminococcaceae* (4% of

reads), *Clostridiales* (3% of reads) of the *Firmicutes* phylum, *Desulfovibrionaceae* (*Proteobacteria*

phylum, 19% of reads), *Eggerthellaceae* (*Actinobacteria* phylum, 12% of reads), *Muribaculaceae*

and *Bacteroidales F1* (*Bacteroidetes* phylum*,* 8% and 1% of reads respectively), *Helicobacteraceae*

(*Epsilonbacteraeota,* 12% of reads) and *Spirochaetaceae* (*Spirochaetes* phylum*,* 3% of reads).

We distinguished two enterotypes from the DMM approach. One (enterotype 1) was mainly

composed of the families *Helicobacteraceae*, *Lachnospiraceae*, *Muribaculaceae and*

Desulfovibrionaceae, while the other (enterotype 2) mainly included *Lactobacillaceae*,

Desulfovibrionaceae and *Eggerthellaceae* (Figure 1B).

Lastly, we found that the *Firmicutes / Bateroidetes* ratio varied significantly between localities

(Figure 1C). Overall, northern localities exhibited lower F/B ratio than southern ones, with all

pairwise comparisons being significant except Chatillon *versus* Chaux-des-Crotenay. Individual

characteristics did not influence this ratio (Table S1).

Figure 1- Composition of the intestinal bacteriome. A) The Bar plot shows the individual variations of 11 bacterial families (F= Unknown family) belonging to 6 phyla and representing 93% of the total composition. Individuals (bars) are grouped by sampling localities, which are ordered from north to south. Each color represents a taxon. B) The composition of the two enterotypes identified using Dirichlet multinomial mixtures (DMMs), at family rank, is shown. Bacterial families are represented using the same colors as in A. and C) the ratio (Firmicutes / Bacteroidetes) is shown for each sampling locality. Box and whisker plots represent the median and interquartile values. Black dots correspond to the mean value, and colored dots correspond to individuals. Different letters indicate statistically significant differences at P < 0.05, with pairwise Tukey post hoc adjustments.

Pathogenic bacteria

Once the filtration steps were applied, the dataset corresponding to potentially pathogenic bacteria included 138 *M. glareolus*. Ten genus/species were considered as potentially pathogens. Four of them had prevalence levels greater than 10%: *Bartonella* sp *(73%), Mycoplasma haemomuris (36%), Anaplasma phagocytophilum (14%), Candidatus Neoehrlichia mikurensis (13%)* (Table 1). Most *M. glareolus* individuals were infected with at least one bacteria species (92%), and about half bank voles (49%) were infected with two or more bacteria species (Table 1).

Gastro-intestinal helminths

We detected eight species of gastro-intestinal helminths among the 153 *M. glareolus* analysed. Four helminth species exhibited prevalence levels higher than 10%: the nematodes *Heligmosomoides glareoli* (prevalence: 48%), *Heligmosomum mixtum* (prevalence: 41%) and the cestodes

Aonchotheca murissylvatici (prevalence: 35%) and *Catenotaenia henttoneni* (prevalence: 10%)

(Table 1). Note that the prevalence of *Paranoplocephala omphaloides* and *Arostrilepis horrida*

were greater than 10% in Chaux-des-Crotenay and Cormaranche. Most bank voles (85%) were

infected with at least one helminth species and almost half (42%) were infected with at least two

different helminth species (Table 1).

Factors influencing the alpha diversity within intra-host communities

Gut bacteriome

At the phylum level, none of the factors significantly explained the variations of specific richness of

the gut bacteria community between bank voles (Table S2A; Figure S3A). When considering

Shannon index, we found that the sampling locality had a significant effect on the alpha diversity of

the gut bacteriome (ΔAICc = 8.70, R^2_{GLM} = 0.09, Table S2A; Figure S3B). After post-hoc

corrections, Mont-sous-Vaudrey and Cormaranche were the only localities that differed

significantly from the others (P_{adj} < 0.001). Mont-sous-Vaudrey showed higher levels of diversity

than Cormaranche (β = -0.19; 95% CI = [-0.29, -0.10]). The other localities also exhibited lower

levels of alpha diversity than Mont-sous-Vaudrey (Chaux-des-Crotenay: β = -0.12; 95% CI = [-

0.22, -0.03]; P = 0.01; P_{adj} = 0.05; Chatillon: β = -0.11; 95% CI = [-0.21, -0.01]; P = 0.02; P_{adj} =

0.10) but these differences were not significant after correcting for multiple tests.

At the family level, the sampling locality had a significant effect, whatever the alpha diversity

metrics used (Specific richness: ΔAICc = 14.28; R^2_{GLM} = 0.21; Figure 2A; Shannon index: ΔAICc =

7.65; R^2_{GLM} = 0.08; Figure 2B; Table S2B). The locality Cormaranche exhibited a higher specific

richness than all other localities (Tukey post hoc test: Mont-sous-Vaudrey: β = 0.22; 95% CI =

[0.10, 0.30]; P_{adj} < 10^{-3}, Chaux-des-Crotenay: β = 0.19; P_{adj} < 10^{-3} and Chatillon: β =0.15; P_{adj} = 1.7

x 10^{-3}) but a lower level of diversity than Mont-sous-Vaudrey when correcting for rare taxa (Tukey

post hoc test, Shannon index: β = -0.76; 95% CI = [-1.2; -0.4]; P_{adj} = 2 x 10^{-3}, Figure 2B). Body

condition (BMI) was also found to have a significant effect, but only when considering specific

richness (ΔAICc=1.99; β = 0.18; 95% CI = [0.1, 0.3], P = 4 x 10^{-3}) – with higher values of BMI

associated with increasing species richness. All these results are detailed in Table S2B and Figure

S4A,D,G).

Pathogenic bacteria

Host gender was the only factor that significantly influenced variations in alpha diversity regardless

of the index considered (Specific richness: ΔAICc = 0.32; R^2_{GLM} =0.04; Shannon index: ΔAICc =

3.4; R^2_{GLM} =0.05; Figure 2C; Figure S4B,E,H). Males had a higher diversity of pathogenic bacteria

than females (Specific richness: $\beta = 0.22$; 95% CI = [0.03,042]; $P= 0.02$; Shannon index: $\beta = 0.17$;

95% CI = [0.03,0.32]; $P=0.02$). All these results are detailed in Table S2C.

Gastro-intestinal helminths

We found a significant effect of the locality on the alpha diversity of the gastro-intestinal helminth

community, whatever the metrics used (Specific richness: $\Delta AICc=24.53$, $R^2_{GLM} = 0.30$; Shannon

index: $\Delta AICc= 41.11$; $R^2_{GLM} = 0.30$; Table S2D). When considering the specific richness, voles

from Mont-sous-Vaudrey showed lower value than in all other localities (Tukey post hoc test,

Figure 2D, Table S2D). In addition, bank voles from Chatillon had lower specific richness than

Cormaranche (Tukey post hoc test, $\beta = 0.61$; $P_{adj} < 10^{-3}$, Figure 2D). When considering Shannon

index, we found that bank voles from Cormaranche exhibited significantly higher levels compared

with the other localities (Table S2D). Moreover, bank voles from Chaux–des-Crotenay exhibited

higher levels of Shannon index than those from Mont-sous-Vaudrey ($\beta = 0.38$, $P_{adj} < 10^{-3}$).

We also found that the age of bank voles significantly influenced the specific richness of the

helminth community, with adults being infected by a higher diversity of helminths than juveniles

(Specific richness: $\Delta AICc=0.49$; $\beta = 0.31$; 95% CI = [0.03, 0.61]; $P = 0.03$; Table S2D, Figure

S4C,F,I).

Relationships between the alpha diversity of the three intra-host communities

Gut bacteriome as the dependant variable. For the species richness, a significant relationship was

only found at the family level between the gut bacteriome and the helminth community ($\Delta AICc =$

8.43; $P < 10^{-3}$), with a more diverse gut bacteriome associated with a greater number of helminth

species infecting bank voles ($\beta = 0.07$; 95% CI = [0.03, 0.10]; $R^2 = 0.10$; Figure 2E; Table S2E,F).

For the Shannon index, we did not find any effect between the alpha diversity estimates of each

intra-host community, whatever the taxonomic rank considered (Figure 2E; Tables S2E,F).

Pathogenic bacteria as the dependant variable. For both species richness and Shannon index, we

found a positive relationship with, respectively, the specific richness ($\Delta AICc = 1.02$; $P < 10^{-3}$)

($\Delta AICc = 1.02$; $P < 10^{-3;}$ Figure 2E) and the Shannon index ($\Delta AICc = 3.47$; $\beta=0.19$; $P = 0.04$,

$R^2=0.02$; Figure 2E; Table S2E,F) of the gastro-intestinal helminth community.

Gastro-intestinal helminths as the dependant variable. For the specific richness, we found a

significant effect of the specific richness of the gut bacteriome (Family level only) and of

pathogenic bacteria ($R^2 = 0.14$; gut bacteriome $\Delta AICc = 6.29$; $P =0.02$; $\beta = 0.06$; pathogenic

bacteria $\Delta AICc = 1.82$; $P = 0.03$; $\beta = 0.17$; Figure 2E, Table S2F). For Shannon index, only the alpa

diversity of pathogenic bacteria remained significant (ΔAICc = 3.29; β = 0.23; P = 0.02; Figure 2E,

Table S2F).

434
Figure 2- Variations of alpha diversity between localities, for the intestinal bacteria (family rank), pathogenic
bacteria and gastro-intestinal helminth communities of bank voles. Alpha diversity is represented using A) the
specific richness of the gut bacteriome, B) the Shannon index of the gut bacteriome, C) the specific richness of the
pathogenic bacteria and D) of the gastro-intestinal helminths. Results are shown per locality, ordered from North to
South. Each colored point represents an individual. Black points indicate the average alpha diversity per locality. Box-
and-whisker plots represent the median and interquartile values. Different letters denote statistically significant
differences at $P < 0.05$, with pairwise post-hoc Tukey adjustments. E) Diagram showing the relationships between the
alpha diversity estimates of the three communities, while taking into account the influence of the sampling localities.
The effect size and the direction of the relationship between communities are represented using an arrow, its width
corresponding to estimate (β) x 10. B is given for the specific richness metrics, and for the Shannon index metrics (in
brackets). Only significant effects are represented.
446

Factors influencing the beta diversity of intra-host communities

Gut bacteriome

Mantel tests revealed a significant positive relationship between the dissimilarities in gut

bacteriome composition and the geographic distance, whatever the taxonomic rank considered

(Phylum: r = 0.19; P = 10^{-4}; Family: r = 0.24; P = 10^{-4}).

At the phylum level, the distance-based redundancy analyses (db-RDA) revealed a significant effect

of the sampling localities (P = 1 x 10^{-3}) and host gender (P = 8 x 10^{-3}) on gut microbiome

composition. Age class and BMI had no significant effect (Table S3A). The best model explained

18% of the total variance, with most of the variance being explained by localities (locality: R^2_{adj} =

0.16; gender: R^2_{adj} = 0.02). The variables constrained on two dimensions explained 15.40% of the

variance in phylum composition (CAP1 = 10.8%; CAP2 = 4.6%). The localities that contributed the

most to the CAP1 axis were Mont-sous-Vaudrey (negative values) and Cormaranche (positive

values) (Figure 3A). The CAP2 axis discriminated the gut bacteriome of male bank voles from

Chaux-des-Crotenay (negative values) and females from Chatillon (positive values). Similar results

were obtained at the family level (locality: $P = 1$ x 10^{-3}; $R^2_{adj} = 0.16$; gender: $P =0.027$; $R^2_{adj} = 0.01$),

with the CAP1 axis strongly discriminating Mont-sous-Vaudrey and Cormaranche localities (12.7%

of the total variance, Figure 3B). However, these results have to be taken cautiously as significant

differences of data dispersion were detected between localities (*betadisper,* Phylum: $P = 1$ x 10^{-3};

Family: $P = 1$ x 10^{-3}). The locality Cormaranche showed a lower dispersion compared to all other

localities (Tukey multiple comparisons, Table S3A).

Figure 3- Distance-based redundancy analysis (db-RDA) of the intestinal bacteria (phylum (A) and family (B)),
pathogenic bacteria (C) and gastro-intestinal helminth (D) communities, based on Bray–Curtis dissimilarities.
Dots represent individuals. The colors and shapes of the dots are associated with different factors : Localities from
North to South (Mont-sous-Vaudrey: Mont, Chaux-des-Crotenay: Chau, Chatillon: Chat and Cormaranche: Corm), age
class (juvenils: J, adults: A) and gender (females: F and males: M). Significatif factors based on the ORDIR2STEP
analysis are shown as arrows. Ellipses represent a 80% confidence interval around the centroid of the clusters, for each
locality.

476 Significant differences in the relative abundance of specific taxa were detected at the phylum and

477 family levels using DeSeq2 (Table 2; Table S3B). The main changes (Log2 fold values higher than

478 20) were detected between the northern (Mont-sous-Vaudrey) and southern localities. The northern

479 population was involved in 75% of all significant pairwise differences (Log2 fold change in

composition > 10). The gut bacteriome of these bank voles includes less Clostridiales (one unknown family; Firmicutes), Bifidobacteriaceae (Actinobacteria) and Desulfovibrionales (two unknown families; Proteobacteria) than in all the three other southern populations. The gut bacteriome of bank voles from Cormaranche (South) is characterized by less Erysipelotrichaceae (Firmicutes), than in the three northern populations.

Table 2- Pairwise comparisons of the relative abundance of the gut bacteriome (phylum and family levels) between sampling localities. Mont = Mont-sous-Vaudrey, Chau = Chaux-des-Croteay, Chat = Chatillon, Corm = Cormaranche. The Log² fold value is indicated for significant changes in abundance between two localities. Blue and red colors respectively correspond to negative and positive values. Higher absolute changes in Log² fold are emphasized with darker colors. The notation "order_fx" or "class_fx" is used when there was no assignation at the family level with the SILVA database.

	Sampling localities					
	Mont vs Chau	Mont vs Chat	Mont vs Corm	Chau vs Chat	Chau vs Corm	Chat vs Corm
Epsilonbacteraeota						
All	0	1.91	3.64	0	2.18	0
Helicobacteraceae	0	0	1.91	0	0	0
Proteobacteria						
All	0	0	0	0	0	0
Rhizobiaceae	10.26	0	0	0	0	0
Desulfovibrionales_f3	−21.32	−26.19	−26.04	−4.8	−4.72	0
Desulfovibrionales_f1	−20.96	−21.26	−24.22	0	0	0
Desulfovibrionales_f2	−4.23	−5.58	−5.1	0	0	0
Desulfovibrionaceae	−1.3	−1.63	−1.54	0	0	0
Burkholderiaceae	0	0	0	0	0.67	0
Firmicutes						
All	0	0.76	0.97	0	0	0
Erysipelotrichaceae	0	0	19.57	0	21.81	25.96
Christensenellaceae	0	0	−3.04	0	0	0
Ruminococcaceae	0	0	1.48	0	1.5	1.08
Streptococcaceae	0	−5.57	0	−6.1	0	0
Clostridiaceae	0	0	−5.24	0	0	0
Lactobacillaceae	0	0	−1.81	0	0	0
Clostridiales	0	0	2.16	0	2.27	1.43
Clostridiales_f4	0	0	−3.29	0	0	0
Clostridiales_f2	−26.15	−26.31	−26.54	0	0	0
Tenericutes						
All	2.74	4.25	3.13	0	0	0
Mollicutes_f1	−3.43	0	−2.93	0	0	0
Patescibacteria						
All	0	−2.94	−2.83	0	0	0
Saccharimonadaceae	−5.82	−5.77	−5.45	0	0	0
Spirochaetes						
All	2.52	2.65	4.52	0	1.99	1.87
Spirochaetaceae	0	0	2.39	0	0	0
Actinobacteria						
All	0	0	−0.89	0	0	0
Bifidobacteriaceae	−20.6	−24.84	−23.21	0	0	0
Eggerthellaceae	0	−1.31	−2.42	0	−1.74	0
Bacteroidetes						
All	1.29	1.73	3.28	0	1.99	1.54
Bacteroidales_f4	0	0	0	0	0	−3.59
Muribaculaceae	0	0	1.74	0	1.93	1.64
Rikenellaceae	0	1.73	0	0	0	0
Marinifilaceae	−2.57	0	0	0	0	0

Besides, differences in the composition of the gut bacteriome between males and females bank

voles were driven by the phylum *Firmicutes*, with males exhibiting higher relative abundance of

this taxa than females (Table S3B).

Pathogenic bacteria

Mantel test indicated no significant relationship between the dissimilarities in the composition of

the pathogenic bacteria community and the geographical distances ($r = 0.24$; $P = 0.34$).

The db-RDA showed significant variations of the pathogenic bacteria community composition with

individual characteristics (age class: $P = 0.01$; gender: $P = 1 \times 10^{-3}$) and localities ($P = 4 \times 10^{-3}$). The

best model explained 15% of the total variance, with all factors explaining only a small part of the

variance (locality: 5%; gender: 5%; age class: 3.5%). Data dispersion was similar between groups

(Table S3A). The db-RDA plot based on the two first axes only represented 11.1% of the total

variance (Figure 3C). CAP1 (7.6%) discriminated individuals mainly according to age class and

gender (male adults were opposed to juveniles and females), while CAP2 (3.5%) discriminated

Chaux-des-Crotenay from all other localities.

Gastro-intestinal helminths

Mantel test revealed a significant positive relationship between the dissimilarities in the

composition of the gastro-intestinal helminth community and the geographic distances ($r = 0.16$, P

$= 10^{-4}$).

The db-RDA emphasized a significant effect of the locality on the composition of the gastro-

intestinal helminth community (db-RDA, $P = 10^{-3}$; Table S3A). Individual characteristics (host

gender, age class and BMI) had no significant effect ($P = 0.27$, $P = 0.43$, $P = 0.15$, respectively).

Locality explained 41% of the total variance. The variables constrained on two dimensions

explained 32.4% of the variance (CAP1 = 24.4 % and CAP2 = 8%). CAP1 discriminated the

localities Mont-sous-Vaudrey and Chatillon from Chaux-des-Crotenay and Cormaranche, while

CAP2 discriminated Mont-sous-Vaudrey from Chatillon (Figure 3D). We found no significant

difference in the dispersion of data between localities ($P = 0.06$).

Interactions between intra-host communities' composition

Partial Mantel tests revealed a positive relationship between dissimilarities in gut bacteriome and

dissimilarities in gastro-intestinal helminth community composition (multiple Mantel test, phylum:

estimate = 0.18; $P = 0.015$; family: estimate = 0.22; $P = 4 \times 10-3$), but not with dissimilarities in

pathogenic bacteria community composition (multiple Mantel test: $P = 0.14$). After controlling for

geographic distances, dissimilarities in gut bacteriome composition remained significantly

correlated with dissimilarities in helminth community composition at the family level only (partial

Mantel test, phylum: $r = 0.03$, $P = 0.18$; family: $r = 0.06$, $P = 0.03$). Other results are detailed in

Table S3C.

The db-RDA and model selection procedures showed that the whole composition of the gut

bacteriome was significantly associated with the presence / absence of three pathogens:

Aonchotheca murissylvatici (ordistep db-RDA, Phylum: $P = 0.002$, $R^2_{adj} = 0.02$ and Family: $P =$

0.002, $R^2_{adj} = 0.04$), *Bartonella* sp (ordistep db-RDA, Phylum: $P = 0.022$, $R^2_{adj} = 0.02$; Family: $P =$

0.04, $R^2_{adj} = 0.01$) and *Heligmosomum mixtum* (ordistep db-RDA, Phylum: $P = 0.002$, $R^2_{adj} = 0.04$

and Family: $P = 0.002$, $R^2_{adj} = 0.06$). Neither the specific richness of pathogenic bacteria nor the

specific richness of the gastro-intestinal helminth community had a significant effect on the gut

bacteriome (Table S3D). The db-RDA triplots based on the two first axes only represented 9.5% at

phylum level and 10.2% at family level of the total variance (Figure 4; Figure S5). It showed that

individuals infected with *Aonchotheca murissylvatici* had more Proteobacteria and Actinobacteria,

but less Spirochaetes, Bacteroidetes and Epsilonbacteraeota. It also revealed contrasted patterns for

individuals either infected with *Bartonella* sp but not with *Heligmosomum mixtum*, or infected with

H. mixtum and not *Bartonella* sp. The former had more Bacteroidetes while the latter had more

Firmicutes.

Deseq2 analysis showed that infections with *A. murissylvatici* and *H. mixtum* were specifically

associated with a slightly different relative abundance (Log2 fold changes did not exceed 3.5) of

three main phyla (Epsilonbacteraeota, Spirochaetes and Bacteroidetes), and of Firmicutes and

Patescibacteria for *A. murissylvatici* only (Table 3). Slight changes (mostly increases) in relative

abundance of one specific family per phylum were also detected for *A. murissylvatici* infection (see

details in Table S3E), and in a lesser extent for *H. mixtum*. It was the case for an undetermined

family of Mollicutes (Tenericutes).

Moreover, this analysis revealed that higher levels of alpha diversity of the helminth community

were associated with slightly lower relative abundance of the phylum Epsilonbacteraeota,

Firmicutes, Spirochaetes and Bacteroidetes, of the family *Rhizobiaceae* (Proteobacteria) and with

slightly higher relative abundance of the family *Eggerthellaceae* (Actinobacteria) and *Mollicutes*

(Tenericutes) (Table 3; Table S3E).

Table 3- Changes in relative abundance of the gut bacteriome at the phylum ad family levels, with regard to
infectious status (helminths and pathogenic bacteria) and alpha diversity. The Log^2 fold change in relative
abundance is indicated for significant values only. Negative values are represented with blue colors, positive
values with red colors. Higher absolute changes in Log^2 fold are emphasized with darker colors.

	Phylum	Family	log2FoldChange
GI helminths (Specific Richness)			
	Spirochaetes		−1.02
	Bacteroidetes		−0.55
	Epsilonbacteraeota		−0.87
	Firmicutes		−0.25
	Actinobacteria		0.27
	Actinobacteria	*Eggerthellaceae*	0.67
	Proteobacteria	*Rhizobiaceae*	−3.79
	Epsilonbacteraeota	*Helicobacteraceae*	−0.89
GI helminths (Shannon Index)			
	Bacteroidetes		−1.26
	Spirochaetes		−2.16
	Epsilonbacteraeota		−1.93
	Firmicutes		−0.5
	Actinobacteria	*Eggerthellaceae*	1.76
	Proteobacteria	*Rhizobiaceae*	−8.08
	Tenericutes	*Mollicutes_f1*	2.8
Aonchotheca murissylvatici			
	Spirochaetes		−2.78
	Epsilonbacteraeota		−1.95
	Firmicutes		−0.55
	Bacteroidetes		−0.93
	Patescibacteria		1.73
	Proteobacteria	*Desulfovibrionaceae*	1.13
	Tenericutes	*Mollicutes_f1*	3.07
	Patescibacteria	*Saccharimonadaceae*	3.03
	Spirochaetes	*Spirochaetaceae*	−1.75
	Actinobacteria	*Eggerthellaceae*	1.4
Heligmosomum mixtum			
	Bacteroidetes		−1.36
	Epsilonbacteraeota		−1.88
	Spirochaetes		−1.93
	Tenericutes	*Mollicutes1*	3.34
Mastophorus muris			
	Proteobacteria	*Desulfovibrionales3*	−29.98
	Firmicutes	*Clostridiales_f1*	−21.41
	Firmicutes	*Clostridiales_f2*	−23.73
Catenotaenia henttoneni			
	Proteobacteria	*Rhizobiaceae*	−24.71
	Proteobacteria	*Burkholderiaceae*	−23.96
	Firmicutes	*Erysipelotrichaceae*	−20.6
Paranoplocephala omphaloides			
	Firmicutes	*Clostridiales_f2*	−25.37
	Proteobacteria	*Desulfovibrionales_f3*	−23.74
	Proteobacteria	*Rhizobiaceae*	−24.86
	Proteobacteria	*Burkholderiaceae*	−22.69
	Firmicutes	*Erysipelotrichaceae*	−22.4
	Firmicutes	*Christensenellaceae*	−5.24
Trichuris muris			
	Firmicutes	*Christensenellaceae*	−20.69
	Firmicutes	*Clostridiales_f1*	−18.62
	Bacteroidetes	*Bacteroidales_f4*	−18.38
Pathogenic bacteria (Specific Richness)			
	Bacteroidetes	*Bacteroidales_f3*	−28.51
Neoehrlichia mikurensis			
	Bacteroidetes		−1.43
	Firmicutes	*Erysipelotrichaceae*	−23.42
	Firmicutes	*Clostridiales*	−2.72
	Bacteroidetes	*Muribaculaceae*	−1.76
Orientia tsutsugamushi			
	Bacteroidetes		−2.8
	Firmicutes	*Erysipelotrichaceae*	−22.95
	Proteobacteria	*Desulfomicrobiaceae*	−5.71
	Bacteroidetes	*Muribaculaceae*	−2.45
Mycoplasma haemomuris			
	Spirochaetes		−1.99
Anaplasma phagocytophilum			
	Firmicutes	*Ruminococcaceae*	−1.75
Rickettsia sp			
	Firmicutes	*Erysipelotrichaceae*	−21.81

Besides, the most important changes observed in the relative abundance of gut bacteriome families

(Log2 fold higher than 18) were specifically associated with all other helminth species:

Mastophorus muris, Catenotaenia henttoneni, Paranoplocephala omphaloides and *Trichuris*

arvicolae. These associations were all strongly negative and concerned a wide array of gut

bacteriome families (Table S3E), most of them belonging to the phylum Firmicutes

(Christensenellaceae, two undetermined families of Clostridiales and Erysipelotrichaceae),

Proteobacteria (Burkholderiaceae and an undetermined family of Desulfovibrionales) and

Bacteroidetes (an undetermined family of Bacteroidales).

Considering pathogenic bacteria, we found that higher levels of alpha diversity (specific richness)

of the pathogenic bacteria community was associated with lower relative abundance of an

undetermined Bacteroidales family (Bacteroidetes), and that *Neoehrlichia mikurensis, Orientia*

tsutsugamushi and *Rickettsia* sp infections were associated with strong decreases in relative

abundance of *Erysipelotrichaceae* (Firmicutes) (Table 3; Table S3E). Other associations between

bacterial infections and changes in relative abundance of specific gut bacteriome taxa were

detected, but with little size effect (log2 fold changes lower than 5).

Discussion

Understanding the complex interlinkages between host microbiome, host-pathogen interactions and

health in wild animal populations has become a key topic in disease ecology, with consequences for

population dynamics, zoonotic risk management or biodiversity conservation. Here, we *(i)* use a

combination of morphological and metabarcoding approaches to describe intra-host communities of

wild rodent populations at a small geographical scale, *(ii)* provide evidence of intricate

interrelationships between the gut bacteriome and communities of gastro-intestinal helminths and

pathogenic bacteria, and *(iii)* investigate the influence of host individual and spatial factors on these

three intra-host communities, in a concomitant way.

A diverse intra-host community

This study brings a new light to the gut bacteriome composition in wild bank voles, which are

important zoonotic reservoirs in Europe. Their gut microbiome has been mainly studied in the

context of exposure to radioactive pollutants (e.g., Lavrinienko (2018), but see Knowles et al.

(2019)). Although we detect similar phyla and families than these previous studies, our in-depth

data allow us to detect other taxa as well as differences in their relative abundance. These

differences likely depend on the nature of the samples considered (faecal, colon or caecum

samples), given there is strong variation of the gut bacteriome both along the rodent digestive tract (e.g., in mice Anders *et al.*, 2021; Gu *et al.*, 2013) and between digestive tract and faeces (Anders *et al.*, 2021) of a single host individual.

Besides, our results confirm that helminths infections are the rule in natural populations of bank vole. Indeed, 85% of vole individuals are infected by at least one helminth. This high prevalence level is consistent with studies conducted on wild rodent populations (e.g., for bank voles (Grzybek *et al.*, 2015; Ribas Salvador *et al.*, 2011) or other rodent species (Behnke *et al.*, 1999; Vanden Broecke *et al.*, 2021)). It likely reflects host tolerance to helminths - as a result of a balance between infection costs and immunopathologies-, or the ability of helminths to manipulate the host immune system (Babayan *et al.*, 2018).

Our findings also confirm the status of bank voles as hyper-reservoirs of zoonotic agents, with at least six zoonotic bacteria being detected using the 16S V4 metabarcoding approach. This complements previous studies that specifically triggered particular bacteria, including *Leptospira* sp., tick-borne bacteria or *Francisella tularensis* (Fischer *et al.*, 2018; Jeske *et al.*, 2019; Perez *et al.*, 2017).

Intrinsic and extrinsic factors influencing the gut bacteriome

We provide evidence of significant inter-individual variations in gut bacteriome composition. Factors intrinsic to bank voles only weakly explain these variations. We find that host gender slightly influences the composition of bank voles' gut bacteriome, males having 1.4 times more Firmicutes than females. Age has no influence on gut bacteriome diversity neither composition, as already found in wild rodents (Goertz *et al.*, 2019; Maurice *et al.*, 2015). However, our sampling did not allow to fully address this issue, given most individuals were adults.

Besides, we show that this inter-individual variability of gut bacteriome composition can be categorized in two compositionally different clusters equivalent to enterotypes (Arumugam *et al.*, 2011). Enterotypes have already been described in wild rodents (Goertz *et al.*, 2019; Li *et al.*, 2016), and they might reflect distinct ways of generating energy from substrates available in the digestive tract, as well as differences in diet (Rinninella *et al.*, 2019; Wang *et al.*, 2014). In bank voles, these enterotypes could be associated with different diets, one oriented toward seeds and plants and another one toward insects and berries. Indeed, enterotype 1 is characterized by families (namely, Helicobacteraceae, Lachnospiraceae and Muribaculaceae) that are involved in the breakdown of carbohydrates, fermentation of plant saccharid and degradation of glycan (see refs in Goertz *et al.*, 2019). These families are also predictive signals of a high-fat diet in mice (Bowerman *et al.*, 2021; Rodriguez-Daza *et al.*, 2020). In the opposite, enterotype 2 is characterized by families

(namely, Lactobacillae and Eggerthellaceae) that can be involved in the digestion of fermented food

(e.g., rodent food store over winter), and insect skeleton (see refs in Maurice *et al.*, 2015) or in the

degradation of polyphenol (Rodriguez-Daza *et al.*, 2020). All these aliments have varying

nutritional and chemical composition and may be part of bank vole diet (Ecke *et al.*, 2018). The

fraction of these different types of resources in bank vole diet may vary with resource preference or

availabilities, reproductive status, sampling date and location (e.g., Maurice *et al.*, 2015).

Spatial variations in gut bacteriome and their potential causes

There are now many evidence that geographical location is likely to shape variations in gut

bacteriome composition between localities sampled and studies. Previous works have already

shown that the structure of rodent gut microbiome varied between localities at large spatial scales

due to biogeographic or genetic factors (Linnenbrink *et al.*, 2013). Geographic variability has also

been found at smaller spatial scales (e.g., few km Goertz *et al.*, 2019). Here, our results provide

significant evidence for spatial structure of gut bacteriome between bank vole populations that are

between 50 and 130 km away, with no obvious barrier to dispersal or gene flow (Dubois *et al.*,

2018).

We observe gradual changes in terms of gut bacteriome richness, evenness and

Firmicutes/Bacteroidetes ratio, between bank voles from the northern and southern populations.

Although the links between the diversity and functional capacity of the gut bacteriome are still not

fully understood (Worsley *et al.*, 2021), it is largely assumed that changes in diversity are associated

with shifts in metabolism (Reese & Dunn, 2018). Bank voles from southern populations exhibit

higher specific richness and lower evenness of the gut bacteriome, as well as lower dispersion of

gut bacteriome composition. They have higher levels of body condition and F/B ratio, that are

indicative of an optimisation of calorie intake and absorption, weight gain and fat storage (see refs

in Wolf *et al.*, 2021). Altogether, these results could suggest strong constraints on gut bacteriome

function to maximise energy extraction. The northern populations show the opposite patterns.

Lower BMI and lower levels of F/B ratio might reflect energy production and conversion, amino

acid transport and metabolism, while diversity patterns (higher evenness and lower specific richness

of the gut bacteriome) could suggest lower stochasticity and/or directional selection. Further studies

are required to investigate the ecological processes driving these changes in gut bacteriome.

We also show that gut microbiome dissimilarities increased with distance between populations,

marked differences being noticeable between the two more distant localities sampled, Mont-sous-

Vaudrey in the North and Cormaranche in the South. These geographic differences might be

explained by dietary variations, and variable proportions of plants, seeds, berries and insects in bank

vole diet. For example, Erysipelotrichaceae are less represented in the South. These taxa are

involved in the metabolism of products of cellulolysis (Bermingham *et al.*, 2017; Kaakoush, 2015).

Their expansion may be promoted by plant sterols (Harris *et al.*, 2014) and may be negatively

correlated with protein content of the diet (Bermingham *et al.*, 2017).

Lastly, these differences in gut microbiome composition between the northern and southern

populations might also reflect physiological variations related to physiology, health, and potentially

to immunity. Clostridiales and Bifidobacteriaceae participate in the maintenance of intestinal

homeostasis, and in the regulation of inflammation or in the gut barrier function (Arboleya *et al.*,

2016; Hakansson & Molin, 2011; Lopetuso *et al.*, 2013), while specific taxa within

Erysipelotrichaceae may be correlated with inflammation or have immunogenic potential

(Kaakoush, 2015; Zhai *et al.*, 2019). Desulfovibrionales activities result in the production of H_2S,

that in turn, leads to damages of the gut barrier, production of endotoxins and pro-inflammatory

cytokines (Hu *et al.*, 2022). Our previous work revealed that bank voles from these southern

populations had lower basal level of *Tnf-a* (a pro-inflammatory cytokine) and higher level of *Mx2*

antiviral gene expression than those from these northern populations (Dubois *et al.*, 2018). Future

studies should assess the potential relationships between variations in gut bacteriome composition

and the capacities to regulate or mount immune responses and inflammation in these bank vole

populations.

Marked relationships between gut bacteriome and helminth communities

While we did not detect any relationship between pathogenic bacteria and gut bacteriome diversity

or composition, we found evidence for strong association between helminth community and gut

bacteriome. The former pattern might be explained by the fact that the pathogenic bacteria detected

here do not constitute a functional community, in particular because their ecological niche can be

very different. For example, hemotrophic *Mycoplasma* parasitizes erythrocytes (Alabi *et al.*, 2020)

while *Borrelia* disseminates through the bloodstream and/or lymphatic system to invade and

colonize various tissue (Zeidner *et al.*, 2001). The latter pattern may be interpreted under two

perspectives.

First, the strong positive associations between the diversity of helminth community and gut

bacteriome might corroborate the hypothesis and experimental evidence showing that helminths

have the capacity to maintain higher gut microbiota diversity and may represent gut homoeostasis

(Kreisinger *et al.*, 2015). Indeed, low-intensity, chronic helminth infections are commonly linked to

high microbial diversity and predominance of bacteria typically associated with gut health (Peachey

et al., 2017). Nevertheless, this interpretation has to be taken cautiously as the diversity of both

communities was strongly influenced by the localities of sampling. The environment might

therefore shape similarly gut bacteriome and helminth community diversity.

Second significant associations between helminth community and gut bacteriome composition -

which remain significant even when potential geographic confounding effects were removed - may

be linked to the fact that both communities reside in the same environmental niche (host intestines).

From there, they likely experience similar selection pressure (e.g., host immune responses), with

potentially strong interactions and reciprocal influence expected between them, which could shape

their composition (Glendinning *et al.*, 2014). Unfortunately, the causal processes behind these gut

microbiome and helminths interactions are complex, multifaceted and difficult to assess, in

particular because experimental studies can only focus on single helminth infections– while

interactions between/within community are the rule within the host organism. The field of

microbiome research would thus benefit from taking into account the whole composition of gastro

intestinal helminth community rather than single helminth infections only.

Three-way relationships within intra-host communities

The associations detected between the gut microbiome diversity or composition and the

presence/absence of infections (helminths and bacteria) are of two types, and might surprisingly not

always reflect gut homeostasis/dysbiosis.

First, some infections are significantly associated with the global composition of the gut

bacteriome, but have only little impact on specific gut bacterial taxa. This pattern concerns

Heligmosomum mixtum, *Bartonella* and *Aonchothecha murisylvatici*. Such associations could

suggest long co-adaptation/coevolution between these taxa and the gut bacteriome. Some evidence

suggests that *Bartonella* may be acting as a symbiont more than a pathogen (refs in Lei & Olival,

2014). Significant coevolutionary congruence has been found between *Bartonella* species and their

rodent hosts, and *Bartonella* infections in rodents lead to an asymptomatic long lasting intra-

erythrocytic bacteraemia (Deng *et al.*, 2012; Lei & Olival, 2014). Besides, *Heligmosomum* species

have been found to have only relatively low pathogenic impact on voles (e.g., Janova *et al.*, 2010).

The negative associations detected here between *Bartonella* and *Heligmosomum mixtum,* and their

respective links with Bacteroidetes and Firmicutes, deserve further attention to decipher the

potential direct and indirect interactions between these taxa, bank vole immunity and gut

bacteriome. Lastly, too little is known on *Aonchotheca murisylvatici* and its impacts on rodent

fitness to interpret the relationships observed between this helminth and gut bacteriome

composition.

Second, other infections are strongly associated with large changes in the relative abundance of one

or few specific taxa from the gut bacteriome, but not with the global composition of this later.

These species-specific associations concern *Trichuris muris, Caetenotaenia henttoneni,*

Paranoplocephala omphaloides and *Mastophorus muris* for the helminths and *Neoehrlichia*

mikurensis, Orientia tsutsugamushi, Mycoplasma haemomuris, Anaplasma phagocytophilum and

Rickettsia for the bacteria.

The bacterial infections were associated with a lower relative abundance of Erysipelotrichaceae.

This is striking because observed changes in the gut microbiota during infection are rarely

consistent, even with respect to single pathogens (Sabey *et al.*, 2021). In humans,

Erysipelotrichaceae have been associated with a number of diseases such as HIV and norovirus

infections, inflammation-related intestinal disease and metabolic disorders (Kaakoush, 2015). No

study has yet investigated the relationships between Erysipelotrichaceae, gut dysbiosis and

infections in rodents. We also find that a larger number of bacterial infections is correlated with

lower relative abundance of an unknown family of Bacteroidales. These later have been shown to

promote epithelial barrier function and maintain epithelial homeostasis (Kuhn *et al.*, 2018). It would

be interesting to investigate whether this family could be an indicator of rodent health, lower

abundance reflecting individuals in poor health and more susceptible to opportunistic infections.

The species-specific associations between helminths infections and gut bacteria included

Proteobacteria (Desulfovibiolales, Burkholderiaceae), Firmicutes (Clostridiales,

Erysipelotrichaceae) and Bacteroidetes (Bacteroidales). High-intensity, acute helminth infections

may correlate with changes in hosts gut microbiota and gut dysbiosis, through direct and indirect

(immune or other processes such as malnutrition) interactions (Peachey *et al.*, 2017). Nevertheless,

up to now, the patterns of shifts in gut bacteriome associated with helminth infections remain hardly

predictable. Research that addressed this issue, using laboratory or wild animals, have provided

variable and sometimes even contradictory conclusions, even for single host-helminth models. A

potential explanation is that these infection-associated microbiota shifts could depend on the

presence of other helminths and the duration of infection (e.g. Sabey *et al.*, 2021). Local

interactions between helminths and between helminths and gut bacteria could mediate changes in

infection outcomes as well as the gut bacteria and helminth populations themselves (Glendinning *et*

al., 2014).

Conclusion

Altogether, these results emphasize complex interlinkages between gut bacteriome, gastro-intestinal

helminths and bacterial infections in wild animal populations. We emphasize the strong impact of

environment, even at fine geographical scales, on these interactions. Shifts in diet or host genetics could mediate the spatial changes observed in gut bacteriome. However, the processes shaping gut bacteriome diversity and composition are many and complex, and further investigations are required to decipher the relative importance of drift, dispersal or selection on bank vole gut bacteriome in the populations studied here. Besides, we find a diverse array of associations between gut bacteriome and gastro-intestinal helminths or pathogenic bacteria, some being significant at the scale of the whole community and other being species-specific only. Whether these patterns reflect coadaptation, dysbiosis or indirect interactions with host immunity and coinfections should now be considered to better understand the spatial variations observed in the relationships between gut bacteriome and health.

Acknowledgement

Data used in this work were partly produced through the genotyping and sequencing facilities of ISEM (Institut des Sciences de l'Evolution-Montpellier) and Labex CeMEB (Centre Méditerranéen Environnement Biodiversité).

Funding

This research was funded through the INRAE metaprogram MEM Hantagulumic and the 2018-2019 BiodivERsA joint call for research proposals, under the BiodivERsA3 ERA-Net COFUND programme, and with the funding organization ANR. A. Dubois PhD was funded by an INRA-EFPA/ANSES fellowship.

Conflict of interest disclosure

The authors declare that they have no financial conflict of interest with the content of this article. N. C. is one of the PCI Inf recommenders. B.R. is part of the managing board of PCI Inf.

Data and script availability

Raw data and scripts are available on zenodo: https://doi.org/10.5281/zenodo.6573141

Supplementary information

Supplementary Figure S1. Maps showing the sampling area (left) and localities (right) in France. Forests are indicated in green and water in blue. The four sampling localities are represented with a colored polygon. The arrow indicates the North.

Supplementary Figure S2. Composition of the gut bacteriome. The relative abundance of six phyla representing 99% of the total composition is represented. Individuals are grouped by sampling localities, which are ordered from North to South. (A) Bar graph shows individual variation in phyla composition (phylum=color). (B) Box and whisker plots represent median and interquartile values for each phylum. Black dots correspond to mean values, and colored dots correspond to individuals.

Supplementary Figure S3. Variations of alpha diversity between localities, for the gut bacteriome (phylum level), pathogenic bacteria and gastro-intestinal helminth communities. Alpha diversity is represented using the specific richness of the gut bacteriome (A) or the Shannon index of the gut bacteriome (B). Results are shown per localities, ordered from North to South. Each colored point represents an individual. Black point indicates the average alpha diversity per locality. Box-and-whisker plots represent the median and interquartile values. Different letters denote statistically significant differences at $P < 0.05$, after pairwise post-hoc Tukey adjustments.

Supplementary Figure S4. Variations of alpha diversity with individual factors, for the gut bacteriome (family level), pathogenic bacteria and gastro-intestinal helminths of bank voles. Alpha diversity is estimated using the Shannon index (A and D) or the specific richness (G) of the gut bacteriome. In all other graphs, the specific richness of the pathogenic bacteria and gastro-intestinal helminths communities is used. In G, H, I, the blue line corresponds to the linear regression line.

Supplementary Figure S5. Relationships between the composition of the gut bacteriome, pathogenic bacteria and gastro-intestinal helminth communities: A) db-RDA triplot shows the structure of the gut microbiome at the family level and the correlations with the pathogen communities. The arrows correspond to the significant explanatory variables. Each point corresponds to an individual, and the colors correspond to the different sampling localities.

Supplementary Table S1. Variation of the Firmicutes/Bacteroidetes ratio with localities and individual factors.

Supplementary Table S2. Alpha diversity metrics and statistics for the gut bacteriome, pathogenic bacteria and helminth communities of bank voles.

Supplementary Table S3. Beta diversity metrics and statistics for the gut bacteriome, pathogenic bacteria and helminth communities of bank voles.

AUTHOR CONTRIBUTIONS

M.B.: Data curation; Formal analysis; Methodology; Writing – original draft

M.G.: Conceptualization; Data curation; Formal analysis; Methodology; Supervision; Writing – review and editing

A.D.: Conceptualization; Data curation

C.A.D.: Data curation; Writing – review and editing

P.M.: Conceptualization; Funding acquisition; Investigation; Supervision; Writing – review and editing

B.R.: Conceptualization; Methodology; Supervision; Writing – review and editing

N.C.: Conceptualization; Funding acquisition; Investigation; Methodology; Project administration; Resources; Supervision; Writing – original draft

REFERENCES

Abbate, J., Galan, M., Razzauti, M.*, et al.* (In_revision). Pathogen community composition and co-infection patterns in a wild community of rodents. *PCI Ecology.* doi:10.1101/2020.02.09.940494

Adair, K. L.,Douglas, A. E. (2017). Making a microbiome: the many determinants of host-associated microbial community composition. *Curr Opin Microbiol, 35*, 23-29. doi:10.1016/j.mib.2016.11.002

Alabi, A. S., Monti, G., Otth, C.*, et al.* (2020). Molecular survey and genetic diversity of Hemoplasmas in rodents from Chile. *Microorganisms, 8*(10). doi:10.3390/microorganisms8101493

Anders, J. L., Moustafa, M. A. M., Mohamed, W. M. A.*, et al.* (2021). Comparing the gut microbiome along the gastrointestinal tract of three sympatric species of wild rodents. *Scientific Reports, 11*(1). doi:10.1038/s41598-021-99379-6

Anderson, R. C., Chabaud, A. G.,Willmott, S. (2009). *Keys to the nematode parasites of vertebrates: archival volume.* CABI, Wallingford, Oxon, UK.

Arboleya, S., Watkins, C., Stanton, C.*, et al.* (2016). Gut Bifidobacteria populations in human health and aging. *Frontiers in Microbiology, 7*. doi:10.3389/fmicb.2016.01204

Arumugam, M., Raes, J., Pelletier, E.*, et al.* (2011). Enterotypes of the human gut microbiome. *Nature, 473*(7346), 174-180. doi:10.1038/nature09944

Babayan, S. A., Liu, W., Hamilton, G.*, et al.* (2018). The immune and non-immune pathways that drive chronic gastrointestinal helminth burdens in the wild. *Frontiers in Immunology, 9*, 56. doi:10.3389/fimmu.2018.00056

Behnke, J. M., Lewis, J. W., Zain, S. N.*, et al.* (1999). Helminth infections in *Apodemus sylvaticus* in southern England: interactive effects of host age, sex and year on the prevalence and abundance of infections. *Journal of Helminthology, 73*(1), 31-44.

Belkaid, Y.,Hand, T. W. (2014). Role of the microbiota in immunity and inflammation. *Cell, 157*(1), 121-141. doi:10.1016/j.cell.2014.03.011

Bermingham, E. N., Maclean, P., Thomas, D. G.*, et al.* (2017). Key bacterial families
(Clostridiaceae, Erysipelotrichaceae and Bacteroidaceae) are related to the digestion of
protein and energy in dogs. *Peerj, 5*. doi:10.7717/peerj.3019

Bowerman, K. L., Knowles, S. C. L., Bradley, J. E.*, et al.* (2021). Effects of laboratory
domestication on the rodent gut microbiome. *ISME Communications, 1*(1), 49.
doi:10.1038/s43705-021-00053-9

Camacho, C., Coulouris, G., Avagyan, V.*, et al.* (2009). BLAST+: architecture and applications.
BMC Bioinformatics, 10, 421. doi:10.1186/1471-2105-10-421

Clemente, J. C., Ursell, L. K., Parfrey, L. W.*, et al.* (2012). The impact of the gut microbiota on
human health: an integrative view. *Cell, 148*(6), 1258-1270. doi:10.1016/j.cell.2012.01.035

Deng, H. K., Le Rhun, D., Buffet, J. P. R.*, et al.* (2012). Strategies of exploitation of mammalian
reservoirs by *Bartonella* species. *Veterinary Research, 43*. doi:10.1186/1297-9716-43-15

Diagne, C. A., Charbonnel, N., Henttonen, H.*, et al.* (2017). Serological survey of zoonotic viruses
in invasive and native commensal rodents in Senegal, West Africa. *Vector Borne and*
Zoonotic Diseases, 17(10), 730-733. doi:10.1089/vbz.2017.2135

Dubois, A., Castel, G., Murri, S.*, et al.* (2017). Experimental infections of wild bank voles (*Myodes*
glareolus) from nephropatia epidemica endemic and non-endemic regions revealed slight
differences in Puumala virological course and immunological responses. *Virus Research,*
235, 67-72. doi:10.1016/j.virusres.2017.04.004

Dubois, A., Castel, G., Murri, S.*, et al.* (2018). Bank vole immunoheterogeneity may limit
nephropatia epidemica emergence in a French non-endemic region. *Parasitology, 145*, 393–
407. doi:10.1017/S0031182017001548

Ecke, F., Berglund, A. M., Rodushkin, I.*, et al.* (2018). Seasonal shift of diet in bank voles explains
trophic fate of anthropogenic osmium? *Science of the Total Environment, 624*, 1634-1639.
doi:10.1016/j.scitotenv.2017.10.056

Escudié, F., Auer, L., Bernard, M.*, et al.* (2018). FROGS: Find Rapidly OTU with Galaxy Solution.
Bioinformatics, 34, 1287–1294.

Fischer, S., Mayer-Scholl, A., Imholt, C.*, et al.* (2018). *Leptospira* genomospecies and sequence
type prevalence in small mammal populations in Germany. *Vector Borne and Zoonotic*
Diseases, 18(4), 188-199. doi:10.1089/vbz.2017.2140

Foster, Z. S. L., Sharpton, T. J.,Grunwald, N. J. (2017). Metacoder: An R package for visualization
and manipulation of community taxonomic diversity data. *Plos Computational Biology,*
13(2). doi:10.1371/journal.pcbi.1005404

Galan, M., Razzauti, M., Bard, E.*, et al.* (2016). 16S metagenomics for epidemiological survey of
bacteria in wildlife. *mSystem, 1*(4), e00032-00016. doi:10.1128/mSystems.00032-16

Glendinning, L., Nausch, N., Free, A.*, et al.* (2014). The microbiota and helminths: sharing the
same niche in the human host. *Parasitology, 141*(10), 1255-1271.
doi:10.1017/S0031182014000699

Goertz, S., de Menezes, A. B., Birtles, R. J.*, et al.* (2019). Geographical location influences the
composition of the gut microbiota in wild house mice (*Mus musculus domesticus*) at a fine
spatial scale. *Plos One, 14*(9). doi:10.1371/journal.pone.0222501

Grzybek, M., Bajer, A., Bednarska, M.*, et al.* (2015). Long-term spatiotemporal stability and
dynamic changes in helminth infracommunities of bank voles (Myodes glareolus) in NE
Poland. *Parasitology, 142*(14), 1722-1743. doi:10.1017/S0031182015001225

Gu, S. H., Chen, D. D., Zhang, J. N.*, et al.* (2013). Bacterial community mapping of the mouse
gastrointestinal tract. *Plos One, 8*(10). doi:10.1371/journal.pone.0074957

Haegeman, B., Hamelin, J., Moriarty, J.*, et al.* (2013). Robust estimation of microbial diversity in
theory and in practice. *Isme Journal, 7*(6), 1092-1101. doi:10.1038/ismej.2013.10

Hakansson, A.,Molin, G. (2011). Gut microbiota and inflammation. *Nutrients, 3*(6), 637-682.
doi:10.3390/nu3060637

Harris, J. K., El Kasmi, K. C., Anderson, A. L*., et al.* (2014). Specific microbiome changes in a
mouse model of parenteral nutrition associated liver injury and intestinal inflammation. *Plos*
One, 9(10). doi:10.1371/journal.pone.0110396
Hoarau, A. O. G., Mavingui, P.,Lebarbenchon, C. (2020). Coinfections in wildlife: Focus on a
neglected aspect of infectious disease epidemiology. *Plos Pathogens, 16*(9).
doi:10.1371/journal.ppat.1008790
Holmes, I., Harris, K.,Quince, C. (2012). Dirichlet Multinomial Mixtures: Generative Models for
Microbial Metagenomics. *Plos One, 7*(2). doi:10.1371/journal.pone.0030126
Honda, K.,Littman, D. R. (2016). The microbiota in adaptive immune homeostasis and disease.
Nature, 535(7610), 75-84. doi:10.1038/nature18848
Hothorn, T., Bretz, F.,Westfall, P. (2008). Simultaneous inference in general parametric models.
Biometrical Journal, 50(3), 346-363. doi:10.1002/bimj.200810425
Hu, H., Shao, W. T., Liu, Q*., et al.* (2022). Gut microbiota promotes cholesterol gallstone formation
by modulating bile acid composition and biliary cholesterol secretion. *Nature*
Communications, 13(1). doi:10.1038/s41467-021-27758-8
Janova, E., Skoric, M., Heroldova, M*., et al.* (2010). Determinants of the prevalence of
Heligmosomum costellatum (Heligmosomidae: Trichostrongyloidea) in a common vole
population in southern Moravia, Czech Republic. *Journal of Helminthology, 84*(4), 410-414.
doi:10.1017/s0022149x1000009x
Jeske, K., Tomaso, H., Imholt, C*., et al.* (2019). Detection of *Francisella tularensis* in three vole
species in Central Europe. *Transboundary Emerging Diseases, 66*(2), 1029-1032.
doi:10.1111/tbed.13078
Johnson, J. B.,Omland, K. S. (2004). Model selection in ecology and evolution. *Trends in Ecology*
and Evolution, 19, 101–108.
Johnson, P. T., de Roode, J. C.,Fenton, A. (2015). Why infectious disease research needs
community ecology. *Science, 349*(6252), 1259504. doi:10.1126/science.1259504
Kaakoush, N. O. (2015). Insights into the role of Erysipelotrichaceae in the human host. *Frontiers*
in Cellular and Infection Microbiology, 5. doi:10.3309/fcimb.2015.00004
Kamada, N., Chen, G. Y., Inohara, N*., et al.* (2013). Control of pathogens and pathobionts by the
gut microbiota. *Nature Immunology, 14*(7), 685-690. doi:10.1038/ni.2608
Khalil, L. F., Jones, A.,Bray, R. A. (1994). *Keys to the cestode parasites of Vertebrates.* CAB
International, Wallingford, Oxon, UK.
Khosravi, A.,Mazmanian, S. K. (2013). Disruption of the gut microbiome as a risk factor for
microbial infections. *Current Opinion in Microbiology, 16*(2), 221-227.
doi:10.1016/j.mib.2013.03.009
Kim, C. H. (2021). Control of lymphocyte functions by gut microbiota-derived short-chain fatty
acids. *Cellular & Molecular Immunology, 18*(5), 1161-1171. doi:10.1038/s41423-020-
00625-0
Knowles, S. C. L., Eccles, R. M.,Baltrunaite, L. (2019). Species identity dominates over
environment in shaping the microbiota of small mammals. *Ecology Letters, 22*(5), 826-837.
doi:10.1111/ele.13240
Kozich, J. J., Westcott, S. L., Baxter, N. T*., et al.* (2013). Development of a dual-index sequencing
strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina
sequencing platform. *Applied and Environmental Microbiology, 79*(17), 5112-5120.
doi:10.1128/aem.01043-13
Kreisinger, J., Bastien, G., Hauffe, H. C*., et al.* (2015). Interactions between multiple helminths and
the gut microbiota in wild rodents. *Philosophical Transactions of the Royal Society B-*
Biological Sciences, 370(1675). doi:10.1098/rstb.2014.0295
Kuhn, K. A., Schulz, H. M., Regner, E. H*., et al.* (2018). Bacteroidales recruit IL-6-producing
intraepithelial lymphocytes in the colon to promote barrier integrity. *Mucosal Immunology,*
11(2), 357-368. doi:10.1038/mi.2017.55

Lahti, L.,Shetty, S. (2017). Tools for microbiome analysis in R.

Lavrinienko, A., Mappes, T., Tukalenko, E.*, et al.* (2018). Environmental radiation alters the gut
microbiome of the bank vole *Myodes glareolus*. *Isme Journal, 12*(11), 2801-2806.
doi:10.1038/s41396-018-0214-x

Lei, B. N. R.,Olival, K. J. (2014). Contrasting patterns in mammal-bacteria coevolution: *Bartonella*
and *Leptospira* in bats and rodents. *Plos Neglected Tropical Diseases, 8*(3).
doi:10.1371/journal.pntd.0002738

Leung, J. M., Graham, A. L.,Knowles, S. C. L. (2018). Parasite-microbiota interactions with the
vertebrate gut: Synthesis through an ecological lens. *Frontiers in Microbiology, 9*.
doi:10.3389/fmicb.2018.00843

Ley, R. E., Hamady, M., Lozupone, C.*, et al.* (2008). Evolution of mammals and their gut microbes.
Science, 320(5883), 1647-1651. doi:10.1126/science.1155725

Li, H., Li, T. T., Beasley, D. E.*, et al.* (2016). Diet diversity is associated with beta but not alpha
diversity of pika gut microbiota. *Frontiers in Microbiology, 7*.
doi:10.3389/fmicb.2016.01169

Linnenbrink, M., Wang, J., Hardouin, E. A.*, et al.* (2013). The role of biogeography in shaping
diversity of the intestinal microbiota in house mice. *Molecular Ecology, 22*(7), 1904-1916.
doi:10.1111/mec.12206

Loke, P.,Lim, Y. A. L. (2015). Helminths and the microbiota: parts of the hygiene hypothesis.
Parasite Immunology, 37(6), 314-323. doi:10.1111/pim.12193

Lopetuso, L. R., Scaldaferri, F., Petito, V.*, et al.* (2013). Commensal Clostridia: leading players in
the maintenance of gut homeostasis. *Gut Pathogens, 5*. doi:10.1186/1757-4749-5-23

Love, M. I., Huber, W.,Anders, S. (2014). Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biology, 15*(12). doi:10.1186/s13059-014-0550-8

Magoc, T.,Salzberg, S. L. (2011). FLASH: fast length adjustment of short reads to improve genome
assemblies. *Bioinformatics, 27*(21), 2957-2963. doi:10.1093/bioinformatics/btr507

Mahe, F., Rognes, T., Quince, C.*, et al.* (2014). Swarm: robust and fast clustering method for
amplicon-based studies. *Peerj, 2*. doi:10.7717/peerj.593

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal, 17, 10-12. doi:10.14806/ej.17.1.200

Maurice, C. F., Knowles, S. C. L., Ladau, J.*, et al.* (2015). Marked seasonal variation in the wild
mouse gut microbiota. *Isme Journal, 9*(11), 2423-2434. doi:10.1038/ismej.2015.53

McKnight, D. T., Huerlimann, R., Bower, D. S.*, et al.* (2019). Methods for normalizing microbiome
data: An ecological perspective. *Methods in Ecology and Evolution, 10*(3), 389-400.
doi:10.1111/2041-210x.13115

McMurdie, P. J.,Holmes, S. (2013). phyloseq: an R package for reproducible interactive analysis
and graphics of microbiome census data. *Plos One, 8*(4), e61217.
doi:10.1371/journal.pone.0061217

Mills, J. N., Childs, J., Ksiazek, T. G.*, et al.* (1995). *Methods for trapping and sampling small*
mammals for virologic testing. Atlanta: Centers for Disease Control and Prevention.

Moran, N. A., Ochman, H.,Hammer, T. J. (2019). Evolutionary and ecological consequences of gut
microbial communities. *Annu Rev Ecol Evol Syst, 50*(1), 451-475. doi:10.1146/annurev-
ecolsys-110617-062453

Morgan, M. (2021). Dirichlet-Multinomial Mixture Model Machine Learning for Microbiome Data.

Oksanen, J., Blanchet, F. G., Friendly, M.*, et al.* (2020). vegan: Community Ecology Package.

Pascoe, E. L., Hauffe, H. C., Marchesi, J. R.*, et al.* (2017). Network analysis of gut microbiota
literature: an overview of the research landscape in non-human animal studies. *Isme*
Journal, 11(12), 2644-2651. doi:10.1038/ismej.2017.133

Peachey, L. E., Jenkins, T. P.,Cantacessi, C. (2017). This gut ain't big enough for both of us. Or is
it? Helminth-microbiota interactions in veterinary species. *Trends in Parasitology, 33*(8),
619-632. doi:10.1016/j.pt.2017.04.004

Perez, G., Bastian, S., Chastagner, A*., et al.* (2017). Ecological factors influencing small mammal
infection by *Anaplasma phagocytophilum* and *Borrelia burgdorferi* s.l. in agricultural and
forest landscapes. *Environmental Microbiology, 19*(10), 4205-4219. doi:10.1111/1462-
2920.13885
Reese, A. T.,Dunn, R. R. (2018). Drivers of microbiome biodiversity: A review of general rules,
feces, and ignorance. *Mbio, 9*(4). doi:10.1128/mBio.01294-18
Revell, L. J. (2012). phytools: an R package for phylogenetic comparative biology (and other
things). *Methods in Ecology and Evolution, 3*(2), 217-223. doi:10.1111/j.2041-
210X.2011.00169.x
Ribas Salvador, A., Guivier, E., Chaval, Y*., et al.* (2011). Concomitant influence of helminth
infection and landscape on the distribution of Puumala hantavirus in its reservoir, *Myodes*
glareolus. *BMC Microbiology, 11*(30), 1-13. doi:http://www.biomedcentral.com/1471-
2180/11/30
Rinninella, E., Raoul, P., Cintoni, M*., et al.* (2019). What is the healthy gut microbiota
composition? A changing ecosystem across age, environment, diet, and diseases.
Microorganisms, 7(1). doi:10.3390/microorganisms7010014
Rodriguez-Daza, M. C., Roquim, M., Dudonne, S*., et al.* (2020). Berry polyphenols and fibers
modulate distinct microbial metabolic functions and gut microbiota enterotype-like
clustering in obese mice. *Frontiers in Microbiology, 11*. doi:10.3389/fmicb.2020.02032
Rolhion, N.,Chassaing, B. (2016). When pathogenic bacteria meet the intestinal microbiota.
Philosophical Transactions of the Royal Society B-Biological Sciences, 371(1707).
doi:10.1098/rstb.2015.0504
Rosenfeld, C. S. (2017). Gut dysbiosis in animals due to environmental chemical exposures.
Frontiers in Cellular and Infection Microbiology, 7. doi:10.3389/fcimb.2017.00396
Round, J. L.,Mazmanian, S. K. (2009). The gut microbiota shapes intestinal immune responses
during health and disease. *Nature Reviews Immunology, 9*(5), 313-323. doi:10.1038/nri2515
Sabey, K. A., Song, S. J., Jolles, A*., et al.* (2021). Coinfection and infection duration shape how
pathogens affect the African buffalo gut microbiota. *Isme Journal, 15*(5), 1359-1371.
doi:10.1038/s41396-020-00855-0
Salter, S. J., Cox, M. J., Turek, E. M*., et al.* (2014). Reagent and laboratory contamination can
critically impact sequence-based microbiome analyses. *Bmc Biology, 12*, 87.
doi:10.1186/s12915-014-0087-z
Suzuki, T. A.,Nachman, M. W. (2016). Spatial heterogeneity of gut microbial composition along
the gastrointestinal tract in natural populations of house mice. *Plos One, 11*(9).
doi:10.1371/journal.pone.0163720
team, R. c. (2020). A language and environment for statistical computing.: Austria, Vienna.
Telfer, S., Lambin, X., Birtles, R*., et al.* (2010). Species interactions in a parasite community drive
infection risk in a wildlife population. *Science, 330*(6001), 243-246.
doi:10.1126/science.1190333
Trevelline, B. K., Fontaine, S. S., Hartup, B. K*., et al.* (2019). Conservation biology needs a
microbial renaissance: a call for the consideration of host-associated microbiota in wildlife
management practices. *Proceedings of the Royal Society B-Biological Sciences, 286*(1895).
doi:10.1098/rspb.2018.2448
Vanden Broecke, B., Bernaerts, L., Ribas, A*., et al.* (2021). Linking behavior, co-infection patterns,
and viral infection risk with the whole gastrointestinal helminth community structure in
M*astomys natalensis*. *Frontiers in Veterinary Science, 8*, 669058.
doi:10.3389/fvets.2021.669058
Vujkovic-Cvijin, I., Sklar, J., Jiang, L*., et al.* (2020). Host variables confound gut microbiota
studies of human disease. *Nature, 587*(7834), 448-454. doi:10.1038/s41586-020-2881-9
Wang, J., Linnenbrink, M., Kunzel, S*., et al.* (2014). Dietary history contributes to enterotype-like
clustering and functional metagenomic content in the intestinal microbiome of wild mice.

Proceedings of the National Academy of Sciences of the United States of America, 111(26), E2703-E2710. doi:10.1073/pnas.1402342111

Wolf, J. F., Kriss, K. D., MacAulay, K. M.*, et al.* (2021). Gut microbiome composition predicts summer core range size in two divergent ungulates. *Fems Microbiology Ecology, 97*(5). doi:10.1093/femsec/fiab048

Worsley, S. F., Davies, C. S., Mannarelli, M. E.*, et al.* (2021). Gut microbiome composition, not alpha diversity, is associated with survival in a natural vertebrate population. *Animal Microbiome, 3*(1). doi:10.1186/s42523-021-00149-6

Zeidner, N. S., Schneider, B. S., Dolan, M. C.*, et al.* (2001). An analysis of spirochete load, strain, and pathology in a model of tick-transmitted Lyme borreliosis. *Vector-Borne and Zoonotic Diseases, 1*(1), 35-+. doi:10.1089/153036601750137642

Zhai, S., Qin, S., Li, L.*, et al.* (2019). Dietary butyrate suppresses inflammation through modulating gut microbiota in high-fat diet-fed mice. *FEMS Microbiology Letters, 366*(13). doi:10.1093/femsle/fnz153