The effects of combined action observation and motor imagery on corticospinal excitability and

movement outcomes: Two meta-analyses

Samantha Chye [a*], Ashika Chembila Valappil [a*], David J. Wright [b], Cornelia Frank [c], David A. Shearer [d], Christopher J. Tyler [a], Ceri E. Diss [a], Omar S. Mian [a], Neale A. Tillin [a] & Adam M. Bruton [a**]

[a] Centre for Integrated Research in Life and Health Sciences, School of Life and Health Sciences, University of Roehampton, London, UK, SW15 4JD. E-mail: chyes1@roehampton.ac.uk, chembila@roehampton.ac.uk, chris.tyler@roehampton.ac.uk, c.diss@roehampton.ac.uk, omar.mian@roehampton.ac.uk, neale.tillin@roehampton.ac.uk, adam.bruton@roehampton.ac.uk.

[b] Research Centre for Health, Psychology and Communities, Department of Psychology, Manchester Metropolitan University, Manchester, UK, M15 6GX. E-mail: d.j.wright@mmu.ac.uk.

[c] Sports and Movement Research Group, Department of Sports and Movement Science, School of Educational and Cultural Studies, Osnabrück University, Osnabrück, Germany. Email: cornelia.frank@uni-osnabrueck.de.

[d] Faculty of Life Science and Education, University of South Wales, UK, CF37 1DL. Email: david.shearer@southwales.ac.uk.

Author Note: *Samantha Chye and Ashika Chembila Valappil are shared first authors for this manuscript. **Adam M. Bruton is the corresponding author for this manuscript: adam.bruton@roehampton.ac.uk, Room 1051 Parkstead House, Whitelands College, School of Life and Health Sciences, University of Roehampton.

Highlights

- Motor imagery (MI) and action observation (AO) can be combined (AOMI)

- This paper synthesizes neurophysiological and behavioral evidence for AOMI

- AOMI had increased corticospinal excitability compared to AO and control but not MI

- AOMI led to improved movement outcomes compared to AO and control but not MI

- The reported effects of AOMI were maintained across all moderators

Abstract

Motor simulation interventions involving motor imagery (MI) and action observation (AO) have received considerable interest in the behavioral sciences. A growing body of research has focused on using AO and MI simultaneously, termed 'combined action observation and motor imagery' (AOMI). The current paper includes two meta-analyses that quantify changes in corticospinal excitability and motor skill performance for AOMI compared to AO, MI and control conditions. Specifically, the first meta-analysis collated and synthesized existing motor evoked potential (MEP) amplitude data from transcranial magnetic stimulation studies and the second meta-analysis collated and synthesized existing movement outcome data from behavioral studies. AOMI had a positive effect compared to control and AO but not MI conditions for both MEP amplitudes and movement outcomes. No methodological factors moderated the effects of AOMI, indicating a robust effect of AOMI across the two outcome variables. The results of the meta-analyses are discussed in relation to existing literature on motor simulation and skill acquisition, before providing viable directions for future research on this topic.

Keywords: AOMI, motor evoked potentials, dual action simulation, motor execution, motor imagery during action observation, transcranial magnetic stimulation

1. Introduction

According to motor simulation theory (Jeannerod, 1994, 2001, 2006), it is possible to cognitively

rehearse an action both overtly and covertly through action observation (AO) and motor imagery

(MI), with this cognitive simulation activating motor regions of the brain in a similar manner to

physical execution of the action. AO is a bottom-up process that involves the deliberate and

structured observation of human movement (Neuman & Gray, 2013), whereas MI is a top-down

process that involves the internal generation of the visual and kinesthetic elements of movement

(Macintyre et al., 2013). Literature has reported positive behavioral outcomes for AO- and MI-based

practice in sport (e.g., Guillot & Collet, 2008; Ste-Marie et al., 2012) and neurorehabilitation settings

(e.g., Buccino, 2014; De Vries & Mulder, 2007). There is preliminary evidence indicating that plastic

changes in the primary motor system may underpin these behavioral improvements (e.g., Yoxon and

Welsh, 2020). A recent large-scale meta-analysis of functional magnetic resonance imaging (fMRI)

data reported a shared network including premotor, rostral parietal, and somatosensory areas of the

brain active during AO, MI, and movement execution (Hardwick et al., 2018). Notably, there were

differences in the neural regions activated during AO and MI, and the brain activity for these

overlapped differently with brain activity for physically performed actions. It is possible that using

AO and MI simultaneously, typically labelled 'combined action observation and motor imagery'

(AOMI), may hold greater neural overlap with physical execution.

AOMI refers to a person watching a video or live demonstration of a movement while

simultaneously generating, maintaining, and transforming a time-synchronized kinesthetic

representation of the same action (Eaves, Riach et al., 2016; Vogt et al., 2013). AOMI has received

growing research interest over the last decade and two hypotheses have been proposed to explain

why AOMI may be more effective as a motor skill intervention than independent AO or MI

interventions. First, Eaves and colleagues (2012, 2014, 2016) suggested the dual action simulation

hypothesis (DASH), which proposes that a person will generate separate motor representations for

the observed and imagined actions and maintain these as two parallel sensorimotor streams when

they engage in AOMI. If a person is simultaneously observing and imagining the same action, these

two motor representations are likely to merge as one sensorimotor stream, producing more

widespread activity in the premotor cortex compared to AO or MI alone. This is likely due to AOMI

increasing activity in shared brain areas for AO and MI, as well as increasing activity in areas solely

recruited during AO (e.g., inferior frontal gyrus, ventral premotor area) and MI (e.g., angular gyrus,

dorsal premotor area) of an action (Filimon et al., 2015; Hardwick et al., 2018). Second, Meers et al.

(2020) introduced the visual guidance hypothesis (VGH) as an alternative account of how AOMI may

influence action. They suggest that MI is prioritized during AOMI, and that the AO component might

merely serve as an external visual guide that facilitates more vivid MI generation. In contrast to

DASH, this would mean that AO does not activate a separate motor representation during AOMI, but

rather strengthens the motor representation resulting from MI. Irrespective of the stance taken,

both the DASH and VGH suggest that AOMI has the capacity to influence motor skill execution above

and beyond AO or MI in isolation through increased activity in motor regions of the brain.

Current neuroscientific evidence using a range of modalities supports this notion, as cortico-

motor activity is increased during AOMI of an action compared to independent AO or MI of that

same action (Eaves, Riach et al., 2016). Studies using fMRI report distinct neural signatures for AO,

MI, and AOMI whereby the blood-oxygen-level-dependent (BOLD) signal is increased and more

widespread in the brain regions involved in movement execution when an individual engages in

AOMI (e.g., Nedelko et al., 2012; Taube et al., 2015; Villiger et al., 2013). For example, Taube et al.

(2015) found greater activation in the supplementary motor area, basal ganglia and cerebellum

during AOMI compared to AO, and greater bilateral activity in the cerebellum and greater activation

in the precuneus compared to MI. Studies using electroencephalography (EEG) report that AOMI

leads to significantly larger event-related desynchronization in the mu/alpha and beta frequency

bands, indicative of increased activity over the primary sensorimotor areas of the brain compared to

both AO and MI alone (Berends et al., 2013; Eaves, Behmer et al., 2016). These fMRI and EEG

findings have important implications for applied practice, where the use of AOMI may prove

beneficial in reinforcing motor (re)learning. The increased neural activity during AOMI has the

potential to support repetitive Hebbian modulation of intracortical and subcortical excitatory

mechanisms through synaptic plasticity, in a similar manner to physical practice, and thus may be an

effective method for behavior change (Holmes & Calmels, 2008).

From a neurophysiological perspective, the first meta-analysis in this paper will focus on

studies adopting single-pulse transcranial magnetic stimulation (TMS) during AOMI, as this is the

most prevalent neuroscientific modality adopted in the literature to date (study count: n = 19 TMS, n

= 14 fMRI, n = 10 EEG). When applied to a muscle representation of the primary motor cortex, TMS

produces a twitch response in the corresponding muscles called motor evoked potentials (MEPs).

MEPs are measured using surface electromyography (EMG) and provide a marker of corticospinal

excitability during the time of stimulation (Rothwell, 1997). This approach has been used extensively

when studying the neural mechanisms for motor imagery (see e.g., Grosprêtre et al., 2016) and

action observation (see e.g., Naish et al., 2014) as it permits a non-invasive assessment of muscle-

specific M1 activity and excitability of the whole cortico-spinal pathway that is specific to the final

motor command for the action being simulated. Studies using TMS during AOMI have explored

changes in MEP amplitudes across a range of movements, including finger movements (Bruton et al.,

2020), basketball free throws (Wright, Wood et al., 2018a), walking (Kaneko et al., 2018), and

balance movements (Mouthon et al., 2016). Current literature predominantly shows increased

corticospinal excitability during AOMI compared to baseline conditions (e.g., Bruton et al., 2020;

Wright et al., 2014, 2016). However, studies comparing AOMI against AO or MI have reported

increased (e.g., Mouthon et al., 2016; Wright et al., 2016), as well as no differences (e.g., Castro et

al., 2021; Mouthon et al., 2015) in corticospinal excitability during AOMI. Given the prevalence of

TMS studies exploring the neurophysiological mechanisms underpinning AOMI engagement, it is

now possible to synthesize the available MEP amplitude data to quantify the effects of AOMI on

corticospinal excitability compared to AO, MI, and control conditions.

AOMI investigations have explored a range of movement outcomes and types, such as

movement time for ball rotations (Kawasaki et al., 2018), upper-limb kinematics for dart throwing

(Romano-Smith et al., 2019), force production in Nordic hamstring curls (Scott et al., 2018), and

radial error from the hole in golf-putting (Marshall & Wright, 2016). This research has been

conducted in sport (e.g., Romano-Smith et al., 2019) and rehabilitation (e.g., Scott et al., 2018)

contexts, with neurotypical (e.g., Di Rienzo et al., 2019) and neurodivergent (e.g., Marshall et al.,

2020) populations. The existing literature has almost exclusively demonstrated that movement

outcomes are improved for different motor skills after AOMI interventions when compared to

control conditions (e.g., Marshall et al., 2019; Romano-Smith et al., 2018; Shimada et al., 2019).

However, comparisons with AO and MI interventions are equivocal, with some studies showing

greater improvements in movement outcomes for AOMI compared to AO (e.g., Bek et al., 2019) and

MI interventions (Scott et al., 2018), and other studies showing no such effects (e.g., Romano-Smith

et al., 2019) or greater improvements for MI-related interventions (e.g., Marshall & Wright, 2016).

Therefore, it is unclear if AOMI should be recommended as the optimal simulation approach when

attempting to improve motor skill performance, highlighting the need to synthesize available

movement outcome data for AOMI interventions.

Early reviews on AOMI (Eaves, Riach et al., 2016; Vogt et al., 2013) summarized the

behavioral neuroscience literature contrasting AO and MI and drew on early AOMI research to

support its use as a motor skill intervention. Since then, research has explored changes in MEP

amplitudes and movement outcomes associated with engagement in AOMI. This has led to

population-specific reviews outlining how AOMI can be used to address sensorimotor deficits for

individuals with Parkinson's disease (Caligiore et al., 2017), developmental coordination disorder

(Scott et al., 2021), or during post-stroke rehabilitation (Emerson et al., 2018). Systematic reviews

and meta-analyses have synthesized the respective effects of MI and AO on corticospinal excitability

(e.g., Grosprêtre et al., 2016; Naish et al., 2014) and motor skill performance (e.g., Ashford et al.,

2006; Simonsmeier et al., 2021, Toth et al., 2020) when used in isolation, however, to our

knowledge, no such meta-analyses exist for AOMI interventions.

The current paper included two meta-analyses to quantify changes in corticospinal

excitability and motor skill performance for AOMI. The first meta-analysis (referred to as MEP meta-

analysis) collated and synthesized existing motor evoked potential (MEP) amplitude data from

transcranial magnetic stimulation studies as an indicator of corticospinal excitability during AOMI

engagement. The second meta-analysis (referred to as Movement meta-analysis) collated and

synthesized existing movement outcome data from behavioral studies to assess changes in motor

skill performance that result from AOMI interventions. The primary aim for both meta-analyses was

to establish the effectiveness of AOMI by comparing its effects on MEP amplitudes (first meta-

analysis) and movement outcomes (second meta-analysis) against those for AO, MI, and control

conditions. Based on previous literature (see Eaves, Riach et al., 2016), it was hypothesized that

AOMI would have a small positive effect when compared to independent AO or MI, and a medium

positive effect when compared to control conditions for both meta-analyses. The secondary aim of

this paper was to explore several methodological parameters hypothesized to have a moderating

effect on the impact of AOMI interventions on MEP amplitudes or movement outcomes across the

two meta-analyses. Accounting for the influence of these methodological aspects directly addresses

questions raised in early reviews on AOMI (Eaves, Riach et al., 2016; Vogt et al., 2013) and may

permit optimal delivery of AOMI interventions in the future.

138   **2. Method**

2.1. Study Protocol

The study procedures were performed according to the methodological guidelines

highlighted by the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA)

statement (Page et al., 2021). The justification for the two meta-analyses and proposed procedures

were documented in a pre-registration protocol document

(https://osf.io/c68ju?view_only=e6ab97909a6f4f4c8f4323390b3b3c76) that is stored alongside

additional supplementary files (https://osf.io/9yebv) on the Open Science Framework. The approach

to the two meta-analyses were similar, and the pre-screening literature search was identical (2.1.1).

However, the two meta-analyses differed in several ways, including but not limited to the inclusion

criteria, effect size preparation, and consideration of moderator effects. A PRISMA flow chart (Figure

1) outlines the study search and selection process for the two meta-analyses.

150   **[INSERT FIGURE 1 HERE]**

2.1.1. Literature Search and Selection

The literature search was performed using three online databases: PubMed, Web of Science,

and PsycINFO. The original search was run in August 2020 and the final search was run in June 2021.

Consistent search terms were decided upon and adapted for each database based on the search

string requirements (see supplementary files for full search string information). Web of Science was

used as an initial search point, followed by PubMed and PsycINFO to cover the biomedical and

psychological literature, respectively. The search was limited to studies with human populations that

had been published in a peer-reviewed journal in English language, without any limitations to

publication years. The literature search with the respective keyword combinations and restrictions

provided 396 hits in total, after 244 duplicates were removed. An exploratory search of review

articles and prior knowledge of research led to 8 additional papers being added, which took the final

number of papers being screened to 404 across the two meta-analyses.

2.1.2. Inclusion of Studies

The decision to include a study was based on five criteria. (1) The study had to either test the

effects of AOMI on corticospinal excitability by recording MEP amplitudes during single pulse TMS

(MEP meta-analysis) or test the effects of AOMI on movement outcomes by recording parameters

related to motor task performance (Movement meta-analysis). Participants were required to engage

in an AOMI task as their intervention or experimental method, and AOMI had to be delivered in a

synchronous manner (i.e., engagement in both forms of simulation at the same time). Studies with

independent AO or MI conditions were included, but only if there was a comparison with an AOMI

condition. Studies that included AOMI as an adjunct to physical practice were also included for the

Movement meta-analysis, but the inclusion of physical practice was not mandatory, and these

studies needed to incorporate physical practice in the control condition to permit assessment of

AOMI effects. (2) The study had to adopt a controlled experimental design where the unconfounded

effects of the AOMI intervention could be compared within- or between- groups. As such, studies

using a correlational or survey design, qualitative research, or case study/single-subject designs were

excluded. Studies that did not report a control group or baseline condition were also excluded,

unless they contained data from AO or MI conditions that could be compared to the AOMI

condition. (3) The study had to include an MEP amplitude (MEP meta-analysis) or movement

outcome measure (Movement meta-analysis) with a specific experimental group, or across all study

groups. Studies that did not have distinguishable MEP amplitude or movement outcome measures

or used neurophysiological modalities other than/combined with TMS (e.g., fMRI, EEG, fNIRS) were

excluded unless they also had independent MEP amplitude or movement outcome measures. (4)

The study had to report original statistics (i.e., not a re-analysis of already published findings or a

review) of the intervention on MEP amplitude or movement outcome measures for the respective

meta-analyses. (5) The study had to be published in English.

2.2. Screening Process

A three-stage screening process was adopted in this study (i.e., title, abstract, and full-text

screening). At all screening stages, the two co-first authors independently screened all articles.

Conflicts were resolved by discussion between the co-first authors in the title screen phase, with

additional input from the final author in the abstract and full text screen stages. First, all 404 titles

were screened to evaluate relevance to the MEP meta-analysis or Movement meta-analysis, with

119 titles classified as eligible for further inspection of abstracts (intercoder reliability = 94.55%).

Second, the 119 abstracts were assessed for eligibility based on the indicative content provided,

with 47 abstracts classified as eligible for further inspection as full texts (intercoder reliability =

96.64%). Third, the full text articles were assessed for eligibility based on the study inclusion criteria

for the two meta-analyses, with 34 studies progressing to data extraction (intercoder reliability =

97.87%). The mean intercoder agreement across the three stages of the screening process was

96.35%. Whenever data were missing to compute the effect sizes of interest, this information was

requested from the corresponding author for the respective article.

2.3. Data Processing

2.3.1. Data Extraction

The final 34 studies were split between the two co-first authors based on study focus, with

author two (ACV) extracting MEP data from 17 studies for the MEP meta-analysis and author one

(SC) extracting movement outcome data from 17 studies for the Movement meta-analysis,

respectively. The co-first authors extracted the primary data (i.e., means, standards deviations, and

samples sizes) and necessary methodological information to investigate the proposed moderator

effects for the present meta-analysis. Once complete, the final author blind-checked the primary

data for all studies and contacted the corresponding authors for studies with missing primary data

for the two meta-analyses. If the authors did not respond within one month, or prior to the study

data analysis cut-off date (January 2021), the article was excluded from the respective meta-

analysis. One study was excluded from each of the meta-analyses, leaving 16 studies in the final MEP

meta-analysis and 16 studies in the final Movement meta-analysis (see supplementary files for

recorded methodological information and primary data for all included articles). No studies met the

inclusion criteria for both meta-analyses, meaning the 16 studies synthesized for the MEP meta-

analysis were not included in the 16 studies synthesized for the Movement meta-analysis, and vice

versa.

2.3.2. Effect Size Preparation

For both meta-analyses, Cohen's *d* effect sizes were calculated using the mean, standard

deviation, and sample size values for the relevant outcome measures prior to conducting the main

data analyses using the 'robu' function from the 'robumeta' package (Fisher et al., 2017) in R studio

statistical software (version 2021.09.2 Build 382). The effect size preparation process varied for the

MEP meta-analysis and Movement meta-analysis. All studies included in the MEP meta-analysis

adopted a repeated measures design, meaning raw data was recorded for all conditions of interest

(i.e., AOMI, AO, MI, control). Additional steps were required for studies included in the Movement

meta-analysis to account for differences in the movement outcome variables and study designs

adopted. When a reduction in a movement outcome measure compared to control was considered

an improvement (e.g., reaction time, movement time, mean radial error), the polarity of the

calculated effect size was inverted (i.e., positive reversed to negative and vice versa) to ensure an

increased value always indicated an improvement.

231   In order to produce standardized effect sizes that could be compared across the different

study designs included in the Movement meta-analysis, effect sizes were calculated for either pre- vs

post-test gain comparisons or post- vs post-test comparisons (see supplementary materials for all

effect size calculation formulae). Pre-vs-post-test gain comparisons were used to compute effect

sizes in studies that adopted mixed experimental designs (i.e., studies that allocated participants to a

specific intervention condition and collected data pre- and post-test) as this controls for any pre-test

differences that exist between intervention condition groups (Durlak, 2009). Post- vs post-test

comparisons were used to compute effect sizes in studies that adopted within-subject (i.e.,

repeated-measures) or between-subject (i.e., independent intervention condition groups)

experimental designs. The number of effect sizes included from each study was not limited as most

studies recorded more than one outcome measure for each comparison of interest (i.e., AOMI vs

AO, MI, or control). Multiple effect sizes were extracted from studies included in the MEP meta-

analysis when MEP amplitudes were collected from more than one target muscle. Multiple effect

sizes were extracted from studies included in the Movement meta-analysis when multiple

movement outcome measures were recorded. Cohen's *d* effect size values were converted into

Hedge's *g* effect size values using a small sample size correction formula (Hedges, 1981) for

sensitivity analysis.

2.4. Data Analysis

Robust variance estimation (RVE) was used to analyze the primary effects for the MEP meta-

analysis and Movement meta-analysis. RVE was used because several studies included in each of the

two meta-analyses reported multiple relevant effect sizes which were not statistically independent

of each other (cf. Tanner-Smith & Tipton, 2014). RVE provides a method for pooling dependent

effect size estimates in the absence of any covariance values, mathematically adjusting the standard

errors of the effect sizes to account for their dependency (Tanner-Smith & Tipton, 2014; Tanner-

Smith et al., 2016). All analyses were run on RStudio using the 'metafor' package (Viechtbauer, 2010)

and the 'robumeta' package (Fisher et al., 2017).

2.4.1. Data Screening

Meta-analyses may be subject to multiple biases (Harrer et al., 2021). To address concerns

about publication bias, visual analyses were conducted on the data using funnel plots (Lau et al.,

2006), and subsequent statistical analyses were run with the Precision Effect Test (PET) using the

RVE approach (Alinaghi & Reed, 2018). For any data that showed publication bias, the trim-and-fill

procedure (Duval & Tweedie, 2000) was used to determine the number of unpublished studies

required to produce an unbiased estimate of the actual effect size. To address concerns about

sample size bias, a sensitivity analysis was run by repeating the main analyses for the two meta-

analyses with previously calculated Hedge's g values. This found no meaningful differences in effect

size estimate for the primary comparisons, so Cohen's d values are reported in both meta-analyses

(cf. Simonsmeier et al., 2021). To address concerns about between-study heterogeneity, outlier

diagnostics were completed using the FIND.OUTLIERS function and influence analyses were run

using the INFLUENCEANALYSIS function from the 'dmetar' package in R (Harrer et al., 2019). This

involved visual inspection of "baujat", "influence" for effect sizes, and "leave-one-out" plots for both

effect size and I^2 values. Potential outliers and influential effect sizes were identified across the

primary comparisons (i.e., AOMI vs AO, AOMI vs MI, AOMI vs Control) for the MEP meta-analysis and

Movement meta-analysis, respectively (see supplementary files for detailed overview of data

screening results). Removal of the outliers and influential cases had minimal impact on the pooled

effect and heterogeneity estimates for all but one comparison across the two meta-analyses. Two

effect sizes were removed from the AOMI vs AO comparison in the Movement meta-analysis ('delta'

from Frenkel-Toledo et al., 2020; 'peak angular velocity of the elbow' from Romano-Smith et al.,

2019) as the data points were deemed outliers, influential, and resulted in a meaningful change to

the pooled effect and heterogeneity when removed from the RVE data analysis for this comparison.

All other effect sizes were retained in the two meta-analyses to preserve the richness of the data.

2.4.2. Quality Assessment

Study quality was assessed to identify if the studies included in the two meta-analyses

provide reliably reported data, as well as indicating whether these studies reach an acceptable

scientific standard (Borenstein et al., 2021). The first author subjectively assessed the quality of each

study using an assessment scale employed in recent meta-analyses (see e.g., Harris et al., 2021). This

quality assessment scale was adapted from the Quality Index (Downs & Black, 1998), the Checklist

for the Evaluation of Research Articles (Durant, 1994), and the Appraisal Instrument (Genaidy et al.,

2007). The quality assessment checklist and individual scores for each study are provided in the

supplementary materials.

2.4.3. Primary and Moderator Effects

The primary data analysis for both the MEP meta-analysis and Movement meta-analysis

involved correlational RVE models run using the 'robumeta' package (Fisher et al., 2017) in RStudio.

Overall, eleven moderators were chosen across the two meta-analyses based on previous meta-

analyses and reviews focused on the effects of MI and AO on corticospinal excitability (e.g.,

Grosprêtre et al., 2016; Naish et al., 2014) and motor skill performance (e.g., Ashford et al., 2006;

Simonsmeier et al., 2021, Toth et al., 2020). Five moderators were shared across the two meta-

analyses (*action observation perspective, skill classification, guided attentional focus, kinesthetic*

imagery ability, and age). Three moderators were specific to the MEP meta-analysis (*timing of TMS*

delivery, number of TMS trials, intensity of TMS pulses). Five moderators were specific to the

Movement meta-analysis (*population type, physical practice, incorporation of PETTLEP principles,*

context, and *intervention volume*)*.* Subgroup analyses and meta regressions were used to examine if

these moderators influenced the effects of AOMI compared to aggregate data from the AO, MI, and

control groups/conditions for the two meta-analyses. Subgroup analyses were used to compare the

effects of AOMI based on moderators that consisted of nominal data across the two meta-analyses.

The sub-group analyses for the MEP meta-analysis included *action observation perspective, skill*

classification, guided attentional focus, timing of TMS delivery moderators*. The sub-group analyses*

for the Movement meta-analysis included population type, action observation perspective, skill

classification, guided attentional focus, physical practice, incorporation of PETTLEP principles, and

context moderators. Meta regression analyses were used to assess if moderators that consisted of

interval data predicted the effects of AOMI. Meta regression analyses for the MEP meta-analysis

included kinesthetic *imagery ability, age, number of TMS trials, and intensity of TMS pulses.* Meta

regression analyses for the Movement meta-analysis included kinesthetic *imagery ability, age, and*

intervention volume.

2.4.3.1. Moderators for Both Meta-Analyses.

2.4.3.1.1. Action Observation Perspective. Studies have used first-person perspective AO

(e.g., Bruton et al., 2020; Romano-Smith et al., 2019) and third-person perspective AO (e.g., Taube et

al., 2014; Wright, Wood et al., 2018a) to examine the effects of AOMI on MEP amplitudes and

movement outcomes. Sub-group analyses were used to compare the effects of AOMI using first-

person vs third-person AO perspectives for both meta-analyses. First-person perspective AO

involved the participant viewing the action as if they were performing it (i.e., through their own

eyes) and third-person perspective AO involved the participant viewing the action as if another

person video recorded them or another person was performing the action (i.e., filmed from a

vantage point away from the body). This was determined by checking written text and visual stimuli

included in the article.

2.4.3.1.2. Skill Classification. Diverse motor tasks ranging from finger movements (e.g.,

Meers et al., 2020) to walking (e.g., Kaneko et al., 2018) have been used in previous AOMI literature.

Sub-group analyses were used to compare the effects of AOMI for fine vs gross and continuous vs

discrete motor tasks for both meta-analyses. The target movement presented in the AOMI stimuli

was classified using a one-dimensional skill classification approach (Spittle, 2021, p.23). Based on

this approach, intricate and precise movements using smaller muscle groups (e.g., finger

movements) were classed as fine motor tasks; larger muscle movements typically based on

fundamental movement patterns (e.g., balance tasks) were classed as gross motor tasks; repetitive

movements that have no distinct beginning or end (e.g., walking) were classed as continuous motor

tasks; and movements that have an identifiable beginning and end (e.g., putting a golf ball) were

classed as discrete motor tasks. Other skill classification comparisons (e.g., open vs closed skills)

were not considered in this moderator category due to a lack of coverage in the synthesized

literature.

2.4.3.1.3. Guided Attentional Focus. Studies on AO have demonstrated different effects on

movement outcomes (e.g., D'Innocenzo et al., 2016) and MEP amplitudes (e.g., Wright, Wood et al.,

2018b) when visual attention is directed, or not, to a specific component of the observed action.

More recently, Bruton et al. (2020) showed that allocation of visual attention modulates the effects

of AOMI on MEP amplitudes for a finger movement task. Sub-group analyses were used to compare

the effects of AOMI for guided attentional focus (i.e., use of instructions to direct attention towards

a specific aspect of the observed movement) vs unguided attentional focus (i.e., no such

instructions) for both meta-analyses. Studies that did not explicitly state if visual attention was

manipulated during AOMI were included in the unguided attentional focus sub-group for each meta-

analysis.

2.4.3.1.4. Kinesthetic Imagery Ability. Kinesthetic imagery is the imagery modality

instructed during AOMI, and the effects of AOMI on movement outcomes reportedly vary as a

function of kinesthetic imagery ability (McNeill et al., 2020). Meta regression analyses were used to

assess if there was a relationship between kinesthetic imagery ability score and the effects of AOMI

for both meta-analyses. Kinesthetic imagery ability data recorded using valid self-report

psychometric scales including the Vividness of Movement Imagery Questionnaire -2 (Roberts et al.,

2008), Movement Imagery Questionnaire-3 (Williams et al., 2012) and the Movement Imagery

Questionnaire-Revised (Hall & Martin, 1997) were included for moderator analyses. Any studies that

used non-validated scales such as a visual analogue scale were excluded from the moderator

analyses. The imagery ability data was extracted from the studies and standardized by reverse-

scoring any measures that adopted an inverse scoring system such that higher numbers meant

better imagery ability, before converting all scores to percentage values based on the range of

values attainable for each scale.

2.4.3.1.5. Age. Studies have typically recruited adults ranging from early to middle

adulthood when assessing MEP amplitudes during AOMI (mean sample age = 27.07 ± 13.48 years)

and movement outcomes after AOMI (mean sample age = 30.89 ± 20.24 years). Studies have shown

age-related differences in MEP amplitudes during simulation of actions (e.g., Mouthon et al., 2016)

and imagery ability is proposed to decline across the lifespan (e.g., Gulyás et al., 2022), suggesting

that age may moderate the effects of AOMI on MEP amplitudes and movement outcomes. Meta

regression analyses were used to assess if there was a relationship between participant age and the

effects of AOMI for both meta-analyses.

2.4.3.2. Moderators for MEP Meta-Analysis.

2.4.3.2.1. Timing of Transcranial Magnetic Stimulation Delivery. AO and MI cause phase-

specific changes in corticospinal excitability (see e.g., Grosprêtre et al., 2016; Naish et al., 2014 for

reviews). Sub-group analysis was used in the MEP meta-analysis to compare the effects of AOMI on

MEP amplitudes for TMS delivered at a random point after movement onset against TMS delivered

at a targeted point after movement onset (e.g., at the point of maximum movement of the limb).

Timing of TMS stimulation delivery refers to the point at which the single pulse is delivered based on

the movement displayed in the visual stimuli during AOMI.

2.4.3.2.2. Number of Transcranial Magnetic Stimulation Trials. The number of TMS trials

impacts the reliability of the MEP's evoked during single-pulse TMS (Goldsworthy et al., 2016). Meta

regression analysis was used in the MEP meta-analysis to assess if there was a relationship between

the number of TMS trials and the effects of AOMI on MEP amplitudes. This was calculated by

recording the number of trials where single-pulse TMS was applied to the participant during AOMI

for each study.

2.4.3.2.3. Intensity of Transcranial Magnetic Stimulation Pulses. The intensity of TMS

pulses impacts the reliability of the MEP's evoked during single-pulse TMS (Pellegrini et al., 2018).

Meta regression analysis was used in the MEP meta-analysis to assess if there was a relationship

between intensity of TMS pulses and the effects of AOMI on MEP amplitudes. TMS stimulation

intensity refers to the intensity of the TMS stimulator output relative to the resting motor threshold,

that is applied to the participant during AOMI.

2.4.3.3. Moderators for Movement Meta-Analysis.

2.4.3.3.1. Population Type. AOMI interventions have been shown to benefit movement

outcomes in both neurotypical and neurodivergent populations (e.g., Scott et al., 2019). Sub-group

analysis was used in the Movement meta-analysis to compare the effects of AOMI on movement

outcomes for these two population types. Neurotypical populations included individuals who are not

characterized by neurologically atypical patterns, thoughts, behavior, or diagnoses, and

neurodivergent populations included individuals whose neurological development and state are

considered atypical.

2.4.3.3.2. Physical Practice. Studies have explored the effects of AOMI interventions on

movement outcomes with (e.g., Marshall & Wright, 2016) and without (e.g., Taube et al., 2014)

physical practice. Sub-group analysis was used in the Movement meta-analysis to compare the

effects of AOMI on movement outcomes when used with physical practice vs without physical

practice.

2.4.3.3.3. Incorporation of PETTLEP Principles. Some studies have adhered to PETTLEP

principles (Holmes & Collins, 2001) when developing and delivering AOMI interventions (e.g.,

Romano-Smith et al., 2019). Sub-group analysis was used in the Movement meta-analysis to

compare the effects of AOMI on movement outcomes with the inclusion of PETTLEP principles vs

without inclusion of PETTLEP principles.

407   *2.4.3.3.4. Context.* AOMI interventions have been used to target changes in movement

outcomes in sport (e.g., Romano-Smith et al., 2018) and rehabilitation (e.g., Marusic et al., 2018)

contexts. Sub-group analysis was used in the Movement meta-analysis to compare the effects of

AOMI on movement outcomes for sport vs rehabilitation vs other contexts. Studies were classified

as sport- or rehabilitation-focused based on the movement being simulated and performed. Studies

including movements that did not clearly fall into sports or rehabilitation contexts (e.g., finger

movements, ball rotation tasks) were classified as 'other'.

414   *2.4.3.3.5. Intervention Volume.* Studies have delivered AOMI interventions over short- (e.g.,

415   Bek et al., 2019) and longer-term (e.g., Shimada et al., 2019) periods when investigating their effects

416   on movement outcomes. Meta regression analysis was used in the Movement meta-analysis to

417   assess the relationship between intervention volume (total minutes) and the effects of AOMI on

418   movement outcomes.

419   **3. Results**

420   **3.1. Study Characteristics**

421   Overall, the two meta-analyses synthesized 111 effect sizes from 32 studies. Of these, the

MEP meta-analysis included 54 effect sizes (n = 16 studies) and the Movement meta-analysis

included 57 effect sizes (n = 16 studies). Studies included across the two meta-analyses were

published between 2009 and 2021, with a total sample size of 823 participants split across studies

included in the MEP meta-analysis (n = 234, 77 females, 92 males, 65 undisclosed) and studies

included in the Movement meta-analysis (n = 589, 281 females, 308 males). The mean age of

participants was 27.07 ± 13.48 years and 30.89 ± 20.24 for the two meta-analyses, respectively.

3.2. Study Quality

429   The study quality assessment indicated that all studies included in the two meta-analyses

displayed a high degree of rigor. For studies included in the MEP meta-analysis, the quality

assessment scores ranged from 18.75-100%, with a mean of 89.58 ± 22.99%. The most poorly

addressed items were 'providing details of a priori sample size determination' and 'consistently

reporting effect sizes' with 18.75% and 43.75% of studies satisfying these criteria, respectively

(Figure 2a). For studies included in the Movement meta-analysis, the quality assessment scores

ranged from 31.25-100%, with a mean of 92.36 ± 17.36%. The most poorly addressed items were

'providing details of a priori sample size determination' and 'applicability of study results to other

relevant populations' with 31.25% and 68.75% of studies satisfying these criteria, respectively

(Figure 2b).

To assess whether studies with smaller samples sizes or lower quality studies were likely to

bias the results, meta-regression analyses were run between quality assessment scores and effect

size, and sample size and effect size using the '*robumeta*' and 'metafor' packages in R. For studies

included in the MEP meta-analysis, the analysis used 54 effect sizes from 16 studies and reported

that neither quality assessment score (b = -0.02, p = .25), nor sample size (b = -0.01, p = .13)

predicted the overall effect of AOMI on MEP amplitudes. Similarly, for studies included in the

Movement meta-analysis, the analysis used 57 effect sizes from 16 studies and reported that neither

quality assessment score (b = 0.01, p = .76), nor sample size (b = -0.03, p = .06) predicted the overall

effect of AOMI on movement outcomes. The non-significant relationships between quality

assessment score, sample size and effect size indicate a low risk of bias for the studies included

across the two meta-analyses (*cf*. Harris et al., 2021).

450   **[INSERT FIGURE 2 HERE]**

3.3. MEP Meta-Analysis

Fifty-four effect sizes from sixteen studies were used in the MEP meta-analysis to determine

the overall effect of AOMI on MEP amplitudes. AOMI had a small to medium positive overall effect

on MEP amplitudes compared to the control, AO and MI conditions in combination (d = 0.48, 95% CI

[0.35, 0.61], p < .001). The between-study heterogeneity variance was estimated at τ^2 = 0.00, with

an I^2 value of 1.23%. The MEP meta-analysis reported no significant moderators (Tables 1 and 2),

demonstrating a robust effect of AOMI on MEP amplitudes irrespective of kinesthetic imagery

ability, sample age, intensity of TMS pulses, number of TMS trials, sample age, AO perspective,

attentional focus strategy, skill classification, and the timing of TMS delivery.

460   **[INSERT TABLE 1 HERE]**

461   **[INSERT TABLE 2 HERE]**

462   ***3.3.1. AOMI vs Control***

463   Nineteen effect sizes from thirteen studies were used to compare AOMI vs control

464   conditions in the MEP meta-analysis. AOMI had a medium positive effect on MEP amplitudes

465   compared to control conditions (d = 0.54, 95% CI [0.41, 0.66], p < .001). The between-study

466   heterogeneity variance was estimated at τ^2 = 0.00, with an I^2 value of 0.00%.

467   **[INSERT FIGURE 3 HERE]**

468   ***3.3.2. AOMI vs AO***

469   Twenty-three effect sizes from thirteen studies were used to compare AOMI vs AO

470   conditions in the MEP meta-analysis. AOMI had a small to medium positive effect on MEP

471   amplitudes compared to AO conditions (d = 0.45, 95% CI [0.27, 0.63], p < .001). The between-study

472   heterogeneity variance was estimated at τ^2 0.02, with an I^2 value of 10.32%.

473   **[INSERT FIGURE 4 HERE]**

474   ***3.3.3. AOMI vs MI***

475   Twelve effect sizes from six studies were used to compare AOMI vs MI conditions in the MEP

476   meta-analysis. AOMI had no significant effect on MEP amplitudes compared to MI conditions (d =

477   0.25, 95% CI [-0.13, 0.63], p = .14). The between-study heterogeneity variance was estimated at τ^2=

478   0.11, with an I^2 value of 42.92%.

479   **[INSERT FIGURE 5 HERE]**

480   **3.4. Movement Meta-Analysis**

481   Fifty-seven effect sizes from sixteen studies were used in the Movement meta-analysis to

determine the overall effect of AOMI on movement outcomes. AOMI had a small to medium positive

overall effect on movement outcomes compared to the control, AO and MI conditions in

combination (d = 0.48, 95% CI [0.18, 0.78], p < .01). The between-study heterogeneity variance was

estimated at τ^2 = 0.39 , with an I^2 value of 69.68%. The Movement meta-analysis reported no

significant moderators (Tables 1 and 3), demonstrating a robust effect of AOMI on movement

outcomes irrespective of intervention volume, kinesthetic imagery ability, sample age, AO

perspective, study context, attentional focus strategy, incorporation of PETTLEP, physical practice,

population type, and skill classification.

490   **[INSERT TABLE 3 HERE]**

491   ***3.4.1. AOMI vs Control***

492   Twenty-seven effect sizes from twelve studies were used to compare AOMI vs control

493   conditions in the Movement meta-analysis. AOMI had a medium to large positive effect on

494   movement outcomes compared to control conditions (d = 0.67, 95% CI [0.16, 1.18], p = .02). The

495   between-study heterogeneity variance was estimated at τ^2 = 0.48, with an I^2 value of 70.74%.

496   **[INSERT FIGURE 6 HERE]**

497   ***3.4.2. AOMI vs AO***

498   Nineteen effect sizes from nine studies were used to compare AOMI vs AO conditions in the

499   Movement meta-analysis. AOMI had a small to medium positive effect on movement outcomes

500   compared to AO conditions (d = 0.44, 95% CI [0.07, 0.81], p = .03). The between-study heterogeneity

501   variance was estimated at τ^2= 0.24, with an I^2 value of 63.23%.

502   **[INSERT FIGURE 7 HERE]**

503   **3.3.2.3. AOMI vs MI.** Eleven effect sizes from six studies were used to compare AOMI vs MI

504   conditions in the Movement meta-analysis. AOMI had no significant effect on movement outcomes

505   compared to MI conditions (d = 0.53, 95% CI [-0.59, 1.66], p = .28). The between-study heterogeneity

506   variance was estimated at τ^2 = 1.30, with an I^2 value of 84.46%.

507   **[INSERT FIGURE 8 HERE]**

3.5. Publication Bias

Based on the funnel plots of effect size level (see supplementary files), publication bias was

identified to be unlikely for all comparisons across the two meta-analyses. Regardless, we ran trim

and fill and PET analyses using the RVE method to retrieve an unbiased effect size estimate

corrected for publication bias for all comparisons made in the MEP meta-analysis and Movement

meta-analysis.

For the control comparison in the MEP meta-analysis, both the PET-intercept (b_0 = 0.09, p =

0.83) and the PET-slope (b_1 = 1.21, p = 0.32) were not statistically significant, suggesting that

publication bias was unlikely. For the AO comparison in the MEP meta-analysis, the funnel plot of

effect size level (Figure 9, left) indicated asymmetry. Trim-and-fill analysis proposed 4 missing values

(Figure 9, right), and the effect size changed from a medium (d = 0.54) to a small to medium positive

effect (d = 0.43), suggesting minimal effects of publication bias for this dataset. For the PET, both the

PET-intercept (b_0 = -1.00, p = 0.16) and the PET-slope (b_1 = 3.85, p = 0.07) were not statistically

significant, suggesting that publication bias was unlikely. For the MI comparison in the MEP meta-

analysis, the funnel plots indicated no funnel asymmetry and the trim-and-fill analysis reported zero

missing values. For the PET, both the PET-intercept (b_0 = -1.14, p = 0.43) and the PET-slope (b_1 = 3.59,

p = 0.41) were not statistically significant, suggesting that publication bias was unlikely.

For the control comparison in the Movement meta-analysis, the funnel plots and trim and fill

analysis showed no signs of asymmetry. Both the PET-intercept (b_0 = -0.38, p = 0.56) and the PET-

slope (b_1 = 4.34, p = 0.26) were not statistically significant, suggesting that publication bias was

unlikely. For the AO comparison in the Movement meta-analysis, the funnel plots and trim-and-fill

analysis showed no signs of asymmetry. Both the PET-intercept (b_0 = -0.08, p = .88) and the PET-

slope (b_1 = 2.04, p = .43) were not statistically significant, suggesting that publication bias was

unlikely. For the MI comparison in the Movement meta-analysis, the funnel plots and trim and fill

analysis showed no signs of asymmetry. Both the PET-intercept (b_0 = -0.51, p = .62) and the PET-

slope (b_1 = 3.65, p = .48) were not statistically significant, suggesting that publication bias was

unlikely.

535   **[INSERT FIGURE 9 HERE]**

536   **4. Discussion**

Since the early reviews introducing AOMI (e.g., Eaves, Riach et al., 2016; Vogt et al., 2013),

researchers have studied both its neurophysiological and behavioral effects as a motor simulation

intervention. The current paper included two meta-analyses to quantify changes in corticospinal

excitability and motor skill performance for AOMI. The MEP meta-analysis collated and synthesized

existing MEP amplitude data from transcranial magnetic stimulation studies as an indicator of

corticospinal excitability during AOMI engagement. The Movement meta-analysis collated and

synthesized existing movement outcome data from behavioral studies to assess changes in motor

skill performance that result from AOMI interventions. The primary aim for both meta-analyses was

to establish the effectiveness of AOMI by comparing its effects on MEP amplitudes (first meta-

analysis) and movement outcomes (second meta-analysis) against those for AO, MI, and control

conditions. Based on previous literature (see Eaves, Riach et al., 2016), it was hypothesized that

AOMI would have a small positive effect compared to independent AO or MI, and a moderate

positive effect compared to control conditions, for both outcome variables. The results of the two

meta-analyses partially support this hypothesis. For the MEP meta-analysis, AOMI had a small to

medium positive overall effect, a medium positive effect compared to control conditions, a small to

medium positive effect compared to AO, and no significant effect compared to MI. For the

movement outcome data, AOMI had a small to medium positive overall effect, a medium to large

positive effect compared to control, a small to medium positive effect compared to AO, and no

significant effect compared to MI conditions.

4.1. MEP Meta-Analysis

In the MEP meta-analysis, AOMI had a medium positive effect compared to control

conditions and a small to medium positive effect compared to AO, but showed no effect compared

to MI. TMS studies have consistently reported increased corticospinal facilitation for AOMI across

diverse motor tasks such as simple finger movements (Bruton et al., 2020), walking (Kaneko et al.,

2018), and basketball free throws (Wright, Wood et al., 2018a). From a theoretical standpoint, the

finding that corticospinal excitability was facilitated for AOMI compared to control and AO but not

MI conditions aligns with the propositions of the VGH that AOMI is driven by MI but may oppose the

sentiments of the DASH (Eaves et al., 2012, 2014). The VGH suggests that observed and imagined

actions are not co-represented, and MI is the driver for increases in motor activity during AOMI.

Specifically, Meers et al. (2020) suggest the AO component acts as a visual primer, facilitating the

production of more vivid images during AOMI compared to AO and MI conditions. Alternatively, the

DASH proposes that concurrent representations of observed and imagined actions can be

maintained as two quasi-encapsulated sensorimotor streams and that these will merge, rather than

compete, when a person is overtly and covertly simulating the same action during AOMI. The

merging of these two sensorimotor streams is likely to produce more widespread activity in the

premotor cortex (see Filimon et al., 2015) than the AO, MI and control conditions, contributing to

increased corticospinal excitability via cortico-cortical connections linking premotor and motor

cortices (Fadiga et al., 2005).

575   The non-significant increase in MEP amplitudes for AOMI compared to MI reported in the

MEP meta-analysis may be explained by the propositions of the VGH, as the increased imagery

vividness for AOMI vs MI could be expected to be represented by a smaller difference in MEP

amplitude between these two conditions. This difference could be expected to be negligible if those

simulating actions were able to generate clear and vivid kinesthetic imagery without a visual primer,

as was the case for the participants synthesized in the MEP meta-analysis (mean normalized

kinesthetic imagery ability score = 67.41%, median = 70.83%, range = 55% - 76.33%). Current

evidence is conflicting for the VGH and DASH accounts of AOMI (see Bruton et al., 2020; Meers et al.,

2020), but both hypotheses offer feasible explanations for the impact of AOMI on the motor system

and thus warrant further systematic investigation.

4.2. Movement Meta-Analysis

In the Movement meta-analysis, AOMI had a medium to large positive effect on movement

outcomes compared to control conditions and a small to medium positive effect compared to AO

conditions. Such positive effects are evidenced across most studies included in the Movement meta-

analysis, with movements ranging from dart throwing (Romano-Smith et al., 2018, 2019) to whole-

body balance tasks (Taube et al., 2014) in both neurotypical (e.g., Aoyama et al., 2020) and

neurodivergent populations (e.g., Marshall et al., 2020). The increased motor activity during AOMI,

as discussed in the previous section, is a possible neurophysiological mechanism for this effect on

movement outcomes. Repeated engagement in AOMI, and thus activation of the motor system, has

the potential to support repetitive Hebbian modulation of intracortical and subcortical excitatory

mechanisms through synaptic plasticity, in a similar manner to physical practice (Holmes & Calmels,

2008). From a cognitive perspective, AO and MI help develop mental representations that comprise

cognitive information relating to movement execution (Frank et al., 2020). When executing a motor

task, a person recalls the relevant mental representation and uses this to guide their movement

(Frank et al., 2020). AO and MI are proposed to contribute differently to the development of such

mental representations, with AO providing sequential and timing information and MI providing

sensory information related to the movement (Kim et al., 2017). It is possible combining the two

forms of motor simulation during AOMI allows for the effective development of mental

representations of action in the long-term memory, benefitting the physical execution of a motor

task (Frank et al., 2020; Kim et al., 2017; Wright et al., 2021).

In contrast, the results of the Movement meta-analysis showed that AOMI had no significant

effect on movement outcomes compared to MI conditions. Robust evidence supports the efficacy of

MI as an intervention to improve motor performance across settings (e.g., MI: Guillot & Collet,

2008). This null finding aligns with the effects of AOMI on corticospinal excitability when compared

to MI in the MEP meta-analysis. Specifically, AOMI did not increase corticospinal excitability or

improve movement outcomes when compared to MI conditions across the two meta-analyses

conducted in this paper. This provides further support for the VGH account for AOMI (Meers et al.,

2020), suggesting that the imagery component drives the effects of AOMI on both the motor system

of the brain and subsequent adaptations to physical movement. The sample synthesized for the MI

comparison in the Movement meta-analysis were more able imagers (mean normalized kinesthetic

imagery ability score = 74.78%, range = 56.67% - 96.17%) than the sample synthesized for the MI

comparison in the MEP meta-analysis (mean normalized kinesthetic imagery ability score = 67.41%,

range = 55.00% - 76.33%). This could suggest that individuals with high imagery ability benefit less

from AOMI because the visual primer provided by AO does not improve the vividness or clarity of

their MI during AOMI.

The findings of the Movement meta-analysis promote AOMI as an effective alternative

intervention to AO and MI as well-established approaches, but do not indicate that combining AO

and MI simultaneously (i.e., AOMI) has an additive benefit towards motor performance compared to

MI. It is worthwhile noting that AOMI had a small to medium positive effect on movement outcomes

compared to MI (d = 0.53) despite the lack of significant differences reported in the Movement

meta-analysis. This is an important consideration in applied settings, such as sport and

neurorehabilitation, where marginal improvements in motor performance can have practical

significance (Lakens, 2013). AOMI interventions are a suitable alternative to AO and MI interventions

as this combined approach can address the reported limitations of using either simulation technique

in isolation. The capacity to generate and maintain mental images, termed 'imagery ability', is a

complex cognitive process that is variable within- and between-populations (Cumming & Eaves,

2018). Individuals with low imagery ability typically find it difficult to generate and control imagined

content during MI interventions, an issue that is not present for AO interventions as specific

movement content can be displayed via video (Holmes & Calmels, 2008). However, the effectiveness

of AO interventions is dependent on the observer's ability to attend to the most important aspects

of the motor task being performed (D'Innocenzo et al., 2016). Based on current recommendations

for delivering AOMI interventions (see Wright et al., 2021), AOMI has the capacity to control the

visual information displayed via AO whilst directing the individual's attention by getting them to

focus on kinesthetic aspects of the movement, subsequently reducing the complexity of MI.

4.3. Limitations and Future Research Recommendations

4.4.1. Study Reporting

A secondary aim of this paper was to explore several methodological parameters

hypothesized to have a moderating effect on the impact of AOMI interventions on MEP amplitudes

or movement outcomes across the two meta-analyses. This was conducted to try and understand

the influence of key methodological aspects raised in early reviews on AOMI (Eaves, Riach et al.,

2016; Vogt et al., 2013) and to provide recommendations for future research and delivery of AOMI

interventions. Whilst moderator analyses were run in the form of meta-regression and sub-group

analyses, only the sub-group analyses included the full sets of studies for each meta-analysis (i.e., 16

studies per analysis), with missing study information meaning that meta-regression analyses

included 75% of the studies on average across the two meta-analyses (mean = 12 studies, min = 6

studies for kinesthetic imagery ability in the MEP meta-analysis, max = 16 studies for sample age in

the Movement meta-analysis). The issue of inadequate reporting has been raised in recent meta-

analyses focusing on imagery interventions, with both noting issues related to imagery ability and

assessment and reporting across studies (see e.g., Simonsmeier et al., 2021; Toth et al., 2020). A

recent article has provided guidance for authors to standardise and improve the quality of reporting

for action simulation studies (Moreno-Verdú et al., 2022). Alongside adhering to these useful

guidelines, we also recommend that authors address the apparent biases in AOMI literature made

evident by the unbalanced population sizes compared in the sub-group analyses for the two meta-

analyses reported in this paper. For TMS studies, we suggest researchers employ guided attentional

focus, include more diverse motor tasks (i.e., gross/continuous/serial/open skills), and test the

impact of the timing of TMS delivery on the effects of AOMI on MEP amplitudes. For behavioral

studies, we recommend that researchers employ guided attentional focus, adhere to PETTLEP

guidelines, incorporate physical practice, recruit neurodivergent populations, and include more

diverse motor tasks (i.e., gross/continuous/serial/open skills) when studying the effects of AOMI on

movement outcomes.

4.4.1. Neurophysiological Modality

The MEP meta-analysis included in this paper synthesized MEP amplitude data from AOMI

studies using single-pulse TMS. TMS has been widely used as a neurophysiological modality with

AOMI as it permits the recording of muscle-specific facilitation in corticospinal excitability, an effect

that has been demonstrated robustly for AO and MI (Naish et al., 2014; Grosprêtre et al., 2016).

Whilst single-pulse TMS provides an indication of activity within the motor and premotor cortices of

the brain during AOMI, EEG and fMRI can be used to provide complimentary knowledge about the

roles of other cortical regions during AOMI as they can measure whole-brain activity and have high

temporal and spatial accuracy, respectively (Holmes & Wright, 2017). Studies have shown that AOMI

leads to activity in brain regions that would not be activated directly during the delivery of TMS to

the primary motor cortex (e.g., rostral prefrontal cortex; Eaves, Behmer, et al., 2016). Activity for

these brain areas is not represented within the MEP meta-analysis conducted in this paper.

Consequently, there is a need to collate and synthesize data on the precise anatomical substrates

involved in AOMI using neuroscientific methods with increased spatial resolution. Hardwick et al.

(2018) recently performed a large-scale activation likelihood estimation meta-analysis on fMRI data

for AO, MI, and movement execution to identify distinct and shared neural regions for these three

action states. This approach could be adopted to advance understanding of the neural mechanisms

underpinning engagement in AOMI once additional fMRI data is available for this form of simulation.

4.4.2. Study Designs

To-date, the AOMI studies synthesized in the Movement meta-analysis have almost entirely

explored the short-term effects of this intervention on movement outcomes, using a between-

groups comparison at one time point or adopting a pre- vs post-test study design. Whilst this

approach is typical in randomized controlled trials of interventions, this does not permit accurate

assessment of the long-term changes that result from AOMI engagement. The benefits of AO and MI

on movement outcomes are reportedly greatest during or immediately after training, with the

positive outcomes gradually disappearing in the absence of simulated practice (Stevens et al., 2003;

Zhang et al., 2019). However, the performance benefits of MI training are retained beyond the

intervention period (Simonsmeier et al., 2021), with repetitive engagement in MI inducing neural

plasticity during recovery phases when this technique is used to acquire a skill (Ruffino et al., 2017).

It remains unclear if the improvements in movement outcomes associated with AOMI are

maintained after the intervention is withdrawn. Future studies should draw from the methodological

approaches adopted in motor learning literature (e.g., Krakauer et al., 2019) to comprehensively

examine the effectiveness of AOMI when learning and improving movement outcomes for different

populations and motor tasks.

4.4.3. Brain-Behavior Interactions

The MEP meta-analysis synthesized data on MEP amplitudes during AOMI engagement and

the Movement meta-analysis synthesized data on changes in movement outcomes after AOMI

interventions. The crossover between these effects could not be analyzed in the current paper as no

studies to our knowledge have collected both measures for AOMI whilst satisfying the inclusion

criteria for the two meta-analyses reported in this paper. The results of the two meta-analyses show

that AOMI had a positive effect on MEP amplitudes and movement outcomes compared to control

and AO conditions. Collecting neurophysiological responses, such as MEP amplitudes, alongside

movement outcome measures will advance our understanding of the relationship between the

changes in brain activity and motor skill performance for this intervention. Such integration of

measures has shown that corticospinal excitability during MI is related to the magnitude of motor

cortical adaptations after MI training (Yoxon & Welsh, 2020), and that MI and physical practice lead

to different changes in brain activity and subsequent movement execution after equal training bouts

(Kraeutner et al., 2020). Future research should combine neuroscientific and human movement

science methods to elucidate the neural mechanisms underlying improvements in motor skill

performance and learning through AOMI.

5. Conclusion

The two meta-analyses included in this paper synthesize the existing MEP amplitude and

movement outcome data for AOMI to compare its effectiveness against AO, MI, and control

conditions. The results of the MEP meta-analysis report that AOMI has a small to medium positive

overall effect on MEP amplitudes, as an indicator of corticospinal excitability. When compared to

different conditions, AOMI has a medium positive effect on MEP amplitudes compared to control, a

small to medium positive effect compared to AO, and no significant effect compared to MI

conditions. For the Movement meta-analysis, AOMI had a small to medium positive overall effect on

movement outcomes. When compared to different conditions, AOMI has a medium to large positive

effect on movement outcomes compared to control, a small to medium positive effect compared to

AO, and no significant effect compared to MI conditions. No methodological factors moderated the

effects of AOMI in either of the two meta-analyses, indicating a robust effect of AOMI for both

outcome variables. However, it should be noted that inadequate reporting of methodological

information as well as limited variation in the current literature on AOMI may have resulted in

biased comparisons being made between moderator sub-groups and low powered assessments of

relationships across the two meta-analyses. Overall, the results of both meta-analyses support the

effectiveness of AOMI as an alternative intervention to AO and MI, two well established

interventions, as AOMI engagement addresses the limitations of using these approaches in isolation

when targeting increased activity in motor regions of the brain and improvements in motor skill

performance. A more methodologically diverse approach that integrates brain and behavior is

needed in future AOMI research to advance the current state of knowledge for this intervention.

Funding statement: This research did not receive any specific grant from funding agencies in the

public, commercial, or not-for-profit sectors.

6. References

Alinaghi, N., & Reed, W. R. (2018). Meta-analysis and publication bias: How well does the FAT-PET-

PEESE procedure work? Research Synthesis Methods, 9(2), 285–311.

https://doi.org/10.1002/jrsm.1298

Aoyama, T., Kaneko, F., & Kohno, Y. (2020). Motor imagery combined with action observation

training optimized for individual motor skills further improves motor skills close to a

plateau. *Human Movement Science*, *73*, 102683.

https://doi.org/10.1016/j.humov.2020.102683

Ashford, D., Bennett, S. J., & Davids, K. (2006). Observational modeling effects for movement

dynamics and movement outcome measures across differing task constraints: A meta-

analysis. *Journal of Motor Behavior*, *38*(3), 185-205. https://doi.org/10.3200/JMBR.38.3.185-

205

Bek, J., Gowen, E., Vogt, S., Crawford, T. J. & Poliakoff, E. (2019). Combined action observation and

motor imagery influences hand movement amplitude in Parkinson's disease. *Parkinsonism*

and Related Disorders, 61, 126-131. https://doi.org/10.1016/j.parkreldis.2018.11.001

Berends, H., Wolkorte, R., Ijzerman, M., & van Putten, M. (2013). Differential cortical activation

during observation and observation-and-imagination. *Experimental Brain Research*, *229*(3),

337-345. https://doi.org/10.1007/s00221-013-3571-8

Borenstein, M., Hedges, L. V., Higgins, J. P., & Rothstein, H. R. (2021). *Introduction to meta-analysis*.

John Wiley & Sons.

Bruton, A., Holmes, P., Eaves, D., Franklin, Z., & Wright, D. (2020). Neurophysiological markers

discriminate different forms of motor imagery during action observation. *Cortex*, *124*, 119-

136. https://doi.org/10.1016/j.cortex.2019.10.016

Buccino, G. (2014). Action observation treatment: A novel tool in neurorehabilitation. *Philosophical*

Transactions of the Royal Society B: Biological Sciences, *369*(1644), 20130185.

https://doi.org/10.1098/rstb.2013.0185

Caligiore, D., Mustile, M., Spalletta, G., & Baldassarre, G. (2017). Action observation and motor

imagery for rehabilitation in Parkinson's disease: A systematic review and an integrative

hypothesis. *Neuroscience & Biobehavioral Reviews*, *72*, 210-222.

https://doi.org/10.1016/j.neubiorev.2016.11.005

Castro, F., Bryjka, P.A., Di Pino, G., Vuckovic, A., Nowicky, A. & Bishop, D. (2021). *Brain and Cognition*,

152, 105768. https://doi.org/10.1016/j.bandc.2021.105768

Cumming, J., & Eaves, D. L. (2018). The nature, measurement, and development of imagery

ability. *Imagination, Cognition and Personality*, *37*(4), 375-393.

https://doi.org/10.1177%2F0276236617752439

D'Innocenzo, G., Gonzalez, C. C., Williams, A. M., & Bishop, D. T. (2016). Looking to learn: the effects

of visual guidance on observational learning of the golf swing. *PloS one, 11*(5), e0155442.

https://doi.org/10.1371/journal.pone.0155442

De Vries, S., & Mulder, T. (2007). Motor imagery and stroke rehabilitation: A critical

discussion. *Journal of Rehabilitation Medicine*, *39*(1), 5-13.

https://doi.org10.2340/16501977-0020

Di Rienzo, F., Joassy, P., Kanthack, T., MacIntyre, T., Debarnot, U., & Blache, Y. et al. (2019). Effects of

Action Observation and Action Observation Combined with Motor Imagery on Maximal

Isometric Strength. *Neuroscience, 418*, 82-95.

https://doi.org/10.1016/j.neuroscience.2019.08.025

Downs, S. H., & Black, N. (1998). The feasibility of creating a checklist for the assessment of the

methodological quality both of randomised and non-randomised studies of health care

interventions. *Journal of Epidemiology & Community Health*, *52*(6), 377-384.

http://dx.doi.org/10.1136/jech.52.6.377

Durant, J. (1994). What is scientific literacy? *European Review*, *2*(1), 83-89.

https://doi.org/10.1017/S1062798700000922

Durlak, J. A. (2009). How to select, calculate, and interpret effect sizes. *Journal of Pediatric*

Psychology, *34*(9), 917-928. https://doi.org/10.1093/jpepsy/jsp004

Duval, S., & Tweedie, R. (2000). A nonparametric "trim and fill" method of accounting for publication

bias in meta-analysis. *Journal of the American Statistical Association*, *95*(449), 89-98.

https://doi.org/10.1080/01621459.2000.10473905

Eaves, D. L., Behmer, L., & Vogt, S. (2016). EEG and behavioural correlates of different forms of

motor imagery during action observation in rhythmical actions. *Brain and Cognition*, *106*, 90-

103. https://doi.org/10.1016/j.bandc.2016.04.013

Eaves, D. L., Haythornthwaite, L., & Vogt, S. (2014). Motor imagery during action observation

modulates automatic imitation effects in rhythmical actions. *Frontiers in Human*

Neuroscience, *8*. https://doi.org/10.3389/fnhum.2014.00028

Eaves, D. L., Riach, M., Holmes, P. S., & Wright, D. J. (2016). Motor imagery during action

observation: a brief review of evidence, theory and future research opportunities. *Frontiers*

in Neuroscience, *10*, 514. https://doi.org/10.3389/fnins.2016.00514

Eaves, D. L., Turgeon, M., & Vogt, S. (2012). Automatic imitation in rhythmical actions: Kinematic

fidelity and the effects of compatibility, delay, and visual monitoring. *PLoS One*, *7*(10),

e46728. https://doi.org/10.1371/journal.pone.0046728

Emerson, J. R., Binks, J. A., Scott, M. W., Kenny, R. P., & Eaves, D. L. (2018). Combined action

observation and motor imagery therapy: A novel method for post-stroke motor

rehabilitation. *AIMS Neuroscience*, *5*(4), 236-252.

https://doi.org/10.3934/Neuroscience.2018.4.236

Fadiga, L., Craighero, L., & Olivier, E. (2005). Human motor cortex excitability during the perception

of others' action. *Current Opinion in Neurobiology*, *15*(2), 213-218.

https://doi.org/10.1016/j.conb.2005.03.013

Filimon, F., Rieth, C., Sereno, M., & Cottrell, G. (2015). Observed, executed, and imagined action

representations can be decoded from ventral and dorsal areas. *Cerebral Cortex*, *25*(9), 3144-

3158. https://doi.org/10.1093/cercor/bhu110

Fisher, Z., Tipton, E., Zhipeng, H., & Fisher, M. Z. (2017). Package 'robumeta'.

Frank, C., Wright, D., & Holmes, P. (2020). Mental simulation and neurocognition: Advances for

motor imagery and action observation training in sport. In D. Hackfort & R. J. Schinke (Eds.),

Routledge International Encyclopaedia of Sport and Exercise Psychology (Vol. 2., pp. 411-

428). Routledge. https://doi.org/10.4324/9781315187228

Genaidy, A. M., Lemasters, G. K., Lockey, J., Succop, P., Deddens, J., Sobeih, T., & Dunning, K. (2007).

An epidemiological appraisal instrument – a tool for evaluation of epidemiological

studies. *Ergonomics*, *50*(6), 920-960. https://doi.org/10.1080/00140130701237667

Goldsworthy, M. R., Hordacre, B., & Ridding, M. C. (2016). Minimum number of trials required for

within-and between-session reliability of TMS measures of corticospinal excitability.

Neuroscience, 320, 205-209. https://doi.org/10.1016/j.neuroscience.2016.02.012.

Grosprêtre, S., Ruffino, C., & Lebon, F. (2016). Motor imagery and cortico-spinal excitability: A

review. *European Journal of Sport Science*, *16*(3), 317-324.

https://doi.org/10.1080/17461391.2015.1024756

Guillot, A., & Collet, C. (2008). Construction of the motor imagery integrative model in sport: A

review and theoretical investigation of motor imagery use. *International Review of Sport and*

Exercise Psychology, *1*(1), 31-44. https://doi.org/10.1080/17509840701823139

Hall, C. R., & Martin, K. A. (1997). Measuring movement imagery abilities: a revision of the

movement imagery questionnaire. *Journal of Mental Imagery*.

Hardwick, R., Caspers, S., Eickhoff, S., & Swinnen, S. (2018). Neural correlates of action: Comparing

meta-analyses of imagery, observation, and execution. *Neuroscience & Biobehavioral*

Reviews, *94*, 31-44. https://doi.org/10.1016/j.neubiorev.2018.08.003

Harrer, M., Cuijpers, P., Furukawa, T. & Ebert, D. D. (2019). dmetar: Companion R Package for the

840   Guide 'DoingMeta-Analysis in R'. R package version 0.0.9000. URL:

841   http://dmetar.protectlab.org/.

842   Harrer, M., Cuijpers, P., Furukawa, T. A., & Ebert, D. D. (2021). *Doing Meta-Analysis with R: A Hands-*

843   *On Guide.* Chapmann & Hall; CRC Press. URL:

844   https://bookdown.org/MathiasHarrer/Doing_Meta_Analysis_in_R/

845   Harris, D. J., Allen, K. L., Vine, S. J., & Wilson, M. R. (2021). A systematic review and meta-analysis of

846   the relationship between flow states and performance. *International Review of Sport and*

847   *Exercise Psychology*, 1-29. https://doi.org/10.1080/1750984X.2021.1929402

848   Hedges, L. V. (1981). Distribution theory for Glass's estimator of effect size and related

849   estimators. *Journal of Educational Statistics*, *6*(2), 107-128.

850   https://doi.org/10.3102/10769986006002107

851   Holmes, P. S., & Calmels, C. (2008). A neuroscientific review of imagery and observation use in

852   sport. *Journal of Motor Behavior*, *40*(5), 433-445. https://doi.org/10.3200/jmbr.40.5.433-445

853   Holmes, P. S., & Collins, D. J. (2001). The PETTLEP approach to motor imagery: A functional

854   equivalence model for sport psychologists. *Journal of Applied Sport Psychology, 13*(1), 60-83.

855   https://doi.org/10.1080/10413200109339004

856   Holmes, P. S., & Wright, D. J. (2017). Motor cognition and neuroscience in sport psychology. *Current*

857   *Opinion in Psychology*, *16*, 43-47. https://doi.org/10.1016/j.copsyc.2017.03.009

858   Jeannerod, M. (1994). Motor representations and reality. *Behavioral and Brain Sciences*, *17*(2), 229-

859   245. https://doi.org/10.1017/s0140525x0003435x

860   Jeannerod, M. (2001). Neural simulation of action: A unifying mechanism for motor

861   cognition. *Neuroimage*, *14*(1), S103-S109. https://doi.org/10.1006/nimg.2001.0832

862   Jeannerod, M. (2006). The origin of voluntary action. History of a physiological concept. *Comptes*

863   *Rendus Biologies*, *329*(5-6), 354-362. https://doi.org/10.1016/j.crvi.2006.03.017

864   Kaneko, N., Masugi, Y., Yokoyama, H., & Nakazawa, K. (2018). Difference in phase modulation of

865   corticospinal excitability during the observation of the action of walking, with and without

866   motor imagery. *NeuroReport*, *29*(3), 169-173.

867   https://doi.org/10.1097/wnr.0000000000000941

868   Kawasaki, T., Tozawa, R., & Aramaki, H. (2018). Effectiveness of using an unskilled model in action

869   observation combined with motor imagery training for early motor learning in elderly

870   people: A preliminary study. *Somatosensory & Motor Research*, *35*(3-4), 204-211.

871   https://doi.org/10.1080/08990220.2018.1527760

872   Kim, T., Frank, C., & Schack, T. (2017). A systematic investigation of the effect of action observation

873   training and motor imagery training on the development of mental representation structure

874   and skill performance. *Frontiers in Human Neuroscience*, *11*, 499.

875   https://doi.org/10.3389/fnhum.2017.00499

876   Kraeutner, S. N., McWhinney, S. R., Solomon, J. P., Dithurbide, L., & Boe, S. G. (2018). Experience

877   modulates motor imagery-based brain activity. *European Journal of Neuroscience*, *47*(10),

878   1221-1229. https://doi.org/10.1111/ejn.13900

879   Kraeutner, S. N., Stratas, A., McArthur, J. L., Helmick, C. A., Westwood, D. A., & Boe, S. G. (2020).

880   Neural and behavioral outcomes differ following equivalent bouts of motor imagery or

881   physical practice. *Journal of Cognitive Neuroscience*, *32*(8), 1590-1606.

882   https://doi.org/10.1162/jocn_a_01575

883   Krakauer, J. W., Hadjiosif, A. M., Xu, J., Wong, A. L., & Haith, A. M. (2019). Motor

884   Learning. *Comprehensive Physiology*, *9*(2), 613–663. https://doi.org/10.1002/cphy.c170043

885   Lakens, D. (2013). Calculating and reporting effect sizes to facilitate cumulative science: a practical

886   primer for t-tests and ANOVAs. *Frontiers in Psychology*, *4*, 863.

887   https://doi.org/10.3389/fpsyg.2013.00863

888   Lau, J., Ioannidis, J. P., Terrin, N., Schmid, C. H., & Olkin, I. (2006). The case of the misleading funnel

889   plot. *British Medical Journal*, *333*(7568), 597-600. https://doi.org/10.1136/bmj.333.7568.597

890   MacIntyre, T., Moran, A., Collet, C., & Guillot, A. (2013). An emerging paradigm: A strength-based

891   approach to exploring mental imagery. *Frontiers in Human Neuroscience*, *7,* 104.

892   https://doi.org/10.3389/fnhum.2013.00104

893   Marshall, B., & Wright, D. (2016). Layered stimulus response training versus combined action

894   observation and imagery: Effects on golf putting performance and imagery ability

895   characteristics. *Journal of Imagery Research in Sport and Physical Activity*, *11*(1), 35-46.

896   https://doi.org/10.1515/jirspa-2016-0007

897   Marshall, B., Wright, D., Holmes, P., & Wood, G. (2019). Combining action observation and motor

898   imagery improves eye-hand coordination during novel visuomotor task performance. *Journal

899   of Motor Behavior*, *52*(3), 333-341. https://doi.org/10.1080/00222895.2019.1626337

900   Marshall, B., Wright, D., Holmes, P., Williams, J., & Wood, G. (2020). Combined action observation

901   and motor imagery facilitates visuomotor adaptation in children with developmental

902   coordination disorder. *Research in Developmental Disabilities*, *98*, 103570.

903   https://doi.org/10.1016/j.ridd.2019.103570

904   Marusic, U., Grosprêtre, S., Paravlic, A., Kovač, S., Pišot, R., & Taube, W. (2018). Motor imagery

905   during action observation of locomotor tasks improves rehabilitation outcome in older

906   adults after total hip arthroplasty. *Neural Plasticity, 2018,* 1-9.

907   https://doi.org/10.1155/2018/5651391

908   McNeill, E., Ramsbottom, N., Toth, A. J., & Campbell, M. J. (2020). Kinaesthetic imagery ability

909   moderates the effect of an AO+ MI intervention on golf putt performance: A pilot

910   study. *Psychology of Sport and Exercise*, *46*, 101610.

911   https://doi.org/10.1016/j.psychsport.2019.101610

912   Meers, R., Nuttall, H., & Vogt, S. (2020). Motor imagery alone drives corticospinal excitability during

913   concurrent action observation and motor imagery. *Cortex*, *126*, 322-333.

914   https://doi.org/10.1016/j.cortex.2020.01.012

915   Moreno-Verdú, M., Hamoline, G., Van Caenegem, E. E., Waltzing, B. M., Forest, S., Chembila-

916   Valappil, A., … Hardwick, R. (2022, September 1). Guidelines for Reporting Action Simulation

917   Studies (GRASS): Proposals to improve reporting of research in motor imagery and action

918   observation. *PsyArXiv*. https://doi.org/10.31234/osf.io/9vywr

919   Mouthon, A., Ruffieux, J., Keller, M., & Taube, W. (2016). Age-related differences in corticospinal

920   excitability during observation and motor imagery of balance tasks. *Frontiers in Aging*

921   *Neuroscience*, *8,* 317. https://doi.org/10.3389/fnagi.2016.00317

922   Mouthon, A., Ruffieux, J., Wälchli, M., Keller, M. & Taube, W. (2015). Task-dependent changes of

923   corticospinal excitability during observation and motor imagery of balance tasks.

924   *Neuroscience, 10*(303), 535-543. https://doi.org/10.1016/j.neuroscience.2015.07.031

925   Naish, K., Houston-Price, C., Bremner, A., & Holmes, N. (2014). Effects of action observation on

926   corticospinal excitability: Muscle specificity, direction, and timing of the mirror

927   response. *Neuropsychologia*, *64*, 331-348.

928   https://doi.org/10.1016/j.neuropsychologia.2014.09.034

929   Nedelko, V., Hassa, T., Hamzei, F., Schoenfeld, M., & Dettmers, C. (2012). Action imagery combined

930   with action observation activates more corticomotor regions than action observation

931   alone. *Journal of Neurologic Physical Therapy*, *36*(4), 182-188.

932   https://doi.org/10.1097/npt.0b013e318272cad1

933   Neuman, B., & Gray, R. (2013). A direct comparison of the effects of imagery and action observation

934   on hitting performance. *Movement & Sport Sciences - Science & Motricité*, (79), 11-21.

935   https://doi.org/10.1051/sm/2012034

936   Pellegrini, M., Zoghi, M., & Jaberzadeh, S. (2018). The effect of transcranial magnetic stimulation test

937   intensity on the amplitude, variability and reliability of motor evoked potentials. *Brain*

938   *Research*, *1700*, 190-198. https://doi.org/10.1016/j.brainres.2018.09.002

Roberts, R., Callow, N., Hardy, L., Markland, D., & Bringer, J. (2008). Movement imagery ability:

Development and assessment of a revised version of the vividness of movement imagery

questionnaire. *Journal of Sport & Exercise Psychology, 30*(2), 200-221.

https://doi.org/10.1123/jsep.30.2.200

Romano Smith, S., Wood, G., Coyles, G., Roberts, J., & Wakefield, C. (2019). The effect of action

observation and motor imagery combinations on upper limb kinematics and EMG during

dart-throwing. *Scandinavian Journal of Medicine & Science in Sports*, *29*(12), 1917-1929.

https://doi.org/10.1111/sms.13534

Romano-Smith, S., Wood, G., Wright, D., & Wakefield, C. (2018). Simultaneous and alternate action

observation and motor imagery combinations improve aiming performance. *Psychology of

Sport and Exercise*, *38*, 100-106. https://doi.org/10.1016/j.psychsport.2018.06.003

Rothwell, J. C. (1997). Techniques and mechanisms of action of transcranial stimulation of the

human motor cortex. *Journal of Neuroscience Methods*, *74*(2), 113-122.

https://doi.org/10.1016/S0165-0270(97)02242-5

Ruffino, C., Papaxanthis, C., & Lebon, F. (2017). Neural plasticity during motor learning with motor

imagery practice: Review and perspectives. *Neuroscience*, *341*, 61-78.

https://doi.org/10.1016/j.neuroscience.2016.11.023

Scott, M. W., Wood, G., Holmes, P. S., Williams, J., Marshall, B., & Wright, D. J. (2021). Combined

action observation and motor imagery: An intervention to combat the neural and

behavioural deficits associated with developmental coordination disorder. *Neuroscience &

Biobehavioral Reviews*. *127*, 638-646. https://doi.org/10.1016/j.neubiorev.2021.05.015

Scott, M., Taylor, S., Chesterton, P., Vogt, S., & Eaves, D. (2018). Motor imagery during action

observation increases eccentric hamstring force: an acute non-physical

intervention. *Disability and Rehabilitation*, *40*(12), 1443-

1451. https://doi.org/10.1080/09638288.2017.1300333

Scott, M.W., Emerson, J.R., Dixon, J., Tayler, M. A. & Eaves, D. L. (2019). Motor imagery during action

observation enhances automatic imitation in children with and without developmental

coordination disorder. *Journal of Experimental Child Psychology, 183*, 242-260.

https://doi.org/10.1016/j.jecp.2019.03.001

Shimada, K., Onishi, T., Ogawa, Y., Yamauchi, J., & Kawada, S. (2019). Effects of motor imagery

combined with action observation training on the lateral specificity of muscle strength in

healthy subjects. *Biomedical Research, 40*(3), 107-113.

https://doi.org/10.2220/biomedres.40.107

Simonsmeier, B., Androniea, M., Buecker, S., & Frank, C. (2021). The effects of imagery interventions

in sports: A meta-analysis. *International Review of Sport and Exercise Psychology*, 1-22.

https://doi.org/10.1080/1750984x.2020.1780627

Spittle, M. (2021). *Motor learning and skill acquisition: Applications for physical education and sport*.

Bloomsbury Publishing.

Ste-Marie, D. M., Lelievre, N., & St. Germain, L. (2020). Revisiting the Applied Model for the Use of

Observation: A Review of Articles Spanning 2011–2018. *Research quarterly for exercise and

sport*, *91*(4), 594-617.

Ste-Marie, D., Law, B., Rymal, A., Jenny, O., Hall, C., & McCullagh, P. (2012). Observation

interventions for motor skill learning and performance: an applied model for the use of

observation. *International Review of Sport and Exercise Psychology*, *5*(2), 145-176.

https://doi.org/10.1080/1750984x.2012.665076

Stevens, J. A., & Stoykov, M. E. (2003). Using motor imagery in the rehabilitation of

hemiparesis. *Archives of Physical Medicine and Rehabilitation*, *84*(7), 1090–1092.

https://doi.org/10.1016/s0003-9993(03)00042-x

Tanner-Smith, E. E., & Tipton, E. (2014). Robust variance estimation with dependent effect sizes:

practical considerations including a software tutorial in Stata and spss. *Research Synthesis

Methods, 5*(1), 13–30. https://doi.org/10.1002/jrsm.1091

Tanner-Smith, E.E., Tipton, E. & Polanin, J.R. (2016). Handling complex meta-analytic data structures

using Robust Variance Estimates: a tutorial in R. *Journal of Developmental Life Course*

Criminology 2, 85–112. https://doi.org/10.1007/s40865-016-0026-5

Taube, W., Lorch, M., Zeiter, S. & Keller, M. (2014). Non-physical practice improves task performance

in an unstable, perturbed environment: Motor imagery and observational balance

training. *Frontiers in Human Neuroscience, 8,* 972.

https://doi.org/10.3389/fnhum.2014.00972

Taube, W., Mouthon, M., Leukel, C., Hoogewoud, H., Annoni, J., & Keller, M. (2015). Brain activity

during observation and motor imagery of different balance tasks: An fMRI study. *Cortex*, *64*,

102-114. https://doi.org/10.1016/j.cortex.2014.09.022

Toth, A., McNeill, E., Hayes, K., Moran, A., & Campbell, M. (2020). Does mental practice still enhance

performance? A 24 Year follow-up and meta-analytic replication and extension. *Psychology*

of Sport and Exercise, *48*, 101672. https://doi.org/10.1016/j.psychsport.2020.101672

Viechtbauer, W. (2010). Conducting meta-analyses in R with the metafor package. *Journal of*

Statistical Software, 36(3), 1-48. https://doi.org/10.18637/jss.v036.i03

Villiger, M., Estévez, N., Hepp-Reymond, M., Kiper, D., Kollias, S., Eng, K., & Hotz-Boendermaker, S.

(2013). Enhanced activation of motor execution networks using action observation

combined with imagination of lower limb movements. *PLoS One*, *8*(8), e72403.

https://doi.org/10.1371/journal.pone.0072403

Vogt, S., Rienzo, F., Collet, C., Collins, A., & Guillot, A. (2013). Multiple roles of motor imagery during

action observation. *Frontiers in Human Neuroscience*, *7,* 807.

https://doi.org/10.3389/fnhum.2013.00807

Williams, S. E., Cumming, J., Ntoumanis, N., Nordin-Bates, S. M., Ramsey, R., & Hall, C. (2012).

Further validation and development of the movement imagery questionnaire. *Journal of*

Sport & Exercise Psychology, *34*(5), 621–646. https://doi.org/10.1123/jsep.34.5.621

Wright D. J., Williams J., Holmes P. S. (2014) Combined action observation and imagery facilitates

corticospinal excitability. *Frontiers in Human Neuroscience,* 27(8), 951.

https://doi.org/10.3389/fnhum.2014.00951

Wright, D. J., Frank, C., & Bruton, A. M. (2021). Recommendations for combining action observation

and motor imagery interventions in sport. *Journal of Sport Psychology in Action,* 1-13.

https://doi.org/10.1080/21520704.2021.1971810

Wright, D.J., McCormick, S.A., Williams, J. & Holmes, P.S. (2016). Viewing instructions accompanying

action observation modulate corticospinal excitability. *Frontiers in Human Neuroscience*, 1,

10(17). https://doi.org/10.3389/fnhum.2016.00017

Wright, D., Wood, G., Eaves, D., Bruton, A., Frank, C., & Franklin, Z. (2018a). Corticospinal excitability

is facilitated by combined action observation and motor imagery of a basketball free

throw. *Psychology of Sport and Exercise, 39*, 114-121.

https://doi.org/10.1016/j.psychsport.2018.08.006

Wright, D. J., Wood, G., Franklin, Z. C., Marshall, B., Riach, M., & Holmes, P. S. (2018b). Directing

visual attention during action observation modulates corticospinal excitability. *Plos One,*

13(1), e0190165. https://doi.org/10.1371/journal.pone.0190165.

Yoxon, E., & Welsh, T. (2020). Motor system activation during motor imagery is positively related to

the magnitude of cortical plastic changes following motor imagery training. *Behavioural*

Brain Research, *390*, 112685. https://doi.org/10.1016/j.bbr.2020.112685

Zhang B, Kan L, Dong A, Zhang J, Bai Z, Xie Y, et al. (2019) The effects of action observation training

on improving upper limb motor functions in people with stroke: A systematic review and

meta-analysis. *PLoS One*, *14*(8), e0221166. https://doi.org/10.1371/journal.pone.0221166

1037

1038

1039

1040

1041

1042

1043

1044

7. Figure Captions

Figure 1*. *Flow chart of the literature search and selection procedures adopted for the MEP meta-analysis and Movement meta-analysis

Figure 2*. *Bar chart displaying quality assessment items and scores for all studies included in the (A) MEP meta-analysis and (B) Movement meta-analysis. The blue bar indicates the number of studies that satisfied, and the red bar indicates the number of studies that did not satisfy, each of the respective quality assessment criteria.

Figure 3. Forest plot of effect sizes (*d*) for all studies included in the AOMI vs control condition comparison for the MEP meta-analysis. The combined estimate (dashed vertical line) and 95% confidence interval (hollow diamond) indicates AOMI has a medium positive effect on MEP amplitudes compared to control conditions. The size of each black square indicates the weight of the study effect size in the combined analysis. Multiple effect sizes are reported for a study if it recorded MEP amplitude data from more than one target muscle.

Figure 4*. *Forest plot of effect sizes (*d*) for all studies included in the AOMI vs AO condition comparison for the MEP meta-analysis. The combined estimate (dashed vertical line) and 95% confidence interval (hollow diamond) indicates AOMI has a small to medium positive effect on MEP amplitudes compared to AO conditions. The size of each black square indicates the weight of the study effect size in the combined analysis. Multiple effect sizes are reported for a study if it recorded MEP amplitude data from more than one target muscle.

Figure 5. Forest plot of effect sizes (*d*) for all studies included in the AOMI vs MI condition comparison for the MEP meta-analysis. The combined estimate (dashed vertical line) and 95% confidence interval (hollow diamond) indicates AOMI has no significant effect on MEP amplitudes compared to MI conditions. The size of each black square indicates the weight of the study effect size in the combined analysis. Multiple effect sizes are reported for a study if it recorded MEP amplitude data from more than one target muscle.

Figure 6*. *Forest plot of effect sizes (*d*) for all studies included in the AOMI vs control condition comparison for the Movement meta-analysis. The combined estimate (dashed vertical line) and 95% confidence interval (hollow diamond) indicates AOMI has a medium to large positive effect on movement outcomes compared to control conditions. The size of each black square indicates the weight of the study effect size in the combined analysis. Multiple effect sizes are reported for a study if it recorded more than one movement outcome variable.

Figure 7. Forest plot of effect sizes (*d*) for all studies included in the AOMI vs AO condition comparison for the Movement meta-analysis. The combined estimate (dashed vertical line) and 95% confidence interval (hollow diamond) indicates AOMI has a small to medium positive effect on movement outcomes compared to AO conditions. The size of each black square indicates the weight of the study effect size in the combined analysis. Multiple effect sizes are reported for a study if it recorded more than one movement outcome variable.

Figure 8*. *Forest plot of effect sizes (*d*) for all studies included in the AOMI vs MI condition comparison for the Movement meta-analysis. The combined estimate (dashed vertical line) and 95% confidence interval (hollow diamond) indicates AOMI has no significant effect on movement outcomes compared to MI conditions. The size of each black square indicates the weight of the

study effect size in the combined analysis. Multiple effect sizes are reported for a study if it recorded
more than one movement outcome variable.

Figure 9. Funnel plot of effect sizes (Cohen's *d*) versus standard error before (left) and after (right)
performing Duval and Tweedie's Trim-and-fill analysis for the AOMI vs AO comparison in the MEP
meta-analysis. Black circles represent existing effects included in the MEP meta-analysis and white
circles represent potential unpublished effects. The contour-enhanced funnel plots display the
significance of the effects for the AOMI vs AO comparison in this meta-analysis. Individual effect
sizes falling inside the white ($p < .05$) and orange ($p < .01$) funnel boundaries represent significant
effects for the AOMI vs AO comparison in the MEP meta-analysis.

1095

1096

1097 **8. Tables**

1098 **Table 1. Meta-regression analyses for the MEP amplitude data synthesized in the MEP meta-**
1099 **analysis and movement outcome data synthesized in the Movement meta-analysis**

Moderator	N	K	Beta	P	Sig.	τ^2	I^2
MEP Meta-Analysis							
Kinesthetic Imagery Ability	6	19	-0.01	**.90**	**No**	0.00	0.00
Intensity of TMS Pulses	15	53	0.06	**.13**	**No**	0.02	5.71
Number of TMS Trials	15	53	0.00	**.85**	**No**	0.02	11.8
Age	13	47	0.00	**.61**	**No**	0.00	0.00
Movement Meta-Analysis							
Intervention Volume	14	52	0.00	**.72**	**No**	0.40	71.03
Kinesthetic Imagery Ability	5	21	-0.01	**.86**	**No**	2.40	88.53
Age	16	57	-0.01	**.19**	**No**	0.38	68.73

1100 *Note.* N = number of studies, K = number of effect sizes, Beta = regression coefficient, P =
1101 significance value, Sig. = significance threshold met or not, τ^2 = measure of variation around average,
1102 I^2 = measure of proportion of observed variance (%).

1103 **Table 2**. **Sub-group analyses for the MEP amplitude data synthesized in the MEP meta-analysis**

Sub-group	N	K	SMD	95% CI		P	Sig.	τ^2	I^2	Sig. Mod.
Action Observation Perspective										
First-person	9	23	0.50	0.33	0.67	**<.001**	**Yes**	0.00	0.00	**ref**
Third-person	7	31	0.48	0.12	0.84	**.02**	**Yes**	0.15	45.41	*ns*
Directed Attentional Focus										
Yes	1	2	0.38	NA	NA	**NA**	**NA**	0.00	0.00	*ref*
No	15	52	0.49	0.34	0.64	**<.001**	**Yes**	0.02	7.15	ns
Skill Classification 1										
Fine	10	24	0.48	0.33	0.64	**<.001**	**Yes**	0.00	0.00	ns
Gross	6	30	0.52	0.07	0.97	**.03**	**Yes**	0.19	52.98	*ref*
Skill Classification 2										
Discrete	12	32	0.52	0.35	0.70	**<.001**	**Yes**	0.00	0.00	*ns*
Continuous	4	22	0.34	0.16	0.52	**<.01**	**Yes**	0.09	33.19	ref
Timing of TMS Delivery										
Targeted	12	35	0.47	0.30	0.64	**<.001**	**Yes**	0.00	0.00	**ref**
Random	4	19	0.53	0.28	0.77	**<.001**	**Yes**	0.08	28.61	*ns*

1104 *Note.* N = number of studies, K = number of effect sizes, SMD = standardized mean difference, 95% CI = lower and upper confidence intervals, P =
1105 significance value, Sig. = significance threshold met or not, τ^2 = measure of variation around average, I^2 = measure of proportion of observed variance (%),
1106 Sig. Mod. = comparison between sub-group categories. NA = insufficient number of data points for the analysis, ref = reference category, *ns* = no significant
1107 difference compared to reference category

1108

1109 **Table 3**. Sub-group analyses for the movement outcome data synthesized in the Movement meta-analysis

Sub-group	N	K	SMD	95% CI		P	Sig.	τ²	I²	Sig. Mod.
Action Observation Perspective										
First-person	8	29	0.59	0.08	1.09	.03	Yes	0.61	73.68	*ref*
Third-person	8	28	0.39	-0.07	0.84	.08	No	0.29	67.28	ns
Context										
Sport	4	17	0.68	-0.65	2.02	.20	No	1.38	86.11	ns
Rehabilitation	7	28	0.43	-0.13	1.00	.11	No	0.42	70.93	*ns*
Other	5	12	0.43	-0.03	0.88	.06	No	0.07	31.62	ref
Directed Attentional Focus										
Yes	4	15	0.38	-0.91	1.68	.41	No	0.59	82.46	*ref*
No	12	42	0.52	0.22	0.82	<.01	Yes	0.33	63.07	ns
PETTLEP										
Yes	5	19	0.69	-.039	1.76	.15	No	0.83	80.52	*ref*
No	11	38	0.39	0.11	0.67	.01	Yes	0.28	63.71	ns
Physical Practice										
Yes	2	7	0.75	-6.12	7.63	.40	No	3.96	91.29	*ref*
No	14	50	0.45	0.13	0.77	.01	Yes	0.28	64.11	ns
Population										
Neurotypical	13	46	0.47	0.19	0.75	<.01	Yes	0.29	62.26	*ref*
Neurodivergent	3	11	0.55	-2.05	3.14	.46	No	1.27	87.65	ns
Skill Classification 1										
Fine	12	41	0.45	0.07	0.83	.03	Yes	0.32	67.52	ns
Gross	4	16	0.59	-0.12	1.31	.08	No	0.98	79.46	*ref*
Skill Classification 2										
Continuous	2	10	0.47	-1.66	2.60	.22	No	0.28	56.32	ref
Discrete	14	47	0.48	0.14	0.83	.01	Yes	0.43	72.18	*ns*

1110   *Note.* N = number of studies, K = number of effect sizes, SMD = standardized mean difference, 95% CI = lower and upper confidence intervals, P =

1111   significance value, Sig. = significance threshold met or not, τ^2 = measure of variation around average, I^2 = measure of proportion of observed variance (%),

1112   Sig. Mod. = comparison between sub-group categories. NA = insufficient number of data points for the analysis, ref = reference category, *ns* = no significant

1113   difference compared to reference category

1114

(A)

Criterion	
Clear aims and hypotheses	
Theoretical framework	
Appropriate design	
Clear outcome measures	
Participant characteristics	
Representative sample	
Sample size determination	
Attention to ethics	
Task clearly described	
Appropriate statistics	
Parameter estimates reported	
Exact p-values reported	
Effect sizes always reported	
Conclusions supported by data	
Findings clearly described	
Generalisable to other populations	
Compared to previous results	
Valid outcome measures	

Count: 0, 4, 8, 12, 16

(B)

Criterion	
Clear aims and hypotheses	
Theoretical framework	
Appropriate design	
Clear outcome measures	
Participant characteristics	
Representative sample	
Sample size determination	
Attention to ethics	
Task clearly described	
Appropriate statistics	
Parameter estimates reported	
Exact p-values reported	
Effect sizes always reported	
Conclusions supported by data	
Findings clearly described	
Generalisable to other populations	
Compared to previous results	
Valid outcome measures	

Count: 0, 4, 8, 12, 16

Study	Muscle	Effect Size
Bruton et al., 2020	First Dorsal Interosseous	0.246
Castro et al., 2021	First Dorsal Interosseous	0.244
Kaneko et al., 2018	Tibialis Anterior	0.237
	Soleus	1.195
Meers et al., 2020	First Dorsal Interosseous	0.309
Mouthon et al., 2015	Soleus	0.207
Mouthon et al., 2016	Soleus	0.432
Ohno et al., 2011	First Dorsal Interosseous	0.773
	Thenar	0.543
Sakamoto et al., 2009	Biceps Brachii	2.100
Tibialis Anteriornaka et al., 2015	First Dorsal Interosseous	0.220
Tsukazaki et al., 2018	First Dorsal Interosseous	1.209
	Abductor Digiti Minimi	0.858
	Abductor Pollicis Brevis	1.023
	First Dorsal Interosseous	0.837
	Abductor Pollicis Brevis	0.352
	Abductor Digiti Minimi	0.914
Wright et al., 2014	First Dorsal Interosseous	0.236
Wright et al., 2016	Opponens Pollicis	0.315
	First Dorsal Interosseous	0.474
	Abductor Digiti Minimi	0.375
Wright et al., 2018	Flexor Carpi Ulnaris	0.755
	Extensor Carpi Ulnaris	0.262

Effect Size

Bek et al., 2016
Duration (s) — 0.028
Peak velocity (mm/s) — 0.490
Time to peak velocity (time/duration) — 0.000
Vertical amplitude (mm) — 0.609
Horizontal amplitude (mm) — -0.098

Bek et al., 2019
Direct vertical amplitude (mm) — 0.070
Elevated vertical amplitude (mm) — -0.606
Direct vertical amplitude (mm) — -0.126
Elevated vertical amplitude (mm) — -0.336

De Lucia et al., 2016
Accuracy — 0.525

Di Rienzo et al., 2019
Total force (%) — 0.016

Frenkel-Toledo et al., 2020
Sequence (ms) — 0.861
Delta (ms) — -0.845
Fail (%) — 0.228

Marshall et al., 2019
Completion time (sec) — 0.824

Romano-Smith et al., 2018
Performance — 1.028

Romano-Smith et al., 2019
Performance score — 0.586
Movement time — 0.000
Peak angular velocity — -0.784

Scott et al., 2019
Mean cycle time ratio td (%) — 2.488
Mean cycle time ratio dcd (%) — 0.000

Effect Size