The effects of combined action observation and motor imagery on corticospinal excitability and

movement outcomes: A two-pronged meta-analysis

Samantha Chye [a*], Ashika Chembila Valappil [a*], David J. Wright [b], Cornelia Frank [c], David A. Shearer [d],
Christopher J. Tyler [a], Ceri E. Diss [a], Omar S. Mian [a], Neale A. Tillin [a] & Adam M. Bruton [a**]

[a] Centre for Integrated Research in Life and Health Sciences, School of Life and Health Sciences,
University of Roehampton, London, UK, SW15 4JD. E-mail: chyes1@roehampton.ac.uk,
chembila@roehampton.ac.uk, chris.tyler@roehampton.ac.uk, c.diss@roehampton.ac.uk,
omar.mian@roehampton.ac.uk, neale.tillin@roehampton.ac.uk, adam.bruton@roehampton.ac.uk.

[b] Research Centre for Health, Psychology and Communities, Department of Psychology, Manchester
Metropolitan University, Manchester, UK, M15 6GX. E-mail: d.j.wright@mmu.ac.uk.

[c] Sports and Movement Research Group, Department of Sports and Movement Science, School of
Educational and Cultural Studies, Osnabrück University, Osnabrück, Germany. Email:
cornelia.frank@uni-osnabrueck.de.

[d] Faculty of Life Science and Education, University of South Wales, UK, CF37 1DL. Email:
david.shearer@southwales.ac.uk.

Author Note: *Samantha Chye and Ashika Chembila Valappil are shared first authors for this
manuscript. **Adam M. Bruton is the corresponding author for this manuscript:
adam.bruton@roehampton.ac.uk, Room 1051 Parkstead House, Whitelands College, School of Life
and Health Sciences, University of Roehampton.

<u>Highlights</u>

• Motor imagery (MI) and action observation (AO) can be combined (AOMI)

• This meta-analysis collates neurophysiological and behavioral evidence for AOMI

• AOMI had positive neurophysiological effects compared to AO and control but not MI

• AOMI had positive behavioral effects compared to AO and control but not MI

• The positive effects of AOMI were maintained across all moderators

64   **Abstract**

Motor simulation interventions involving motor imagery (MI) and action observation (AO) have

received considerable interest in the behavioral sciences. A growing body of research has focused on

using AO and MI simultaneously, termed 'combined action observation and motor imagery' (AOMI).

The current meta-analysis adopted a two-pronged approach by collating and synthesizing existing

motor evoked potential (MEP) amplitude data from transcranial magnetic stimulation studies and

movement outcome data from behavioral studies to quantify changes in corticospinal excitability

and motor skill performance for AOMI compared to AO, MI and control conditions. AOMI had a

positive effect compared to control and AO but not MI conditions for both MEP amplitudes and

movement outcomes. No methodological factors moderated the effects of AOMI, indicating a robust

effect of AOMI across the two outcome variables. The results of this meta-analysis are discussed in

relation to existing literature on motor simulation and skill acquisition, before providing viable

directions for future research on this topic.

Keywords: AOMI, motor evoked potentials, dual action simulation, motor execution, motor imagery

during action observation, transcranial magnetic stimulation

1. Introduction

According to motor simulation theory (Jeannerod, 1994, 2001, 2006), it is possible to cognitively

rehearse an action both overtly and covertly through action observation (AO) and motor imagery

(MI), with this cognitive simulation activating motor regions of the brain in a similar manner to

physical execution of the action. AO is a bottom-up process that involves the deliberate and

structured observation of human movement (Neuman & Gray, 2013), whereas MI is a top-down

process that involves the internal generation of the visual and kinesthetic elements of movement

(Macintyre et al., 2013). Literature has reported positive behavioral outcomes for AO- and MI-based

practice in sport (e.g., Guillot & Collet, 2008; Ste-Marie et al., 2012) and neurorehabilitation settings

(e.g., Buccino, 2014; De Vries & Mulder, 2007). There is preliminary evidence indicating that plastic

changes in the primary motor system may underpin these behavioral improvements (e.g., Yoxon and

Welsh, 2020). A recent large-scale meta-analysis of functional magnetic resonance imaging (fMRI)

data reported a shared network including premotor, rostral parietal, and somatosensory areas of the

brain active during AO, MI, and movement execution (Hardwick et al., 2018). Notably, there were

differences in the neural regions activated during AO and MI, and the brain activity for these

overlapped differently with brain activity for physically performed actions. It is possible that using

AO and MI simultaneously, typically labelled 'combined action observation and motor imagery'

(AOMI), may hold greater neural overlap with physical execution.

AOMI refers to a person watching a video or live demonstration of a movement while

simultaneously generating, maintaining, and transforming a time-synchronized kinesthetic

representation of the same action (Eaves, Riach et al., 2016; Vogt et al., 2013). AOMI has received

growing research interest over the last decade and two hypotheses have been proposed to explain

why AOMI may be more effective as a motor skill intervention than independent AO or MI

interventions. First, Eaves and colleagues (2012, 2014, 2016) suggested the dual action simulation

hypothesis (DASH), which proposes that a person will generate separate motor representations for

the observed and imagined actions and maintain these as two parallel sensorimotor streams when

they engage in AOMI. If a person is simultaneously observing and imagining the same action, these

two motor representations are likely to merge as one sensorimotor stream, producing more

widespread activity in the premotor cortex compared to AO or MI alone. This is likely due to AOMI

increasing activity in shared brain areas for AO and MI, as well as increasing activity in areas solely

recruited during AO (e.g., inferior frontal gyrus, ventral premotor area) and MI (e.g., angular gyrus,

dorsal premotor area) of an action (Filimon et al., 2015; Hardwick et al., 2018). Second, Meers et al.

(2020) introduced the visual guidance hypothesis (VGH) as an alternative account of how AOMI may

influence action. They suggest that MI is prioritized during AOMI, and that the AO component might

merely serve as an external visual guide that facilitates more vivid MI generation. In contrast to

DASH, this would mean that AO does not activate a separate motor representation during AOMI, but

rather strengthens the motor representation resulting from MI. Irrespective of the stance taken,

both the DASH and VGH suggest that AOMI has the capacity to influence motor skill execution above

and beyond AO or MI in isolation through increased activity in motor regions of the brain.

118   Current neuroscientific evidence using a range of modalities supports this notion, as cortico-

motor activity is increased during AOMI of an action compared to independent AO or MI of that

same action (Eaves, Riach et al., 2016). Studies using fMRI report distinct neural signatures for AO,

MI, and AOMI whereby the blood-oxygen-level-dependent (BOLD) signal is increased and more

widespread in the brain regions involved in movement execution when an individual engages in

AOMI (e.g., Nedelko et al., 2012; Taube et al., 2015; Villiger et al., 2013). For example, Taube et al.

(2015) found greater activation in the supplementary motor area, basal ganglia and cerebellum

during AOMI compared to AO, and greater bilateral activity in the cerebellum and greater activation

in the precuneus compared to MI. Studies using electroencephalography (EEG) report that AOMI

leads to significantly larger event-related desynchronization in the mu/alpha and beta frequency

bands, indicative of increased activity over the primary sensorimotor areas of the brain compared to

both AO and MI alone (Berends et al., 2013; Eaves, Behmer et al., 2016). These fMRI and EEG

findings have important implications for applied practice, where the use of AOMI may prove

beneficial in reinforcing motor (re)learning. The increased neural activity during AOMI has the

potential to support repetitive Hebbian modulation of intracortical and subcortical excitatory

mechanisms through synaptic plasticity, in a similar manner to physical practice, and thus may be an

effective method for behavior change (Holmes & Calmels, 2008).

135   From a neurophysiological perspective, this meta-analysis will focus on studies adopting

single-pulse transcranial magnetic stimulation (TMS) during AOMI, as this is the most prevalent

neuroscientific modality adopted in the literature to date (study count: n = 19 TMS, n = 14 fMRI, n =

10 EEG). When applied to a muscle representation of the primary motor cortex, TMS produces a

twitch response in the corresponding muscles called motor evoked potentials (MEPs). MEPs are

measured using surface electromyography (EMG) and provide a marker of corticospinal excitability

during the time of stimulation (Rothwell, 1997). This approach has been used extensively when

studying the neural mechanisms for motor imagery (see e.g., Grosprêtre et al., 2016) and action

observation (see e.g., Naish et al., 2014) as it permits a non-invasive assessment of muscle-specific

M1 activity and excitability of the whole cortico-spinal pathway that is specific to the final motor

command for the action being simulated. Studies using TMS during AOMI have explored changes

in MEP amplitudes across a range of movements, including finger movements (Bruton et al., 2020),

basketball free throws (Wright, Wood et al., 2018a), walking (Kaneko et al., 2018), and balance

movements (Mouthon et al., 2016). Current literature predominantly shows increased corticospinal

excitability during AOMI compared to baseline conditions (e.g., Bruton et al., 2020; Wright et al.,

2014, 2016). However, studies comparing AOMI against AO or MI have reported increased (e.g.,

Mouthon et al., 2016; Wright et al., 2016), as well as no differences (e.g., Castro et al., 2021;

Mouthon et al., 2015) in corticospinal excitability during AOMI. Given the prevalence of TMS studies

exploring the neurophysiological mechanisms underpinning AOMI engagement, it is now possible to

synthesize the available MEP amplitude data to quantify the effects of AOMI on corticospinal

excitability compared to AO, MI, and control conditions.

AOMI investigations have explored a range of movement outcomes and types, such as

movement time for ball rotations (Kawasaki et al., 2018), upper-limb kinematics for dart throwing

(Romano-Smith et al., 2019), force production in Nordic hamstring curls (Scott et al., 2018), and

radial error from the hole in golf-putting (Marshall & Wright, 2016). This research has been

conducted in sport (e.g., Romano-Smith et al., 2019) and rehabilitation (e.g., Scott et al., 2018)

contexts, with neurotypical (e.g., Di Rienzo et al., 2019) and neurodivergent (e.g., Marshall et al.,

2020) populations. The existing literature has almost exclusively demonstrated that movement

outcomes are improved for different motor skills after AOMI interventions when compared to

control conditions (e.g., Marshall et al., 2019; Romano-Smith et al., 2018; Shimada et al., 2019).

However, comparisons with AO and MI interventions are equivocal, with some studies showing

greater improvements in movement outcomes for AOMI compared to AO (e.g., Bek et al., 2019) and

MI interventions (Scott et al., 2018), and other studies showing no such effects (e.g., Romano-Smith

et al., 2019) or greater improvements for MI-related interventions (e.g., Marshall & Wright, 2016).

Therefore, it is unclear if AOMI should be recommended as the optimal simulation approach when

attempting to improve motor skill performance, highlighting the need to synthesize available

movement outcome data for AOMI interventions.

Early reviews on AOMI (Eaves, Riach et al., 2016; Vogt et al., 2013) summarized the

behavioral neuroscience literature contrasting AO and MI and drew on early AOMI research to

support its use as a motor skill intervention. Since then, research has explored changes in MEP

amplitudes and movement outcomes associated with engagement in AOMI. This has led to

population-specific reviews outlining how AOMI can be used to address sensorimotor deficits for

individuals with Parkinson's disease (Caligiore et al., 2017), developmental coordination disorder

(Scott et al., 2021), or during post-stroke rehabilitation (Emerson et al., 2018). Systematic reviews

and meta-analyses have synthesized the respective effects of MI and AO on corticospinal excitability

(e.g., Grosprêtre et al., 2016; Naish et al., 2014) and motor skill performance (e.g., Ashford et al.,

2006; Simonsmeier et al., 2021, Toth et al., 2020) when used in isolation, however, to our

knowledge, no such meta-analyses exist for AOMI interventions.

The purpose of the current meta-analysis was two-pronged; first, to methodically collate and

synthesize the available MEP amplitude data as an indicator of corticospinal excitability during AOMI

engagement, and second, to methodically collate and synthesize the available movement outcome

data to assess changes in motor skill performance that result from AOMI interventions. Our primary

aim was to establish the effectiveness of AOMI by comparing its effects on MEP amplitudes and

movement outcomes against those for AO, MI, and control conditions. Based on previous literature

(see Eaves, Riach et al., 2016), it was hypothesized that AOMI would have a small positive effect

when compared to independent AO or MI, and a medium positive effect when compared to control

conditions for both outcome variables. Our secondary aim was to explore several methodological

parameters hypothesized to have a moderating effect on the impact of AOMI interventions on MEP

amplitudes or movement outcomes. Considering the influence of these methodological aspects

directly addresses questions raised in early reviews on AOMI (Eaves, Riach et al., 2016; Vogt et al.,

2013) and may permit optimal delivery of AOMI interventions in the future.

196   **2. Method**

2.1. Study Protocol

The study procedures were performed according to the methodological guidelines

highlighted by the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA)

statement (Page et al., 2021). The meta-analysis justification and proposed procedures were

documented in a pre-registration protocol document (https://osf.io/9yebv) that is stored alongside

additional supplementary files (https://osf.io/9yebv) on the Open Science Framework. A PRISMA

flow chart (Figure 1) outlines the study search and selection process for the current meta-analysis.

Figure 1. Flow chart of the meta-analysis literature search and selection procedures

2.1.1. Literature Search and Selection

The literature search was performed using three online databases: PubMed, Web of Science,

and PsycINFO. The original search was run in August 2020 and the final search was run in June 2021.

Consistent search terms were decided upon and adapted for each database based on the search

string requirements (see supplementary files for full search string information). Web of Science was

used as an initial search point, followed by PubMed and PsycINFO to cover the biomedical and

psychological literature, respectively. The search was limited to studies with human populations that

had been published in a peer-reviewed journal in English language, without any limitations to

publication years. The literature search with the respective keyword combinations and restrictions

provided 396 hits in total, after 244 duplicates were removed. An exploratory search of review

articles and prior knowledge of research led to 8 additional papers being added, which took the final

number of papers being screened to 404.

2.1.2. Inclusion of Studies

The decision to include a study was based on five criteria. (1) The study had to either test the

effects of AOMI on corticospinal excitability by recording MEP amplitudes during single pulse TMS or

test the effects of AOMI on movement outcomes by recording parameters related to motor task

performance. Participants were required to engage in an AOMI task as their intervention or

experimental method, and AOMI had to be delivered in a synchronous manner (i.e., engagement in

both forms of simulation at the same time). Studies with independent AO or MI conditions were

included, but only if there was a comparison with an AOMI condition. Studies that included AOMI as

an adjunct to physical practice were also included, but the inclusion of physical practice was not

mandatory, and these studies needed to incorporate physical practice in the control condition to

permit assessment of AOMI effects. (2) The study had to adopt a controlled experimental design

where the unconfounded effects of the AOMI intervention could be compared within- or between-

groups. As such, studies using a correlational or survey design, qualitative research, or case

study/single-subject designs were excluded. Studies that did not report a control group or baseline

condition were also excluded, unless they contained data from AO or MI conditions that could be

compared to the AOMI condition. (3) The study had to include an MEP amplitude or movement

outcome measure with a specific experimental group, or across all study groups. Studies that did not

have distinguishable MEP amplitude or movement outcome measures or used neurophysiological

modalities other than/combined with TMS (e.g., fMRI, EEG, fNIRS) were excluded unless they also

had independent MEP amplitude or movement outcome measures. (4) The study had to report

original statistics (i.e., not a re-analysis of already published findings or a review) of the intervention

on MEP amplitude or movement outcome measures. (5) The study had to be published in English.

2.2. Screening Process

A three-stage screening process was adopted in this study (i.e., title, abstract, and full-text

screening). At all screening stages, the two co-first authors independently screened all articles.

Conflicts were resolved by discussion between the co-first authors in the title screen phase, with

additional input from the final author in the abstract and full text screen stages. First, all 404 titles

were screened to evaluate relevance to the current meta-analysis, with 119 titles classified as

eligible for further inspection of abstracts (intercoder reliability = 94.55%). Second, the 119 abstracts

were assessed for eligibility based on the indicative content provided, with 47 abstracts classified as

eligible for further inspection as full texts (intercoder reliability = 96.64%). Third, the full text articles

were assessed for eligibility based on the study inclusion criteria, with 34 studies progressing to data

extraction (intercoder reliability = 97.87%). The mean intercoder agreement across the three stages

of the screening process was 96.35%. Whenever data were missing to compute the effect sizes of

interest, this information was requested from the corresponding author for the respective article.

2.3. Data Processing

2.3.1. Data Extraction

The final 34 studies were split between the two co-first authors based on study focus, with

author two (ACV) extracting MEP data from 17 studies and author one (SC) extracting movement

outcome data from 17 studies, respectively. The co-first authors extracted the primary data (i.e.,

means, standards deviations, and samples sizes) and necessary methodological information to

investigate the proposed moderator effects for the present meta-analysis. Once complete, the final

author blind-checked the primary data for all studies and contacted the corresponding authors for

studies with missing primary data. If the authors did not respond within one month, or prior to the

study data analysis cut-off date (January 2021), the article was excluded from the present meta-

analysis. One study that collected MEP data and one study that collected movement outcome data

were excluded, leaving 32 studies in the final meta-analysis (see supplementary files for recorded

methodological information and primary data for all included articles).

2.3.2. Effect Size Preparation

For both sets of data, Cohen's *d* effect sizes were calculated using the mean, standard

deviation, and sample size values for the relevant outcome measures prior to conducting the main

data analyses using the 'robu' function from the 'robumeta' package (Fisher et al., 2017) in R studio

statistical software (version 2021.09.2 Build 382). The effect size preparation process varied

between the studies that collected MEP amplitude data and movement outcome data. All studies

that collected MEP amplitude data adopted a repeated measures design, meaning raw data was

recorded for all conditions of interest (i.e., AOMI, AO, MI, control). Additional steps were required

for studies that collected movement outcome data to account for differences in the movement

outcome variables and study designs adopted. When a reduction in a movement outcome measure

compared to control was considered an improvement (e.g., reaction time, movement time, mean

radial error), the polarity of the calculated effect size was inverted (i.e., positive reversed to negative

and vice versa) to ensure an increased value always indicated an improvement.

In order to produce standardized effect sizes that could be compared across the different

study designs, effect sizes were calculated for either pre- vs post-test gain comparisons or post- vs

post-test comparisons (see supplementary materials for all effect size calculation formulae). Pre-vs-

post-test gain comparisons were used to compute effect sizes in studies that adopted mixed

experimental designs (i.e., studies that allocated participants to a specific intervention condition and

collected data pre- and post-test) as this controls for any pre-test differences that exist between

intervention condition groups (Durlak, 2009). Post- vs post-test comparisons were used to compute

effect sizes in studies that adopted within-subject (i.e., repeated-measures) or between-subject (i.e.,

independent intervention condition groups) experimental designs. The number of effect sizes

included from each study was not limited as most studies recorded more than one outcome

measure for each comparison of interest (i.e., AOMI vs AO, MI, or control). Multiple effect sizes were

extracted from studies when MEP amplitudes were collected from more than one target muscle or

multiple movement outcome measures were recorded. Cohen's *d* effect size values were converted

into Hedge's *g* effect size values using a small sample size correction formula (Hedges, 1981) for

sensitivity analysis.

2.4. Data Analysis

Robust variance estimation (RVE) was used to analyze the primary effects for the MEP

amplitude data and movement outcome date in this meta-analysis. RVE was used because several

studies included across the two prongs of this meta-analysis reported multiple relevant effect sizes

which were not statistically independent of each other (cf. Tipton). RVE provides a method for

pooling dependent effect size estimates in the absence of any covariance values, mathematically

adjusting the standard errors of the effect sizes to account for their dependency (Tanner-Smith &

Tipton, 2014; Tanner-Smith et al., 2016). All analyses were run on RStudio using the 'metafor'

package (Viechtbauer, 2010) and the 'robumeta' package (Fisher et al., 2017).

2.4.1. Data Screening

Meta-analyses may be subject to multiple biases (Harrer et al., 2021). To address concerns

about publication bias, visual analyses were conducted on the data using funnel plots (Lau et al.,

2006), and subsequent statistical analyses were run with the Precision Effect Test (PET) using the

RVE approach (Alinaghi & Reed, 2018). For any data that showed publication bias, the trim-and-fill

procedure (Duval & Tweedie, 2000) was used to determine the number of unpublished studies

required to produce an unbiased estimate of the actual effect size. To address concerns about

sample size bias, a sensitivity analysis was run by repeating the main analyses with previously

calculated Hedge's *g* values. This found no meaningful differences in effect size estimate for the

primary comparisons, so Cohen's *d* values are reported in this meta-analysis (cf. Simonsmeier et al.,

2021). To address concerns about between-study heterogeneity, outlier diagnostics were completed

using the FIND.OUTLIERS function and influence analyses were run using the INFLUENCEANALYSIS

function from the 'dmetar' package in R (Harrer et al., 2019). This involved visual inspection of

"baujat", "influence" for effect sizes, and "leave-one-out" plots for both effect size and I^2 values.

Potential outliers and influential effect sizes were identified across the six primary comparisons for

the MEP amplitude and movement outcome data (see supplementary files for detailed overview of

data screening results). Removal of the outliers and influential cases had minimal impact on the

pooled effect and heterogeneity estimates for all but one comparison. Two effect sizes were

removed from the AO comparison for the movement outcome data (Frenkel-Toledo et al., 2020;

delta & Romano-Smith et al., 2019; peak angular velocity of the elbow) as the data points were

deemed outliers, influential, and resulted in a meaningful change to the pooled effect and

heterogeneity when removed from the RVE data analysis. All other effect sizes were retained in the

meta-analysis to preserve the richness of the data.

2.4.2. Quality Assessment

Study quality was assessed to identify if the studies included in this meta-analysis provide

reliably reported data, as well as indicating whether these studies reach an acceptable scientific

standard (Borenstein et al., 2021). To address these, the first author subjectively assessed the quality

of each study using an assessment scale employed in recent meta-analyses (see e.g., Harris et al.,

2021). This quality assessment scale was adapted from the Quality Index (Downs & Black, 1998), the

Checklist for the Evaluation of Research Articles (Durant, 1994), and the Appraisal Instrument

(Genaidy et al., 2007). The quality assessment checklist and individual scores for each study are

provided in the supplementary materials.

2.4.3. Primary and Moderator Effects

The primary meta-analyses for both the MEP amplitude and movement outcome data

involved correlational RVE models run using the 'robumeta' package (Fisher et al., 2017) in RStudio.

Overall, eleven moderators were chosen across both datasets based on previous meta-analyses and

reviews focused on the effects of MI and AO on corticospinal excitability (e.g., Grosprêtre et al.,

2016; Naish et al., 2014) and motor skill performance (e.g., Ashford et al., 2006; Simonsmeier et al.,

2021, Toth et al., 2020). Five moderators were shared across the MEP amplitude and movement

outcome data (*action observation perspective, skill classification, guided attentional focus,*

kinesthetic imagery ability, and age). Three moderators were specific to the MEP amplitude data

(*timing of TMS delivery, number of TMS trials, intensity of TMS pulses*). Five moderators were

specific to the movement outcome data (*population type, physical practice, incorporation of PETTLEP*

principles, context, and *intervention volume*). Subgroup analyses and meta regressions were used to

examine if these moderators influenced the effects of AOMI on the MEP amplitude and movement

outcome data compared to aggregate data from the AO, MI, and control groups. Subgroup analyses

were used to compare the effects of AOMI on MEP amplitude and movement outcome data for

moderators that permitted sub-division of the primary data sets based on nominal data (i.e.,

population type, action observation perspective, skill classification, guided attentional focus, timing

of TMS delivery, physical practice, incorporation of PETTLEP principles, and context). Regression

analyses were used to assess if moderators that consisted of interval data (i.e., kinesthetic *imagery*

ability, age, number of TMS trials, intensity of TMS pulses, and intervention volume) predicted the

effects of AOMI on MEP amplitude and movement outcome data.

2.4.3.1. Moderators for Both Sets of Data.

2.4.3.1.1. Action Observation Perspective. Studies have used first-person perspective AO

(e.g., Bruton et al., 2020; Romano-Smith et al., 2019) and third-person perspective AO (e.g., Taube et

al., 2014; Wright, Wood et al., 2018a) to examine the effects of AOMI on MEP amplitude and

movement outcome data. Sub-group analysis was used to compare the effects of AOMI on MEP

amplitude and movement outcome data between these two AO perspectives. First-person

perspective AO involved the participant viewing the action as if they were performing it (i.e.,

through their own eyes) and third-person perspective AO involved the participant viewing the action

as if another person video recorded them or another person was performing the action (i.e., filmed

from a vantage point away from the body). This was determined by checking written text and visual

stimuli included in the article.

372   *2.4.3.1.2. Skill Classification.* Diverse motor tasks ranging from finger movements (e.g.,

Meers et al., 2020) to walking (e.g., Kaneko et al., 2018) have been used in previous AOMI literature.

Sub-group analysis was used to compare the effects of AOMI on MEP amplitude and movement

outcome data for fine vs gross and continuous vs discrete motor tasks. The target movement

presented in the AOMI stimuli was classified using a one-dimensional skill classification approach

(Spittle, 2021, p.23). Based on this approach, intricate and precise movements using smaller muscle

groups (e.g., finger movements) were classed as fine motor tasks; larger muscle movements typically

based on fundamental movement patterns (e.g., balance tasks) were classed as gross motor tasks;

repetitive movements that have no distinct beginning or end (e.g., walking) were classed as

continuous motor tasks; and movements that have an identifiable beginning and end (e.g., putting a

golf ball) were classed as discrete motor tasks. Other skill classification comparisons (e.g., open vs

closed skills) were not considered in this moderator category if both motor skill categories were not

covered in the synthesized literature.

385   *2.4.3.1.3. Guided Attentional Focus.* Studies on AO have demonstrated different effects on

movement outcomes (e.g., D'Innocenzo et al., 2016) and MEP amplitudes (e.g., Wright, Wood et al.,

2018b) when visual attention is directed, or not, to a specific component of the observed action.

More recently, Bruton et al. (2020) showed that allocation of visual attention modulates the effects

of AOMI on MEP amplitudes for a finger movement task. Sub-group analysis was used to compare

the effects of AOMI on MEP amplitude and movement outcome data for guided attentional focus

(i.e., use of instructions to direct attention towards a specific aspect of the observed movement) vs

unguided attentional focus (i.e., no such instructions). Studies that did not explicitly state if visual

attention was manipulated during AOMI were included in the unguided attentional focus sub-group.

2.4.3.1.4. Kinesthetic Imagery Ability. Kinesthetic imagery is the imagery modality

instructed during AOMI and the effects of AOMI on movement outcomes reportedly vary as a

function of kinesthetic imagery ability (McNeill et al., 2020). Meta regression analyses were used to

assess if there was a relationship between kinesthetic imagery ability score and the effects of AOMI

on MEP amplitude and movement outcome data. Kinesthetic imagery ability data recorded using

valid self-report psychometric scales including the Vividness of Movement Imagery Questionnaire -2

(Roberts et al., 2008), Movement Imagery Questionnaire-3 (Williams et al., 2012) and the Movement

Imagery Questionnaire-Revised (Hall & Martin, 1997) were included for moderator analyses. Any

studies that used non-validated scales such as a visual analogue scale were excluded from the

moderator analyses. The imagery ability data was extracted from the studies and standardized by

reverse-scoring any measures that adopted an inverse scoring system such that higher numbers

meant better imagery ability, before converting all scores to percentage values based on the range

of values attainable for each scale.

2.4.3.1.5. Age. Studies have typically recruited adults ranging from early to middle

adulthood when assessing MEP amplitudes during AOMI (mean age = 27.07 ± 13.48 years) and

movement outcomes after AOMI (mean age = 30.89 ± 20.24 years). Studies have shown age-related

differences in MEP amplitudes during simulation of actions (e.g., Mouthon et al., 2016) and imagery

ability is proposed to decline across the lifespan (e.g., Gulyás et al., 2022), suggesting that age may

moderate the effects of AOMI on MEP amplitudes and movement outcomes. Meta regression

analysis was used to assess if there was a relationship between participant age and the effects of

AOMI on MEP amplitude and movement outcome data.

2.4.3.2. Moderators for MEP Amplitude Data.

2.4.3.2.1. Timing of Transcranial Magnetic Stimulation Delivery. AO and MI cause phase-

specific changes in corticospinal excitability (see e.g., Grosprêtre et al., 2016; Naish et al., 2014 for

reviews). Sub-group analysis was used to compare the effects of AOMI on MEP amplitude data for

TMS delivered at a random point after movement onset against TMS delivered at a targeted point

after movement onset (e.g., at the point of maximum movement of the limb). Timing of TMS

stimulation delivery refers to the point at which the single pulse is delivered based on the movement

displayed in the visual stimuli during AOMI.

2.4.3.2.2. Number of Transcranial Magnetic Stimulation Trials. The number of TMS trials

impacts the reliability of the MEP's evoked during single-pulse TMS (Goldsworthy et al., 2016). Meta

regression analysis was used to assess if there was a relationship between the number of TMS trials

and the effects of AOMI on MEP amplitude data. This was calculated by recording the number of

trials where single-pulse TMS was applied to the participant during AOMI for each study.

2.4.3.2.3. Intensity of Transcranial Magnetic Stimulation Pulses. The intensity of TMS

pulses impacts the reliability of the MEP's evoked during single-pulse TMS (Pellegrini et al., 2018).

Moderator analyses were used to assess if there was a relationship between number and intensity of

TMS trials and the effects of AOMI on MEP amplitude data. TMS stimulation intensity refers to the

intensity of the TMS stimulator output relative to the resting motor threshold, that is applied to the

participant during AOMI.

2.4.3.3. Moderators for Movement Outcome Data.

2.4.3.3.1. Population Type. AOMI interventions have been shown to benefit movement

outcomes in both neurotypical and neurodivergent populations (e.g., Scott et al., 2019). Sub-group

analysis was used to compare the effects of AOMI on movement outcome data for these two

population types. Neurotypical populations included individuals who are not characterized by

neurologically atypical patterns, thoughts, behavior, or diagnoses, and neurodivergent populations

included individuals whose neurological development and state are considered atypical.

2.4.3.3.2. Physical Practice. Studies have explored the effects of AOMI interventions on

movement outcomes with (e.g., Marshall & Wright, 2016) and without (e.g., Taube et al., 2014)

physical practice. Sub-group analysis was used to compare the effects of AOMI on movement

outcome data when used with physical practice vs without physical practice.

2.4.3.3.3. Incorporation of PETTLEP Principles. Some studies have adhered to PETTLEP

principles (Holmes & Collins, 2001) when developing and delivering AOMI interventions (e.g.,

Romano-Smith et al., 2019). Sub-group analysis was used to compare the effects of AOMI on

movement outcome data with the inclusion of PETTLEP principles vs without inclusion of PETTLEP

principles.

2.4.3.3.4. Context. AOMI interventions have been used to target changes in movement

outcomes in sport (e.g., Romano-Smith et al., 2018) and rehabilitation (e.g., Marusic et al., 2018)

contexts. Sub-group analysis compared the effects of AOMI on movement outcome data for sport vs

rehabilitation vs other contexts. Studies were classified as sport- or rehabilitation-focused. Studies

including movements that did not clearly fall into sports or rehabilitation contexts (e.g., finger

movements, ball rotation tasks) were classified as 'other'.

2.4.3.3.5. Intervention Volume. Studies have delivered AOMI interventions over short- (e.g.,

Bek et al., 2019) and longer-term (e.g., Shimada et al., 2019) periods when investigating their effects

on movement outcomes. Moderator analysis was used to assess the relationship between

intervention volume (total minutes) and the effects of AOMI on movement outcome data.

3. Results

3.1. Study Characteristics

Overall, this two-pronged meta-analysis analyzed 111 effect sizes from 32 studies. Of these,

54 effect sizes were from MEP amplitude data (n = 16 studies) and 57 effect sizes were from

movement outcome data (n = 16 studies). Studies included in this meta-analysis were published

between 2009 and 2021, with a total sample size of 823 participants split across studies that

collected MEP amplitude data (n = 234, 77 females, 92 males, 65 undisclosed) and studies that

collected movement outcome data (n = 589, 281 females, 308 males). The mean age of participants

was 27.07 ± 13.48 years and 30.89 ± 20.24 for the two respective prongs of the meta-analysis.

3.2. Study Quality

The study quality assessment indicated that all studies included in the two-pronged meta-

analysis displayed a high degree of rigor. For studies that collected MEP amplitude data, the quality

assessment scores ranged from 18.75-100%, with a mean of 89.58 ± 22.99%. The most poorly

addressed items were 'providing details of a priori sample size determination' and 'consistently

reporting effect sizes' with 18.75% and 43.75% of studies satisfying these criteria, respectively

(Figure 2a). For studies that collected movement outcome data, quality assessment scores ranged

from 31.25-100%, with a mean of 92.36 ± 17.36%. The most poorly addressed items were 'providing

details of a priori sample size determination' and 'applicability of study results to other relevant

populations' with 31.25% and 68.75% of studies satisfying these respective criteria (Figure 2b).

To assess whether studies with smaller samples sizes or lower quality studies were likely to

bias the results, meta-regression analyses were run between quality assessment scores and effect

size, and sample size and effect size using the '*robumeta*' and 'metafor' packages in R. For studies

that collected MEP amplitude data, the analysis used 54 effect sizes from 16 studies and reported

that neither the quality assessment scores (b = -0.02, p = .25), nor sample size (b = -0.01, p = .13)

predicted the overall effect of AOMI on MEP amplitudes. Similarly, for studies that collected

movement outcome data, the analysis used 57 effect sizes from 16 studies and reported that neither

quality assessment score (b = 0.01, p = .76) or sample size (b = -0.03, p = .06) predicted the overall

effect of AOMI on movement outcomes. The non-significant relationships between quality

assessment score, sample size and effect size indicate a low risk of bias for the studies included in

this meta-analysis (*cf*. Harris et al., 2021).

(A)

(B)

Figure 2. Bar chart displaying quality assessment items and scores for all studies that collected (A) MEP amplitude data and (B) movement outcome data
included in the meta-analysis. The blue bar indicates the number of studies that satisfied, and the red bar indicates the number of studies that did not
satisfy, each of the respective quality assessment criteria.

3.3. Primary Effects

3.3.1. MEP Amplitudes

Fifty-four effect sizes from sixteen studies were used to determine the overall effect of AOMI on MEP amplitudes. AOMI had a small to medium positive overall effect on MEP amplitudes compared to the control, AO and MI conditions in combination (d = 0.48, 95% CI [0.35, 0.61], $p <$ .001). The between-study heterogeneity variance was estimated at τ^2 = 0.00, with an I^2 value of 1.23%. The MEP amplitude data had no significant moderators (Tables 1 and 2), demonstrating a robust effect of AOMI on MEP amplitudes irrespective of kinesthetic imagery ability, sample age, intensity of TMS pulses, number of TMS trials, sample age, AO perspective, attentional focus strategy, skill classification, and the timing of TMS delivery.

Table 1. Meta-regression analyses for the MEP amplitude and movement outcome data

Moderator	N	K	Beta	P	Sig.	τ^2	I^2
MEP Amplitude							
Kinesthetic Imagery Ability	6	19	-0.01	**.90**	**No**	0.00	0.00
Intensity of TMS Pulses	15	53	0.06	**.13**	**No**	0.02	5.71
Number of TMS Trials	15	53	0.00	**.85**	**No**	0.02	11.8
Age	13	47	0.00	**.61**	**No**	0.00	0.00
Movement Outcome							
Intervention Volume	14	52	0.00	**.72**	**No**	0.40	71.03
Kinesthetic Imagery Ability	5	21	-0.01	**.86**	**No**	2.40	88.53
Age	16	57	-0.01	**.19**	**No**	0.38	68.73

Note. N = number of studies, K = number of effect sizes, Beta = regression coefficient, P = significance value, Sig. = significance threshold met or not, τ^2 = measure of variation around average, I^2 = measure of proportion of observed variance (%).

513 **Table 2**. **Sub-group analyses for the MEP amplitude data**

Sub-group	N	K	SMD	95% CI		P	Sig.	τ^2	I^2	Sig. Mod.
Action Observation Perspective										
First-person	9	23	0.50	0.33	0.67	**<.001**	**Yes**	0.00	0.00	**ref**
Third-person	7	31	0.48	0.12	0.84	**.02**	**Yes**	0.15	45.41	*ns*
Directed Attentional Focus										
Yes	1	2	0.38	NA	NA	**NA**	**NA**	0.00	0.00	*ref*
No	15	52	0.49	0.34	0.64	**<.001**	**Yes**	0.02	7.15	ns
Skill Classification 1										
Fine	10	24	0.48	0.33	0.64	**<.001**	**Yes**	0.00	0.00	ns
Gross	6	30	0.52	0.07	0.97	**.03**	**Yes**	0.19	52.98	*ref*
Skill Classification 2										
Discrete	12	32	0.52	0.35	0.70	**<.001**	**Yes**	0.00	0.00	*ns*
Continuous	4	22	0.34	0.16	0.52	**<.01**	**Yes**	0.09	33.19	ref
Timing of TMS Delivery										
Targeted	12	35	0.47	0.30	0.64	**<.001**	**Yes**	0.00	0.00	**ref**
Random	4	19	0.53	0.28	0.77	**<.001**	**Yes**	0.08	28.61	*ns*

Note. N = number of studies, K = number of effect sizes, SMD = standardized mean difference, 95% CI = lower and upper confidence intervals, P =
significance value, Sig. = significance threshold met or not, τ^2 = measure of variation around average, I^2 = measure of proportion of observed variance (%),
Sig. Mod. = comparison between sub-group categories. NA = insufficient number of data points for the analysis, ref = reference category, *ns* = no significant
difference compared to reference category

3.3.1.1. AOMI vs Control. Nineteen effect sizes from thirteen studies were used in this

analysis. AOMI had a medium positive effect on MEP amplitudes compared to control conditions (d =

0.54, 95% CI [0.41, 0.66], p < .001). The between-study heterogeneity variance was estimated at τ^2 =

0.00, with an I^2 value of 0.00%.

Figure 3. Forest plot of effect sizes (*d*) for all studies included in the meta-analysis that compared
MEP amplitude data for AOMI vs control conditions. The combined estimate and 95% confidence
interval (hollow diamond) indicates AOMI has a medium positive effect on MEP amplitudes
compared to control conditions. The size of each black square indicates the weight of the study
effect size in the combined analysis. Multiple effect sizes are reported for a study if it recorded MEP
amplitude data from more than one target muscle.

3.3.1.2. AOMI vs AO. Twenty-three effect sizes from thirteen studies were used in this analysis.

AOMI had a small to medium positive effect on MEP amplitudes compared to AO conditions (d =

0.45, 95% CI [0.27, 0.63], p < .001). The between-study heterogeneity variance was estimated at τ^2

0.02, with an I^2 value of 10.32%.

Figure 4. Forest plot of effect sizes (d) for all studies included in the meta-analysis that compared

MEP amplitude data for AOMI vs AO conditions. The combined estimate and 95% confidence

interval (hollow diamond) indicates AOMI has a small to medium positive effect on MEP amplitudes

compared to AO conditions. The size of each black square indicates the weight of the study effect

size in the combined analysis. Multiple effect sizes are reported for a study if it recorded MEP

amplitude data from more than one target muscle.

3.3.1.3. AOMI vs MI. Twelve effect sizes from six studies were used in this analysis. AOMI

had no significant effect on MEP amplitudes compared to MI conditions (d = 0.25, 95% CI [-0.13,

0.63], p = .14). The between-study heterogeneity variance was estimated at τ^2= 0.11, with an I^2 value

of 42.92%.

Figure 5. Forest plot of effect sizes (d) for all studies included in the meta-analysis that compared
MEP amplitude data for AOMI vs MI conditions. The combined estimate and 95% confidence
interval (hollow diamond) indicates AOMI has no significant effect on MEP amplitudes compared to
MI conditions. The size of each black square indicates the weight of the study effect size in the
combined analysis. Multiple effect sizes are reported for a study if it recorded MEP amplitude data
from more than one target muscle.

3.3.2. Movement Outcomes

Fifty-seven effect sizes from sixteen studies were used to determine the overall effect of

AOMI on movement outcomes. AOMI had a small to medium positive overall effect on movement

outcomes compared to the control, AO and MI conditions in combination (d = 0.48, 95% CI [0.18,

0.78], p < .01). The between-study heterogeneity variance was estimated at τ^2 = 0.39 , with

an I^2 value of 69.68%. The movement outcome data had no significant moderators (Tables 1 and 3),

demonstrating a robust effect of AOMI on movement outcomes irrespective of intervention volume,

kinesthetic imagery ability, sample age, AO perspective, study context, attentional focus strategy,

incorporation of PETTLEP, physical practice, population type, and skill classification.

Table 3. **Sub-group analyses for the movement outcome data**

Sub-group	N	K	SMD	95% CI		P	Sig.	τ^2	I^2	Sig. Mod.
Action Observation Perspective										
First-person	8	29	0.59	0.08	1.09	**.03**	**Yes**	0.61	73.68	*ref*
Third-person	8	28	0.39	-0.07	0.84	**.08**	**No**	0.29	67.28	**ns**
Context										
Sport	4	17	0.68	-0.65	2.02	**.20**	**No**	1.38	86.11	**ns**
Rehabilitation	7	28	0.43	-0.13	1.00	**.11**	**No**	0.42	70.93	*ns*
Other	5	12	0.43	-0.03	0.88	**.06**	**No**	0.07	31.62	**ref**
Directed Attentional Focus										
Yes	4	15	0.38	-0.91	1.68	**.41**	**No**	0.59	82.46	*ref*
No	12	42	0.52	0.22	0.82	**<.01**	**Yes**	0.33	63.07	**ns**
PETTLEP										
Yes	5	19	0.69	-.039	1.76	**.15**	**No**	0.83	80.52	*ref*
No	11	38	0.39	0.11	0.67	**.01**	**Yes**	0.28	63.71	**ns**
Physical Practice										
Yes	2	7	0.75	-6.12	7.63	**.40**	**No**	3.96	91.29	*ref*
No	14	50	0.45	0.13	0.77	**.01**	**Yes**	0.28	64.11	**ns**
Population										
Neurotypical	13	46	0.47	0.19	0.75	**<.01**	**Yes**	0.29	62.26	*ref*
Neurodivergent	3	11	0.55	-2.05	3.14	**.46**	**No**	1.27	87.65	**ns**
Skill Classification 1										
Fine	12	41	0.45	0.07	0.83	**.03**	**Yes**	0.32	67.52	**ns**
Gross	4	16	0.59	-0.12	1.31	**.08**	**No**	0.98	79.46	*ref*
Skill Classification 2										
Continuous	2	10	0.47	-1.66	2.60	**.22**	**No**	0.28	56.32	**ref**
Discrete	14	47	0.48	0.14	0.83	**.01**	**Yes**	0.43	72.18	*ns*

Note. N = number of studies, K = number of effect sizes, SMD = standardized mean difference, 95% CI = lower and upper confidence intervals, P =
significance value, Sig. = significance threshold met or not, τ^2 = measure of variation around average, I^2 = measure of proportion of observed variance (%),
Sig. Mod. = comparison between sub-group categories. NA = insufficient number of data points for the analysis, ref = reference category, *ns* = no significant
difference compared to reference category

3.3.2.1. AOMI vs Control. Twenty-seven effect sizes from twelve studies were used in this

analysis. AOMI had a medium to large positive effect on movement outcomes compared to control

conditions (*d* = 0.67, 95% CI [0.16, 1.18], *p* = .02). The between-study heterogeneity variance was

estimated at τ^2 = 0.48, with an I^2 value of 70.74%.

Figure 6. Forest plot of effect sizes (*d*) for all studies included in the meta-analysis that compared
movement outcome data for AOMI vs control conditions. The combined estimate and 95%
confidence interval (hollow diamond) indicates AOMI has a medium to large positive effect on
movement outcomes compared to control conditions. The size of each black square indicates the
weight of the study effect size in the combined analysis. Multiple effect sizes are reported for a study
if it recorded more than one movement outcome variable.

3.3.2.2. AOMI vs AO. Nineteen effect sizes from nine studies were used in this analysis.

AOMI had a small to medium positive effect on movement outcomes compared to AO conditions (d

= 0.44, 95% CI [0.07, 0.81], *p* = .03). The between-study heterogeneity variance was estimated at τ^2=

0.24, with an I^2 value of 63.23%.

Figure 7. Forest plot of effect sizes (*d*) for all studies included in the meta-analysis that compared

movement outcome data for AOMI vs AO conditions. The combined estimate and 95% confidence

interval (hollow diamond) indicates AOMI has a small to medium positive effect on movement

outcomes compared to AO conditions. The size of each black square indicates the weight of the

study effect size in the combined analysis. Multiple effect sizes are reported for a study if it recorded

more than one movement outcome variable.

3.3.2.3. AOMI vs MI. Eleven effect sizes from six studies were used in this analysis. AOMI

had no significant effect on movement outcomes compared to MI conditions (d = 0.53, 95% CI [-

0.59, 1.66], p = .28). The between-study heterogeneity variance was estimated at τ^2 = 1.30, with

an I^2 value of 84.46%.

Figure 8. Forest plot of effect sizes (d) for all studies included in the meta-analysis that compared
movement outcome data for AOMI vs MI conditions. The combined estimate and 95% confidence
interval (hollow diamond) indicates AOMI has no significant effect on movement outcomes
compared to MI conditions. The size of each black square indicates the weight of the study effect
size in the combined analysis. Multiple effect sizes are reported for a study if it recorded more than
one movement outcome variable.

3.4. Publication Bias

Based on the funnel plots of effect size level (see supplementary files), publication bias was

identified to be unlikely for all comparisons in the movement outcome and MEP data. Regardless,

we ran trim and fill and PET analyses using the RVE method to retrieve an unbiased effect size

estimate corrected for publication bias for all comparisons.

For the control comparison in the MEP amplitude data, both the PET-intercept (b_0 = 0.09, p =

0.83) and the PET-slope (b_1 = 1.21, p = 0.32) were not statistically significant, suggesting that

publication bias was unlikely. For the AO comparison in the MEP amplitude data, the funnel plot of

effect size level (Figure 9, left) indicated asymmetry. Trim-and-fill analysis proposed 4 missing values

(Figure 9, right), and the effect size changed from a medium (d = 0.54) to a small to medium positive

effect (d = 0.43), suggesting minimal effects of publication bias for this dataset. For the PET, both the

PET-intercept (b_0 = -1.00, p = 0.16) and the PET-slope (b_1 = 3.85, p = 0.07) were not statistically

significant, suggesting that publication bias was unlikely. For the MI comparison in the MEP

amplitude data, the funnel plots indicated no funnel asymmetry and the trim-and-fill analysis

reported zero missing values. For the PET, both the PET-intercept (b_0 = -1.14, p = 0.43) and the PET-

slope (b_1 = 3.59, p = 0.41) were not statistically significant, suggesting that publication bias was

unlikely.

For the control comparison in the movement outcome data, the funnel plots and trim and

fill analysis, showed no signs of asymmetry. Both the PET-intercept (b_0 = -0.38, p = 0.56) and the PET-

slope (b_1 = 4.34, p = 0.26) were not statistically significant, suggesting that publication bias was

unlikely. For the AO comparison in the movement outcome data, the funnel plots and trim-and-fill

analysis showed no signs of asymmetry. Both the PET-intercept (b_0 = -0.08, p = .88) and the PET-

slope (b_1 = 2.04, p = .43) were not statistically significant, suggesting that publication bias was

unlikely. For the MI comparison in the movement outcome data, the funnel plots and trim and fill

analysis showed no signs of asymmetry. Both the PET-intercept (b_0 = -0.51, p = .62) and the PET-

slope (b_1 = 3.65, p = .48) were not statistically significant, suggesting that publication bias was

unlikely.

Figure 9. Funnel plot of effect sizes (Cohen's *d*) versus standard error before (left) and after (right) performing Duval and Tweedie's Trim-and-fill analysis
for the AO comparison in the MEP amplitude data. Black circles represent existing effects included in the meta-analysis and white circles represent
potential unpublished effects. The contour-enhanced funnel plots display the significance of the effects for the AO comparison. Individual effect sizes falling
inside the white (*p* < .05) and orange (*p* < .01) funnel boundaries represent significant effects for the AO comparison

4. Discussion

Since the early reviews introducing AOMI (e.g., Eaves, Riach et al., 2016; Vogt et al., 2013),

researchers have studied its effects as a motor simulation intervention. The purpose of the current

study was to methodically collate and synthesize the available MEP amplitude data as an indicator of

corticospinal excitability during AOMI engagement, and movement outcome data to assess changes

in behavior that result from AOMI interventions. The purpose of the current meta-analysis was two-

pronged; first, to methodically collate and synthesize the available MEP amplitude data as an

indicator of corticospinal excitability during AOMI engagement, and second, to methodically collate

and synthesize the available movement outcome data to assess changes in motor skill performance

that result from AOMI interventions. Based on previous literature (see Eaves, Riach et al., 2016), it

was hypothesized that AOMI would have a small positive effect compared to independent AO or MI,

and a moderate positive effect compared to control conditions, for both outcome variables. The

results of this meta-analysis partially support this hypothesis. For the MEP amplitude data, AOMI

had a small to medium positive overall effect, a medium positive effect compared to control

conditions, a small to medium positive effect compared to AO, and no significant effect compared to

MI. For the movement outcome data, AOMI had a small to medium positive overall effect, a medium

to large positive effect compared to control, a small to medium positive effect compared to AO, and

no significant effect compared to MI conditions.

4.1. MEP Amplitudes

In this meta-analysis, AOMI had a medium positive effect compared to control conditions

and a small to medium positive effect compared to AO, but showed no effect compared to MI. TMS

studies have consistently reported increased corticospinal facilitation for AOMI across diverse motor

tasks such as simple finger movements (Bruton et al., 2020), walking (Kaneko et al., 2018), and

basketball free throws (Wright, Wood et al., 2018a). From a theoretical standpoint, the finding that

corticospinal excitability was facilitated for AOMI compared to control and AO but not MI conditions

aligns with the propositions of the VGH that AOMI is driven by MI but may oppose the sentiments of

the DASH (Eaves et al., 2012, 2014). The VGH suggests that observed and imagined actions are not

co-represented, and MI is the driver for increases in motor activity during AOMI. Specifically, Meers

et al. (2020) suggest the AO component acts as a visual primer, facilitating the production of more

vivid images during AOMI compared to AO and MI conditions. Alternatively, the DASH proposes that

concurrent representations of observed and imagined actions can be maintained as two quasi-

encapsulated sensorimotor streams and that these will merge, rather than compete, when a person

is overtly and covertly simulating the same action during AOMI. The merging of these two

sensorimotor streams is likely to produce more widespread activity in the premotor cortex (see

Filimon et al., 2015) than the AO, MI and control conditions, contributing to increased corticospinal

excitability via cortico-cortical connections linking premotor and motor cortices (Fadiga et al., 2005).

The non-significant increase in MEP amplitudes for AOMI compared to MI reported in this

meta-analysis may be explained by the propositions of the VGH, as the increased imagery vividness

for AOMI vs MI could be expected to be represented by a smaller difference in MEP amplitude

between these two conditions. This difference could be expected to be negligible if those simulating

actions were able to generate clear and vivid kinesthetic imagery without a visual primer, as was the

case for the participants synthesized in this meta-analysis (mean normalized kinesthetic imagery

ability score = 67.41%, median = 70.83%, range = 55% - 76.33%). Current evidence is conflicting for

the VGH and DASH accounts of AOMI (see Bruton et al., 2020; Meers et al., 2020), but both

hypotheses offer feasible explanations for the impact of AOMI on the motor system and thus

warrant further systematic investigation.

4.2. Movement Outcomes

AOMI had a medium to large positive effect on movement outcomes compared to control

and a small to medium positive effect compared to AO conditions. Such positive effects are

evidenced across most studies included in this meta-analysis, with movements ranging from dart

throwing (Romano-Smith et al., 2018, 2019) to whole-body balance tasks (Taube et al., 2014), in

both neurotypical (e.g., Aoyama et al., 2020) and neurodivergent populations (e.g., Marshall et al.,

2020). The increased motor activity during AOMI, as discussed in the previous section, is a possible

neurophysiological mechanism for this effect on movement outcomes. Repeated engagement in

AOMI, and thus activation of the motor system, has the potential to support repetitive Hebbian

modulation of intracortical and subcortical excitatory mechanisms through synaptic plasticity, in a

similar manner to physical practice (Holmes & Calmels, 2008). From a cognitive perspective, AO and

MI help develop mental representations that comprise cognitive information relating to movement

execution (Frank et al., 2020). When executing a motor task, a person recalls the relevant mental

representation and uses this to guide their movement (Frank et al., 2020). AO and MI are proposed

to contribute differently to the development of such mental representations, with AO providing

sequential and timing information and MI providing sensory information related to the movement

(Kim et al., 2017). It is possible combining the two forms of motor simulation during AOMI allows for

the effective development of mental representations of action in the long-term memory, benefitting

the physical execution of a motor task (Frank et al., 2020; Kim et al., 2017; Wright et al., 2021).

705   In contrast, AOMI had no significant effect on movement outcomes compared to MI

conditions. Robust evidence supports the efficacy of MI as an intervention to improve motor

performance across settings (e.g., MI: Guillot & Collet, 2008). This null finding aligns with the effects

of AOMI on corticospinal excitability when compared to MI in this meta-analysis. Specifically, AOMI

did not increase corticospinal excitability or improve movement outcomes when compared to MI

conditions. This provides further support for the VGH account for AOMI (Meers et al., 2020),

suggesting that the imagery component drives the effects of AOMI on both the motor system of the

brain and subsequent adaptations to physical movement. The sample synthesized for the MI

comparison in the movement outcome data were more able imagers than the sample synthesized

for the MI comparison in the MEP amplitude data in this meta-analysis (mean normalized kinesthetic

imagery ability score = 74.78%, range = 56.67% - 96.17%). This could suggest that individuals with

high imagery ability do not benefit from AOMI because the visual primer provided by AO does not

make their MI more vivid or clear during AOMI.

The findings of this meta-analysis promote AOMI as an effective alternative intervention to

AO and MI as well-established approaches, but do not indicate that combining AO and MI

simultaneously (i.e., AOMI) has an additive benefit towards motor performance compared to MI. It is

worthwhile noting that AOMI had a small to medium positive effect on movement outcomes

compared to MI (d = 0.53) despite the lack of significant differences. This is an important

consideration in applied settings, such as sport and neurorehabilitation, where marginal

improvements in motor performance can have practical significance (Lakens, 2013). AOMI

interventions are a suitable alternative to AO and MI interventions as this combined approach can

address the reported limitations of using either simulation technique in isolation. The capacity to

generate and maintain mental images, termed 'imagery ability', is a complex cognitive process that

is variable within- and between-populations (Cumming & Eaves, 2018). Individuals with low imagery

ability typically find it difficult to generate and control imagined content during MI interventions, an

issue that is not present for AO interventions as specific movement content can be displayed via

video (Holmes & Calmels, 2008). However, the effectiveness of AO interventions is dependent on

the observer's ability to attend to the most important aspects of the motor task being performed

(D'Innocenzo et al., 2016). Based on current recommendations for delivering AOMI interventions

(see Wright et al., 2021), AOMI has the capacity to control the visual information displayed via AO

whilst directing the individual's attention by getting them to focus on kinesthetic aspects of the

movement, subsequently reducing the complexity of MI.

4.4. Limitations and Future Research Recommendations

4.4.1. Study Reporting

A secondary aim of this meta-analysis was to explore several methodological parameters

hypothesized to have a moderating effect on the impact of AOMI interventions on MEP amplitudes

or movement outcomes. This was conducted to try and understand the influence of key

methodological aspects raised in early reviews on AOMI (Eaves, Riach et al., 2016; Vogt et al., 2013)

and to provide recommendations for future research and delivery of AOMI interventions. Whilst

moderator analyses were run in the form of meta-regression and sub-group analyses, only the sub-

group analyses included the full sets of studies (i.e., 16 studies per analysis), with missing study

information meaning that meta-regression analyses included 75% of the studies on average (mean =

12 studies, min = 6 studies for kinesthetic imagery ability, max = 16 studies for sample age). The

issue of inadequate reporting has been raised in recent meta-analyses focusing on imagery

interventions, with both noting issues related to imagery ability and assessment and reporting across

studies (see e.g., Simonsmeier et al., 2021; Toth et al., 2020). A recent article has provided guidance

for authors to standardise and improve the quality of reporting for action simulation studies

(Hardwick et al., 2022). Alongside adhering to these useful guidelines, we also recommend that

authors consider attempting to address the biases made apparent by the sub-group analyses for

AOMI literature in the current meta-analysis. For TMS studies, we suggest researchers employ

guided attentional focus, include more diverse motor tasks (i.e., gross/continuous/serial/open skills),

and test the impact of the timing of TMS delivery on the effects of AOMI on MEP amplitudes. For

behavioral studies, we recommend that researchers employ guided attentional focus, adhere to

PETTLEP guidelines, incorporate physical practice, recruit neurodivergent populations, and include

more diverse motor tasks (i.e., gross/continuous/serial/open skills) when studying the effects of

AOMI on movement outcomes.

4.4.1. Neurophysiological Modality

This meta-analysis synthesized MEP amplitude data from AOMI studies using single-pulse

TMS. TMS has been widely used as a neurophysiological modality with AOMI as it permits the

recording of muscle-specific facilitation in corticospinal excitability, an effect that has been

demonstrated robustly for AO and MI (Naish et al., 2014; Grosprêtre et al., 2016). Whilst single-pulse

TMS provides an indication of activity within the motor and premotor cortices of the brain during

AOMI, EEG and fMRI can be used to provide complimentary knowledge about the roles of other

cortical regions during AOMI as they can measure whole-brain activity and have high temporal and

spatial accuracy, respectively (Holmes & Wright, 2017). However, studies have shown that AOMI

leads to activity in brain regions that would not be activated directly during the delivery of TMS to

the primary motor cortex (e.g., rostral prefrontal cortex; Eaves, Behmer, et al., 2016), and therefore

not considered within this meta-analysis. Consequently, there is a need to collate and synthesize

data on the precise anatomical substrates involved in AOMI using neuroscientific methods with

increased spatial resolution. Hardwick et al. (2018) recently performed a large-scale activation

likelihood estimation meta-analysis on fMRI data for AO, MI, and movement execution to identify

distinct and shared neural regions for these three action states. This approach could be adopted

once additional fMRI data is available for AOMI to advance understanding of the neural mechanisms

underpinning engagement in this form of simulation.

4.4.2. Study Designs

To-date, AOMI studies have almost entirely explored the short- term effects of this

intervention on movement outcomes, using a between-groups comparison at one time point or

adopting a pre- vs post-test study design. Whilst this approach is typical in randomized controlled

trials of interventions, this does not permit accurate assessment of the long-term changes that result

from AOMI engagement. The benefits of AO and MI on movement outcomes are reportedly greatest

during or immediately after training, with the positive outcomes gradually disappearing in the

absence of simulated practice (Stevens et al., 2003; Zhang et al., 2019). However, the performance

benefits of MI training are retained beyond the intervention period (Simonsmeier et al., 2021), with

repetitive engagement in MI inducing neural plasticity during recovery phases when this technique is

used to acquire a skill (Ruffino et al., 2017). It remains unclear if the improvements in movement

outcome associated with AOMI are maintained after the intervention is withdrawn. Future studies

should draw from the methodological approaches adopted in motor learning literature (e.g.,

Krakauer et al., 2019) to comprehensively examine the effectiveness of AOMI when learning and

improving movement outcomes for different populations and motor tasks.

4.4.3. Brain-Behavior Interactions

797   This meta-analysis synthesized data on MEP amplitudes during AOMI engagement and

changes in movement outcomes after AOMI interventions. The crossover between these effects

could not be analyzed in the current meta-analysis as no studies to our knowledge have collected

both measures for AOMI whilst satisfying the inclusion criteria of this meta-analysis. The results of

this meta-analysis showed that AOMI had a positive effect on MEP amplitudes and movement

outcomes compared to control and AO conditions. Collecting neurophysiological responses, such as

MEP amplitudes, alongside movement outcome measures will advance our understanding of the

relationship between the changes in brain activity and motor skill performance for this intervention.

Such integration of measures has shown that corticospinal excitability during MI is related to the

magnitude of motor cortical adaptations after MI training (Yoxon & Welsh, 2020), and that MI and

physical practice lead to different changes in brain activity and subsequent movement execution

after equal training bouts (Kraeutner et al., 2020). Future research should combine neuroscientific

and human movement science methods to elucidate the neural mechanisms underlying

improvements in motor skill performance and learning through AOMI.

811   **5. Conclusion**

812   This two-pronged meta-analysis synthesizes the existing MEP amplitude and movement

outcome data for AOMI to compare its effectiveness against AO, MI, and control conditions. The

results of the present meta-analysis report that AOMI had a small to medium positive overall effect

on MEP amplitudes, as an indicator of corticospinal excitability. When compared to different

conditions, AOMI has a medium positive effect compared to control, a small to medium positive

effect compared to AO, and no significant effect compared to MI conditions. For the movement

outcome data, AOMI had a small to medium positive overall effect, a medium to large positive effect

compared to control, a small to medium positive effect compared to AO, and no significant effect

compared to MI conditions. No methodological factors moderated the effects of AOMI on MEP

amplitudes or movement outcomes, indicating a robust effect of AOMI for both outcome variables.

However, it should be noted that inadequate reporting of methodological information as well as

limited variation in the current literature on AOMI may have resulted in biased comparisons being

made between moderator sub-groups and low powered assessments of relationships. Overall, the

findings of this meta-analysis support the effectiveness of AOMI as an alternative intervention to AO

and MI, two well established interventions, as it addresses the limitations of these approaches in

isolation when targeting increased activity in motor regions of the brain and improvements in motor

skill performance. A more methodologically diverse approach that integrates brain and behavior is

needed in future AOMI research to advance the current state of knowledge for this intervention.

Funding statement: This research did not receive any specific grant from funding agencies in the

public, commercial, or not-for-profit sectors.

833   **6. References**

Alinaghi, N., & Reed, W. R. (2018). Meta-analysis and publication bias: How well does the FAT-PET-

PEESE procedure work? Research Synthesis Methods, 9(2), 285–311.

https://doi.org/10.1002/jrsm.1298

Aoyama, T., Kaneko, F., & Kohno, Y. (2020). Motor imagery combined with action observation

training optimized for individual motor skills further improves motor skills close to a

plateau. *Human Movement Science*, *73*, 102683.

https://doi.org/10.1016/j.humov.2020.102683

Ashford, D., Bennett, S. J., & Davids, K. (2006). Observational modeling effects for movement

dynamics and movement outcome measures across differing task constraints: A meta-

analysis. *Journal of Motor Behavior*, *38*(3), 185-205. https://doi.org/10.3200/JMBR.38.3.185-

205

Bek, J., Gowen, E., Vogt, S., Crawford, T. J. & Poliakoff, E. (2019). Combined action observation and

motor imagery influences hand movement amplitude in Parkinson's disease. *Parkinsonism*

and Related Disorders, 61, 126-131. https://doi.org/10.1016/j.parkreldis.2018.11.001

Berends, H., Wolkorte, R., Ijzerman, M., & van Putten, M. (2013). Differential cortical activation

during observation and observation-and-imagination. *Experimental Brain Research*, *229*(3),

337-345. https://doi.org/10.1007/s00221-013-3571-8

Borenstein, M., Hedges, L. V., Higgins, J. P., & Rothstein, H. R. (2021). *Introduction to meta-analysis*.

John Wiley & Sons.

Bruton, A., Holmes, P., Eaves, D., Franklin, Z., & Wright, D. (2020). Neurophysiological markers

discriminate different forms of motor imagery during action observation. *Cortex*, *124*, 119-

136. https://doi.org/10.1016/j.cortex.2019.10.016

Buccino, G. (2014). Action observation treatment: A novel tool in neurorehabilitation. *Philosophical*

Transactions of the Royal Society B: Biological Sciences, *369*(1644), 20130185.

https://doi.org/10.1098/rstb.2013.0185

Caligiore, D., Mustile, M., Spalletta, G., & Baldassarre, G. (2017). Action observation and motor

imagery for rehabilitation in Parkinson's disease: A systematic review and an integrative

hypothesis. *Neuroscience & Biobehavioral Reviews*, *72*, 210-222.

https://doi.org/10.1016/j.neubiorev.2016.11.005

Castro, F., Bryjka, P.A., Di Pino, G., Vuckovic, A., Nowicky, A. & Bishop, D. (2021). *Brain and Cognition*,

152, 105768. https://doi.org/10.1016/j.bandc.2021.105768

Cumming, J., & Eaves, D. L. (2018). The nature, measurement, and development of imagery

ability. *Imagination, Cognition and Personality*, *37*(4), 375-393.

https://doi.org/10.1177%2F0276236617752439

D'Innocenzo, G., Gonzalez, C. C., Williams, A. M., & Bishop, D. T. (2016). Looking to learn: the effects

of visual guidance on observational learning of the golf swing. *PloS one, 11*(5), e0155442.

https://doi.org/10.1371/journal.pone.0155442

De Vries, S., & Mulder, T. (2007). Motor imagery and stroke rehabilitation: A critical

discussion. *Journal of Rehabilitation Medicine*, *39*(1), 5-13.

https://doi.org10.2340/16501977-0020

Di Rienzo, F., Joassy, P., Kanthack, T., MacIntyre, T., Debarnot, U., & Blache, Y. et al. (2019). Effects of

Action Observation and Action Observation Combined with Motor Imagery on Maximal

Isometric Strength. *Neuroscience, 418*, 82-95.

https://doi.org/10.1016/j.neuroscience.2019.08.025

Downs, S. H., & Black, N. (1998). The feasibility of creating a checklist for the assessment of the

methodological quality both of randomised and non-randomised studies of health care

interventions. *Journal of Epidemiology & Community Health*, *52*(6), 377-384.

http://dx.doi.org/10.1136/jech.52.6.377

Durant, J. (1994). What is scientific literacy? *European Review*, *2*(1), 83-89.

https://doi.org/10.1017/S1062798700000922

Durlak, J. A. (2009). How to select, calculate, and interpret effect sizes. *Journal of Pediatric*

Psychology, *34*(9), 917-928. https://doi.org/10.1093/jpepsy/jsp004

Duval, S., & Tweedie, R. (2000). A nonparametric "trim and fill" method of accounting for publication

bias in meta-analysis. *Journal of the American Statistical Association*, *95*(449), 89-98.

https://doi.org/10.1080/01621459.2000.10473905

Eaves, D. L., Behmer, L., & Vogt, S. (2016). EEG and behavioural correlates of different forms of

motor imagery during action observation in rhythmical actions. *Brain and Cognition*, *106*, 90-

103. https://doi.org/10.1016/j.bandc.2016.04.013

Eaves, D. L., Haythornthwaite, L., & Vogt, S. (2014). Motor imagery during action observation

modulates automatic imitation effects in rhythmical actions. *Frontiers in Human*

Neuroscience, *8*. https://doi.org/10.3389/fnhum.2014.00028

Eaves, D. L., Riach, M., Holmes, P. S., & Wright, D. J. (2016). Motor imagery during action

observation: a brief review of evidence, theory and future research opportunities. *Frontiers*

in Neuroscience, *10*, 514. https://doi.org/10.3389/fnins.2016.00514

Eaves, D. L., Turgeon, M., & Vogt, S. (2012). Automatic imitation in rhythmical actions: Kinematic

fidelity and the effects of compatibility, delay, and visual monitoring. *PLoS One*, *7*(10),

e46728. https://doi.org/10.1371/journal.pone.0046728

Emerson, J. R., Binks, J. A., Scott, M. W., Kenny, R. P., & Eaves, D. L. (2018). Combined action

observation and motor imagery therapy: A novel method for post-stroke motor

rehabilitation. *AIMS Neuroscience*, *5*(4), 236-252.

https://doi.org/10.3934/Neuroscience.2018.4.236

Fadiga, L., Craighero, L., & Olivier, E. (2005). Human motor cortex excitability during the perception

of others' action. *Current Opinion in Neurobiology*, *15*(2), 213-218.

https://doi.org/10.1016/j.conb.2005.03.013

Filimon, F., Rieth, C., Sereno, M., & Cottrell, G. (2015). Observed, executed, and imagined action

representations can be decoded from ventral and dorsal areas. *Cerebral Cortex*, *25*(9), 3144-

3158. https://doi.org/10.1093/cercor/bhu110

Fisher, Z., Tipton, E., Zhipeng, H., & Fisher, M. Z. (2017). Package 'robumeta'.

Frank, C., Wright, D., & Holmes, P. (2020). Mental simulation and neurocognition: Advances for

motor imagery and action observation training in sport. In D. Hackfort & R. J. Schinke (Eds.),

Routledge International Encyclopaedia of Sport and Exercise Psychology (Vol. 2., pp. 411-

428). Routledge. https://doi.org/10.4324/9781315187228

Genaidy, A. M., Lemasters, G. K., Lockey, J., Succop, P., Deddens, J., Sobeih, T., & Dunning, K. (2007).

An epidemiological appraisal instrument – a tool for evaluation of epidemiological

studies. *Ergonomics*, *50*(6), 920-960. https://doi.org/10.1080/00140130701237667

Goldsworthy, M. R., Hordacre, B., & Ridding, M. C. (2016). Minimum number of trials required for

within-and between-session reliability of TMS measures of corticospinal excitability.

Neuroscience, 320, 205-209. https://doi.org/10.1016/j.neuroscience.2016.02.012.

Grosprêtre, S., Ruffino, C., & Lebon, F. (2016). Motor imagery and cortico-spinal excitability: A

review. *European Journal of Sport Science*, *16*(3), 317-324.

https://doi.org/10.1080/17461391.2015.1024756

Guillot, A., & Collet, C. (2008). Construction of the motor imagery integrative model in sport: A

review and theoretical investigation of motor imagery use. *International Review of Sport and*

Exercise Psychology, *1*(1), 31-44. https://doi.org/10.1080/17509840701823139

Hall, C. R., & Martin, K. A. (1997). Measuring movement imagery abilities: a revision of the

movement imagery questionnaire. *Journal of Mental Imagery*.

Hardwick, R., Caspers, S., Eickhoff, S., & Swinnen, S. (2018). Neural correlates of action: Comparing

meta-analyses of imagery, observation, and execution. *Neuroscience & Biobehavioral*

Reviews, *94*, 31-44. https://doi.org/10.1016/j.neubiorev.2018.08.003

Harrer, M., Cuijpers, P., Furukawa, T. & Ebert, D. D. (2019). dmetar: Companion R Package for the

Guide 'DoingMeta-Analysis in R'. R package version 0.0.9000. URL:

http://dmetar.protectlab.org/.

Harrer, M., Cuijpers, P., Furukawa, T. A., & Ebert, D. D. (2021). *Doing Meta-Analysis with R: A Hands-*

On Guide. Chapmann & Hall; CRC Press. URL:

https://bookdown.org/MathiasHarrer/Doing_Meta_Analysis_in_R/

Harris, D. J., Allen, K. L., Vine, S. J., & Wilson, M. R. (2021). A systematic review and meta-analysis of

the relationship between flow states and performance. *International Review of Sport and*

Exercise Psychology, 1-29. https://doi.org/10.1080/1750984X.2021.1929402

Hedges, L. V. (1981). Distribution theory for Glass's estimator of effect size and related

estimators. *Journal of Educational Statistics*, *6*(2), 107-128.

https://doi.org/10.3102/10769986006002107

Holmes, P. S., & Calmels, C. (2008). A neuroscientific review of imagery and observation use in

sport. *Journal of Motor Behavior*, *40*(5), 433-445. https://doi.org/10.3200/jmbr.40.5.433-445

Holmes, P. S., & Collins, D. J. (2001). The PETTLEP approach to motor imagery: A functional

equivalence model for sport psychologists. *Journal of Applied Sport Psychology, 13*(1), 60-83.

https://doi.org/10.1080/10413200109339004

Holmes, P. S., & Wright, D. J. (2017). Motor cognition and neuroscience in sport psychology. *Current*

Opinion in Psychology, *16*, 43-47. https://doi.org/10.1016/j.copsyc.2017.03.009

Jeannerod, M. (1994). Motor representations and reality. *Behavioral and Brain Sciences*, *17*(2), 229-

245. https://doi.org/10.1017/s0140525x0003435x

Jeannerod, M. (2001). Neural simulation of action: A unifying mechanism for motor

cognition. *Neuroimage*, *14*(1), S103-S109. https://doi.org/10.1006/nimg.2001.0832

Jeannerod, M. (2006). The origin of voluntary action. History of a physiological concept. *Comptes*

Rendus Biologies, *329*(5-6), 354-362. https://doi.org/10.1016/j.crvi.2006.03.017

Kaneko, N., Masugi, Y., Yokoyama, H., & Nakazawa, K. (2018). Difference in phase modulation of

corticospinal excitability during the observation of the action of walking, with and without

motor imagery. *NeuroReport*, *29*(3), 169-173.

https://doi.org/10.1097/wnr.0000000000000941

Kawasaki, T., Tozawa, R., & Aramaki, H. (2018). Effectiveness of using an unskilled model in action

observation combined with motor imagery training for early motor learning in elderly

people: A preliminary study. *Somatosensory & Motor Research*, *35*(3-4), 204-211.

https://doi.org/10.1080/08990220.2018.1527760

Kim, T., Frank, C., & Schack, T. (2017). A systematic investigation of the effect of action observation

training and motor imagery training on the development of mental representation structure

and skill performance. *Frontiers in Human Neuroscience*, *11*, 499.

https://doi.org/10.3389/fnhum.2017.00499

Kraeutner, S. N., McWhinney, S. R., Solomon, J. P., Dithurbide, L., & Boe, S. G. (2018). Experience

modulates motor imagery-based brain activity. *European Journal of Neuroscience*, *47*(10),

1221-1229. https://doi.org/10.1111/ejn.13900

Kraeutner, S. N., Stratas, A., McArthur, J. L., Helmick, C. A., Westwood, D. A., & Boe, S. G. (2020).

Neural and behavioral outcomes differ following equivalent bouts of motor imagery or

physical practice. *Journal of Cognitive Neuroscience*, *32*(8), 1590-1606.

https://doi.org/10.1162/jocn_a_01575

Krakauer, J. W., Hadjiosif, A. M., Xu, J., Wong, A. L., & Haith, A. M. (2019). Motor

Learning. *Comprehensive Physiology*, *9*(2), 613–663. https://doi.org/10.1002/cphy.c170043

Lakens, D. (2013). Calculating and reporting effect sizes to facilitate cumulative science: a practical

primer for t-tests and ANOVAs. *Frontiers in Psychology*, *4*, 863.

https://doi.org/10.3389/fpsyg.2013.00863

Lau, J., Ioannidis, J. P., Terrin, N., Schmid, C. H., & Olkin, I. (2006). The case of the misleading funnel

plot. *British Medical Journal*, *333*(7568), 597-600. https://doi.org/10.1136/bmj.333.7568.597

MacIntyre, T., Moran, A., Collet, C., & Guillot, A. (2013). An emerging paradigm: A strength-based

approach to exploring mental imagery. *Frontiers in Human Neuroscience*, *7,* 104.

https://doi.org/10.3389/fnhum.2013.00104

Marshall, B., & Wright, D. (2016). Layered stimulus response training versus combined action

observation and imagery: Effects on golf putting performance and imagery ability

characteristics. *Journal of Imagery Research in Sport and Physical Activity*, *11*(1), 35-46.

https://doi.org/10.1515/jirspa-2016-0007

Marshall, B., Wright, D., Holmes, P., & Wood, G. (2019). Combining action observation and motor

imagery improves eye-hand coordination during novel visuomotor task performance. *Journal

of Motor Behavior*, *52*(3), 333-341. https://doi.org/10.1080/00222895.2019.1626337

Marshall, B., Wright, D., Holmes, P., Williams, J., & Wood, G. (2020). Combined action observation

and motor imagery facilitates visuomotor adaptation in children with developmental

coordination disorder. *Research in Developmental Disabilities*, *98*, 103570.

https://doi.org/10.1016/j.ridd.2019.103570

Marusic, U., Grosprêtre, S., Paravlic, A., Kovač, S., Pišot, R., & Taube, W. (2018). Motor imagery

during action observation of locomotor tasks improves rehabilitation outcome in older

adults after total hip arthroplasty. *Neural Plasticity, 2018,* 1-9.

https://doi.org/10.1155/2018/5651391

McNeill, E., Ramsbottom, N., Toth, A. J., & Campbell, M. J. (2020). Kinaesthetic imagery ability

moderates the effect of an AO+ MI intervention on golf putt performance: A pilot

study. *Psychology of Sport and Exercise*, *46*, 101610.

https://doi.org/10.1016/j.psychsport.2019.101610

Meers, R., Nuttall, H., & Vogt, S. (2020). Motor imagery alone drives corticospinal excitability during

concurrent action observation and motor imagery. *Cortex*, *126*, 322-333.

https://doi.org/10.1016/j.cortex.2020.01.012

Mouthon, A., Ruffieux, J., Keller, M., & Taube, W. (2016). Age-related differences in corticospinal

excitability during observation and motor imagery of balance tasks. *Frontiers in Aging*

Neuroscience, *8,* 317. https://doi.org/10.3389/fnagi.2016.00317

Mouthon, A., Ruffieux, J., Wälchli, M., Keller, M. & Taube, W. (2015). Task-dependent changes of

corticospinal excitability during observation and motor imagery of balance tasks.

Neuroscience, 10(303), 535-543. https://doi.org/10.1016/j.neuroscience.2015.07.031

Naish, K., Houston-Price, C., Bremner, A., & Holmes, N. (2014). Effects of action observation on

corticospinal excitability: Muscle specificity, direction, and timing of the mirror

response. *Neuropsychologia*, *64*, 331-348.

https://doi.org/10.1016/j.neuropsychologia.2014.09.034

Nedelko, V., Hassa, T., Hamzei, F., Schoenfeld, M., & Dettmers, C. (2012). Action imagery combined

with action observation activates more corticomotor regions than action observation

alone. *Journal of Neurologic Physical Therapy*, *36*(4), 182-188.

https://doi.org/10.1097/npt.0b013e318272cad1

Neuman, B., & Gray, R. (2013). A direct comparison of the effects of imagery and action observation

on hitting performance. *Movement & Sport Sciences - Science & Motricité*, (79), 11-21.

https://doi.org/10.1051/sm/2012034

Pellegrini, M., Zoghi, M., & Jaberzadeh, S. (2018). The effect of transcranial magnetic stimulation test

intensity on the amplitude, variability and reliability of motor evoked potentials. *Brain*

Research, *1700*, 190-198. https://doi.org/10.1016/j.brainres.2018.09.002

Roberts, R., Callow, N., Hardy, L., Markland, D., & Bringer, J. (2008). Movement imagery ability:

Development and assessment of a revised version of the vividness of movement imagery

questionnaire. *Journal of Sport & Exercise Psychology, 30*(2), 200-221.

https://doi.org/10.1123/jsep.30.2.200

Romano Smith, S., Wood, G., Coyles, G., Roberts, J., & Wakefield, C. (2019). The effect of action

observation and motor imagery combinations on upper limb kinematics and EMG during

dart-throwing. *Scandinavian Journal of Medicine & Science in Sports*, *29*(12), 1917-1929.

https://doi.org/10.1111/sms.13534

Romano-Smith, S., Wood, G., Wright, D., & Wakefield, C. (2018). Simultaneous and alternate action

observation and motor imagery combinations improve aiming performance. *Psychology of*

Sport and Exercise, *38*, 100-106. https://doi.org/10.1016/j.psychsport.2018.06.003

Rothwell, J. C. (1997). Techniques and mechanisms of action of transcranial stimulation of the

human motor cortex. *Journal of Neuroscience Methods*, *74*(2), 113-122.

https://doi.org/10.1016/S0165-0270(97)02242-5

Ruffino, C., Papaxanthis, C., & Lebon, F. (2017). Neural plasticity during motor learning with motor

imagery practice: Review and perspectives. *Neuroscience*, *341*, 61-78.

https://doi.org/10.1016/j.neuroscience.2016.11.023

Scott, M. W., Wood, G., Holmes, P. S., Williams, J., Marshall, B., & Wright, D. J. (2021). Combined

action observation and motor imagery: An intervention to combat the neural and

behavioural deficits associated with developmental coordination disorder. *Neuroscience &*

Biobehavioral Reviews. *127*, 638-646. https://doi.org/10.1016/j.neubiorev.2021.05.015

Scott, M., Taylor, S., Chesterton, P., Vogt, S., & Eaves, D. (2018). Motor imagery during action

observation increases eccentric hamstring force: an acute non-physical

intervention. *Disability and Rehabilitation*, *40*(12), 1443-

1451. https://doi.org/10.1080/09638288.2017.1300333

Scott, M.W., Emerson, J.R., Dixon, J., Tayler, M. A. & Eaves, D. L. (2019). Motor imagery during action

observation enhances automatic imitation in children with and without developmental

coordination disorder. *Journal of Experimental Child Psychology, 183*, 242-260.

https://doi.org/10.1016/j.jecp.2019.03.001

Shimada, K., Onishi, T., Ogawa, Y., Yamauchi, J., & Kawada, S. (2019). Effects of motor imagery

combined with action observation training on the lateral specificity of muscle strength in

healthy subjects. *Biomedical Research, 40*(3), 107-113.

https://doi.org/10.2220/biomedres.40.107

Simonsmeier, B., Androniea, M., Buecker, S., & Frank, C. (2021). The effects of imagery interventions

in sports: A meta-analysis. *International Review of Sport and Exercise Psychology*, 1-22.

https://doi.org/10.1080/1750984x.2020.1780627

Spittle, M. (2021). *Motor learning and skill acquisition: Applications for physical education and sport*.

Bloomsbury Publishing.

Ste-Marie, D. M., Lelievre, N., & St. Germain, L. (2020). Revisiting the Applied Model for the Use of

Observation: A Review of Articles Spanning 2011–2018. *Research quarterly for exercise and*

sport, *91*(4), 594-617.

Ste-Marie, D., Law, B., Rymal, A., Jenny, O., Hall, C., & McCullagh, P. (2012). Observation

interventions for motor skill learning and performance: an applied model for the use of

observation. *International Review of Sport and Exercise Psychology*, *5*(2), 145-176.

https://doi.org/10.1080/1750984x.2012.665076

Stevens, J. A., & Stoykov, M. E. (2003). Using motor imagery in the rehabilitation of

hemiparesis. *Archives of Physical Medicine and Rehabilitation*, *84*(7), 1090–1092.

https://doi.org/10.1016/s0003-9993(03)00042-x

Tanner-Smith, E. E., & Tipton, E. (2014). Robust variance estimation with dependent effect sizes:

practical considerations including a software tutorial in Stata and spss. *Research Synthesis*

Methods, 5(1), 13–30. https://doi.org/10.1002/jrsm.1091

Tanner-Smith, E.E., Tipton, E. & Polanin, J.R. (2016). Handling complex meta-analytic data structures

using Robust Variance Estimates: a tutorial in R. *Journal of Developmental Life Course*

Criminology 2, 85–112. https://doi.org/10.1007/s40865-016-0026-5

Taube, W., Lorch, M., Zeiter, S. & Keller, M. (2014). Non-physical practice improves task performance

in an unstable, perturbed environment: Motor imagery and observational balance

training. *Frontiers in Human Neuroscience, 8,* 972.

https://doi.org/10.3389/fnhum.2014.00972

Taube, W., Mouthon, M., Leukel, C., Hoogewoud, H., Annoni, J., & Keller, M. (2015). Brain activity

during observation and motor imagery of different balance tasks: An fMRI study. *Cortex*, *64*,

102-114. https://doi.org/10.1016/j.cortex.2014.09.022

Toth, A., McNeill, E., Hayes, K., Moran, A., & Campbell, M. (2020). Does mental practice still enhance

performance? A 24 Year follow-up and meta-analytic replication and extension. *Psychology*

of Sport and Exercise, *48*, 101672. https://doi.org/10.1016/j.psychsport.2020.101672

Viechtbauer, W. (2010). Conducting meta-analyses in R with the metafor package. *Journal of*

Statistical Software, 36(3), 1-48. https://doi.org/10.18637/jss.v036.i03

Villiger, M., Estévez, N., Hepp-Reymond, M., Kiper, D., Kollias, S., Eng, K., & Hotz-Boendermaker, S.

(2013). Enhanced activation of motor execution networks using action observation

combined with imagination of lower limb movements. *PLoS One*, *8*(8), e72403.

https://doi.org/10.1371/journal.pone.0072403

Vogt, S., Rienzo, F., Collet, C., Collins, A., & Guillot, A. (2013). Multiple roles of motor imagery during

action observation. *Frontiers in Human Neuroscience*, *7,* 807.

https://doi.org/10.3389/fnhum.2013.00807

Williams, S. E., Cumming, J., Ntoumanis, N., Nordin-Bates, S. M., Ramsey, R., & Hall, C. (2012).

Further validation and development of the movement imagery questionnaire. *Journal of*

Sport & Exercise Psychology, *34*(5), 621–646. https://doi.org/10.1123/jsep.34.5.621

Wright D. J., Williams J., Holmes P. S. (2014) Combined action observation and imagery facilitates

corticospinal excitability. *Frontiers in Human Neuroscience,* 27(8), 951.

https://doi.org/10.3389/fnhum.2014.00951

Wright, D. J., Frank, C., & Bruton, A. M. (2021). Recommendations for combining action observation

and motor imagery interventions in sport. *Journal of Sport Psychology in Action,* 1-13.

https://doi.org/10.1080/21520704.2021.1971810

Wright, D.J., McCormick, S.A., Williams, J. & Holmes, P.S. (2016). Viewing instructions accompanying

action observation modulate corticospinal excitability. *Frontiers in Human Neuroscience*, 1,

10(17). https://doi.org/10.3389/fnhum.2016.00017

Wright, D., Wood, G., Eaves, D., Bruton, A., Frank, C., & Franklin, Z. (2018a). Corticospinal excitability

is facilitated by combined action observation and motor imagery of a basketball free

throw. *Psychology of Sport and Exercise, 39*, 114-121.

https://doi.org/10.1016/j.psychsport.2018.08.006

Wright, D. J., Wood, G., Franklin, Z. C., Marshall, B., Riach, M., & Holmes, P. S. (2018b). Directing

visual attention during action observation modulates corticospinal excitability. *Plos One,*

13(1), e0190165. https://doi.org/10.1371/journal.pone.0190165.

Yoxon, E., & Welsh, T. (2020). Motor system activation during motor imagery is positively related to

the magnitude of cortical plastic changes following motor imagery training. *Behavioural*

Brain Research, *390*, 112685. https://doi.org/10.1016/j.bbr.2020.112685

Zhang B, Kan L, Dong A, Zhang J, Bai Z, Xie Y, et al. (2019) The effects of action observation training

on improving upper limb motor functions in people with stroke: A systematic review and

meta-analysis. *PLoS One*, *14*(8), e0221166. https://doi.org/10.1371/journal.pone.0221166