Accurate inference of genome-wide spatial expression with iSpatial

Chao Zhang[1,2,3,] †, Renchao Chen[1,2,3,] †, and Yi Zhang[1,2,3,4,5,] ‡

[1]Howard Hughes Medical Institute, Boston Children's Hospital, Boston, Massachusetts 02115,

USA; [2]Program in Cellular and Molecular Medicine, Boston Children's Hospital, Boston,

Massachusetts 02115, USA; [3]Division of Hematology/Oncology, Department of Pediatrics,

Boston Children's Hospital, Boston, Massachusetts 02115, USA; [4]Department of Genetics,

Harvard Medical School, Boston, Massachusetts 02115, USA; [5]Harvard Stem Cell Institute,

WAB-149G, 200 Longwood Avenue, Boston, Massachusetts 02115, USA; †Co-first authors.

‡To whom correspondence should be addressed

14   E-mail: yzhang@genetics.med.harvard.edu

Keywords: Spatial transcriptomics, scRNA-seq, MERFISH, Slide-seq, STARmap, iSpatial

Manuscript information: 52 pages, 6 Figures, 11 Supplementary figures

HIGHLIGHTS

• iSpatial infers genome-wide spatial gene expression pattern by integrating spatial

transcriptomic and scRNA-seq data

• iSpatial outperforms existing approaches in inferring spatial gene expression patterns

• iSpatial reduces false-positive/negative signals of the original spatial transcriptome

• iSpatial is applicable to spatial transcriptomic datasets from different tissues and techniques

ABSTRACT

Spatially resolved transcriptomic analyses can reveal molecular insights underlying tissue

structure and context-dependent cell-cell or cell-environment interaction. Due to the current

technical limitation, obtaining genome-wide spatial transcriptome at single-cell resolution is

challenging. Here we developed a new algorithm named iSpatial to derive spatial pattern of the

entire transcriptome by integrating spatial transcriptomic and single-cell RNA-seq datasets.

Compared to other existing methods, iSpatial has higher accuracy in predicting gene expression

and their spatial distribution. Furthermore, it reduces false-positive and false-negative signals in

the original datasets. By testing iSpatial with multiple spatial transcriptomic datasets, we

demonstrate its wide applicability to datasets from different tissues and by different techniques.

Thus, we provide a computational approach to reveal spatial organization of the entire

transcriptome at single cell resolution. With numerous high-quality datasets available in the

public domain, iSpatial provides a unique way for understanding the structure, function of

complex tissues and disease processes.

INTRODUCTION

In the past decade, single-cell RNA-sequencing (scRNA-seq) has transformed our understanding on the cellular heterogeneity of various tissues/organs in multicellular organisms (*1-4*). With current scRNA-seq techniques, obtaining whole transcriptomic profiles of tens to hundreds of thousands of single cells has become a routine. However, most high-throughput scRNA-seq methods use dissociated cells and consequently the spatial information of the analyzed cells is lost, which prevents directly connecting the molecular feature of the analyzed cell types to their anatomic and functional features. On the other hand, the development of spatially resolved transcriptomic assays have enabled the transcripts/cells location analysis in the tissue context, which has the potential to reveal how single-cell gene activity orchestrates the structure and function of complex tissues in health and disease (*5*).

In the past few years, different methods for spatial transcriptomic (ST) assays have been developed (*6-13*). Ideally, the spatial transcriptome data should provide genome-wide and spatially resolved expression measurements at single-cell resolution. However, due to technical limitations, either spatial resolution or gene coverage is compromised in most ST assays. For example, *in situ* capture and sequencing-based techniques are able to capture any mRNA molecules without pre-knowledge, but the spatial resolution is not at single-cell level (*6, 14*). On the other hand, *in situ* sequencing and FISH-based mRNA measurement can achieve cellular or subcellular resolution, but most such assays are limited with their throughput on genes that can be detected (usually 30-500), and require pre-knowledge for probe design (*7-9*).

With the rapid development of scRNA-seq and ST technologies, new bioinformatic tools have been developed to overcome the challenges in single-cell or spatial transcriptomic data analysis (*15-20*). Several imputation methods for scRNA-seq data have emerged, including MAGIC (*21*), scImpute (*22*), DrImpute (*23*), ALRA (*24*). However, methods developed for inferring ST data are still limit. Notably, by integrating scRNA-seq and spatial resolved profiling data, recent computational methods have leveraged the strength of different datasets and revealed information that otherwise cannot be obtained from a single experimental paradigm. For example, when the corresponding scRNA-seq is available, SpatialDWLS and RCTD could

perform deconvolution on the "low-resolution" spatial dataset to estimate the cell

type/proportion in each spatially resolved spot (*25, 26*). On the other hand, Tangram and

Cell2location could predict the spatial location of molecularly defined cell types (from scRNA-

seq) based on ST data (*27, 28*). Seurat and Liger could impute transcriptome-wide spatial

expression by integrating with corresponding scRNA-seq data and transferring the expression

data to spatial transcriptome (*29, 30*). Despite these existing tools, inferring expression patterns

of all genes at high spatial resolution by integrating scRNA-seq and ST data is not

straightforward, and different approaches have variable performance when applied to different

datasets. Since the performance of this task is critical for downstream spatial analysis, a robust

and convenient tool for spatial pattern prediction is highly desirable.

Here, we present iSpatial, an R-based bioinformatic tool that integrates scRNA-seq and ST

profiling data to infer the expression pattern of each gene at high spatial resolution. We show

that iSpatial outperforms existing approaches on its accuracy and it can also reduce false-positive

and false-negative signals in the original data. By applying iSpatial to datasets from different

tissues (hippocampus, hemibrain, cortex, striatum, liver) and generated with different techniques

(Slide-seq, Stereo-seq, MERFISH, STARmap), we have revealed both known and novel spatial

expression patterns in each dataset, indicating iSpatial is broadly applicable for analyzing

different ST datasets. Collectively, our analyses demonstrate that iSpatial is a useful tool for

resolving transcriptome-wide spatial expression patterns at single cell resolution in complex

tissues.

RESULTS

Overview of iSpatial

The FISH and *in situ* sequencing-based ST techniques, such as MERFISH (*7*), seqFISH (*8*),

osmFISH (*12*) and STARmap (*9*), can simultaneously reveal gene expression and location at

single-cell resolution, but with limited pre-defined gene targets (**Fig. 1A**, left). On the other

hand, scRNA-seq can unbiasedly profile the whole transcriptome, but without providing spatial

information (**Fig. 1A**, middle). We reasoned that by integrating the single-cell gene expression

profiles (the gene by cell matrices) of the two methods, the missing information of non-targeted

genes in each spatially profiled cells could be inferred based on scRNA-seq data, resulting in

genome-wide spatial expression information of the profiled cells (**Fig. 1A**, right).

To this end, we first performed dimension reduction on scRNA-seq and ST data separately,

followed by expression stabilization, which removes potential noise/background expression

based on the expression level of adjacent cells in PCA space. The two datasets were then

normalized and embedded into a common space with two sequential rounds of integration: first

by reciprocal PCA (*29*) and then through Harmony (*31*). Based on the common embedding, the

expression value of each gene in each cell of the ST dataset is then inferred using a weighted k-

nearest neighbors (KNN) model. Since the physical locations of these cells have been resolved in

spatial profiling, the results represent a new single-cell gene expression profile with both

genome-wide coverage and single-cell spatial resolution, which could be used for downstream

analyses including identification of spatially variable genes (SVGs) (**Fig. 1B**).

iSpatial outperforms existing tools in its accuracy on predicting spatial expression pattern

To evaluate the performance of iSpatial and compare it with existing tools, we used a mouse

hippocampal dataset generated from Slide-seq V2 (*32*). Since this dataset includes the spatial

expression of all genes, it can be used for evaluating the prediction performance (**Fig. 2A**).

Specifically, we divided the dataset into the training and validation groups, which contain 3,000

and ~20,000 genes, respectively. The training data group (mimic a ST dataset) was integrated

with a scRNA-seq dataset covering the same brain region (hippocampus) (*1, 33*) (but by a

different method) to infer the expression level and spatial patterns of the genes in the validation

data group. By comparing the inferred expression patterns with the "truth" determined by Slide-

seq (validation data group), we found that iSpatial could predict the spatial expression pattern

with high accuracy. For example, iSpatial inferred the expression of *Atp2b1*, *Prox1* and *Fibcd1*,

which were not included in the training data group, across the entire hippocampus, dentate gyrus

(DG), and CA1, respectively, consistent with the Slide-seq validation data and ISH results from

Allen Brain Atlas (ABA) (*33*) (**Fig. 2B**). Interestingly, we found that iSpatial could "enhance"

the signals not well detected in the original data. For example, *Slit1*, *Tspan18*, *Efnb2*, *Car12* and

others were barely detectable in hippocampal cells by Slide-seq, thus difficult to determine their

spatial pattern. With iSpatial, the expression of these genes was clearly visible, thus their spatial

pattern could be clearly recognized. This is unlikely an artifact of imputation, as the spatial

expression inferred by iSpatial is consistent with that of the ABA data (**Fig. 2C and fig. S1A**).

We further compared the performance of iSpatial with another two existing tools, Liger (*30*) and

Seurat (*29*), on the same task using the Slide-seq dataset. Although these two methods could also

infer the expression patterns of genes not included in the training data group, compared with

iSpatial, the spatial patterns obtained from Liger and Seurat were more ambiguous with higher

background in general (**Fig. 2B, 2C and fig. S1A**). To quantitatively benchmark these different

methods, we calculated the expression and spatial correlation between Slide-seq data (regarded

as ground truth) and inferred results from iSpatial, Liger or Seurat on each gene of the validation

dataset. The results showed iSpatial exhibited significantly higher accuracy than the other

methods across all gene groups with different expression levels, and the accuracy of prediction is

positively correlated with the gene expression level (**Fig. 2D and 2E**). In addition, cell type

specifically expressed genes exhibit higher prediction accuracy (**fig. S1B**). This result suggests

that iSpatial achieves higher prediction accuracy on functionally relevant genes. Furthermore,

when compared the SVGs identified from the original Slide-seq data with those identified from

inferred data of different methods (**fig. S1C**), we found that iSpatial has the highest prediction

power with area under the curve (AUC) over 0.84 on SVGs among the three methods (**Fig. 2F**).

We also used Stereo-seq data of an adult mouse coronal hemibrain section (*34*) to benchmark the

performance of our method. Similar to Slide-seq V2, we randomly sampled 3,000 genes as

training dataset, and other genes as validation dataset (**fig. S2A**). After integrating with a single

cell dataset of corresponding brain regions (*2*), we compared the performance of Liger, Seurat

and iSpatial on predicting the gene expression levels and patterns. The results showed that

iSpatial achieved higher correlation than other methods on validation datasets (**fig. S2B-E**).

Collectively, these analyses indicate that iSpatial outperforms existing tools in terms of accuracy

on predicting spatial expression pattern.

iSpatial is broadly applicable to different spatial transcriptomic datasets

After validating the performance of iSpatial with Slide-seq and Stereo-seq data, we further tested

whether iSpatial can be applied to other ST datasets generated from different tissues and with

different techniques. To this end, we first used iSpatial to analyze a STARmap dataset which

covered the primary visual cortex (V1) of mouse brain (*9*) (**Fig. 3A**). Although the original

STARmap data only included 1,020 gene targets, iSpatial successfully inferred the expression of

over 20,000 genes by integrating a single cell smart-seq dataset from ABA (*35*) (**Fig. 3B**).

Importantly, the spatial expression patterns of genes not included in the original STARmap data

could be faithfully inferred by iSpatial. For example, the layer-specific expression of a number of

genes were accurately predicted as evidenced by its similarity to that of the ABA ISH results

(**Fig. 3C**). Notably, iSpatial not only correctly predicted the layer distribution of non-targeted

genes, but also detected the expression variation of certain genes based on their spatial locations.

For example, it predicted: 1) a high-to-low gradient of *Pvrl3* across upper cortical layers, and 2)

strong expression of *Serinc2* and *Col5a1* in cortical layer VI but relatively weak expression in

upper layer V, both of which were confirmed by ISH data (**Fig. 3C**).

In additional to the STARmap dataset, we analyzed a recently published MERFISH dataset of

mouse striatum (*36*) with iSpatial (**Fig. 3D**). The original MERFISH dataset contained 253 target

genes which allowed the identification of 9 major cell types in striatum (**fig. S3A**), with 175

target genes exhibit significant enrichment in certain cell types (**fig. S3B**). By integrating this

dataset with corresponding scRNA-seq data, iSpatial could infer the expression and location of

~9,000 genes at single-cell resolution (**Fig. 3E and fig. S3C**), with over 2,200 genes identified as

cell type specific expressed genes (**fig. S3D and S3E**). Importantly, the spatial patterns of

inferred genes were largely consistent with those determined by ISH. For example, *Gpr37* was

highly enriched in the anterior commissure (AC), *Coch* formed a high-to-low gradient from the

dorsolateral to the ventromedial striatum, and *Stard5* was specifically expressed in the medial

nucleus accumbens (NAc), all these iSpatial inferred expression patterns are consistent with the

results from ABA (**Fig. 3F**).

To globally evaluate the predication accuracy, we adopted a 10-fold cross validation approach

and found that iSpatial showed significantly higher accuracy than Liger and Seurat in predicting

the expression and spatial location of genes in both datasets (**fig. S4**). Collectively, these results

demonstrated the capacity of iSpatial in predicting the expression and location of genes using ST

dataset generated from different tissues with different techniques.

iSpatial reduces false-positive and false-negative signals from spatial transcriptome

Although imaging-based transcriptomic assays have a higher detection efficiency when

compared to that of sequencing-based methods, their performance is highly variable depending

on the specific gene probes. For example, some transcripts are too short to be targeted by enough

probes, which may lead to false-negative (dropout). On the other hand, some other genes may

have close homologs which are difficult to distinguish with hybridization, leading to false-

positive (background). We hypothesized that iSpatial could reduce these false signals by giving

higher weights to cells of scRNA-seq when performing expression prediction, which were

insensitive to gene length and could also unambiguously distinguish similar transcripts based on

sequence differences. To test this hypothesis, we first compared the expression pattern of some

well-establish cell type markers on the UMAP between the original STARmap data and iSpatial

inferred data. We found that although these cell type-specific markers exhibited high enrichment

in corresponding cell types, there were often false-positive signal in other cell types when

analyzed by STARmap (**Fig. 4A**, upper panels, *Slc17a7*, *Gad1*, *Plp1* and *Cldn5*). In some cases,

the expected expression patterns were not observed, likely due to false-negative (**Fig. 4A**, upper

panels, *Aqp4*). Consistent with our hypothesis, iSpatial could remove most false-positive signals

from irrelevant cell types, without affecting the true positive signals (**Fig. 4A**, lower panels,

Slc17a7, *Gad1*, *Plp1* and *Cldn5*). Furthermore, iSpatial successfully inferred gene expression

pattern that was missed in the original STARmap analysis (**Fig. 4A**, lower panels, *Aqp4*),

suggesting that iSpatial can also reduce false-negative results. Based on these findings, we

further asked whether iSpatial could infer spatial patterns which were not well detected by

STARmap. Indeed, we found a number of genes, whose expected layer-specific expression

patterns were not detected in the original STARmap analysis. For example, *Nov*, *Rorb*, *Rspo1*,

Fezf2, *Foxp2* are established markers of different cortical layers. However, STARmap only

detected sparse signals or even failed to detect real signals of these genes across all cortical

layers (**Fig. 4B**). In contrast, iSpatial accurately captured the layer-specific expression patterns of

these genes that were also detected by ISH (**Fig. 4B**).

In addition to the STARmap cortical dataset, a similar effect of iSpatial in correcting false-

positive and false-negative expression on the MERFISH striatum data is also observed.

Specifically, iSpatial removed most false-positive noise of known cell type-specific markers (**fig.**

S5A and B). It also accurately predicted the expression and spatial pattern of genes not well

detected by MERFISH, such as *Tac2*, *Serpinb2* and *Kctd4* (**Fig. 4C and fig. S5C**). Compared to

the original MERFISH data, the spatial patterns inferred by iSpatial showed higher consistency

with those determined by ISH (**Fig. 4C and fig. S5C**). For example, MERFISH indicates a broad

distribution of *Serpinb2* across the striatum, but iSpatial suggested it was selectively expressed in

a small group of cells located in the medial shell of NAc (**Fig. 4C**). ISH from ABA confirmed

the accuracy of iSpatial's prediction (**Fig. 4C**), suggesting that iSpatial is capable of reducing

noise. Collectively, these results showed that iSpatial can reduce false-positive and false-

negative signals in the original ST data from different tissues generated by different techniques.

iSpatial enables whole-transcriptome level spatial analysis

One major goal of ST analysis is identifying SVGs, which are the molecular basis of

structural/functional heterogeneity in different tissues. Since iSpatial could reliably infer

genome-wide gene expression and their spatial locations, we sought to test whether iSpatial

could augment the capability of certain ST dataset in detecting SVGs and spatial gene expression

patterns. To this end, we applied iSpatial to the STARmap cortex dataset to identify SVGs. We

found that iSpatial inferred data dramatically increased the number of detected SVGs (from 21 to

2,122, **fig. S6A**). Clustering analysis of the SVGs revealed six major spatial patterns (**fig. S6B**),

which resemble the known layer organization of mouse cortex. Notably, even when we restricted

the analysis to the target genes of STARmap, iSpatial still identified more SVGs (162 in inferred

data and 21 in original data), likely due to the correction of false-positive and false-negative

signals in the original data (see above).

In addition to STARmap, we performed parallel analysis to evaluate iSpatial's effect on SVG

identification using the MERFISH striatum dataset (**Fig. 5A**) and observed a similar increase in

the SVGs number and statistic power of spatially variable test. Specifically, the SVG number

increased by > 20 folds (from 94 in the original data to 1,968 in the inferred data) (**fig. S6C**).

Compared to cerebral cortex, the anatomic organization of striatum is more ambiguous and less

well understood, although recent studies have suggested that distinct transcriptional features and

cell types underlying its anatomic heterogeneity (*36, 37*). By unbiased clustering analysis of the

SVGs obtained from the iSpatial inferred data, we identified 12 distinct spatial patterns of SVGs

(**Fig. 5B**). Importantly, many of these patterns closely resemble the known anatomic subregions

in the striatum. For example, the C12 cluster is mainly expressed in the dorsal striatum, while C1

and C4 clusters are highly enriched in the NAc (**Fig. 5B and 5C**). Additionally, C7 and C11

clusters correspond to the core region of the NAc, while the C2, C9 and C10 clusters represent

the shell region (*38*) (**Fig. 5B and 5C**). Interestingly, the C8 cluster is specifically expressed in

the medial shell (**Fig. 5B, 5C and fig. S6D**), a NAc subregion known to possess distinct

anatomic and functional features (*39-41*). These results indicate that iSpatial can facilitate

identification of biologically relevant spatial gene expression patterns.

iSpatial improves analysis of spatial transcriptomic data from liver

Having demonstrated the utility of iSpatial in the analysis of ST data from different brain

regions, we next sought to test iSpatial's performance with ST data from other tissues. To this

end, we analyzed a Vizgen MERFISH Mouse Liver Map dataset with 347 target genes included

in the original data (https://vizgen.com/data-release-program/). By integrating the MERFISH

data with a liver scRNA-seq dataset (*42*), iSpatial successfully inferred the expression of over

6,000 genes on average in each single cell (**Fig. 6A and 6B**), which increased by > 20 folds from

the original data. Importantly, the inferred spatial patterns were largely consistent with

established knowledge. For example, iSpatial predicted selective expression of *Slc1a2* and

Aldh1b1 in cells around the central vein (CV) and portal vein (PV), respectively (**Fig. 6C**).

Similarly, *Cyp2e1* and *Cyp2f2* were predicted to be biased to CV and PV, but they have broader

distribution than *Slc1a2* and *Aldh1b1* (**Fig. 6C**). All these spatial patterns were confirmed in

previous studies (*43, 44*). We further generated UMAP based on iSpatial inferred expression

profile and found a close correlation between cells' positions on UMAP and their *in-situ*

distribution along the CV-PV axis (**Fig. 6C and 6D**), revealing a gradient expression profile

along the CV-PV axis. Notably, although Liger and Seurat can also reveal a similar gradient

expression patten, a comparison among the three methods indicated that iSpatial achieved a

higher specificity and accuracy, especially on genes with more spatially restricted expression

patterns. For example, *Slc1a2* is selectively expressed in a monolayer surrounding the CV, which

was accurately predicted by iSpatial (**Fig. 6C**), while Liger and Seurat revealed a more broader

expression pattern (**fig. S7A and S7B**). Based on the observed relationship between gene

expression and spatial location of liver cells, we calculated a CV score for each cell (see

Methods) to reflect its relative position to CV/PV, with a high/low CV score indicating close to

CV/PV, respectively. As expected, the CV score showed gradual increase from PV to CV in both

the liver tissue and the UMAP space (**Fig. 6E and 6F**). Then all the genes in the inferred dataset

(from iSpatial) were ranked according to their correlation with the CV score, which enabled us to

systematically identify genes strongly related to cell's spatial distribution (**Fig. 6G**). From this

analysis, 141 and 692 genes with correlation to CV score > 0.3 or < -0.3 were predicted to be

strongly enriched in cells close to CV or PV. In contrast, only 3 and 17 of these spatially variable

genes were included in the original MERFISH dataset. As expected, many genes known to be

biased to CV or PV were found, including *Gulo* and *Cyp2a5* enriched in cells adjacent to CV,

Cdh1 and *Etnppl* mainly expressed in cells around PV (**Fig. 6H**). Furthermore, by applying

KEGG pathway analysis to CV and PV enriched genes, we found that they are involved in

different biological functions (**Fig. 6I**). For example, the genes highly expressed around CV were

enriched in "drug metabolism" and "PPAR signaling pathway", while the genes highly expressed

in PV were enriched in "protein processing in ER", and "complement and coagulation cascades"

(**Fig. 6I and fig. S7C**). These findings were consistent with previous reports (*43, 44*). Taken

together, the above analyses demonstrate that iSpatial can overcome the limited target gene

numbers of various ST analyses to the whole-transcriptome level with high accuracy in different

tissues.

DISCUSSION

Spatial transcriptomic assays simultaneously profile gene expression and their spatial location in tissue context, with the potential to unveil transcriptional features associated with tissue organization, cell-cell interaction, and region-specific physiological/pathological changes (*45*). Although sequencing and imaging-based ST techniques have been rapidly evolving (*5*), obtaining a genome-wide expression profile with single-cell spatial resolution is still challenging. To overcome this limitation, we developed a computational tool iSpatial to infer the genome-wide spatially resolved transcriptional information. iSpatial is especially useful for imaging-based ST analysis (such as MERFISH, seqFISH, STARmap), which in general has high detection efficiency and single-cell/subcellular spatial resolution, but is usually limited by the pre-defined gene targets. By integrating such kind of ST data with corresponding scRNA-seq profiles, the expression levels of untargeted genes could be inferred from scRNA-seq data, while the spatial information is directly inherited from the ST data, enabling high-resolution spatial analysis at the whole-transcriptome scale.

To ensure accurate expression imputation, it is critical to account for intrinsic noise in different original datasets. Specifically, high-throughput scRNA-seq has low capture efficiency on mRNA molecules, leading to a large proportion of zero counts for expressed genes (dropout). On the other hand, due to the variable performance of different gene probes, imaging-based ST analysis may generate both false-positive and false-negative signals. iSpatial includes an expression stabilization step, which borrows the information from cells with similar global expression pattern to minimize random noise in the original data (**fig. S8A and B**). The proper alignment of scRNA and ST is critical for this method. Harmony integration bias to keeping the global heterogeneity instead of local heterogeneity (**fig. S8C and D**). iSpatial uses two-rounds integration to reduce potential technology bias and batch effect on PCA space, allowing accurate integration of ST and scRNA-seq datasets (**fig. S8C and D**). A comparative analysis indicated that two-round integration resulted in more accurate prediction than one-round integration (**fig. S9**). As a result, iSpatial exhibits a significantly higher accuracy in both benchmark analysis with Slide-seq data and cross-validation with image-based ST data when compared with other existing imputation methods. Notably, iSpatial is not only able to faithfully predict the spatial expression

of genes out of the original ST data, but also shows robust performance on reducing the false-

positive/negative signals in the raw data. In both cortical STARmap and striatal MERFISH

datasets, iSpatial is able to remove random expression of cell type-specific markers in non-

relevant cell types and correct some non-specific background in the raw data (probably caused

by poor performance of certain probes) to generate spatial expression patterns that are consistent

with established knowledge. Since it is difficult to directly validate the detected mRNA

sequences that generate the signals in imaging-based ST techniques, iSpatial provides a useful

method to evaluate the performance of different probes.

iSpatial is based on a KNN approach, where the setting of K value is important. Theoretically,

large K value will dilute the signals for rare cell types, while small K value can increase

specificity but also reduce coverage. In iSpatial, we used weighted KNN when performing

expression inference: the neighbors close to the inquired cell will be assigned higher weights

than neighbors far from the cell in expression imputation. This should reduce the over-smoothing

effect for rare cell types when relatively large K is used, as the neighbors relatively far away

from the rare cell types will have less impact on the inferred expression (**fig. S10**). Nevertheless,

the optimal K value could vary for different datasets, thus we implemented a function

"recommend_k" in the iSpatial package to help the user determine the optimal K value.

In the real situation, the scRNA-seq and ST dataset would not always match exactly. To exam

the performance of iSpatial under such situation, we compared Liger, Seurat and iSpatial with

unmatched scRNA-seq and ST datasets. Specifically, we used a MERFISH data covering a

whole coronal mouse brain section (https://vizgen.com/data-release-program/) and a scRNA-seq

dataset (*46*) generated from mouse prefrontal cortex (**fig. S11A**). The results indicate that

iSpatial can specifically infer the expression of cells in ST by inputting the corresponding

scRNA-seq data (**fig. S11B**). *Cux1* and *Tle4* are established markers for cortical layer 2/3 and

layer 6, respectively. iSpatial accurately inferred the expression patterns (**fig. S11C and D**),

demonstrating iSpatial can accurately impute the gene expression pattern in relevant cell types

even when unmatched ST and scRNA-seq datasets were used.

We have tested iSpatial with multiple ST datasets generated from different tissues and

techniques. In all the applications, iSpatial is able to expand spatial information from a pre-

defined gene panel in the original ST data to the whole transcriptome, which renders several

benefits for downstream analysis: First, it enables systematic identification of SVGs. In both the

brain and liver datasets, we found that the number of SVGs increased from a few hundreds to

several thousand after iSpatial imputation. Second, iSpatial enables the discovery of distinct

spatial expression patterns across the tissue, which is achieved by spatial clustering analysis of

SVGs. Importantly, many such unbiasedly identified expression patterns are biologically

relevant. For example, we found that the SVG groups are organized into layer structure in the

cortex and they exhibit core/shell enrichment in the striatum, indicating a tight relationship

between gene expression and tissue organization. Third, iSpatial enables bioinformatic analysis

requiring sufficient gene number or high statistical power, such as KEGG analysis, to be

performed on SVGs or SVG subgroups. For example, by inferring the transcriptome-wide spatial

pattern in liver, we found genes enriched in CV and PV are involved in distinct KEGG terms,

suggesting a likely link between region-specific gene expression and function.

A potential limitation of iSpatial is that it requires corresponding ST and scRNA-seq data, which

may not be always available. But given the rapid development of ST and scRNA-seq techniques,

and ongoing effort in large single-cell consortium and spatial transcriptomics (*4, 47, 48*), we

anticipate that iSpatial will be widely used to help understanding the molecular basis of structural

and functional heterogeneity in complex tissues of diverse organs in normal and disease states.

To facilitate iSpatial implementation, we have made the iSpatial R package available at

https://github.com/YiZhang-lab/iSpatial, where a tutorial on how to use iSpatial to integrate

striatal scRNA-seq and MERFISH data to infer genome-wide spatial expression patterns can be

found.

MATERIALS AND METHODS

iSpatial workflow

The workflow of iSpatial R-package contains the following steps: 1) Expression stabilization,

which removes possible false positive and false negative gene expression signals at single cell

level; 2) Integration. Two rounds of integration are performed to achieve accurate mapping of

single cell RNA-seq data and spatial transcriptome data; 3) Infer expression according to the

weighted k-nearest neighbors (KNN); 4) Downstream analysis. Detection of spatial variable

genes and patterns.

Expression stabilization

In the probe-based spatial transcriptome, we observed some cell type markers are not fully

detected in corresponding cell cluster. However, these markers are indeed expressed across the

cluster based on the scRNA-seq data. This indicates some probes have low binding affinities,

which generates false negative (FN) signals. We also found some markers could be detected in

some cell types which should not be expressed (random distributed in whole slice). This can be

caused by nonspecific binding of some probes, which cause false positive (FP) signals. iSpatial

try to remove these FN/FP signals to correct the expression in each cell. We assume that these

FN/FP signals are randomly distributed. Thus, it can be corrected using the cells show similar

global expression patterns, but don't exhibit FN/FP on the same gene. To remove these FN and

FP signals in each single cell i, we first find the K-nearest neighbors ($KNN_{i,k}$) which most

correlate with the cell i based on the global expression pattern.

441   E_i: the vector of gene expression in cell i

442            $\hat{E}_i$: the vector of corrected gene expression in cell i

443            $KNN_{i,k}$: $\{KNN_{i,1}, KNN_{i,2}, \dots, KNN_{i,k}\}$: the set of k-nearest neighbor of cell i

444   Then, we correct the gene expression $\hat{E}_i$ by the expression of KNNs:

445            $$\hat{E}_i = \alpha E_i + (1 - \alpha) \sum_k E_{KNN_{i,k}}$$

where α controls the weight of expression from cell i and neighbors. By default, we set $\alpha = 0.5$.

Higher value strongly corrects the noise by the KNNs, but loss cell specificity, which may cause

lower detection power of some rarely expressed genes.

Integration of spatial transcriptomic data and scRNA-seq data

To achieve accurate integration of spatial transcriptomic data and scRNA-seq data, iSpatial uses

a two-round integration approach. In the first round, we adopted a reciprocal principal

component analysis (RPCA) method from Seurat. At this step, the PCA spaces are calculated in

both datasets. Then, one dataset is projected onto the other's PCA space and constrain the cell

anchors with the same mutual neighbors. We use the "FindIntegrationAnchors" function with the

parameter reduction = "rpca" in Seurat (version 4.0.5) to find the anchors and use

"IntegrateData" to get the integrated data. For the second-round integration, we project the cells

from both spatial transcriptome and scRNA-seq into a shared PCA embedding. Then an iterative

clustering method is used to remove the technology bias and batch effect on PCA space, which

get harmonious PCA embeddings for both datasets. Harmony (version 0.1.0) is then used to

generate normalized PCA embeddings in this step.

Infer spatial expression

After integrating two datasets into one reduced dimension space, iSpatial uses a weighted KNN

approach to infer the expression of nontargeted genes. For each cell t in spatial transcriptome

data, iSpatial searches the K-nearest neighbors ($KNN_{t,k}$).

t: cell in spatial transcriptome data

c: cell in scRNA-seq data

$KNN_{t,k}$: $\{KNN_{t,1}, KNN_{t,2}, \dots, KNN_{t,k}\}$: the set of k-nearest neighbor of cell t

Then, $KNN_{t,k}$ are restricted to the cells from scRNA-seq data. Because the cells from scRNA-

seq data contain the expression information of the whole transcriptome.

$KNN'_{t,k}$: $KNN_{t,k} \in c$

The final inferred expression $\widehat{E}_t$ of cell t is calculated by the expression of cell t itself and the

expression of $KNN'_{t,k}$:

$$\widehat{E}_t = (1 - \beta)E_t + \beta\left(\sum_k \omega_{t,k} E_{KNN'_{t,k}}\right)$$

where ω is the weight of each neighbor k of cell t. For the genes targeted in cell t, β balance the

expression from spatial transcriptome and scRNA-seq. $\beta \in [0,1]$. For genes not measured in ST

data, the $\beta = 1$. The weights ω between cell t and its neighbor KNN'_t are defined by the

normalized transcriptional distance $d_{t,k}$. Here, iSpatial use 1 - Pearson's correlation coefficient

to measure the distance:

$$d_{t,k} = dist\left(E_t, E_{KNN'_{t,k}}\right)$$

$$\omega_{t,k} = \frac{d_{t,k}^2}{\sum_k d_{t,k}^2}$$

And,

$$dist\left(E_t, E_{KNN'_{t,k}}\right) = 1 - cor\left(E_t, E_{KNN'_{t,k}}\right)$$

$$= \frac{\sum(E_t - \bar{E}_t)(E_{KNN'_{t,k}} - \bar{E}_{KNN'_{t,k}})}{\sqrt{\sum(E_t - \bar{E}_t)^2}\sqrt{\sum(E_{KNN'_{t,k}} - \bar{E}_{KNN'_{t,k}})^2}}$$

489   ***Identifying spatially variable genes***

To identify significant spatially variable genes, the x and y axes of the spatial location are evenly

divided into n bins, then the spatial location is further grided into $n \times n$ grids. For each gene j,

we calculate the average expression values E_j over the n^2 grids. We then randomly sample the

spatial location of each cell and calculate the average expression E'_j over the randomly sampled

n^2 grids. If a gene j has no specific spatial expression pattern, the distribution of observed E_j

should not be different from that of random E'_j. On the contrary, if a gene has strong spatial

expression pattern, the distribution E_j should be significantly different from E'_j. Thus, whether a

gene exhibits spatial expression pattern depends on whether there is a difference between the

distribution of E_j and E'_j. Here, we apply a non-parametric two-sides Mann–Whitney U test to

determine the difference between E_j and E'_j. We also offer the Kolmogorov-Smirnov test to test

the distribution differences.

Some studies found that a gene with spatial expression pattern always display a specific

expression bias on scRNA-seq UMAP/tSNE projection. iSpatial also integrates scRNA-seq

information into spatial variable gene detection. If a gene not only displays spatial expression

pattern on spatial transcriptome data but also exhibits expression specificity on scRNA-seq

UMAP/tSNE projection, this gene has a higher confidence of spatial expression pattern. Similar

to the detection of spatial expression gene on spatial location, iSpatial uses the same method to

detect whether a gene displays a specific expression location on scRNA-seq UMAP/tSNE. To

integrate these two lines of information, the final p value is equal to the p value from the spatial

transcriptomic data multiplied by the p value from the scRNA-seq data. Then the adjusted p

values are calculated to control the false discovery for multiple comparisons.

Spatially variable gene clustering

The genes with spatial expression patterns could be grouped into clusters. For each gene, iSpatial

captures the spatial expression features according to the average expression value over the n^2

grids, which was described before. Based on these features, Pearson's correlation coefficients are

calculated for pairs of genes. Then, distances among genes are measured by 1 minus Pearson's

correlation coefficients. Finally, Hierarchical clustering is performed using "hclust" in R to

spatial variable genes. "cutree" function from R is used to group these genes into desired number

of groups.

Single-cell RNA-seq data processing

The initial gene×cell matrix for each study was downloaded according to the original papers. The

expression matrix was then transferred into Seurat object for downstream analysis. The raw

counts of gene expression profile of each cell were normalized to 10,000 counts and natural log

transformed using the Seurat function "NormalizeData". To generate UMAP, we used standard

pipeline from Seuart. In short, "FindVariableFeatures" was used to identify top variable genes,

"ScaleData" was used to scale and center these genes in the data, then PCA was performed by

"RunPCA" based on the selected features. Finally, the top 30 principal components from PCA

were used to generate the UMAP projection by "RunUMAP" with the parameters "dims=1:30".

Mouse hippocampus data processing

The Slide-seq V2 of mouse hippocampus was download from the Broad Institute single cell

portal website (https://singlecell.broadinstitute.org/single_cell/study/SCP815). Here, we only use

the "Puck_200115_08" dataset (*32*). This dataset contains two files, the raw expression matrix

and the barcode locations. The data processing of Slide-seq V2 followed the same procedure as

that of single cell RNA-seq. The main difference is that Slide-seq contains spatial location of

each cell. According to the vignettes of Seurat, the coordinate of each cell is stored as a

"SlideSeq" class in Seurat object. For the scRNA-seq data of mouse hippocampus, we used a

published dataset (*1*). To facilitate analysis, we used a pre-processed Seurat object

(https://www.dropbox.com/s/cs6pii5my4p3ke3/mouse_hippocampus_reference.rds?dl=0)

offered by the Satija Lab.

Mouse hemibrain data processing

The single cell resolution Stereo-seq data of mouse hemibrains is download from

(https://db.cngb.org/stomics/mosta/) (*34*). We first transfer single cell level expression matrix to

Seurat. And the spatial locations of each cell are inputted into Seurat object. Then, the cells with

expressed genes over 500 are kept. The expression was normalized by NormalizeData. The

corresponding scRNAseq data was download from Linnarsson lab

(http://mousebrain.org/adolescent/downloads.html) (*2*). We only used cells from central nervous

system, and removed cells not in the brain regions profiled by Stereo-seq.

Mouse cortex data processing

The STARmap data of mouse visual cortex was download from the STARmap resource website

(https://www.starmapresources.org/data). We chose the dataset "20180505_BY3_1kgenes"

which profiles 1020 genes. The expression matrix data "cell_barcode_count.csv" was imported

to Seurat object. The spatial location coordinate of each cell was extracted from "labels.npz"

according to the method provide by the original paper (https://github.com/weallen/STARmap).

The cell coordinates were integrated into Seurat object. The single cell RNA-seq was download

from the Allen Brain Atlas (https://portal.brain-map.org/atlases-and-data/rnaseq/mouse-v1-and-

alm-smart-seq). The "mouse_VISp_2018-06-14_exon-matrix.csv" file was used to generate the

expression profile. Low-quality cells were removed according to the meta data

"mouse_VISp_2018-06-14_samples-columns.csv".

Mouse striatum data processing

The mouse striatum merFISH data was processed and normalized as described in the original

paper (*36*). Briefly, the expression of each cell was normalized by cell size and total RNA

counts.   Then, the log-transformed data was applied to the expression matrix. In this manuscript,

we used the data from a representative slice (slice 10). We used the pre-process Seurat object

data in https://www.dropbox.com/s/ghkcovukgtctm76/NA_merFISH.RDS?dl=0. A down-sample

version of this data are provided by iSpatial package as a test dataset. After installing the R

package, this command "data(NA_merFISH)" could load the merFISH data into R environment.

The scRNA-seq data of mouse striatum was downloaded from GEO under accession

GSE118020. To speeding up the analysis, we down-sampled the full dataset to 10,000 cells using

R function "sample". Then this data was used to infer the spatial expression of all genes.

Mouse liver data processing

The mouse liver merFISH data is download from Vizgen MERFISH Mouse Liver Map

(https://vizgen.com/data-release-program/), which targets 347 genes. This dataset contains

multiply slices. We only used slice 3 from the first replicate in this manuscript. The merFISH

data processing is the same as described above. The matched scRNA-seq data was download

from GEO under the accession GSE166504 (*42*). The original data set not only contains wild-

type healthy livers but also livers with nonalcoholic fatty liver disease (NAFLD). Three healthy

samples (Hepatocyte_Chow_Animal1_Capture1, Hepatocyte_Chow_Animal2_Capture1,

Hepatocyte_Chow_Animal3_Capture1) were used in this analysis.

CV score calculation

To measure the distance between each cell and central vein (CV), we calculated a CV score of

each cell based on well-known CV markers (*Slc1a2*, *Cyp2e1*) and PV markers (*Cyp2f2*,

Aldh1b1). For each cell, the CV score was defined by the mean expression level of *Slc1a2*,

Cyp2e1 minus mean expression level of *Cyp2f2*, *Aldh1b1*. A positive value means more likely

that the cell is located near the CV. On the contrary, a negative value indicates that the cell is

located near the PV. After defining the CV score, we could then identify the genes specifically

expressed in CV/PV. For each gene, we calculated the Pearson's correlation between the

corresponding expression value and CV score across all cells. A positive correlation coefficient

represents gene preferentially expressed in CV, and a negative one represents gene preferentially

expressed in PV. According to the distribution of the correlation coefficient among all genes, we

manually chose > 0.3 or < -0.3 as cutoffs to define genes with most CV/PV bias.

Visualize the spatial expression

601   Visualization of gene spatial expression was achieved using the "SpatialFeaturePlot" in the

602   Seurat package. iSpatial provides a function "spatial_signature_plot" for spatial visualization of

603   the mean expression of a group of genes.

Benchmark methods

We chose two popular methods for comparison: Seurat (version 4.0.5) and rliger (version 1.0.0).

The Seurat provides a "canonical correlation analysis" (CCA) method to integrate the spatial

transcriptome and scRNA-seq. Here we used the function "FindTransferAnchors" with

parameter "reduction = "cca"" to identify the integrated anchors. And then genes expression

values from scRNA-seq data were transferred to spatial transcriptome data using "TransferData"

with the parameter "weight.reduction = "cca"". Different from Seurat, Liger uses an integrative

non-negative matrix factorization (iNMF) method to integrate the datasets, which is embedded in

the function "optimizeALS". For imputing the non-targeted genes by rliger, we performed

following steps according to the rliger vignettes: "createLiger", "normalize", "scaleNotCenter",

"optimizeALS", and "quantile_norm". Finally, "imputeKNN" was used to search the nearest

neighbors and impute the non-targeted genes.

Performance evaluation

The Slide-seq V2 data contains a total of 23,264 genes. We randomly sampled 3,000 genes as the

training dataset and used others as validation. After performing different methods to infer the

spatial expression of all genes, we conducted the comparisons between the inferred expression

values and the raw expression values. At cell level, we calculated the Pearson's correlation

coefficient of each gene across all cells, then compared the overall differences of correlation

coefficients among all genes of different methods. At spatial expression level, we grided the

spatial location into 50×50 bricks. Then, we calculated the mean expression in each brick and

compared the differences between raw expression and inferred one using 50×50 bricks as

features. In addition, we compared the detection accuracies of spatial variable genes among three

methods. The spatially variable genes were also called using validation dataset as the ground

truth. Receiver operating characteristic (ROC) curves and area under the curve (AUC) were

performed by R package pROC (version 1.18.0) (*49*).

Ten-fold cross validation

For the datasets other than Slide-seq which we don't have the ground truth, we used 10-fold

cross validation to evaluate the prediction performance. For the targeted genes in spatial

transcriptome, we randomly separated it into ten groups. Each time we chose nine groups to infer

the expression and used the left one to validate. After ten rounds of inferring and validation,

every gene was used to validate the prediction performance. Then, correlation of gene level and

spatial expression level were calculated as described above.

Data and Software Availability

iSpatial is available as an R package on github (https://github.com/YiZhang-lab/iSpatial).

Acknowledgements

We thank Dr. Aritra Bhattacherjee for some initial discussion and for critical reading of the

manuscript. This project was supported by 1R01DA042283, 1R01DA050589, and HHMI. R.C.

is partially supported by the 2020 NARSAD Young Investigator Grant from the Brain and

Behavior Research Foundation. Y.Z. is an investigator of the Howard Hughes Medical Institute.

Author Contributions

Y.Z. and C. Z. conceived the project. C.Z. developed the iSpatial package and performed the data

analysis. C.Z., R.C., and Y.Z. designed the analyses, interpreted the results, and wrote the

manuscript.

Competing Interests

The authors declare no competing interests.

References

659   1.   A. Saunders, E. Z. Macosko, A. Wysoker, M. Goldman, F. M. Krienen, H. de Rivera, E. Bien, M.
660   Baum, L. Bortolin, S. Wang, A. Goeva, J. Nemesh, N. Kamitaki, S. Brumbaugh, D. Kulp, S. A.
661   McCarroll, Molecular Diversity and Specializations among the Cells of the Adult Mouse Brain.
662   *Cell* **174**, 1015-1030 e1016 (2018).
663   2.   A. Zeisel, H. Hochgerner, P. Lonnerberg, A. Johnsson, F. Memic, J. van der Zwan, M. Haring, E.
664   Braun, L. E. Borm, G. La Manno, S. Codeluppi, A. Furlan, K. Lee, N. Skene, K. D. Harris, J. Hjerling-
665   Leffler, E. Arenas, P. Ernfors, U. Marklund, S. Linnarsson, Molecular Architecture of the Mouse
666   Nervous System. *Cell* **174**, 999-1014 e1022 (2018).
667   3.   X. Han, R. Wang, Y. Zhou, L. Fei, H. Sun, S. Lai, A. Saadatpour, Z. Zhou, H. Chen, F. Ye, D. Huang,
668   Y. Xu, W. Huang, M. Jiang, X. Jiang, J. Mao, Y. Chen, C. Lu, J. Xie, Q. Fang, Y. Wang, R. Yue, T. Li, H.
669   Huang, S. H. Orkin, G. C. Yuan, M. Chen, G. Guo, Mapping the Mouse Cell Atlas by Microwell-
670   Seq. *Cell* **173**, 1307 (2018).
671   4.   O. Rozenblatt-Rosen, M. J. T. Stubbington, A. Regev, S. A. Teichmann, The Human Cell Atlas:
672   from vision to reality. *Nature* **550**, 451-453 (2017).
673   5.   V. Marx, Method of the Year: spatially resolved transcriptomics. *Nat Methods* **18**, 9-14 (2021).
674   6.   P. L. Stahl, F. Salmen, S. Vickovic, A. Lundmark, J. F. Navarro, J. Magnusson, S. Giacomello, M.
675   Asp, J. O. Westholm, M. Huss, A. Mollbrink, S. Linnarsson, S. Codeluppi, A. Borg, F. Ponten, P. I.
676   Costea, P. Sahlen, J. Mulder, O. Bergmann, J. Lundeberg, J. Frisen, Visualization and analysis of
677   gene expression in tissue sections by spatial transcriptomics. *Science* **353**, 78-82 (2016).
678   7.   K. H. Chen, A. N. Boettiger, J. R. Moffitt, S. Wang, X. Zhuang, RNA imaging. Spatially resolved,
679   highly multiplexed RNA profiling in single cells. *Science* **348**, aaa6090 (2015).
680   8.   S. Shah, E. Lubeck, W. Zhou, L. Cai, seqFISH Accurately Detects Transcripts in Single Cells and
681   Reveals Robust Spatial Organization in the Hippocampus. *Neuron* **94**, 752-758 e751 (2017).
682   9.   X. Wang, W. E. Allen, M. A. Wright, E. L. Sylwestrak, N. Samusik, S. Vesuna, K. Evans, C. Liu, C.
683   Ramakrishnan, J. Liu, G. P. Nolan, F. A. Bava, K. Deisseroth, Three-dimensional intact-tissue
684   sequencing of single-cell transcriptional states. *Science* **361**, (2018).
685   10.   C. L. Eng, M. Lawson, Q. Zhu, R. Dries, N. Koulena, Y. Takei, J. Yun, C. Cronin, C. Karp, G. C. Yuan,
686   L. Cai, Transcriptome-scale super-resolved imaging in tissues by RNA seqFISH. *Nature* **568**, 235-
687   239 (2019).
688   11.   Y. Liu, M. Yang, Y. Deng, G. Su, A. Enninful, C. C. Guo, T. Tebaldi, D. Zhang, D. Kim, Z. Bai, E.
689   Norris, A. Pan, J. Li, Y. Xiao, S. Halene, R. Fan, High-Spatial-Resolution Multi-Omics Sequencing
690   via Deterministic Barcoding in Tissue. *Cell* **183**, 1665-1681 e1618 (2020).
691   12.   S. Codeluppi, L. E. Borm, A. Zeisel, G. La Manno, J. A. van Lunteren, C. I. Svensson, S. Linnarsson,
692   Spatial organization of the somatosensory cortex revealed by osmFISH. *Nat Methods* **15**, 932-
693   935 (2018).
694   13.   L. E. Borm, A. M. Albiach, C. C. A. Mannens, J. Janusauskas, C. Özgün, D. Fernández-García, R.
695   Hodge, E. S. Lein, S. Codeluppi, S. Linnarsson, Scalable in situ single-cell profiling by
696   electrophoretic capture of mRNA. *bioRxiv*, 2022.2001.2012.476082 (2022).
697   14.   M. Asp, S. Giacomello, L. Larsson, C. Wu, D. Furth, X. Qian, E. Wardell, J. Custodio, J. Reimegard,
698   F. Salmen, C. Osterholm, P. L. Stahl, E. Sundstrom, E. Akesson, O. Bergmann, M. Bienko, A.
699   Mansson-Broberg, M. Nilsson, C. Sylven, J. Lundeberg, A Spatiotemporal Organ-Wide Gene
700   Expression and Cell Atlas of the Developing Human Heart. *Cell* **179**, 1647-1660 e1619 (2019).
701   15.   R. Dries, J. Chen, N. Del Rossi, M. M. Khan, A. Sistig, G. C. Yuan, Advances in spatial
702   transcriptomic data analysis. *Genome Res* **31**, 1706-1718 (2021).
703   16.   B. Hie, J. Peters, S. K. Nyquist, A. K. Shalek, B. Berger, B. D. Bryson, Computational Methods for
704   Single-Cell RNA Sequencing. *Annual Review of Biomedical Data Science* **3**, 339-364 (2020).

17.   Z. Hu, A. A. Ahmed, C. Yau, CIDER: an interpretable meta-clustering framework for single-cell
RNA-seq data integration and evaluation. *Genome Biol* **22**, 337 (2021).
18.   K. Polanski, M. D. Young, Z. Miao, K. B. Meyer, S. A. Teichmann, J. E. Park, BBKNN: fast batch
alignment of single cell transcriptomes. *Bioinformatics* **36**, 964-965 (2020).
19.   B. Hie, B. Bryson, B. Berger, Efficient integration of heterogeneous single-cell transcriptomes
using Scanorama. *Nat Biotechnol* **37**, 685-691 (2019).
20.   L. Haghverdi, A. T. L. Lun, M. D. Morgan, J. C. Marioni, Batch effects in single-cell RNA-
sequencing data are corrected by matching mutual nearest neighbors. *Nat Biotechnol* **36**, 421-
427 (2018).
21.   D. van Dijk, R. Sharma, J. Nainys, K. Yim, P. Kathail, A. J. Carr, C. Burdziak, K. R. Moon, C. L.
Chaffer, D. Pattabiraman, B. Bierie, L. Mazutis, G. Wolf, S. Krishnaswamy, D. Pe'er, Recovering
Gene Interactions from Single-Cell Data Using Data Diffusion. *Cell* **174**, 716-729 e727 (2018).
22.   W. V. Li, J. J. Li, An accurate and robust imputation method scImpute for single-cell RNA-seq
data. *Nat Commun* **9**, 997 (2018).
23.   W. Gong, I. Y. Kwak, P. Pota, N. Koyano-Nakagawa, D. J. Garry, DrImpute: imputing dropout
events in single cell RNA sequencing data. *BMC Bioinformatics* **19**, 220 (2018).
24.   G. C. Linderman, J. Zhao, M. Roulis, P. Bielecki, R. A. Flavell, B. Nadler, Y. Kluger, Zero-preserving
imputation of single-cell RNA-seq data. *Nat Commun* **13**, 192 (2022).
25.   R. Dong, G. C. Yuan, SpatialDWLS: accurate deconvolution of spatial transcriptomic data.
Genome Biol **22**, 145 (2021).
26.   D. M. Cable, E. Murray, L. S. Zou, A. Goeva, E. Z. Macosko, F. Chen, R. A. Irizarry, Robust
decomposition of cell type mixtures in spatial transcriptomics. *Nat Biotechnol*, (2021).
27.   T. Biancalani, G. Scalia, L. Buffoni, R. Avasthi, Z. Lu, A. Sanger, N. Tokcan, C. R. Vanderburg, A.
Segerstolpe, M. Zhang, I. Avraham-Davidi, S. Vickovic, M. Nitzan, S. Ma, A. Subramanian, M.
Lipinski, J. Buenrostro, N. B. Brown, D. Fanelli, X. Zhuang, E. Z. Macosko, A. Regev, Deep learning
and alignment of spatially resolved single-cell transcriptomes with Tangram. *Nat Methods* **18**,
1352-1362 (2021).
28.   V. Kleshchevnikov, A. Shmatko, E. Dann, A. Aivazidis, H. W. King, T. Li, R. Elmentaite, A. Lomakin,
V. Kedlian, A. Gayoso, M. S. Jain, J. S. Park, L. Ramona, E. Tuck, A. Arutyunyan, R. Vento-Tormo,
M. Gerstung, L. James, O. Stegle, O. A. Bayraktar, Cell2location maps fine-grained cell types in
spatial transcriptomics. *Nat Biotechnol*, (2022).
29.   T. Stuart, A. Butler, P. Hoffman, C. Hafemeister, E. Papalexi, W. M. Mauck, 3rd, Y. Hao, M.
Stoeckius, P. Smibert, R. Satija, Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-
1902 e1821 (2019).
30.   J. D. Welch, V. Kozareva, A. Ferreira, C. Vanderburg, C. Martin, E. Z. Macosko, Single-Cell Multi-
omic Integration Compares and Contrasts Features of Brain Cell Identity. *Cell* **177**, 1873-1887
e1817 (2019).
31.   I. Korsunsky, N. Millard, J. Fan, K. Slowikowski, F. Zhang, K. Wei, Y. Baglaenko, M. Brenner, P. R.
Loh, S. Raychaudhuri, Fast, sensitive and accurate integration of single-cell data with Harmony.
Nat Methods **16**, 1289-1296 (2019).
32.   R. R. Stickels, E. Murray, P. Kumar, J. Li, J. L. Marshall, D. J. Di Bella, P. Arlotta, E. Z. Macosko, F.
Chen, Highly sensitive spatial transcriptomics at near-cellular resolution with Slide-seqV2. *Nat
Biotechnol* **39**, 313-319 (2021).
33.   E. S. Lein, M. J. Hawrylycz, N. Ao, M. Ayres, A. Bensinger, A. Bernard, A. F. Boe, M. S. Boguski, K.
S. Brockway, E. J. Byrnes, L. Chen, L. Chen, T. M. Chen, M. C. Chin, J. Chong, B. E. Crook, A.
Czaplinska, C. N. Dang, S. Datta, N. R. Dee, A. L. Desaki, T. Desta, E. Diep, T. A. Dolbeare, M. J.
Donelan, H. W. Dong, J. G. Dougherty, B. J. Duncan, A. J. Ebbert, G. Eichele, L. K. Estin, C. Faber,
B. A. Facer, R. Fields, S. R. Fischer, T. P. Fliss, C. Frensley, S. N. Gates, K. J. Glattfelder, K. R.

Halverson, M. R. Hart, J. G. Hohmann, M. P. Howell, D. P. Jeung, R. A. Johnson, P. T. Karr, R.
Kawal, J. M. Kidney, R. H. Knapik, C. L. Kuan, J. H. Lake, A. R. Laramee, K. D. Larsen, C. Lau, T. A.
Lemon, A. J. Liang, Y. Liu, L. T. Luong, J. Michaels, J. J. Morgan, R. J. Morgan, M. T. Mortrud, N. F.
Mosqueda, L. L. Ng, R. Ng, G. J. Orta, C. C. Overly, T. H. Pak, S. E. Parry, S. D. Pathak, O. C.
Pearson, R. B. Puchalski, Z. L. Riley, H. R. Rockett, S. A. Rowland, J. J. Royall, M. J. Ruiz, N. R.
Sarno, K. Schaffnit, N. V. Shapovalova, T. Sivisay, C. R. Slaughterbeck, S. C. Smith, K. A. Smith, B. I.
Smith, A. J. Sodt, N. N. Stewart, K. R. Stumpf, S. M. Sunkin, M. Sutram, A. Tam, C. D. Teemer, C.
Thaller, C. L. Thompson, L. R. Varnam, A. Visel, R. M. Whitlock, P. E. Wohnoutka, C. K. Wolkey, V.
Y. Wong, M. Wood, M. B. Yaylaoglu, R. C. Young, B. L. Youngstrom, X. F. Yuan, B. Zhang, T. A.
Zwingman, A. R. Jones, Genome-wide atlas of gene expression in the adult mouse brain. *Nature*
445, 168-176 (2007).
34.   A. Chen, S. Liao, M. Cheng, K. Ma, L. Wu, Y. Lai, X. Qiu, J. Yang, J. Xu, S. Hao, X. Wang, H. Lu, X.
Chen, X. Liu, X. Huang, Z. Li, Y. Hong, Y. Jiang, J. Peng, S. Liu, M. Shen, C. Liu, Q. Li, Y. Yuan, X.
Wei, H. Zheng, W. Feng, Z. Wang, Y. Liu, Z. Wang, Y. Yang, H. Xiang, L. Han, B. Qin, P. Guo, G. Lai,
P. Munoz-Canoves, P. H. Maxwell, J. P. Thiery, Q. F. Wu, F. Zhao, B. Chen, M. Li, X. Dai, S. Wang,
H. Kuang, J. Hui, L. Wang, J. F. Fei, O. Wang, X. Wei, H. Lu, B. Wang, S. Liu, Y. Gu, M. Ni, W.
Zhang, F. Mu, Y. Yin, H. Yang, M. Lisby, R. J. Cornall, J. Mulder, M. Uhlen, M. A. Esteban, Y. Li, L.
Liu, X. Xu, J. Wang, Spatiotemporal transcriptomic atlas of mouse organogenesis using DNA
nanoball-patterned arrays. *Cell* **185**, 1777-1792 e1721 (2022).
35.   B. Tasic, Z. Yao, L. T. Graybuck, K. A. Smith, T. N. Nguyen, D. Bertagnolli, J. Goldy, E. Garren, M.
N. Economo, S. Viswanathan, O. Penn, T. Bakken, V. Menon, J. Miller, O. Fong, K. E. Hirokawa, K.
Lathia, C. Rimorin, M. Tieu, R. Larsen, T. Casper, E. Barkan, M. Kroll, S. Parry, N. V. Shapovalova,
D. Hirschstein, J. Pendergraft, H. A. Sullivan, T. K. Kim, A. Szafer, N. Dee, P. Groblewski, I.
Wickersham, A. Cetin, J. A. Harris, B. P. Levi, S. M. Sunkin, L. Madisen, T. L. Daigle, L. Looger, A.
Bernard, J. Phillips, E. Lein, M. Hawrylycz, K. Svoboda, A. R. Jones, C. Koch, H. Zeng, Shared and
distinct transcriptomic cell types across neocortical areas. *Nature* **563**, 72-78 (2018).
36.   R. Chen, T. R. Blosser, M. N. Djekidel, J. Hao, A. Bhattacherjee, W. Chen, L. M. Tuesta, X. Zhuang,
Y. Zhang, Decoding molecular and cellular heterogeneity of mouse nucleus accumbens. *Nat*
Neurosci **24**, 1757-1771 (2021).
37.   G. Stanley, O. Gokce, R. C. Malenka, T. C. Sudhof, S. R. Quake, Continuous and Discrete Neuron
Types of the Adult Murine Striatum. *Neuron* **105**, 688-699 e688 (2020).
38.   P. Voorn, L. J. Vanderschuren, H. J. Groenewegen, T. W. Robbins, C. M. Pennartz, Putting a spin
on the dorsal-ventral divide of the striatum. *Trends Neurosci* **27**, 468-474 (2004).
39.   D. C. Castro, K. C. Berridge, Opioid hedonic hotspot in nucleus accumbens shell: mu, delta, and
kappa maps for enhancement of sweetness "liking" and "wanting". *J Neurosci* **34**, 4239-4250
(2014).
40.   R. Al-Hasani, J. G. McCall, G. Shin, A. M. Gomez, G. P. Schmitz, J. M. Bernardi, C. O. Pyo, S. I. Park,
C. M. Marcinkiewcz, N. A. Crowley, M. J. Krashes, B. B. Lowell, T. L. Kash, J. A. Rogers, M. R.
Bruchas, Distinct Subpopulations of Nucleus Accumbens Dynorphin Neurons Drive Aversion and
Reward. *Neuron* **87**, 1063-1077 (2015).
41.   J. W. de Jong, S. A. Afjei, I. Pollak Dorocic, J. R. Peck, C. Liu, C. K. Kim, L. Tian, K. Deisseroth, S.
Lammel, A Neural Circuit Mechanism for Encoding Aversive Stimuli in the Mesolimbic Dopamine
System. *Neuron* **101**, 133-151 e137 (2019).
42.   Q. Su, S. Y. Kim, F. Adewale, Y. Zhou, C. Aldler, M. Ni, Y. Wei, M. E. Burczynski, G. S. Atwal, M. W.
Sleeman, A. J. Murphy, Y. Xin, X. Cheng, Single-cell RNA transcriptome landscape of hepatocytes
and non-parenchymal cells in healthy and NAFLD mouse liver. *iScience* **24**, 103233 (2021).
43.   K. B. Halpern, R. Shenhav, O. Matcovitch-Natan, B. Toth, D. Lemze, M. Golan, E. E. Massasa, S.
Baydatch, S. Landen, A. E. Moor, A. Brandis, A. Giladi, A. S. Avihail, E. David, I. Amit, S. Itzkovitz,

Single-cell spatial reconstruction reveals global division of labour in the mammalian liver. *Nature*
542, 352-356 (2017).
44.   F. Hildebrandt, A. Andersson, S. Saarenpaa, L. Larsson, N. Van Hul, S. Kanatani, J. Masek, E. Ellis,
A. Barragan, A. Mollbrink, E. R. Andersson, J. Lundeberg, J. Ankarklev, Spatial Transcriptomics to
define transcriptional patterns of zonation and structural components in the mouse liver. *Nat
Commun* **12**, 7046 (2021).
45.   A. Rao, D. Barkley, G. S. Franca, I. Yanai, Exploring tissue architecture using spatial
transcriptomics. *Nature* **596**, 211-220 (2021).
46.   A. Bhattacherjee, M. N. Djekidel, R. Chen, W. Chen, L. M. Tuesta, Y. Zhang, Cell type-specific
transcriptional programs in mouse prefrontal cortex during adolescence and addiction. *Nat
Commun* **10**, 4169 (2019).
47.   B. I. C. C. Network, A multimodal cell census and atlas of the mammalian primary motor cortex.
Nature **598**, 86-102 (2021).
48.   A. Regev, S. A. Teichmann, E. S. Lander, I. Amit, C. Benoist, E. Birney, B. Bodenmiller, P.
Campbell, P. Carninci, M. Clatworthy, H. Clevers, B. Deplancke, I. Dunham, J. Eberwine, R. Eils,
W. Enard, A. Farmer, L. Fugger, B. Gottgens, N. Hacohen, M. Haniffa, M. Hemberg, S. Kim, P.
Klenerman, A. Kriegstein, E. Lein, S. Linnarsson, E. Lundberg, J. Lundeberg, P. Majumder, J. C.
Marioni, M. Merad, M. Mhlanga, M. Nawijn, M. Netea, G. Nolan, D. Pe'er, A. Phillipakis, C. P.
Ponting, S. Quake, W. Reik, O. Rozenblatt-Rosen, J. Sanes, R. Satija, T. N. Schumacher, A. Shalek,
E. Shapiro, P. Sharma, J. W. Shin, O. Stegle, M. Stratton, M. J. T. Stubbington, F. J. Theis, M.
Uhlen, A. van Oudenaarden, A. Wagner, F. Watt, J. Weissman, B. Wold, R. Xavier, N. Yosef, P.
Human Cell Atlas Meeting, The Human Cell Atlas. *Elife* **6**, (2017).
49.   X. Robin, N. Turck, A. Hainard, N. Tiberti, F. Lisacek, J. C. Sanchez, M. Muller, pROC: an open-
source package for R and S+ to analyze and compare ROC curves. *BMC Bioinformatics* **12**, 77
(2011).

Figure

Fig. 1: Overview of iSpatial. **A**, A diagram showing the rationale of iSpatial. The probe-based

spatial transcriptome includes $t \times n$ (*t*, genes; *n*, cells) expression matrix and location of each

cell. After integrating with $m \times n'$ scRNA-seq data, iSpatial infers the genome-wide

transcriptional expression of all *n* cells. **B**, The iSpatial pipeline is consisted of: 1) Dimension

reduction and expression stabilization; 2) Expression normalization; 3) Inferring transcriptional

expression; and 4) Downstream analysis such as spatial variable gene detection.

Fig. 2: Benchmarking the performance of iSpatial in inferring genome-wide spatial

transcriptome. **A**, A graphical illustration of the evaluation procedure. **B**, Representative

examples showing the performance of Liger, Seurat and iSpatial in inferring spatial

transcriptome. **C**, Representative examples of inferred expression of genes barely detectable in

raw Slide-seq data. **D**, The gene expression correlation between Slide-seq raw data and Liger,

Seurat, or iSpatial inferred data. The validation genes are divided into 3 groups based on their

expression levels. Two-side Wilcoxon Rank Sum test was used. **E**, The spatial expression

correlation between inferred and raw Slide-seq data. Two-side Wilcoxon Rank Sum test was

used. **F**, The ROC curves comparing the prediction power of spatial variable genes by Liger,

Seurat, and iSpatial. AUCs (Area under the curves) are indicated.

Fig. 3: iSpatial accurately infers the genome-wide spatial transcriptomes in mouse cortex and striatum. **A**, Schematic of the anatomic region of mouse V1 cortex. **B**, The numbers of detectable genes in each cell in raw STARmap (left) and after inferring by iSpatial (right). The mean number of detected genes in each cell are shown in brackets. **C**, Inferred layer expression patterns of representative genes not targeted in the original STARmap library compared with the ISH data from the ABA. **D**, Schematic of the anatomic region of mouse striatum. **E**, The numbers of detectable genes in each cell in raw MERFISH (left) and after inferring by iSpatial

(right). **F**, Inferred spatial expression patterns of representative genes not targeted in the original

MERFISH library compared with the ISH data from the ABA.

Fig. 4: iSpatial can reduce false-positive and false-negative signals in the original spatial

transcriptomic data. **A**, UMAPs showing the expression levels of representative cell type

markers in raw STARmap (top panels) and iSpatial inferred (bottom panels) data. Excitatory

neuron (*Slc17a7*), Inhibitory neuron (*Gad1*), Oligodendrocyte (*Plp1*), Astrocyte (*Aqp4*), and

Endothelial cell (*Cldn5*). **B**, The spatial expression of cortex layer markers in the raw STARmap

(top panels) and inferred by iSpatial (middle panels) compared with the ISH data from the ABA

(bottom panels). Layer information: "L1 to L6", the six cortical layers; "cc", corpus callosum;

"HPC", hippocampus. **C**, The UMAP and spatial expression of *Tac2* and *Serpinb2* in the raw

MERFISH (top panels) and inferred by iSpatial (middle panels) compared with the ISH data

from the ABA (bottom panels).

Fig. 5: iSpatial enables whole-transcriptome level spatial analysis. **A**, Scatterplot showing genes with expression specificity in spatial location and UMAP projection of the corresponding scRNA-seq data. **B**, The 12 clusters of the spatial variable genes in mouse striatum. The plot is colored coded by the average gene expression in each cluster. **C**, Inferred spatial expression signals of representative genes in different clusters. C1: *Dio3* and *Grm8*, C5: *Olfml1* and *Slc12a2*, C7: *Acbd7* and *Myo3b*, C8: *Sdk1* and *Trhr*, C11: *Camk1g* and *Tmco5*.

Fig. 6: iSpatial infers the spatial expression patterns in liver. **A**, The numbers of detectable

genes in each cell in raw MERFISH (left) and after inferring by iSpatial (right). **B**, Boxplot

showing the detectable gene numbers without or with interfering by iSpatial. **C**, Representative

examples of genes exhibiting spatial patterns enriched in CV (*Slc1a2*, *Cyp2e1*) or PV (*Cyp2f2*,

Aldh1b1). **D**, UMAPs showing the expression levels of the genes showed in (C). The UMAPs

were generated with iSpatial inferred expression. **E**, The spatial location of each cell colored by

CV score. **F**, The UMAP of all cells colored by CV score. **G**, The scatterplot showing the ranked

correlations between gene expression level and CV score. The names of top 10 enriched genes in

PV or CV are listed. **H**, Examples of genes selectively expressed nearby CV or PV. **I**, The top 10

enriched KEGG pathways of genes selectively expressed near CV or PV.

Supplementary Figures

fig. S1: Benchmarking the performance of iSpatial in inferring genome-wide spatial

transriptome. **A**, Representative examples showing the performance of Liger, Seurat and

iSpatial in inferring spatial transcriptome. **B**, The gene expression correlation between raw data

and Liger, Seurat, or iSpatial inferred data of cell type specific genes or other genes. **C**, The

overlaps of the spatial variable genes detected by different methods.

fig. S2: Benchmarking the performance on Stereo-seq data. A, A graphical illustration of the

evaluation procedure. **B**, Representative examples showing the raw or inferred spatial expression

by Liger, Seurat and iSpatial. Allen brain ISH data are also shown. **C**, The gene expression

correlation between Stereo-seq raw data and Liger, Seurat, or iSpatial inferred data. The

validation genes are divided into 3 groups based on their expression levels. **D**, The spatial

expression correlation between inferred and raw Stereo-seq data. **E**, The gene expression

correlation between Stereo-seq raw data and Liger, Seurat, or iSpatial inferred data of cell type

specific genes or other genes. Two-side Wilcoxon Rank Sum test was used.

fig. S3: iSpatial is applicable to MERFISH dataset. **A**, Spatial distribution of the cell types

detected by MERFISH. **B**, The heatmap of the 175 cell type markers detected in the raw

MERFISH data. **C**, The numbers of detected genes in raw MERFISH data and after interfering

by iSpatial. **D**, The number of cell type specific expressed genes in each cell type detected in raw

and iSpatial inferred data. **E**, The heatmap of all maker genes detected after inferring by iSpatial.

fig. S4: Ten-fold cross validation on the performance of the three methods in mouse cortex

(A) and striatum (B). Left panels: gene expression correlation between raw expression data and

inferred data. Right panels: spatial expression correlation between raw expression data and

inferred data.

fig. S5: iSpatial can reduce false-positive and false-negative signals in the original

MERFISH data. A, Cell type clusters on the UMAPs derived from raw or inferred data. **B**,

UMAPs showing the expression of representative cell type marker genes. The UMAPs are

generated from raw or iSpatial inferred expression profiles, respectively. Marker genes for

neuron (*Snap25*), D1 medium spiny neuron (*Drd1*), D2 medium spiny neuron (*Drd2*), and

astrocytes (*Gja1*) are presented. **C**, The UMAP and spatial expression of *Kctd4* in the raw

MERFISH (top panels) and inferred by iSpatial (middle panels) compared with the ISH data

from the ABA (bottom panel).

fig. S6: iSpatial detects spatial variable genes in mouse cortex (A, B) and striatum (C, D).

A, The number of spatial variable genes in mouse V1 cortex detected by STARmap and after

iSpatial interferring. **B**, The layer distribution of the 6 clusters spatial variable genes. The plot is

colored by the average expression level of the genes in each cluster. **C**, The number of spatial

variable genes in mouse striatum detected by MERFISH and after iSpatial interfering. **D**,

Inferred spatial expression signals of representative genes in the C8 cluster.

A

B

C

fig. S7: iSpatial infers the spatial expression patterns in liver. **A, B**, Examples of spatial

expressed genes enriched in CV (*Slc1a2*, *Cyp2e1*) or PV (*Cyp2f2*, *Aldh1b1*), inferred by Seurat

(A) and Liger (B). The upper panels show the spatial expression patterns. The bottom panels

show the inferred genes' expression by the indicated method projects on the iSpatial UMAP. **C**,

The top 20 enriched Gene Ontology terms of genes selectively expressed near CV or PV.

fig. S8: Stabilization and two-rounds integration steps contribute to the improved

performance of iSpatial. A, B, UMAPs showing the expression signals of representative

markers before (top panels) and after (bottom panels) expression stabilization in mouse striatum

(**A**) or cortex (**B**) data. **C, D**, The co-embed UMAPs showing the integration of ST and scRNA-

seq data in mouse striatum (**C**) or cortex (**D**) data.

fig S9: Comparing the two-rounds integration with one round. The spatial expression of the representative genes of the mouse cortex raw STARmap data inferred by one round (top two rows) or two rounds (third row) of integration by iSpatial compared with the ISH data from the ABA (bottom row). Reciprocal PCA is used for the first-round integration, and Harmony is used for the second-round integration.

969
fig S10: Comparing the performance of weighted or un-weighted KNN. A, B, The inferred
spatial expression of *Serpinb2* (**A**) and *Tac2* (**B**) in rare cell population using un-weighted KNN
or weighted KNN under different K value. Allen brain ISH data were showed on right. Arrows
indicate the location of the rare cell population.
974

976
fig. S11: Infer spatial expression by unmatched ST and scRNAseq data. A, Schematic of the
anatomic region of mouse cortex and striatum. **B**, Varied number of measured genes in each cell
of a whole brain slice inferred by Liger, Seurat and iSpatial. Unmatched MERFISH and scRNA-
seq data are used. MERFISH data covers the entire slice. scRNA-seq data only covers cortex
region. **C and D**, Representative examples showing the performance of Liger, Seurat and
iSpatial in inferring spatial transcriptome with unmatched scRNA-seq data. Allen brain ISH data
were showed in left.