Episodic memory formation in unrestricted viewing

Andrey R. Nikolaev[1,2]*, Inês Bramão[1], Roger Johansson[1], Mikael Johansson[1]

[1]Lund Memory Lab, Department of Psychology, Lund University, Lund, Sweden

[2] Brain & Cognition Research Unit, KU Leuven, Leuven, Belgium

* Corresponding author

Department of Psychology,

Lund University,

Box 213

22100 Lund

Sweden

andrey.nikolaev@psy.lu.se

Abstract

The brain systems of episodic memory and oculomotor control are tightly linked, suggesting a crucial role of eye movements in memory. But little is known about the neural mechanisms of memory formation across eye movements in unrestricted viewing behavior. Here, we leverage simultaneous eye tracking and EEG recording to examine episodic memory formation in free viewing. Participants memorized multi-element events while their EEG and eye movements were concurrently recorded. Each event comprised elements from three categories (face, object, place), with two exemplars from each category, in different locations on the screen. A subsequent associative memory test assessed participants' memory for the between-category associations that specified each event. We used a deconvolution approach to overcome the problem of overlapping EEG responses to sequential saccades in free viewing. Brain activity was time-locked to the fixation onsets, and we examined EEG power in the theta and alpha frequency bands, the putative oscillatory correlates of episodic encoding mechanisms. Three modulations of fixation-related EEG predicted high subsequent memory performance: 1) theta increase at fixations after *between-category* gaze transitions, 2) theta and alpha increase at fixations after *within-element* gaze transitions, 3) alpha decrease at fixations after *between-exemplar* gaze transitions. Thus, event encoding with unrestricted viewing behavior was characterized by three neural mechanisms, manifested in fixation-locked theta and alpha EEG activity that rapidly turned on and off during the unfolding eye movement sequences. These three distinct neural mechanisms may be the essential building blocks that subserve the buildup of coherent episodic memories during unrestricted viewing behavior.

Keywords: EEG, eye movement, unrestricted viewing, episodic memory, deconvolution modeling

1 Introduction

Life events consist of visual, semantic, contextual, and other associated features stored as coherent episodic memory representations. Episodic memory allows us to mentally travel in time to relive past events with considerable detail and specificity (Tulving, 1983). A critical aspect when building such memory representations is that the visual field of our perception is spatially limited. At any given time, we can only apprehend information in full acuity in a small spot, determined by the size of the fovea. We overcome this limitation by constantly shifting our visual focus with eye movements. In effect, these visual "samples" of the world constitute the pieces we successively bind together into meaningful, coherent episodic memories (e.g., Ryan et al., 2020; Voss et al., 2017). Nonetheless, in the neuroscience of human memory, memory formation has primarily been studied in experimental paradigms where the study material is presented in a single location on the screen and where eye movements are typically treated as artifacts. Thus, little is currently known about the neural mechanisms of episodic memory formation across eye movements. In the present study, we set out to capture the neural signatures of building coherent episodic memories via eye movements at the level of individual gaze fixations. To this end, we apply a state-of-the-art approach of simultaneous eye-tracking and EEG recording and analysis in a free-viewing memory task.

Episodic memory formation critically hinges upon binding together the different elements that constitute an event. Sequential saccadic eye movements organize these relationships as a mechanism that links different aspects of the environment into coherent memory representations (for overviews, see Ryan et al., 2020; Voss et al., 2017). Eye movements also serve an important role during memory retrieval (Damiano and Walther, 2019; Johansson and Johansson, 2014; Johansson et al., 2022; Wynn et al., 2020), and a longstanding theory holds that the stored representations include the sequences of eye movements established during memory formation, which supposedly carry information about linked event elements (Brandt and Stark, 1997; Noton and Stark, 1971).

The clock frequency of this 'linking' is the theta rhythm (4-7 Hz). We typically make about 3-4 saccades in natural viewing per second. Such visual sampling frequency corresponds to the theta rhythm (Amit et al., 2017; Otero-Millan et al., 2008). The saccadic theta rhythm is part of a periodic mechanism of attentional deployment (Hogendoorn, 2016). It also corresponds to theta oscillations of the hippocampus, which is of crucial importance for associative memory formation (Cohen and Eichenbaum, 1993). The hippocampus also has tight anatomical connections with brain structures that control oculomotor behavior (Pierrot-Deseilligny et al., 2004; Shen et al., 2016). Recent data suggest that hippocampal theta oscillations support saccade guidance during memory formation on

a fixation-by-fixation basis (Jutras et al., 2013; Kota et al., 2020; Kragel et al., 2021; Kragel et al., 2020). In this way, the theta cycle may organize sequential visual inputs and enables the binding of event elements into a coherent memory representation (Herweg et al., 2020).

The relevance of theta oscillations for memory processes has long been established in studies using stimulus-response paradigms with fixed stimulus presentations. For instance, increased frontal theta power during episodic memory encoding predicts subsequent retrieval performance (Hsieh and Ranganath, 2014; Klimesch, 1996). The increase in theta power, i.e., theta synchronization, is often accompanied by a decrease in alpha (8-13 Hz) power, i.e., alpha desynchronization (Hanslmayr et al., 2016; Waldhauser et al., 2012). Alpha desynchronization may represent the fidelity of memory representations for the constituent elements of episodes (Hanslmayr et al., 2016). Thus, alpha and theta oscillations serve essential but distinct functions of episodic memory formation: representing elements (alpha) and binding them together into a coherent whole (theta). It is currently unknown how these mechanisms engage during the sequential sampling of event elements across saccades to allow later episodic remembering of the event as a whole. In the current study, we track how these neural mechanisms subserve the buildup of episodic memories during unrestricted viewing behavior.

Brain activity and eye movements must be examined simultaneously to allow insight into neural mechanisms involved in unrestricted viewing. Over the last decade, the combination of EEG and eye-tracking has become a powerful tool for studying perceptual and cognitive processes in unrestricted viewing (Coco et al., 2020; Devillez et al., 2015; Dimigen et al., 2011; Fudali-Czyz et al., 2018; Körner et al., 2014; Tyson-Carr et al., 2020), including attention and memory across saccades. For instance, presaccadic EEG activity has been found to predict successful working memory of task-relevant scene elements in a change blindness task (Nikolaev et al., 2011). However, despite the apparent benefit of utilizing EEG-eye movement coregistration, attempts to apply it are virtually absent in the episodic memory literature. One explanation may be the vast methodological challenges when combining these techniques, including the effects of overlapping saccades on EEG activity in free viewing (Dimigen and Ehinger, 2021; Dimigen et al., 2011; Nikolaev et al., 2016). Moreover, low-level oculomotor effects of eye movement characteristics can systematically differ between experimental conditions and distort the EEG, creating differences that are likely to be confounded with the effects of the experimental conditions. To overcome these and other challenges as described below, the present study applied a regression-based deconvolution approach that recovers unknown isolated neural responses given the measured EEG and the latencies of experimental events (Dimigen and

Ehinger, 2021; Ehinger and Dimigen, 2019). This approach ultimately allowed us to isolate and track

the neural correlates of episodic memory formation across eye movements.

In the present study, we investigate how theta and alpha activity subserve the binding of separate

event elements into a coherent episodic memory across eye movements. To this end, we introduced

a free-viewing episodic memory task in which participants encoded multiple events. Each event was

defined by six distinct event elements (images), which always comprised two exemplars from each

of three categories (faces, places, objects) presented in different locations on the screen. The two

exemplars from the same category always appeared close together, forming a pair, whereas the

distance between categories (i.e., between the pairs) was much greater (Fig. 1). Participants were

asked to memorize the images that appeared together on the same screen, i.e., belonged to the

same event. In a subsequent associative memory task, we tested participants' episodic memory for

the between-category associations (i.e., associations between elements from different categories)

within each event (Fig. 1).

To form a coherent event representation during encoding, participants critically need to engage in

sequential eye movements, where each event element gets visually attended to on one or multiple

occasions. Thus, to successfully establish the element associations specifying each event, we

expected the encoding process to critically depend on participants' gaze transitions between the

elements (cf. Kragel et al., 2021), as well as on the visual sampling of each individual element (cf. Liu

et al., 2017; Loftus, 1972; Olsen et al., 2016). As the ensuing associative memory task concerned

between-category associations, we expected gaze transitions between categories to be of particular

importance and, thus, more predictive of memory performance than transitions between exemplars

from the same category (associations that participants were not asked to retrieve). Based on

previous research by Kragel et al. (2021), we also expected repeated cross-category gaze transitions

to facilitate successful event formation.

We simultaneously recorded EEG and eye movements during event encoding to examine theta and

alpha activity related to such gaze behaviors during memory formation. In our analytical approach,

we first sought to verify that the oscillatory nature of the EEG signals coregistered with eye

movements in our free-viewing paradigm corresponds to the encoding-related neural signatures

typically found in comparable studies with fixed stimulus presentation (Hanslmayr et al., 2016). To

achieve the central goals of the present study, we then corrected EEG from overlapping effects of

eye movements using a deconvolution approach (Ehinger and Dimigen, 2019) and extracted the EEG

power relative to fixation onset. Since this was the first study to examine scalp-recorded EEG

coregistered with eye movements during a free-viewing associative memory task, we first

investigated the time course of fixation-related EEG during the whole 10-s interval of event

encoding. Then, we examined EEG power related to three types of fixations: (1) fixations succeeding

a *between-category* saccade (i.e., a saccade moving from one element to another across categories);

(2) fixations succeeding a *between-exemplar* saccade (i.e., a saccade moving from one element to

another within the same category); and (3) fixations succeeding a *within-element* saccade (i.e., a

saccade moving from one point to another within the same element). In general, we expect all

saccades between event elements to be important for building a coherent episodic representation.

However, as our memory task required the retrieval of between-category associations, we reasoned

that saccades between categories should be vital and predictive of subsequent performance, in

contrast to between-exemplar saccades supporting associations that were not called for. Thus, by

analyzing both saccade types, we can examine the specificity of gaze transitions between event

elements upon subsequent memory performance as a function of task relevance. This approach

allowed us to examine alpha and theta activity related to functionally distinct gaze behaviors

contributing with complementary aspects to the episodic memory representation: eye movements

that serve to (1) sample visual information from the individual event elements and crucially (2) link

the different event elements into a coherent whole.

2   Methods

2.1   Participants

Thirty-six healthy adults participated in the experiment in exchange for a gift card in a shopping mall

(approx. €10). Three participants were excluded because of system crashes during the data

collection and another three due to below-chance memory performance (the chance level threshold

was set at 53.2%, which was obtained from the permutation procedure indicating chance

performance in a two-alternative forced choice (2AFC) task ('left' or 'right' responses in a memory

test) for the 5% confidence level and for a total number of 648 test trials). Finally, two participants

were excluded due to ceiling performance (> 90%), i.e., too few incorrect trials for the planned

subsequent memory analysis. The final group included 28 participants (20 females; mean age 23.3;

age range 18-32).

The study was conducted in accordance with the Swedish Act concerning the Ethical Review of

Research involving humans (2003:460). All participants gave written informed consent, and the

study followed the local ethical guidelines at Lund University. Moreover, the study followed the Lund

180 University safety protocol for conducting EEG, eye-tracking, and behavioral experiments in the

181 context of the COVID-19 pandemic.

2.2 Stimuli

A total of 324 photo images from three different categories: faces, places, and objects, were used.

Faces were selected from the Oslo Face Database (Chelnokova et al., 2014), which provided consent

for the publication of the faces that were used in the figures. Objects were selected from the Bank of

Standardized Stimuli (Brodeur et al., 2010). Places were selected from the Michael Tarr lab scene

database (http://www.tarrlab.org/). The images were converted to black-and-white, and each

image's pixel intensity was normalized to the average pixel intensity of all images. In the present

study, multiple simultaneously presented images defined an 'event'. Each event comprised three

image pairs (two exemplars from each category) distributed across distinct locations on the display

(Fig. 1). Two exemplars of each a category were necessary for sufficient task complexity with

respect to the target-distractor dynamics (i.e., exemplars from the same category both within and

across events). Moreover, this particular configuration was found to give optimal memory

performance (60-70% correct) for nine events in the block (see below) in pilot behavioral studies,

where we manipulated the number of exemplars (one or three) per category together with the

number of events per block. Exemplars from the same category were close to each other, whereas

the different categories were more separated. Specifically, each image had a square shape with a

side of 4.2 degrees of visual angle (dva). The width of a pair (two exemplars from one category) was

8.5 dva, and the distance between the centers of the three pairs was 17.9 dva. The three pairs were

placed around the perimeter of an invisible circle with a radius of 9.2 dva centered at the center of

the screen. The location of the pairs on the circle was random for each event, but the pairs were

always separated by arcs of 120 degrees. The distribution of images and their pairing across events

was random for each participant. The images were presented on a grey background.

2.3 Procedure

Participants sat in a dimly lit room, with their heads stabilized with a chin rest. Stimuli were

presented at a viewing distance of 62 cm on an EIZO FlexScan EV2451 monitor with a resolution of

1920 x 1080 pixels and a refresh rate 60 Hz, which was part of the Tobii Pro Spectrum eye tracker.

The experiment comprised six blocks with the possibility to take a short break between each block.

Each block included two main stages: memory encoding and retrieval, with a distractor task in

between (Fig. 1). Participants self-initiated each block by pressing the space bar. During encoding,

nine events were presented for 10 s each. Before each event, a fixation cross was presented for a

random interval between 1 and 1.5 s. Participants were instructed to memorize the faces, places,

and objects that appeared together in the same event. The encoding stage was followed by a

distractor task, where participants were asked to count backward in steps of 7 from a random 3-digit

number for 30 s. Pilot experiments indicated that the selected number of exemplars, events, and

blocks gave rise to an optimal memory performance (60-70% correct).

A cued recall task followed three seconds after the distractor task, which assessed episodic memory

by testing retrieval of the between-category associations of each event. A test screen with three

images, one at the top and two at the bottom, was presented for 2 s (Fig. 1). Each test screen was

preceded by a fixation cross randomly jittered between 1 and 1.5 s. The image at the top served as

the retrieval cue. Participants were asked to indicate which of the two images at the bottom had

appeared together with the cue during encoding, i.e., to indicate the target. The distractor was an

exemplar from the same category as the target but selected from a different event studied no more

than two events before or after the target event. Participants responded by pressing the left or right

arrow keys of a standard keyboard. Participants were thereafter asked to indicate how confident

they were in their response. The question "Are you sure?" was displayed on the screen, and

participants responded: "Sure", "Maybe", or "Guess", by pressing the left, down, and right arrow

keys, respectively.

Between-category associations were tested for each image in each event on four trials. For example,

a face image was tested twice with place images (once as cue and once as target), and twice with

object images (once as cue and once as target). The cue-target status of categories was

counterbalanced across blocks, such that in the odd blocks, the cue-target status was faces-places,

places-objects, objects-faces, and in the even blocks, the corresponding cue-target status was

places-faces, objects-places, faces-objects. Thus, memory for each event was assessed by tests of 12

different associations. In total, 108 associations were tested in each block (12 x 9 = 108 tests for the

nine events within a block) with randomized order of presentation. Participants were not explicitly

instructed that only the between-category associations were tested but could infer that from the

exhaustive practice sessions (described below).

Memory performance for the associates may depend on varying encoding difficulty for the three

categories. We eliminated such potential confound by the counterbalancing of individual exemplars

and their screen location across events, blocks, and participants, during both encoding and testing

screens (Fig. 1). We also evaluated performance as a function of the type of cross-category

association within an event: faces – places, faces – objects, places – objects (reported in 3.1).

Prior to the main experiment, two practice sessions with increasing difficulty (three events in the

first session, six events in the second session) were conducted in which participants received

feedback on their performance after each test trial. The gradual increase in difficulty over the two

practice sessions served to prepare participants for what they could expect in the main experiment

(where nine events per block were presented). Ten associative memory tests were delivered in the

practice sessions, in which between-category associations were evenly distributed over events and

categories. Images in the practice sessions were not repeated in the main experiment and were not

repeated over the two practice sessions. The overall duration of the experiment was about two

hours. The experiment was programmed in PsychoPy v.2020.2.4 (Peirce et al., 2019).

Fig. 1. At encoding, nine events were sequentially presented. Each event comprised two exemplar images from

three different categories (faces, places, objects) and was presented for 10 s. After a 30-s distractor task,

episodic memory was assessed by testing all between-category associations from each event. The test screen

(presented for 2 s) included a cue image at the top with two images at the bottom from a different category.

Participants were asked to indicate which of the two images had appeared together with the cue in the

previously encountered events. In this example, the correct response is the lock to the right, as it appeared

together with the face cue in Event 1, whereas the binoculars to the left appeared in Event 2. Note that the

angular distances between categories were much larger on the actual encoding screen.

2.4 Simultaneous EEG and eye movement recording

EEG and eye movements were recorded concurrently throughout the experiment. EEG data were recorded using a Grael 4K EEG amplifier (Neuroscan, Compumedics Limited, Australia) at a sampling frequency of 2048 Hz using 31 Ag/AgCl electrodes positioned according to the extended 10-20 International system (Easycap GmbH, Germany). Two electrodes were mounted over the left and right mastoids, using the latter as the recording reference. The ground electrode was AFz. Additional electrodes recorded the vertical and horizontal electrooculograms (VEOG and HEOG): the VEOG electrodes were placed above and below the right eye, the HEOG electrodes were placed at the left and right outer canthi of two eyes. Electrode impedances were kept below 5 kΩ.

The Tobii Pro Spectrum eye tracker (Tobii, Stockholm, Sweden) recorded movements of both eyes with the sampling frequency of 600 Hz. 9-point calibration and validation routines were conducted before the first block. An error of 1 dva between calibration and validation was tolerated: if it was larger, the calibration was repeated. The eye tracking was controlled from the PsychoPy experimental program via the open-source toolbox Titta (Niehorster et al., 2020). The experimental program ran on the same computer that controlled the eye tracker. To synchronize stimulus presentation and EEG recording, a transistor-transistor logic (TTL) signal was sent via a parallel port from the stimulus presentation computer to the EEG system at the beginning and end of each block, encoding event, and memory test.

2.5 Data analysis

Only data from the encoding stage were analyzed. Analysis was conducted with custom scripts for R (Team, 2020) as well as for MATLAB R2020a (The MathWorks Inc., Natick, Massachusetts), using the EEGLAB toolbox (Delorme and Makeig, 2004), the EYE-EEG extension (Dimigen et al., 2011) for EEGLAB, and the Unfold toolbox (Ehinger and Dimigen, 2019).

2.5.1 Behavioral analysis

Memory performance for each event was assessed by retrieval accuracy weighted by confidence, such that correct responses were scored as 3 for "Sure" responses, as 2 for "Maybe" responses, and as 1 for "Guess" responses. Each image from each event occurred either as a cue or a target in four memory tests trials. Thus, each image can in total receive a score ranging from 0 (incorrect responses in all four tests) to 12 (correct responses accompanied by four "sure" responses in all four tests). To calculate a memory score for the whole event, we then summed the scores for the six elements of each event (ranging from 0 to 72). For the EEG analysis, events were divided into "high memory" and "low memory," according to the subsequent memory scores. For each participant,

high and low memory was calculated by a median split of that participant's memory scores for the

whole events.

2.5.2 Eye movement analysis

Fixations and saccades were detected from participants' right eye using the velocity-based algorithm

for saccade detection (Engbert and Mergenthaler, 2006) implemented in the EYE-EEG extension.

Fixations were considered located on an image if within 0.5 dva of the outer contour of the image.

Since participants cannot adequately perform a memory test if they do not visually process (fixate)

all images while event encoding, we excluded from all analyses the events where fixations were

detected only on two of the three categories (on average 3±4 (mean ± SD) events per participant).

Such events occurred because of eye-tracking issues mainly towards the end of the experiment (see

the Supplementary materials and Fig. S1 for details). Fixations outside the images on different parts

of the display were assigned to an 'other' saccade type, which was used as a reference level in the

deconvolution modeling (2.5.5).

Saccades were classified according to whether they occurred between categories (between-category

saccades), between the two exemplars of a category (between-exemplar saccades), and within an

exemplar (within-element saccades). In addition, we distinguished between the first visit to an image

and revisits to it. Altogether, we defined five different types of saccades (Fig. 3A):

1. Between-category saccades: first visits

2. Between-category saccades: revisits

3. Between-exemplar saccades: first visits

4. Between-exemplar saccades: revisits

5. Within-element saccades

We expected these saccade types to differentially contribute to forming a coherent event

representation. Specifically, in our associative memory task, we tested episodic memory for

between-category associations (i.e., associations between exemplars from different categories)

within each event. The formation of these associations supposedly depends on the corresponding

between-category saccades during encoding. In contrast, associations established by between-

exemplar saccades (i.e., gaze transitions between associations incidental to the task) were not

tested and did not expect to predict subsequent memory performance. Thus, between-category and

between-exemplar saccades support associations between event elements that were or were not

critical to the success of the subsequent memory test, respectively.

Revisits, i.e., refixations of previously visited locations, have proven to be of particular interest in memory-related gaze behavior. Specifically, refixations can increase the sampling of information necessary for successful memory formation (Mirza et al., 2016; Turk-Browne, 2019; Voss et al., 2011) and recover information lost or missed during scanning (Gilchrist and Harvey, 2000; Körner and Gilchrist, 2008; Meghanathan et al., 2019; Tatler et al., 2005). Our previous coregistration studies have shown that refixations differ in the allocation of attention during saccade planning from ordinary fixations to locations visited once (Nikolaev et al., 2018) and that refixation planning is goal-dependent (Meghanathan et al., 2020). Moreover, we found that information acquisition at fixations to locations that are revisited later is distinct when compared to the other fixations (Nikolaev et al., 2022). Refixations may also play an important role in the formation of episodic memories. A recent study combining eye-tracking with intracranial EEG (Kragel et al., 2021) demonstrated that presaccadic hippocampal theta activity predicts whether an upcoming saccade will be a first visit or a revisit. Based on these previous findings, we expected EEG activity during revisits to be predictive of the successful binding of the event elements.

Additionally, we examined scrutinizing saccades within an exemplar (*within-element*), as it is well-known that the cumulative number of fixations on a visual stimulus predicts subsequent memory of that stimulus (Liu et al., 2017; Loftus, 1972; Olsen et al., 2016). This allowed us to examine two types of memory mechanisms: the first related to eye movements that bind the elements into a coherent event, and the second related to eye movements that sample visual information from individual elements.

2.5.3 EEG preprocessing

The goal of our study required fixation-related analysis of EEG in short epochs time-locked to fixation onsets in the encoding interval. This analysis used deconvolution modeling to correct for the overlapping effects of saccades on EEG. Deconvolution modeling involves time regression, which requires continuous EEG; thus, preprocessing was performed on continuous EEG. In addition, we analyzed EEG time-locked to the event onset during the entire 10-s encoding interval. The preprocessing for this analysis was the same as for the fixation-related analysis. We used the following preprocessing and deconvolution pipelines developed in our previous research (Nikolaev et al., 2022).

We down-sampled EEG signals to 256 Hz. The EEG and eye-tracking data were synchronized using the function *pop_importeyetracker* from the EYE-EEG extension. Saccades, fixations, event onsets, and bad eye-tracking intervals were inserted into the EEGLAB data structure.

The preprocessing pipeline included a series of cleaning functions from EEGLAB. The *pop_cleanline*

function removed power line noise using multi-tapering and a Thompson F-statistic. The

clean_artifacts function removed flat-line channels, low-frequency drifts, noisy channels, and short-

time bursts. This function is based on artifact subspace reconstruction (ASR), which compares the

structure of the artifactual EEG to that of known artifact-free reference data (Kothe and Jung, 2016).

The tradeoff between artifactual and retaining brain activities depends on the ASR parameter, which

we set to 20 according to the recommendations by Chang and colleagues (Chang et al., 2020).

Ocular artifacts were removed with the *OPTICAT* function (Dimigen, 2020). First, EEG was high-pass

filtered at 2 Hz to suppress large deviations from baseline due to summation of the corneo-retinal

artifacts during sequential eye movements. Next, 30-ms segments around saccade onsets were

obtained and re-appended to EEG to 'overweight' the contribution of the saccadic spike activity in

the EEG input to independent component analysis (ICA). Then, the ICA weights obtained after ICA

training on these filtered data were transferred to the unfiltered version of the same data. Finally,

the ratio between the mean variance of independent components during saccade and fixation

intervals was calculated. If the ratio was higher than 1.1, the corresponding independent

components were considered saccade-related and removed (Plöchl et al., 2012).

The automatic classifier (*pop_iclabel*) with the following settings (the probability ranges of ICs to be

an artifact and excluded are given in brackets): Brain [0 0], Muscle [0.4 1], Eye [0.9 1], Heart [0.05 1],

Line Noise [0.4 1], Channel Noise [0.4 1], Other [0.4 1] revealed independent components related to

remaining artifacts (Pion-Tonachini et al., 2019). On average, 16.9 (SD = 3.3, range 10-25)

components per participant were removed. EEG was re-referenced to average reference. The

removed channels (mean = 1.1, SD = 1 per participant) were interpolated with spherical spline

interpolation.

Theta and alpha power were extracted by applying the procedure from (Ossandón et al., 2020). We

filtered the continuous EEG with a low cut-off of 4 Hz and high cut-off of 7 for the theta band and

with a low cut-off of 8 Hz and high cut-off of 13 for the alpha band using the *pop_eegfiltnew*

function with default settings. We Hilbert-transformed the filtered signal and obtained the

instantaneous power. We normalized the EEG power at each sampling point to its ratio with the

respective channel mean power in a baseline interval. In the analysis of the encoding interval, the

baseline interval was set from -200 to 0 ms before the onset of the corresponding event. In the

fixation-related analysis, to avoid an interval of saccade execution just before fixation onset, the

baseline interval was set from -200 to -100 ms before the onset of the corresponding fixation

interval. The power values were log-transformed.

2.5.4 Oscillation detection

To test the presence of theta and alpha oscillations in the encoding interval we used the approach

for detection of transient oscillatory bursts, which separates periodic (oscillatory) from aperiodic

activity. This approach is based on comparison of the observed power spectrum with the aperiodic

background 1/f activity. It has been implemented in a number of methods, such as better oscillation

detection (BOSC) (Caplan et al., 2001) and its further developments, such as extended BOSC (eBOSC)

(Kosciessa et al., 2020) and the fBOSC (Seymour et al., 2022), which includes the 'fitting oscillations

and one over f' (FOOOF) algorithm (Donoghue et al., 2020). We searched for theta and alpha

oscillations in the 10-s encoding interval using the fBOSC toolbox for MATLAB (Seymour et al., 2022).

The fBOSC method was particularly suitable for our goals because it assumes that the 1/f fit is not

necessarily linear across all frequencies in log-log space, and allowed us to fit an aperiodic 'knee' in

the power spectrum. The presence of such a 'knee' is especially likely in data with pronounced theta

rhythmicity (Seymour et al., 2022), which is quite expected in our data.

To detect oscillations, fBOSC computes two thresholds: a power threshold estimated from the

aperiodic background spectrum, which we set at the 95th percentile of the theoretical probability

(chi-squared) distribution of power, and a duration threshold, which we set at 3 oscillatory cycles.

Since oscillations may distort the background spectrum, fBOSC excludes oscillatory peaks from its

calculation. To do this, fBOSC performs an initial 1/f fit, then iteratively models oscillatory peaks

above this background and removes them from the spectrum. Next, fBOSC models the 1/f power

spectrum as nonlinear by using a variable knee parameter (see Seymour et al. (2022) for details on

its estimation). We applied fBOSC to single EEG trials of the 10-s encoding interval. The results of the

single-trail modeling were averaged across all participants for low and high memory events

separately. To analyze the frequency distribution of the detected oscillations and their relationship

to memory, we used 'abundance', which is considered the most informative metric in burst analysis

(Kosciessa et al., 2020). Abundance is the duration of a rhythmic episode relative to the length of the

analyzed segment, which can range from 0 to 1, where 0 means no burst and 1 means a single

continuous burst.

2.5.5 Deconvolution

To overcome the problems associated with EEG-eye movement coregistration in unrestricted

viewing behavior described above, we used deconvolution modeling implemented in the Unfold

toolbox (Ehinger and Dimigen, 2019). The toolbox allowed us to correct overlapping EEG responses to sequential saccades in free viewing, as well as to account for several possible detrimental covariates. The deconvolution modeling is based on multiple regression, as traditional mass-univariate ERP modeling, which has been used for post-hoc control of multiple simultaneous discrete and continuous covariates (Smith and Kutas, 2015). However, mass-univariate modeling requires a prespecified analysis window. Therefore, when applied to correct the overlapping effects of unrestricted eye movements on EEG, it cannot account for the varying temporal overlap caused by variable fixation durations between adjacent eye movements (Dimigen and Ehinger, 2021). Deconvolution corrects for overlapping effects with variable fixation intervals by time-expansion of the design matrix in the duration of the entire (continuous) EEG recording (see Fig. 4 in Dimigen and Ehinger (2021) for an illustration of time-expansion). Furthermore, the correction method implemented in the Unfold toolbox accounts for the nonlinear effects of oculomotor covariates (e.g., saccade size) on EEG by the combination of linear deconvolution with nonlinear spline regression, as used in the generalized additive model (GAM) (Wood, 2017). The resulting time series of regression coefficients, or betas, represent non-overlapping brain responses for each predictor and correspond to subject-level ERP waveforms in traditional ERP analysis. Thus, the outputs of the toolbox are beta coefficients, which represent partial effects for the predictors of interest, adjusted for the covariates included in the model.

These covariates included the low-level oculomotor effects of eye movement characteristics and the ordering factors in time courses of an encoding trial, a block, and an entire experiment. Specifically, we considered the low-level oculomotor characteristics of fixation duration and saccade size and direction, which are known to affect the fixation-related EEG (Dimigen et al., 2011; Nikolaev et al., 2016). Furthermore, we considered three ordering factors. Since the time course of a trial with multiple eye movements may affect the fixation-related EEG activity (Fischer et al., 2013; Kamienkowski et al., 2018), we considered the rank of a fixation in relation to all other fixations that occurred during the entire encoding interval of an event. The rank was assigned to each fixation as an ordinal number of this fixation in the encoding interval. Behavioral results (see below) revealed a nonlinear effect of event order within a block and a strong linear effect of block order on subsequent memory. We accounted for these factors by including them in the deconvolution model.

The Unfold toolbox involved the following procedures, which were sequentially applied to the continuous EEG of each participant.

480   To account for the experimental conditions of interest and possible confounding factors in the same

481   model, we used the following Wilkinson notation of the model formula:

483   ```
Fixation: y ~ 1 + cat(MemoryPerformance) + rank + iBlock + spl(iEventPerBlock,4) +
484   spl(duration,5) + spl(sac_amplitude,5) + circspl(sac_angle,5,-180,180)
485   Event: y ~ 1
486   Levels of MemoryPerformance: 'other' (reference level), 'low memory', 'high memory'
```

488    This formula includes the multiple effects at fixations, as well as the onset of the event screen.

489    Specifically, for fixations, the formula considers the fixation onset (y ~ 1, i.e., the intercept), a

490    categorical predictor of the subsequent memory effect ('MemoryPerformance') and continuous

491    predictors: fixation rank ('rank'), a block order in an experiment ('iBlock'), an event order in a block

492    ('iEventPerBlock'), fixation duration ('duration'), size ('sac_amplitude') and direction ('sac_angle') of

493    the preceding saccade. In this example formula, the 'MemoryPerformance' predictor includes three

494    levels: 'other' fixations (see Eye movement analysis above), 'low memory' and 'high memory' (see

495    Behavioral analysis above). We used treatment (dummy) coding with 'other' fixations as the

496    reference level (intercept) relatively to which other levels (i.e., 'low memory' and 'high memory') of

497    the 'MemoryPerformance' predictor were estimated. The inclusion of 'other' fixations allowed us to

498    account for the overlapping effects on EEG produced at the latencies of 'other' eye movements. To

499    effectively account and control for these overlapping effects, we need to input in the model all

500    fixations that occurred in the experiment, without exception. Thus, the "other" fixations, together

501    with the fixations assigned to low and high memory levels, were essential levels of the

502    'MemoryPerformance' predictor. However, because the "other" fixations cannot explain memory for

503    the event, they were never used in further comparisons. Since fixation duration, and saccade size

504    and direction have nonlinear effects on EEG (Dimigen and Ehinger, 2021; Nikolaev et al., 2016), we

505    modeled them with a basis set of five spline predictors (circular splines were used to model saccade

506    direction). Since we found that the event order in the block has a nonlinear effect on the memory

507    score (Fig. 2B), and we assumed its nonlinear effect on the EEG, we modeled it also with spline

508    predictors. Spline knots were placed on the percentiles of the covariates for each participant. For

509    events, we considered only their onsets (y ~ 1), i.e., the intercept that described the overall shape of

510    EEG evoked by the event screens. In further analyses, we modified the 'MemoryPerformance'

511    predictor in the formula by inserting predictor levels depending on the analysis goals, as specified

512    below in each section of Results. All other predictors were constant in all analyses.

514    To recover isolated EEG responses (betas) to each fixation and to each event screen best explaining

515    the continuous EEG, we created a design matrix (Fig. S3A) and time-expanded it in a window

516  between −200 and +400 ms around fixation and event onsets. This length of the time window was

517  motivated by the need to keep the number of iterations for fitting the deconvolution model before

518  model convergence reasonably small (e.g., <400) to avoid overfitting. The number of iterations

519  depends on the number (and quality: spline or not) of predictors and the number of sampling points.

520  At the pilot stage of the project, our experimentation with the window length using the selected

521  predictors showed that the length of -200+400 ms is optimal. Time expansion involved duplicating

522  and shifting the predictors of the mass univariate linear regression design matrix for each time lag.

523  The time-expanded design matrix (Fig. S3B) spanned the duration of the entire EEG recording. As a

524  result, instead of estimating multiple linear models for all time points, we estimated one linear

525  model, simultaneously estimating all fixation and event betas. The variable temporal distance

526  between sequential fixations allowed the separation of the overlapping effects.

528  Before fitting the model, we excluded irrelevant intervals from modeling by filling them with zeros in

529  the time-expanded design matrix. Filling with zeros preserved the timing of continuous EEG. If we

530  would remove these intervals completely, we might remove along with them the eye movements

531  that produce overlapping effects on clean EEG parts used in the analyses of interest. Specifically,

532  these intervals corresponded to the distractor task, memory tests, intervals between events, breaks

533  between blocks, and bad eye-tracking intervals, leaving only intervals of free viewing exploration of

534  event screens during encoding. We fitted the deconvolution model to each of the 31 electrodes

535  using the iterative Least Squares Minimal Residual algorithm for sparse design matrices (Fong and

536  Saunders, 2011).

538  The raw beta coefficients obtained for categorical predictors of interest after fitting the

539  deconvolution model represent the "pure" effects of the predictors, which could be already

540  interpreted and compared statistically. The effects could be understood as a rough analog of

541  difference waves relative to the reference level of the categorical predictor. Consequently, the

542  curves of the "pure" effects have highly wiggly shapes, similar to the difference waves, oscillating

543  around zero, as illustrated in Fig. S4. This makes it difficult to compare deconvolution results with

544  the waveforms in the traditional EEG literature. To solve this problem, we need to add the model

545  intercept onto the difference wave. However, in the case of nonlinear modeling with splines,

546  calculating the intercept is not straightforward as it would be in the case of linear models.

547  Specifically, in nonlinear modeling with splines, the classical model-intercept is ill-defined, since

548  there is no clear reference level for the nonlinear basis set (the set which transforms predictor

549  values by weighting beta coefficients). To resolve this issue, we estimated all nonlinear effects at

550  their respective mean predictor levels and added these estimates to the beta coefficients of the

categorical predictors. For example, if we aim to adjust our memory effect to saccade amplitude, and the mean saccade amplitude for a given participant is 7 dva, we evaluated the nonlinear effect at 7 dva and added the resulting value to the beta coefficients of the memory effect. This procedure ensures that all levels of the memory effect are visualized at the same estimated saccade amplitude of 7 dva. The described solutions are implemented in the functions *uf_predictContinuous* and *uf_addmarginal* of the Unfold toolbox, which we applied to the raw beta coefficients for each participant. The resulting waveforms then became comparable to the participant-level averages in traditional EEG analysis.

### 2.5.6   Statistical analysis

Unless indicated otherwise, we used a repeated measures ANOVA for the statistical analysis. To handle violation of sphericity, we applied the Huynh-Feldt correction for p-values associated with two or more degrees of freedom. Tukey HSD test was used for post-hoc analyses. The statistical analyses were performed with STATISTICA 10 (StatSoft Inc., Tulsa, OK, USA) and R (R Core Team, 2020).

Data analysis involved three major stages. First, we aimed to examine whether the theta and alpha activity obtained from EEG signals coregistered with eye movements in unrestricted viewing would yield results consistent with the previous episodic memory studies with fixed stimulus presentation (Hanslmayr et al., 2016). Since theta and alpha activity represent distinct functions of episodic memory formation and have complex relationships, we analyzed them in separate ANOVA models. We segmented theta and alpha EEG power before deconvolution into 10-s trials relative to the onsets of the encoding screens (events). We compared the mean EEG power in five 2-s time bins of 10-s trials between events with low and high subsequent memory performance. Furthermore, we detected the presence of theta and alpha oscillations in the encoding interval with the fBOSC method (Seymour et al., 2022). We considered the duration ('abundance') of theta and alpha oscillations for low and high subsequent memory. In these and all subsequent EEG analyses, to capture effects with potentially different topographical distributions, we grouped electrodes in eight regions of interest (ROIs). Specifically, we selected three electrodes around the landmark electrodes of the International 10-20 System of Electrode Placement: F3, F4, C3, C4, P3, P4, O1, O2 (the inset in Fig. 4). These groups of electrodes constituted eight ROIs over frontal, central, parietal and occipital brain regions of the left and right hemisphere (FL, FR, CL, CR, PL, PR, OL, OR). For each ROI, we averaged power waveforms across three electrodes since regional averaging of electrodes provides a more reliable estimate of activity in a region than a single measurement (Dien and Santuzzi, 2005). Eight ROIs were symmetrically about the sagittal axis and systematically distributed over the scalp.

586  Thus, these analyses allowed us to evaluate memory-related changes of theta and alpha power over

587  the entire encoding interval.

589  Second, we evaluated the general characteristics of fixation-related power extracted from the

590  deconvolved EEG. We estimated the theta and alpha power at the first peak of their cycle about 100

591  ms after the fixation onset (see 4.3 for justification of such selection). We averaged the power values

592  40 ms before and after the peak, that is, between 60 and 140 ms after the fixation onset. To

593  evaluate the time course of fixation-related EEG power during the 10-s encoding interval, we divided

594  EEG power related to successive eye movements into three bins according to fixation rank. We

595  compared memory-related changes of theta and alpha power across rank bins in eight ROIs. Thus,

596  this analysis was analogous at the fixation level to the first analysis at the event level.

598  The third analysis was devoted to the main goal of our study – to compare fixation-related EEG *after*

599  the five types of saccades described in section 2.5.2. We performed three separate analyses on each

600  saccade type because not all conditions were matched for all saccade subtypes. Specifically, we

601  could distinguish between First visits and Revisits for Between-category and Between-exemplar

602  saccades but not for Within-element saccades. All analyses compared the fixation-related theta and

603  alpha power between low and high memory for the same time window as in the second analysis and

604  eight ROIs. This analysis provided a detailed picture of associations during the formation of coherent

605  episodic memory at the fixation level. To emphasize findings directly relevant to the goals of our

606  study, we report effects and interactions in the Results section only when they involved the factors

607  of memory performance (High vs. Low) and gaze transition status (First visits vs. Revisit). Complete

608  statistical results, including effects and interactions between ROIs and hemispheres, are reported in

609  Table S2 in Supplementary materials.

611  # 3  Results

612  ## 3.1  Behavioral results

613  Average memory performance (regardless of confidence) across 28 participants was 68.6 ± 8.33 %

614  (mean ± SD). The memory score weighted by confidence for the low memory condition was 27.8 +/-

615  6.8 and for the high memory condition 48.8 +/- 9 (mean +/- SD). Memory performance changed in

616  the time course of a block and experiment (Fig. 2). We evaluated these changes with a repeated

617  measures ANOVA on the memory scores weighted by confidence with the factors of Event (nine

618  levels) and Block (six levels). There were significant main effects of Event ($F_{(8, 216)}$ = 3.7, p = .004, ε

619  = .62) and Block ($F_{(5, 135)}$ = 9.8, p < .001, ε = .95)), but no interaction between them. To characterize

620    the effects of Event and Block, we used planned comparisons with the linear and quadratic

621    contrasts. The contrast analysis revealed significant effects for the linear contrast for Block ($F_{(1, 27)}$

622    = 31.2, $p < .001$) and for the quadratic contrast for Event ($F_{(1, 27)}$ = 16.8, $p < .001$), but not for the

623    linear contrast for Event ($F_{(1, 27)}$ = 0.7, $p = .4$). These results indicate a prominent learning effect

624    during the experiment, as well as possible primacy and recency effects in the time course of a block.

625    To evaluate the contribution of confidence to memory scores, we repeated the above analyses on

626    memory scores that were not weighted by confidence. The results were qualitatively the same

627    (Supplementary materials and Fig. S2), indicating that confidence contributes little to our measure of

628    memory performance.

630    To compare encoding of different categories, we computed three separate memory scores

631    (weighted by confidence), considering only paired associations within an event: faces – places, faces

632    – objects, places – objects. Then we divided these memory scores according to low and high memory

633    performance in the entire event (as in the EEG analysis). We applied a repeated-measures ANOVA

634    on the memory scores of category associations with factors of Memory (Low vs. High) and

635    Association type (faces – places, faces – objects, places – objects). We found effects of Memory ($F_{(1,}$

636    $_{27)}$ = 375, $p < .001$) and Association type ($F_{(2, 54)}$ = 10.5, $p = .001$, $\varepsilon = .66$), and no interaction ($F_{(2,}$

637    $_{54)}$ = 0.5, $p = .63$) (Fig. 2C). The post-hoc test showed similar results for low and high memory

638    performance: no difference in memory scores between associations of faces – places and faces –

639    objects (both $p > .06$) and significant differences in memory scores between associations of faces –

640    places and places – objects (both $p < .001$), as well as between associations of faces – objects and

641    places – objects (both $p < .021$). These results statistically confirmed the low recall of faces reported

642    by most participants in post-experiment interviews. More importantly, differences in performance

643    across categories were equally distributed between the low and high memory events used as

644    conditions in the EEG analyses and, therefore, could not bias the EEG results.

646 Fig. 2. Memory performance weighted by confidence in the time course of an experiment (A) and a block (B).

647 The thin lines indicate significant linear (A) and quadratic (B) fits. (C) Memory performance weighted by

648 confidence for paired associations between particular categories within an event and divided according to low

649 and high memory performance in the entire event (as in the EEG analysis).

651 ## 3.2 Eye movement results

652 The dynamics of eye movement behavior during the 10-s encoding interval is illustrated in Fig. 3B by

653 the fixation rank probability density for each type of preceding saccade. At first, participants tended

654 to visit all categories quickly with between-category saccades. Then, between-exemplar saccades

655 became predominant. The number of within-element saccades was relatively stable throughout the

656     whole encoding interval. Towards the latter part of the interval, between-category revisits became

657     more frequent.

659     To determine if, and to what degree, the saccade categories predicted subsequent memory

660     performance, we analyzed the relationship between the cumulative number of saccades (in each

661     category) within an event and subsequent memory performance for that event. The between-

662     category saccades for first visits were not analyzed since there were always 3 per event. We

663     computed the number of the remaining four saccade types for each event and pooled these

664     numbers with the corresponding event memory scores over 28 participants. Since the number of

665     saccades for different types varied quite widely (Fig. 3C), we could not establish a single categorical

666     predictor of the number of saccades for all saccade types to include them in the same ANOVA

667     design. Therefore, we applied a one-way ANOVA on the event memory score with the number of the

668     saccades as the predictor for each saccade type separately (bins with less than 20 values at the left

669     and right ends of the distributions were excluded). The results showed that memory for the event

670     significantly increased with the cumulative number of between-category revisits ($F_{(8, 1392)} = 3.8$, $p$

671     $< .001$) and with the cumulative number of within-element saccades ($F_{(14, 1360)} = 3.1$, $p < .001$) (Fig.

672     3C).

674     To be successful in the retrieval task, association from all three categories need to be considered

675     during event encoding. Two different gaze patterns can achieve this: (1) engaging in active gaze

676     transitions between each of the three categories, or (2) engaging in active gaze transitions between

677     only two of the categories, i.e., when the association between the first visited category and the last

678     visited category is inferred without an active gaze transition between the categories. In the present

679     data set, gaze transitions involving only two categories characterized approximately one-third of the

680     events. These events were distributed between participants approximately normally (the Lilliefors

681     test could not reject a null hypothesis of normality of the distribution: $d = 0.12$, $p > .2$). To evaluate

682     whether the two eye movement patterns are associated with different subsequent memory

683     performance, we compared event memory scores between these pattern types with a paired t-test.

684     We found no difference in memory between the two patterns ($t_{(27)} = 0.05$, $p = .96$). The memory

685     score and the number of events involving only two categories did not correlate across participants ($r$

686     $= .25$, $p = .2$). These findings suggest that event memory does not require gaze transitions between

687     all three categories.

689     We also compared the rate of eye movements between low and high subsequent memory

690     performance. The eye movement rate normalized to the total number of saccades for each

691 participant was much higher in the low (.53, .025) than high (.47, .025) (mean, SD per participant)

692 memory condition (t(27) = 5.7, p < .001).

695 Fig. 3. A: an example scanpath illustrating five types of saccades. B: Probability density estimation of the

696 fixation rank in a 10-s event interval pooled across 28 participants. The vertical lines indicate mean ranks of the

697 corresponding fixation type. C: Dependence between the occurrence frequency of saccade types and the

698 memory score. Significance of the dependence obtained with one-way ANOVA is indicated in each panel. Error

699 bars indicate standard errors of means across 28 participants.

701 In sum, the eye movement results show that between-category saccades (revisits) and within-

702 element saccades predict subsequent memory performance, whereas between-exemplar saccades

703    do not. The associative memory task determined these dependencies, which required strong

704    associations between the categories, whereas associations between the exemplars of a category

705    were less important to the task. Within-element saccades were necessary for acquiring sufficient

706    visual information about each image, a prerequisite for establishing a strong representation of each

707    element, and for recognizing them during retrieval.

709    ## 3.3   EEG results at the event level
710

711    ### 3.3.1   Subsequent memory effects at the event level
712

713    To assess how theta and alpha power at encoding predicts subsequent memory at the time scale of

714    the entire encoding interval, we segmented the EEG power (after removing all artifacts but before

715    deconvolution) from -0.2 to 10 s relative to the event onset and baseline corrected it to the interval

716    from – 0.2 to 0 s before the event onset (Fig. 4A, B). We smoothed the individual power waveforms

717    with a Savitzky-Golay filter, using filter order 3 and filter length 201 sampling points (R package

718    'signal'). Events were split between low and high memory performance (the number of low memory

719    events was 28.5 (1.35); the number of high memory events was 25.1 (1.74) (mean (SD) per

720    participant). To investigate the modulation across a 10-s encoding interval, we created five time-bins

721    of 2 s each and averaged the power values in these bins (Fig. 4C, D). We applied a repeated-

722    measures ANOVA on the power values with the factors of Memory (Low vs. High), ROI (Frontal,

723    Central, Parietal, Occipital), Hemisphere (Left vs. Right), and Time (5 bins). For theta power, we

724    found an interaction between Memory and Time ($F_{(4, 108)} = 4.9$, $p = .001$, $\varepsilon = .99$). Post-hoc test

725    revealed that this interaction occurred due to higher power for the high than low memory events for

726    the 4th and 5th time bins (all $p < .02$), i.e., between 6 and 10 s at the end of the encoding interval (Fig.

727    4C).

729    For alpha power, there was a clear trend towards decreased power in high-memory versus low-

730    memory events, most noticeably over the occipital areas (Fig. 4B, D), which, however, did not reach

731    significance. The change of alpha power over time appears to be nonlinear. To test the properties of

732    this change statistically, we applied linear and quadratic contrasts to the time bins as planned

733    comparisons. For this analysis, we used only the data from the occipital ROIs, where the memory

734    effect on alpha power was most prominent (Fig. 4B). We found that the quadratic fit more

735    adequately describes the memory effect on the alpha activity over time, revealing a significant

736    difference between high and low memory events ($F_{(1, 27)} = 5.0$, $p = .03$), than the linear fit, which

737    did not reveal a significant effect between high and low memory events ($F_{(1, 27)} = 0.5$, $p = .47$). For

738  completeness, we also applied the linear and quadratic contrasts to the time bins for theta power

739  (all ROIs). Here the results were the opposite: there was a significant difference between low and

740  high memory events for the linear fit (F(1, 27) = 17.5, p < .001) but not for the quadratic fit (F(1, 27) =

741  2.8, p = .1).

743  Fig. 4. Theta (A) and alpha (B) power time-locked to the event onset during a 10-s encoding interval for low

744  and high subsequent memory performance. The power is grand averaged across 28 participants in 8 ROIs

745  selected as indicated in the inset between panels A and B. (C – D) The time course of the grand-averaged theta

746  (C) and alpha (D) power waveforms and the mean-error plots (mean +/- SEM) for five 2-s time bins. Significant

results of the post-hoc test for the interaction between Memory and Time bins are indicated with asterisks. In (C) the straight lines indicate a linear fit, whereas in (D) the curved lines indicate a quadratic fit, best reflecting the power changes over time in the respective bands. (E) Log-log plot of the EEG power spectrum averaged across 28 participants over the left parietal area. (F – G) The duration of the rhythmic episode relative to the length of the analyzed segment ('abundance') averaged across 28 participants over the left parietal area for low and high memory performance. (F) (top) Abundance for the frequency range 3-20 Hz. (bottom) Abundance in the theta band (close-up of the plot above). The arrows indicate the abundance increase for high memory performance. (G) The mean-error plots (mean +/- SEM) of abundance in the theta band for 8 ROIs.

In sum, EEG analysis of the entire encoding interval revealed the expected neural signatures, where higher subsequent memory performance was associated with increased theta synchronization and alpha desynchronization during encoding (Hanslmayr et al., 2016). Consistent with the idea of a gradual buildup of an episodic memory representation, theta increase was more prominent towards the end of the encoding interval. Theta increase was accompanied by alpha decrease, whose relationship with subsequent memory performance over time was nonlinear and best described by a quadratic fit.

### 3.3.2   Detection of theta and alpha oscillations at the event level

The finding that subsequent memory performance is associated with theta and alpha activity raises the question of whether memory-related changes in spectral pattern result from neural oscillations pertinent to encoding (Hanslmayr et al., 2016) or instead reflect increases in broadband activity or shifts in spectral tilt. We detected theta and alpha oscillations using the fBOSC method, which considered the possible nonlinearity of the 1/f power spectrum (Seymour et al., 2022). Fig. 4E shows the averaged power spectrum, the aperiodic 1/f component, and the FOOOF model fit for the left parietal area for high memory events (the results for all ROIs and both events are shown in Fig. S5A). The grand-averaged results were about the same for low- and high-memory events. The power spectrum had a clear peak in the alpha range and a deviation from the aperiodic fit at low frequencies.

Although the grand-averaged power did not reach the power threshold, multiple oscillation episodes were detected at the single-trial level. We compared the duration of these oscillations ('abundance') between low and high memory. Fig. 4F shows the abundance averaged across participants for the left parietal area for the frequency range 3-20 Hz (top) and for the close-up in the theta range (bottom). The abundance for all ROIs is shown in Fig. S5B. The arrows point to the frequency interval

of about 5-6 Hz, where the abundance was larger for the high than low memory events. The

abundance at the peak alpha frequency was visually larger for the low than high memory events. To

test these observations statistically, we averaged the abundance within the theta and alpha bands

for each participant and analyzed it using a repeated-measures ANOVA with the factors of Memory

(low vs. high), ROI (frontal, central, parietal, occipital), and Hemisphere (left vs. right). For the theta

band, we found main effects of Memory ($F(1, 27) = 4.8$, $p = .038$) and a tendency for an interaction

between Memory, ROI, and Hemisphere ($F(3, 81) = 2.3$, $p = .086$, $\varepsilon = .98$). Fig. 4G suggests that this

interaction tendency may occur because of the increased abundance over the parietal regions for

high memory events. For the alpha band, there was no Memory effect ($F(1, 27) = 0.4$, $p = .56$).

Thus, the abundance analysis shows that a long duration of theta oscillations throughout the

encoding interval predicts high memory performance. The duration of alpha oscillations is much

longer than the duration of theta oscillations, as indicated by the prominent increase of abundance

in the alpha band (Fig. 4F). But the duration of alpha oscillations throughout the encoding interval is

not predictive of memory performance. This, however, does not exclude their involvement in

memory processes at the fixation level.

## 3.4   EEG results at the fixation level

Since the deconvolution framework is a novel approach to free-viewing behavior in tasks of episodic

memory formation, we first evaluated how it changes the power waveforms (see Supplementary

materials and Fig. S6 for details).

### 3.4.1   Subsequent memory effects at the fixation level

Before analyzing the data for the different saccade categories, we wanted to assess if the EEG

findings at the event level could be reproduced at the fixation level. Thus, we tested the subsequent

memory effect on fixation-related theta and alpha power by contrasting the fixation-related EEG

overall eye movements within an event. We divided the memory scores into three bins according to

low, medium, and high memory events. We divided the fixations into three bins according to fixation

rank. In the formula for deconvolution modeling, we used ten levels of categorical predictors: all

possible combinations of three memory and three rank bins and 'other'. The number of fixations

used in the deconvolution modeling of each memory and rank bin condition is presented in Table S1.

We applied a repeated-measures ANOVA on the EEG power averaged in the window from 60 to 140

ms after the fixation onset with the factors of Memory (low vs. high, we excluded the medium

818    memory bin to increase the memory contrast), Rank (3 bins), ROI (frontal, central, parietal, occipital)

819    and Hemisphere (left vs. right). For theta power, we found interactions between Memory and Rank

820    $(F(2, 54) = 3.6, p = .038, \varepsilon = .96)$ (Fig. 5B) and between Memory and ROI $(F(3, 81) = 3.6, p = .023, \varepsilon =$

821    $.86)$ (Fig. 5C). Post-hoc tests revealed that the Memory-Rank interaction occurred due to higher

822    power for the high than low memory in rank bin 3 $(p = .02)$. Fig. 5A shows power waveforms for this

823    rank bin. The Memory-ROI interaction occurred due to higher power for the high than low memory

824    over the frontal $(p = .006)$ and central $(p = .001)$ areas. The difference maps indicate that this effect

825    was largest in rank bin 3 (Fig. 5D).

827    For alpha power, we found an interaction between Memory, Rank, and Hemisphere $(F(2, 54) = 5.0, p$

828    $= .01, \varepsilon = .98)$ (Fig. 5E-G). Post-hoc tests revealed that the interaction occurred due to a power

829    decrease for the high than low memory in rank bin 2 (Fig. 5F) over the left hemisphere $(p = .002)$; the

830    similar trend in rank bin 3 did not reach post-hoc significance $(p = .14)$. There was also an interaction

831    between Memory, Rank, ROI, and Hemisphere $(F(6, 162) = 2.4, p = .04, \varepsilon = .84)$, but the post-hoc

832    tests showed no significant differences.

835 Fig. 5. Theta and alpha fixation-related grand-averaged (N=28) power in the analysis of three fixation rank bins.

836 (A) Theta power in 8 ROIs for rank bin 3. (B – C) The mean-error (mean +/- SEM) plots of the theta power in the

837 interval from 60 to 140 ms after the fixation onset. (B) Memory-Rank interaction. (C) Memory-ROI interaction.

838 (D) The theta power difference maps: high minus low memory. (E) Alpha power in 8 ROIs for rank bin 2. (F) The

839 Memory-Rank interaction over the left hemisphere. (G) The alpha power difference maps: high minus low

840 memory conditions. Significant results of post-hoc tests are indicated with asterisks.

842 To summarize, the fixation-related EEG results corroborate and extend what was found at the event

843 level, i.e., theta increase and alpha decrease during encoding covary with subsequent memory

844 performance. These effects were also more prominent toward the end of the encoding interval.

845 Next, we analyzed EEG power at fixations following the five saccade types of interest.

847 3.4.2    Subsequent memory effects across saccade types

849 We focused on EEG power in the fixation intervals succeeding the five saccade types (Fig. 3A). For

850 this purpose, we specified ten levels of categorical predictors in the deconvolution model formula:

851 the five saccade types x memory performance (low, high). The number of fixations that were used in

852 the deconvolution modeling of each saccade and memory condition is presented in Table 1. "Other"

853 fixations were the reference level, as in the previous model.

855 Table 1. The number of fixations used in the deconvolution modeling of each saccade type per

856 participant: mean (range, SD) across 28 participants.

| | low memory | high memory |
|---|---|---|
| Between-category saccades: first visits | 83 (57-96, 8.6) | 69 (57-78, 7.1) |
| Between-category saccades: revisits | 82 (28-158, 35.4) | 74 (18-156, 32.6) |
| Between-exemplar saccades: first visits | 64 (39-83, 12.3) | 55 (37-75, 11.9) |
| Between-exemplar saccades: revisits | 115 (47-214, 51.7) | 95 (31-177, 44.1) |
| Within-element saccades | 187 (101-321, 67.4) | 164 (88-288, 60.5) |

858 *3.4.2.1    Between-category saccades*

859 Here, we examined EEG power in the fixation intervals succeeding the between-category saccades

860 that we predicted to support tested associations between categories (Fig. 6A). We applied a

861 repeated measures ANOVA on fixation-related theta and alpha power with the factors of Gaze

862 transition status (First visits vs. Revisit), Memory (Low vs. High), ROI (Frontal, Central, Parietal,

863 Occipital), and Hemisphere (Left vs. Right). For theta power, we found a significant main effect of

864 Gaze: theta power was higher during first visits than revisits ($F(1, 27) = 5.0$, $p = .03$). We also found

865 an interaction between Gaze and ROI ($F(3, 81) = 4.2$, $p = .01$, $\varepsilon = .94$). More importantly, we found an

866 interaction between Gaze, Memory, and Hemisphere ($F(1, 27) = 11.8$, $p = .002$) (Fig. 6B-E). Post-hoc

867 tests revealed that this interaction occurred due to higher power for the high than low memory for

868 revisits over the left hemisphere ($p < .001$) (Fig. 6E). For alpha power, no main effects of Memory

869    (F(1, 27) = 3.1, p = .09) or Gaze (F(1,27) = 1.7, p = .2) emerged (nor an interaction between them,

870    F(1, 27) = 0.5, p = .5).

872    Thus, theta power increase over the left hemisphere after between-category revisits predicts

873    subsequent event memory performance. This theta memory effect is consistent with the eye-

874    movement analyses, where subsequent memory performance improved with the cumulative

875    number of between-category revisits during encoding (Fig. 3C).

878    Fig. 6. Fixation-related EEG after between-category saccades. (A) An example scanpath illustrating a subset of

879    between-category (colored) saccades preceding the fixation intervals used in the EEG analysis. The grey arrows

are saccades not included in this analysis. (B, D) The grand-averaged (N=28) theta power over the left hemisphere for first visits (B) and revisits (D). (C) The power difference maps: high minus low memory conditions. (E) The mean-error (mean +/- SEM) plot of the power over the left hemisphere in the interval from 60 to 140 ms after the fixation onset for first visits and revisits. Significant results of post-hoc tests are indicated with an asterisk.

### 3.4.2.2 Between-exemplar saccades

Next, we examined EEG power in the fixation intervals succeeding between-exemplar saccades, i.e., gaze transitions establishing element associations that were not tested in the subsequent memory test (Fig. 7A). As in the previous analysis, we used a repeated measures ANOVA with a 2x2 design: Gaze transition status (First visit vs. Revisit) x Memory (Low vs. High), ROI (Frontal, Central, Parietal, Occipital), and Hemisphere (Left vs. Right). For alpha power, we found the main effects of Memory (Fig. 7B-D), where power was lower for high than low memory performance ($F(1,27) = 5.1$, $p = .03$). For theta power, no significant effects of Memory ($F(1,27) = 0.3$, $p = .6$) or Gaze ($F(1,27) = 2.1$, $p = .16$) emerged (nor an interaction between them $F(1,27) = 1.5$, $p = .24$).

As expected, theta power after between-exemplar saccades did not predict subsequent event memory performance for the tested between-category associations. This is consistent with the eye-movement result, where the cumulative number of between-exemplar saccades did not influence subsequent memory performance (Fig. 3C). The effect of lower alpha power for high memory was unexpected, but may reflect other mechanisms, apart from binding *per se*, that are important for optimal memory formation.

904 Fig. 7. Fixation-related EEG after between-exemplar saccades. (A) An example scanpath illustrating a subset of

905 between-exemplar (colored) saccades preceding the fixation intervals used in the EEG analysis. The grey

906 arrows are saccades not included in this analysis. (B) The grand-averaged (N=28) alpha power for fixations

907 after between-exemplar saccades. (C) The difference alpha power maps: high minus low memory conditions.

908 (D) The mean-error (mean +/- SEM) plot of the alpha power in the interval from 60 to 140 ms after the fixation

909 onset.

911 *3.4.2.3    Within-element saccades*

912 Finally, we analyzed theta and alpha power in the fixation intervals after within-element saccades

913 (Fig. 8A). For this saccade type, we did not distinguish between first visits and revisits, so the ANOVA

914 design included three factors: Memory (Low vs. High), ROI (Frontal, Central, Parietal, Occipital), and

Hemisphere (Left vs. Right). For theta power, we found an interaction between Memory and ROI ($F_{(3, 81)}$ = 3.4, p = .025, ε = .95) (Fig. 8B, C). Post-hoc tests revealed that this interaction occurred due to higher theta power for high than low memory over the frontal (p = .002) and central (p = .03) areas (Fig. 8D). For alpha power, we found an interaction between Memory and Hemisphere ($F_{(1, 27)}$ = 4.9, p = .035) (Fig. 8E-F). Post-hoc tests revealed that this interaction occurred due to higher alpha power for high than low memory over the right hemisphere (p = .01) (Fig. 8G).

Fig. 8. Fixation-related EEG after within-element saccades. (A) An example scanpath illustrating a subset of within-element (colored) saccades preceding the fixation intervals used in the EEG analysis. The grey arrows are saccades not included in this analysis. The grand-averaged (N=28) theta (B) and alpha (E) power for 8 ROIs. The theta (C) and alpha (F) power difference maps: high minus low memory conditions. (D, G) The mean-error (mean +/- SEM) plot of the power in the interval from 60 to 140 ms after the fixation onset. (D) Memory-ROI interaction for theta power. (G) Memory-Hemisphere interaction for alpha power. Significant results of post-hoc tests are indicated with asterisks.

Thus, the theta power related to within-element saccades predicts subsequent memory performance similarly as for the between-category revisits. Moreover, for both saccade types, subsequent memory increased as a function of the cumulative number of saccades during encoding (Fig. 3C). Importantly, however, the topographies of the theta memory effects for the two saccade types were *distinct*: a left central maximum for between-category revisits (Fig. 6C) and a frontal maximum for within-element saccades (Fig. 8C), suggesting dissociable neural mechanisms.

To test the observed topographical dissociation of the two theta effects, we extracted the 'within-element effect' as the difference between theta power in high and low memory performance for fixations after within-element saccades. We extracted the 'between-category effect' as the difference between theta power in high and low memory performance for fixations after between-category revisits. A repeated-measures ANOVA on the theta power difference with factors of Theta effect ('within-element' vs. 'between-category'), ROI (Frontal, Central, Parietal, Occipital), and Hemisphere (Left vs. Right) revealed an interaction between Theta effect and Hemisphere ($F(1, 27) = 9.0$, $p = .006$). The post-hoc test revealed higher power for the between-category effect over the left hemisphere ($p = .03$) (Fig. 9). This finding indicates topographical differences for the two theta effects and suggests that they are associated with distinct neural mechanisms operating in the same frequency band.

In sum, the correspondence of theta and eye movement results for between-category saccades and within-element saccades, which may be based on distinct mechanisms, indicates a diversity of relationships between theta activity, eye movements, and episodic memory formation, as will be discussed below.

957 Fig. 9. Comparison of the 'within-element' and 'between-category' theta memory effects. (A) The mean-error

958 (mean ± SEM) plot of the power difference in the interval from 60 to 140 ms after the fixation onset.

959 Significant result of the post-hoc test is indicated with an asterisk. (B) The power difference map: 'between-

960 category' effect minus 'within-element' effect.

## 4   Discussion

965 The present study set out to advance our current understanding of the relationship between eye

966 movements and episodic memory encoding. In particular, we targeted the neural mechanisms that

967 mediate the formation of coherent episodic memories during consecutive saccades to event

968 elements during unrestricted viewing behavior. To this end, we applied a state-of-the-art analytical

969 approach to EEG coregistered with eye movements during a free-viewing episodic memory task. Our

970 approach provided a unique possibility to capture neural encoding mechanisms across eye

971 movements at the level of gaze fixations while overcoming the confounding effects of sequential

972 saccades on brain activity. We identified neural signatures that are associated with the buildup of

973 episodic memories as a function of visual sampling behavior. The encoding of associations between

974 event elements was accompanied by simultaneous modulation of eye movements and theta activity

975 that were predictive of subsequent memory. This modulation likely reflects the binding of event

976 elements into a coherent representation across eye movements. In addition, a subsequent memory

977 effect was also observed for theta and alpha activity related to small scrutinizing saccades within

978 elements. Although the association between exemplars within categories was incidental to the task,

979    alpha activity related to between-exemplar saccades also proved predictive of later episodic

980    remembering. Overall, we have identified the engagement of distinct neural mechanisms that may

981    be the essential building blocks in forming episodic memory during unrestricted viewing.

983    Theta and alpha activity are considered to serve important but distinct functions of episodic memory

984    formation: binding individual elements together into a whole (theta) and representing individual

985    elements (alpha) (Hanslmayr et al., 2016). But it has not yet been established how these

986    mechanisms are engaged when coherent episodic memories are formed in a piecemeal procedure

987    across eye movements. In the current study, we aimed to fill this gap by tracking how these neural

988    mechanisms subserve the buildup of episodic memories during unrestricted viewing behavior.

990    ## 4.1   Saccade types that predict subsequent memory

991    It is well known that eye movement behavior during free viewing is important for the successful

992    formation of episodic memory (for overviews, see Ryan et al., 2020; Wynn et al., 2019). We replicate

993    and extend these findings by showing that the number of within-element saccades and between-

994    category revisits positively correlate with performance in an associative memory task (Fig. 3C). Short

995    within-element saccades scrutinize the visual features of single exemplars, increasing the visual

996    sampling of different details, which leads to better subsequent memory of the stimuli (Liu et al.,

997    2017; Loftus, 1972; Olsen et al., 2016).

999    As expected, the cumulative number of gaze transitions between categories during encoding

1000   increases memory performance. The between-category saccades presumably contribute to the

1001   binding of separate elements into a coherent representation of the event in which the hippocampus

1002   is involved (Herweg et al., 2020). Previous studies corroborate this idea by reporting that

1003   hippocampal activity and memory performance increase with the number of gaze fixations (Liu et al.,

1004   2017) and especially with the number of revisits (Kragel et al., 2021; Voss et al., 2011), as in our

1005   study.

1006

1007   In contrast to between-category saccades, the between-exemplar saccades did not predict

1008   subsequent memory performance. This was expected, as the memory test concerned between-

1009   category associations. This does, however, not mean that the between-exemplar saccades were not

1010   useful in forming a coherent event representation.

1011

1012

## 4.2 Theta and alpha modulations during the event encoding interval predict subsequent memory

First, the EEG analysis confirmed the presence of theta and alpha activity predictive of episodic remembering, extending previous work where the to-be-remembered material was presented centrally without the need for visual exploration via eye movements (Hanslmayr and Staudigl, 2014; Klimesch, 1996). Our results revealed that towards the end of the encoding interval (6-10 s), the theta power was higher for high than low subsequent event memory (Fig. 4A, C).

Consistent with previous research (Griffiths et al., 2019; Hanslmayr and Staudigl, 2014; Klimesch et al., 1996), we observed that high subsequent associative memory performance is related to a decrease in alpha power, mostly noticeable over the occipital areas (Fig. 4B). This decrease, however, became statistically significant only when we used a quadratic fit over time in the contrast analysis (Fig. 4D) while the subsequent memory effect for theta activity was linear.

Overall, in good agreement with the expected role of theta and alpha activity in episodic memory formation, we observe opposite effects predictive of subsequent memory in theta and alpha power. The theta increase and alpha decrease complement each other, reflecting the functioning of the hippocampal and cortical memory systems, respectively (Hanslmayr et al., 2016).

Additionally, we used the fBOSC method for oscillation detection (Seymour et al., 2022) to confirm the presence of true oscillatory activity in the data. In particular, the longest duration of oscillations ('abundance') in the encoding interval occurred around the alpha peak frequency. However, the abundance of alpha oscillatory activity did not predict subsequent memory performance (Fig. 4F). The duration of theta oscillations was shorter than the alpha oscillations; however, the theta oscillatory activity was predictive of memory performance. Oscillatory theta activity has been previously associated with memory performance (Hanslmayr et al., 2016), and our study replicates these previous findings. Alternatively, the duration of the theta oscillations may be associated with the eye movement rate since unrestricted eye movements are made at the theta frequency (Amit et al., 2017; Otero-Millan et al., 2008). Although oculomotor artifacts were eliminated from EEG during preprocessing, some residual artifacts at the rate of eye movements still could give rise to a pattern that appears as theta oscillations. However, the observed eye movement rate was much higher in low than high memory performance. Consequently, this eliminates the possibility that the eye movement rhythmicity could mechanistically determine the long duration of theta oscillatory activity, as well as the large power of theta oscillations that characterized high memory

1047 performance. Instead, we propose that the 'long' duration of memory-related theta oscillations is

1048 associated with the modulation of ongoing or saccade-induced theta oscillations (see below). Such

1049 oscillations may overlap, which would require correction by deconvolution. Perhaps this can be done

1050 by combining the oscillation detection methods (such as those used in 2.5.4), providing latencies of

1051 oscillation onsets, with the deconvolution technique. However, solutions to this intricate issue await

1052 method development far beyond the scope of the present study.

1053

1054 ## 4.3   Fixation-related EEG predicts subsequent memory

1055 The fixation-related EEG corroborates the findings reported at the event level. We observed reliably

1056 higher theta and lower alpha power for high than low memory performance (Fig. 5). In line with the

1057 findings at the event level, theta and alpha memory effects became greater over time of the

1058 encoding interval (Fig. 5B, F), which may reflect the buildup of an event representation across a

1059 sequence of eye movements. The predominantly anterior topography of the theta memory effect

1060 (Fig. 5D) suggests the contribution of frontal midline theta oscillations, which are related to intrinsic

1061 episodic memory processes (Hsieh and Ranganath, 2014). The alpha memory effect was

1062 predominantly distributed over the left hemisphere.

1063

1064 Memory processes in natural viewing are much more complex and evolve at a much higher rate than

1065 is considered by the design of traditional paradigms with fixed stimulus presentation. Depending on

1066 the fixation location, learning with unrestricted eye movements may involve encoding *and* retrieval,

1067 just as later remembering may trigger retrieval *and* encoding processes (Kragel and Voss, 2022).

1068 These dynamic memory processes across eye movements produce multiple effects on the

1069 underlying brain activity. The ongoing memory-related theta and alpha oscillations described above

1070 interfere with the activity elicited by saccade execution and information acquisition at new fixation

1071 locations. Each saccade evokes a broadband neural response (Dimigen et al., 2009; Ossandón et al.,

1072 2010) that lasts longer than 200-300 ms typical for fixation duration in natural viewing and therefore

1073 overlaps each other (Dimigen and Ehinger, 2021; Dimigen et al., 2011; Nikolaev et al., 2016).

1074 Moreover, each saccade aligns the phase of ongoing oscillations observed in the visual cortex of

1075 monkeys (Ito et al., 2011; Rajkai et al., 2008), in the human and macaque hippocampus (Hoffman et

1076 al., 2013), as well as in the scalp EEG in humans (Nikolaev et al., 2016). Such phase resetting

1077 supports successful memory formation (Jutras et al., 2013; Kota et al., 2020; Kragel et al., 2020) by

1078 synchronizing brain areas and optimizing the processing and encoding of visual information (Jutras

1079 and Buffalo, 2010; Voss et al., 2017). Thus, the neural activity during unrestricted viewing behavior

1080 constitutes a combination of overlapping evoked responses and the phase-reset of ongoing

oscillations (Ito et al., 2011). In such a situation, mapping external stimuli, mental processes, and neural activity is not straightforward, as in the traditional stimulus-response paradigm. In particular, ongoing oscillations may simultaneously represent the current and previous memory items (Herweg et al., 2020), and overlapping responses evoked by a series of eye movements may represent information about several previous fixation locations. The relative contribution of saccade-evoked responses and phase-reset oscillations to the resulting neural activity at fixation is challenging to estimate. Accordingly, the question remains open as to how the ongoing memory-related oscillations interact with the fixation-related activity representing visual processing at the current fixation location. It is clear, however, that neural activity time-locked to fixation onsets in a free-viewing memory task may reflect not only information acquisition at the currently fixated location but also preceding mental representations and more sustained higher-order memory processes.

The time window in the fixation-related EEG analysis was centered on the first power peak about 100 ms after the fixation onset. The latency of this peak coincides with the latency of lambda activity, which is associated with early perceptual processes at gaze fixation (Kazai and Yagi, 2003; Ossandón et al., 2010; Ries et al., 2016; Thickbroom et al., 1991). However, given that observed memory-related EEG effects have topographies that do not overlap with the typical occipital prominence of lambda activity (Ossandón et al., 2010; Thickbroom et al., 1991), it seems unlikely that the perceptual lambda activity would explain them. This interpretation is corroborated by previous research, where a comparable memory-related peak latency has been reported for frontal theta power (Adam et al., 2018). Ongoing theta oscillations supposedly drive perceptual sampling by organizing sequential inputs: they represent several items within the same cycle, where the current item is strongly represented and previous items are concurrently represented in a weaker form (Herweg et al., 2020). Such theta oscillations may thus be further modulated by fixation-related activity, where not only the currently fixated information is represented but also previously visited elements from the ongoing event memorization process (as well as previous events). Thus, our time window may capture individual perceptual moments at fixations and the sum of active and latent representations accumulated over a series of eye movements.

## 4.4   EEG memory effects after saccades between event elements

Fascinating results were obtained when analyzing EEG in fixation intervals after different saccade types during the 10-s encoding interval: fixations following between-category, between-exemplar, and within-element saccades. In our paradigm, between-category saccades supported associations between event elements that were directly targeted in the episodic memory task. The theta power

1115 was higher over the left hemisphere for high than low subsequent memory after between-category

1116 revisits (Fig. 6D, E), and the cumulative number of such revisits predicted subsequent memory (Fig.

1117 3C). The alpha power was lower for high than low memory at fixations after between-exemplar

1118 saccades, i.e., the associations that were not tested (Fig. 7). The cumulative number of such gaze

1119 transitions did not predict subsequent memory (Fig. 3C).

1120

1121 It is conceivable that gaze transitions between categories link the event elements together, creating

1122 a holistic representation of the entire event. These gaze transitions are thus critical to episodic

1123 memory as they support the binding of information experienced separately in space and time

1124 (Cohen and Eichenbaum, 1993; Voss et al., 2017). Previous research has shown that this binding

1125 mechanism relies on the transfer of information between distant brain regions and is promoted by

1126 synchronizing their activity at a theta rhythm (Clouter et al., 2017; Fell and Axmacher, 2011). Our

1127 results fit well with this previous literature. We observed high theta power after between-category

1128 saccades, predictive of high event memory, suggesting the involvement of theta in the binding of

1129 elements into a coherent episodic representation. It may be the case that theta reflects an

1130 optimized oculomotor control that guides this binding mechanism. In fact, it has been shown that

1131 the theta rhythm also underlies viewing behavior optimal for memory formation (Herweg et al.,

1132 2020; Jutras and Buffalo, 2010; Voss et al., 2017).

1133

1134 The crucial role of the hippocampal theta oscillations in memory-related gaze behavior has recently

1135 been demonstrated using intracranial EEG recording combined with eye tracking in free-viewing

1136 memory tasks (Kragel et al., 2021; Kragel et al., 2020). Kragel and colleagues (2021) described how

1137 revisits support the efficiency of visual sampling in the formation of memory for scenes.

1138 Hippocampal theta power decreased before revisits and increased after revisits, indicating rapid

1139 switches between memory-based saccade guidance and perceptual processing at fixation. Thus, the

1140 theta increase after revisits may reflect the mechanism binding information encoded at multiple

1141 fixations into a coherent representation. The present study supports the crucial role of revisits in

1142 successful memory formation: the number of between-category revisits and succeeding theta power

1143 predicts subsequent associative memory performance. These results demonstrate the feasibility of

1144 applying our approach in free-viewing studies of episodic memory in broad cohorts of healthy

1145 participants.

1146

1147 As expected, the cumulative number of between-exemplar saccades that were incidental to the task

1148 did not predict subsequent memory. Nevertheless, alpha power at fixations after these saccades was

1149 lower for high than low subsequent memory (Fig. 7). Although this alpha effect thus reflects a

1150 mechanism that supports successful memory formation, its exact nature remains unclear. For

1151 example, in episodic memory encoding, alpha activity indicates cortical information processing

1152 during the perception of an event (Jensen and Mazaheri, 2010; Klimesch, 2012), and its decrease

1153 predicts the successful formation of episodic memory (Hanslmayr et al., 2016). As discussed above,

1154 alpha desynchronization is thought to increase the capacity of local cell ensembles to process

1155 information by reducing local synchronization (Hanslmayr et al., 2016). Thus, the occipital

1156 predominance of memory-related alpha decrease in our visual memory task may reflect an increase

1157 in the capacity of cell ensembles in the visual system, which is beneficial for category-specific

1158 processing. Visual perception in our task involves discriminating between two similar exemplars

1159 from the same category, and the between-exemplar saccades may facilitate a comparison of the two

1160 exemplars. Alpha activity has been shown to predict behavioral performance in visual discrimination

1161 tasks (Hanslmayr et al., 2005; Nelli et al., 2017). Thus, the here observed alpha decrease after

1162 between-exemplar saccades may indicate efficient discrimination of two similar exemplars that

1163 promote successful encoding.

1164

1165 ## 4.5  EEG memory effects after saccades within event elements

1166 Within-element saccades are the only saccade type for which memory-related effects were found in

1167 both theta and alpha bands. They had the same direction in both frequency bands: high power

1168 predicted high memory. After within-element saccades, alpha power was higher for high than low

1169 subsequent memory performance over the right hemisphere (Fig. 8E-G), whereas theta power was

1170 higher for high than low memory performance over the frontal and central areas. Short within-

1171 element saccades that serve to scrutinize the visual features of images are known to increase visual

1172 sampling, which leads to better performance on subsequent item recognition tests (Loftus, 1972;

1173 Olsen et al., 2016). Accordingly, it is conceivable that a higher number of within-element saccades

1174 helps establish strong individual element representations to form part of multi-element event

1175 representations. Increased visual sampling results in a high memory load, reflected in high frontal

1176 theta power (Jensen and Tesche, 2002; Sederberg et al., 2006). In our paradigm, an increased

1177 memory load across encoding time corresponds to the accumulation of more visual details about

1178 event exemplars, which is favorable for later recall. Alternatively, frontal theta power has also been

1179 associated with detecting interference during selective retrieval in episodic memory (Ferreira et al.,

1180 2014; Kerrén et al., 2021; Staudigl et al., 2010) and during multiple-list learning in working memory

1181 (Kliegl and Bauml, 2021). In the present study, encoding exemplars from multiple events may be

1182 accompanied by interference from exemplars of the same categories in previously acquired events.

1183 Therefore, detecting interference may be crucial for encoding distinct event memories and may

1184   trigger adaptive inhibition of exemplars from previous events, ensuring that each event is stored

1185   separately to promote task performance. It is also conceivable that the frontal theta reflects the

1186   complementary effects of interference detection and memory load, as high memory load can lead to

1187   increased interference. The topography of the theta memory effect for within-element saccades

1188   differs from that of the between-category saccades, which was more prominent over the left

1189   hemisphere (Fig. 9), suggesting that those two theta effects represent distinct neural mechanisms.

1190

1191   There is overwhelming evidence that task-related increase in alpha power often corresponds to

1192   inhibitory neural processes (Jensen and Mazaheri, 2010). In studies of competitive memory retrieval,

1193   it has been demonstrated that high alpha power is associated with the inhibition of goal-irrelevant

1194   memories (Fellner et al., 2020; Waldhauser et al., 2012). Considering the likely involvement of both

1195   encoding and retrieval processes during free-viewing memory formation (Kragel and Voss, 2022), the

1196   observed alpha power increase may indicate inhibition of competing information (within and across

1197   events) that may interfere with building a coherent event representation across momentary

1198   sequences of visual sampling.

1199

## 1200  5  Conclusions

1201   We used a novel method of EEG-eye movement coregistration to examine episodic memory

1202   encoding in unrestricted viewing with a precision unavailable thus far. Our findings confirm eye

1203   movements' crucial role in forming episodic memory (Voss et al., 2017). We demonstrate that the

1204   functional specialization of eye movements manifested in various types of saccades is associated

1205   with different perceptual and cognitive processes, as evidenced by fixation-related theta and alpha

1206   EEG power. Episodic memory about an event is constructed through rapid switching between these

1207   processes in a sequence of fixations. These processes may indicate the complementary functioning

1208   of the hippocampal theta system for binding episodes and the cortical alpha system for representing

1209   the content of these episodes (Hanslmayr et al., 2016). Our study extends previous knowledge by

1210   elucidating how these systems engage across eye movements in the service of building coherent

1211   episodic memories for personal experiences.

1212

## 1213  Acknowledgments

1214   We thank Benedikt Ehinger for his help with deconvolution and Axel Ekström for his assistance with

1215   data collection. We are grateful to Marcus Nyström and Diederick Niehorster for their help in

1216   developing the EEG-eye movement coregistration system.

## Funding

This work was supported by a grant from the Marcus and Amelia Wallenberg Foundation (MAW2015.0043) to MJ.

## Data and code availability statement

The data and main codes used in this article are available at Open Science Framework:

https://osf.io/nad4y/?view_only=df274a4eaecc4ef6b9ec5053a0d2b1f0

## Declaration of Competing Interest

The authors declare no competing financial interests.

## CRediT authorship contribution statement

**Andrey R. Nikolaev**: Conceptualization, Methodology, Software, Formal analysis, Writing - Original Draft, Writing - Review & Editing. **Inês Bramão**: Conceptualization, Methodology, Writing - Review & Editing. **Roger Johansson**: Conceptualization, Methodology, Writing - Review & Editing. **Mikael Johansson**: Conceptualization, Methodology, Writing - Review & Editing, Supervision, Funding acquisition.

## References

Adam, K.C.S., Robison, M.K., Vogel, E.K., 2018. Contralateral Delay Activity Tracks Fluctuations in Working Memory Performance. J Cogn Neurosci 30, 1229-1240. 10.1162/jocn_a_01233.

Amit, R., Abeles, D., Bar-Gad, I., Yuval-Greenberg, S., 2017. Temporal dynamics of saccades explained by a self-paced process. Sci Rep 7, 886. 10.1038/s41598-017-00881-7.

Brandt, S.A., Stark, L.W., 1997. Spontaneous eye movements during visual imagery reflect the content of the visual scene. J Cogn Neurosci 9, 27-38. 10.1162/jocn.1997.9.1.27.

Brodeur, M.B., Dionne-Dostie, E., Montreuil, T., Lepage, M., 2010. The Bank of Standardized Stimuli (BOSS), a New Set of 480 Normative Photos of Objects to Be Used as Visual Stimuli in Cognitive Research. Plos One 5. 10.1371/journal.pone.0010773.

Caplan, J.B., Madsen, J.R., Raghavachari, S., Kahana, M.J., 2001. Distinct patterns of brain oscillations underlie two basic parameters of human maze learning. J Neurophysiol 86, 368-380. 10.1152/jn.2001.86.1.368.

Chang, C.Y., Hsu, S.H., Pion-Tonachini, L., Jung, T.P., 2020. Evaluation of Artifact Subspace Reconstruction for Automatic Artifact Components Removal in Multi-Channel EEG Recordings. Ieee T Bio-Med Eng 67, 1114-1121. 10.1109/Tbme.2019.2930186.

1249    Chelnokova, O., Laeng, B., Eikemo, M., Riegels, J., Loseth, G., Maurud, H., Willoch, F., Leknes, S.,
1250    2014. Rewards of beauty: the opioid system mediates social motivation in humans. Mol Psychiatr 19,
1251    746-747.  10.1038/mp.2014.1.

1252    Clouter, A., Shapiro, K.L., Hanslmayr, S., 2017. Theta Phase Synchronization Is the Glue that Binds
1253    Human Associative Memory. Curr Biol 27, 3143-3148.  10.1016/j.cub.2017.09.001.

1254    Coco, M.I., Nuthmann, A., Dimigen, O., 2020. Fixation-related Brain Potentials during Semantic
1255    Integration of Object-Scene Information. J Cogn Neurosci 32, 571-589.  10.1162/jocn_a_01504.

1256    Cohen, N.J., Eichenbaum, H., 1993. Memory, Amnesia, and The Hippocampal System. MIT Press,
1257    Cambridge, MA.

1258    Damiano, C., Walther, D.B., 2019. Distinct roles of eye movements during memory encoding and
1259    retrieval. Cognition 184, 119-129.  10.1016/j.cognition.2018.12.014.

1260    Delorme, A., Makeig, S., 2004. EEGLAB: an open source toolbox for analysis of single-trial EEG
1261    dynamics including independent component analysis. J Neurosci Meth 134, 9-21.
1262    10.1016/j.jneumeth.2003.10.009.

1263    Devillez, H., Guyader, N., Guerin-Dugue, A., 2015. An eye fixation-related potentials analysis of the
1264    P300 potential for fixations onto a target object when exploring natural scenes. J Vis 15, 20.
1265    https://doi.org/10.1167/15.13.20.

1266    Dien, J., Santuzzi, A.M., 2005. Application of repeated measures ANOVA to high density ERP
1267    datasets: A review and tutorial. In: Handy, T.C. (Ed.), Event-related potentials: A methods handbook
1268    MIT Press, Cambridge, MA, pp. 57– 82.

1269    Dimigen, O., 2020. Optimizing the ICA-based removal of ocular EEG artifacts from free viewing
1270    experiments. Neuroimage 207, 116117.  10.1016/j.neuroimage.2019.116117.

1271    Dimigen, O., Ehinger, B.V., 2021. Regression-based analysis of combined EEG and eye-tracking data:
1272    Theory and applications. J Vis 21.  10.1167/jov.21.1.3.

1273    Dimigen, O., Sommer, W., Hohlfeld, A., Jacobs, A.M., Kliegl, R., 2011. Coregistration of eye
1274    movements and EEG in natural reading: analyses and review. J Exp Psychol Gen 140, 552-572.
1275    https://doi.org/10.1037/a0023885.

1276    Donoghue, T., Haller, M., Peterson, E.J., Varma, P., Sebastian, P., Gao, R., Noto, T., Lara, A.H., Wallis,
1277    J.D., Knight, R.T., Shestyuk, A., Voytek, B., 2020. Parameterizing neural power spectra into periodic
1278    and aperiodic components. Nat Neurosci 23, 1655-1665.  10.1038/s41593-020-00744-x.

1279    Ehinger, B.V., Dimigen, O., 2019. Unfold: an integrated toolbox for overlap correction, non-linear
1280    modeling, and regression-based EEG analysis. PeerJ 7, e7838.  10.7717/peerj.7838.

1281    Engbert, R., Mergenthaler, K., 2006. Microsaccades are triggered by low retinal image slip. P Natl
1282    Acad Sci USA 103, 7192-7197.  10.1073/pnas.0509557103.

1283    Fell, J., Axmacher, N., 2011. The role of phase synchronization in memory processes. Nat Rev
1284    Neurosci 12, 105-118.  10.1038/nrn2979.

1285 Fellner, M.C., Waldhauser, G.T., Axmacher, N., 2020. Tracking Selective Rehearsal and Active
1286 Inhibition of Memory Traces in Directed Forgetting. Curr Biol 30, 2638-2644 e2634.
1287 10.1016/j.cub.2020.04.091.

1288 Ferreira, C.S., Marful, A., Staudigl, T., Bajo, T., Hanslmayr, S., 2014. Medial prefrontal theta
1289 oscillations track the time course of interference during selective memory retrieval. J Cogn Neurosci
1290 26, 777-791. 10.1162/jocn_a_00523.

1291 Fischer, T., Graupner, S.T., Velichkovsky, B.M., Pannasch, S., 2013. Attentional dynamics during free
1292 picture viewing: evidence from oculomotor behavior and electrocortical activity. Front Syst Neurosci
1293 7, 17. https://doi.org/10.3389/fnsys.2013.00017.

1294 Fong, D.C.L., Saunders, M., 2011. Lsmr: An Iterative Algorithm for Sparse Least-Squares Problems.
1295 Siam J Sci Comput 33, 2950-2971. Doi 10.1137/10079687x.

1296 Fudali-Czyz, A., Francuz, P., Augustynowicz, P., 2018. The Effect of Art Expertise on Eye Fixation-
1297 Related Potentials During Aesthetic Judgment Task in Focal and Ambient Modes. Front Psychol 9,
1298 1972. 10.3389/fpsyg.2018.01972.

1299 Gilchrist, I.D., Harvey, M., 2000. Refixation frequency and memory mechanisms in visual search. Curr
1300 Biol 10, 1209-1212. 10.1016/s0960-9822(00)00729-6.

1301 Griffiths, B.J., Parish, G., Roux, F., Michelmann, S., van der Plas, M., Kolibius, L.D., Chelvarajah, R.,
1302 Rollings, D.T., Sawlani, V., Hamer, H., Gollwitzer, S., Kreiselmeyer, G., Staresina, B., Wimber, M.,
1303 Hanslmayr, S., 2019. Directional coupling of slow and fast hippocampal gamma with neocortical
1304 alpha/beta oscillations in human episodic memory. Proc Natl Acad Sci U S A 116, 21834-21842.
1305 10.1073/pnas.1914180116.

1306 Hanslmayr, S., Klimesch, W., Sauseng, P., Gruber, W., Doppelmayr, M., Freunberger, R.,
1307 Pecherstorfer, T., 2005. Visual discrimination performance is related to decreased alpha amplitude
1308 but increased phase locking. Neurosci Lett 375, 64-68. 10.1016/j.neulet.2004.10.092.

1309 Hanslmayr, S., Staresina, B.P., Bowman, H., 2016. Oscillations and episodic memory: addressing the
1310 synchronization/desynchronization conundrum. Trends Neurosci 39, 16-25.
1311 10.1016/j.tins.2015.11.004.

1312 Hanslmayr, S., Staudigl, T., 2014. How brain oscillations form memories--a processing based
1313 perspective on oscillatory subsequent memory effects. Neuroimage 85 Pt 2, 648-655.
1314 10.1016/j.neuroimage.2013.05.121.

1315 Herweg, N.A., Solomon, E.A., Kahana, M.J., 2020. Theta oscillations in human memory. Trends Cogn
1316 Sci 24, 208-227. 10.1016/j.tics.2019.12.006.

1317 Hogendoorn, H., 2016. Voluntary saccadic eye movements ride the attentional rhythm. J Cogn
1318 Neurosci 28, 1625-1635. 10.1162/jocn_a_00986.

1319 Hsieh, L.T., Ranganath, C., 2014. Frontal midline theta oscillations during working memory
1320 maintenance and episodic encoding and retrieval. Neuroimage 85 Pt 2, 721-729.
1321 10.1016/j.neuroimage.2013.08.003.

1322 Jensen, O., Mazaheri, A., 2010. Shaping functional architecture by oscillatory alpha activity: gating by
1323 inhibition. Front Hum Neurosci 4, 186. 10.3389/fnhum.2010.00186.

1324 Jensen, O., Tesche, C.D., 2002. Frontal theta activity in humans increases with memory load in a
1325 working memory task. Eur J Neurosci 15, 1395-1399. 10.1046/j.1460-9568.2002.01975.x.

1326 Johansson, R., Johansson, M., 2014. Look here, eye movements play a functional role in memory
1327 retrieval. Psychol. Sci. 25, 236-242. 10.1177/0956797613498260.

1328 Johansson, R., Nystrom, M., Dewhurst, R., Johansson, M., 2022. Eye-movement replay supports
1329 episodic remembering. Proc Biol Sci 289, 20220964. 10.1098/rspb.2022.0964.

1330 Jutras, M.J., Buffalo, E.A., 2010. Synchronous neural activity and memory formation. Curr Opin
1331 Neurobiol 20, 150-155. 10.1016/j.conb.2010.02.006.

1332 Jutras, M.J., Fries, P., Buffalo, E.A., 2013. Oscillatory activity in the monkey hippocampus during
1333 visual exploration and memory formation. Proc Natl Acad Sci U S A 110, 13144-13149.
1334 10.1073/pnas.1302351110.

1335 Kamienkowski, J.E., Varatharajah, A., Sigman, M., Ison, M.J., 2018. Parsing a mental program:
1336 Fixation-related brain signatures of unitary operations and routines in natural visual search.
1337 Neuroimage 183, 73-86. 10.1016/j.neuroimage.2018.08.010.

1338 Kazai, K., Yagi, A., 2003. Comparison between the lambda response of eye-fixation-related potentials
1339 and the P100 component of pattern-reversal visual evoked potentials. Cogn Affect Behav Neurosci 3,
1340 46-56. 10.3758/cabn.3.1.46.

1341 Kerrén, C., Bramão, I., Hellerstedt, R., Johansson, M., 2021. Strategic retrieval prevents memory
1342 interference: The temporal dynamics of retrieval orientation. Neuropsychologia 154, 107776.
1343 10.1016/j.neuropsychologia.2021.107776.

1344 Kliegl, O., Bauml, K.H.T., 2021. The Mechanisms Underlying Interference and Inhibition: A Review of
1345 Current Behavioral and Neuroimaging Research. Brain Sci 11. 10.3390/brainsci11091246.

1346 Klimesch, W., 1996. Memory processes, brain oscillations and EEG synchronization. Int J
1347 Psychophysiol 24, 61-100. 10.1016/s0167-8760(96)00057-8.

1348 Klimesch, W., 2012. alpha-band oscillations, attention, and controlled access to stored information.
1349 Trends Cogn Sci 16, 606-617. 10.1016/j.tics.2012.10.007.

1350 Klimesch, W., Schimke, H., Doppelmayr, M., Ripper, B., Schwaiger, J., Pfurtscheller, G., 1996. Event-
1351 related desynchronization (ERD) and the Dm effect: does alpha desynchronization during encoding
1352 predict later recall performance? Int J Psychophysiol 24, 47-60. 10.1016/s0167-8760(96)00054-2.

1353 Körner, C., Braunstein, V., Stangl, M., Schlogl, A., Neuper, C., Ischebeck, A., 2014. Sequential effects
1354 in continued visual search: using fixation-related potentials to compare distractor processing before
1355 and after target detection. Psychophysiology 51, 385-395. https://doi.org/10.1111/psyp.12062.

1356 Körner, C., Gilchrist, I.D., 2008. Memory processes in multiple-target visual search. Psychological
1357 research 72, 99-105. https://doi.org/10.1007/s00426-006-0075-1.

1358 Kosciessa, J.Q., Grandy, T.H., Garrett, D.D., Werkle-Bergner, M., 2020. Single-trial characterization of
1359 neural rhythms: Potential and challenges. Neuroimage 206, 116331.
1360 10.1016/j.neuroimage.2019.116331.

Kota, S., Rugg, M.D., Lega, B.C., 2020. Hippocampal Theta Oscillations Support Successful Associative Memory Formation. J Neurosci 40, 9507-9518.  10.1523/JNEUROSCI.0767-20.2020.

Kothe, C.A.E., Jung, T.P., 2016. Artifact removal techniques with signal reconstruction. U.S. Patent Application. 14/895,440.

Kragel, J.E., Schuele, S., VanHaerents, S., Rosenow, J.M., Voss, J.L., 2021. Rapid coordination of effective learning by the human hippocampus. Sci Adv 7.  10.1126/sciadv.abf7144.

Kragel, J.E., VanHaerents, S., Templer, J.W., Schuele, S., Rosenow, J.M., Nilakantan, A.S., Bridge, D.J., 2020. Hippocampal theta coordinates memory processing during visual exploration. Elife 9. 10.7554/eLife.52108.

Kragel, J.E., Voss, J.L., 2022. Looking for the neural basis of memory. Trends Cogn Sci 26, 53-65. 10.1016/j.tics.2021.10.010.

Liu, Z.X., Shen, K., Olsen, R.K., Ryan, J.D., 2017. Visual Sampling Predicts Hippocampal Activity. J Neurosci 37, 599-609.  10.1523/JNEUROSCI.2610-16.2016.

Loftus, G.R., 1972. Eye fixations and recognition memory for pictures. Cognitive Psychol 3, 525-551. Doi 10.1016/0010-0285(72)90021-7.

Meghanathan, R.N., Nikolaev, A.R., van Leeuwen, C., 2019. Refixation patterns reveal memory-encoding strategies in free viewing. Atten Percept Psychophys 81, 2499-2516.  10.3758/s13414-019-01735-2.

Meghanathan, R.N., van Leeuwen, C., Giannini, M., Nikolaev, A.R., 2020. Neural correlates of task-related refixation behavior. Vision Res 175, 90-101.  10.1016/j.visres.2020.07.001.

Mirza, M.B., Adams, R.A., Mathys, C.D., Friston, K.J., 2016. Scene Construction, Visual Foraging, and Active Inference. Front Comput Neurosc 10.  10.3389/fncom.2016.00056.

Nelli, S., Itthipuripat, S., Srinivasan, R., Serences, J.T., 2017. Fluctuations in instantaneous frequency predict alpha amplitude during visual perception. Nat Commun 8, 2071.  10.1038/s41467-017-02176-x.

Niehorster, D.C., Andersson, R., Nystrom, M., 2020. Titta: A toolbox for creating PsychToolbox and Psychopy experiments with Tobii eye trackers. Behav Res Methods 52, 1970-1979.  10.3758/s13428-020-01358-8.

Nikolaev, A.R., Ehinger, B.V., Meghanathan, R.N., van Leeuwen, C., 2022. At a second glance: cognitive and oculomotor neural activity of refixation planning. bioRxiv, 660308. https://doi.org/10.1101/660308.

Nikolaev, A.R., Meghanathan, R.N., van Leeuwen, C., 2016. Combining EEG and eye movement recording in free viewing: pitfalls and possibilities. Brain Cogn 107, 55-83. https://doi.org/10.1016/j.bandc.2016.06.004.

Nikolaev, A.R., Meghanathan, R.N., van Leeuwen, C., 2018. Refixation control in free viewing: a specialized mechanism divulged by eye-movement related brain activity. Journal of neurophysiology 120, 2311-2324.  10.1152/jn.00121.2018.

1398  Nikolaev, A.R., Nakatani, C., Plomp, G., Jurica, P., van Leeuwen, C., 2011. Eye fixation-related
1399  potentials in free viewing identify encoding failures in change detection. Neuroimage 56, 1598-1607.
1400  https://doi.org/10.1016/j.neuroimage.2011.03.021.

1401  Noton, D., Stark, L., 1971. Scanpaths in Saccadic Eye Movements While Viewing and Recognizing
1402  Patterns. Vision Research 11, 929-&.  10.1016/0042-6989(71)90213-6.

1403  Olsen, R.K., Sebanayagam, V., Lee, Y., Moscovitch, M., Grady, C.L., Rosenbaum, R.S., Ryan, J.D., 2016.
1404  The relationship between eye movements and subsequent recognition: Evidence from individual
1405  differences and amnesia. Cortex 85, 182-193.  10.1016/j.cortex.2016.10.007.

1406  Ossandón, J.P., Helo, A.V., Montefusco-Siegmund, R., Maldonado, P.E., 2010. Superposition model
1407  predicts EEG occipital activity during free viewing of natural scenes. J Neurosci 30, 4787-4795.
1408  https://doi.org/10.1523/JNEUROSCI.5769-09.2010.

1409  Ossandón, J.P., König, P., Heed, T., 2020. No Evidence for a Role of Spatially Modulated alpha-Band
1410  Activity in Tactile Remapping and Short-Latency, Overt Orienting Behavior. J Neurosci 40, 9088-9102.
1411  10.1523/Jneurosci.0581-19.2020.

1412  Otero-Millan, J., Troncoso, X.G., Macknik, S.L., Serrano-Pedraza, I., Martinez-Conde, S., 2008.
1413  Saccades and microsaccades during visual fixation, exploration, and search: foundations for a
1414  common saccadic generator. J Vis 8, 21 21-18.  10.1167/8.14.21.

1415  Peirce, J., Gray, J.R., Simpson, S., MacAskill, M., Hochenberger, R., Sogo, H., Kastman, E., Lindelov,
1416  J.K., 2019. PsychoPy2: Experiments in behavior made easy. Behav Res Methods 51, 195-203.
1417  10.3758/s13428-018-01193-y.

1418  Pierrot-Deseilligny, C., Milea, D., Muri, R.M., 2004. Eye movement control by the cerebral cortex.
1419  Curr Opin Neurol 17, 17-25.  10.1097/00019052-200402000-00005.

1420  Pion-Tonachini, L., Kreutz-Delgado, K., Makeig, S., 2019. ICLabel: An automated
1421  electroencephalographic independent component classifier, dataset, and website. Neuroimage 198,
1422  181-197.  10.1016/j.neuroimage.2019.05.026.

1423  Plöchl, M., Ossandón, J.P., König, P., 2012. Combining EEG and eye tracking: identification,
1424  characterization, and correction of eye movement artifacts in electroencephalographic data. Front
1425  Hum Neurosci 6, 278.  https://doi.org/10.3389/fnhum.2012.00278.

1426  Ries, A.J., Touryan, J., Ahrens, B., Connolly, P., 2016. The Impact of Task Demands on Fixation-
1427  Related Brain Potentials during Guided Search. Plos One 11, e0157260.
1428  10.1371/journal.pone.0157260.

1429  Ryan, J.D., Shen, K., Liu, Z.X., 2020. The intersection between the oculomotor and hippocampal
1430  memory systems: empirical developments and clinical implications. Ann N Y Acad Sci 1464, 115-141.
1431  10.1111/nyas.14256.

1432  Sederberg, P.B., Gauthier, L.V., Terushkin, V., Miller, J.F., Barnathan, J.A., Kahana, M.J., 2006.
1433  Oscillatory correlates of the primacy effect in episodic memory. Neuroimage 32, 1422-1431.
1434  10.1016/j.neuroimage.2006.04.223.

1435  Seymour, R.A., Alexander, N., Maguire, E.A., 2022. Robust estimation of 1/f activity improves
1436  oscillatory burst detection. Eur J Neurosci.  10.1111/ejn.15829.

1437    Shen, K., Bezgin, G., Selvam, R., McIntosh, A.R., Ryan, J.D., 2016. An Anatomical Interface between
1438    Memory and Oculomotor Systems. J Cognitive Neurosci 28, 1772-1783.  10.1162/jocn_a_01007.

1439    Smith, N.J., Kutas, M., 2015. Regression-based estimation of ERP waveforms: I. The rERP framework.
1440    Psychophysiology 52, 157-168.  10.1111/psyp.12317.

1441    Staudigl, T., Hanslmayr, S., Bauml, K.H., 2010. Theta oscillations reflect the dynamics of interference
1442    in episodic memory retrieval. J Neurosci 30, 11356-11362.  10.1523/JNEUROSCI.0637-10.2010.

1443    Tatler, B.W., Gilchrist, I.D., Land, M.F., 2005. Visual memory for objects in natural scenes: from
1444    fixations to object files. The Quarterly journal of experimental psychology. A, Human experimental
1445    psychology 58, 931-960.  https://doi.org/10.1080/02724980443000430.

1446    Team, R.C., 2020. R: A language and environment for statistical computing. R Foundation for
1447    Statistical Computing, Vienna, Austria.

1448    Thickbroom, G.W., Knezevic, W., Carroll, W.M., Mastaglia, F.L., 1991. Saccade onset and offset
1449    lambda waves: relation to pattern movement visually evoked potentials. Brain Res 551, 150-156.
1450    https://doi.org/10.1016/0006-8993(91)90927-N.

1451    Tulving, E., 1983. Ecphoric Processes in Episodic Memory. Philos T Roy Soc B 302, 361-371.  DOI
1452    10.1098/rstb.1983.0060.

1453    Turk-Browne, N.B., 2019. The hippocampus as a visual area organized by space and time: A
1454    spatiotemporal similarity hypothesis. Vision Research 165, 123-130.  10.1016/j.visres.2019.10.007.

1455    Tyson-Carr, J., Soto, V., Kokmotou, K., Roberts, H., Fallon, N., Byrne, A., Giesbrecht, T., Stancak, A.,
1456    2020. Neural underpinnings of value-guided choice during auction tasks: An eye-fixation related
1457    potentials study. Neuroimage 204, 116213.  10.1016/j.neuroimage.2019.116213.

1458    Voss, J.L., Bridge, D.J., Cohen, N.J., Walker, J.A., 2017. A closer look at the hippocampus and
1459    memory. Trends Cogn Sci 21, 577-588.  10.1016/j.tics.2017.05.008.

1460    Voss, J.L., Warren, D.E., Gonsalves, B.D., Federmeier, K.D., Tranel, D., Cohen, N.J., 2011. Spontaneous
1461    revisitation during visual exploration as a link among strategic behavior, learning, and the
1462    hippocampus. Proc Natl Acad Sci U S A 108, E402-409.  10.1073/pnas.1100225108.

1463    Waldhauser, G.T., Johansson, M., Hanslmayr, S., 2012. Alpha/beta oscillations indicate inhibition of
1464    interfering visual memories. J Neurosci 32, 1953-1961.  10.1523/JNEUROSCI.4201-11.2012.

1465    Wood, S.N., 2017. Generalized additive models: an introduction with R 2nd edition ed. Chapman &
1466    Hall/CRC, New York.

1467    Wynn, J.S., Ryan, J.D., Buchsbaum, B.R., 2020. Eye movements support behavioral pattern
1468    completion. Proc Natl Acad Sci U S A 117, 6246-6254.  10.1073/pnas.1917586117.

1469    Wynn, J.S., Shen, K., Ryan, J.D., 2019. Eye movements actively reinstate spatiotemporal mnemonic
1470    content. Vision (Basel) 3.  10.3390/vision3020021.
1471