Genetic adaptations to SIV across chimpanzee populations

Harvinder Pawar[1,2], Harrison J. Ostridge[1], Joshua M. Schmidt[1,3,4*§], Aida M. Andrés[1*§]

1: UCL Genetics Institute, Department of Genetics, Evolution and Ecology, University College

London, London, UK.

2: Present address: Institute of Evolutionary Biology (UPF-CSIC), Barcelona, Spain.

3: Department of Ecology and Evolution, School of Biological Sciences, University of Adelaide

Adelaide, Australia.

4: Present address: Flinders Centre for Ophthalmology, Eye and Vision Research, Flinders Health

and Medical Research Institute (FHMRI), Flinders University, Adelaide, Australia.

* Equal contribution.

§ Corresponding authors: Joshua M. Schmidt and Aida M. Andrés.

Email: joshmschmidt1@gmail.com (JMS), a.andres@ucl.ac.uk (AMA).

Keywords

Simian Immunodeficiency Virus; host-pathogen coevolution; natural immunity; adaptation

Short title

Adaptation to SIV in chimpanzee populations

Author contributions

JMS and AMA conceived the project. HP and JMS wrote code for simulations. HP performed

simulations and data analyses, under the supervision of JMS and AMA. JMS and HO performed

data analyses. JMS modified code for 3P-CLR, and wrote code for the joint enrichment tests. HP

wrote the first version of the manuscript and all co-authors contributed to writing the manuscript.

Abstract

Central and eastern chimpanzees are infected with Simian Immunodeficiency Virus (SIV) in the wild, typically without developing acute immunodeficiency. Yet the recent zoonotic transmission of chimpanzee SIV to humans, which were naïve to the virus, gave rise to the Human Immunodeficiency Virus (HIV), which causes AIDS and is responsible for one of the deadliest pandemics in human history. Chimpanzees have been infected with SIV for tens of thousands of years and have likely evolved to reduce its pathogenicity, becoming semi-natural hosts that largely tolerate the virus. In support of this view, central and eastern chimpanzees show evidence of positive selection in genes involved in SIV/HIV cell entry and immune response to SIV, respectively. We hypothesise that the first critical adaptations aimed at controlling the lethal potential of zoonotic SIV happened in the ancestors of central-eastern chimpanzees, the chimpanzee population that was first infected with the virus. Under that scenario, studying this population would allow us to identify those first critical host adaptations. Here, we use population genomics with that aim. In support of our hypothesis, the genes with signatures of positive selection in the ancestral population are significantly enriched in SIV-related genes, especially those involved in the immune response to SIV and those encoding for host genes that physically interact with SIV/HIV (VIPs). Interestingly, integrating these genes with candidates of positive selection in the two infected subspecies reveals novel signals of adaptation to SIV/HIV. Specifically, we observe evidence of positive selection in numerous steps of the biological pathway responsible for T-helper cell differentiation, including *CD4* and multiple genes that SIV/HIV use to infect and control host cells. This pathway is active only in CD4+ cells which SIV/HIV infects, and it plays a crucial role in shaping the immune response so it can efficiently control the virus. Our results confirm the importance of SIV as a selective factor, identify specific genetic changes that may have allowed our closest living relatives to reduce SIV's pathogenicity, and demonstrate the potential of population genomics to reveal the evolutionary mechanisms used by naïve hosts to reduce the pathogenicity of zoonotic pathogens.

Author Summary

Chimpanzees are at the origin of HIV-1, a virus that generates an incurable disease and that generated a pandemic that has claimed 35 million lives. Chimpanzees have evolved to control the pathogenicity of the virus, which does not typically develop into AIDS in the same way as in humans. Identifying the genetic adaptations responsible for this process provides critical

61 knowledge about SIV and HIV. Our analysis of chimpanzee genetic adaptations identified specific

62 genes and molecular pathways involved in adaptation to SIV, providing important insights into the

63 mechanisms that likely allowed our closest living relatives to control SIV/HIV. Further, we

64 establish SIV as a strong and recurrent selective pressure in central and eastern chimpanzees,

65 two important subspecies of large mammals that are currently endangered.

Introduction

A typical evolutionary outcome of pathogen infections is the development of host immunity or tolerance to an infectious agent that was initially highly pathogenic, and that remains so in naïve species. Host genetic adaptations underlie the sometimes striking differences in infection outcome in different species infected by the same virus. The lentivirus Simian Immunodeficiency Virus (SIV) provides a fascinating example of this process. Most African primates, the great apes being notable exceptions, have their own endemic SIV strain for which they are considered natural hosts (Chahroudi et al. 2012). When natural hosts (e.g. vervet monkeys) are exposed to their endemic SIV, they have a proportionate immune activation that preserves CD4+ T cell counts and wider immune function, resulting in asymptomatic outcomes and no significant reduction in lifespan despite viral replication (Ma et al. 2013; Ma et al. 2014; Silvestri et al. 2007). In contrast, species that are naïve to SIV, such as non-African primates, are unable to resolve the acute phase of infection and instead generate chronic immune activation, CD4+ T cell depletion and eventual immunodeficiency (Goldstein et al. 2005; Mandell et al. 2014; Barouch et al. 2016; Chahroudi et al. 2012). The course of these infections and the development of clinical disease is almost identical to that of humans infected with HIV, which are also a naïve species and which, without treatment, progress to acquired immunodeficiency syndrome (AIDS). Host genetic adaptation can thus turn a lethal virus into a tolerable infection.

Humans have received multiple zoonotic transmissions of SIV. The most deadly, which introduced HIV-1 group M responsible for the AIDS pandemic, originated when the SIV of a central chimpanzee (*Pan troglodytes troglodytes*) jumped into humans in the twentieth century (Keele et al. 2006; Worobey et al. 2008). HIV is thus genetically very similar to chimpanzee SIV (SIVcpz) (Keele et al. 2006). Also, the chimpanzee genome has a high sequence identity to the human genome (Chimpanzee Sequencing and Analysis Consortium 2005) and the two species share most aspects of their physiology. Therefore, the mechanisms used by chimpanzees to limit the pathogenicity of SIV may be informative about potential mechanisms to control HIV.

Chimpanzees have traditionally been considered natural SIV hosts, because despite infections in zoos and laboratories, infected chimpanzees rarely progress to AIDS-like symptoms (Sharp et al. 2005; de Groot et al. 2017; Gilden et al. 1986). Further, SIV-infected chimpanzees show proportionate patterns of immune activation, similar to those observed in natural hosts, such as vervet monkeys (Greenwood et al. 2015). Yet, recent observations have suggested that chimpanzees are not true natural hosts, instead lying between natural and symptomatic SIV

hosts. Disease symptoms such as CD4+ T-cell depletion and thrombocytopenia have been
observed in infected chimpanzees and associated with reduced fitness, particularly in the wild
(Keele et al. 2009; Greenwood et al. 2015; Etienne et al. 2011). Specifically, Keele *et al.*, observed
in habituated wild eastern chimpanzees that SIV infection increased mortality risk and infected
females had fewer offspring and higher infant mortality rates than uninfected females (Keele et
al. 2009). Similarly, SIV infection has been associated with population decline in the Kalande
(Rudicell et al. 2010). It should be noted that studies investigating the fitness effects of SIVcpz in
the wild have been restricted to small sample sizes from few habituated populations due to the
inherent challenges associated with studying wild populations of endangered large mammals.
Nevertheless, evidence suggests that chimpanzees can be considered 'semi-natural' hosts, as
the virus lacks the dramatic pathogenicity seen in naïve species like humans yet likely has non-
negligible health consequences and fitness effects. This is perhaps expected if chimpanzees have
already acquired adaptations that control the high initial pathogenicity of SIV but have not yet
acquired adaptations that fully control the effects of the infection.

Interestingly, of the four genetically and geographically distinct subspecies of chimpanzee, natural
SIV infection is found only in central (*P. t. troglodytes*) and eastern (*P. t. schweinfurthii*)
chimpanzees, who are most closely related to each other (Prado-Martinez et al. 2013) (Figure 1).
SIV infection has not been detected in the wild in the second clade of chimpanzees, which
includes the western (*P. t. verus*) and Nigeria-Cameroon (*P. t. ellioti*) subspecies (Locatelli et al.
2016; Keele et al. 2006; Prado-Martinez et al. 2013). The uneven distribution of SIV among
chimpanzee subspecies has sparked much interest in the origin of the virus in the *Pan* lineage.

We know that SIVcpz is the result of zoonotic transmission to chimpanzees of at least two African
monkey SIV strains (from the red-capped mangabey and the ancestor of the mona, moustached
and greater spot-nosed monkeys), which recombined and generated SIVcpz, a mosaic lineage
able to infect and transmit in chimpanzees (Bell and Bedford 2017; Bailes et al. 2003). Given this
complex origin, it is most likely that SIVcpz was acquired only once. A number of scenarios are
theoretically compatible with the distribution of SIVcpz in nature, most notably transmission to the
common ancestor of all chimpanzees and subsequent clearance in western and Nigeria-
Cameroon; transmission to the common ancestor of central and eastern chimpanzees; and
transmission to one subspecies (e.g. central) with subsequent transfer via migration to the other
subspecies (e.g. eastern). While this point is not completely settled, the epidemiological
distribution of SIVcpz and phylogenetic evidence both point towards zoonotic transmission into

the common ancestor of central and eastern chimpanzees (hereafter central-eastern ancestor).
An old transmission into the common ancestor of all chimpanzees is highly unlikely as it would
require the virus to be lost in two subspecies and a time of transmission (older than 500 thousand
years ago (kya)) that is incompatible with phylogenetic estimates from SIVcpz sequences
(Wertheim and Worobey 2009). The remaining two scenarios are more likely. The phylogeny of
SIVcpz separates two monophyletic sister clades, one containing all SIVcpzPtt sequences (the
virus infecting central chimpanzees) and one containing all SIVcpzPts sequences (the virus
infecting eastern chimpanzees) (Leitner et al. 2007; Sharp, Shaw, and Hahn 2005), which could
be compatible with both scenarios. Still, SIVcpzPts lineages do not fall within the diversity of
SIVcpzPtt, or vice versa (Leitner et al. 2007), as we would expect if one subspecies (e.g. centrals)
was originally infected and passed the virus to the other subspecies (e.g. easterns). Further, the
most recent common ancestor of all SIVcpz has been inferred to ~450-100 kya (Bailes et al. 2003;
Locatelli et al. 2016), a time before the two subspecies split ~106kya (de Manuel et al. 2016),
further supporting a scenario of zoonotic transmission to the central-eastern ancestor. Thus,
based on current evidence we use this scenario as our working hypothesis.
150
151
152

Figure 1. Phylogeny of chimpanzee subspecies and bonobos with the distribution of natural SIV infection. The dotted red line indicates the central-eastern ancestor where the first SIV infection most likely occurred, for which we investigate positive selection.

SIV is a dangerous pathogen, and as such a strong selective force. Once SIV infects a species, SIV-related adaptation is likely to be pervasive and continuous. In fact even natural hosts such as vervet monkeys, which have been infected for 0.5-3 million years, show genome-wide signatures of positive selection in SIV-related genes (Svardal et al. 2017; Ma et al. 2014). It has long been thought that SIV may be a strong selective force in chimpanzees, posited to drive allele frequency change and adaptation in a few immune and SIV-related genes (de Groot et al. 2002; de Groot et al. 2000; de Groot et al. 2008; Wooding et al. 2005; MacFie et al. 2009). Analysing dozens of genomes, we established how SIV has driven chimpanzee evolution by uncovering the presence of recent adaptations to SIV in the central and eastern subspecies (Schmidt et al. 2019). Using the PBSnj statistic, we identified the SNPs with the greatest allele frequency change in each chimpanzee subspecies, which are the strongest candidate targets of subspecies-specific positive selection. Strikingly, both in central and eastern chimpanzees, these SNPs are enriched in genes related to SIV. As expected under a model of zoonotic transmission into the central-eastern clade, neither western nor Nigeria-Cameroon showed evidence of recent positive selection in SIV-related genes (Schmidt et al. 2019). Although earlier work had reported evidence of low diversity at *CCR5, CXCR4* and *CX3CR1* in Nigeria-Cameroon and western, and suggested that this may be due to positive selection (Wooding et al. 2005; MacFie et al. 2009), it is unclear if SIV was the selective force.

The evidence of genetic adaptation in sets of SIV-related genes is clearer and easy to interpret. Identified candidate targets of recent selection in central chimpanzees are enriched in cytokine coreceptors due to signatures in *CCR3*, *CCR9* and *CXCR6* (Schmidt et al. 2019), which mediate HIV cell entry together with the primary receptor CD4 (Elliott et al. 2015; Wetzel et al. 2017; Nedellec et al. 2009) and are paralogs of the HIV coreceptors *CCR5* and *CXCR4* (Berger 1997; Moore et al. 2004; Blumenthal et al. 2012). In contrast, selection targets in eastern chimpanzees are enriched in "SIV-response genes", genes that upon SIV infection show different expression profiles in CD4+ T lymphocytes in natural host vs. naïve host species (in this case, vervet monkeys vs macaques, (Svardal et al. 2017; Jacquelin et al. 2014; Jacquelin et al. 2009)). These SIV-response genes likely contribute to the finely tuned natural host response to SIV, which is able to control the infection and results in non-pathogenic outcomes (Schmidt et al. 2019). Thus, interestingly, the two subspecies have evolved differential adaptation mechanisms to control SIV.

Under a scenario of zoonotic transmission into the central-eastern ancestor, this naïve population
would have been under the strongest pressure to adapt. We thus hypothesise that positive
selection in the central-eastern ancestor was critical to control the lethal potential of zoonotic SIV
upon a naïve chimpanzee population. Under that hypothesis, we could detect signatures of
positive selection in SIV-related genes in the central-eastern ancestor. We test this hypothesis
and aim to identify these adaptive genetic changes using 47 chimpanzee whole genomes. We
find evidence of adaptation in genes involved in SIV biology, providing further support for SIV
infection in the central-eastern ancestor. Further, candidate targets of positive selection point to
diverse adaptive mechanisms, including host response to infection, viral interactions and cell
entry, and combining information across populations we discover previously known mechanisms
of adaptation to SIV. Specifically, we identify the T helper cell type-1/type-2 (Th1/Th2)
differentiation pathway as a critical player that has been repeatedly targeted by positive selection
at different time points during chimpanzee evolution. This reveals the potential of this functional
pathway, and specific genes within the pathway, to control the pathogenicity of SIV/HIV infection.

Excitingly, population genetics tools allow us to identify the genetic adaptations that happened at
this critical time during which chimpanzees evolved to reduce the pathogenicity of a potentially
deadly virus.
205

Results

Signatures of positive selection in the central-eastern ancestor

We identified genomic regions showing evidence of positive selection in the central-eastern
ancestor using 3P-CLR (Racimo 2016). 3P-CLR tests distortions of the site frequency spectrum
(SFS) due to selective sweeps, by modelling the evolutionary trajectory of alleles in a 3-population
tree and comparing the likelihood of the observed SFS under contrasting hypotheses of neutrality
and positive selection (Racimo 2016). 3P-CLR is ideal for our purpose. First, it has high power to
detect the events we target, as it was designed to identify hard sweeps in ancestral modern
humans using Neanderthals as an outgroup (Racimo 2016), and the modern human-Neanderthal
divergence (Prüfer et al. 2017) is on the order of the inferred split time between chimpanzee
clades (de Manuel et al. 2016). We established with simulations that 3P-CLR also has very high
power to detect selective events in the central-eastern ancestor when Nigeria-Cameroon is used
as outgroup (Figure S1), with power equal or higher than that reported in humans (Racimo 2016)
under the same scenario: strong selection and fixation of the advantageous allele (Supplementary
Material). Second, 3P-CLR has high power to detect selection in the ancestral population while

being largely unaffected by convergent evolution (independent selection in both central and eastern chimpanzees) that does not generate the SFS distortions that translate into high 3P-CLR likelihood ratio scores (Racimo 2016).

Informed by the power analysis, we applied 3P-CLR to the central–eastern–Nigeria-Cameroon 3-population tree, using the genotypes from high-coverage autosomal genomes of 47 chimpanzees (18 central, 19 eastern and 10 Nigeria-Cameroon) (de Manuel et al. 2016), sliding windows of size 0.25 centiMorgans (cM) and the recombination map of Auton *et al.*, (Auton et al. 2012). The windows with the highest 3P-CLR scores in the genome-wide empirical distribution have the strongest evidence of positive selection at this time depth. We thus consider candidate targets of positive selection the windows with the highest 99.5%, 99.9% and 99.95% 3P-CLR likelihood ratio scores in the genome, which correspond to the 0.5% (n=4090), 0.1% (n=818) and 0.05% (n=409) tails of the empirical distribution. The candidate windows have additional signatures that, while not fully independent from the 3P-CLR signatures, are expected under positive selection in the central-eastern ancestor: they contain a marked excess of sites with high derived allele frequencies (DAF) in central and eastern (Figure 2A), and an excess of highly differentiated SNPs between the central-eastern clade and Nigeria-Cameroon (Figure 2B). While we cannot discard the presence of some false positives, these genomic regions are prime candidates to have mediated genetic adaptations in the central-eastern ancestral population.

Figure 2. Site frequency patterns of SNPs in candidate windows. Allele frequencies of SNPs genome-wide (grey) and at the most stringent 3P-CLR tail threshold (0.05% candidates for positive selection in the central-eastern ancestor, orange). **A**: Unfolded SFS for central, eastern and central and eastern combined. The X axis is limited to focus on high-frequency derived alleles, full SFS in Figure S2, SFS at all 3P-CLR tail thresholds in Figure S3. **B**: Absolute DAF difference between central-eastern and Nigeria-Cameroon.

SIV-related selection in the central-eastern ancestor

Under the hypothesis that SIV was a strong driver of adaptive evolution in the central-eastern ancestor, genes with signatures of positive selection would fall in known SIV-related functions more often than expected by chance. We test this expectation with enrichment tests for existing SIV/HIV-related categories on the genes that overlap our candidate windows, using Gowinda (Kofler and Schlotterer 2012). Gowinda tests for overrepresentation of a given gene set in our candidate windows compared to the expectation under neutrality (Kofler and Schlotterer 2012) (see Methods).

First we investigate selection on the immune response to SIV, by testing for an enrichment among the candidates in SIV-response genes, which in natural hosts change expression after SIV infection (Jacquelin et al. 2014; Jacquelin et al. 2009) and whose concerted action is thought to control SIV infection in natural hosts (Jacquelin et al. 2014; Jacquelin et al. 2009). SIV-response genes are enriched in signatures of positive selection in eastern chimpanzees (Schmidt et al. 2019) and vervet monkeys (Svardal et al. 2017), suggesting that they mediate adaptation to SIV. We find that the strongest candidate targets of positive selection in the central-eastern ancestor are modestly enriched in SIV-response genes, (0.05% candidates threshold, 11.8 expected, 18 observed, p-value=0.043; Figure 3). This signature can be refined by exploring enrichment in the 33 modules of differentially expressed genes that co-express temporally during SIV infection, defined by (Svardal et al. 2017). Two SIV co-expression modules are significantly enriched among candidate genes (Figure 3), both of which are defined by an acute response to SIV infection six days post-infection and exhibit strong signatures of positive selection in vervet monkeys (Svardal et al. 2017). Six days post-infection corresponds to when SIV can be first detected and when the immune response is typically initiated in natural hosts (Svardal et al. 2017).

Next we investigate selection on host-virus physical protein interactions, by testing whether the candidates are enriched in genes that encode Viral Interacting Proteins (VIPs) –host proteins that physically interact with viral proteins, viral RNA or viral DNA (Enard et al. 2016; Enard and Petrov

2018). HIV/SIV-interacting VIPs have not been shown to be enriched in signatures of positive
selection in chimpanzees or vervet monkeys, but VIPs are clear mediators of adaptation to viruses
(Enard et al. 2016; Enard and Petrov 2018). Testing enrichment in all the 152 defined VIP
categories, one of the strongest and most consistent enrichment signals is in VIPs that interact
with HIV/SIV, across all candidate cut-offs (0.5% candidates threshold, p-value=0.00046; 0.1%
candidates threshold, p-value=0.00014; 0.05% candidates threshold, p-value=0.00398; FDR
values given in Figure 3). Interestingly, we find similarly strong enrichment in one additional
category: influenza-interacting VIPs (0.5% candidates threshold, p-value=0.00002; 0.1%
candidates threshold, p-value=0.00010; 0.05% candidates threshold, p-value=0.00014; FDR
values given in Figure 3), in agreement with recent work in humans suggesting that RNA viruses
are an important selective force in mammals (Enard et al. 2016). Given the differences in gene
set sizes between the influenza and SIV/HIV (984 vs 810 genes) we consider the enrichment
results largely similar. At the 0.05% tail for influenza we expect 4.568 genes and observe 14 (p-
value = 0.00014, FDR=0.00112), whereas for HIV/SIV we expect 4.323 genes and observe 11
(p-value=0.00398, FDR=0.01854). Of note, these two categories overlap substantially, with 36%
of the 0.05% tail candidate genes in the influenza VIP set being also HIV/SIV VIPs, which makes
it difficult to establish the independence of their signatures. Not surprisingly other VIP categories
also show some evidence of positive selection. Ebola is an interesting example, although it is a
very small category with substantial overlap with HIV/SIV: three of the four 0.05% tail candidate
genes in the ebola VIPs set are also HIV/SIV VIPs. Thus, evidence for other VIP categories exist,
but together our results point to HIV/SIV as a particularly important selective force in
chimpanzees.

Finally, we explore other biological categories in a hypothesis-free analysis. An enrichment test
of GO categories (Ashburner et al. 2000) reveals 25 significantly enriched GO categories, two of
which are relevant to host-viral interactions: 'IκB/NFκB complex' (0.5% candidates threshold, p-
value=0.00008) and 'positive regulation by host of viral transcription' (0.5% candidates threshold,
p-value=0.00002) (see rest of categories and FDR values in Figure S4). Notably, as discussed
below, these two categories are intimately involved in the host biology under SIV/HIV infection.
As expected, this analysis indicates likely adaptations to selective forces beyond SIV, although
no category is as consistently enriched across thresholds as the HIV/SIV VIPs or SIV-response
categories.

Thus, the potential targets of positive selection in the central-eastern ancestral population show
enrichment patterns expected under adaptation to SIV/HIV and point to physical protein

interactions (VIPs) and initial immune reaction (SIV-response) as likely being key in early
adaptations to zoonotic SIV. Interestingly, the candidate windows are not enriched in existing HIV-
related GWAS hits when enrichment is investigated with Gowinda as above. This suggests that
while association studies of AIDS-related traits can identify genetic variants involved in clinical
phenotypes (often with treatment), they do not reveal the genetic bases of the biological
mechanisms that may allow natural hosts to control the pathogenicity of the virus.

Figure 3. Enrichment of HIV/SIV-related and VIP categories in 3P-CLR candidate genes for
the central-eastern ancestor. Columns for the three 3P-CLR quantiles, rows for each category
that is significantly enriched in at least one quantile (FDR<0.05). From top to bottom: SIV
response genes (Jacquelin et al. 2014, 2009), SIV co-expression modules (Svardal et al. 2017)
and VIPs (Enard et al. 2016; Enard and Petrov 2018). For the SIV-response gene set there is
only one category, hence we consider p-value<0.05 as significant, following (Schmidt et al. 2019).
Colours represent FDR values, with grey representing categories undetected in that particular
quantile. Stars mark categories with FDR < 0.05.

Potential sweeps of non-synonymous variants in *CD4*

A fundamental gene in SIV pathology is *CD4*, which encodes the glycoprotein required for
HIV/SIV cell entry, additional to a chemokine coreceptor (Blumenthal et al. 2012). Excitingly, *CD4*
lies within a candidate window at the 0.5% 3P-CLR threshold. In fact, *CD4*'s protein-coding
genomic region shows not only significant 3P-CLR signatures, but also SNPs with high derived
allele frequencies in the central-eastern clade that are also highly differentiated compared with

Nigeria-Cameroon (Figure 4A), as well as multiple PBSnj peaks both in central and eastern
chimpanzees (data from Schmidt et al. (Schmidt et al. 2019)). These signatures provide support
for positive selection driving the evolutionary history of the gene in chimpanzees. Bibollet-Ruche
et al. (Bibollet-Ruche et al. 2019) have shown that CD4 harbours functionally relevant variants in
chimpanzees, so to integrate both types of information and better understand the gene's
signatures of selection we aimed to localise the most likely selected variant(s). We focused on
SNPs with the highest allele frequency differentiation between the two chimpanzee clades
(central-eastern and Nigeria-Cameroon-western), as they are prime candidates to explain the
selective sweeps in the central-eastern ancestor identified by 3P-CLR (see Methods). We
identified these SNPs with PBSnj in the internal branch (representing the central-eastern
ancestor) with data from (Schmidt et al. 2019).

Two SNPs within the candidate window in *CD4* encode amino acids found to determine SIV
infectivity of chimpanzee cells by Bibollet-Ruche *et al.,* variants P68T and V55I (Bibollet-Ruche
et al. 2019). The P68T variant (chr12:6978232) falls in the tail of the empirical PBSnj distribution
(PBSnj p-value=0.0066), and is among the SNPs with the highest PBSnj value in this window. It
is also one of only four CD4 amino acid polymorphisms known to determine SIV infectivity of
chimpanzee cells (Bibollet-Ruche et al. 2019) being located, in the folded protein, near the binding
site between CD4 and SIV's *env* protein (Figure 4B). Central and eastern chimpanzees carry the
ancestral P allele, able to inhibit SIVcpz. The P68T variant is a prime candidate to drive the
selective sweep we observe in the central-eastern ancestor, though it remains possible that
selection acted on a different genetic variant, or on multiple. The V55I variant (chr12:6978193)
(Figure 4A) is not strongly differentiated in this branch (PBSnj p-value=0.1217), but has PBSnj
signatures of positive selection in central chimpanzees (central PBSnj p-value = 0.0014). Like
P68T, in the folded protein V55I is near the binding site between CD4 and SIV's *env* protein
(Figure 4B), and it is likely also functionally relevant, though the evidence from (Bibollet-Ruche et
al. 2019) is weaker than for P68T. We note that in addition, a splice site region synonymous
variant (chr12:6963043) is also among the most highly differentiated SNPs in the internal branch
according to PBSnj (Figure 4B). It seems clear that CD4 being a key player in SIV infection, has
been targeted by positive selection recurrently and in complex ways. Together with the three
chemokine co-receptors showing signatures of positive selection in central chimpanzees
(Schmidt et al. 2019), this indicates that proteins involved in SIV cell entry have evolved under
adaptive evolution across chimpanzee populations.

Figure 4. Signatures of selective sweeps in the extended *CD4* locus. A. Genomic representation of 150 kb and a magnified view of *CD4* with signatures of positive selection from 3P-CLR, PBSnj and clade-DAF. Statistics are coloured by population. The vertical dotted lines indicate best candidate SNPs of interest. The pink line indicates a splice site synonymous variant at position 6963043 with signatures of positive selection in the central-eastern ancestor. The black dotted line indicates two SNPs 39bp apart: a missense SNP in position 6978193 (the V55I SNP) with signatures of positive selection in centrals, and a missense SNP at position 6978232 (the P68T SNP) with signatures of selection in the central-eastern ancestor. For the DAF of these candidate SNPs across chimpanzee subspecies see Figure S5. **B.** Protein structure diagram showing the first two CD4 domains (grey) bound to the HIV envelope protein (yellow) (Kwong et al. 1998) and the amino acids corresponding to the two missense SNPs of interest, 6978193 (V55I) in blue and 6978232 (P68T) in red.

SIV-mediated selection across populations

Our results thus indicate that SIV was a strong selective force in the common ancestor of central
and eastern chimpanzees. This adds to previous evidence of SIV-related adaptation both in the
central and eastern populations (Schmidt et al. 2019). To understand how chimpanzees have
been able to adapt to SIV thus requires integrating information across these populations. We
integrated selection scores across the three populations (3P-CLR for the central-eastern ancestral
population and previously computed PBSnj for the central and eastern subspecies (Schmidt et al.
2019), Figure 1) and tested as above if any SIV-related biological categories have been targeted
by positive selection across populations using Gowinda (Figure S7).

The VIPs HIV/SIV category shows nominal enrichment across all possible configurations of
populations: the central-eastern ancestor and central subspecies (p-value=0.00894, BH-
FDR=0.185535), the central-eastern ancestor and eastern subspecies (p-value=0.00662, BH-
FDR=0.1855350), and the three populations together (central-eastern ancestor, central and
eastern subspecies; p-value=0.0161, BH-FDR=0.2700533). None of these are significant after
correcting FDR values for the number of populations tested using a Benjamini and Hochberg (BH)
correction; however, Gowinda is underpowered in this scenario, as several genes show
signatures of positive selection in multiple lineages, but are counted only once. To address this
limitation, we developed a custom method, '*set_perm*', which extends the functionality of Gowinda
to test if the joint distribution of candidate loci from different populations reveals enrichment for
particular biological functions (Methods, Supplementary Note). Focusing on the three populations
together, *set_perm* reveals significant enrichment (after correcting FDR values for testing multiple
populations using a BH correction) for the VIPs HIV/SIV category (105.37901 expected, 145
observed, p-value=0.00003, BH FDR=0.00159), SIV-response genes (276.44037 expected, 308
observed, p-value=0.01678, BH FDR=0.04604) and the green SIV co-expression module (8.4254
expected, 20 observed, p-value=0.00039, BH FDR=0.02241) (Figure S7). The higher power of
set_perm when genes have signatures in several populations means that additional categories
also become significant, including VIPs for influenza (112.18832 expected, 145 observed, p-
value=0.00061, BH FDR=0.01815) and VIPs for dengue viruses (9.40297 expected, 20 observed,
p-value=0.00083, BH FDR=0.02742).

To explore more generally which biological processes have been targeted by positive selection,
we also tested for enrichment of signatures of positive selection in KEGG pathways (Qiu 2013).

The KEGG pathway database consists of manually drawn molecular interaction diagrams for a
wide range of biological pathways (Qiu 2013). Using Gowinda, only one pathway exhibits nominal
enrichment in the combined candidates from the three populations (central-eastern ancestor,
central and eastern subspecies): "Th1/Th2 cell differentiation pathway" (10.6 expected, 22
observed, p-value=0.00056, BH-FDR=0.25552) (Figure S8 and Table S1). Underpinning this
signal are 22 genes: five with signatures of positive selection in the central-eastern ancestor,
eleven in eastern and nine in central chimpanzees, of which three genes are candidates both in
eastern and central chimpanzees (Figure 5). This result is confirmed and strengthened using
set_perm, where this pathway shows a strongly significant enrichment even after correcting for
testing multiple populations: ("Th1/Th2 cell differentiation pathway", 10.2 expected, 25 observed,
p-value=0.00002, BH-FDR=0.013800) (Figure S9). The similar contribution of signatures of
selection in the three populations shows that the Th1/Th2 pathway has likely repeatedly
undergone selection through time, although the individual genes targeted have largely differed
across the three populations. This pathway is intimately associated with SIV infection, being active
only in CD4+ T cells, which are the only cells that SIV infects (since CD4 is required for cell entry).
Moreover, SIV coordinates its replication with T cell activation, as differentiation of CD4+ T helper
cells involves upregulation of the transcription factors required to transcribe HIV/SIV genes
(Cullen 1991; Mingyan et al. 2009). By this process, HIV replication leads to pathogenesis via the
destruction of CD4+ T cells and the development of AIDS (Okoye and Picker 2013). Finally, the
Th1/Th2 differentiation pathway is critical to generate an efficient immune response against
viruses such as HIV/SIV (see discussion). Of note, multiple genes with signatures of positive
selection in this pathway encode for proteins that play critical roles in SIV/HIV infection (Figure
5). These include the transcription factors Nuclear Factor Kappa B (NFκB) and Nuclear Factor of
Activated T-cells (NFAT) (see Discussion).

The higher power of *set_perm* means that it reveals additional significantly enriched KEGG
categories (Figure S9). For example, in the three populations together the immune-related
'primary immunodeficiency' category becomes significant (2.0645 expected, 10 observed, p-
value=0.00003, BH FDR=0.006900), although it has only eight significant genes, half of them in
centrals. Interestingly 'Epstein-Barr virus infection' (18.09451 expected, 37 observed, p-
value=0.00003, BH FDR=0.013800) and 'leishmaniasis' (5.61591 expected, 17 observed, p-
value=0.00006, BH FDR=0.016848) pathways also show significant enrichment in the combined
three populations. These are not biological networks but rather pathogen-associated categories
that might possibly point to additional selective pressures. Still, we note that 69.2% of the

significant 'leishmaniasis' genes belong also to the larger Th1/Th2 pathway. Thus, to what extent

this signature is fully independent from those in the Th1/Th2 pathway is unclear.

In summary, our genome-wide analysis shows strong evidence of positive selection in the

common ancestor of central and eastern chimpanzees for SIV-response genes, SIV co-

expression modules, and HIV/SIV VIPs (together with VIPs for other viruses). Further, when

combining candidates from the central-eastern ancestor with those of its daughter species, we

find evidence of positive selection in each of these SIV-related categories, together with recurrent

positive selection in immune-related pathways, notably the "Th1/Th2 cell differentiation pathway".

Figure 5. Simplified diagram of the Th1/Th2 cell differentiation pathway, highlighting

proteins encoded by genes with signatures of positive selection. Proteins are coloured

according to the population(s) in which they show signatures of positive selection (see legend)

with white indicating proteins with no signatures of positive selection. Proteins in contact with

other proteins indicate protein complexes. *IL13* is a candidate in both central and eastern but is

not included in this simplified diagram. *RELA* is a candidate gene in the central-eastern ancestor

and codes for a protein subunit of both NFκB and c-Rel. This diagram was created with
BioRender.com.

Discussion

SIV is a formidable selective force, which has seemingly driven genetic adaptation in
chimpanzees since zoonotic infection until recent times. Here we identify adaptations to SIV in
the central-eastern ancestor. Together with our previous work showing evidence of positive
selection in multiple SIV-related genes in central and eastern chimpanzees, but not in western
and Nigeria-Cameroon (Schmidt et al. 2019), this supports the notion that zoonotic transmission
took place in the central-eastern ancestor and that SIV immediately became a strong and
continuous selective pressure in infected populations.

In the central-eastern ancestral population, loci with the strongest evidence of positive selection
in the genome are significantly enriched in two sets of genes that are intimately associated with
SIV, although in substantially different ways: genes that change expression upon SIV infection
('SIV-response genes' and 'SIV co-expression modules') and genes that encode proteins which
physically interact with HIV/SIV ('VIPs'). We also find evidence of positive selection driving
sequence evolution of CD4, a protein with a critical role in immunity which is necessary for SIV
cell entry. This reveals how diverse the initial adaptation to SIV likely was, consisting of genetic
changes in host proteins involved in the host response, physical interaction and cell entry of SIV.

Notably, only the VIPs show an enrichment in signatures of positive selection only in the ancestral
population, with the other categories showing evidence of selection at later times too. It is easy to
imagine why VIPs may be under strong selective pressure after first infection, as they physically
interact with the virus, while other mechanisms of adaptation may become more relevant at later
times.

Genes involved in the host response to SIV in natural hosts (SIV-response genes) show evidence
of positive selection not only in the central-eastern ancestral population, but also in eastern
chimpanzees (Schmidt et al. 2019). Combined, these observations suggest that the modulation
of the host response through gene expression evolves immediately after zoonotic transmission,
and continues over time. This idea agrees well with the fact that SIV-response genes evolve under
positive selection also in vervet monkeys (Svardal et al. 2017), which are long-term SIV natural
hosts. It is interesting that SIV-response genes were not identified experimentally in chimpanzees,

but in vervet monkeys, (Svardal et al. 2017; Jacquelin et al. 2014; Jacquelin et al. 2009). Their
apparent role in adaptation in both species is consistent with the idea that host tolerance to SIV
may evolve via similar mechanisms across distantly related species.

Genes that encode for the few proteins required by SIV to enter the cell also show evidence of
positive selection across populations. Bibollet-Ruche et al. (Bibollet-Ruche et al. 2019; Russell et
al. 2021) showed that CD4, which is the required primary receptor for SIV cell entry, harbours
multiple functionally relevant genetic variants. Complementing this important work, we find
evidence of positive selection in the central-eastern ancestor at CD4 with the most likely selected
variant, P68T, which we hypothesise underwent a soft sweep in the central-eastern ancestor,
being among only four CD4 missense variants shown to modify SIV infectivity in chimpanzee cells
(Bibollet-Ruche et al. 2019). Of note, with both alleles chimpanzee P68T inhibits SIVcpz
compared with the human CD4, though the derived T allele is most inhibitory (Bibollet-Ruche et
al. 2019). The T allele is fixed in western chimpanzees while Nigeria-Cameroon chimpanzees are
polymorphic (0.55) and bonobos and humans harbour the P allele, which is ancestral (Russell et
al. 2021; Bibollet-Ruche et al. 2019). It is thus likely that the site was polymorphic in the common
ancestor of all chimpanzees, making selection on P68 in the central-eastern ancestor likely to
have occurred on standing variation–although it remains possible that selection on a different
variant is responsible for the signatures of selection in CD4. It is nevertheless interesting that the
site is invariant in western chimpanzees. Bibollet-Ruche *et al.,* (Bibollet-Ruche et al. 2019)
proposed that fixation of the T allele in western chimpanzees could be due to adaptation to an
SIV-related retrovirus. This is an interesting hypothesis that we are able to test using a formal test
of positive selection in western chimpanzees, the PBSnj statistic in the western branch from
Schmidt et al., (Schmidt et al. 2019). Under our conservative 0.5% critical value western
chimpanzees show no significant evidence of positive selection at P68T (PBSnj western branch
p-value=0.0218). Still, this SNP falls within the 5% tail of the empirical distribution, and under less
stringent criteria would be considered to harbour signatures of positive selection. Thus, while we
cannot discard the possibility that the lack of diversity at P68T in western chimpanzees is
explained by random genetic drift, which is strong in this subspecies due to their small effective
population size and bottlenecks (de Manuel et al. 2016), our results suggest that the fixation of
the T allele in western chimpanzees could be due to positive selection. CD4 has a critical role in
immunity and has been shown to have some signatures of positive selection in species without
historical infection by SIV (e.g. humans (Meyerson et al. 2014)). Given the absence of natural SIV
infections in western chimpanzees it appears more likely that selection was due to a different
virus, perhaps a related retrovirus. In any case, we note that the allele frequency change in P68T

in western chimpanzees has no effect over the signatures of positive selection in CD4 in the
central-eastern ancestor (we used Nigeria-Cameroon as the outgroup).

Beyond P68T, *CD4* harbours signatures of positive selection in other SNPs and subspecies. As
mentioned in the Results, the V55I variant, which due to its position in the protein is likely to have
functional effects, has PBSnj signatures of positive selection in central chimpanzees. Further, a
C to T SNP within *CD4* (chr12:6973383) has significant signatures of positive selection in Nigeria-
Cameroon (PBSnj p-value = 0.002) (data from (Schmidt et al. 2019)). This adds to a picture of
multiple sweeps, both ancestral and lineage-specific, in *CD4*. Although outside of the scope of
this paper, an extensive analysis of this complex region that integrates functional and evolutionary
information would be extremely interesting. Remarkably, the multiple signatures of positive
selection in CD4 add to existing evidence of adaptation in central chimpanzees in *CCR3*, *CCR9*
and *CXCR6* (Schmidt et al. 2019), which mediate SIV/HIV cell entry (Elliott et al. 2015; Wetzel et
al. 2017; Nedellec et al. 2009) together with the primary receptor CD4*.*

Together, our results reveal that combining information across time and populations is critical to
understand chimpanzee adaptation to SIV. By integrating selection scores from the central-
eastern ancestor with those from the central and eastern subspecies, we identify strong
enrichment in the Th1/Th2 cell differentiation pathway. Twenty-two genes in the Th1/Th2
differentiation pathway have signatures of positive selection across time, three genes in two
separate populations. Together these observations show that this molecular pathway has been
repeatedly hit by positive selection over time. Why this pathway? First, the Th1/Th2 cell
differentiation pathway is critical for immunity against intracellular pathogens, including viruses.
Naïve CD4+ T cells recognise a MHC class II peptide, are activated and divide to give rise to
clone effector CD4+ T cells specific for that antigen (Zhang et al. 2014). CD4+ T cells can
differentiate into T helper type-1 (Th1), Th type-2 (Th2), or other T helper types, each with distinct
cytokine-secretion phenotypes, production of distinct interferons, and different downstream
immune responses (Zhang et al. 2014). By shaping which type of helper cell a CD4+ T cell will
differentiate into, this pathway has critical effects on immunity.

Second, the Th1/Th2 cell differentiation pathway is particularly relevant for HIV/SIV pathogenesis,
especially for control of viral replication. This is illustrated by the fact that a unique subset of
humans, 'HIV controllers', who are able to spontaneously control HIV infection without treatment,
are characterised by a Th1 differentiation bias (Vingert et al. 2012; Gomes et al. 2017).
Specifically, HIV controllers differentiate more naïve CD4+ T cells into Th1 than into other T helper
subsets (Gomes et al. 2017; Vingert et al. 2012). *In vitro* studies demonstrate that Th1 cells are

more resistant to HIV replication than Th2 cells, since their higher expression of *APOBEC3G*
limits reverse transcription and integration of HIV virions in Th1 better than in Th2 cells (Vetter
and D'Aquila 2009; Kervevan and Chakrabarti 2021). It thus seems natural that biasing the
Th1/Th2 differentiation pathway towards Th1 differentiation may be an efficient route to control
SIV/HIV pathogenicity.

Identified candidate targets of selection include key transcription factors required for SIV/HIV
replication: NFκB (in central-eastern ancestor) and NFAT (in easterns). NFAT proteins and HIV-
1 upregulate each other and may allow establishment of HIV-1 in the early stages of infection
(Pessler and Cron 2004). Both NFκB and NFAT bind to the same site which is identical in HIV-1
and HIV-2. We find signatures of positive selection in genes encoding proteins which inhibit (IκB
in centrals and easterns) and activate (IKK in central-eastern ancestor) the NFκB protein complex.
SIV/HIV have evolved to manipulate NFκB and IκB to minimise antiviral gene expression, while
allowing NFκB-induced viral transcription (Hotter et al. 2017; Langer et al. 2019; Sauter et al.
2015). Hence, genes encoding NFκB, IκB and their immediate interactors are plausible targets
for selection (Hotter et al. 2017; Langer et al. 2019; Sauter et al. 2015).

Our results contribute to the growing literature indicating that SIV has been and continues to be
a strong selection pressure in chimpanzee evolution. Previous studies have largely focused on
possible SIV-mediated selection by characterising variation at the MHC. Evidence of an ancient
selective sweep at MHC-I in the ancestor of chimpanzees and bonobos was initially suggested to
have been driven by SIV (de Groot et al. 2000; de Groot et al. 2002; de Groot et al. 2008) but is
now attributed to an 'SIV-like retrovirus', given such an ancient SIVcpz origin (and subsequent
loss of the virus in three lineages) is very unlikely (de Groot et al. 2017; Wroblewski et al. 2019).
Whereas, evidence that SIV drives adaptation at MHC-I in the very recent past has been found
by monitoring allele frequencies through time in eastern communities with high and low SIV loads
(Wroblewski et al. 2015). We do not find evidence of positive selection at MHC-I in the central-
eastern ancestor, although we likely have low power to identify signatures of selection in this
highly complex region. Further, it is likely that balancing selection is acting on the MHC, as
diversity at these sites can be protective against SIV and other viruses (Maibach et al. 2019),
further complicating the picture. We note that a key novelty of our study is that the genomic
dataset allowed us to go beyond the few genes known to play important roles in SIV/HIV infection
and perform formal tests for selection genome-wide. Our approach provides compelling evidence
of SIV-mediated selection only in central-eastern chimpanzees.

Although we have focused on selection in response to SIV, additional selective pressures were
surely relevant for the fitness of the central-eastern ancestor (and later populations). Here, in the
central-eastern ancestor, we also identify significant enrichments of signatures of selection in
VIPs that interact with influenza and other viruses, including ebolaviruses (Figure 3). Ebolaviruses
have been implicated in disease and mortality of wild chimpanzees (Formenty et al. 1999).
Influenza can infect captive chimpanzees (Buitendijk et al. 2014; Snyder et al. 1986; Kalter et al.
1997) and although respiratory and 'flu-like' diseases have been detected in wild chimpanzees,
we are not aware of any study that has specifically detected influenza in such cases (Negrey et
al. 2019; Hanamura et al. 2008; Kaur et al. 2008) and those would in any case represent
contemporary infections. Nevertheless, it is possible that an archaic influenza-like virus infected
chimpanzees thousands of years ago, leaving its mark in the genome in the same way as an
unknown archaic SARS-CoV-2-like virus has been proposed to leave a signature of genetic
adaptation in Asian human populations (Souilmi et al. 2021). In the future it would be interesting
to explore the biological consequences of other targets of selection identified in our candidate
windows.

Still, we find overwhelming evidence that SIV has been a strong selective force in central and
eastern chimpanzees, and their common ancestor. This includes initial adaptations in SIV-
interacting proteins, combined with adaptations on SIV response genes and cell entry genes that
continued into daughter populations, and with adaptations in the Th1/Th2 differentiation pathway
over time. This suggests the development of natural host immunity to SIV likely requires
adaptation in the host-virus interacting factors and the factors that mediate cell entry and the
immune response to infection. All of this strongly implies that SIV mediated selection has and
continues to be important in the evolution of central and eastern chimpanzees, and that we can
identify these adaptations with population genetics.

Methods

Simulations and Power Analysis

For 3P-CLR power analysis, we performed forward-in-time simulations in SLiM v.3.2 (Haller et al.
2019; Haller and Messer 2019). Following the inferred *Pan* demographic model (de Manuel et al.
2016; Won and Hey 2004) we simulated regions of length 1.2Mb, under neutrality and with
positive selection. We assumed a 25-year generation time (Langergraber et al. 2012), 0.96×10^{-8}
recombination rate (de Manuel et al. 2016; Won and Hey 2004) and a Wright-Fisher model. We

performed 1,000 neutral simulations and 1,000 simulations for both selection coefficients ($s = 0.1$,

0.05). Our selection simulations followed a hard sweep, conditional on fixation before the central-

eastern subspecies split (106 kya). Therefore, we sampled only a subset of all possible

evolutionary trajectories of the beneficial mutation. Following Racimo (Racimo 2016) every 10^{th}

SNP was a focal SNP, around which a 0.25cM window was centred, sliding every 10 SNPs. 100

SNPs were randomly sampled per window and used to calculate the 3P-CLR statistic, to test each

window for selection signatures. For computational efficiency we extended 3P-CLR to allow the

method to analyse 5kb segments of each chromosome. Results for the original and extended

source codes show the expected high correlation (Figure S10). For the distribution of 3P-CLR

scores for simulated under neutrality and positive selection see Figure S11. Nigeria-Cameroon

was used as an outgroup to the central and eastern target populations. We used Nigeria-

Cameroon rather than western who exhibit higher levels of drift and lower levels of segregating

polymorphism (de Manuel et al. 2016). Hence using western as the outgroup would reduce the

number of sites we could investigate, as 3P-CLR only considers sites which are segregating in

the outgroup (Racimo 2016). We also note that gene flow between Nigeria-Cameroon and central-

eastern chimpanzees may mask some signatures of selection, but would not generate them. We

produced ROC curves to visualise 3P-CLR's sensitivity and specificity, using pROC (Robin et al.

2011) in R v.3.5.2 [R Core Team 2018].

Identifying selection in the central-eastern ancestor

We analysed genomic data generated by De Manuel et al. (de Manuel et al. 2016), using the

same 3P-CLR parameters from our power analysis. This data consists of 47 genomes sequenced

to high coverage (mean 22.5-fold coverage per individual), which had been sampled from

chimpanzees of known subspecies: 18 central, 19 eastern, 10 Nigeria-Cameroon chimpanzees.

We used the EPO alignment to infer the ancestral allele for calculating derived allele frequencies.

We used the *Pan* diversity recombination map (Auton et al. 2012) to obtain genetic distances from

physical distances. For the empirical distribution of 3P-CLR scores genome-wide and in the tails

of the distribution see Figures S12-S13.

Post-hoc **SFS analysis of candidate windows in the central-eastern ancestor**

We used the VCF file from De Manuel et al. (de Manuel et al. 2016) to calculate the unfolded SFS

for the whole genome and for each 3P-CLR candidate window threshold. Sites were included only

if they were polymorphic when considering central, eastern and Nigeria-Cameroon i.e. western

specific variants were excluded. Combined central and eastern allele frequencies were calculated

by simply pooling the samples as each subspecies had almost identical sample sizes (central:
18, eastern: 19). The difference in allele frequencies between central-eastern and Nigeria-
Cameroon was calculated as the absolute difference in derived allele frequencies (DAF) between
central-eastern and Nigeria-Cameroon for each site. The minimum DAF at which we can identify
a polymorphism depends on the sample size. We have 18 central and 19 eastern samples, and
so the minimum DAFs possible are as follows - central: 0.028, eastern: 0.026, central-eastern
combined: 0.014.

Gene set enrichment analyses in the central-eastern ancestor
We tested our 3P-CLR candidate windows for gene set enrichments, using Gowinda v.1.12 (Kofler
and Schlötterer 2012), at the 3 highest quantiles of likelihood scores (0.5%, 0.1% and 0.05%
candidate windows), as power to detect significant gene enrichment varies with the number of
genes sampled. We tested for enrichment in different gene sets: SIV-response genes, SIV co-
expression modules, VIPs, GO categories and KEGG biological pathways (Svardal et al. 2017;
Enard et al. 2016; Enard and Petrov 2018; Ashburner et al. 2000; Jacquelin et al. 2014; Jacquelin
et al. 2009; Qiu 2013). Gowinda provides a p-value for the likelihood of finding the same evidence
of selection seen in the candidate genes in a random set of genes, when considering genes of
equivalent length and localisation in the genome (Kofler and Schlötterer 2012). These sets of
random genes provide the appropriate negative controls for the analysis. Gowinda also calculates
an FDR for each p-value. All genes tested were 1-1 human homologs and the analysis was run
in 'gene' mode.

Overlap with all PBSnj SNPs in the internal branch
To identify the SNPs with the strongest evidence of selection, we selected those with the largest
allele frequency difference between the central-eastern and the western-Nigeria-Cameroon
clades, within each significant 3P-CLR window at the 0.5% threshold. PBSnj (Schmidt et al. 2019)
was used to assess allele frequency difference between the two clades. We then filtered by the
20 SNPs with the highest PBSnj value per window. This results in 10,676,926 SNPs genome-
wide, which we annotated using ENSEMBL for pantro2.1.4, variant effect predictor (McLaren et
al. 2016) and regulomeDB (Boyle et al. 2012). Functional annotation of the 20 highest PBSnj
SNPs per window at the 0.5% threshold are shown in Figures S14-S15. SNPs of interest,
additional to those identified in *CD4,* are shown in Table S2. We also calculated the difference in
derived allele frequencies between the two clades (clade-DAF), as the difference in mean-

weighted derived allele frequencies. The positions of amino acids within the structure of CD4

coded for by SNPs of interest were plotted using iCn3D v3.1.1 structure viewer (Wang et al. 2020).

Identifying repeated targets of selection across branches

To investigate the targets of selection across time and populations, we combined selection

candidates identified in the central-eastern ancestor with those previously identified in the central

and the eastern subspecies by (Schmidt et al. 2019). To ensure each population is equally

represented in the analysis, we assigned a single empirical p-value to each gene for each

population and used genes in the 0.5% threshold for the central-eastern ancestor, to have an

equal number of candidate genes as for the subspecies. Total number of candidate genes per

population is as follows: in the central-eastern ancestor (817 genes), central $p < 0.000192$ (816

genes), eastern $p < 0.000228$ (806 genes). We tested for enrichment of our combined candidate

sets in the same gene sets as above: SIV-response genes, SIV co-expression modules, VIPs,

GO categories and KEGG biological pathways (Qiu 2013; Svardal et al. 2017; Jacquelin et al.

2014; Jacquelin et al. 2009; Enard et al. 2016; Enard and Petrov 2018; Ashburner et al. 2000)

using Gowinda (Kofler and Schlötterer 2012). However, Gowinda is underpowered when testing

for enrichment in multiple lineages.

In order to perform gene set enrichment tests on more than one lineage, we used a new method

developed by JMS. **Set_perm** implements a permutation-based enrichment test that for a single

lineage is equivalent to Gowinda (Kofler and Schlotterer 2012) but that correctly accounts for

genes showing signatures of natural selection in more than one lineage. Details of the method

can be found in the Supplementary Note and the code is available at

https://github.com/joshuamschmidt/set_perm).

Briefly, a joint test of lineages is performed by summing the number of candidate genes per gene

set across lineages. Thus, the total observed number of genes is simply the sum of the observed

number of genes across the different lineages, per gene set. Note, that this means that the same

gene can be counted more than once, if it is a candidate in more than one lineage.

Permutations are performed by first generating independent random gene sets for each lineage,

and then combining them into a joint set across all lineages. Thus, In this case, for each gene set

(S) and permutation (i), the joint permutation set, $joint_{Si}$, is obtained by summing the number of

genes (n) per lineage (k) i.e.

$joint_{Si} = \sum_1^k n_k$.

The p-values and FDR-corrected significance are calculated from permutation sets as described

for Gowinda (Kofler and Schlotterer 2012).

As above, any gene present in more than one independent permutation test will contribute more

than once to the joint permutation test.

Note, the number of observed genes may differ slightly between Gowinda and *set_perm*. This is

because, in order to perform joint lineage testing with Gowinda we define candidates at the gene

level, whereas with *set_perm* candidates are defined at the SNP level.

Acknowledgments

723   We thank Dean Cornish, Clàudia Fontserè, Sojung Han, Martin Kuhlwilm and Tomàs Marquès-

724   Bonet for helpful comments on the manuscript. HP was supported by a Formació de Personal

725   Investigador fellowship from Generalitat de Catalunya (FI_B100131), HO by a London NERC

726   DTP studentship (NE/S007229/1), and JS and AA by a UCL's Wellcome Trust ISSF3 award

727   (Grant Reference 204841/Z/16/Z).

References

Ashburner, Michael, Catherine A. Ball, Judith A. Blake, David Botstein, Heather Butler, J.
Michael Cherry, Allan P. Davis, et al. 2000. "Gene Ontology: Tool for the Unification of
Biology. The Gene Ontology Consortium." *Nature Genetics* 25 (1): 25–29.
Auton, Adam, Adi Fledel-Alon, Susanne Pfeifer, Oliver Venn, Laure Ségurel, Teresa Street,
Ellen M. Leffler, et al. 2012. "A Fine-Scale Chimpanzee Genetic Map from Population
Sequencing." *Science* 336 (6078): 193–98.
Bailes, Elizabeth, Feng Gao, Frederic Bibollet-Ruche, Valerie Courgnaud, Martine Peeters,
Preston A. Marx, Beatrice H. Hahn, and Paul M. Sharp. 2003. "Hybrid Origin of SIV in
Chimpanzees." *Science* 300 (5626): 1713.
Barouch, Dan H., Khader Ghneim, William J. Bosche, Yuan Li, Brian Berkemeier, Michael Hull,
Sanghamitra Bhattacharyya, et al. 2016. "Rapid Inflammasome Activation Following
Mucosal SIV Infection of Rhesus Monkeys." *Cell* 165 (3): 656–67.
Bell, Sidney M., and Trevor Bedford. 2017. "Modern-Day SIV Viral Diversity Generated by
Extensive Recombination and Cross-Species Transmission." *PLoS Pathogens* 13 (7):
e1006466.
Berger, Edward A. 1997. "HIV Entry and Tropism: The Chemokine Receptor Connection." *AIDS*
11 Suppl A: S3–16.
Bibollet-Ruche, Frederic, Ronnie M. Russell, Weimin Liu, Guillaume B. E. Stewart-Jones, Scott
Sherrill-Mix, Yingying Li, Gerald H. Learn, et al. 2019. "CD4 Receptor Diversity in
Chimpanzees Protects against SIV Infection." *Proceedings of the National Academy of
Sciences of the United States of America* 116 (8): 3229–38.
Blumenthal, Robert, Stewart Durell, and Mathias Viard. 2012. "HIV Entry and Envelope
Glycoprotein-Mediated Fusion." *The Journal of Biological Chemistry* 287 (49): 40841–49.
Boyle, Alan P., Eurie L. Hong, Manoj Hariharan, Yong Cheng, Marc A. Schaub, Maya Kasowski,

Konrad J. Karczewski, et al. 2012. "Annotation of Functional Variation in Personal Genomes Using RegulomeDB." *Genome Research* 22 (9): 1790–97.

Buitendijk, Hester, Zahra Fagrouch, Henk Niphuis, Willy M. Bogers, Kristin S. Warren, and Ernst J. Verschoor. 2014. "Retrospective Serology Study of Respiratory Virus Infections in Captive Great Apes." *Viruses* 6 (3): 1442–53.

Chahroudi, Ann, Steven E. Bosinger, Thomas H. Vanderford, Mirko Paiardini, and Guido Silvestri. 2012. "Natural SIV Hosts: Showing AIDS the Door." *Science* 335 (6073): 1188–93.

Chimpanzee Sequencing and Analysis Consortium. 2005. "Initial Sequence of the Chimpanzee Genome and Comparison with the Human Genome." *Nature* 437 (7055): 69–87.

Cullen, Bryan R. 1991. "Regulation of HIV-1 Gene Expression." *The FASEB Journal* 5 (10): 2361-68.

Elliott, Sarah T. C., Katherine S. Wetzel, Nicholas Francella, Steven Bryan, Dino C. Romero, Nadeene E. Riddick, Farida Shaheen, et al. 2015. "Dualtropic CXCR6/CCR5 Simian Immunodeficiency Virus (SIV) Infection of Sooty Mangabey Primary Lymphocytes: Distinct Coreceptor Use in Natural versus Pathogenic Hosts of SIV." *Journal of Virology* 89 (18): 9252-61.

Enard, David, Le Cai, Carina Gwennap, and Dmitri A. Petrov. 2016. "Viruses Are a Dominant Driver of Protein Adaptation in Mammals." *eLife* 5: e12469.

Enard, David, and Dmitri A. Petrov. 2018. "Evidence That RNA Viruses Drove Adaptive Introgression between Neanderthals and Modern Humans." *Cell* 175 (2): 360–71.e13.

Etienne, Lucie, Eric Nerrienet, Matthew LeBreton, Godwin Tafon Bibila, Yacouba Foupouapouognigni, Dominique Rousset, Ahmadou Nana, et al. 2011. "Characterization of a New Simian Immunodeficiency Virus Strain in a Naturally Infected Pan Troglodytes Troglodytes Chimpanzee with AIDS Related Symptoms." *Retrovirology* 8: 4.

Formenty, Pierre, Christophe Boesch, Monique Wyers, Claudia Steiner, Franca Donati, Frédéric Dind, Francine Walker, and Bernard Le Guenno. 1999. "Ebola Virus Outbreak among Wild Chimpanzees Living in a Rain Forest of Côte d'Ivoire." *The Journal of Infectious Diseases* 179 Suppl 1: S120–26.

Gilden, Raymond V., Larry O. Arthur, W. Gerard Robey, John C. Kelliher, Charles E. Graham, and Peter J. Fischinger. 1986. "HTLV-III Antibody in a Breeding Chimpanzee Not Experimentally Exposed to the Virus." *The Lancet* 1 (8482): 678–79.

Goldstein, Simoy, Ilnour Ourmanov, Charles R. Brown, Ronald Plishka, Alicia Buckler-White, Russell Byrum, and Vanessa M. Hirsch. 2005. "Plateau Levels of Viremia Correlate with the Degree of CD4+-T-Cell Loss in Simian Immunodeficiency Virus SIVagm-Infected Pigtailed Macaques: Variable Pathogenicity of Natural SIVagm Isolates." *Journal of Virology* 79 (8): 5153–62.

Gomes, Samara Tatielle M., Érica R. Gomes, Mike B. Dos Santos, Sandra S. Lima, Maria Alice F. Queiroz, Luiz Fernando A. Machado, Izaura M. V. Cayres-Vallinoto, Antonio Carlos R. Vallinoto, Marluísa de O Guimarães Ishak, and Ricardo Ishak. 2017. "Immunological and Virological Characterization of HIV-1 Viremia Controllers in the North Region of Brazil." *BMC Infectious Diseases* 17 (1): 381.

Greenwood, Edward J. D., Fabian Schmidt, Ivanela Kondova, Henk Niphuis, Vida L. Hodara, Leah Clissold, Kirsten McLay, et al. 2015. "Simian Immunodeficiency Virus Infection of Chimpanzees (Pan Troglodytes) Shares Features of Both Pathogenic and Non-Pathogenic Lentiviral Infections." *PLoS Pathogens* 11 (9): e1005146.

Groot, Natasja G. de, Corinne M. C. Heijmans, and Ronald E. Bontrop. 2017. "AIDS in Chimpanzees: The Role of MHC Genes." *Immunogenetics* 69 (8-9): 499–509.

Groot, Natasja G. de, Corrine M. C. Heijmans, Nanine de Groot, Nel Otting, Annemiek J. M. de Vos-Rouweller, Edmond J. Remarque, Maxime Bonhomme, Gaby G. M. Doxiadis, Brigitte Crouau-Roy, and Ronald E. Bontrop. 2008. "Pinpointing a Selective Sweep to the Chimpanzee MHC Class I Region by Comparative Genomics." *Molecular Ecology* 17 (8):

805   2074–88.

806 Groot, Natasja G. de, Nel Otting, Gaby G. M. Doxiadis, Sunita S. Balla-Jhagjhoorsingh,
807   Jonathan L. Heeney, Jon J. van Rood, Pascal Gagneux, and Ronald E. Bontrop. 2002.
808   "Evidence for an Ancient Selective Sweep in the MHC Class I Gene Repertoire of
809   Chimpanzees." *Proceedings of the National Academy of Sciences of the United States of
810   America* 99 (18): 11748–53.

811 Groot, Natasja G. de, Nel Otting, Rafael Argüello, David I. Watkins, Gaby G. Doxiadis, J.
812   Alejandro Madrigal, and Ronald E. Bontrop. 2000. "Major Histocompatibility Complex Class
813   I Diversity in a West African Chimpanzee Population: Implications for HIV Research."
814   *Immunogenetics* 51 (6): 398–409.

815 Haller, Benjamin C., Jared Galloway, Jerome Kelleher, Philipp W. Messer, and Peter L. Ralph.
816   2019. "Tree-Sequence Recording in SLiM Opens New Horizons for Forward-Time
817   Simulation of Whole Genomes." *Molecular Ecology Resources* 19 (2): 552–66.

818 Haller, Benjamin C., and Philipp W. Messer. 2019. "SLiM 3: Forward Genetic Simulations
819   Beyond the Wright-Fisher Model." *Molecular Biology and Evolution* 36 (3): 632–37.

820 Hanamura, Shunkichi, Mieko Kiyono, Magdalena Lukasik-Braum, Titus Mlengeya, Mariko
821   Fujimoto, Michio Nakamura, and Toshisada Nishida. 2008. "Chimpanzee Deaths at Mahale
822   Caused by a Flu-like Disease." *Primates; Journal of Primatology* 49 (1): 77–80.

823 Hotter, Dominik, Teresa Krabbe, Elisabeth Reith, Ali Gawanbacht, Nadia Rahm, Ahidjo Ayouba,
824   Benoît Van Driessche, et al. 2017. "Primate Lentiviruses Use at Least Three Alternative
825   Strategies to Suppress NF-κB-Mediated Immune Activation." *PLoS Pathogens* 13 (8):
826   e1006598.

827 Jacquelin, Béatrice, Véronique Mayau, Brice Targat, Anne-Sophie Liovat, Désirée Kunkel, Gaël
828   Petitjean, Marie-Agnès Dillies, et al. 2009. "Nonpathogenic SIV Infection of African Green
829   Monkeys Induces a Strong but Rapidly Controlled Type I IFN Response." *Journal of Clinical
830   Investigation* 119 (12): 3544-55.

831 Jacquelin, Béatrice, Gaël Petitjean, Désirée Kunkel, Anne-Sophie Liovat, Simon P. Jochems,
832   Kenneth A. Rogers, Mickaël J. Ploquin, et al. 2014. "Innate Immune Responses and Rapid
833   Control of Inflammation in African Green Monkeys Treated or Not with Interferon-Alpha
834   during Primary SIVagm Infection." *PLoS Pathogens* 10 (7): e1004241.

835 Kalter, Seymour S., Richard L. Heberling, Anthony W. Cooke, John D. Barry, Pei Y. Tian, and
836   William J. Northam. 1997. "Viral Infections of Nonhuman Primates." *Laboratory Animal
837   Science* 47 (5): 461–67.

838 Kaur, Taranjit, Jatinder Singh, Suxiang Tong, Charles Humphrey, Donna Clevenger, Wendy
839   Tan, Brian Szekely, et al. 2008. "Descriptive Epidemiology of Fatal Respiratory Outbreaks
840   and Detection of a Human-Related Metapneumovirus in Wild Chimpanzees (Pan
841   Troglodytes) at Mahale Mountains National Park, Western Tanzania." *American Journal of
842   Primatology* 70 (8): 755–65.

843 Keele, Brandon F., James Holland Jones, Karen A. Terio, Jacob D. Estes, Rebecca S. Rudicell,
844   Michael L. Wilson, Yingying Li, et al. 2009. "Increased Mortality and AIDS-like
845   Immunopathology in Wild Chimpanzees Infected with SIVcpz." *Nature* 460 (7254): 515–19.

846 Keele, Brandon F., Fran Van Heuverswyn, Yingying Li, Elizabeth Bailes, Jun Takehisa, Mario L.
847   Santiago, Frederic Bibollet-Ruche, et al. 2006. "Chimpanzee Reservoirs of Pandemic and
848   Nonpandemic HIV-1." *Science* 313 (5786): 523–26.

849 Kervevan, Jérôme, and Lisa A. Chakrabarti. 2021. "Role of CD4+ T Cells in the Control of Viral
850   Infections: Recent Advances and Open Questions." *International Journal of Molecular
851   Sciences* 22 (2): 523.

852 Kofler, Robert, and Christian Schlötterer. 2012. "Gowinda: Unbiased Analysis of Gene Set
853   Enrichment for Genome-Wide Association Studies." *Bioinformatics* 28 (15): 2084–85.

854 Kwong, Peter D., Richard Wyatt, James Robinson, Raymond W. Sweet, Joseph Sodroski, and
855   Wayne A. Hendrickson. 1998. "Structure of an HIV gp120 Envelope Glycoprotein in

Complex with the CD4 Receptor and a Neutralizing Human Antibody." *Nature* 393 (6686):
648-59.
Langergraber, Kevin E., Kay Prüfer, Carolyn Rowney, Christophe Boesch, Catherine Crockford,
Katie Fawcett, Eiji Inoue, et al. 2012. "Generation Times in Wild Chimpanzees and Gorillas
Suggest Earlier Divergence Times in Great Ape and Human Evolution." *Proceedings of the*
National Academy of Sciences of the United States of America 109 (39): 15716–21.
Langer, Simon, Christian Hammer, Kristina Hopfensperger, Lukas Klein, Dominik Hotter, Paul
D. De Jesus, Kristina M. Herbert, et al. 2019. "HIV-1 Vpu Is a Potent Transcriptional
Suppressor of NF-κB-Elicited Antiviral Immune Responses." *eLife* 8: e41930.
Leitner, Thomas, Marie-Christine Dazza, Michel Ekwalanga, Cristian Apetrei, and Sentob
Saragosti. 2007. "Sequence Diversity among Chimpanzee Simian Immunodeficiency
Viruses (SIVcpz) Suggests That SIVcpzPts Was Derived from SIVcpzPtt through Additional
Recombination Events." *AIDS Research and Human Retroviruses* 23 (9): 1114–18.
Locatelli, Sabrina, Ryan J. Harrigan, Paul R. Sesink Clee, Matthew W. Mitchell, Kurt A. McKean,
Thomas B. Smith, and Mary Katherine Gonder. 2016. "Why Are Nigeria-Cameroon
Chimpanzees (Pan Troglodytes Ellioti) Free of SIVcpz Infection?" *PloS One* 11 (8):
e0160788.
MacFie, Tammie S., Eric Nerrienet, Natasja G. de Groot, Ronald E. Bontrop, and Nicholas I.
Mundy. 2009. "Patterns of Diversity in HIV-Related Loci among Subspecies of
Chimpanzee: Concordance at CCR5 and Differences at CXCR4 and CX3CR1." *Molecular*
Biology and Evolution 26 (4): 719–27.
Ma, Dongzhu, Anna J. Jasinska, Felix Feyertag, Viskam Wijewardana, Jan Kristoff, Tianyu He,
Kevin Raehtz, et al. 2014. "Factors Associated with Siman Immunodeficiency Virus
Transmission in a Natural African Nonhuman Primate Host in the Wild." *Journal of Virology*
88(10): 5687-705.
Ma, Dongzhu, Anna J. Jasinska, Jan Kristoff, J. Paul Grobler, Trudy Turner, Yoon Jung,
Christopher Schmitt, et al. 2013. "SIVagm Infection in Wild African Green Monkeys from
South Africa: Epidemiology, Natural History, and Evolutionary Considerations." *PLoS*
Pathogens 9 (1): e1003011.
Maibach, Vincent, Kevin Langergraber, Fabian H. Leendertz, Roman M. Wittig, and Linda
Vigilant. 2019. "Differences in MHC-B Diversity and KIR Epitopes in Two Populations of
Wild Chimpanzees." *Immunogenetics* 71 (10): 617–33.
Mandell, Daniel T., Jan Kristoff, Thaidra Gaufin, Rajeev Gautam, Dongzhu Ma, Netanya
Sandler, George Haret-Richter, et al. 2014. "Pathogenic Features Associated with
Increased Virulence upon Simian Immunodeficiency Virus Cross-Species Transmission
from Natural Hosts." *Journal of Virology* 88 (12): 6778–92.
Manuel, Marc de, Martin Kuhlwilm, Peter Frandsen, Vitor C. Sousa, Tariq Desai, Javier Prado-
Martinez, Jessica Hernandez-Rodriguez, et al. 2016. "Chimpanzee Genomic Diversity
Reveals Ancient Admixture with Bonobos." *Science* 354 (6311): 477–81.
McLaren, William, Laurent Gil, Sarah E. Hunt, Harpreet Singh Riat, Graham R. S. Ritchie, Anja
Thormann, Paul Flicek, and Fiona Cunningham. 2016. "The Ensembl Variant Effect
Predictor." *Genome Biology* 17 (1): 122.
Meyerson, Nicholas R., Paul A. Rowley, Christina H. Swan, Dona T. Le, Gregory K. Wilkerson,
and Sara L. Sawyer. 2014. "Positive Selection of Primate Genes That Promote HIV-1
Replication." *Virology* 454-455 (April): 291–98.
Mingyan, Yu, Liu Xinyong, and Erik De Clercq. 2009. "NF-kappaB: The Inducible Factors of
HIV-1 Transcription and Their Inhibitors." *Mini Reviews in Medicinal Chemistry* 9 (1): 60–69.
Moore, John P., Scott G. Kitchen, Pavel Pugach, and Jerome A. Zack. 2004. "The CCR5 and
CXCR4 Coreceptors—Central to Understanding the Transmission and Pathogenesis of
Human Immunodeficiency Virus Type 1 Infection." *AIDS Research and Human*
Retroviruses 20 (1): 111-26.

Nedellec, Rebecca, Mia Coetzer, Naoki Shimizu, Hiroo Hoshino, Victoria R. Polonis, Lynn
908  Morris, Ulrika E. A. Mårtensson, James Binley, Julie Overbaugh, and Donald E. Mosier.
909  2009. "Virus Entry via the Alternative Coreceptors CCR3 and FPRL1 Differs by Human
910  Immunodeficiency Virus Type 1 Subtype." *Journal of Virology* 83 (17): 8353–63.
911 Negrey, Jacob D., Rachna B. Reddy, Erik J. Scully, Sarah Phillips-Garcia, Leah A. Owens,
912  Kevin E. Langergraber, John C. Mitani, et al. 2019. "Simultaneous Outbreaks of Respiratory
913  Disease in Wild Chimpanzees Caused by Distinct Viruses of Human Origin." *Emerging*
914  *Microbes & Infections* 8 (1): 139–49.
915 Okoye, Afam A., and Louis J. Picker. 2013. "CD4(+) T-Cell Depletion in HIV Infection:
916  Mechanisms of Immunological Failure." *Immunological Reviews* 254 (1): 54–64.
917 Pessler, Frank, and Randy Q. Cron. 2004. "Reciprocal Regulation of the Nuclear Factor of
918  Activated T Cells and HIV-1." *Genes and Immunity* 5 (3): 158–67.
919 Prado-Martinez, Javier, Peter H. Sudmant, Jeffrey M. Kidd, Heng Li, Joanna L. Kelley, Belen
920  Lorente-Galdos, Krishna R. Veeramah, et al. 2013. "Great Ape Genetic Diversity and
921  Population History." *Nature* 499 (7459): 471–75.
922 Prüfer, Kay, Cesare de Filippo, Steffi Grote, Fabrizio Mafessoni, Petra Korlević, Mateja
923  Hajdinjak, Benjamin Vernot, et al. 2017. "A High-Coverage Neandertal Genome from
924  Vindija Cave in Croatia." *Science* 358 (6363): 655–58.
925 Qiu, Yu-Qing. 2013. "KEGG Pathway Database." *Encyclopedia of Systems Biology*. Springer,
926  New York, NY. https://doi.org/10.1007/978-1-4419-9863-7_472.
927 Racimo, Fernando. 2016. "Testing for Ancient Selection Using Cross-Population Allele
928  Frequency Differentiation." *Genetics* 202 (2): 733–50.
929 Robin, Xavier, Natacha Turck, Alexandre Hainard, Natalia Tiberti, Frédérique Lisacek, Jean-
930  Charles Sanchez, and Markus Müller. 2011. "pROC: An Open-Source Package for R and S
931  to Analyze and Compare ROC Curves." *BMC Bioinformatics* 12: 77.
932 Rudicell, Rebecca S., James Holland Jones, Emily E. Wroblewski, Gerald H. Learn, Yingying Li,
933  Joel D. Robertson, Elizabeth Greengrass, et al. 2010. "Impact of Simian Immunodeficiency
934  Virus Infection on Chimpanzee Population Dynamics." *PLoS Pathogens* 6 (9): e1001116.
935 Russell, Ronnie M., Frederic Bibollet-Ruche, Weimin Liu, Scott Sherrill-Mix, Yingying Li, Jesse
936  Connell, Dorothy E. Loy, et al. 2021. "CD4 Receptor Diversity Represents an Ancient
937  Protection Mechanism against Primate Lentiviruses." *Proceedings of the National Academy*
938  *of Sciences of the United States of America* 118 (13): e2025914118.
939 Sauter, Daniel, Dominik Hotter, Benoît Van Driessche, Christina M. Stürzel, Silvia F. Kluge,
940  Steffen Wildum, Hangxing Yu, et al. 2015. "Differential Regulation of NF-κB-Mediated
941  Proviral and Antiviral Host Gene Expression by Primate Lentiviral Nef and Vpu Proteins."
942  *Cell Reports* 10 (4): 586–99.
943 Schmidt, Joshua M., Marc de Manuel, Tomas Marques-Bonet, Sergi Castellano, and Aida M.
944  Andrés. 2019. "The Impact of Genetic Adaptation on Chimpanzee Subspecies
945  Differentiation." *PLoS Genetics* 15 (11): e1008485.
946 Sharp, Paul M., George M. Shaw, and Beatrice H. Hahn. 2005. "Simian Immunodeficiency Virus
947  Infection of Chimpanzees." *Journal of Virology* 79 (7): 3891–3902.
948 Silvestri, Guido, Mirko Paiardini, Ivona Pandrea, Michael M. Lederman, and Donald L. Sodora.
949  2007. "Understanding the Benign Nature of SIV Infection in Natural Hosts." *The Journal of*
950  *Clinical Investigation* 117 (11): 3148–54.
951 Snyder, Mark H., William T. London, Eveline L. Tierney, Hunein F. Maassab, and Brian R.
952  Murphy. 1986. "Restricted Replication of a Cold-Adapted Reassortant Influenza A Virus in
953  the Lower Respiratory Tract of Chimpanzees." *The Journal of Infectious Diseases* 154 (2):
954  370–71.
955 Souilmi, Yassine, M. Elise Lauterbur, Ray Tobler, Christian D. Huber, Angad S. Johar, Shayli
956  Varasteh Moradi, Wayne A. Johnston, Nevan J. Krogan, Kirill Alexandrov, and David
957  Enard. 2021. "An Ancient Viral Epidemic Involving Host Coronavirus Interacting Genes

More than 20,000 Years Ago in East Asia." *Current Biology: CB* 31 (16): 3704.

Svardal, Hannes, Anna J. Jasinska, Cristian Apetrei, Giovanni Coppola, Yu Huang, Christopher
A. Schmitt, Beatrice Jacquelin, et al. 2017. "Ancient Hybridization and Strong Adaptation to
Viruses across African Vervet Monkey Populations." *Nature Genetics* 49 (12): 1705–13.

Vetter, Michael L., and Richard T. D'Aquila. 2009. "Cytoplasmic APOBEC3G Restricts Incoming
Vif-Positive Human Immunodeficiency Virus Type 1 and Increases Two-Long Terminal
Repeat Circle Formation in Activated T-Helper-Subtype Cells." *Journal of Virology* 83 (17):
8646–54.

Vingert, Benoît, Daniela Benati, Olivier Lambotte, Pierre de Truchis, Laurence Slama, Patricia
Jeannin, Moran Galperin, et al. 2012. "HIV Controllers Maintain a Population of Highly
Efficient Th1 Effector Cells in Contrast to Patients Treated in the Long Term." *Journal of
Virology* 86 (19): 10661-74.

Wang, Jiyao, Philippe Youkharibache, Dachuan Zhang, Christopher J. Lanczycki, Renata C.
Geer, Thomas Madej, Lon Phan, et al. 2020. "iCn3D, a Web-Based 3D Viewer for Sharing
1D/2D/3D Representations of Biomolecular Structures." *Bioinformatics* 36 (1): 131–35.

Wertheim, Joel O., and Michael Worobey. 2009. "Dating the Age of the SIV Lineages That Gave
Rise to HIV-1 and HIV-2." *PLoS Computational Biology* 5 (5): e1000377.

Wetzel, Katherine S., Yanjie Yi, Sarah T. C. Elliott, Dino Romero, Beatrice Jacquelin, Beatrice
H. Hahn, Michaela Muller-Trutwin, Cristian Apetrei, Ivona Pandrea, and Ronald G. Collman.
2017. "CXCR6-Mediated Simian Immunodeficiency Virus SIVagmSab Entry into Sabaeus
African Green Monkey Lymphocytes Implicates Widespread Use of Non-CCR5 Pathways in
Natural Host Infections." *Journal of Virology* 91 (4): e01626-16.

Won, Yong-Jin and Jody Hey 2004. "Divergence Population Genetics of Chimpanzees."
Molecular Biology and Evolution 22 (2): 297-307.

Wooding, Stephen, Anne C. Stone, Diane M. Dunn, Srinivas Mummidi, Lynn B. Jorde, Robert K.
Weiss, Sunil Ahuja, and Michael J. Bamshad. 2005. "Contrasting Effects of Natural
Selection on Human and Chimpanzee CC Chemokine Receptor 5." *American Journal of
Human Genetics* 76 (2): 291–301.

Worobey, Michael, Marlea Gemmel, Dirk E. Teuwen, Tamara Haselkorn, Kevin Kunstman,
Michael Bunce, Jean-Jacques Muyembe, et al. 2008. "Direct Evidence of Extensive
Diversity of HIV-1 in Kinshasa by 1960." *Nature* 455 (7213): 661–64.

Wroblewski, Emily E., Paul J. Norman, Lisbeth A. Guethlein, Rebecca S. Rudicell, Miguel A.
Ramirez, Yingying Li, Beatrice H. Hahn, Anne E. Pusey, and Peter Parham. 2015.
"Signature Patterns of MHC Diversity in Three Gombe Communities of Wild Chimpanzees
Reflect Fitness in Reproduction and Immune Defense against SIVcpz." *PLoS Biology* 13
(5): e1002144.

Wroblewski, Emily E., Peter Parham, and Lisbeth A. Guethlein. 2019. "Two to Tango: Co-
Evolution of Hominid Natural Killer Cell Receptors and MHC." *Frontiers in Immunology* 10:
177.

Zhang, Yuan, Yaguang Zhang, Wangpeng Gu, and Bing Sun. 2014. "TH1/TH2 Cell
Differentiation and Molecular Signals." *Advances in Experimental Medicine and Biology*
841: 15–44.

Supporting Information

1001

1002 **Supplementary Files**

1003

1004 **set_perm_SOM.docx: Supplementary note of *set_perm* method**

1005 **enrichment_tables.xlsx: Supplementary tables of gene set enrichment results**

1006 **SOM_RevisedMS_Pawar_ChimpSIV.docx: Supplementary figures S1-S15 and tables S1,**
1007 **S2**

1008

1009 **Supplementary Figures**
1010

1011 **Figure S1: Power of 3P-CLR in the ancestral central-eastern population**. Each ROC curve
1012 was generated from 1000 neutral simulations and 1000 selection simulations for each s (s=0.05,
1013 0.1).

1014

Figure S2: Full unfolded SFS for the whole genome and each 3P-CLR tail threshold in
centrals (top left), easterns (top right) and central-eastern combined (bottom). The central-
eastern combined SFS was made by simply pooling all the samples as both subspecies have
nearly identical sample sizes (central: 18, eastern: 19). The SFS are all indicative of selective
sweeps.

Figure S3. Site frequency spectrum of SNPs in candidate windows. Allele frequencies of
SNPs genome-wide and at different 3P-CLR tail thresholds. A: Unfolded SFS for central, eastern
and central and eastern combined. The X axis is limited to focus on high-frequency derived alleles
B: Absolute DAF difference between central-eastern and Nigeria-Cameroon.

1024

Figure S4: Enrichment of gene ontology (GO) categories across candidate genes at
different 3P-CLR quantiles in the central-eastern ancestor. Only categories with a significant
enrichment in at least one quantile (FDR<0.05) are shown. Categories are separated by GO
class: Biological Process, Cellular Component and Molecular Function. Colours represent FDR
values (red as highest significance). Grey represents instances where a GO category was
undetected in that particular quantile. Stars indicate a significant enrichment in that 3P-CLR
quantile.

1032

Figure S5: DAF of the three candidate SNPs of interest in *CD4* across chimpanzee
subspecies. These candidate SNPs correspond to those highlighted in Figure 4. SNP at
chr12:6963043 represents a splice variant with signatures of positive selection in the central-
eastern ancestor. SNPs at chr12:6978193 (V55I SNP) and chr12:6978232 (P68T SNP) are
missense variants with signatures of positive selection in centrals and the central-eastern
ancestor respectively.

1039

Figure S6: Enrichment in SIV-related, VIP and GO categories of candidate targets of
positive selection across populations, tested using Gowinda. Abbreviations indicate the
populations tested: central-eastern ancestor + central (A+C), central-eastern ancestor + eastern
(A+E), central-eastern ancestor + central + eastern (A+C+E). Categories are separated by
gene set tested: SIV responsive genes, SIV co-expression modules, VIPs and GO categories

(Svardal et al. 2017; Enard et al. 2016; Enard and Petrov 2018; Ashburner et al. 2000;
Jacquelin et al. 2014, 2009). Colours represent Benjamini and Hochberg corrected-FDR values
(red as highest significance). We note that no category reaches significant enrichment (after
correcting FDR values for the number of populations tested using a BH correction).

1049

Figure S7: Enrichment in SIV-related, and VIP categories of candidate targets of positive
selection across populations, tested using *set_perm*. Abbreviations indicate the populations
tested: central (C), eastern (E), central-eastern ancestor (A), central-eastern ancestor + central
(A+C), central-eastern ancestor + eastern (A+E), central-eastern ancestor + central + eastern
(A+C+E). The following categories are significantly enriched in the three populations together at
BH-corrected FDR < 0.05: SIV responsive genes, SIV co-expression green module, HIV/SIV
VIPs, influenza (IAV) VIPs and dengue (DENV) VIPs .All of these categories are also
significantly enriched when combining the central-eastern ancestor and the eastern subspecies.
The SIV co-expression green module is significantly enriched when combining the central-
eastern ancestor and the central subspecies. In the single lineages, we see significant
enrichment in the eastern subspecies in the SIV-responsive and HIV/SIV VIPs . While the
central-eastern ancestor is significantly enriched in the SIV co-expression green module,
HIV/SIV VIPs and influenza (IAV) VIPs .

1063

Figure S8: Enrichment in KEGG pathways of candidate targets of positive selection
across populations, tested using Gowinda. Abbreviations indicate the population(s) tested:
central-eastern ancestor (A), central-eastern ancestor + central + eastern (A+C+E). For the
central-eastern ancestor we used candidate genes in the least stringent quantile (0.5), to match
the number of candidates for the subspecies. Colours represent BH-corrected FDR values (red
as highest significance). We note that none of the KEGG pathways reaches BH-corrected FDR
< 0.05. The strongest enrichment is for the Th1 and Th2 cell differentiation pathway for A+C+E
with p-value=0.00056 and BH-FDR=0.25552, which represents a nominal enrichment.

1072

Figure S9: Enrichment in KEGG pathways of candidate targets of positive selection
across populations, tested using *set_perm*. Abbreviations indicate the populations tested:
central-eastern ancestor + central (A+C), central-eastern ancestor + eastern (A+E), central-
eastern ancestor + central + eastern (A+C+E). The following KEGG categories are significantly
enriched in the three populations together at BH-corrected FDR < 0.05: 'Th1 and Th2 cell
differentiation', 'primary immunodeficiency', 'Epstein-Barr virus infection' and 'leishmaniasis'. The
'primary immunodeficiency' and 'leishmaniasis' categories are also significantly enriched in the
central subspecies. Hence for the 'primary immunodeficiency' and 'leishmaniasis' gene sets,
the central lineage is likely driving the signal when the three populations are combined. When
the central-eastern ancestor and eastern subspecies are combined, we see significant
enrichment in the 'chromosome and associated proteins' pathway, which is also significantly
enriched in the central-eastern ancestor.

1085

Figure S10: Correlation between the 3P-CLR values with the original and extended code.
Likelihood scores are significantly correlated between the original and modified 3P-CLR source
code (rho=0.4565, p<2.2e-16). The absence of a perfect correlation is not due to differences in
the algorithm, but due to the sampling variance of SNPs in each window. Specifically, if more
than 100 SNPs are present within a given window, 3P-CLR chooses 100 SNPs at random.

Thus, the same SNPs will not be sampled each time the method is run. Variation in 3P-CLR
likelihood ratio scores therefore results if different SNPs were used to calculate the statistic. As
expected, the correlation is weaker for low 3P-CLR but high at higher 3P-CLR values, where
candidates of positive selection fall.

1095

1096 **Figure S11: Distribution of 3P-CLR scores** for data simulated under neutrality (A) and
1097 positive selection with selection coefficients of 0.05 (B) and 0.1 (C), 1000 replicates were
1098 generated in each case.

1099

1100 **Figure S12: Empirical distribution of 3P-CLR scores**. Vertical lines indicate the thresholds
1101 used to define candidate windows (0.5%, 0.1% and 0.05% tails of the empirical distribution).

1102

1103 **Figure S13: Distribution of 3P-CLR scores in the tails of the empirical distribution.** (A) the
1104 0.5%, (B) 0.1% and (C) 0.05% candidate quantiles, corresponding to 4090, 818 and 409
1105 genomic windows respectively.

1106

1107 **Figure S14: Functional annotation of 20 highest PBSnj SNPs per window at the 0.5%**
1108 **threshold using VEP**.

1109

1110 **Figure S15: Functional annotation of 20 highest PBSnj SNPs per window at the 0.5%**
1111 **threshold using regulomeDB**.

1112

1113 **Supplementary Tables**

1114

Table S1: Number of candidate genes from each population which belong to gene
categories of interest. The number of genes which are candidates in central only, eastern
only, and ancestral only are shown in the C, E and A columns respectively. The number of
genes which are candidates in both central and eastern, central and ancestral, eastern and
ancestral, or all three populations are shown in the C+E, C+A, E+A, and C+E+A columns
respectively.

1121

Table S2: SNPs of interest in genes identified within candidate windows of positive
selection at the 0.5% threshold in the central-eastern ancestor, additional to *CD4*. SNPs
with regulomeDB scores of 1f, 2a, 2b, 2c are considered to have putative significant regulatory
function, due to the lack of eQTL data for chimpanzees. Category indicates the gene set each
gene belongs to. Abbreviations: HCMV and ADV indicate human cytomegalovirus and
adenovirus respectively.