Alcohol reverses the effects of *KCNJ6* (GIRK2) noncoding variants on excitability of human glutamatergic neurons

Dina Popova[1], Isabel Gameiro-Ros[2], Mark M. Youssef[3], Petronio Zalamea[1], Ayeshia D. Morris[4], Iya Prytkova[2], Azadeh Jadali[3], Kelvin Y. Kwan[3], Chella Kamarajan[5], Jessica E. Salvatore[6], Xiaoling Xuei[7], David B. Chorlian[5], Bernice Porjesz[5], Samuel Kuperman[8], Danielle M. Dick[9], Alison Goate[10], Howard J. Edenberg[11], Jay A. Tischfield[1], Zhiping P. Pang[1,12], Paul A. Slesinger[2] and Ronald P. Hart[1,3,*]

Affiliations:

[1]Human Genetics Institute, Rutgers University, Piscataway, NJ 08854

[2]Nash Family Department of Neuroscience, Icahn School of Medicine at Mount Sinai, New York, NY 10029

[3]Department of Cell Biology & Neuroscience, Rutgers University, Piscataway, NJ 08854

[4]Joint Program in Toxicology, Rutgers University, Piscataway, NJ 08854

[5]Dept. of Psychiatry & Behavioral Sciences, SUNY Downstate Health Sciences University, Brooklyn, NY 11203

[6]Department of Psychiatry, Rutgers Robert Wood Johnson Medical School, Rutgers University, Piscataway, NJ 08854

[7]Department of Medical and Molecular Genetics, Indiana University School of Medicine, Indianapolis, IN 46202

[8]Department of Psychiatry, University of Iowa Carver College of Medicine, Iowa City, IA 52242

[9]Rutgers Addiction Research Center, Robert Wood Johnson Medical School, Rutgers University, Piscataway, NJ 08854

[10]Department of Genetics & Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, NY 10029

[11]Department of Biochemistry and Molecular Biology, Indiana Univ School of Medicine, Indianapolis, IN 46202

[12]Child Health Institute, Robert Wood Johnson Medical School, Rutgers University, New Brunswick, NJ 08854

*Corresponding author:
Rutgers University
Department of Cell Biology & Neuroscience
604 Allison Rd
Piscataway, NJ 08854

(848) 445-1783
rhart@rutgers.edu

Abstract

Synonymous and noncoding single nucleotide polymorphisms (SNPs) in the *KCNJ6* gene, encoding G protein-gated inwardly rectifying potassium (GIRK2) channel subunit 2, have been linked with increased electroencephalographic frontal theta event-related oscillations (ERO) in subjects diagnosed with alcohol use disorder (AUD). To identify molecular and cellular mechanisms while retaining the appropriate genetic background, we generated induced excitatory glutamatergic neurons (iN) from iPSCs derived from four AUD-diagnosed subjects with *KCNJ6* variants ('Affected: AF') and four control subjects without variants ('Unaffected: UN'). Neurons were analyzed for changes in gene expression, morphology, excitability and physiological properties. Single cell RNA sequencing suggests that *KCNJ6* AF variant neurons have altered patterns of synaptic transmission and cell projection morphogenesis. Results confirm that AF neurons express lower levels of GIRK2, have greater neurite area, and elevated excitability. Interestingly, exposure to intoxicating concentrations of ethanol induces GIRK2 expression and reverses functional effects in AF neurons. Ectopic overexpression of GIRK2 alone mimics the effect of ethanol to normalize induced excitability. We conclude that *KCNJ6* variants decrease GIRK2 expression and increase excitability and that this effect can be minimized or reduced with ethanol.

Introduction

Alcohol use disorder (AUD) is a heritable (h^2 = 0.49) condition characterized by an impaired ability to stop or control alcohol use despite adverse social, occupational, or health consequences[1]. An estimated 14.1 million American adults (ages 18+) were diagnosed with AUD in 2019 and the numbers continue to grow, particularly due to the COVID-19 pandemic[2]. Even though several evidence-based treatments are available for AUD[3-5], there is a high discrepancy in treatment outcomes, suggesting the existence of a variety of traits influencing the development of specific physiological and behavioral responses to alcohol. Understanding biological factors, specifically genetic risk, at the level of neuronal function is fundamental to developing tailored preventive interventions and better matching of patients to treatment.

The Collaborative Study on the Genetics of Alcoholism (COGA) collected a diverse set of phenotypes, including electroencephalogram (EEG) parameters, in the search for genes and endophenotypes associated with AUD[6, 7]. A family-based, genome-wide association study (GWAS) of a frontal theta event related oscillation (ERO) phenotype identified an association with several single nucleotide polymorphisms (SNPs), including a synonymous SNP, rs702859, in the *KCNJ6* gene on chromosome 21[8]. More recent studies concluded that the rs702859 polymorphism influences the magnitude and topography of ERO theta power during reward processing in a monetary gambling task, reflecting a genetic link to neuronal circuits[9, 10]. Two other SNPs (rs702860 and rs2835872) within *KCNJ6* were linked with the ERO endophenotype and AUD, but are noncoding, either intronic or within the 3' untranslated region (3' UTR), respectively. Additional studies identified various genetic loci correlating with selected EEG results but all consistently include alcohol dependence[11, 12]. To understand how these SNPs might influence these phenotypes, it is essential to characterize the pathways and mechanisms by which genetic risk unfolds at molecular, cellular and network resolutions. Furthermore, since the *KCNJ6* SNPs are non-coding or synonymous, the optimal strategy would incorporate human genetic backgrounds of actual subjects to preserve potential effects of noncoding sequences.

KCNJ6 encodes the G protein-gated inwardly rectifying potassium channel subunit 2 (GIRK2), which, when assembled into a GIRK channel, plays a role in regulating cell excitability[13]. The inward rectifying properties of GIRK are based on the ability to conduct potassium ions into the cell more easily than out, leading to a reduction in excitability[14, 15]. GIRK function in neurons is associated with activation of G protein-coupled receptors (GPCRs), which leads to dissociation of Gβγ subunits from the G protein complex and binding to the channel[16]. In addition, this family of GIRK channels can be potentiated by ethanol concentrations (20-50 mM) relevant to human alcohol usage through direct interaction with a hydrophobic pocket on the channel[17, 18]. Studies in model organisms also reveal the importance of GIRK function in AUD: mice lacking GIRK2 demonstrate reduced ethanol analgesia, greater ethanol-stimulated activity in open-field tests, increased self-administration of ethanol, and failure to develop a conditioned place preference for ethanol[19-21]. Thus, GIRK channels not only modulate the excitability of the neurons but also play an essential role in regulating responses to alcohol.

Recent studies have employed the use of subject-specific induced stem cell technology to
model, characterize and elucidate mechanisms underlying various types of addiction disorders
including AUD[22-25]. The generation of specific subtypes of cultured human neurons enables the
study of formerly inaccessible functional properties using standard neuroscience methods[22, 24-
26]. The contribution of various *KCNJ6* SNPs was found to affect a variety of properties
associated with neuronal function or sensitivity to drugs of abuse, suggesting targets for
potential therapeutic interventions. Associating specific genetic variants with mechanisms
contributing to addictive behaviors may produce more effective therapies tailored to individual
genotypes.

We hypothesize that noncoding SNP variants in *KCNJ6* alter neuronal excitability, potentially
contributing directly or indirectly to network-level endophenotypes such as ERO. Furthermore,
since ethanol interacts directly with GIRK2, we predict that excitability will be modulated by
ethanol. To test these hypotheses, we evaluated a human neuron model system that includes
variant genotypes, so that specific intergenic or noncoding sequences as well as modifying
genetic background influencing these functions would be preserved. Therefore, we selected
subjects from the NIAAA/COGA Sharing Repository, four with the ERO-associated allelic variant
in *KCNJ6* and the presence of diagnosed alcohol dependence and four without the variant and
unaffected for AUD and produced eight iPSC lines. iPSCs were reprogrammed into excitatory
human neurons (iN) to investigate the potential contribution of the *KCNJ6* SNPs to AUD
diagnosis and the ERO power endophenotype (Fig. 1A). We found that neurons from *KCNJ6*
variant AUD-affected individuals demonstrated initial transcriptomic and morpho-physiological
differences, mostly affecting excitability of the cells, which were paralleled by differences in
GIRK2 expression levels. Ethanol exposure, conversely, induced GIRK2 expression, ameliorating
differences in excitability. Moreover, by overexpressing *KCNJ6* we replicated effects of ethanol
on neuronal excitability. The results promote a better understanding of the association
between genetic variants and brain-wide changes affecting AUD risk and can potentially be
used for development of personalized interventions.

Methods

Generation of human iPS cells and glutamatergic neurons (iN)

Subjects were selected from the NIAAA/COGA Sharing Repository of the Collaborative Study on the Genetics of Alcoholism (COGA) project with characteristics summarized in Table 1. Selection criteria consisted of *KCNJ6* SNP genotype, DSM IV diagnosis of AUD, sex, and availability of frozen lymphocytes in the repository. All subjects were of European ancestry. Cryopreserved lymphocytes were reprogrammed to iPSC using Sendai-viral-expressed transcription factors (Oct3/4, Sox2, Klf4 and c-Myc; Cytotune™-iPS Reprogramming Kit, ThermoFisher) by RUCDR Infinite Biologics® (now Sampled, Inc.). Pluripotent cells were selected by colony morphology, alkaline phosphatase positivity, and immunocytochemical expression of Oct4 and TRA-1-60. Pluripotency was confirmed by colony morphology and immunocytochemistry for Oct4 and Tra-1-60 markers (Supplemental Fig. 1A). Identity of each iPSC line was confirmed by matching a 96 SNP panel to results obtained from fresh blood upon banking the original sample. Cultures were tested for mycoplasma (all negative) prior to release by RUCDR. Identities of the SNPs listed in Table 1 were confirmed in all iPSC lines by PCR amplification from genomic DNA and Sanger sequencing. Genomic integrity was confirmed using e-Karyotyping[27] using RNAseq data, which detected no anomalies (Supplemental Fig. 1B). Frozen aliquots were obtained for each subject, thawed, expanded, and maintained for neurogenesis. All iPS lines were cultured up to 50 passages for these experiments.

The previously-described protocol for generating glutamatergic human induced neurons (iN)[28-30] was modified. To obtain cultures with enhanced levels of spontaneous activity, we minimized the time of Ngn2 induction to reduce alternate cell identities[31] and fed cultures with a low-molality medium for at least 30 days post-induction[32]. Briefly, iPS cells were plated as dissociated cells on Matrigel® Matrix (Corning Life Sciences)-coated dishes in iPS-Brew-XF (StemMACS) medium supplemented with 2 µM Y-27632 (Peprotech, Inc.) and 2 ng/µl doxycycline, and infected with lentiviruses expressing Ngn2 (AddGene #52047) and rtTA (FUW-M2rtTA; AddGene # 20342) for 10–12h. The next day, the culture medium was replaced with Neurobasal medium/B27/GlutaMAX™ (Life Technologies) supplemented with 2 µg/mL of doxycycline (MP Biomedical), 2 µM Y-27632 and 2 ng/µl puromycin. Puromycin selection was continued for 2 days, and on day 5, the induced neurons were dissociated with Accutase (STEMCELL Technologies) and plated on Matrigel-covered glass coverslips with a monolayer of primary astrocytes (passage 2-3) isolated from postnatal day 0-1 mouse pups, as described previously[33]. Following plating, 50% of Neurobasal medium was changed to Neurobasal Plus culture medium with ascorbic acid, GlutaMAX, B27 Plus and Culture One supplements (Life Technologies). Half of the Neurobasal Plus medium with all supplements was replaced every 2–3 days. Cultures of iN typically exhibited resting membrane potentials averaging -40 mV with the presence of synaptic markers, spontaneous action potentials, and synaptic activity.

Intermittent Ethanol Exposure (IEE).

Since ethanol evaporates during culture incubation (Supplemental Figure 7), we initially added ethanol to 20 mM and then replenished half the medium daily with 40 mM ethanol in medium. Ethanol concentrations were measured using and AM1 Alcohol Analyzer (Analox Instruments, Ltd.), as described previously[25].

Primary mouse cortical neuron cultures

P0-P1 mouse brains were obtained following decapitation on ice, meninges were removed, cortex was dissected and cells were dissociated by incubation in trypsin (Gibco) for 5-7 min at 37°C. The cells were then washed with Hank's balanced salt solution (HBS) and triturated by gentle pipetting. The cell suspension was centrifuged for 5 min at 160 x g and plated on Matrigel-coated glass coverslips in Neurobasal medium/B27/L-Glutamine (Life Technologies). When the density of glia in the culture reached ~40–50% (approximately 2 days after plating), 50% of the conditioned culture medium was replaced with fresh medium containing 4 mM Ara-C (Sigma; 2 mM final concentration). The cultures were maintained in medium containing 2 mM Ara-C until fixation at 13–18 days in vitro (DIV).

Generation of CRISPR/Cas9 Knockout iPSC

As a negative control for GIRK2 antibody validation we generated a *KCNJ6* knockout by CRISPR/Cas9 deletion in WTC-11, a publicly-available, control iPS line obtained from the Gladstone Institute[34]. Two optimal gRNA targeting sequences were selected (using http://crispr.mit.edu) to generate a frameshift 76-bp deletion, introducing three termination codons within 40 bp following the deletion. A well-characterized iPSC line, WTB[35], was obtained from the Gladstone Institute Stem Cell Core Facility (San Francisco, CA). Approximately 3×10^6 WTB iPSC (passage 58) were electroporated (amaxa Nucleofector) in suspension with a mixture of two plasmids constructed from pSpCas9(BB)-T2A-GFP [AddGene #48138[36]], with sequences targeting *KCNJ6*: G1, 5'-GATATCGGTCAGGTAGCGAT-3'; or G2, 5'-TCAGCCGAGATCGGACCAAA-3'. One day after transfection, cells were dissociated with Accutase and sorted to select GFP+ cells into KSR medium (Life Technologies) with 2 µM Y-27632 and FGF (Peprotech, Inc.) and then plated at low density into two wells of a 6-well plate. Approximately 7-10 days later, individual colonies were hand-picked into 96-well plates. Three days after plating, cultures were split to two 96-well plates. One plate was harvested at ~50-100% confluence for screening. Cells were lysed in Quickextract (Lucigen), heated to 68°C for 6 minutes followed by 98°C for 2 minutes. The lysate was PCR amplified (DET3-F: 5'- TGGACCCCAACACAGATTGG-3' and -R: 5'-TGGATCAGGACGTCGAAAGC-3'), treated with exoSAPIT (Life Technologies) and analyzed by agarose gel electrophoresis. From 96 colonies, approximately 19 appeared to have a single-band deletion of the predicted size. Aliquots of the PCR reactions were sequenced with the F primer to confirm. To confirm homozygosity, PCR products from four candidates were cloned into a plasmid (StrataClone Blunt PCR Cloning Kit, Agilent) and individual colonies sequenced to identify each allele. Only one candidate, A7, exhibited the designed homozygous 76 bp deletion. A second colony with no deletion, E11, was selected to serve as unedited control.

Bulk and single cell RNA sequencing

For bulk RNAseq, individual iN cultures were harvested in lysis buffer and column purified
(Quick-RNA kit, Zymo). Purified RNA was sent to Novogene, Inc., for sequencing. Fastq files
were downloaded, filtered with Fastp[37], aligned to human reference genome (hg38) with
HISAT2[38], SNPs in mapped *KCNJ6* sequence were extracted using samtools mpileup[39] and
bcftools to filter for quality. Ensembl VEP[40] was used to annotate SNPs. RNAseq data have been
deposited with the NIH GEO repository (accession number GSE196491).

For single-cell RNA sequencing (scRNAseq), iN cultures were treated with 20 mM ethanol for 7
days using an intermittent ethanol exposure (IEE) protocol where alcohol was replenished daily
(Supplemental Fig. 7)[25]. At 28 days after plating onto glia, iN were dissociated using trypLE
Express (Thermo Fisher) for 5 minutes at 37°C. The dissociated cells were centrifuged at 200 x g
for 5 minutes at 4°C. The supernatant was removed, and cells were washed twice with 1X HBSS
(Thermo Fisher). After washes, cells were resuspended in 1% BSA (Bovine Serum Albumen,
ThermoFisher)/1xPBS (ThermoFisher). Cells were counted and resuspended at 1200 cells/µl in
1% BSA/ 1xPBS and placed on ice. Single-cell cDNA libraries were generated using the
Chromium Next GEM Single Cell 3 GEM Reagent Kits v3.1 (10X Genomics). Approximately
20,000 dissociated cells were loaded onto a cassette in the Chromium Controller with
accompanying reagents to generate Gel Beads in Emulsions (GEMs). GEMs were subjected to
reverse transcription to generate single-cell cDNA libraries before the oil emulsion was
disrupted and cDNA purified using Dynabeads MyOne Silane (Thermo Fisher). cDNA was
amplified by PCR for 11 cycles and purified using SPRIselect reagent (Beckman Coulter). cDNA
fragmentation, A-tailing, and end repair was performed followed by adapter ligation for paired-
end sequencing. An additional 11 cycles of PCR were performed to incorporate the sample
index sequences and to amplify the libraries before samples were purified using SPRIselect
reagent and used for sequencing. Libraries were shipped to Psomagen, Inc., for sequencing
service. Fastq files were downloaded, aligned with reference genome (initially hg38 and mm10)
using CellRanger (10XGenomics) and then imported into a Seurat (v. 4.0.6) object[41]. A summary
of human (GRCh38) and mouse (mm10) characteristics is shown in Supplemental Figure 2A and
CellRanger parameters are provided in Supplemental Table 2. Clusters of cells expressing
primarily mouse genes and therefore consistent with the co-cultured mouse glia was identified
(Supplemental Figure 2B) and removed. Human cells were isolated for further analysis. Cells
from individual subjects were identified using genome-wide SNPs that were expressed in the
mRNA. Demuxlet[42] assigned "best guess" identification by cell barcode, which was imported as
metadata into the human Seurat object. The number of identified cells was plotted for each
cell line and treatment (Supplemental Figures 2C). Seurat clusters consistent with neuronal
identity (Figure 1C, Supplemental Figure 2E) were selected for further analysis. Total numbers
of neurons by subject and treatment are plotted in Supplemental Figure 2D. A minimum of 10
cells per subject and treatment were required for analysis—only subject 351 in the control
condition fell below this threshold.

Differential gene expression was performed on gene counts aggregated by subject

("pseudobulk" analysis). To exclude sporadically-expressed genes, detectable expression was

required in at least 25% of the cells in one group. After filtering, gene counts were summed for

each cell line and condition. Tables of summed gene counts were analyzed by DESeq2[43] using

the likelihood ratio test (LRT), comparing a model of groups to a reduced model. Contrasts

were calculated and significance required an adjusted p-value of 0.05 or less and at least a 1.5-

fold difference. Genes either up- or down-regulated were assessed for enriched gene ontology

biological process terms using the enrichGO function from the clusterProfiler package[44].

Summary data are listed in Supplemental Figure 2. These data have been deposited with the

NIH GEO archive, accession number GSE203530.

Fluorescent in situ hybridization (FISH)

Fluorescent, single molecule *in situ* hybridization was performed with the RNAscope® Multiplex

Fluorescent Detection Kit v2 (Advanced Cell Diagnostics, ACD) following the manufacturer's

instructions. In brief, iN cultures were fixed in 4% paraformaldehyde for 20 min at room

temperature, followed by a series of ethanol dehydration/rehydration steps, permeabilization

(with PBS-0.01% Tween buffer), blocking and overnight incubation with primary antibody

against MAP2 (Millipore AB5543, 1:100 dilution). Prior to probe hybridization, samples were

treated with hydrogen peroxide for 10 min and then with protease III (1:10) for 10 min. Human-

specific *KCNJ6* probe was hybridized for 2 h, followed by standard amplification and color

reaction with 520 Opal dye (Akoya Biosciences). MAP2 antibody was labeled with secondary

antibody (Alexa 633), nuclei were stained with DAPI, and coverslips were mounted on slides

using mounting medium (Thermo Fisher). Images were acquired on a Zeiss LSM 800 confocal

microscope using a 40x objective. *KCNJ6* was quantified by manually counting fluorescent

puncta within the MAP2/DAPI positive regions of individual neurons (defined as "somatic") or

as puncta scattered along the MAP2-associated processes ("non-somatic"). Levels of *KCNJ6*

were binned according to semi-quantitative histological scoring methodology by levels using

Advanced Cell Diagnostics scoring criteria, where low expression of RNA target was defined to

be 1-3 puncta per cell, intermediate 4-9, high 10-15 and very high >15 per cell.

Electrophysiology

Functional analyses of induced human neurons were performed using whole-cell patch-clamp

electrophysiology as previously described[24, 25]. Recordings were done in HEPES buffer consisting

of (in mM): 140 NaCl, 5 KCl, 2 $CaCl_2$, 2 $MgCl_2$, 10 HEPES, 10 Glucose. Osmolality of the buffer

was between 290-305 mOsm and the pH was adjusted to 7.4. Whole-cell patch-clamp

recordings were performed using borosilicate glass electrodes with resistances of 4-8 MΩ and

intracellular solution containing the following (in mM): 126 K-Gluconate, 4 KCl, 10 HEPES, 4

ATP-Mg, 0.3 GTP-Na_2, 10 Phosphocreatine, with pH adjusted to 7.2 and osmolarity to 270–290

mOsm. After patching, neurons were allowed to rest for 5-7 min, or until the RMP drift

stabilized, before recording. For GIRK2 channel recordings, an extracellular solution with high K^+

was used, which consisted of: 20 mM KCl, 140 mM NaCl, 2 mM $CaCl_2$, 2 mM $MgCl_2$ and 10 mM

HEPES (pH 7.4). GIRK-mediated currents were measured at -40 mV membrane potential
holding. Excitability of the neurons was measured by applying ramp (+200 pA) and step (-50 pA
− 100 pA) depolarizing pulses while holding neurons at -65mV (current clamp mode). All
recordings in cultured human neurons were performed at room temperature. Cells were
excluded from analysis if access resistance changed by more than 20% across the duration of
the recording. For current injection studies, neurons were excluded if RMP was close to -63mV
and the required current injection was less than 1 pA (absolute value). In addition, any induced
neurons whose access resistance was greater than 35 MΩ were also excluded. Drugs were
applied through a perfusion system in the following concentrations: 160 nM ML297 or 20 mM
alcohol. Clampfit software (pClamp 11; Molecular Devices) was used for analysis of recordings.

Immunocytochemistry and confocal imaging
iN cultures were fixed for 30 min in ice cold methanol and permeabilized using 0.2% Triton X-
100 in PBS for 15 min at room temperature. Cells were then incubated in blocking buffer (5%
BSA with 5% normal goat serum in PBS) for 30 min at room temperature and then incubated
with primary antibodies diluted in blocking buffer overnight at 4°C, washed with PBS three
times, and subsequently incubated with secondary antibodies for 1 h at room temperature.
Confocal imaging analysis was performed using a Zeiss LSM700. Primary antibodies used: rabbit
anti-GIRK2 (Alomone labs, APC-006, 1:400), mouse anti-βIII-tub (Bio legend, MMS-
435P,1:1000), chicken anti-MAP2 (Millipore AB5543, 1:1000), mouse anti-Syn1 (SYSY,106-011,
1:200), mouse anti-PSD 95 (SYSY, 124-011, 1:2000), mouse anti-mCherry (Thermofisher
Scientific, M11217, 1:100).

Validation of GIRK2 immunoreactivity
To validate the specificity of GIRK2 detection, GIRK2 was overexpressed by transduction with a
ubiquitin C (UbiC) promoter-driven, *KCNJ6*-expressing lentivirus or expression was eliminated
by introducing frameshift deletions into *KCNJ6* using CRISPR/Cas9-editing. Overexpression was
initiated by infection on day 27 post-induction and GIRK2 expression was evaluated 14 days
later. Levels of GIRK2 in iN prepared from the CRISPR/Cas9 knockout line were measured at a
similar time post-induction (~day 40). Viral GIRK2 transduction was confirmed by co-expression
of mCherry.

Image analysis
Cytometric analysis of confocal images of MAP2-stained iN cultures was performed using the
Fiji/ImageJ image analysis program[45]. Regions of Interest (ROIs) were created for each cell by
encapsulating the cell body through use of the cell magic wand tool. The cell body was then
quantified for area and shape descriptors. The NeurphologyJ Fiji macro[46] was utilized for
detection and quantification of extended neuronal morphometry. A threshold was selected
manually so that most neurites are well-defined without significant particle detection or noise.
GIRK2 puncta quantification used trainable Weka Segmentation[47] to segment the image to
establish two classes. One class identified pixels that corresponded to GIRK2 puncta for
intensity, circularity, and size. A second class identified pixels that did not correspond to puncta,

such as noise from neurons and background. The classifier was trained multiple times until
there was a high confidence that the pixels segmented into class one had a high (90%)
confidence interval denoted by probability maps. The segmented image was adjusted to
remove any false negatives. The class one binary image was then used for puncta cytometry.
ImageJ scripts can be downloaded from https://github.com/rhart604/imagej. The person
performing analysis was given coded images with no genotype or line information so that
analysis would be blinded.

Lentiviruses
FSW-hSyn-GCaMP6f assembly was described previously[48]. FUGW-KCNJ6-mCherry was
assembled using lentiviral backbone from FUGW (AddGene #14883), *KCNJ6* coding sequence
amplified from human neuron cDNA (forward primer: agccaggaaaagcacaaaga, reverse primer:
ggggagaagagaagggtttg), and mCherry with a nuclear localization signal from pME-nlsmCherry (a
gift from Dr. Kelvin Kwan). During construction, the GIRK2 protein-coding sequence was tagged
with a 3xHA tag and linked to mCherry with a T2A "self-cleaving" element.

Calcium imaging in iPSC-derived iN populations
For assessing neuronal excitability, iN co-cultured with mouse glia on Matrigel[TM] coated 10 mm
glass coverslips were transduced with lentivirus expressing hSyn-GCaMP6f at least 2 weeks
prior to imaging to ensure robust expression. All fluorescence imaging experiments were
performed in ~50 DIV (days in vitro) iNs. Coverslips were mounted on a diamond-shaped
chamber and placed on a Nikon Eclipse TE2000-U microscope with a 10X objective. GCaMP6f
was excited using a 480 nm (Mic-LED-480A, Prizmatix), a HQ480/40x excitation filter, a Q505LP
dichroic mirror, and a HQ535/50m emission filter (Semrock). Fluorescence was projected onto a
sCMOS Zyla 5.5 camera (Andor) and sampled at a rate of 4.7 fps with a frame exposure of 200
ms at 160x120 pixels (4x4 binning). Light source and sCMOS camera were controlled with the
Nikon Elements software (NIS-Elements AR 5.20.01).

Neurons were continuously perfused during fluorescence recording with ACSF with the
following composition (in mM): NaCl 125, KCl 5, D-Glucose 10, HEPES-Na 10, $CaCl_2$ 3.1 and
$MgCl_2$ 1.3. The pH was adjusted to 7.4 with HCl and osmolarity corrected with sucrose to 290-
300 mOsm. Depolarizing solutions were prepared in ACSF with 10 or 50 µM glutamate, or with
18 mM KCl. Perfusion was gravity fed (flow rate of 0.065 ml/s). Solution exchanges were
controlled with a ValveBank8 II (AutoMate Scientific Inc.). Calcium transients were recorded
using a protocol based on glutamate and KCl perfusion, that consisted of 3 min baseline activity
(ACSF), 3 x 30 s pulses of 10 µM glutamate (30 s ACSF between pulses), one 30 s pulse of 50 µM
glutamate, 2 min of ACSF wash, 3 x 30 s pulses of 18 mM KCl (30 s ACSF between pulses), and a
final ACSF wash for 5 min (Fig. 4C).

ROI segmentation of GCaMP6f-expressing neurons, raw fluorescence extraction and
background correction were performed with Nikon Elements software. ΔF/F was calculated
with the formula $(F_t - F_{min})/F_{min}$, where F_t is the raw fluorescence at time t and F_{min} is the
minimum fluorescence for the entire trace. The resulting trace was then denoised with a low-

pass Butterworth filter and baseline corrected for drift using an adaptative iteratively
reweighted Penalized Least Squares (AirPLS) based algorithm [49], in R-studio (R version 4.0.3).
Peak detection was performed using a custom script in R that considered neuron action
potential-derived Ca^{2+} spikes that had the following criteria (for a framerate acquisition of 4.7
fps): duration < 30 frames, rise phase >= 2 frames, fall phase >= 5 frames, rise phase < fall
phase, peak height > 5*SD, and peak height > 5*max background signal. The analysis pipeline is
depicted in Supp. Fig. 6. Statistical analysis was performed in GraphPad Prism 8.4.3 or R version
4.0.3.

Statistics

Parameters that were sampled in cells from multiple microscopic fields and from multiple cell
lines were fit to a linear mixed-effects model using the lme4 package in R[50]. The model included
the cell line identifier and sex. A Tukey post-hoc test identified pairwise differences in two-
factor models (e.g., combined genotype and ethanol exposure). Electrophysiology results were
modeled using generalized estimating equations (GEE) with the cell line as grouping identifier
using the geepack R package[51]. Where appropriate, ANOVA or Student's t-test was used as
indicated in figure legends. For RNAseq, pseudo-bulk data (single cell reads pooled by subject
identifier) were modeled in DESeq2[43], testing first by likelihood ratio testing (LRT) over all
groups, and then pairwise comparisons were evaluated using Wald tests. After Benjamini-
Hochberg multiple measurements correction, a false discovery rate of 5% was set as a
threshold for significance. Numbers of replicate cells and/or fields per cell line and genotype
used for statistical testing are listed in Supplemental Table 6.

Results

Gene expression patterns predict functional differences between iN generated from AUD
affected and unaffected individuals

To investigate the role of *KCNJ6* gene variants in neuronal function and ethanol response we
selected a balanced set of eight subjects of European ancestry from the COGA cell repository,
with contrasting *KCNJ6* SNPs and the presence or absence of alcohol dependence (Table 1).
Since multiple SNP genotypes were used, we label groups as AUD "affected" (**AF**) or
"unaffected" (**UF**) for simplicity, since in this set of subjects diagnosis correlates with the SNP
haplotype. However, the focus of this study is the *KCNJ6* haplotype.

To confirm and extend SNP genotypes in the two groups, iN cultures from each cell line were
harvested for bulk RNAseq analysis. To identify RNA isoforms, genome-aligned RNAseq data
identified only the longer *KCNJ6* transcript isoform ENST00000609713 (Fig. 1B), which includes
a 3' exon of 18,112 nucleotides that primarily consists of noncoding 3'UTR. Genotypes for the
two exonic SNPs (Table 1) were confirmed for all eight cell lines, but we also identified an
additional 19 noncoding, 3'UTR SNPs linked with the initial three (Supplementary Table 1;
mapped with red lines in Fig. 1B), constituting a haplotype of 22 variants (synonymous and
noncoding) within the expressed isoform of *KCNJ6* mRNA, all in a region of linkage
disequilibrium (LD; Supplemental Figure 1C). There are no nonsynonymous variants in either
haplotype, therefore, any function would be predicted to affect *KCNJ6* mRNA stability,
translation, or other post-transcriptional processes.

Group	ID	Sex	DSMIV Dx	rs702859 Synonymous	rs702860 Intronic	rs2835872 Intronic
AF	376	M	Affected	GG	GG	AA
	246	M	Affected	GG	GG	AA
	351	F	Affected	GG	GG	AA
	233	F	Affected	GG	GG	AA
UN	472	M	Unaffected	AA	AA	GG
	384	M	Unaffected	AA	AA	GG
	451	F	Unaffected	AA	AA	GG
	420	F	Unaffected	AA	AA	GG

Table 1. Subjects selected for iPSC. The ID is a three-digit code that is not traceable to the subject identity.
Genotypes for listed SNPs were obtained from GWAS studies[8]. SNP genotypes were confirmed by iPSC genomic
DNA PCR and Sanger sequencing.

To enable selective analysis of the more appropriately induced neurons, we used single-cell
RNAseq (scRNAseq) of pooled neurons from multiple cell lines, a strategy known as a "cell
village[52]." Based on preliminary results indicating differences in synaptic activity, we cultured
each group separately, to avoid potential secondary effects due to differential synaptic activity,
combining cells from all 4 lines within each group in equivalent proportions. Previous functional
studies suggested that iN cultures may be heterogeneous, potentially confounding a bulk

RNAseq analysis[31]. Following maturation (~30 days after induction), dissociated cells were
processed to generate scRNAseq libraries. Mapping cells by t-distributed stochastic neighbor
embedding (tSNE), a distinct cluster of cells coordinately expressed several markers consistent
with neuron physiological function (Fig. 1C and Supplemental Fig. 2E), including synaptic marker
(*SYP*), voltage-gated sodium channel (*SCN3A*), glutamate transporter (*SLC17A6*), and both
NMDA (*GRIN2A, GRIN2B, GRIA2, GRIA4*) and kainate glutamate receptors (*GRIK2*). This cluster
also expressed both *KCNJ6*, encoding GIRK2, and *KCNJ3*, encoding GIRK1, which are required to
form heterotetrameric, functional channels[14, 53]. Detection of mRNA expression in individual
cells by scRNAseq is likely an underestimate due to the relatively small number of reads per
cells, so the cluster identified as neurons is likely to express markers more uniformly than
observed here. We conclude that this subset of cells represents those expressing components
required for neuron physiological activity, and that other, sporadic cells, expressing subsets of
these neuronal markers, are likely alternate products of induction, which we label as
"transitional" neurons (Fig. 1C), since reprogramming is not identical to differentiation[54, 55].
Therefore, we focused gene expression analysis on the phenotypically neuronal subset of cells.

Cells from individual subjects were distinguished by expressed SNP profiles[42], and sequencing
reads of cells from each subject were combined to create a "pseudo-bulk" analysis, treating
each subject and treatment condition as a single replicate. Comparing the untreated AF group
with the untreated UN group, we identified 797 genes up-regulated, and 596 genes down-
regulated (Fig. 1D; Supplemental Table 3). Examining the results for differences by sex instead
of *KCNJ6* haplotype identified only 6 genes, of which 4 are encoded on X or Y chromosomes,
indicating that the sex of the samples did not substantially contribute to gene expression
differences. Gene ontology analysis of the down-regulated genes (Fig. 1E; Supplemental Table
4-5) predicts several biological processes associated with nervous system development, axonal
transport, trans-synaptic signaling and others. By grouping enriched gene ontology terms into a
tree plot by their parent terms (Supplemental Fig. 3), the major themes in the down-regulated
genes (AF < UN; Supplemental Table 5) are synaptic signaling and neuro projection
morphogenesis. Up-regulated genes (AF > UN; Fig. 1E; Supplemental Table 4) predict functions
associated with protein targeting within the cells, catabolic metabolism, and nonsense-
mediated decay. The level of *KCNJ6* mRNA trended lower in untreated AF than in UN (Fig. 1C; p
= 3.15 x 10^{-5} by LRT over both genotype and ethanol treatment; p = 0.0508 for untreated AF vs.
UN by Wald test). Interestingly, 7 d of 20 mM IEE increased levels of *KCNJ6* mRNA (Fig. 1C; p =
0.0225, Wald test) so that the treated AF group was no longer different from untreated UN (p =
0.322; Wald test). Analysis by sex identified no significant difference in *KCNJ6* mRNA. Results
indicate that the variant *KCNJ6* haplotype leads to differential expression of GIRK2 and that
ethanol exposure will reverse these effects.

GIRK2 expression and function in iPSC-derived induced excitatory human neurons (iN)
Given the differences in gene expression and partial reversal of KCNJ6 downregulation, we
evaluated detailed neuronal phenotypes in UN and AF iN cultures, including GIRK2 expression,
neuronal morphology, and physiological response. Since GIRK2 immunocytochemistry had not

been reported in cultured human neurons, we validated detection using mouse primary cortical
cultures[56]. GIRK2 immunoreactivity in human iNs matched a pattern of process-selective
expression in mouse (Fig. 2A, Supplemental Fig. 4A), where GIRK2 was detected as relatively
small (~ 0.5 μm diameter) puncta scattered primarily along the processes (Fig. 2B). Importantly,
silencing KCNJ6 with shRNAs or frameshift knockout by CRISPR/Cas9 eliminated GIRK2
immunoreactivity (Supplemental Fig. 4B,C). Localization of GIRK2 immunoreactivity in human iN
did not directly colocalize with synaptic vesicle marker VGLUT2 or synaptic marker Syn1
(Supplemental Fig. 5), but instead was found most frequently adjacent to synapses but
overlapping the shafts of the βIII tubulin-positive processes, and less so on MAP2 positive
processes (Figs. 2D.a, 3C, 3E). Cultured neurons express βIII tubulin throughout the cell, but not
as strongly in axonal processes[57]. We previously found that processes in human iN cells stained
for ankyrin G, identifying the axonal initial segment, which similarly lacked βIII-tubulin[58].
Detection of GIRK2 primarily on βIII-tubulin$^+$/MAP$^-$ processes, therefore, suggests pre-axonal,
and likely presynaptic, localization. Results confirm that lentiviral-transduced GIRK2
overexpression increased levels of immunostaining about 3.5-fold (Fig. 2C, mCherry$^+$ cells are
KCNJ6 lentiviral-transduced) which further confirms antibody specificity.

To confirm that GIRK2 expression contributes to potassium channel function, we treated iN
cultures with ML297, a selective activator of the GIRK1/GIRK2 heterotetramer complex[59, 60].
Results show that the magnitude and frequency of native (basal) currents observed in human iN
were relatively small (Fig. 2D.b), with only 6.8% of the neurons responding to a shift in
membrane potential holding current (~10 pA). This agrees well with immunocytochemical data
showing that GIRK2 expression is relatively low and with scRNAseq detection of KCNJ6 in a
subset of cells (Fig. 1C). However, by recording from cells overexpressing GIRK2 (identified by
mCherry co-expression; Fig. 2D.c), we found an increased frequency of response (GIRK currents)
to 30% of the cells after ML297 addition (Fig. 2D.b) without a change in magnitude (Fig. 2D.e).
Importantly, we also found that GIRK activation affected excitability of the neurons by shifting
resting membrane potential to more negative values (Fig. 2D.d), thus affecting the ability of
neurons to fire APs when induced (Fig. 2D.f). We conclude that increased KCNJ6 expression
affects neuronal excitability by altering GIRK channel activity.

KCNJ6 haplotype alters morphology and membrane excitability in human iNs
To identify neuronal properties affected by KCNJ6 haplotype, we focused on three aspects:
morphology, expression of GIRK2, and basal physiological properties. A detailed analysis of
neuronal morphology is essential to rule out differences affecting excitability[61]. Most measures
of basic neuronal morphology did not exhibit statistically significant differences by group,
including soma size, circularity, and solidity, which describe the most fundamental aspects of
neuron shape (Fig. 3B.a-c). However, neurite area, as determined by βIII-tubulin-stained area,
reflecting the number and/or branching of neurites per cell, increased in the AF group (p
=0.018; Fig. 3B.d), with example images shown for individual cell lines in Figure 3C. Results are
plotted for each cell line or aggregated by KCNJ6 haplotype from multiple cell lines. As

predicted by the scRNAseq expression of *KCNJ6* mRNA (Fig. 1J), GIRK2 immunoreactivity was
decreased in the AF group (Fig. 3D), as measured by puncta counts (p = 0.0012; Fig. 3D.a),
puncta circularity (p =0.007; Fig. 3D.c), and solidity (p = 0.037; Fig. 3D.d) but not puncta size
(Fig. 3D.b). No difference was found by sex. Sample GIRK2 puncta are shown for each cell line in
Figure 3E. These results demonstrate an overall decrease in GIRK2 in neuronal processes in the
AF (*KCNJ6* variant allele) group.

Reduced GIRK2 expression is predicted to affect the excitability of neurons. As expected, we
found no differences in membrane capacitance, membrane resistance, spontaneous excitatory
post-synaptic potential (sEPSC) frequencies or amplitudes, or spontaneous action potential (AP)
firing (Fig. 3F). However, we observed a significant increase in excitability in the AF group (p =
1.2×10^{-5}, Fig. 3G). Less current injection, using a ramp technique, was required for the AF
group to shift the resting membrane potential (RMP) of neurons held at -65 mV, producing
more AP firing, compared with the UN group (p = 2.0×10^{-9}, Fig. 3G.d). Therefore, the *KCNJ6*
variant haplotype group (AF) exhibited increased neurite area, reduced neurite GIRK2
expression, and greater excitability. Other neuronal properties were not different across the
two groups, suggesting that overall neuron differentiation status was similar.

Electrophysiology was assessed in multiple individual cells, but since cultures are
heterogeneous (Fig. 1C), we wished to evaluate larger samples of cultures. Therefore, to
confirm that the observed effects are consistent across larger populations of cells, we used
calcium imaging to assess spontaneous and glutamate-stimulated firing. We selected four cell
lines for these studies (AF: 233 and 246, and UN: 420 and 472) and used the viral-transduced,
genetically-encoded calcium indicator GCaMP6f as a proxy for neuronal activity[62]. Example
images show consistent expression in all cell lines (Fig. 4A). Cultures were treated with
repeated pulses of 10 μM glutamate (each followed by wash-out with ACSF), and one pulse of
50 μM glutamate to elicit transient increases in fluorescence, indicating receptor-induced
excitability, followed by pulses of 18 mM KCl, indicating cellular excitability (Fig. 4B-D). Neurons
from both UN and AF groups had heterogeneous spiking patterns, with >50% of the neurons
remaining inactive during the baseline period or after stimulation with glutamate or KCl. This is
consistent with scRNAseq results indicating that only ~23% of the iN cells express genetic
markers consistent with excitability (Fig. 1C). Increased spontaneous and induced spiking (> 1-2
spikes) was apparent in the AF group when viewing individual cell responses in a raster plot
(Fig. 4B-C). Spiking frequencies increased 2.4-fold in baseline conditions (Fig. 4E; p = 1.1×10^{-6})
and 1.3-fold following glutamate stimulation (Fig. 4F; p = 1.6×10^{-3}) in the AF group, but no
difference was found in KCl-induced activity (Fig. 4G; p = 0.85), confirming the above results
from individual neurons. When examining the AF or UN neurons individually, we also observed
higher spontaneous and glutamate elicited excitability for lines 233 and 246 when compared to
line 472, which was found to be the least active spontaneously or under glutamate stimulation
among the four studied with this imaging approach (Supp. Fig. 6). Overall, cells from the AF
KCNJ6 variant allele group were more excitable, with no difference in basal physiological

properties (i.e., KCl-induced activity). Furthermore, not only did the spike rate increase per cell
(bar plots, Spikes/ROI/min), but we also observed increases in the proportion of cells in a field
that responded (pie plots). These results not only confirm earlier findings of differences in
excitability, but also demonstrate that the difference in the AF iNs is found in not only individual
cell activity but also in the frequency of detecting active cells within a population.

Ethanol exposure eliminates differences in neuronal properties affected by *KCNJ6* haplotype
Analysis of *KCNJ6* mRNA expression in selected neurons (Fig. 1C) not only detected decreased
basal levels in the AF group, but also demonstrated increased expression following ethanol
treatment (~2-fold increased, p = 0.0225 Wald test). This result predicts that increasing GIRK2
expression in the AF group following ethanol treatment would eliminate differences from the
UN group in morphology, expression, and physiology.

To control ethanol dosage in culture, we utilized an intermittent ethanol exposure (IEE)
paradigm as in previous studies[25]. We replenished ethanol daily to account for loss by
evaporation, producing measured concentrations with a mean of 15.4 ± 1.16 mM
(Supplemental Fig. 7). We determined that the half-life of ethanol is 14.5 h (with a 95% CI of
13.5 to 15.6 h), so over 24 h the concentration would drop to ~5-8 mM before replenishment.
We targeted a peak concentration of 20 mM ethanol, which is similar to a blood alcohol
concentration (BAC) of 0.08% (17 mM), the legal limit for intoxication. We also selected this
concentration based on the reported concentration-dependent interaction of GIRK2 with
ethanol[17]. After 7 days of IEE, we evaluated neuronal morphological and functional properties.
Focusing on those parameters that were different by genotype without IEE (Fig. 3&4), we found
that ethanol eliminated the differences in total neurite area (not significant; p = 0.98, Fig. 5A.d),
GIRK2 puncta counts (p = 0.46, Fig. 5B.a), excitation following current injection (p = 0.32, Fig.
5E.a), step- (p = 0.48, Fig. 5E.b), and ramp- (p = 0.95, Fig. 5E.d) induced APs. Membrane
capacitance was slightly but significantly reduced in the IEE AF group (p = 9.6×10^{-9}, Fig. 5D.a).
Other measurements were unchanged, again indicating that the cultured neurons exhibited
similar differentiation properties. Results are consistent with the interpretation that neurite
and excitability differences correlate with reduced GIRK2 expression, and that ethanol increases
expression of GIRK2, potentially reversing these effects.

To confirm that both *KCNJ6* haplotype and ethanol exposure affect GIRK2 expression, we
evaluated GIRK2 immunocytochemistry and *KCNJ6* mRNA in neurons using fluorescent in situ
hybridization (FISH). GIRK2 increased following IEE (p=1.0×10^{-20}, Fig. 5G-H). By probing *KCNJ6*
fluorescent puncta colocalized with MAP2 immunocytochemistry, we confirmed that neurons
from AF individuals had lower levels of *KCNJ6* mRNA expression compared to UN neurons (Fig.
5J.a, p = 8.2×10^{-3}). We also found that the proportion of MAP2$^+$ neurons expressing KCNJ6 was
reduced in AF compared with UN (50.1% vs 64.7%, Fig. 5J.b), and the AF neurons had a greater
proportion of cells with low levels of expression (Fig. 5J.c) with no detectable High or Very High
(>9 puncta/cell) expression. Differences in expression were found by haplotype in somatic
regions of neurons (Fig. 5J.f, p = 5.5×10^{-3}) but not in non-somatic processes (Fig. 5J.g, p = 0.23).

Ethanol increased *KCNJ6* mRNA levels in both groups (Fig. 5J.a-e), eliminating the difference
between genotypes. Interestingly, we also observed that the proportions of AF neurons
expressing KCNJ6 was increased (8.3%, Fig. 5J.b) by ethanol and was paralleled by a reduction in
number of low-expressing cells and increased appearance of high-expressing neurons (Fig. 5J.c).
Results indicate that variant *KCNJ6* haplotype leads to reduced *KCNJ6* expression, which is
consistent with RNA sequencing and immunocytochemical results, and that ethanol increases
KCNJ6 expression, eliminating differences between haplotype groups.

GIRK2 overexpression mimics IEE in human neurons with *KCNJ6* variants
To test if ethanol-induced GIRK2 expression could underlie the elimination of differences in
neuronal properties associated with the *KCNJ6* haplotype, we compared ethanol treatment
with virus transduced *KCNJ6* overexpression. Since ethanol has been shown to potentiate
GIRK2-containing channels, we used a single treatment with 20 mM ethanol followed by
fixation 24 h later. As expected, neither ethanol nor GIRK2 overexpression exhibited differences
in passive neuronal properties. However, while current-induced activity increased in control
cultures ($p = 6.8 \times 10^{-6}$, Fig. 6C), no significant difference was found in cultures exposed to 24 h
ethanol ($p = 0.74$) and GIRK2 overexpression ($p = 0.23$). Similarly, while control cultures had
increased ramp-induced APs ($p = 5.9 \times 10^{-4}$; Fig. 6A) neither GIRK overexpression ($p = 0.14$) or
ethanol ($p = 0.10$) were significantly different. As a control, GIRK2 puncta count increased in a
sample cell line after overexpression (line 376, $p = 0.04$, Fig. 6D) without an increase in
circularity or solidity (not shown). These results indicate that the increased excitability in *KCNJ6*
minor allele haplotype cells is due to reduced expression of GIRK2 protein and this effect is at
least partially ameliorated or reversed by exposure to doses of ethanol found in human brain
after moderate to heavy daily drinking.

Discussion

This study focused on the effect of an AUD- and endophenotype-linked *KCNJ6* haplotype on
neuronal function, with a goal of identifying mechanisms that could eventually drive
therapeutic strategies. To retain the appropriate genetic background for individuals with AUD,
we selected subjects from the large NIAAA/COGA Sharing Repository, using both the selected
KCNJ6 haplotype and diagnosed alcohol dependence as criteria (Table 1). By preparing iPSC
from these subjects and inducing them to model excitatory neurons, we identified differences
in *KCNJ6* mRNA levels and GIRK2 immunoreactivity in individuals with variant *KCNJ6*
haplotypes. Gene expression differences predicted effects on neuronal signaling including
synaptic function. The *KCNJ6* haplotype was also associated with differences in the area of βIII-
tubulin^{+} neuron projections and membrane excitability, but not in other measures of neuronal
differentiation. Surprisingly, we found that ethanol exposure led to an increase in *KCNJ6* mRNA
and GIRK2 expression, paralleled by a reduction or elimination of morphological and
physiological differences in these neurons. Reversal of *KCNJ6* haplotype effects by ethanol
could also be mimicked by overexpressing GIRK2. These results demonstrate that genetic
variants enhancing risk of AUD, even if they do not alter protein sequence, can trigger neuronal
mechanisms at the cellular level mirroring or predicting endophenotypes, here ERO, linked with
AUD behavior.

Non-coding *KCNJ6* polymorphisms in AUD

Most GWAS SNPs map to non-coding regions of the genome and 3.7% are found in the UTRs[63,
64]. The 3′ UTRs of transcripts that are expressed in the brain tend to be longer than those
expressed in other tissues and can be involved in post-transcriptional regulation of transcript
abundance by affecting RNA stability, translation and/or localization[65, 66]. Reference genome
annotation of *KCNJ6* identifies two transcripts, one with a short 3′ UTR (1,926 bp; Ensembl
ENST00000645093, RefSeq NM_002240) and one with a substantially longer 3′ UTR (18,112 bp;
Ensembl ENST00000609713). RNAseq coverage analysis in human iN cultures detects only the
longer isoform, although it is possible that a relatively small portion could be the shorter
isoform, but the uniformly distributed coverage (Fig. 1B) does not indicate this. This longer
transcript is orthologous to a 16-kb transcript found in rat brain[67], which includes multiple AU-
rich elements, which may affect mRNA stability and could be affected by *KCNJ6* allelic variation.
We conclude that the vast majority of *KCNJ6* mRNA is consistent with having the extended 3′
UTR, with the potential for variant SNPs to affect mRNA stability or translation efficiency.

In addition to the three synonymous and intronic SNPs used to select subjects (Table 1),
examination of variants in the long 3′UTR by RNAseq alignment predicts linkage of 19 additional
SNPs (Fig. 1B; Supplemental Table 1). None of the 3′ UTR SNPs could be mapped uniquely to
known regulatory sequences such as predicted microRNA target sites. That is, some SNPs are
predicted to destroy or add predicted targeting sites, but all of these potential target sites were
found in multiple locations within the 3′ UTR, not only where they might be altered by a SNP. It
is possible, however, that altered microRNA targeting of one or more sites among many in the

3'UTR could play a role in regulating the stability or translation of the variant *KCNJ6* mRNA, but

no clear candidates could be identified for testing. Furthermore, since ethanol exposure of

neurons has been found to inhibit 1-carbon metabolic pathways, leading to selective changes in

histone methylation[68], it remains possible that a sequence-specific epigenetic mechanism other

than microRNAs may affect *KCNJ6* variant expression.

However, results demonstrate that the non-coding variant haplotype alters expression of GIRK2

protein and *KCNJ6* mRNA. We found that iN from subjects with the homozygous variant *KCNJ6*

haplotype exhibited lower expression of GIRK2 by immunocytochemistry (Fig. 3D), but while

the difference in neuronal mRNA levels as detected by scRNAseq analysis matched this trend, it

did not reach significance (p = 0.0508; Fig. 1C). FISH analysis, however, confirmed that *KCNJ6*

mRNA was reduced in the AF group (Fig. 5J). *KCNJ6* mRNA was also detected in fewer MAP2$^+$

cells and at lower levels per cell (Fig. 5J.b-c). The results of counting GIRK2 puncta and

evaluating *KCNJ6* mRNA by scRNAseq and FISH support a diminished expression of GIRK2 in

neurons from the affected group.

The secondary effects of *KCNJ6* mRNA and GIRK2 protein regulation include many predictors of

altered neuronal connectivity and signaling. Genes that were reduced in the AF group (GO:BP

Down in Fig. 1E; Supplemental Fig. 3B) point to changes in synaptic transmission and projection

morphology. The tree plots in Supp. Fig. 3B are particularly useful to group GO terms by their

parent terms and to judge the numbers of genes affected (by the diameters of the circles

plotted). Since we focused on *KCNJ6* haplotype differences in cells without ethanol, we did not

expect to match findings in, for example, human AUD brains[69], which found prominent effects

in astrocytes and microglia.

Surprisingly, ethanol exposure increased expression of both GIRK2 immunoreactivity and *KCNJ6*

mRNA. Following 7 days of IEE with a peak concentration of 20 mM ethanol, GIRK2 puncta in

the AF group increased to levels significantly higher than the UN group (Fig. 5B). Similarly,

KCNJ6 mRNA levels were shown to be increased above untreated levels by scRNAseq results

(Fig. 1C) as well as FISH analysis (Fig. 5J). Ethanol also increased the proportion of MAP2$^+$ cells

expressing *KCNJ6* as well as the expression level per cell (Fig. 5J.b-c). While previous studies

identified a binding site for ethanol in a defined pocket of GIRK2[70], suggesting that ethanol

could affect protein stabilization, this binding could not explain the increased mRNA level.

There is no indication that ethanol or *KCNJ6* haplotype altered translocation of *KCNJ6* mRNA to

neuronal processes (Fig. 5J.d-e). Ethanol has been found to affect microRNA expression

patterns that regulate mRNA[65, 71-76]. It is intriguing to speculate that ethanol might increase

expression of GIRK2 through this type of mechanism, with polymorphisms in the 3'UTR serving

as the modulator.

GIRK2 in the context of alcohol dependence

GIRK channels play an essential role in maintaining the excitability of neurons. For example, in

Trisomy 21 individuals, whose cells contain an additional copy of *KCNJ6*, neuronal activity was

shown to be reduced due to excessive channel activity[77-79]. On the other hand, loss of GIRK2

activity in knockout mice increases susceptibility to induced seizures[20, 21]. We detected the
presence of GIRK channel function in glutamatergic iNs using the ML297 agonist of GIRK1/2
heterotetramers, showing that activation led to a reduction in the frequency of action
potentials induced by depolarization (Fig. 2D). However, only 6.7% of neurons responded to
ML297, but overexpression of GIRK2 increased the proportion of responding neurons to 30%
(Fig. 2D). We therefore chose to focus on evaluating neuronal excitability as an indirect
response to changes in GIRK function.

By identifying the correlation between GIRK2 expression levels and altered GIRK2-mediated
function, we predict that neuronal excitability will be affected by *KCNJ6* variants and alcohol
exposure. Human iNs generated from subjects with the variant *KCNJ6* haplotype and AUD were
more excitable and had lower levels of GIRK2 expression. Moreover, ethanol exposure, both
shorter (1 d) and longer (7 d) term, diminished these differences, mimicking the effect of GIRK2
overexpression. While GIRK channels play relatively small role in maintaining neuronal
homeostasis, modulating GIRK activity to alter cell excitability is predicted to play a critical role.
Additionally, downstream effectors of the G protein-mediated signaling pathways presumably
regulate neuronal function indirectly[14, 15, 17, 19]. GIRK channels are implicated in several other
disorders with abnormal neuronal excitability, including epilepsy, suggesting that they have
therapeutic potential[19-21, 77-80].

Potential limitations

Several factors beyond *KCNJ6* haplotype may contribute to these interpretations. First, we
selected sex-balanced groups because it is possible that the sex of individual neurons might
contribute to the results or at least affect variability. However, in nearly all parameters tested,
there was no significant difference by sex. The scRNAseq results identified only 6 genes as
different by sex and 4 of these are encoded by sex chromosomes. None of the basal
morphology measurements (Fig. 3B, D) differed by sex, but membrane capacitance (Fig. 3F.a, p
$= 2.6 \times 10^{-4}$), induced APs (step protocol, Fig. 3G.b, p = 0.012), and ethanol-treated membrane
capacitance (Fig. 5D.a, p = 3.1×10^{-8}) all reached significance, but it is difficult to interpret how
sex interacts with neuronal function in the context of artificially cultured neurons. Second, the
choices we made about pooling subjects for "cell village" experiments (Fig. 1C-E) might have
affected results. We chose to mix subjects with the same *KCNJ6* haplotype, fearing a non-cell-
autonomous effect based on altered excitability. Finally, we chose to focus on homozygous
haplotypes and extreme AUD diagnoses in this study to enhance contrasts, however, additional
studies might examine if heterozygotes would exhibit gene dosage effects or be rescued by the
presence of the major alleles, similar to endophenotype studies with the original subjects[8].

From genes to behavior

In this study, we found that the variant *KCNJ6* haplotype affects excitability of neurons (Fig. 3 &
4E). In Ca^{2+} imaging experiments, concomitant with enhanced excitability in neurons from
affected individuals following stimulation with glutamate pulses, we observed an overall
increase in basal activity of the neuronal population (Fig. 4F). This is reminiscent of the original

observation linking *KCNJ6* variants with an EEG endophenotype, where Kamarajan and
colleagues observed that individuals with alcohol dependence and the *KCNJ6* haplotype had an
ERO theta power that varied as a function of the *KCNJ6* haplotype in both loss and gain
conditions[10]. There is an extensive human literature linking impulsivity to alcohol use
problems[81-84]. Impulsivity is elevated in offspring who are at high risk for substance use
disorders and may be a reflection of a genetic vulnerability for substance use problems[85].
Excitability in individual neurons and particularly in a culture dish of neurons is several levels of
complexity removed from EEG patterns in brain. However, our study points to a mechanistic
underpinning of how heritable risk traits are likely to play a role in development of unique
physiological response to alcohol in the brain.

Acknowledgements

The Collaborative Study on the Genetics of Alcoholism (COGA), Principal Investigators B. Porjesz, V. Hesselbrock, T. Foroud; Scientific Director, A. Agrawal; Translational Director, D. Dick, includes eleven different centers: University of Connecticut (V. Hesselbrock); Indiana University (H.J. Edenberg, T. Foroud, Y. Liu, M. Plawecki); University of Iowa Carver College of Medicine (S. Kuperman, J. Kramer); SUNY Downstate Health Sciences University (B. Porjesz, J. Meyers, C. Kamarajan, A. Pandey); Washington University in St. Louis (L. Bierut, J. Rice, K. Bucholz, A. Agrawal); University of California at San Diego (M. Schuckit); Rutgers University (J. Tischfield, R. Hart, J. Salvatore); The Children's Hospital of Philadelphia, University of Pennsylvania (L. Almasy); Virginia Commonwealth University (D. Dick); Icahn School of Medicine at Mount Sinai (A. Goate, P. Slesinger); and Howard University (D. Scott). Other COGA collaborators include: L. Bauer (University of Connecticut); J. Nurnberger Jr., L. Wetherill, X., Xuei, D. Lai, S. O'Connor, (Indiana University); G. Chan (University of Iowa; University of Connecticut); D.B. Chorlian, J. Zhang, P. Barr, S. Kinreich, G. Pandey (SUNY Downstate); N. Mullins (Icahn School of Medicine at Mount Sinai); A. Anokhin, S. Hartz, E. Johnson, V. McCutcheon, S. Saccone (Washington University); J. Moore, Z. Pang, S. Kuo (Rutgers University); A. Merikangas (The Children's Hospital of Philadelphia and University of Pennsylvania); F. Aliev (Virginia Commonwealth University); H. Chin and A. Parsian are the NIAAA Staff Collaborators. We continue to be inspired by our memories of Henri Begleiter and Theodore Reich, founding PI and Co-PI of COGA, and also owe a debt of gratitude to other past organizers of COGA, including Ting- Kai Li, P. Michael Conneally, Raymond Crowe, and Wendy Reich, for their critical contributions. This national collaborative study is supported by NIH Grant U10AA008401 from the National Institute on Alcohol Abuse and Alcoholism (NIAAA) and the National Institute on Drug abuse (NIDA).

746 ## Conflict of Interest

747 The authors declare no conflict of interest.

References

1. Verhulst B, Neale MC, Kendler KS. The heritability of alcohol use disorders: a meta-analysis of twin and adoption studies. *Psychol Med* 2015; **45**(5)**:** 1061-1072.

2. Pollard MS, Tucker JS, Green HD, Jr. Changes in Adult Alcohol Use and Consequences During the COVID-19 Pandemic in the US. *JAMA Network Open* 2020; **3**(9)**:** e2022942-e2022942.

3. Bouza C, Angeles M, Muñoz A, Amate JM. Efficacy and safety of naltrexone and acamprosate in the treatment of alcohol dependence: a systematic review. *Addiction* 2004; **99**(7)**:** 811-828.

4. Heilig M, Egli M. Pharmacological treatment of alcohol dependence: Target symptoms and target mechanisms. *Pharmacology & Therapeutics* 2006; **111**(3)**:** 855-876.

5. Skinner MD, Lahmek P, Pham H, Aubin HJ. Disulfiram efficacy in the treatment of alcohol dependence: a meta-analysis. *PLoS One* 2014; **9**(2)**:** e87366.

6. Salvatore JE, Gottesman, II, Dick DM. Endophenotypes for Alcohol Use Disorder: An Update on the Field. *Curr Addict Rep* 2015; **2**(1)**:** 76-90.

7. Chorlian DB, Rangaswamy M, Manz N, Meyers JL, Kang SJ, Kamarajan C *et al.* Genetic correlates of the development of theta event related oscillations in adolescents and young adults. *Int J Psychophysiol* 2017; **115:** 24-39.

8. Kang SJ, Rangaswamy M, Manz N, Wang JC, Wetherill L, Hinrichs T *et al.* Family-based genome-wide association study of frontal θ oscillations identifies potassium channel gene KCNJ6. *Genes Brain Behav* 2012; **11**(6)**:** 712-719.

9. Clarke TK, Laucht M, Ridinger M, Wodarz N, Rietschel M, Maier W *et al.* KCNJ6 is associated with adult alcohol dependence and involved in gene × early life stress interactions in adolescent alcohol drinking. *Neuropsychopharmacology* 2011; **36**(6)**:** 1142-1148.

10. Kamarajan C, Pandey AK, Chorlian DB, Manz N, Stimus AT, Edenberg HJ *et al.* A KCNJ6 gene polymorphism modulates theta oscillations during reward processing. *Int J Psychophysiol* 2017; **115:** 13-23.

11. Andrew C, Fein G. Event-related oscillations versus event-related potentials in a P300 task as biomarkers for alcoholism. *Alcohol Clin Exp Res* 2010; **34**(4)**:** 669-680.

790 12. Kamarajan C, Rangaswamy M, Tang Y, Chorlian DB, Pandey AK, Roopesh BN *et al.*
791   Dysfunctional reward processing in male alcoholics: an ERP study during a gambling
792   task. *J Psychiatr Res* 2010; **44**(9)**:** 576-590.

793
794 13. Glaaser IW, Slesinger PA. Structural Insights into GIRK Channel Function. *International*
795   *review of neurobiology* 2015; **123:** 117-160.

796
797 14. Luscher C, Slesinger PA. Emerging roles for G protein-gated inwardly rectifying
798   potassium (GIRK) channels in health and disease. *Nat Rev Neurosci* 2010; **11**(5)**:** 301-
799   315.

800
801 15. Zhao Y, Gameiro-Ros I, Glaaser IW, Slesinger PA. Advances in Targeting GIRK
802   Channels in Disease. *Trends Pharmacol Sci* 2021; **42**(3)**:** 203-215.

803
804 16. Reuveny E, Slesinger PA, Inglese J, Morales JM, Iñiguez-Lluhi JA, Lefkowitz RJ *et al.*
805   Activation of the cloned muscarinic potassium channel by G protein beta gamma
806   subunits. *Nature* 1994; **370**(6485)**:** 143-146.

807
808 17. Aryal P, Dvir H, Choe S, Slesinger PA. A discrete alcohol pocket involved in GIRK
809   channel activation. *Nat Neurosci* 2009; **12**(8)**:** 988-995.

810
811 18. Bodhinathan K, Slesinger PA. Alcohol modulation of G-protein-gated inwardly rectifying
812   potassium channels: from binding to therapeutics. *Front Physiol* 2014; **5:** 76.

813
814 19. Blednov YA, Stoffel M, Chang SR, Harris RA. Potassium channels as targets for ethanol:
815   studies of G-protein-coupled inwardly rectifying potassium channel 2 (GIRK2) null
816   mutant mice. *J Pharmacol Exp Ther* 2001; **298**(2)**:** 521-530.

817
818 20. Blednov YA, Stoffel M, Chang SR, Harris RA. GIRK2 deficient mice. Evidence for
819   hyperactivity and reduced anxiety. *Physiol Behav* 2001; **74**(1-2)**:** 109-117.

820
821 21. Hill KG, Alva H, Blednov YA, Cunningham CL. Reduced ethanol-induced conditioned
822   taste aversion and conditioned place preference in GIRK2 null mutant mice.
823   *Psychopharmacology (Berl)* 2003; **169**(1)**:** 108-114.

824
825 22. Lieberman R, Levine ES, Kranzler HR, Abreu C, Covault J. Pilot study of iPS-derived
826   neural cells to examine biologic effects of alcohol on human neurons in vitro. *Alcohol*
827   *Clin Exp Res* 2012; **36**(10)**:** 1678-1687.

828
829 23. Lieberman R, Kranzler HR, Levine ES, Covault J. Examining the effects of alcohol on
830   GABA(A) receptor mRNA expression and function in neural cultures generated from
831   control and alcohol dependent donor induced pluripotent stem cells. *Alcohol* 2018; **66:**
832   45-53.

834   24.   Halikere A, Popova D, Scarnati MS, Hamod A, Swerdel MR, Moore JC *et al.* Addiction
835   associated N40D mu-opioid receptor variant modulates synaptic function in human
836   neurons. *Mol Psychiatry* 2020; **25**(7)**:** 1406-1419.

838   25.   Scarnati MS, Boreland AJ, Joel M, Hart RP, Pang ZP. Differential sensitivity of human
839   neurons carrying mu opioid receptor (MOR) N40D variants in response to ethanol.
840   *Alcohol* 2020; **87:** 97-109.

842   26.   Patzke C, Dai J, Brockmann MM, Sun Z, Fenske P, Rosenmund C *et al.* Cannabinoid
843   receptor activation acutely increases synaptic vesicle numbers by activating synapsins in
844   human synapses. *Mol Psychiatry* 2021.

846   27.   Weissbein U, Schachter M, Egli D, Benvenisty N. Analysis of chromosomal aberrations
847   and recombination by allelic bias in RNA-Seq. *Nature communications* 2016; **7:** 12144.

849   28.   Vierbuchen T, Ostermeier A, Pang ZP, Kokubu Y, Sudhof TC, Wernig M. Direct
850   conversion of fibroblasts to functional neurons by defined factors. *Nature* 2010;
851   **463**(7284)**:** 1035-1041.

853   29.   Pang ZP, Yang N, Vierbuchen T, Ostermeier A, Fuentes DR, Yang TQ *et al.* Induction of
854   human neuronal cells by defined transcription factors. *Nature* 2011; **476**(7359)**:** 220-223.

856   30.   Zhang Y, Pak C, Han Y, Ahlenius H, Zhang Z, Chanda S *et al.* Rapid single-step
857   induction of functional neurons from human pluripotent stem cells. *Neuron* 2013; **78**(5)**:**
858   785-798.

860   31.   Lin HC, He Z, Ebert S, Schörnig M, Santel M, Nikolova MT *et al.* NGN2 induces diverse
861   neuron types from human pluripotency. *Stem Cell Reports* 2021; **16**(9)**:** 2118-2127.

863   32.   Bardy C, van den Hurk M, Eames T, Marchand C, Hernandez RV, Kellogg M *et al.*
864   Neuronal medium that supports basic synaptic functions and activity of human neurons
865   in vitro. *Proc Natl Acad Sci U S A* 2015; **112**(20)**:** E2725-2734.

867   33.   Yang N, Chanda S, Marro S, Ng Y-H, Janas JA, Haag D *et al.* Generation of pure
868   GABAergic neurons by transcription factor programming. *Nature Methods* 2017; **14**(6)**:**
869   621-628.

871   34.   Kreitzer FR, Salomonis N, Sheehan A, Huang M, Park JS, Spindler MJ *et al.* A robust
872   method to derive functional neural crest cells from human pluripotent stem cells. *Am J*
873   *Stem Cells* 2013; **2**(2)**:** 119-131.

35. Miyaoka Y, Chan AH, Judge LM, Yoo J, Huang M, Nguyen TD *et al.* Isolation of single-base genome-edited human iPS cells without antibiotic selection. *Nature methods* 2014; **11**(3)**:** 291-293.

36. Ran FA, Hsu PD, Wright J, Agarwala V, Scott DA, Zhang F. Genome engineering using the CRISPR-Cas9 system. *Nature Protocols* 2013; **8**(11)**:** 2281-2308.

37. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics (Oxford, England)* 2018; **34**(17)**:** i884-i890.

38. Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology* 2019; **37**(8)**:** 907-915.

39. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO *et al.* Twelve years of SAMtools and BCFtools. *GigaScience* 2021; **10**(2).

40. McLaren W, Gil L, Hunt SE, Riat HS, Ritchie GRS, Thormann A *et al.* The Ensembl Variant Effect Predictor. *Genome Biology* 2016; **17**(1)**:** 122.

41. Butler A, Hoffman P, Smibert P, Papalexi E, Satija R. Integrating single-cell transcriptomic data across different conditions, technologies, and species. *Nat Biotechnol* 2018; **36**(5)**:** 411-420.

42. Kang HM, Subramaniam M, Targ S, Nguyen M, Maliskova L, McCarthy E *et al.* Multiplexed droplet single-cell RNA-sequencing using natural genetic variation. *Nat Biotechnol* 2018; **36**(1)**:** 89-94.

43. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* 2014; **15**(12)**:** 550.

44. Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. *Innovation (N Y)* 2021; **2**(3)**:** 100141.

45. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. *Nature methods* 2012; **9**(7)**:** 671-675.

46. Ho S-Y, Chao C-Y, Huang H-L, Chiu T-W, Charoenkwan P, Hwang E. NeurphologyJ: An automatic neuronal morphology quantification method and its application in pharmacological discovery. *BMC Bioinformatics* 2011; **12**(1)**:** 230.

916 47. Arganda-Carreras I, Kaynig V, Rueden C, Eliceiri KW, Schindelin J, Cardona A *et al.*
917 Trainable Weka Segmentation: a machine learning tool for microscopy pixel
918 classification. *Bioinformatics* 2017; **33**(15)**:** 2424-2426.

919
920 48. Fantuzzo JA, Robles DA, Mirabella VR, Hart RP, Pang ZP, Zahn JD. Development of a
921 high-throughput arrayed neural circuitry platform using human induced neurons for drug
922 screening applications. *Lab Chip* 2020; **20**(6)**:** 1140-1152.

923
924 49. Zhang ZM, Chen S, Liang YZ. Baseline correction using adaptive iteratively reweighted
925 penalized least squares. *Analyst* 2010; **135**(5)**:** 1138-1146.

926
927 50. Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using
928 lme4. *Journal of Statistical Software* 2015; **67**(1)**:** 1 - 48.

929
930 51. Højsgaard S, Halekoh U, Yan J. The R Package geepack for Generalized Estimating
931 Equations. *Journal of Statistical Software* 2005; **15**(2)**:** 1 - 11.

932
933 52. Mitchell JM, Nemesh J, Ghosh S, Handsaker RE, Mello CJ, Meyer D *et al.* Mapping
934 genetic effects on cellular phenotypes with "cell villages". *bioRxiv* 2020**:**
935 2020.2006.2029.174383.

936
937 53. Lesage F, Guillemare E, Fink M, Duprat F, Heurteaux C, Fosset M *et al.* Molecular
938 properties of neuronal G-protein-activated inwardly rectifying K+ channels. *J Biol Chem*
939 1995; **270**(48)**:** 28660-28667.

940
941 54. Ang CE, Wernig M. Induced neuronal reprogramming. *J Comp Neurol* 2014; **522**(12)**:**
942 2877-2886.

943
944 55. Shelby H, Shelby T, Wernig M. Somatic Lineage Reprogramming. *Cold Spring Harb*
945 *Perspect Biol* 2021.

946
947 56. Marron Fernandez de Velasco E, Zhang L, B NV, Tipps M, Farris S, Xia Z *et al.* GIRK2
948 splice variants and neuronal G protein-gated K(+) channels: implications for channel
949 function and behavior. *Sci Rep* 2017; **7**(1)**:** 1639.

950
951 57. Cáceres A, Banker GA, Binder L. Immunocytochemical localization of tubulin and
952 microtubule-associated protein 2 during the development of hippocampal neurons in
953 culture. *J Neurosci* 1986; **6**(3)**:** 714-722.

954
955 58. Ma F, Xu J, Liu Y, Popova D, Youssef MM, Hart RP *et al.* The amyloid precursor protein
956 modulates the position and length of the axon initial segment offering a new perspective
957 on Alzheimer's disease genetics. *bioRxiv* 2022**:** 2022.2001.2023.477413.

59. Wydeven N, Marron Fernandez de Velasco E, Du Y, Benneyworth MA, Hearing MC, Fischer RA *et al.* Mechanisms underlying the activation of G-protein-gated inwardly rectifying K+ (GIRK) channels by the novel anxiolytic drug, ML297. *Proc Natl Acad Sci U S A* 2014; **111**(29)**:** 10755-10760.

60. Days E, Kaufmann K, Romaine I, Niswender C, Lewis M, Utley T *et al.* Discovery and Characterization of a Selective Activator of the G-Protein Activated Inward-Rectifying Potassium (GIRK) Channel. *Probe Reports from the NIH Molecular Libraries Program.* National Center for Biotechnology Information (US): Bethesda (MD), 2010.

61. Kang S, Chen X, Gong S, Yu P, Yau S, Su Z *et al.* Characteristic analyses of a neural differentiation model from iPSC-derived neuron according to morphology, physiology, and global gene expression pattern. *Scientific reports* 2017; **7**(1)**:** 12233.

62. Chen TW, Wardill TJ, Sun Y, Pulver SR, Renninger SL, Baohan A *et al.* Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature* 2013; **499**(7458)**:** 295-300.

63. Steri M, Idda ML, Whalen MB, Orrù V. Genetic variants in mRNA untranslated regions. *Wiley Interdiscip Rev RNA* 2018; **9**(4)**:** e1474.

64. Wang D, Liu S, Warrell J, Won H, Shi X, Navarro FCP *et al.* Comprehensive functional genomic resource and integrative model for the human brain. *Science* 2018; **362**(6420).

65. Nunez YO, Mayfield RD. Understanding Alcoholism Through microRNA Signatures in Brains of Human Alcoholics. *Front Genet* 2012; **3:** 43.

66. Wehrspaun C, Ponting C, Marques A. Brain-expressed 3′UTR extensions strengthen miRNA cross-talk between ion channel/transporter encoding mRNAs. *Frontiers in Genetics* 2014; **5**(41).

67. Suda S, Nibuya M, Suda H, Takamatsu K, Miyazaki T, Nomura S *et al.* Potassium channel mRNAs with AU-rich elements and brain-specific expression. *Biochem Biophys Res Commun* 2002; **291**(5)**:** 1265-1271.

68. Sun JW, D.; Wong, G.; Lau, T.-M.; Yang, M.; Hart, R.P.; Kwan, K.-M.; Chan, H.Y.E.; Chow, K.H. Chronic alcohol metabolism results in DNA repair infidelity and cell-cycle-induced senescence in neurons. Submitted 2022.

69. Brenner E, Tiwari GR, Kapoor M, Liu Y, Brock A, Mayfield RD. Single cell transcriptome profiling of the human alcohol-dependent brain. *Hum Mol Genet* 2020; **29**(7)**:** 1144-1153.

70. Bodhinathan K, Slesinger PA. Molecular mechanism underlying ethanol activation of G-protein-gated inwardly rectifying potassium channels. *Proc Natl Acad Sci U S A* 2013; **110**(45)**:** 18309-18314.

71. Sathyan P, Golden HB, Miranda RC. Competing interactions between micro-RNAs determine neural progenitor survival and proliferation after ethanol exposure: evidence from an ex vivo model of the fetal cerebral cortical neuroepithelium. *J Neurosci* 2007; **27**(32)**:** 8546-8557.

72. Pietrzykowski AZ, Friesen RM, Martin GE, Puig SI, Nowak CL, Wynne PM *et al.* Posttranscriptional regulation of BK channel splice variant stability by miR-9 underlies neuroadaptation to alcohol. *Neuron* 2008; **59**(2)**:** 274-287.

73. Lewohl JM, Nunez YO, Dodd PR, Tiwari GR, Harris RA, Mayfield RD. Up-regulation of microRNAs in brain of human alcoholics. *Alcohol Clin Exp Res* 2011; **35**(11)**:** 1928-1937.

74. Osterndorff-Kahanek EA, Tiwari GR, Lopez MF, Becker HC, Harris RA, Mayfield RD. Long-term ethanol exposure: Temporal pattern of microRNA expression and associated mRNA gene networks in mouse brain. *PLoS One* 2018; **13**(1)**:** e0190841.

75. Lim Y, Beane-Ebel JE, Tanaka Y, Ning B, Husted CR, Henderson DC *et al.* Exploration of alcohol use disorder-associated brain miRNA-mRNA regulatory networks. *Translational psychiatry* 2021; **11**(1)**:** 504.

76. Zhu S, Wu J, Hu J. Non-coding RNA in alcohol use disorder by affecting synaptic plasticity. *Exp Brain Res* 2022.

77. Best TK, Siarey RJ, Galdzicki Z. Ts65Dn, a mouse model of Down syndrome, exhibits increased GABAB-induced potassium current. *J Neurophysiol* 2007; **97**(1)**:** 892-900.

78. Harashima C, Jacobowitz DM, Witta J, Borke RC, Best TK, Siarey RJ *et al.* Abnormal expression of the G-protein-activated inwardly rectifying potassium channel 2 (GIRK2) in hippocampus, frontal cortex, and substantia nigra of Ts65Dn mouse: a model of Down syndrome. *J Comp Neurol* 2006; **494**(5)**:** 815-833.

79. Reeves RH, Irving NG, Moran TH, Wohn A, Kitt C, Sisodia SS *et al.* A mouse model for Down syndrome exhibits learning and behaviour deficits. *Nat Genet* 1995; **11**(2)**:** 177-184.

80. Kleschevnikov AM, Yu J, Kim J, Lysenko LV, Zeng Z, Yu YE *et al.* Evidence that increased Kcnj6 gene dose is necessary for deficits in behavior and dentate gyrus synaptic plasticity in the Ts65Dn mouse model of Down syndrome. *Neurobiol Dis* 2017; **103:** 1-10.

81. Congdon E, Canli T. The endophenotype of impulsivity: reaching consilience through behavioral, genetic, and neuroimaging approaches. *Behav Cogn Neurosci Rev* 2005; **4**(4): 262-281.

1047
1048   82.   Dick DM, Smith G, Olausson P, Mitchell SH, Leeman RF, O'Malley SS *et al.*
1049   Understanding the construct of impulsivity and its relationship to alcohol use disorders.
1050   *Addict Biol* 2010; **15**(2)**:** 217-226.

1051
1052   83.   Sher KJ, Trull TJ. Personality and disinhibitory psychopathology: alcoholism and
1053   antisocial personality disorder. *J Abnorm Psychol* 1994; **103**(1)**:** 92-102.

1054
1055   84.   Verdejo-García A, Lawrence AJ, Clark L. Impulsivity as a vulnerability marker for
1056   substance-use disorders: review of findings from high-risk research, problem gamblers
1057   and genetic association studies. *Neurosci Biobehav Rev* 2008; **32**(4)**:** 777-810.

1058
1059   85.   Polich J, Bloom FE. P300, alcoholism heritability, and stimulus modality. *Alcohol* 1999;
1060   **17**(2)**:** 149-156.

1061

1062

1063

Figure Legends

Figure 1. Experimental design and gene expression analysis

A. Diagram outlining experimental design. Lymphocytes from subjects with or without AUD diagnosis and *KCNJ6* haplotype variants were selected, reprogrammed into iPSC, induced into excitatory iNs, and analyzed by morphometry, immunocytochemistry, gene expression, and electrophysiology. **B.** Sequencing alignment and depth analysis of bulk RNA sequencing confirmed expression of *KCNJ6* mRNA in iN cultures, specifically the ENST00000609713 isoform, containing an 18.1 kilobase 3'UTR region. *KCNJ6* exons are mapped to chromosome locations (marked in mb, megabases) and the position of the gene is indicated by the red box on the chromosome 21 pictogram, top, with transcription direction on the minus strand indicated by the broken arrow. Variant analysis of RNA sequences predicts a region of linkage disequilibrium of 22 SNPs, including the 3 SNPs used to select subjects (red, Table 1), and 19 additional SNPs (blue, Supplemental Table 1). Depth: number of sequencing reads per base aligned by position. Frequency: thickness of curved lines represents the relative frequency of splice site utilization between exons. **C.** Single-cell RNAseq identifies a cluster of induced neurons (lower left), expressing markers consistent with neuronal function including *SYP, SLC17A6, GRIN2B, SCN3A, KCNJ3*, and *KCNJ6*; distinct from "transition neurons" that either do not express these markers or express sporadically. Additional markers are plotted in Supplemental Fig. 2E. Isolating *KCNJ6* mRNA expression, aggregated by subject and treatment, AF neurons expressed a trend towards lower levels than UN neurons (p = 0.0508; Wald test), but treatment of 7d with IEE at 20 mM peak concentration increased AF expression above untreated (p = 0.0225) to levels similar to UN control (p = 0.322, not denoted on figure). **D.** Volcano plot for untreated UN vs AF neurons, highlighting genes significantly different (FDR > 0.05) and at least 1.5-fold changed (red dots). Genes below the fold-change cut-off are marked in green, and those not significantly different are marked in blue. Significantly different genes are listed in Supplemental Table 3. **F.** Gene ontology (GO) enrichment of top 10 biological process (BP) terms for up- or down-regulated genes. Plots indicate the number of regulated genes from the term and the color indicates the adjusted p-value (q-value; key). All enriched terms are listed in Supplemental Tables 4-5.

Figure 2. Validation of GIRK2 expression and function in human induced neurons.

A. Representative confocal images of GIRK2 immunoreactivity in mouse cortical neurons. Arrowheads indicate locations of GIRK2-staining puncta, with an example punctum enlarged in the inset, overlapping or adjacent to βIII-tubulin-positive processes. We observed two cellular expression patterns – one where the entire neuron is decorated with GIRK2 antibody (Supplemental Fig. 4), or another where GIRK2 expression is relatively faint and observed mostly on neuronal processes, shown here. **B.** GIRK2 expression patterns in human induced neurons (iNs), showing representative confocal images from line 420. Inset shows two adjacent puncta. **C.** Following infection of iN cultures with lentivirus expressing both *KCNJ6* and mCherry, large numbers of GIRK2$^+$ puncta are seen in representative images (line 420). **D.** Evaluation of GIRK2 function in iNs, (a) quantification of GIRK2 expression on MAP2$^+$ vs. βIII-tubulin$^+$ neuronal

processes. GIRK2 is more abundant on βIII-tubulin processes (p=0.0006, one-tailed Student's t-
test, n=15 cells per group, cell line 420). (b) Basal levels (upper pie plot) of the GIRK current in
iNs as percent of neurons responding with hyperpolarization to the selective GIRK activator
(160 nM ML297); compared with responding percentage when GIRK2 is overexpressed (lower
pie chart). (c) Representative image of iN overexpressing GIRK2, as confirmed with mCherry
fluorescence. (d) Representative traces of induced action potential firing before and after GIRK
activation, demonstrating the contribution of GIRK function to cell excitability. (e)
Representative trace of spontaneous postsynaptic potential (sEPSCs) recordings during ML297
(160 nM) GIRK activator wash-in, demonstrating a shift of 7mV holding current (amplifier-
dependent compensation of GIRK-mediated membrane hyperpolarization). (f) Quantification of
neuronal excitability at baseline and following GIRK activation with 160 nM ML297 (p=0.015,
paired Student's t-test, n=9 cells before/after ML297, cell line 376).

Figure 3. Impact of AUD-associated *KCNJ6* haplotype on neuronal properties.
A. Principles of morphological analysis of induced neurons: (a) Neurite area was the total TuJ1^{+}
(βIII-tubulin)$^{+}$ staining area outside the cell soma. (b) Solidity is the area of the soma divided by
its convex hull area. (c) Soma size was the area of the MAP2^{+} cell body. (d) Circularity compared
the perimeter to the area. **B**. Morphometry of iNs from *KCNJ6* haplotype variant and affected
(**AF**, cyan) or unaffected (**UN**, grey) individuals. Results are summed by group (left) or plotted
individually by cell line (right), with subjects identified by line number (see Table 1—females
identified with grey numbers). Individual cells are plotted as dots with the bar showing the
mean, with error bars indicating the standard error of the mean (SEM). No significant
differences were found in (a) soma size, (b) circularity, or (c) soma solidity, but total neurite
area was increased in the AF group (p =0.018). **C**. Representative images of iNs from individual
lines, with arrows identifying individual GIRK2 puncta (red) localized on βIII-tubulin^{+} processes
(gray). **D**. GIRK2 expression was decreased in the AF as measured by puncta counts (a, p =
0.0012), circularity (c, p = 0.007), or solidity (d, p = 0.037), while there was no difference in
puncta size (b). **E**. Representative images of individual GIRK2 puncta (red) localized on βIII-
tubulin^{+} processes (gray). **F**. Electrophysiological analysis of passive neuronal properties,
showing no difference in (a) membrane capacitance (b) membrane resistance, or (c)
spontaneous EPSCs frequency. (d) Representative sEPSCs traces for each line. (e) Spontaneous
EPSCs amplitude. **G**. Electrophysiological analysis of active neuronal properties. (a)
Quantification of current required to shift resting membrane potential to -65mV in pA:
difference by group p = 1.2×10^{-5}; (b) quantification of maximum number of action potentials
(APs) induced with the "step" protocol, p = 0.086; (c) representative traces of APs induced with
the "step" protocol; (d) quantification of number of action potentials (APs) induced with the
"ramp" protocol, p = 2.0×10^{-9}; (e) representative traces of APs induced with the "ramp"
protocol. A generalized linear model was used to evaluate group differences for morphometry
and GIRK2 expression, and generalized estimation equations was used for electrophysiology
results. Numbers of cell lines and replicates for each experiment are shown in Supplemental
Table 6.

Figure 4. Evaluation of neuronal excitability in human induced neuron populations.
A. Representative epifluorescence images of GcaMP6f expression in iNs from affected (233, 246) and unaffected individuals (420, 472). **B, C.** Raster plots of the calcium spiking pattern of representative experiments from unaffected neurons (**B**, individual 472) and affected neurons (**C**, individual 233), under the stimulation protocol used. Green and purple bars show the application of 30 s pulses of Glu 10-50 µM and KCl 18 mM, respectively, on the neuron populations during the stimulation protocol. Arrows show the epochs of the calcium imaging recordings selected for calcium spike quantification: baseline, glutamate pulse 1, and KCl pulse 1. **D.** Stimulation protocol (upper bar) and representative raw fluorescence trace (AF individual 233) from a complete recording. The zoom-ins on the first glutamate and KCl pulses depict glutamate and KCl elicited activity, respectively. The vertical dotted lines correspond to the calcium spikes detected after analysis. **E, F, G.** Number of spikes per ROI per minute during baseline (**E**, spontaneous activity), glutamate pulse 1 (**F**, glutamate-elicited activity) and KCl pulse 1 (**G**, KCl-elicited activity). The pie charts represent the proportion of neurons of UN (grey) and AF (blue) individuals that exhibited a certain number of calcium spikes during each three epochs of the recording. The bar plots show the number of spikes per ROI per minute fired by unaffected individuals (grey) and affected individuals (blue) during baseline, glutamate pulse 1 and KCl pulse 1. The sample size is depicted in the bar of each group as number of iN batches/number of experiments/number of neurons. Differences between affected and unaffected groups were evaluated by two-tailed unpaired Student's t-test (Welch's correction for non-equal SD, **$p < 0.01$, ****$p < 0.0001$).

Figure 5. Ethanol treatment reduced *KCNJ6* haplotype differences in iN excitability and GIRK2 expression.
A. Morphological analysis of IEE iNs generated from affected and unaffected individuals, showing no difference in: (a) neuronal soma size ($p = 0.32$); (b) soma circularity ($p = 0.54$); (c) soma solidity ($p = 0.92$); and (d) total neurite area ($p = 0.98$). **B.** There was no difference in GIRK2 expression in the IEE AF group iNs compared with UN by (a) puncta counts ($p = 0.46$), (b) puncta size ($p = 0.36$), (c) puncta circularity ($p = 0.052$), or (d) solidity ($p = 0.47$). **C.** Representative images of individual GIRK2 puncta (red) localized on βIII-Tubulin positive processes (gray) for each cell line. **D.** Electrophysiological analysis of passive neuronal properties in IEE iNs, showing (a) a small decrease in AF membrane capacitance ($p = 9.6 \times 10^{-9}$), but no difference in (b) membrane resistance ($p = 0.63$), (c) spontaneous EPSCs frequency ($p = 0.32$), or (d) spontaneous EPSCs amplitude ($p = 0.19$). (e) Representative sEPSCs traces for each cell line. (f) The AF group exhibited no change in resting membrane potential after IEE ($p = 0.79$). **E.** Electrophysiological analysis of active neuronal properties found no difference in IEE iNs for (a) current required to shift resting membrane potential to -65mV ($p = 0.32$), (b) maximum number of action potentials (APs) induced with the "step" protocol ($p = 0.48$), with (c) representative traces of APs induced with the "step" protocol, (d) number of action potentials (APs) induced with the "ramp" protocol ($p = 0.95$), with (e) representative traces of APs induced with the "ramp" protocol. **F.** Representative images from individual lines of iNs,

exposed to 7d of IEE, marked with arrows pointing to individual GIRK2 puncta (red) localized on
βIII-Tubulin positive processes (gray). **G.** Summarized results from all lines showing differences
in GIRK2 expression levels before and after 7 days of 20 mM IEE with ethanol (EtOH; p = 1.0 x
10^{-20}). **H.** Representative images of individual GIRK2 puncta (red) localized on βIII-Tubulin
positive processes (gray) prior and following 7 days 20 mM IEE with ethanol. Sample images are
from line 246. **I.** Representative images of FISH detection of *KCNJ6* mRNA for each cell line. **J.**
Quantification of FISH. (a) The number of *KCNJ6* puncta normalized to the number of cells in an
image shows decreased expression in control AF compared with UN (p = 8.2 x 10^{-3}), increased
expression following IEE (p = 8.2 x 10^{-3}; using Tukey's pairwise comparisons). (b) The percentage
of *KCNJ6*-expressing MAP2$^+$ cells substantially increase in the (c) AF group but not in the (b) UN
group. Numbers of *KCNJ6* puncta were analyzed by expression levels per cell, as recommended
by the FISH manufacturer in (d) UN or (e) AF cells. (f) *KCNJ6* puncta within the neuronal soma
show increases following IEE in both UN and AF groups (p = 0.006 for genotype, Tukey's post-
hoc for UN, p = 0.01, for AF, p = 0.01). (g) A similar analysis of non-somatic puncta, presumably
within neurites, showed no differences following IEE between genotypes (p=0.23).

Figure 6. GIRK2 overexpression mimics ethanol response
A. Current required to shift resting membrane potential to -65mV, in untreated (control, p = 5.9
x 10^{-4}), lentiviral *KCNJ6* overexpression (over., p = 0.23), or 1 d 20 mM IEE (EtOH, p = 0.75)
cultures. **B.** Representative traces of APs induced with the "ramp" protocol. **C.** Quantification of
maximum number of action potentials (APs) induced with "ramp" protocol, control (p = 6.8 x
10^{-6}), overexpression (p = 0.014), or 1 d 20 mM IEE (p = 0.10). **D.** quantification of GIRK2 puncta
after overexpression (line 376, one-tailed Student's t-test p = 0.04).

1205

Supplementary Materials

1206

Supplemental Tables 1-6: https://doi.org/10.6084/m9.figshare.19798882.v2

Supplemental Table 1. SNPs identified in RNAseq samples. Variants were identified in the
region surrounding KCNJ6 (chr21: 37499112-38245792) using samtools mpileup to bcftools call,
followed by filtering to remove low quality. Results were loaded into the Ensembl VEP tool. A
dot (".") indicates missing data and/or low quality. The reference allele and alternate alleles
were obtained from the UCSC Genome Table Browser.

Supplemental Table 2: CellRanger parameters for scRNAseq libraries. Output from CellRanger
shows the alignment statistics and numbers of reads confidently mapped.

Supplemental Table 3. Differentially expressed genes (DEG) comparing AF to UN without
exposure to ethanol. The table contains all output from DESeq2 results. Excel filters are set to
show only genes significantly different (padj <= 0.05) and at least 1.5-fold different
(abs(log2FoldChange) > 0.585). Release the filters to see the full list.

Supplemental Table 4. Enriched Gene Ontology-Biological Process (GO-BP) terms from genes
increased in AF relative to UN.

Supplemental Table 5. Enriched Gene Ontology-Biological Process (GO-BP) terms from genes
decreased in AF relative to UN.

Supplemental Table 6. Numbers of replicates. For each parameter tested with statistics, the
numbers of cells and/or microscope fields is indicated and labeled by Figure numbers/letters.

Supplemental Figures: https://doi.org/10.6084/m9.figshare.19798873.v3

1226

Figure 1, Popova et al., 2022

A GIRK2 expression - mouse cortical neurons

B GIRK2 expression - human induced neurons (iN)

C GIRK2 overexpression in human iN

D GIRK2 function in human iN

Figure 2, Popova et al., 2022

A Principles of morphological analysis
B Morphometry

C Representative images

D GIRK2 expression
E GIRK2 expression

F Basal neuronal properties

G Induced neuronal properties

Figure 3, Popova et al., 2022

A GCaMP6f expression

B Peak activity raster plot **UF**

C Peak activity raster plot **AF**

D Stim. protocol / representative traces

E Spontaneous activity

F Glutamate-elicited activity

G KCl-elicited activity

Figure 4, Popova et al., 2022

Figure 5, Popova et al., 2022

Active neuronal properties

Figure 6, Popova et al., 2022