TELOMERASE-DEPENDENT AGEING IN THE ZEBRAFISH BRAIN

Raquel R. Martins[1,2,4], Savandara Besse[3], Pam S. Ellis[1, 2], Naomi Hartopp[4], Nadiyah Mughal[1,

2], Owain Evans[5], Mina HF Wahib[5], Zerkif Yazigan[5], Heather Mortiboys[4], Michael Rera[3] and

Catarina M. Henriques[1, 2]

1.   Division of Clinical Medicine, Healthy Lifespan Institute and MRC-Arthritis Research UK Centre for

Integrated research into Musculoskeletal Ageing, University of Sheffield, Sheffield, UK

2.   Bateson Centre, University of Sheffield, Sheffield, S10 2TN, UK.

3.   Institut Jacques Monod (UMR7592), Paris, France.

4.   Sheffield Institute for Translational Neuroscience (SITraN), University of Sheffield, 385a Glossop Road,

Sheffield S10 2HQ, U.K

5.   MBChB Medicine Degree, University of Sheffield

Corresponding author: c.m.henriques@sheffield.ac.uk

GRAPHICAL ABSTRACT

Graphical Abstract: The zebrafish brain displays telomerase-dependent transcriptional hallmarks of ageing

which associate with telomerase-dependent accumulation of senescence-associated markers and

inflammation in the brain. Importantly, *in vivo*, telomerase deficiency accelerates dysfunction of the blood

brain barrier and behavioural change with ageing.

ABSTRACT

Decreased telomerase expression, telomere shortening, senescence-associated markers and

inflammation have all been independently observed in the ageing brain and associated with

disease. However, causality between limited telomerase expression and brain senescence
and neuro-inflammation in the natural ageing setting is yet to be established. Here, we
address these questions using the zebrafish as an ageing model which, akin to humans,
displays premature ageing and death in the absence of telomerase and where telomere
shortening is a driver of cellular senescence.

Our work shows for the first time that telomerase deficiency (*tert*$^{-/-}$) accelerates key
hallmarks of ageing identified in the Wild Type (WT) zebrafish brain at transcriptional,
cellular, tissue and functional levels. We show that Tert-dependent transcriptomic changes
associated with dysregulation of gene expression, stress response and dysregulation of
immune genes are accompanied by accelerated accumulation of senescence-associated
markers and inflammation in the aged brain. Importantly, *In vivo*, these changes correlate
with increased blood-brain barrier permeability and altered cognitive behaviour. Of note,
telomerase-dependent accumulation of senescence-associated markers in the brain occurs
not only in the expected proliferative areas but also in non-proliferative ones, where it is
unlikely due to telomere-dependent replicative exhaustion, suggesting that non-canonical
roles of telomerase may be involved.

Together, our work suggests that telomerase has a protective role in the zebrafish brain
against the accumulation of senescence and neuro-inflammation and is required for blood
brain barrier integrity.

KEY WORDS: Zebrafish, telomerase, TERT, ageing, brain, inflammation, Blood Brain Barrier
(BBB), senescence, behaviour, transcriptomics

ABBREVIATIONS:

AD, Alzheimer's disease

ALS, amyotrophic lateral sclerosis

ANOVA, analysis of variance

ATH, ageing transcriptional hallmarks

BBB, blood brain barrier

BCA, bicinchoninic acid

bp, base pair

BSA, bovine serum albumin

Ce, cerebellum

Cent, centromere

CNS, central nervous system

DAPI, 4',6-diamidino-2-phenylindole

DAVID, database for annotation, visualization, and integrated discovery

DDR, DNA damage response

DEG, differential expression gene

Die, diencephalon

dH2O, distilled water

DNA, deoxyribonucleic acid

Dpf, day post fertilisation

dscDNA, double-stranded cDNA

EDTA, ethylenediaminetetraacetic acid

ENU, N-ethyl-N-nitrosourea

FAM, carboxyfluorescein

FISH, fluorescence *in situ* hybridisation

GEO, gene expression omnibus

GOBP, gene ontology biological pathway

h, hours

HBSS, Hanks buffered solution

IF, immunofluorescence

IP, intraperitoneal injection

IR, irradiation

KCN, potassium cyanide

KEEG, Kyoto encyclopaedia of genes and genomes

KH_2PO_4, potassium dihydrogen phosphate

KO, knock-out

LCP1, Lplastin

LMF, Wolfson light microscopy facility

LPS, lipopolysaccharide

MgCl2, magnesium chloride

Min, minutes

MO, medulla oblongata

NaCl, sodium chloride

NADH, nicotinamide adenine dinucleotide reduced disodium salt

NFkβ, nuclear factor kappa β

nm, nanometers

NO, novel object

NPCs, neural progenitor cells

NSCs, neural stem cells

O.C.T., mounting media

OT, optic tectum

PBS, phosphate buffered saline

PCA, principal component analysis

PCNA, proliferating cell nuclear antigen

PFA, paraformaldehyde

PNA, peptide nucleic acid

RIN, RNA integrity number

RNA, ribonucleic acid

RPM, revolutions per minute

RT, room temperature

SASP, senescence-associated secretory phenotype

sec, seconds

SITraN, Sheffield Institute for Translational Neuroscience

SSC, saline-sodium citrate

TE, Tris-EDTA

Tel, telencephalon

Telo, telomere

Tert, telomerase

TERT, telomerase reverse transcriptase

TNFα, tumour necrosis factor alpha

WT, wild-typeZIRC, Zebrafish international resource center

4-MU-chitotrioside , 4–methylumbelliferyl–β-D-N, N', N''-triacetylchitotriose

2,6-DCPIP, 2,6-dichlorobenzenone-indophenol sodium salt

1 INTRODUCTION

Brain ageing and related neurodegenerative diseases have dramatic consequences on

quality of life and medical care costs. Dementia alone, one of the most prevalent

manifestations of neurodegenerative disease affects 23 million people only in the UK and it

costs around £26 billion a year to the NHS (Collins et al., 2022; Panca et al., 2019; Wittenberg

et al., 2019). Understanding the mechanisms underlying brain ageing will target multiple age-

associated brain diseases rather than one disease, aiming to ameliorate quality of life in the

elderly.

There are well-known key hallmarks of ageing, which have recently been expanded and

consist of genomic instability, telomere attrition, epigenetic alterations, loss of proteostasis,

disabled macroautophagy, deregulated nutrient sensing, mitochondrial dysfunction, cellular

senescence, stem cell exhaustion, altered intercellular communication, chronic

inflammation and dysbiosis (Lemoine, 2021; Lopez-Otin, Blasco, Partridge, Serrano, &

Kroemer, 2023). Telomeres are (TTAGGG)n DNA repeats that together with a complex of

proteins (known as Shelterin) create a "cap-like" structure at the end of linear chromosomes

(de Lange, 2004), preventing the ends of linear chromosomes from being recognised as

deleterious DNA double strand breaks (M. G. Ferreira, Miller, & Cooper, 2004). However, in

humans, due to time- and cell-specific-limited telomerase expression, telomeres shorten

with each cell division, eventually leading to proliferative exhaustion and replicative

senescence (Bodnar, 1998; d'Adda di Fagagna et al., 2003). Even though senescence cells do

not proliferate, they are metabolically active, releasing several inflammatory factors, tissue

remodelling factors and growth factors, known as Senescence Associated Secretory

Phenotype (SASP) (Coppe et al., 2010). Consequently, accumulation of senescent cells

contributes to impairment of tissue homeostasis and a low level of chronic inflammation.

Importantly, there is accumulation of cellular senescence, which can be thought of as a

general stress response (Birch & Gil, 2020; Fridlyanskaya, Alekseenko, & Nikolsky, 2015), with

ageing in humans (Dimri et al., 1995) and senescence has been linked to several age-

associated diseases (Ovadya & Krizhanovsky, 2014; Tuttle, Luesken, Waaijer, & Maier, 2021).

Restricted telomerase expression and consequent telomere dysfunction function are key

hallmarks of natural ageing in humans, underpinning multiple age-related diseases

(Blackburn, Epel, & Lin, 2015). As tissues with high cellular turnover present accelerated

telomere erosion (Bodnar, 1998; H. W. Lee et al., 1998), it is reasonable to speculate that

telomerase functions are likely to primarily affect highly proliferative tissues. Accordingly,

premature accumulation of critically short telomeres has been identified in highly

proliferative tissues such as the gut, in tert-deficient animal models, including late

generation TERT KO mice and zebrafish *tert*$^{-/-}$ mutants (Anchelin et al., 2013; Carneiro et al.,

2016; Henriques, Carneiro, Tenente, Jacinto, & Ferreira, 2013; H. W. Lee et al., 1998).

Nevertheless, because senescence has been shown to "spread", in a phenomenon described

as paracrine senescence, telomere-dependent genome instability and replicative senescence

may lead to non-cell autonomous increase in senescence-associated markers and

inflammation (Lex et al., 2020). This means that the role of telomerase and telomeres is not

restricted to highly proliferative tissues. To complicate things further, growing evidence

suggests that the Tert component of telomerase also has non-canonical activities,

independent of its action at telomeres (Goodman & Jain, 2011; Romaniuk et al., 2018; Segal-

Bendirdjian & Geli, 2019; Sung, Ali, & Lee, 2014). In the nucleus, these non-canonical

functions include transcriptional regulation of genes involved in inflammation, including

nuclear factor kappa β (NFkβ) and tumour necrosis factor alpha (TNFα) (Deacon & Knox,

2018; Ghosh et al., 2012; Mattiussi, Tilman, Lenglez, & Decottignies, 2012), as well as genes

involved in cell proliferation (Choi et al., 2008; Sarin et al., 2005) and cell survival (Cao, Li,

Deb, & Liu, 2002; Rahman, Latonen, & Wiman, 2005). Telomerase can also translocate to the

mitochondria, where it has been shown to play a protective role against DNA damage and

oxidative stress (Ahmed et al., 2008; Haendeler et al., 2009). In the brain, considered a

predominantly post-mitotic tissue, telomerase has been shown to have a protective role

against excitotoxicity (Eitan et al., 2012), oxidative stress (Spilsbury, Miwa, Attems, &

Saretzki, 2015), and neuronal death (J. Lee et al., 2010), all involved in neurodegenerative

diseases. Studies in late-generation telomerase-deficient mice have suggested that limited

telomerase expression is associated with premature accumulation of senescence-associated

markers in different cell populations including Purkinje neurons, cortical neurons and

microglia (De Felice et al., 2014; Jaskelioff et al., 2011; Jurk et al., 2012; Raj et al., 2015).

However, it is still unclear whether this is due to telomere length maintenance or non-

canonical functions of telomerase (Choi et al., 2008; Sarin et al., 2005).

There are many different stimuli that can cause cellular senescence, including epigenetic

alterations, oxidative stress, mitochondrial dysfunction, inactivation of tumour suppressor

genes, mechanical or shear stress, pathogens, and activation of oncogenes (Hernandez-

Segura, Nehme, & Demaria, 2018; Tripathi, Misra, Tchkonia, & Kirkland, 2021). What causal

role telomerase and telomere dysfunction plays in CNS senescence in humans is, however,

not clear, even if implicated (Yu et al., 2023). Nevertheless, telomere shortening has been

associated with neurodegenerative diseases, including Parkinson's and AD
(Papageorgakopoulou, Bania, Lagogianni, Birmpas, & Assimakopoulou, 2024), including via
glia activation and pathogenesis (Vellingiri et al., 2023). Of note, a recent study analysing
human brain biobank data from over 30,000 patients identified a positive correlation
between shorter leukocyte telomere length (used as a proxy for the patient's general
telomere length) and neurodegenerative phenotypes (Topiwala et al., 2023). Additionally,
cellular senescence-associated markers have been reported in human neuroinflammation
and neurodegenerative diseases. In particular, increased levels of DNA damage checkpoints
such as p53, p21 and p16, as well as SASP factors have been observed in the spinal cord and
brain of patients with amyotrophic lateral sclerosis (ALS) (De Felice et al., 2014) and
Alzheimer's disease (AD) (Bhat et al., 2012), respectively. Higher mRNA levels of p16INK4a in
the substantia nigra pars compacta (SNpc) were also observed in post-mortem tissue of
patients with Parkinson's disease (PD) compared to healthy controls (Chinta et al., 2018) .
Moreover, senescence, neuroinflammation and oligomeric tau have been suggested to
associated with AD pathology, recently reviewed here (Gaikwad, Senapati, Haque, & Kayed,
2024). Hence, the literature shows that despite being a low proliferative tissue, the CNS
accumulates senescence or at least senescence-associated markers, and that this may be
associated with brain diseases of ageing. However, causality between limited telomerase
expression and increased senescence and neuro-inflammation in the natural ageing setting
is yet to be established. Nevertheless, mice models where putative senescent cells have been
cleared (Bussian et al., 2018) show an improvement of the cognitive decline in aged mice,
suggesting that senescence can actively contribute to it. It remains to be determined which
cell populations undergo senescence in natural ageing, in the CNS, and what type of
senescence this is, i.e., replicative (telomere-dependent) or stress induced senescence.
Understanding whether and how telomerase restriction may contribute to ageing-associated
brain degenerative phenotypes would help understand the fundamental mechanisms of
brain functional decline and potentially contribute to the uncovering of new therapeutic
targets to delay or prevent disease.
223
224 The role of telomerase restriction and its consequences in natural ageing may benefit from
225 additional and complementary models, beyond the mouse, where most lab-strains have
226 significant longer telomeres than humans and show little evidence of replicative senescence

(Forsyth, Wright, & Shay, 2002; Gomes et al., 2011; Sullivan et al., 2021). The zebrafish may offer such a model since, like humans, they depend on telomerase for health and lifespan (Anchelin et al., 2013; Carneiro et al., 2016; Henriques et al., 2013; Henriques & Ferreira, 2012). As it was shown that the zebrafish brain presents low levels of Tert in the adult brain, which tend to decrease with ageing (Sprungala, 2009; Zambusi, Pelin Burhan, Di Giaimo, Schmid, & Ninkovic, 2020), we hypothesised that limited expression of Tert in the aged zebrafish brain contributes to brain degeneration with ageing. To our knowledge, this manuscript represents the first broad characterisation of zebrafish brain ageing in the presence and absence of telomerase (Tert).

Our data show that the zebrafish brain undergoes Tert-dependent age-associated changes at both transcriptional and functional levels. In specific, our transcriptomic data and enrichment analysis identified dysregulation of gene expression, stress response and dysregulation of immune genes as the top 3 significantly enriched transcriptional hallmarks of old age (Frenk & Houseley, 2018) in WT. Within these differentially expressed genes (DEGs) of old age (WT young *VS* WT old), we identified about 24% which were accelerated in the absence of Tert. When enriched and mapped to the previously described transcriptomic hallmarks of ageing, these Tert-dependent genes of old age identified the same top 3 significantly enriched ageing transcriptional hallmarks (ATH) (Frenk & Houseley, 2018) as the overall DEGs of aged WT. Notably, most of the Tert-dependent hallmarks of ageing already significantly accelerated by the age of 9 months in $tert^{-/-}$ zebrafish, contrasting to >30 months in WT. At the tissue level, telomerase-dependent stress response and immune dysregulation transcriptional changes with ageing were accompanied by an accelerated accumulation of senescence-associated markers and inflammation. *In vivo*, these changes correlated with increased blood-brain barrier permeability and altered cognitive behaviour. Together, our work suggests that telomerase has a protective role in the zebrafish brain against the accumulation of senescence and neuro-inflammation and is required for blood brain barrier integrity.

2 RESULTS

2.1 Tert-dependent transcriptomic changes of ageing in the zebrafish brain

Tissue-specific transcriptomic analysis over the life course can offer important insights into the downstream molecular mechanisms underpinning age-associated pathologies. Significant research is being dedicated to these approaches in different animal models, including in mice (Schaum et al., 2020; Tabula Muris, 2020; Zhang, Pisco, Darmanis, & Zou, 2021). To identify telomerase-dependent and -independent transcriptional signatures of ageing in the zebrafish brain, we performed RNA-Sequencing of whole brain tissue, throughout the lifespan of WT and telomerase-deficient (*tert*$^{-/-}$) fish (**Fig. 1** and supplementary source data files). While *tert*$^{-/-}$ fish have a lifespan of c. 12-20 months, WT fish typically die between c. 36-42 months of age (Carneiro et al., 2016). The data shown here include 4 age-groups of WT (2, 9, 22 and >30 months), corresponding to young, adult, median lifespan and old, respectively. As the *tert*$^{-/-}$ fish have a shorter lifespan compared with their WT siblings, the data include 3 age-groups of telomerase-deficient fish (2, 9, and 22 months), which correspond to young, medium lifespan and old (see Materials & Methods for further explanation of genotypes and ages chosen) (**Fig 1A**). The reads were aligned to the latest zebrafish genome build GRCz11 (Lawson et al., 2020) and resulted in uniquely mapped read percentages ranging from 92.1% to 94.4%, which is considered good quality (Lawson et al., 2020). To analyse the overall impact of the genotype and age on transcriptomic regulation, we performed a Principal Component Analysis (PCA). We observed that the samples cluster mainly *per* age, but within each age group we observe a distinction between genotypes (**Supp Fig 1A**). This provided the first hint of a change in the ageing transcriptomic profile in the *tert*$^{-/-}$. A summary of the number of significant differentially expressed genes (DEGs) over the life course of WT and *tert*$^{-/-}$ is represented in **Supp Fig 1B-D** (for further details see supplementary source data files).

To identify a broad transcriptomic signature of old age, we compared gene expression changes between young (c.2 months) and old (>30 months) WT (**Fig 1B**). Once these DEGs of old age were identified (for further details see supplementary source data files), we performed Gene Ontology Biological Pathway (GOBP) and Kyoto Encyclopaedia of Genes and Genomes (KEGG) enrichment analysis (**Fig 1B1**) using DAVID software. GOBP analysis of DEGs of old age identified cell cycle, division, chromatin-organisation and chromosome

segregation and downregulation of DNA transcription as the top 5 enriched pathways. KEGG analysis mainly identified pathways involved in immune responses (phagosome, infection, PPAR signalling), cell cycle (cell cycle, cellular senescence) and gene regulation (polycomb repressive complex). To contextualise these pathway enrichments within the hallmarks of ageing, we used a previously described strategy (Zane et al., 2023) (see additional details in Materials and Methods Section 9), whereby known GOBP terms are mapped to experimentally identified ageing transcriptomic hallmarks (ATH) (Frenk & Houseley, 2018). ATH mapping identified dysregulation of gene expression, stress response and dysregulation of immune genes as the top 3 significantly enriched transcriptional hallmarks of old age (Frenk & Houseley, 2018) in WT, followed by growth factor reduction, downregulation of mitochondrial proteins and ribosome downregulation(**Fig 1B2**, **Supp Fig 1C** and supplementary data source files). The specific genes associated with each ATH in WT and *tert*$^{-/-}$ over-time are described in **Supp Fig 3** and supplementary data source files.

To identify how many of the DEGs of old age were Tert-dependent, we compared DEGs in *tert*$^{-/-}$ zebrafish brains over time with WT DEGs of old age. We selected only the ones that changed in *tert*$^{-/-}$, in the same direction as WT, i.e., either up or downregulated. We then compared these with WT DEGs at each timepoint, to ensure they were indeed accelerated in *tert*$^{-/-}$, i.e., changed at younger ages than WT (**Supp Fig 2**). Using this strategy, we identified 704 *tert*-dependent DEGs (**Fig 1 B3**), which comprised c.24% of the total DEGs of old age (WT young *VS* WT old). Most GOBP enriched terms of *tert*-dependent DEGs of old age related to pathways involved in gene transcription and neuronal homeostasis. KEGG analysis identified changes in metabolism and immunity (such as insulin and PPAR signalling pathways), actin cytoskeleton and neuronal stem cell function (hedgehog signalling pathway). When these enrichments were mapped to the previously described ATH (**Fig 1 B4**), the *tert*-dependent gene changes of old age identified dysregulation of gene expression, stress response and dysregulation of immune genes as the top 3 significantly enriched transcriptional hallmarks (Frenk & Houseley, 2018), similarly to the overall DEGs of WT old, and to similar extents. These were followed by growth factor reduction, downregulation of mitochondrial proteins and ribosome downregulation. The specific genes associated with each ATH and whether they were up or downregulated are described in **Fig 1 B5 and** supplementary data source files. Notably, most of the *tert*-dependent DEGs of old age, irrespectively of whether

323 enriched and identified as an ageing transcriptomic hallmark, are already changed by the age

324 of 9 months in *tert*^{-/-} zebrafish, contrasting to >30 months in WT (**Supp Fig2**), highlighting a

325 significant acceleration of these hallmarks of ageing in the absence of telomerase (Tert).

A) Study design for transcriptomics

Brain

Analysis of significantly differentially expressed genes (DEGs) between genotypes at the different ages

Categorise according to Ageing Transcriptomic Hallmarks (ATH) as described by Frenk & Houseley, 2018

Determine telomerase (*tert*)-dependencies

Age (months)

B) Differential Gene Expression (DEGs) and enrichment analysis

B2)

GO BP enrichment analysis WT 2 *VS* 35 months

B1) **WT**

Youngest *VS* Oldest

2 ⌒ 35 months

2887 DEGs

KEGG enrichment analysis WT 2 *VS* 35 months

ATH2 Stress response 22.7%

ATH3 Downregulation of mitochondrial proteins 3.4%

ATH1 Dysregulation of immune genes 23.5%

ATH4 Ribosome downregulation 0.8%

ATH6 Dysregulation of gene expression 34.5%

ATH5 Growth factor reduction 15.1%

B3) Which gene changes of WT ageing are *tert*-dependent (accelerated in absence of tert)?

WT Youngest *VS* Oldest

2 ⌒ 35 months

GO BP enrichment analysis tert-dependent genes

B4)

2183 75.6%   704 24.4%

tert-independent genes of ageing

tert-dependent genes of ageing

KEGG enrichment analysis tert-dependent genes

ATH2 Stress response 15.8%

ATH3 Downregulation of mitochondrial proteins 5.3%

ATH1 Dysregulation of immune genes 22.1%

ATH4 Ribosome downregulation 2.1%

ATH6 Dysregulation of gene expression 40%

ATH5 Growth factor reduction 14.7%

B5) Upregulated   Downregulated

ATH1 Dysregulation of immune genes		ATH2 Stress response		ATH3 Downregulation of mitochondrial proteins		ATH4 Ribosome downregulation		ATH5 Growth factor reduction		ATH6 Dysregulation of gene expression			
c8b *cd74a* *zbtb38* *tfe3b* *mhc2d4b*	NA	*ino80db* *macrod1* *parp3* *polh*	*ccnd1* *e2f8* *hpf1* *nabp1a* *rev3l* *smc1al* *spata18* *uba1*	*sdhb* *sdhc*	*idh1*	*riok1*	*adh5*	*glg1a* *ncl*	*marcksl1a* *pdgfbb*	*bsdc1* *eef2l2* *eif1axb* *eif3hb* *eif4a1a* *eif5* *fdxacb1* *mbd1a*	*mrps12* *trim2b* *trim3a* *trim3b*	*dhx29* *eef1b2* *igf2bp3* *mettl13* *nanos1* *rpl32* *sh3bgrl* *ubxn6*	*vars2* *ybx1*

327 **Figure 1. Transcriptomic signatures of ageing in the presence and absence of telomerase in the zebrafish**

328 **brain.** (A) Schematic figure of the study design for transcriptomics: RNA from whole brain tissue from young

(ca. 2 months), young adult (ca. 9 months), middle aged (ca. 22 months) and old (time of death, >30 months) WT fish, and young (ca. 2 months), middle aged (ca. 9 months) and old (time of death, ca. 22 months) $tert^{-/-}$ fish was used for RNA sequencing. N=3 per group. (B1) Differential gene expression (DEGs) analysis using DESeq2 identified a total of 2887 DEGs with natural ageing (WT). Heatmaps show the top 10 enriched pathways identified in the GOBP and KEGG databases (analysis performed in DAVID). (B2) The enriched terms and pathways identified to be associated with natural ageing were mapped to the ageing transcriptomic hallmarks (ATH), as previously described (Frenk & Houseley, 2018). (B3) Within the DEGs associated with natural ageing, 24.4% (704) were accelerated in the absence of telomerase ($tert^{-/-}$). Heatmaps show the top 10 enriched pathways (GOBP and KEGG databases) associated with tert-dependent genes of ageing. (B4) The mapping of the enriched pathways associated with tert-dependent ageing to the ATH showed the same pattern identified in natural ageing, with dysregulation of gene expression, dysregulation of immune system, and stress response being the top 3 significantly changed transcriptional hallmarks of ageing. (B5) Table with the specific genes associated with each ATH.

2.2 Tert-dependent stress response with ageing in the zebrafish brain

Given that the top 3 main biological pathways identified by our transcriptomic analysis of old age and telomerase-dependent pathways of ageing included stress response and immune dysfunction, we started by assessing senescence-associated markers in the zebrafish brain, over the life course, in the presence and absence of telomerase (**Fig 2A**). There is not a single marker that can be used on its own to irrefutably identify a senescent cell and markers often described to be associated with senescence in proliferative tissues, such as P16 and P21, are highly expressed in terminally differentiated neurons (Huang, Hickson, Eirin, Kirkland, & Lerman, 2022; Vazquez-Villasenor et al., 2020). This may confound identification of putative senescent cells. Nevertheless, markers such as P16 have previously identified as a good marker for predicting disease progression in Parkinson's Disease (PD) patients (Martin-Ruiz et al., 2020), and with ageing (Idda et al., 2020). Acknowledging these limitations, we performed a broad characterisation of putative cellular senescence as a stress response using four main readouts (Hernandez-Segura et al., 2018): accumulation of Senescence-Associated β Galactosidase (SA-β-Gal) a known biomarker of cellular senescence, at least in most of the cell types tested (Dimri et al., 1995), and often present in different types of senescence (**Fig 2B**); levels of DNA Damage Response (DDR) by γH2AX expression levels (**Fig 2C and Supp Fig 4A**); levels of expression of cell cycle kinase inhibitors *p21* and *p16* genes (**Fig 2D**); and proliferation, by Proliferating Cell Nuclear Antigen (PCNA) expression (**Supp Fig 4D**).

Our initial whole brain SA-β-Gal staining in WT zebrafish brains detected more intense blue staining in WT old (>30 months), when compared to young (c. 5 months) (**Fig 2B and B1**). We then dissected the brains to determine which regions had increased SA-β-Gal staining. At this point, we asked whether SA-β-Gal staining was accelerated in the absence of telomerase, by comparing the staining in WT *versus* $tert^{-/-}$ zebrafish brains. Our results showed a significant increase in SA-β-Gal with ageing in specific regions of the WT zebrafish brain, such as in the midline of the telencephalon, diencephalon, cerebellum and medulla oblongata (**Fig 2B2**). Accumulation of SA-β-Gal in both the diencephalon (quantification shown as a representative example in **Fig 2B3**) and the cerebellum was significantly accelerated in the absence of telomerase ($tert^{-/-}$), where, at 5 months, the $tert^{-/-}$ brain is already displaying the pattern and levels of SA-β-Gal observed in old WT brains of >30 months of age. Another common marker of senescent cells is the accumulation of DNA damage response (DDR) markers, such as γH2AX foci. Our data show that very few cells accumulate γH2AX foci in the aged brain, in all macroareas tested (never more than 2% of cells). Nevertheless, we did identify a significant increase in the % of γH2AX foci$^+$ cells in the diencephalon and cerebellum of old WT, when compared to middle-aged adults, and this was accelerated in $tert^{-/-}$ fish (**Fig 2C** and **Supp Fig 4A**). To test whether increase in senescence associated markers correlated with shorter telomeres (i.e. putative replicative senescence), we measured relative telomere length in brain sections of WT fish over time, alongside their $tert^{-/-}$ counterparts as previously described by us, using a telomeric Peptide Nucleic Acid (PNA) probe alongside a centromeric PNA probe in Fluorescence *in situ* Hybridisation (FISH) (Ellis et al., 2022) (**Fig 2E** and **Supp Fig 4C**). We show that relative telomere length decreases with ageing in WT brains and is significantly shorter in $tert^{-/-}$ from an earlier age, as has been shown before for multiple tissues (Carneiro et al., 2016). However, as expected, relative telomere length in the brain is significantly longer than in the highly proliferative gut epithelia of the same individuals, both in WT and $tert^{-/-}$, at all ages tested. Despite observing telomere shortening in the brain, we did not detect significant changes in cell proliferation, as assessed by PCNA expression, in any of the areas tested, including in the known proliferative niches present in parts of the diencephalon and midline of the telencephalon (**Supp Fig 4B**). Proliferation levels, however, are very low to start with, and we never detected more than 6% of cells proliferating (**Supp Fig 4D**). Increased levels of p16 and *p21* are usually observed in senescent cells, depending on the stimuli and cell type (Dulic, Beney,

Frebourg, Drullinger, & Stein, 2000). We therefore tested whether increased levels of these
two cyclin inhibitors correlate with increased levels of SA-β-Gal in different brain regions.
Levels of *p16* and *p21* were assessed by RNA *in-situ* hybridisation. We detected increased
levels of *p16*, but not *p21*, in the proliferative grey area of the optic tectum and in some
regions of the diencephalon of old WT fish, where SA-β-Gal is also observed. Increased SA-
β-Gal and *p16* expression was observed at an earlier age, in *tert-/-* fish (**Fig 2D**).

Figure 2. Telomerase-dependent stress response in the zebrafish brain with ageing. (A) Schematic figure of
the study design for stress response: brain tissue collected from young (ca. 5 months), middle aged (ca. 12

months) and old (time of death, >30 months) WT fish, and young (ca. 5 months) and old (time of death, ca. 12 months) $tert^{-/-}$ fish were used to assess stress response through SA-β-Gal staining (in whole brain and coronal cryosections of telencephalon, optic tectum and rombencephalon), telomere length and DNA damage (in longitudinal paraffin sections), and metabolomics (in whole brain tissue). (B) Representative images of SA-B-Gal staining in the whole WT zebrafish brain and (B1) respective quantifications show an increased expression of SA- β -Gal in old WT brains compared to young ones. N=5-6 per group. (B2) Representative images of SA-B-Gal staining in coronal cryosections of WT and $tert^{-/-}$ zebrafish show that there is an increased expression of SA-B-Gal in old WT zebrafish and that this is anticipated in young $tert^{-/-}$ (ca. 5 months), particularly in the diencephalon for which (B3) quantifications are shown. N=8-14 per group. (C) DNA damage response assessed by γH2AX staining in longitudinal paraffin sections of zebrafish brain show increased DNA damage from 12 to >30 months of age in WT fish, and this increase is anticipated in the absence of telomerase at 12 months of age. N=3-7 per group. (D) Representative images from in-situ hybridisation in coronal cryosections show an increased expression of *p16* but not *p21* in old WT zebrafish optic tectum and diencephalon, in areas where SA- β -Gal expression is observed. This increase in *p16* expression is already significant at 12 months of age in $tert^{-/-}$. (E)Telomere/centromere ratio assessed by telo-FISH in longitudinal paraffin sections of zebrafish brain (white) and gut (yellow) show that telomere length decreases with ageing in WT fish and that this is anticipated in young $tert^{-/-}$ fish (ca. 5 months). N=4-5 per group. (B1, B3, C) Each dot represents one animal. (E) Each dot represents one cell. Error bars: SEM. * $p<0.05$; ** $p<0.01$; *** $p<0.001$. Abbreviations: OB, olfactive bulb; Tel, telencephalon; Die, diencephalon; OT, optic tectum; Ce, cerebellum; MO, medulla oblongata.

Different types of senescence stimuli have been identified to express different SASP factors and an attempt has been made to collate the variety of SASP factors assembled by proteomic analysis of conditioned media from different types of *in vitro* senescence (Basisty et al., 2020), some of which can be found in http://www.saspatlas.com/. Additionally, the same work identified 3 key secreted proteins in common to some DDR and DDR-independent types of senescence, namely CXCL1, MMP1 and STC1. Moreover, GDF15, STC1 and SERPINs were identified as SASP-associated factors in aged human plasma (Basisty et al., 2020). Our RNA sequencing data identified 13 putative SASP genes upregulated in aged WT brains, using SASPatlas.com (**Supp. Fig 5**), namely *arpc1b, b2m, ccdc88b, dcn, fhod1, fkbp3, flnb, ifi35, itga5, lmna, myo9b, naga, pfn1, psme1 and 2, rbp4, scin and thy1*. Of these, 5 were *Tert*-dependent, namely *arpc1b, fhod1, psme1* and *2 and rbp4*. Interestingly, most of the SASP molecules up-regulated (8 out of 13) in WT aged brains were also upregulated in oncogene induced senescence of fibroblasts, whereas the genotoxic stress SASP in the same cells had less in common with WT aged brains, according to saspatlas.com (**Supp Fig 5**). We therefore searched for these genes in our transcriptomic data and found related molecules upregulated

in aged WT zebrafish brains, namely *cxcl12a*, *cxcl32b.1* and *cxcl18b* (zebrafish *cxcl1 per se* has not yet been described); *mmp13b*, *stc2a* and serpina10a, *serpinb1l3* and *serpinb1l1*. Of these, *mmp13b* upregulation was significantly accelerated in *tert$^{-/-}$* fish, where upregulation is already significant at the age of 2 months, contrasting to >30 months in WT. Additionally, mmp13 is 5 times more upregulated in *tert$^{-/-}$* at 2 months than WT at >30 months, with log2 expression of 5.3 *versus* 1.2, respectively. Expression of this putative SASP gene is not only accelerated but also exacerbated in the absence of Tert (supplementary source data).

Together, these data show a telomerase-dependent accumulation of cellular senescence-associated markers with ageing in the zebrafish brain (increased SA-β-Gal, DNA damage response, and *p16* expression, and up-regulation of some putative SASP factors), suggesting that Tert has a protective effect against accumulation of senescence in the brain. Interestingly, accumulation of senescence-associated markers was detected in both proliferative and non-proliferative regions of the brain (**Supp Fig 4B**). This could suggest that there are other types of telomerase-dependent senescence accumulating with ageing in addition to classic replicative senescence. One hypothesis is that non-canonical, i.e. telomere-length independent functions of telomerase may also have a protective role in the brain (Segal-Bendirdjian & Geli, 2019).

2.3 Tert-dependent inflammation with ageing in the zebrafish brain

Within its non-canonical functions, the Tert component of telomerase (i.e telomerase reverse transcriptase) has been reported to play a role in transcriptional regulation of inflammatory-associated genes (Choi et al., 2008; Ghosh et al., 2012). Moreover, telomere dysfunction has been shown to contribute to a dysfunctional inflammatory environment in the zebrafish model (Lex et al., 2020). Accordingly, our RNA sequencing data identifies immune dysregulation as one of the top 3 telomerase-dependent transcriptional hallmarks of ageing and we wanted to investigate this further at tissue level. We started by assessing overall levels of inflammation in the zebrafish brain, in young and old animals, in the presence and absence of telomerase. Because there is no agreed specific microglia marker in zebrafish, we combined the tg(*mpeg*1.1-mCherry-caax) transgenic zebrafish line (Ellett, Pase, Hayman, Andrianopoulos, & Lieschke, 2011), which has been shown to identify macrophages and microglia in the brain (Oosterhof et al., 2017), with a well-described pan-

immune cell marker (L-plastin or LCP-1) (Cvejic et al., 2008; Mathias et al., 2010; Redd, Kelly,

Dunn, Way, & Martin, 2006; Van Houcke et al., 2017) and use immunofluorescence (IF) to

quantify immune cell populations in the brain (Martins, Ellis, MacDonald, Richardson, &

Henriques, 2019) as a readout for inflammation. Using these two markers allowed us to

identify macrophages/microglia as L-plastin$^+$*mpeg*$^+$ cells, and distinguish them from any

other immune cells, which would be L-plastin$^+$*mpeg*$^-$ (**Fig 3A).** Our data show that there is an

increased number of plastin$^+$*mpeg*$^+$ cells (**Fig 3A1**) but not of L-plastin$^+$*mpeg*$^-$ cells (**Fig 3A2**)

with ageing, in most of the zebrafish brain (WT 5 months *VS* WT 30 months) (**Supp Fig 6**),

suggesting a macrophage/microglia specific increase. The macrophage/microglia increase in

the brain was accelerated in the absence of telomerase already at the early age of 5 months

(WT 5 months *VS tert*$^{-/-}$ 5 months), suggesting that neuroinflammation is as a telomerase-

dependent ageing phenotype (**Fig 3A1**).

To understand whether macrophage/microglia increase with ageing was associated with

immune activation, we used a chitotriosidase activity assay. Chitotriosidase is an enzyme

expressed by activated macrophages and other phagocytic cells and can therefore be used

as a biomarker of microglia/macrophage activation, including in zebrafish (Keatinge et al.,

2015). Our data show that there are increased levels of chitotriosidase in the naturally aged

brain (WT 4-6 *VS* 14-16 *VS* 30 months) and that this is accelerated in the absence of

telomerase from the age of 14-16 months, suggesting that neuronal immune activation is a

telomerase-dependent phenotype of ageing (**Fig. 3B**).

Accumulation of cellular senescence is thought to contribute to chronic inflammation in

ageing, often termed "inflammageing" (Deleidi, Jäggle, & Rubino, 2015; Franceschi &

Campisi, 2014; M, Selman;, Lene, & Bellantuono, 2016). On one hand, in some contexts,

immune cells have been shown to be recruited to sites of cellular senescence (Demaria et

al., 2014; Krizhanovsky et al., 2008). On the other hand, senescent cells have been shown to

release pro-inflammatory factors (part of SASP), which can contribute to paracrine

senescence (Acosta et al., 2013) as well as to tissue inflammation (Franceschi & Campisi,

2014). We therefore set out to determine how the increased levels of senescence-associated

markers we detected in the zebrafish brain correlated with immune cell numbers. To do this,

we performed IF staining with the pan-immune cell marker L-plastin on SA-β-Gal-stained

Figure 3. Telomerase (Tert)-dependent increase in inflammation with ageing in the zebrafish brain. (A)

Representative images of mpeg and L-plastin staining in the diencephalon of adult zebrafish. The yellow arrow

highlights a Lplastin+; mpeg- cell. (A1) Quantifications of L-plastin-positive; mpeg-positive cells (orange; from

red and green co-localisation) and (A2) L-plastin-positive; mpeg-negative cells (red) in the whole zebrafish brain

show an increased number of mpeg$^+$ cells with natural ageing, and this is accelerated in the absence of

telomerase (in $tert^{-/-}$ at 5 months of age). N=-6 per group. (B) Schematic figure of the chitotriosidase assay (left).

Chitotriosidase activity increases with natural ageing at 30 months of age, and this is accelerated in the $tert^{-/-}$

at 14-16 months (right). (C) Representative staining of SA-β-Gal (blue) and L-plastin (white) shows that most of

the L-plastin-positive cells do not co-localise with the SA-β-Gal staining (left). Staining quantification shows a

correlation between increased number of immune cells and increased expression of SA-B-Gal with ageing in

both WT and $tert^{-/-}$ (right). (A1, A2, B) Each dot represents one animal. (A1, A2, B, C) Bar errors represent the

SEM. * <0.05; ** <0.01; *** <0.001.

brain tissue sections. Interestingly, even though we cannot exclude that some of these

immune cells may also be SA-β-Gal$^+$ themselves, most of the immune cells appear as swarms

surrounding SA-β-Gal$^+$ areas (**Fig 3C**). Moreover, there is a statistically significant positive

correlation between % of SA-β-Gal$^+$ areas and % of immune cells with ageing in the WT

zebrafish brain (R^2=0.99), p(two-tailed) =0.0313. Importantly, this correlation seems to be

lost in the absence of telomerase ($tert^{-/-}$), where, for example, at c.5 months, the % SA-β-Gal$^+$

areas is more than double that of the WT at the same age and yet the % of immune cells is

the same. This could suggest that the inflammatory response to raised levels of senescence-

associated markers is deficient in the absence of telomerase. Another, not mutually exclusive

possibility is that $tert^{-/-}$ SA-β-Gal$^+$ cells are more immune evasive.

2.4 Tert-dependent blood-brain barrier dysfunction with ageing

Because our data identified a telomerase-dependent increased number of immune cells in

the zebrafish brain with ageing (**Fig 3A**), which was not explained by increased immune cell

proliferation (**Supp Fig 6**), we hypothesised that there could be immune infiltration from the

periphery. Immune cells should not cross the Blood Brain Barrier (BBB) and invade the brain

in a healthy condition unless the BBB is compromised (reviewed in (Muldoon et al., 2013)).

We therefore sought out to test BBB integrity in young and old zebrafish, in the presence

and absence of telomerase ($tert^{-/-}$).

It has been described that zebrafish have a functional BBB from the 10th day post fertilisation

(dpf) (Fleming, Diekmann, & Goldsmith, 2013), after which molecules such as horseradish

peroxidase (HRP, 44,000Da), evans blue (961Da), and sodium fluorescein (376Da) do not

cross this barrier (Fleming et al., 2013; Jeong et al., 2008). BBB permeability has been described to increase with ageing, becoming "leaky" (Pakulski, Drobnik, & Millo, 2000) and is associated with neuropathology and neuroinflammation in humans (Andjelkovic et al., 2023; Knox, Aburto, Clarke, Cryan, & O'Driscoll, 2022; Sweeney, Zhao, Montagne, Nelson, & Zlokovic, 2019), including in Alzheimer's disease (Bruno et al., 2024). To test whether zebrafish BBB becomes leaky with ageing and whether telomerase plays a role in it, we set up a BBB permeability assay in both WT and *tert*[-/-] at similar ages where increased immune activation (chitotriosidase activity) was identified in the brain. To do this, we performed intraperitoneal injections (IP) of an infrared fluorescent dye IRDye 680RD Dextran diluted in saline (Hanks buffered solution-HBSS). We confirmed this dye's suitability for assessing BBB by performing a pilot study comparing dispersion of conjugated molecules of different sizes throughout the body at different time points and determined that 4kDa Dextran conjugated with fluorescent dye was optimal for assessment of tissue dispersal 3 hrs post-injection (**Supp Fig 7**). Three hours after injection, fish were culled, and dissected brain fluorescence was measured (using a Li-cor Odyssey imaging system) to assess BBB permeability. Our data show increased 4kDa dextran-FITC-associated fluorescence in the naturally aged zebrafish brain at >30 months, when compared to 12 months brains, normalised to age-matched vehicle (HBSS) injected animals. In the absence of telomerase (*tert*[-/-]), however, there is already a significant increase in dextran-associated fluorescence at 12 months when compared to WT, suggesting that BBB "leakiness" increase with old age in zebrafish is a telomerase-dependent phenotype and could be a means for immune cell infiltration from the periphery (Deverman & Patterson, 2009).

Figure 4. Telomerase-dependent increased blood brain barrier permeability with ageing in the zebrafish brain. (A) Schematic figure of the blood brain barrier permeability assay. Dextran-FITC 4kDa or vehicle control

(HBSS) were injected by intraperitoneal injection and 3 hours later the fish were imaged live in the Odyssey® CLx (Li-cor) to confirm the presence of Dextran-FITC in the whole body. (A1) Then the brains were dissected and imaged to assess the presence of Dextran-FITC. Quantification of the green fluorescence in the fish brains shows higher FITC expression in old WT (>30 months) compared to middle aged ones (12 months). This increased presence of dextran-FITC in *tert*[-/-] is already observed at 12 months, overall suggesting that there is increased brain permeability with natural ageing and that this is anticipated in the absence of telomerase. N=9-20 per group in the tested animals; N=3-6 per group in the vehicle control animals. Bar errors represent the SEM. Statistics: unpaired t-tests. * $p<0.05$; ** $p<0.01$; *** $p<0.001$.

2.5 Tert-dependent behavioural changes with ageing

It is known that inflammation can influence behaviour. For example, lipopolysaccharide (LPS)-induced peripheral inflammation has been shown to lead to behavioural alterations in rodents, in a dose-dependent manner, including changes in anxiety-like behaviour (Bassi et al., 2012; Tarr, Chen, Wang, Sheridan, & Quan, 2012). Given that we observed a telomerase-dependent increase in neuroinflammation with ageing, we set out to test whether it associated with any alterations in behaviour. To do this, we assessed anxiety-like behaviour using a novel object test (Maximino et al., 2010; Pegram et al., 1991; Wright, Nakamichi, Krause, & Butlin, 2006), in WT and *tert*[-/-] zebrafish at young and old ages (**Fig 5A**). The novel object test showed that, with ageing, WT fish tend to spend more time exploring the area near the object in the centre of the tank (higher number of entries in the area next to the novel object in 14-17 *versus* >30m), and this behaviour is already detected in *tert*[-/-] at the earlier age of 14-17 months (**Fig 5B1**). Increased exploration of the novel object suggests decreased anxiety-like behaviour with natural ageing, which our data suggest is a telomerase-dependent phenotype of old age.

Figure 5: Telomerase-dependent behaviour alterations with ageing in zebrafish. (A) Representative images of the setup used during the novel object test. (B) Heatmap representing young vs old WT and *tert*[-/-] fish locomotion in the tank throughout the NO test. (B1) Quantifications show that old WT (>30 months) spend

longer periods of time near the novel object compared to younger siblings, and that this behaviour is already

observed in *tert*[-/-] at 14-17 months. N=7-9. Bar errors represent the SEM. * p<0.05; ** p<0.01; *** p<0.001

3 DISCUSSION

Brain ageing and related neurodegenerative diseases have dramatic consequences in quality

of life and medical care costs. Understanding the fundamental mechanisms underlying brain

ageing has the potential to identify therapeutic targets for multiple age-associated brain

diseases, which hold the promise of ameliorating quality of life in the elderly and extending

healthspan. Decreased telomerase expression, telomere shortening, senescence-associated

markers and inflammation have all been independently observed in the ageing brain and

associated with human neurodegeneration. However, causality between limited telomerase

expression and increased brain senescence and neuro-inflammation in the natural ageing

setting is yet to be established. Here, we address these questions using the zebrafish as an

ageing model which, akin to humans, displays premature ageing and death in the absence of

telomerase and where telomere shortening is a driver of cellular senescence. Our broad

transcriptomic data and enrichment analysis of whole zebrafish brains throughout the life-

course identified dysregulation of gene expression, stress response and dysregulation of

immune genes as the top 3 significantly enriched transcriptional hallmarks of old age (Frenk

& Houseley, 2018) in WT and these were all accelerated in the absence of telomerase (*tert*[-/-]

). We decided to focus on senescence as a stress response and inflammation as proxy for

dysregulated immune genes, as these are amenable for further cellular and molecular

analysis at tissue level.

3.1 Tert dependent increase in senescence-associated markers with ageing in the zebrafish

brain

Proof of principle experiments have shown that cellular senescence is a contributor to age-

associated dysfunction in mice, and that selective killing of putative senescent, p16

expressing cells significantly ameliorated age-associated dysfunction, including cognitive

decline (Baker et al., 2016; Bussian et al., 2018). Further research on cellular senescence and

SASP have identified senolytic molecules capable removing putative senescent cells, many

of which are in current clinical trials, recently reviewed in (Huang et al., 2022). Importantly,

it is now appreciated that SASP itself plays key roles in the pathogenesis of cellular

senescence and SASP factors have been described to potentially contribute to the "spreading" of senescence phenotypes to nearby cells, i.e, "paracrine senescence"(Acosta et al., 2013). Therefore, senomorphic molecules (i.e., SASP modulating molecules) represent a promising therapeutic strategy for age-related diseases. One example of such senomorphic compounds is ruxolitinib, a JAK inhibitor shown to reduce putative SASP-associated inflammation and alleviate frailty in old mice (Xu et al., 2015). In fact, the SASP has been associated with several age-associated phenotypes including neurodegeneration (Wang et al., 2024).

The transcriptional changes of old age (WT young *VS* old DEGs) that we mapped to cellular senescence (enriched KEGG profile) included well-known cell signalling and cell-cycle genes such *foxo1a*, *pik3*, *mapk1*, *mapk11*, multiple cyclins (*ccnd2a, ccnb3, ccnb1, ccnd1, ccna1*), cyclin dependent kinases (*cdk1, cdk2, cdk6*), *chek1*, the transcription factor *e2f1*, autophagy regulator sqstm1 amongst others (see supplementary data source files). We also identified a number of yet to be described zebrafish genes associated with senescence to be investigated in future work, namely ZGC:56235, ZGC:171775, ZFP36L1A and ZGC:114130. When searching specifically for handpicked senescence-associated genes, namely *p21* and *p16 (Huang et al., 2022)*, we identified *p16* as significantly up-regulated in WT aged brains (WT 2m *VS* 30m, log2 fold 0.983 and p adjusted of 0.00025), which was exacerbated in the absence of telomerase (tert 2m *VS* 22 months, log2 fold 1.09 and p adjusted of 0.000000002). In contrast, *p21* gene expression was not significantly changed either in WT or *tert-/-* aged brains.

Cellular senescence can be induced by different factors causing cellular impairment, including DNA damage, telomere shortening or dysfunction, oncogene activation or loss of tumour suppressor functions, epigenetic changes, and organelle damage, recently reviewed here (Huang et al., 2022). Notably, different types of cellular senescence will express different senescence-associated markers and SASP factors (Basisty et al., 2020; Hernandez-Segura et al., 2018). For example, whereas DNA damage response (DDR)-induced cellular senescence is often characterised by p53-p21 upregulation and pathway activation (Rossiello, Herbig, Longhese, Fumagalli, & d'Adda di Fagagna, 2014), epigenetic (Petrova, Velichko, Razin, & Kantidze, 2016) or mitochondria dysfunction-driven cellular senescence usually upregulates

p16. Moreover, p21 expression has been associated with more initial phases of cellular

senescence, whereas p16 seems to be involved in maintaining the senescence phenotype

(Dulic et al., 2000).

The most used cellular senescence-associated markers are not only the expression of p16 and

p21 genes but also accumulation of senescence-associated β-Galactosidase (SA- β-Gal)

(Huang et al., 2022). However, there is no single marker that can be exclusively and 100

percent relied upon to identify cellular senescence, as this likely represents a gradient of

phenotypes, dependent on cell type and senescence stimuli, as mentioned above (Basisty et

al., 2020; Hernandez-Segura et al., 2018). Acknowledging this limitation, we decided to

investigate the expression of several senescence-associated markers at tissue level, in aged

WT and *tert*$^{-/-}$ zebrafish brains. We identified an accumulation of SA-β-Gal in WT zebrafish

brains with ageing, which was accelerated in the absence of telomerase (*tert*$^{-/-}$)*. This was

accompanied by an increase in *p16* gene expression as assessed by RNA *in situ* hybridisation.

Consistent with our RNA sequencing data, we did not detect any changes in *p21* gene

expression as assessed by RNA *in situ* hybridisation in either WT or *tert*$^{-/-}$ aged brains. We

detected only a mild increase in the expression of the DDR molecule γH2AX with ageing.

Together with the fact that there were no significant changes observed in proliferation,

despite a decrease in telomere length in aged and *tert*$^{-/-}$ brains, and the identification of SA-

β-Gal in both known proliferative and non-proliferative areas of the zebrafish brain, our data

suggest that there is a largely DDR-independent, *p16*-dependent increase in putative

senescent cells in the aged zebrafish brain, which is accelerated in the absence of telomerase

(*tert*$^{-/-}$). Of note, even though telomere length is shorter in the *tert*$^{-/-}$ when compared to WT

siblings, significant even at the early age of 5 months, we did not detect significant further

telomere shortening from 5 to 12 months old in the *tert*$^{-/-}$. This is consistent with very low

levels of proliferation in the brain, but we cannot exclude the possibility that alternative

lengthening of telomeres (ALT) may be present, counteracting telomere shortening (Idilli et

al., 2020; Recagni, Bidzinska, Zaffaroni, & Folini, 2020).

Consistent with DDR-independent the SASP factors we identified in aged zebrafish brains had

more in common with oncogene- than genotoxic stress-induced senescence, according to

saspatlas.com, which has data from irradiation-induced (IR) genotoxic stress and oncogene

(RAS)-induced senescence in epithelial and fibroblast cells, *in vitro* (Basisty et al., 2020).

Which cells are undergoing senescence in this context remains to be determined and will

require future work. It has been reported that several cell populations in the CNS can undergo

senescence, depending on the context. Neural stem cells (NSCs) and neural progenitor cells

(NPCs) have been shown to undergo stress-induced senescence when exposed to ionized

radiation, characterised by increased SA-β-Gal (Mason, 2012) and p16 expression (Le et al.,

2018). Microglia, the specialised immune cells of the CNS, seem to undergo replicative

senescence, *in vitro* and *in vivo* (Flanary & Streit, 2004; Raj et al., 2015). *In vivo,* this was

demonstrated in a progeroid mice model where microglia display short telomeres and

modest increased expression of p21 and SASP-factors (Raj et al., 2015). Astrocytes, the most

abundant glial cell in the CNS, also seem to undergo stress-induced senescence, in humans

(Crowe et al., 2016; Mason, 2012), exhibiting high levels of SASP factors when exposed to

irradiation (Mason, 2012), as well as SA-β-gal activity, increased expression of DNA damage

checkpoints (Bitto et al., 2010) and SASP-factors (Crowe et al., 2016), when exposed to H_2O_2.

However, *in vitro* studies have reported controversial results regarding whether these cells

also undergo replicative senescence. Whilst Flanary & Streit (2004) did not identify replicative

senescence in rat astrocytes (Flanary & Streit, 2004), another two studies reported that

astrocytes from humans (Evans et al., 2003), mice and rats undergo replicative senescence

(Evans et al., 2003). Oligodendrocytes are the glial cells responsible for producing the sheath

of myelin around the neuronal axons. Post-mortem studies have reported increased DNA

damage in the white matter, together with oxidative damage and SA-β-Gal in this cell

population (Nasrabady et al., 2018). Importantly, neurons, considered largely non-

proliferative cells, have also been reported to undergo senescence. Le et al. showed increased

expression of p16 in the hippocampus, sub-ventricular zone (SVZ) and cortex of mice, after

irradiation (Le et al., 2018). Similarly, cortical neurons and Purkinje neurons of aged mice

present DNA damage, SA-β-Gal and SASP factors. Myenteric ganglia also undergo senescence

in the aged mice, presenting DNA damage, together with activated P38MAPK p38 mitogen-

activated protein kinases (p38MAPK), SA-β-Gal and ROS (Jurk et al., 2012). Our data show that

the majority of SA-β–Gal⁺ cells are not of immune origin (as assessed by L-plastin expression) and are too abundant and widely distributed to be restricted to neural stem or progenitor stem cells and are therefore likely to be a mixture of cell types. Future work will be required to investigate this further.

3.2 Tert-dependent neuroinflammation with ageing

The Tert component of telomerase is known to play telomere-independent roles, including in the modulation of mitochondria function and inflammation. Because we did not identify significant changes in mitochondria complex I or II function in WT aged zebrafish brains, and considering our data described so far, our current hypothesis is that non-canonical functions of telomerase involved in inflammation may be contributing to the accumulation of cellular senescence associated markers in the zebrafish brain, which we aim to test in future work.

Accordingly, immune dysregulation was one of the top 3 enriched ageing associated transcriptomic hallmarks in both WT and *tert⁻ᐟ⁻* aged zebrafish brains. Furthermore, at tissue level, we identified a telomerase-dependent increase in the number of macrophages/microglia with ageing in the zebrafish brain. This agrees with previous literature, since neuro-inflammation in humans is known to involve increased number of microglia as well as increased immune activation (Damani et al., 2011; Crowe et al., 2016). Similarly, two recent studies in zebrafish reported an increased number of microglia/ macrophages in the aged brain, more specifically in the telencephalon and optic nerve (Van Houcke et al., 2017; Zambusi et al., 2020). In the telencephalon, microglia were also shown to display an ameboid shape in the aged fish, suggesting increased activation (Zambusi et al., 2020). Furthermore, chitotriosidase activity has been proposed as a putative marker of microglial/ macrophage activation in mammals. In humans, increased levels of chitotriosidase activity have been reported to increase with normal ageing (in serum) (Ramanathan et al., 2013) and exacerbated in pathologies associated with chronic inflammation, including in ALS (Steinacker et al., 2018). Accordingly, our data show that Tert-dependent increased inflammation in the aged zebrafish brain was accompanied by increased chitotriosidase activity, which is consistent with increased immune activation.

Whether this increased neuroinflammation is detrimental remains to be determined, and it

is not entirely clear in human neurodegeneration. Neuroinflammation is associated with

neurodegeneration and disease in humans (Gaikwad et al., 2024; Righi et al., 2024; Samant,

Standaert, & Harms, 2024), but causality is not clear. Our data show that SA-β–Gal$^+$, putative

senescent cells are surrounded by swarms of cells of immune origin (L-plastin$^+$) and there is a

significant positive correlation between increased β–Gal$^+$ areas and increased % of immune

cells in the brain. However, this is not the case in the absence of telomerase. For example, in

young tert$^{-/-}$ fish (c.5 months), the % SA-β-Gal$^+$ area is more than double that of the WT at the

same age and yet the % of immune cells is the same. This could suggest that the inflammatory

response to raised levels of senescence-associated markers is deficient in the absence of

telomerase. Our previous work identified unexpected roles for telomerase in macrophage

function, affecting immune activation and phagocytic ability in the zebrafish gut (Ellis et al.,

2022). Whether telomerase could also impact on immune recruitment and clearance of

senescent cells remains to be determined. Another, not mutually exclusive possibility is that

tert$^{-/-}$ SA-β-Gal$^+$ cells are more immune evasive and mechanisms similar to cancer evasion

from immune surveillance (Pereira et al., 2019) may be at play. Future work will be required

to test these hypotheses.

Finally, it is not clear whether neuroinflammation in the aged zebrafish brain is contributing

to the senescence phenotypes via paracrine senescence mechanisms (Acosta et al., 2013), or

whether immune cells are actually trying to clear senescent cells (Lujambio, 2016).

Nevertheless, the fact that in the absence of telomerase we have double the amount of SA-

β-Gal$^+$ area when compared to WT, yet the same % of immune cells as in the WT, suggests

that it is not inflammation that is driving it, at least in the tert$^{-/-}$ context. Experimental work

manipulating senescence and inflammation will be required to determine this. However,

modulating senescence and inflammation separately will present its own challenges since

many of the senolytic drugs currently in use also have senescence-independent anti-

inflammatory effects (Hernandez-Silva et al., 2022).

3.3 Tert-dependent BBB dysfunction with ageing

Our data show that neuroinflammation in the aged zebrafish brain was not explained by

increased levels of proliferation. We therefore hypothesised that perhaps there is increased

immune infiltration from the periphery due to blood brain barrier permeability or "leakiness"

with ageing, as has been described before with human ageing (Pakulski et al., 2000). The BBB

constitutes a protective barrier between the blood and the CNS, avoiding the entrance of

foreign molecules into the brain, including the entrance of peripheral immune cells.

Increased permeability can therefore lead to the infiltration of peripheral immune cells in

the brain (Deverman & Patterson, 2009). Moreover, the choroid plexus holds some

populations of resident immune cells, mostly including T-cells and monocytes but also B-

cells, dendritic cells and natural-killer cells (Meeker, Williams, Killebrew, & Hudson, 2012).

These resident immune cells have been shown to be able to invade the brain parenchyma in

response to inflammation (Strominger et al., 2018). Moreover, it has been described that

auto-reactive T-cells (Coder, Wang, Ruan, & Su, 2015) are capable of penetrating the brain

with ageing, both through the bloodstream and lymphatic vessels (Louveau et al., 2015).

Additionally, circumventricular organs constitute another mechanism by which peripheral

immune cells could invade the brain. These are brain structures that lack the classical BBB

and display high permeability (Ganong, 2000), constituting vulnerable regions to the

infiltration of large molecules and possibly, cells, into the brain parenchyma. Therefore,

increased number of immune cells in the aged brain is likely to be associated with the

following mechanisms: (1) increased proliferation of immune cells, particularly of microglia;

(2) recruitment of immune cells from the periphery, through the BBB; (3) crossing of resident

immune cells from the cerebrospinal fluid (choroid plexus and meninges) into the brain

parenchyma; and/or (4) infiltration of peripheral immune cells, through the

circumventricular organs.

Our data show that there is a telomerase-dependent increase in BBB permeability, therefore

supporting the second mechanism. However, cell tracking *in vivo* experiments would be

required to directly show that there is indeed immune infiltration from the periphery.

Mechanistically, endothelial cells and pericytes, two major components of the blood brain

barrier (BBB), have been reported to undergo senescence with ageing in mice, *in vitro* and *in*

vivo, expressing elevated levels of p16, p21 and SA-β-Gal (Yamazaki et al., 2016). Whether

BBB permeability increase with ageing in the zebrafish brain is due to endothelial senescence

and, if so, whether this is cell autonomous or paracrine-induced senescence remains to be

tested.

3.4 Tert dependent changes in anxiety like behaviour with ageing

Given that we observed a telomerase-dependent increase in neuroinflammation with ageing, and anxiety-like behaviour has previously been reported to increase in response to inflammatory stimuli (Bassi et al., 2012; Tarr et al., 2012), we sought to test whether there are telomerase-dependent behavioural changes with ageing in zebrafish. Surprisingly, against our initial hypothesis that increased inflammation would lead to increase anxiety-like behaviour, our data revealed a decrease in anxiety-like behaviour with ageing, which was accelerated in the absence of telomerase. Both age and levels of inflammation are likely to play a role in anxiety-like behaviour and interact with one another.

Anxiety-like behaviour has been reported to increase with ageing in rodents (Nolte, Nolte, & Yan, 2019) and zebrafish (Evans, Torres-Perez, Miletto Petrazzini, Riley, & Brennan, 2021; Kacprzak et al., 2017). Nonetheless, contradictory results have been observed, and this is likely to be associated with differences in animal strains used and/or differences in behaviours assessed. For example, a study in zebrafish has shown that differences in strains and specific measures assessed lead to different results in exploratory behaviour (Hudock & Kenney, 2024).

Similarly, a study in rodents (Bassi et al., 2012) has reported that anxiety-like behaviours were only detected in some of the tests performed, suggesting that the results may be variable depending on the behaviour test used and/or measures assessed. Another variable likely to influence the results is the animal age assessed, as age-associated behavioural alteration may not be linear throughout lifespan. Indeed, Ferreira et al (2023) (A. C. Ferreira, Sousa, Sousa, & Marques, 2023) observed that although there is an increase in anxiety-like behaviour in mice at 12-months of age when compared to younger mice, this was reduced later in life, at 18 months of age. In zebrafish, anxiety-like behaviour has been shown to increase with ageing, up to 18 (Evans et al., 2021; Kacprzak et al., 2017) and 24 months of age (just over "middle aged" adults) (Vasconcelos, Gordillo-Martinez, Ramos, & Lau, 2023). However, to our knowledge, our study is the first one assessing anxiety-like behaviour in old adults (i.e. >30 months old). In hand with what has been shown in mice, by Ferreira et al

(2023) (A. C. Ferreira et al., 2023), our study seems to suggest that there is a slight increase

in anxiety-like behaviour in adults (14-17 months) and a significant reduction later on in life.

Similarly to what has been reported with ageing, behaviour changes in response to

inflammation do not appear to be linear. Bassi et al 2012 (Bassi et al., 2012) reported a dose-

dependent increase in anxiety-like behaviour in response to an inflammatory stimulus, which

followed an inverted U curve, with slight reduced levels of anxiety-like behaviour in response

to intermediate doses, and increased levels of anxiety-like behaviour in response to higher

doses. The reduced anxiety-like behaviour we observed in old fish may be explained by

chronic, low levels of inflammation characteristic of ageing, in opposition to high

inflammatory levels of a response to an acute stimulus.

CONCLUSIONS

Our work shows for the first time that the zebrafish brain undergoes key telomerase (Tert)-

dependent age-associated changes at transcriptional, cellular, tissue and functional levels.

We show that Tert-dependent transcriptomic changes associated with dysregulation of gene

expression, stress response and dysregulation of immune genes are accompanied by

accelerated accumulation of senescence-associated markers and inflammation in the aged

brain. *In vivo*, these changes correlated with increased blood-brain barrier permeability and

altered cognitive behaviour. Together, our work suggests that telomerase has a protective

role in the zebrafish brain against the accumulation of senescence and neuro-inflammation

and is required for blood brain barrier integrity.

MATERIALS AND METHODS

1. Animal husbandry, ethics, genotypes and ages

Zebrafish were maintained at the standard conditions of 27-28ºC, in a 14:10 hour light-dark

cycle, and fed twice a day with Artemia (live rotifers) and Sparus (dry food). All the

experiments were performed in the University of Sheffield. All animal work was approved by

local animal review boards, including the Local Ethical Review Committee at the University

of Sheffield (performed according to the protocols of Project Licence 70/8681). Two strains

of adult zebrafish (Danio rerio) were used for these studies: wild-type (WT; AB strain) and

tert$^{-/}$ $^{(tertAB/hu3430)}$. Wild-type fish were obtained from the Zebrafish International Resource Center (ZIRC). The telomerase mutant line tert $^{AB/hu3430}$ was generated by N-ethyl-nitrosourea mutagenesis (Utrecht University, Netherlands (Wienholds et al., 2003)). It has a T→A point mutation in the tert gene and is available at the ZFIN repository, ZFIN ID: ZDB-GENO-100412–50, from ZIRC. The fish used in this study are direct descendants of the ones used in the first studies describing this line(Carneiro et al., 2016; Henriques et al., 2013), by which point it had been subsequently outcrossed five times with WT AB for clearing of potential background mutations derived from the random ENU mutagenesis from which this line was originated.

The tert $^{hu3430/hu3430}$ homozygous mutant is referred to in the paper as *tert$^{-/-}$* and was obtained by in-crossing our tert heterozygous AB/hu3430 strain, following gold standard guidelines to use this line, as recently reviewed here(Henriques & Ferreira, 2024). Genotyping was performed by PCR of the tert gene as before. To study age-related phenotypes in the zebrafish brain, we used >30 months old fish for what we consider old in WT (in the last 25-30% of their lifespan), and we considered the *tert$^{-/-}$* old fish at the equivalent age (>22 months), which approximately corresponds to the last 25-30% of their lifespan. In specific, 'Old' was defined as the age at which most of the fish present age-associated phenotypes, such as cachexia, loss of body mass and curvature of the spine. These phenotypes develop close to the time of death and are observed at >30 months of age in WT and at >22 months in *tert$^{-/-}$* (Madalena C Carneiro et al., 2016; Henriques et al., 2013).

Note on Zebrafish gene and protein nomenclature

This manuscript followed the agreed convention as described in:

https://zfin.atlassian.net/wiki/spaces/general/pages/1818394635/ZFIN+Zebrafish+Nomenclature+Conventions

In summary:

Gene nomenclature

Genes should be named after the mammalian orthologue whenever possible. When mammalian orthologues are known, the same name and abbreviation should be used, except all letters are italicized and lower case. Members of a gene family are sequentially numbered.

Examples:

Names - engrailed 1a, engrailed 2b

Symbols - eng1a, eng2b

Protein nomenclature

The protein symbol is the same as the gene symbol, but non-italic and the first letter is

uppercase.

Note the differences between zebrafish and mammalian naming conventions:

species / gene / protein

zebrafish /*shha*/ Shha

human / *SHH* / SHH

mouse / *Shh* / SHH

2. Tissue fixation

<u>Paraffin-embedded sections</u>: Fish were culled by immersion in concentrated tricaine (MS-

222). After terminal anaesthesia, the animals were fixed by immersion in 50 ml of 4%

paraformaldehyde (PFA) buffered at pH 7.0 (VWR International, Radnor, PA, USA) at 4ºC for

48h-72h, and decalcified in 50 ml of 0.5M ethylenediaminetetraacetic acid (EDTA), pH 8, for

additional 48-72h. Then, the whole fish were paraffin-embedded (in formalin I for 10min,

formalin II for 50min, ethanol 50% for 1h, ethanol 70% for 1h, ethanol 95% for 1h 30min,

ethanol 100% for 2h, ethanol 100% for 2h 30min, 50%:50% of ethanol 100% : Xilol for 1h

30min, xylene I for 3h, Xylene II for 3h, paraffin I for 3h, paraffin II for 4.30h), longitudinally

sliced (sagittal slices, 3 µm thick, using the Leica TP 1020), (3 µm thick, using the Leica TP

1020), and stored at room temperature (RT). To do this, diluted ethanol absolute (VWR

International) and paraffin histosec pastilles (Merck & Co, Kenilworth, NJ, USA) were used.

<u>Cryosections</u>: Fish were culled by immersion in concentrated MS-222, followed by

immediate decapitation (Schedule 1 method). The brain was dissected, transferred to and

rinsed in cold L15 (Thermo Fisher Scientific, Waltham, MA, USA). Afterwards, the brain was

fixed in 4% PFA (SIGMA-ALDRICH, St. Louis, MO, USA) overnight at 4°C, washed in 3ml cold

Phosphate Buffered Saline (PBS, 1x), and immersed in 3ml of 30% sucrose in PBS overnight

at 4ºC for cryoprotection. Before embedding in mounting media (O.C.T. compound, VWR

International), the tissues were immersed in 50%:50% solution of 30% sucrose:O.C.T. for

30mins at 4°C. Finally, the tissue was mounted in a base mould (Electron Microscopy

Sciences, Hatfield, PA, USA), with O.C.T., snap frozen, and stored at -20°C. Prior to use, the

tissues were sliced in 13μm-thick coronal sections using a cryostat.

3. Immunofluorescence (IF)

Paraffin sections with whole fish slices were dewaxed and hydrated through the following

sequence of washes: histoclear (Scientific Laboratory Supplies, Wilford, Nottingham, UK, 2x5

min), 100% ethanol (Thermo Fisher Scientific, 2x5 min), 90% ethanol (5 min), 70% ethanol (5

min), and distilled water (2x5 min). Cryosections were hydrated in PBS for 10 min before

proceeding. After antigen retrieval in 0.01M citrate buffer at pH 6.0 (sub-boiling at 800 W in

a microwave for 10 min), the tissue sections were permeabilized in PBS 0.5% Triton X-100

(for 10 min) and blocked in 3% bovine serum albumin (BSA), 5% Goat Serum in 0.3% Tween-

20 in PBS (for 1 h). Then, primary antibody incubation (overnight at 4ºC) was followed by

washes in PBS 0.1% Tween-20 and secondary antibody incubation (1 h at RT or overnight at

4ºC). Finally, the tissues were incubated in DAPI for nuclear staining (Thermo Fisher

Scientific), 1:10000 dilution in PBS from a 10mg/ml stock, for 10 min; washed in PBS 1x,

mounted with vectashield (an antifade mounting medium, from Vector Laboratories,

Burlingame, CA, USA), covered with coverslip (Scientific Laboratory Supplies), and sealed

with clear nail varnish. For details on the primary and secondary antibodies refer to Table 1

and Table 2, respectively.

4. Telomere-fluorescence *in situ* hybridization (telo-FISH)

Telo-FISH was performed after IF. Briefly, after incubation with the secondary antibody and

respective washes, the tissues were fixed in 4% PFA (SIGMA-ALDRICH) for 20min. Washes in

PBS, 3x 10min were followed by a dehydration step consisting of washes with ice cold 70%,

90% and 100% ethanol (Thermo Fisher Scientific, 3min each). After this, the slides were left

to dry for 1h, 10μl of hybridisation solution was pipetted on top of each tissue section and

the slides were covered with a coverslip. Hybridisation solution (for a total volume of 100μl):

8.56μl MgCl2 buffer pH 7.0, and 82mM sodium hydrogen phosphate (SIGMA-ALDRICH), 70μl

of deionised formamide (Thermo Fisher Scientific), 1μl of 1mM Tris, pH 7.5 (SIGMA-

ALDRICH), 5μl of 1x blocking buffer (10x blocking reagent (Merck & Co.) diluted to 1x in

maleic acid (SIGMA-ALDRICH)), 2μl of PNA probe (Telo-Cy3, zebrafish, 568; Cambridge

Research Biochemicals, Cleveland, UK), 2µl of Cent-Fam probe (488; Eurogentec, Liège,

Belgium), and 11.44µl of dH2O.

The slides were incubated for 10min at 80°C for DNA denaturation, followed by incubation

for 2h at RT, in the dark, to allow the probe to hybridise with the complementary DNA. After

the incubation period, the coverslips were gently removed and the slides were washed for

10min in a solution made of 70% formamide (SIGMA-ALDRICH), 10% 2x saline-sodium citrate

(SSC, SIGMA-ALDRICH) and 20% dH2O. This was followed by two more washes (10min each),

in 2x SSC. SSC was diluted from a 20x stock solution consisting of 88.3g of Tri- sodium citrate

dihydrate (SIGMA-ALDRICH) and 175.3g of NaCl (SIGMA-ALDRICH), diluted in dH2O to a total

volume of 1L, adjusted to pH of 7.0 and autoclaved. Finally, the slides were washed in PBS,

incubated in DAPI (Thermo Fisher Scientific) for 10min and mounted as previously described

in section 2.3.2.

5. IF and telo-FISH imaging and quantification

IF and telo-FISH staining was imaged using the DeltaVision microscope at the Wolfson Light

Microscopy Facility (LMF), using a 40x objective.t. An average of 20 images were acquired

from the brain of each animal (approximately 7 images per macro-area).

After acquisition, the images were analysed by manual counting: (1) the number of cells with

more than 5 foci of γH2AX (considered γH2AX+ cells) PCNA-, (2) the number of PCNA+ cells,

(3) the number of L-plastin+, (4) the number of mpeg+ cells, and (5) the total number of cells.

Telo-FISH staining was quantified by the intensity of the mean of the nucleus. The ratio

'telomeres/centromeres' was calculated by dividing the intensity of the mean of telomeres

(568 channel) by the intensity of the mean of centromeres (488 channel), which normalises

the telomeric probe signal by the centromeric probe signal within individual cells.

The analysis was performed on raw images, using the ImageJ Fiji (v. 1.51). Images were then

processed with Abode Illustrator 21.0.2 for display purposes.

1000   Table 1. Primary antibodies used for immunostaining.

Antibody, species, and type	Dilution factor	Catalogue number; company; country
PCNA mouse monoclonal	1:500	NB500-106; Novus Biologicals, Littleton, CO, USA
γH2AX, rabbit polyclonal	1:500	GTX127342; GeneTex, Irvine, CA, USA
L-plastin (Lcp1), rabbit polyclonal	1:200	GTX124420; Novus Biologicals, Littleton, CO, USA
RFP, mouse monoclonal	1:500	GTX82561; GeneTex, Irvine, CA, USA

1001

1002   Table 2. Secondary antibodies used for immunostaining.

Antibody, species, and type	Dilution factor	Catalogue number; company; country
Goat anti-rabbit IgG Alexa Fluor® 488	1:500	A11008; Invitrogen, Carlsbad, CA, USA
Goat anti-rabbit IgG Alexa Fluor® 568	1:500	10032302; Thermo Fisher Scientific, Waltham, MA, USA
Goat anti-mouse IgG Alexa Fluor® 647	1:500	A21235; Thermo Fisher Scientific, Waltham, MA, USA

1003

6. SA-β-Gal

The Senescence-Associated β-Galactosidase Staining Kit (Cell Signalling Technology, Danvers,

MA, USA) was used to perform the SA-β-Gal staining, following manufacturer's instructions.

Whole brain SA-β-Gal staining: Fish were culled with neat Tricaine and decapitated. Whole

heads were fixed in 4ml PFA overnight at 4c with shaking. The following day, whole heads

were washed 3 x for 1h in PBS at RT with slow shaking. Whole brains were then dissected in

fresh PBS and immersed in 2ml in SA-β-Gal staining solution mix, consisting in 1x staining

solution (pH 5.9), 1x solution A, 1x Solution B, 1mg/ml Xgal (SA-β-Gal staining kit #9860 by

Cell Signaling Technology). and incubated at 37°C overnight with slow shaking in the dark.

The following day, brains were washed 3 x for 1h in PBS at RT with slow shaking, and imaged

using a Nikon SMZ1500 Stereomicroscope (2000).

<u>SA-β-Gal staining on cryosections</u>: Briefly, after rinsing the cryosections in 1x PBS for 5 min,

we added 200ul of β-Gal staining solution mix as above, per slide, Then, the slides were

coverslipped and incubated at 37ºC overnight in a humidified chamber box. On the next day,

cryosections were rinsed in 1x PBS (3x1min), and mounted in glycergel (Agilent, Santa Clara,

CA, USA). Slides were then imaged using an Olympus BX60 microscope and analysed using

ImageJ Fiji. Images were then processed with Abode Illustrator 21.0.2 for display purposes.

1021

7. RNA Extraction

Fish were culled by Schedule 1 method (as described before in section 2.2). Zebrafish brains

were dissected, transferred to a microcentrifuge tube with 100 µl of Trizol (Thermo Fisher

Scientific), snap frozen in dry ice and stored at -80ºC until RNA extraction.

For RNA extraction, the tissue was homogenised with a mechanical homogenizer (VWR

International) and 1.5 pestle (Kimble Chase, Vineland, NJ, USA), and an extra 50 µl of fresh

Trizol (Thermo Fisher Scientific) was added. After 5 min of incubation at RT, 30 µl of

chloroform (1:5, VWR International) was added. The samples were then incubated for

further 3 min at RT and centrifuged (at 13,000g for 30 min at 4ºC). The resulting upper

aqueous phase was transferred to a new microcentrifuge tube, and isopropanol (Thermo

Fisher Scientific) was added (approximately 83% of the amount of aqueous phase). After an

incubation of 10 min at RT, the samples were centrifuged (13,000g for 15 min at 4ºC) and

the supernatant was discarded. The pellets were washed in 250 µl of 75% ethanol twice and

then left to air dry. In the end, the pellets were resuspended in 14 µl of nuclease-free water

(VWR International) and stored at -80ºC. All the samples were analysed by Nanodrop 1000

Spectrophotometer to measure the concentration of RNA. Some samples (randomly chosen)

were also analysed in the Bioanalyzer to assess RNA integrity. All the samples included in this

study had an RNA integrity number (RIN) of >9.

1040

1041 **8. RNA sequencing**

1042 RNA-Sequencing was performed at the Genomics and Sequencing facility of Sheffield

1043 Institute for Translational Neuroscience (SITraN). Mature RNA was isolated and fragmented

1044 using the NEBNext Poly(A) mRNA Magnetic Isolation Module (New England Biolabs Inc),

following manufacturer's instructions. The libraries were prepared using the NEBNext Ultra Directional RNA Library Prep Kit (Illumina), following manufacturer's instructions. The NEBNext Multiplex Oligos from Illumina (Index Primers Set 1, New England Biolabs Inc) was used for the library preparation using the following PCR program: 98ºC for 30 sec (1 cycle), 98ºC for 10 sec followed by 65ºC for 75 sec (8 cycles), and 65ºC for 5min. The PCR product (≈50µl) was then purified using 45µl of Oligo dT Beads (Agencourt AMPure XP), followed by three washes with ethanol 80%. After air-drying, the dscDNA was eluted by adding 23µl of 0.1x TE, and the resulting 20µl of the library were collected. Sequencing was run on an Illumina HiScan SQ sequencer, using a high output run and sequencing by synthesis V3 chemistry on a single 100 base pair (bp) run. The sequencing process was monitored in real-time using Sequencing Analysis Viewer v1.8.37 Software, and therefore quality control was performed during and after sequencing. Then, the data was imported into a FASTA file format to perform the analysis.

9. RNA Sequencing analysis

Data processing:

Quality control was performed using MultiQC version 1.9.Cutadapt version 3.0 was used for trimming the first 13 bases from the reads to remove poor quality base pairs in the reads. Read alignment was performed as follows. Single-end reads were aligned against the reference genome Danio_rerio.GRCz11.dna. primary_assembly.fa using STAR. A bespoke alignment index was built using annotation file Danio_rerio.GRCz11.103.gtf and an expected read length of 88bp. Ht-seq count was run in non-stranded mode to obtain per gene read counts.

Differential expression: To identify signatures of ageing, WT brain samples were subjected separately to DESEq2 analysis, comparing the time points 9, 22 and >30 months with the time-point of 2 months. Then, $tert^{-/-}$ samples at 2, 9 and 22 months were compared with WT at 2 months, to identify telomerase-dependent ageing processes.

Enrichment analysis: Pathway enrichment analysis was performed using the Functional Annotation Bioinformatics Microarray Analysis programme (https://david.ncifcrf.gov/). Pathways identified in both Gene Ontology Biological Process (GO-BP) and KEGG databases were included in the analysis. The top 10 GO-BP and KEGG terms were then plotted into a heatmap using graphpad Prism.

<u>Mapping to the hallmarks of ageing:</u> We analysed the recent Frenk & Houseley's paper (Frenk

& Houseley, 2018), which describes the different transcriptional hallmarks of aging. In this

paper, the different hallmarks are defined within specific text sections. For each hallmark,

we collected groups of keywords that could be within GO term names. The complete list of

GO term names was extracted from the file go-basic.obo from the Gene Ontology website

(https://geneontology.org/). We then established lists of GO term names for each

transcriptional hallmark of aging. The mapping table between the hallmarks, the keywords,

and the lists of GO term names is available in the table GO_to_HA-THA_V1.csv.

1085

Zebrafish genes and their associated GO term names were retrieved from the mapping

function Gene2GoReader() from the Python library goatools. We built 3 distinct dictionaries

for each GO category (Biological Process, Molecular Function, and Cellular Component),

where the entrezgene IDs are associated with their GO term names. From the differential

analyses, we obtained different lists of gene symbols that must first be mapped to

entrezgene ID, using the Python library mygene. Each gene ID is associated with a list of GO

term names. To assess a particular transcriptional hallmark of aging to a given gene, we

parsed their associated list of GO term names and mapped them to their corresponding

hallmark of aging.

1095

Depending on the situation, the attribution of the hallmark of ageing is done as follows: 1)

all the GO term names are mapped to the same transcriptional hallmark of ageing, 2) the GO

term names are mapped to several transcriptional hallmarks of ageing, the hallmark with the

highest number of occurrence is chosen. Otherwise, the gene is assigned with an

"ambiguous" hallmark of ageing if 3) none of the GO term names are mapped to any

transcriptional hallmarks of ageing, then the gene is not annotated with a transcriptional

hallmark of aging.

1103

10. Protein extraction and quantification

Fish were culled by Schedule 1 method as previously described. Brain tissue was dissected,

transferred to a microcentrifuge tube and immediately snap-frozen in liquid nitrogen. The

samples were stored at -80°C until protein extraction.

The tissue was homogenised in 100μl of ultra-pure water using a mechanical homogenizer

(VWR International) and RNAse-free disposable pestles (Thermo Fisher Scientific) for

approximately 30sec, on ice. Then, the samples were spun down at 4°C, at 2,000rpm, for

5min. After transferring the supernatant to a new tube, samples were stored at -80°C until

further use.

1113

Protein extracts were quantified by the bicinchoninic acid (BCA) method, using the Pierce™

BCA Protein Assay Kit (Thermo Fisher Scientific), following manufacturer's instructions.

Briefly, a standard curve was prepared with a range of 0.2-2μm of BSA, along with a blank

control (containing ultra-pure water only). A total of 5μl of each protein extract was mixed

with 1ml of BCA, in microcentrifugation tubes, and incubated at 37°C for 10min in a water

bath. After adding an additional 20μl of copper sulphate, the solution was mixed and

incubated at 37°C for 20min. Then, 200μl of each sample (or BSA standard curve) were

transferred to a Corning™ Costar™ Brand 96-Well EIA/RIA Plate (Thermo Fisher Scientific), in

duplicates, and absorbances were read in a Varioskan Plate Reader at a wavelength of

562nm.

1124

1125   **11. Chitotriosidase activity assay**

1126   Chitotriosidase activity was measured using the protocol previously described by Keatinge

1127   et al., (2015). This method allows assessment of the hydrolysis capacity of 4–

1128   methylumbelliferyl–β-D-N, N', N"-triacetylchitotriose (4-MU-chitotrioside), by

1129   chitotriosidase enzyme.

1130

Briefly, a total of 10μl of protein extract per sample was mixed with 100μl of substrate

solution and incubated at 28°C for 6 min. A blank control containing 10μl of ultra-pure water

was prepared alongside the samples. Substrate solution consisted in 4-MU-chitotrioside

(SIGMA-ALDRICH) diluted in McIlvaine solution at pH 5.2 (1.73mg of 4-MU-chitotrioside per

100 ml of McIlvaine solution). To make the McIlvaine solution, stock solutions of 0.1M citric

acid (SIGMA-ALDRICH) and 0.2M di-sodium hydrogen phosphate (SIGMA-ALDRICH) were

mixed at a ratio of 45% and 55%, respectively, obtaining a final pH of 5.2. After incubation,

1ml of stop solution (0.25M glycine buffer, pH 10.4, containing 0.01% Triton X-100 (SIGMA-

ALDRICH) was added to each sample (and blank control), and 200μl of the final solution was

transferred to a Corning™ Costar™ Brand 96-Well EIA/RIA Plate (Thermo Fisher Scientific).

Duplicates of each sample were used. An extra microcentrifuge tube was prepared with

200µl of standard solution (5µm 1,4-Methylumbelliferone, SIGMA-ALDRICH) and 900µl of

stop solution. A volume of 200µl of the final solution was transferred to the 96-well plate, in

duplicates. The fluorescence was then read in a Varioskan Plate Reader (excitation: 365nm;

emission: 450nm). Finally, the chitotriosidase activity was calculated in nM/h/ml, using

Formula 4.

1147

1148 **T**, fluorescence of test samples

1149 **B**, fluorescence of blank

1150 **Std**, standard

1151
$$\frac{(\boldsymbol{T} - \boldsymbol{B})}{\boldsymbol{std}} \; x \; \frac{60}{incubation\ time\ [min]} \; x \; \frac{1000}{volume\ [\mu l]}$$

1152

12. Blood Brain Barrier (BBB) permeability assay

To test BBB permeability, a total of 5µl of 0.1% IRDye 680RD Dextran (LI-COR Biosciences,

Lincoln, NE, USA) diluted in HBSS (Thermo Fisher Scientific) were injected by intraperitoneal

injection (IP). Two different sizes of the compound were tested, with the following molecular

weights: 4kDa and 70kDa. A group of fish injected with 5µl of HBSS alone was used as a

vehicle control group. The fish were culled 3h after IP, and the brains were collected and

imaged using a LI-COR Odyssey CLx machine (LI-COR). The brain fluorescence was then

quantified using the Image Studio 4.0 Software for Odyssey CLx (LI-COR).

1161

13. Mitochondrial complex assays

Homogenised brain tissue was diluted to approximately 0.7mg protein/ml and subjected to

3 cycles of rapid freeze-thawing in liquid nitrogen. All spectrophotometric assays were

performed at 28°C in 200µl final volume using a Fluostar Omega plate reader. For assessing

complex I activity, the assay buffer was 25mM KH_2PO_4, 5mM $MgCl_2$, (pH7.2), 3mM KCN,

2.5mg/ml BSA, 50µM ubiquinone, 2µg/ml antimycin A. Complex I activity was determined by

measuring the oxidation of nicotinamide adenine dinucleotide reduced disodium salt

(NADH) at 340nm with a reference wavelength of 425nm ($\varepsilon = 6.22 mM^{-1}\ cm^{-1}$). 125µM NADH

started the reaction, and 3µg/ml rotenone was used to inhibit the reaction. For assessing

Complex II activity the brain homogenate was incubated for 10min in assay buffer containing

25mM KH_2PO_4, 5mM $MgCl_2$, (pH 7.2), 3mM KCN, 20 mM succinate. 50µM 2,6-DCPIP, 2µg/ml

antimycin A and 3µg/ml rotenone were then added, and baseline measurements recorded.

The reaction was started with the addition of 50µM ubiquinone Q1 and complex II activity

determined by measuring the reduction of 2,6-dichlorobenzenone-indophenol sodium salt

(2,6-DCPIP) at 600nm ($\varepsilon = 19.2\text{mM}^{-1}\text{ cm}^{-1}$). Complex I and complex II activity rates are

normalised to citrate synthase activity.

1178

14. Novel object (NO) test

Before the experiment, all fish were acclimatised to the behaviour room for at least 1h. The

NO test was performed using a 27x14x14cm tank (length x width x height) containing 7cm of

water depth, with a red 3x1.5x1.5cm object (length x width x height) placed in the centre.

The walls of the test tank were covered in white paper to avoid external visual stimuli. A fish

was gently placed into the test tank with a net and the recording of the test behaviour started

immediately, using a top camera, for a total period of 10min. In the end, the fish was

returned to its tank. The water in the test tank was replaced in between every fish. The assay

was performed for 3 consecutive days and 10 to 20 fish were tested per day. The day and

time of experiment were semi-randomly determined so that all the fish from each group

were equally distributed for the morning and afternoon sessions of each day of experiment

to avoid confounding factors related to the day and schedule of the experiment. Later, fish

behaviour was analysed using the video tracking software EthoVision® XT (Noldus

Information Technology Inc., Lessburg, VA, USA). Anxiety-like behaviour was measured by

the number of entries in the areanear the novel object. Two times the width of the fish was

the measure used to delimit the area considered to be 'near the object'.

1195

15. Statistical analysis

All the graphs and respective statistical analysis were performed with GraphPad Prism 7.0.

Normality was assessed by comparing group variances using either the Brown-Forsythe test

or the Shapiro-Wilk normality test. Unpaired t-test or Mann Whitney's unpaired t-test were

used to compare 2 groups, depending on whether the samples presented a Gaussian

distribution or not, respectively. For groups with normal distribution, comparisons over-time

were performed using ANOVA, followed by Turkey test for multiple comparisons. For groups

that failed the normality tests, non-parametric tests were performed, followed by Dunnett's
test for multiple comparisons. A critical value for significance of $p < 0.05$ was used in all the
analysis.

1206

ACKNOWLEDGEMENTS

We would like to acknowledge and thank Dr Marcus Keatinge for helping with the
chitotriosidase activity assay. We would like to acknowledge and thank Dr William Norton
from the University of Leicester for kindly allowing us to use his EthoVision® XT software
(Noldus). We would like to acknowledge and thank the University of Sheffield Aquarium staff
for the great care they provide to our fish and help with protocols, fish selection and
breeding. We would like to acknowledge and thank the Wolfson Light Microscopy Facility
staff for their support throughout the years with all our imaging needs and challenges.

This work was generously funded by a University of Sheffield PhD studentship to RRM. This
research was funded in whole, or in part, by Wellcome [206224/Z/17/Z/WT_/Wellcome
Trust/United Kingdom]. For the purpose of open access, the author has applied a CC BY
public copyright licence to any Author Accepted Manuscript version arising from this
submission. CMH was also funded by a Vice Chancellor's Fellowship, The University of
Sheffield and a Dunhill Medical Trust grant AISRPG2305\32, which cover PSE salary.

MR is funded by the CNRS and the ANR ADAGIO (ANR-20-CE44-0010). Thanks to the
Bettencourt Schueller Foundation long-term partnership, this work was partly supported by
the CRI Core Research Fellowship to Michael Rera for SB salary.

1224

1225 **CONFLICT OF INTEREST**

1226 The authors declare no competing interests.

1227

1228 **AUTHOR CONTRIBUTIONS**

1229 *Author contributions according to [CRediT](#):*

1230 **RRM**: conceptualisation, data curation, formal analysis, funding acquisition, investigation,
1231 methodology, project administration, validation, visualisation, writing

SB: data curation, formal analysis, investigation, methodology, software, validation, visualisation, writing

PSE: data curation, formal analysis, investigation, methodology, validation, writing

NH: data curation, formal analysis, investigation, methodology

NM: investigation

OE: investigation

MHFW: investigation

ZY: investigation

HM: data curation, formal analysis, funding acquisition, investigation, methodology

MR: data curation, formal analysis, funding acquisition, investigation, methodology, project administration, software, validation, visualisation, writing

CMH: conceptualisation, data curation, formal analysis, funding acquisition, investigation, methodology, project administration, resources, supervision, validation, writing

DATA AVAILABILITY STATEMENT

The dataset and source data generated during and/or analysed during the current study is either shared as supplementary and source data or is available from the corresponding author upon reasonable request. The RNA sequencing data from this experiment were deposited in gene expression omnibus (GEO) and will be made available when in peer review.

ORCID

Raquel R. Martins: https://orcid.org/0000-0003-3952-8649

Savandara Besse: https://orcid.org/0000-0003-2448-4022

Pam S. Ellis: https://orcid.org/0000-0002-5915-4739

Naomi Hartopp: https://orcid.org/0000-0003-4069-6324

Owain Evans: https://orcid.org/0009-0000-8493-4983

1258   *Heather Mortiboys:* https://orcid.org/0000-0001-6439-0579

1259   *Michael Rera:* https://orcid.org/0000-0002-6574-6511

1260   *Catarina M. Henriques:* https://orcid.org/0000-0003-1882-756X

1261

1262   *TOTAL CHARACTERS WITH SPACES: 87566 (includes figure legends. For final submission we will move most of*
1263   *the details in methodology to supplementary information)*

1264

1265

REFERENCES

Acosta, J. C., Banito, A., Wuestefeld, T., Georgilis, A., Janich, P., Morton, J. P., . . . Gil, J. (2013).
A complex secretory program orchestrated by the inflammasome controls paracrine
senescence. *Nat Cell Biol, 15*(8), 978-990. doi:10.1038/ncb2784

Ahmed, S., Passos, J. F., Birket, M. J., Beckmann, T., Brings, S., Peters, H., . . . Saretzki, G. (2008).
Telomerase does not counteract telomere shortening but protects mitochondrial
function under oxidative stress. *J Cell Sci, 121*(Pt 7), 1046-1053.
doi:10.1242/jcs.019372

Anchelin, M., Alcaraz-Perez, F., Martinez, C. M., Bernabe-Garcia, M., Mulero, V., & Cayuela,
M. L. (2013). Premature aging in telomerase-deficient zebrafish. *Dis Model Mech, 6*(5),
1101-1112. doi:10.1242/dmm.011635

Andjelkovic, A. V., Situ, M., Citalan-Madrid, A. F., Stamatovic, S. M., Xiang, J., & Keep, R. F.
(2023). Blood-Brain Barrier Dysfunction in Normal Aging and Neurodegeneration:
Mechanisms, Impact, and Treatments. *Stroke, 54*(3), 661-672.
doi:10.1161/STROKEAHA.122.040578

Baker, D. J., Childs, B. G., Durik, M., Wijers, M. E., Sieben, C. J., Zhong, J., . . . van Deursen, J.
M. (2016). Naturally occurring p16(Ink4a)-positive cells shorten healthy lifespan.
Nature, 530(7589), 184-189. doi:10.1038/nature16932

Basisty, N., Kale, A., Jeon, O. H., Kuehnemann, C., Payne, T., Rao, C., . . . Schilling, B. (2020). A
proteomic atlas of senescence-associated secretomes for aging biomarker
development. *PLoS Biol, 18*(1), e3000599. doi:10.1371/journal.pbio.3000599

Bassi, G. S., Kanashiro, A., Santin, F. M., de Souza, G. E., Nobre, M. J., & Coimbra, N. C. (2012).
Lipopolysaccharide-induced sickness behaviour evaluated in different models of

anxiety and innate fear in rats. *Basic Clin Pharmacol Toxicol, 110*(4), 359-369.

doi:10.1111/j.1742-7843.2011.00824.x

Bhat, R., Crowe, E. P., Bitto, A., Moh, M., Katsetos, C. D., Garcia, F. U., . . . Torres, C. (2012).

Astrocyte senescence as a component of Alzheimer's disease. *PLoS One, 7*(9), e45069.

doi:10.1371/journal.pone.0045069

Birch, J., & Gil, J. (2020). Senescence and the SASP: many therapeutic avenues. *Genes Dev,*

34(23-24), 1565-1576. doi:10.1101/gad.343129.120

Blackburn, E. H., Epel, E. S., & Lin, J. (2015). Human telomere biology: A contributory and

interactive factor in aging, disease risks, and protection. *Science, 350*, 1193-1198.

doi:10.1126/science.aab3389

Bodnar, A. G. (1998). Extension of Life-Span by Introduction of Telomerase into Normal

Human Cells. *Science, 279*(5349), 349-352. doi:10.1126/science.279.5349.349

Bruno, M., Bonomi, C. G., Ricci, F., Di Donna, M. G., Mercuri, N. B., Koch, G., . . . Motta, C.

(2024). Blood-brain barrier permeability is associated with different

neuroinflammatory profiles in Alzheimer's disease. *Eur J Neurol, 31*(1), e16095.

doi:10.1111/ene.16095

Bussian, T. J., Aziz, A., Meyer, C. F., Swenson, B. L., van Deursen, J. M., & Baker, D. J. (2018).

Clearance of senescent glial cells prevents tau-dependent pathology and cognitive

decline. *Nature, 562*(7728), 578-582. doi:10.1038/s41586-018-0543-y

Cao, Y., Li, H., Deb, S., & Liu, J. P. (2002). TERT regulates cell survival independent of

telomerase enzymatic activity. *Oncogene, 21*(20), 3130-3138.

doi:10.1038/sj.onc.1205419

Carneiro, M. C., Henriques, C. M., Nabais, J., Ferreira, T., Carvalho, T., & Ferreira, M. G. (2016).

Short Telomeres in Key Tissues Initiate Local and Systemic Aging in Zebrafish. *PLoS*

Genetics, 12, e1005798. doi:10.1371/journal.pgen.1005798

Chinta, S. J., Woods, G., Demaria, M., Rane, A., Zou, Y., McQuade, A., . . . Andersen, J. K. (2018).

Cellular Senescence Is Induced by the Environmental Neurotoxin Paraquat and

Contributes to Neuropathology Linked to Parkinson's Disease. *Cell Rep, 22*(4), 930-

940. doi:10.1016/j.celrep.2017.12.092

Choi, J., Southworth, L. K., Sarin, K. Y., Venteicher, A. S., Ma, W., Chang, W., . . . Artandi, S. E.

(2008). TERT promotes epithelial proliferation through transcriptional control of a

Myc- and Wnt-related developmental program. *PLoS Genet, 4*(1), e10. doi:10.1371/journal.pgen.0040010

Coder, B. D., Wang, H., Ruan, L., & Su, D. M. (2015). Thymic involution perturbs negative selection leading to autoreactive T cells that induce chronic inflammation. *J Immunol, 194*(12), 5825-5837. doi:10.4049/jimmunol.1500082

Collins, B., Bandosz, P., Guzman-Castillo, M., Pearson-Stuttard, J., Stoye, G., McCauley, J., . . . O'Flaherty, M. (2022). What will the cardiovascular disease slowdown cost? Modelling the impact of CVD trends on dementia, disability, and economic costs in England and Wales from 2020-2029. *PLoS One, 17*(6), e0268766. doi:10.1371/journal.pone.0268766

Coppe, J. P., Desprez, P.-Y. Y., Krtolica, A., Campisi, J., Coppé, J.-P., Desprez, P.-Y. Y., . . . Campisi, J. (2010). The senescence-associated secretory phenotype: the dark side of tumor suppression. *Annual review of pathology, 5*, 99-118. doi:10.1146/annurev-pathol-121808-102144

Cvejic, A., Hall, C., Bak-Maier, M., Flores, M. V., Crosier, P., Redd, M. J., & Martin, P. (2008). Analysis of WASp function during the wound inflammatory response--live-imaging studies in zebrafish larvae. *J Cell Sci, 121*(Pt 19), 3196-3206. doi:10.1242/jcs.032235

d'Adda di Fagagna, F., Reaper, P. M., Clay-Farrace, L., Fiegler, H., Carr, P., Von Zglinicki, T., . . . Jackson, S. P. (2003). A DNA damage checkpoint response in telomere-initiated senescence. *Nature, 426*(6963), 194-198. doi:10.1038/nature02118

De Felice, B., Annunziata, A., Fiorentino, G., Manfellotto, F., D'Alessandro, R., Marino, R., . . . Biffali, E. (2014). Telomerase expression in amyotrophic lateral sclerosis (ALS) patients. *J Hum Genet, 59*(10), 555-561. doi:10.1038/jhg.2014.72

de Lange, T. (2004). T-loops and the origin of telomeres. *Nature reviews. Molecular cell biology, 5*, 323-329. doi:10.1038/nrm1422

Deacon, K., & Knox, A. J. (2018). PINX1 and TERT Are Required for TNF-alpha-Induced Airway Smooth Muscle Chemokine Gene Expression. *J Immunol, 200*(4), 1283-1294. doi:10.4049/jimmunol.1700414

Deleidi, M., Jäggle, M., & Rubino, G. (2015). Immune aging, dysmetabolism, and inflammation in neurological diseases. *Frontiers in neuroscience, 9*, 172. doi:10.3389/fnins.2015.00172

Deverman, B. E., & Patterson, P. H. (2009). Cytokines and CNS development. *Neuron, 64*(1), 61-78. doi:10.1016/j.neuron.2009.09.002

Dimri, G. P., Lee, X., Basile, G., Acosta, M., Scott, G., Roskelley, C., . . . et al. (1995). A biomarker that identifies senescent human cells in culture and in aging skin in vivo. *Proc Natl Acad Sci U S A, 92*(20), 9363-9367.

Dulic, V., Beney, G. E., Frebourg, G., Drullinger, L. F., & Stein, G. H. (2000). Uncoupling between phenotypic senescence and cell cycle arrest in aging p21-deficient fibroblasts. *Mol Cell Biol, 20*(18), 6741-6754. doi:10.1128/MCB.20.18.6741-6754.2000

Eitan, E., Tichon, A., Gazit, A., Gitler, D., Slavin, S., & Priel, E. (2012). Novel telomerase-increasing compound in mouse brain delays the onset of amyotrophic lateral sclerosis. *EMBO Mol Med, 4*(4), 313-329. doi:10.1002/emmm.201200212

Ellett, F., Pase, L., Hayman, J. W., Andrianopoulos, A., & Lieschke, G. J. (2011). mpeg1 promoter transgenes direct macrophage-lineage expression in zebrafish. *Blood, 117*(4), e49-56. doi:10.1182/blood-2010-10-314120

Ellis, P. S., Martins, R. R., Thompson, E. J., Farhat, A., Renshaw, S. A., & Henriques, C. M. (2022). A subset of gut leukocytes has telomerase-dependent "hyper-long" telomeres and require telomerase for function in zebrafish. *Immun Ageing, 19*(1), 31. doi:10.1186/s12979-022-00287-8

Evans, J. R., Torres-Perez, J. V., Miletto Petrazzini, M. E., Riley, R., & Brennan, C. H. (2021). Stress reactivity elicits a tissue-specific reduction in telomere length in aging zebrafish (Danio rerio). *Sci Rep, 11*(1), 339. doi:10.1038/s41598-020-79615-1

Ferreira, A. C., Sousa, N., Sousa, J. C., & Marques, F. (2023). Age-related changes in mice behavior and the contribution of lipocalin-2. *Front Aging Neurosci, 15*, 1179302. doi:10.3389/fnagi.2023.1179302

Ferreira, M. G., Miller, K. M., & Cooper, J. P. (2004). Indecent exposure: when telomeres become uncapped. *Mol Cell, 13*(1), 7-18.

Fleming, A., Diekmann, H., & Goldsmith, P. (2013). Functional characterisation of the maturation of the blood-brain barrier in larval zebrafish. *PLoS One, 8*(10), e77548. doi:10.1371/journal.pone.0077548

Forsyth, N. R., Wright, W. E., & Shay, J. W. (2002). Telomerase and differentiation in multicellular organisms: turn it off, turn it on, and turn it off again. *Differentiation, 69*(4-5), 188-197. doi:10.1046/j.1432-0436.2002.690412.x

Franceschi, C., & Campisi, J. (2014). Chronic inflammation (inflammaging) and its potential contribution to age-associated diseases. *J Gerontol A Biol Sci Med Sci, 69 Suppl 1*, S4-9. doi:10.1093/gerona/glu057

Frenk, S., & Houseley, J. (2018). Gene expression hallmarks of cellular ageing. *Biogerontology, 19*(6), 547-566. doi:10.1007/s10522-018-9750-z

Fridlyanskaya, I., Alekseenko, L., & Nikolsky, N. (2015). Senescence as a general cellular response to stress: A mini-review. *Exp Gerontol, 72*, 124-128. doi:10.1016/j.exger.2015.09.021

Gaikwad, S., Senapati, S., Haque, M. A., & Kayed, R. (2024). Senescence, brain inflammation, and oligomeric tau drive cognitive decline in Alzheimer's disease: Evidence from clinical and preclinical studies. *Alzheimers Dement, 20*(1), 709-727. doi:10.1002/alz.13490

Ganong, W. F. (2000). Circumventricular organs: definition and role in the regulation of endocrine and autonomic function. *Clin Exp Pharmacol Physiol, 27*(5-6), 422-427. doi:10.1046/j.1440-1681.2000.03259.x

Ghosh, A., Saginc, G., Leow, S. C., Khattar, E., Shin, E. M., Yan, T. D., . . . Tergaonkar, V. (2012). Telomerase directly regulates NF-kappaB-dependent transcription. *Nat Cell Biol, 14*(12), 1270-1281. doi:10.1038/ncb2621

Gomes, N. M. V., Ryder, O. a., Houck, M. L., Charter, S. J., Walker, W., Forsyth, N. R., . . . Wright, W. E. (2011). Comparative biology of mammalian telomeres: hypotheses on ancestral states and the roles of telomeres in longevity determination. *Aging Cell, 10*, 761-768. doi:10.1111/j.1474-9726.2011.00718.x

Goodman, W. A., & Jain, M. K. (2011). Length does not matter: A new take on telomerase reverse transcriptase. *Arteriosclerosis, Thrombosis, and Vascular Biology, 31*, 235-236. doi:10.1161/ATVBAHA.110.220343

Haendeler, J., Drose, S., Buchner, N., Jakob, S., Altschmied, J., Goy, C., . . . Dimmeler, S. (2009). Mitochondrial telomerase reverse transcriptase binds to and protects mitochondrial DNA and function from damage. *Arterioscler Thromb Vasc Biol, 29*(6), 929-935. doi:10.1161/ATVBAHA.109.185546

Henriques, C. M., Carneiro, M. C., Tenente, I. M., Jacinto, A., & Ferreira, M. G. (2013). Telomerase is required for zebrafish lifespan. *PLoS Genet, 9*(1), e1003214. doi:10.1371/journal.pgen.1003214

Henriques, C. M., & Ferreira, M. G. (2012). Consequences of telomere shortening during
lifespan. *Curr Opin Cell Biol, 24*(6), 804-808. doi:10.1016/j.ceb.2012.09.007

Henriques, C. M., & Ferreira, M. G. (2024). Telomere length is an epigenetic trait - Implications
for the use of telomerase-deficient organisms to model human disease. *Dis Model*
Mech, 17(3). doi:10.1242/dmm.050581

Hernandez-Segura, A., Nehme, J., & Demaria, M. (2018). Hallmarks of Cellular Senescence.
Trends Cell Biol, 28(6), 436-453. doi:10.1016/j.tcb.2018.02.001

Hernandez-Silva, D., Canton-Sandoval, J., Martinez-Navarro, F. J., Perez-Sanchez, H., de
Oliveira, S., Mulero, V., . . . Cayuela, M. L. (2022). Senescence-Independent Anti-
Inflammatory Activity of the Senolytic Drugs Dasatinib, Navitoclax, and Venetoclax in
Zebrafish Models of Chronic Inflammation. *Int J Mol Sci, 23*(18).
doi:10.3390/ijms231810468

Huang, W., Hickson, L. J., Eirin, A., Kirkland, J. L., & Lerman, L. O. (2022). Cellular senescence:
the good, the bad and the unknown. *Nat Rev Nephrol, 18*(10), 611-627.
doi:10.1038/s41581-022-00601-z

Hudock, J., & Kenney, J. W. (2024). Aging in zebrafish is associated with reduced locomotor
activity and strain dependent changes in bottom dwelling and thigmotaxis. *PLoS One,*
19(5), e0300227. doi:10.1371/journal.pone.0300227

Idda, M. L., McClusky, W. G., Lodde, V., Munk, R., Abdelmohsen, K., Rossi, M., & Gorospe, M.
(2020). Survey of senescent cell markers with age in human tissues. *Aging (Albany NY),*
12(5), 4052-4066. doi:10.18632/aging.102903

Idilli, A. I., Cusanelli, E., Pagani, F., Berardinelli, F., Bernabe, M., Cayuela, M. L., . . . Mione, M.
C. (2020). Expression of tert Prevents ALT in Zebrafish Brain Tumors. *Front Cell Dev*
Biol, 8, 65. doi:10.3389/fcell.2020.00065

Jaskelioff, M., Muller, F. L., Paik, J.-H., Thomas, E., Jiang, S., Adams, A. C., . . . Depinho, R. a.
(2011). Telomerase reactivation reverses tissue degeneration in aged telomerase-
deficient mice. *Nature, 469*, 102-106. doi:10.1038/nature09603

Jeong, J. Y., Kwon, H. B., Ahn, J. C., Kang, D., Kwon, S. H., Park, J. A., & Kim, K. W. (2008).
Functional and developmental analysis of the blood-brain barrier in zebrafish. *Brain*
Res Bull, 75(5), 619-628. doi:10.1016/j.brainresbull.2007.10.043

Jurk, D., Wang, C., Miwa, S., Maddick, M., Korolchuk, V., Tsolou, A., . . . von Zglinicki, T. (2012).
Postmitotic neurons develop a p21-dependent senescence-like phenotype driven by

a DNA damage response. *Aging Cell, 11*(6), 996-1004. doi:10.1111/j.1474-
9726.2012.00870.x

Kacprzak, V., Patel, N. A., Riley, E., Yu, L., Yeh, J. J., & Zhdanova, I. V. (2017). Dopaminergic
control of anxiety in young and aged zebrafish. *Pharmacol Biochem Behav, 157*, 1-8.
doi:10.1016/j.pbb.2017.01.005

Keatinge, M., Bui, H., Menke, A., Chen, Y. C., Sokol, A. M., Bai, Q., . . . Bandmann, O. (2015).
Glucocerebrosidase 1 deficient Danio rerio mirror key pathological aspects of human
Gaucher disease and provide evidence of early microglial activation preceding alpha-
synuclein-independent neuronal cell death. *Hum Mol Genet*.
doi:10.1093/hmg/ddv369

Knox, E. G., Aburto, M. R., Clarke, G., Cryan, J. F., & O'Driscoll, C. M. (2022). The blood-brain
barrier in aging and neurodegeneration. *Mol Psychiatry, 27*(6), 2659-2673.
doi:10.1038/s41380-022-01511-z

Lawson, N. D., Li, R., Shin, M., Grosse, A., Yukselen, O., Stone, O. A., . . . Zhu, L. (2020). An
improved zebrafish transcriptome annotation for sensitive and comprehensive
detection of cell type-specific genes. *eLife, 9*. doi:10.7554/eLife.55792

Lee, H. W., Blasco, M. A., Gottlieb, G. J., Horner, J. W., 2nd, Greider, C. W., & DePinho, R. A.
(1998). Essential role of mouse telomerase in highly proliferative organs. *Nature,*
392(6676), 569-574. doi:10.1038/33345

Lee, J., Jo, Y. S., Sung, Y. H., Hwang, I. K., Kim, H., Kim, S. Y., . . . Lee, H. W. (2010). Telomerase
deficiency affects normal brain functions in mice. *Neurochem Res, 35*(2), 211-218.
doi:10.1007/s11064-009-0044-3

Lemoine, M. (2021). The Evolution of the Hallmarks of Aging. *Front Genet, 12*, 693071.
doi:10.3389/fgene.2021.693071

Lex, K., Maia Gil, M., Lopes-Bastos, B., Figueira, M., Marzullo, M., Giannetti, K., . . . Ferreira,
M. G. (2020). Telomere shortening produces an inflammatory environment that
increases tumor incidence in zebrafish. *Proc Natl Acad Sci U S A, 117*(26), 15066-
15074. doi:10.1073/pnas.1920049117

Lopez-Otin, C., Blasco, M. A., Partridge, L., Serrano, M., & Kroemer, G. (2023). Hallmarks of
aging: An expanding universe. *Cell, 186*(2), 243-278. doi:10.1016/j.cell.2022.11.001

Louveau, A., Smirnov, I., Keyes, T. J., Eccles, J. D., Rouhani, S. J., Peske, J. D., . . . Kipnis, J. (2015). Structural and functional features of central nervous system lymphatic vessels. *Nature, 523*(7560), 337-341. doi:10.1038/nature14432

Lujambio, A. (2016). To clear, or not to clear (senescent cells)? That is the question. *Bioessays, 38 Suppl 1*, S56-64. doi:10.1002/bies.201670910

M, I. F. A. F. C., Selman;, H. A. M. A. L.-C. C., Lene, E. F. E. S. G. J. L. T. J. M., & Bellantuono, J. R. S. X. T. E. I. (2016). Interventions for age-related diseases: shifting the paradigm. *Mech Ageing Dev*(in press).

Martin-Ruiz, C., Williams-Gray, C. H., Yarnall, A. J., Boucher, J. J., Lawson, R. A., Wijeyekoon, R. S., . . . Saretzki, G. (2020). Senescence and Inflammatory Markers for Predicting Clinical Progression in Parkinson's Disease: The ICICLE-PD Study. *J Parkinsons Dis, 10*(1), 193-206. doi:10.3233/JPD-191724

Martins, R. R., Ellis, P. S., MacDonald, R. B., Richardson, R. J., & Henriques, C. M. (2019). Resident Immunity in Tissue Repair and Maintenance: The Zebrafish Model Coming of Age. *Front Cell Dev Biol, 7*, 12. doi:10.3389/fcell.2019.00012

Mathias, J. R., Dodd, M. E., Walters, K. B., Yoo, S. K., Erik, A., & Huttenlocher, A. (2010). Characterization of Zebrafish Larval Inflammatory Macrophages Jonathan. *Developmental and comparative immunology, 33*, 1212-1217. doi:10.1016/j.dci.2009.07.003.Characterization

Mattiussi, M., Tilman, G., Lenglez, S., & Decottignies, A. (2012). Human telomerase represses ROS-dependent cellular responses to Tumor Necrosis Factor-α without affecting NF-κB activation. *Cellular signalling, 24*, 708-717. doi:10.1016/j.cellsig.2011.11.004

Maximino, C., de Brito, T. M., da Silva Batista, A. W., Herculano, A. M., Morato, S., & Gouveia, A., Jr. (2010). Measuring anxiety in zebrafish: a critical review. *Behav Brain Res, 214*(2), 157-171. doi:10.1016/j.bbr.2010.05.031

Meeker, R. B., Williams, K., Killebrew, D. A., & Hudson, L. C. (2012). Cell trafficking through the choroid plexus. *Cell Adh Migr, 6*(5), 390-396. doi:10.4161/cam.21054

Muldoon, L. L., Alvarez, J. I., Begley, D. J., Boado, R. J., Del Zoppo, G. J., Doolittle, N. D., . . . Neuwelt, E. A. (2013). Immunologic privilege in the central nervous system and the blood-brain barrier. *J Cereb Blood Flow Metab, 33*(1), 13-21. doi:10.1038/jcbfm.2012.153

Nolte, E. D., Nolte, K. A., & Yan, S. S. (2019). Anxiety and task performance changes in an aging mouse model. *Biochem Biophys Res Commun, 514*(1), 246-251. doi:10.1016/j.bbrc.2019.04.049

Oosterhof, N., Holtman, I. R., Kuil, L. E., van der Linde, H. C., Boddeke, E. W., Eggen, B. J., & van Ham, T. J. (2017). Identification of a conserved and acute neurodegeneration-specific microglial transcriptome in the zebrafish. *Glia, 65*(1), 138-149. doi:10.1002/glia.23083

Ovadya, Y., & Krizhanovsky, V. (2014). Senescent cells: SASPected drivers of age-related pathologies. *Biogerontology, 15*, 627-642. doi:10.1007/s10522-014-9529-9

Pakulski, C., Drobnik, L., & Millo, B. (2000). Age and sex as factors modifying the function of the blood-cerebrospinal fluid barrier. *Med Sci Monit, 6*(2), 314-318.

Panca, M., Livingston, G., Barber, J., Cooper, C., La Frenais, F., Marston, L., . . . Hunter, R. M. (2019). Healthcare resource utilisation and costs of agitation in people with dementia living in care homes in England - The Managing Agitation and Raising QUality of LifE in Dementia (MARQUE) study. *PLoS One, 14*(2), e0211953. doi:10.1371/journal.pone.0211953

Papageorgakopoulou, M. A., Bania, A., Lagogianni, I. A., Birmpas, K., & Assimakopoulou, M. (2024). The Role of Glia Telomere Dysfunction in the Pathogenesis of Central Nervous System Diseases. *Mol Neurobiol, 61*(8), 5868-5881. doi:10.1007/s12035-024-03947-6

Pegram, R. G., Smith, R., Franklin, R., Gallagher, T., Oosterwijk, G., & Wilsmore, A. J. (1991). Comparison of the calf suckling technique and milk oxytocin test for estimation of milk yields. *Trop Anim Health Prod, 23*(2), 99-102. doi:10.1007/BF02361190

Pereira, B. I., Devine, O. P., Vukmanovic-Stejic, M., Chambers, E. S., Subramanian, P., Patel, N., . . . Akbar, A. N. (2019). Senescent cells evade immune clearance via HLA-E-mediated NK and CD8(+) T cell inhibition. *Nat Commun, 10*(1), 2387. doi:10.1038/s41467-019-10335-5

Petrova, N. V., Velichko, A. K., Razin, S. V., & Kantidze, O. L. (2016). Small molecule compounds that induce cellular senescence. *Aging Cell, 15*(6), 999-1017. doi:10.1111/acel.12518

Rahman, R., Latonen, L., & Wiman, K. G. (2005). hTERT antagonizes p53-induced apoptosis independently of telomerase activity. *Oncogene, 24*(8), 1320-1327. doi:10.1038/sj.onc.1208232

Raj, D. D. A., Moser, J., van der Pol, S. M. A., van Os, R. P., Holtman, I. R., Brouwer, N., . . . Boddeke, H. W. G. M. (2015). Enhanced microglial pro-inflammatory response to lipopolysaccharide correlates with brain infiltration and blood-brain barrier dysregulation in a mouse model of telomere shortening. *Aging Cell*, 1003-1013. doi:10.1111/acel.12370

Ramanathan, R., Kohli, A., Ingaramo, M. C., Jain, A., Leng, S. X., Punjabi, N. M., . . . Fedarko, N. S. (2013). Serum chitotriosidase, a putative marker of chronically activated macrophages, increases with normal aging. *J Gerontol A Biol Sci Med Sci, 68*(10), 1303-1309. doi:10.1093/gerona/glt022

Recagni, M., Bidzinska, J., Zaffaroni, N., & Folini, M. (2020). The Role of Alternative Lengthening of Telomeres Mechanism in Cancer: Translational and Therapeutic Implications. *Cancers (Basel), 12*(4). doi:10.3390/cancers12040949

Redd, M. J., Kelly, G., Dunn, G., Way, M., & Martin, P. (2006). Imaging macrophage chemotaxis in vivo: studies of microtubule function in zebrafish wound inflammation. *Cell motility and the cytoskeleton, 63*, 415-422. doi:10.1002/cm.20133

Righi, D., Manco, C., Pardini, M., Stufano, A., Schino, V., Pelagotti, V., . . . Plantone, D. (2024). Investigating interleukin-8 in Alzheimer's disease: A comprehensive review. *J Alzheimers Dis*, 13872877241298973. doi:10.1177/13872877241298973

Romaniuk, A., Paszel-Jaworska, A., Toton, E., Lisiak, N., Holysz, H., Krolak, A., . . . Rubis, B. (2018). The non-canonical functions of telomerase: to turn off or not to turn off. *Mol Biol Rep*. doi:10.1007/s11033-018-4496-x

Rossiello, F., Herbig, U., Longhese, M. P., Fumagalli, M., & d'Adda di Fagagna, F. (2014). Irreparable telomeric DNA damage and persistent DDR signalling as a shared causative mechanism of cellular senescence and ageing. *Curr Opin Genet Dev, 26*, 89-95. doi:10.1016/j.gde.2014.06.009

Samant, R. R., Standaert, D. G., & Harms, A. S. (2024). The emerging role of disease-associated microglia in Parkinson's disease. *Front Cell Neurosci, 18*, 1476461. doi:10.3389/fncel.2024.1476461

Sarin, K. Y., Cheung, P., Gilison, D., Lee, E., Tennen, R. I., Wang, E., . . . Artandi, S. E. (2005). Conditional telomerase induction causes proliferation of hair follicle stem cells. *Nature, 436*(7053), 1048-1052. doi:10.1038/nature03836

Schaum, N., Lehallier, B., Hahn, O., Palovics, R., Hosseinzadeh, S., Lee, S. E., . . . Wyss-Coray, T. (2020). Ageing hallmarks exhibit organ-specific temporal signatures. *Nature, 583*(7817), 596-602. doi:10.1038/s41586-020-2499-y

Segal-Bendirdjian, E., & Geli, V. (2019). Non-canonical Roles of Telomerase: Unraveling the Imbroglio. *Front Cell Dev Biol, 7*, 332. doi:10.3389/fcell.2019.00332

Spilsbury, A., Miwa, S., Attems, J., & Saretzki, G. (2015). The role of telomerase protein TERT in Alzheimer's disease and in tau-related pathology in vitro. *Journal of Neuroscience, 35*(4), 1659-1674. doi:10.1523/JNEUROSCI.2925-14.2015

Sprungala, S. (2009). Telomere length, telomerase and maintenance of stem cells in the adult zebrafish brain. In. TECHNISCHE UNIVERSITÄT MÜNCHEN.

Steinacker, P., Verde, F., Fang, L., Feneberg, E., Oeckl, P., Roeber, S., . . . group, F. T. s. (2018). Chitotriosidase (CHIT1) is increased in microglia and macrophages in spinal cord of amyotrophic lateral sclerosis and cerebrospinal fluid levels correlate with disease severity and progression. *J Neurol Neurosurg Psychiatry, 89*(3), 239-247. doi:10.1136/jnnp-2017-317138

Strominger, I., Elyahu, Y., Berner, O., Reckhow, J., Mittal, K., Nemirovsky, A., & Monsonego, A. (2018). The Choroid Plexus Functions as a Niche for T-Cell Stimulation Within the Central Nervous System. *Front Immunol, 9*, 1066. doi:10.3389/fimmu.2018.01066

Sullivan, D. I., Jiang, M., Hinchie, A. M., Roth, M. G., Bahudhanapati, H., Nouraie, M., . . . Alder, J. K. (2021). Transcriptional and Proteomic Characterization of Telomere-Induced Senescence in a Human Alveolar Epithelial Cell Line. *Front Med (Lausanne), 8*, 600626. doi:10.3389/fmed.2021.600626

Sung, Y. H., Ali, M., & Lee, H. W. (2014). Extracting extra-telomeric phenotypes from telomerase mouse models. *Yonsei Medical Journal, 55*, 1-8. doi:10.3349/ymj.2014.55.1.1

Sweeney, M. D., Zhao, Z., Montagne, A., Nelson, A. R., & Zlokovic, B. V. (2019). Blood-Brain Barrier: From Physiology to Disease and Back. *Physiol Rev, 99*(1), 21-78. doi:10.1152/physrev.00050.2017

Tabula Muris, C. (2020). A single-cell transcriptomic atlas characterizes ageing tissues in the mouse. *Nature, 583*(7817), 590-595. doi:10.1038/s41586-020-2496-1

Tarr, A. J., Chen, Q., Wang, Y., Sheridan, J. F., & Quan, N. (2012). Neural and behavioral
responses to low-grade inflammation. *Behav Brain Res, 235*(2), 334-341.
doi:10.1016/j.bbr.2012.07.038

Topiwala, A., Nichols, T. E., Williams, L. Z. J., Robinson, E. C., Alfaro-Almagro, F., Taschler, B., .
. . Smith, S. M. (2023). Telomere length and brain imaging phenotypes in UK Biobank.
PLoS One, 18(3), e0282363. doi:10.1371/journal.pone.0282363

Tripathi, U., Misra, A., Tchkonia, T., & Kirkland, J. L. (2021). Impact of Senescent Cell Subtypes
on Tissue Dysfunction and Repair: Importance and Research Questions. *Mech Ageing
Dev, 198*, 111548. doi:10.1016/j.mad.2021.111548

Tuttle, C. S. L., Luesken, S. W. M., Waaijer, M. E. C., & Maier, A. B. (2021). Senescence in tissue
samples of humans with age-related diseases: A systematic review. *Ageing Res Rev,
68*, 101334. doi:10.1016/j.arr.2021.101334

Van Houcke, J., Bollaerts, I., Geeraerts, E., Davis, B., Beckers, A., Van Hove, I., . . . Moons, L.
(2017). Successful optic nerve regeneration in the senescent zebrafish despite age-
related decline of cell intrinsic and extrinsic response processes. *Neurobiol Aging, 60*,
1-10. doi:10.1016/j.neurobiolaging.2017.08.013

Vasconcelos, R. O., Gordillo-Martinez, F., Ramos, A., & Lau, I. H. (2023). Effects of Noise
Exposure and Ageing on Anxiety and Social Behaviour in Zebrafish. *Biology (Basel),
12*(9). doi:10.3390/biology12091165

Vazquez-Villasenor, I., Garwood, C. J., Heath, P. R., Simpson, J. E., Ince, P. G., & Wharton, S. B.
(2020). Expression of p16 and p21 in the frontal association cortex of ALS/MND brains
suggests neuronal cell cycle dysregulation and astrocyte senescence in early stages of
the disease. *Neuropathol Appl Neurobiol, 46*(2), 171-185. doi:10.1111/nan.12559

Vellingiri, B., Balasubramani, K., Iyer, M., Raj, N., Elangovan, A., Song, K., . . . Cho, S. G. (2023).
Role of Telomeres and Telomerase in Parkinson's Disease-A New Theranostics? *Adv
Biol (Weinh), 7*(12), e2300097. doi:10.1002/adbi.202300097

Wang, Y., Kuca, K., You, L., Nepovimova, E., Heger, Z., Valko, M., . . . Jomova, K. (2024). The
role of cellular senescence in neurodegenerative diseases. *Arch Toxicol, 98*(8), 2393-
2408. doi:10.1007/s00204-024-03768-5

Wittenberg, R., Knapp, M., Hu, B., Comas-Herrera, A., King, D., Rehill, A., . . . Kingston, A.
(2019). The costs of dementia in England. *Int J Geriatr Psychiatry, 34*(7), 1095-1103.
doi:10.1002/gps.5113

Wright, D., Nakamichi, R., Krause, J., & Butlin, R. K. (2006). QTL analysis of behavioral and
morphological differentiation between wild and laboratory zebrafish (Danio rerio).
Behav Genet, 36(2), 271-284. doi:10.1007/s10519-005-9029-4

Xu, M., Tchkonia, T., Ding, H., Ogrodnik, M., Lubbers, E. R., Pirtskhalava, T., . . . Kirkland, J. L.
(2015). JAK inhibition alleviates the cellular senescence-associated secretory
phenotype and frailty in old age. *Proc Natl Acad Sci U S A, 112*(46), E6301-6310.
doi:10.1073/pnas.1515386112

Yu, X., Liu, M. M., Zheng, C. Y., Liu, Y. T., Wang, Z., & Wang, Z. Y. (2023). Telomerase reverse
transcriptase and neurodegenerative diseases. *Front Immunol, 14*, 1165632.
doi:10.3389/fimmu.2023.1165632

Zambusi, A., Pelin Burhan, O., Di Giaimo, R., Schmid, B., & Ninkovic, J. (2020). Granulins
Regulate Aging Kinetics in the Adult Zebrafish Telencephalon. *Cells, 9*(2).
doi:10.3390/cells9020350

Zane, F., Bouzid, H., Sosa Marmol, S., Brazane, M., Besse, S., Molina, J. L., . . . Rera, M. (2023).
Smurfness-based two-phase model of ageing helps deconvolve the ageing
transcriptional signature. *Aging Cell, 22*(11), e13946. doi:10.1111/acel.13946

Zhang, M. J., Pisco, A. O., Darmanis, S., & Zou, J. (2021). Mouse aging cell atlas analysis reveals
global and cell type-specific aging signatures. *eLife, 10*. doi:10.7554/eLife.62293

1650